Using eDNA to elucidate Silver and Bighead Carp range expansion in two Missouri River

tributaries in eastern South Dakota

Lindsey A. P. LaBrie[1], Jeff S. Wesner, Hugh B. Britten

Department of Biology, 414 E Clark St., University of South Dakota, Vermillion, SD 57069

Corresponding Author: llabrie@uark.edu

[1] Current Address: Department of Biological Sciences, 507 Science Engineering Hall, University

of Arkansas, Fayetteville, AR 72701

Running head: Invasive carp range expansion in eastern South Dakota

Abstract

Objective: This study used environmental DNA (eDNA) sampling methods to determine the

possibility of Bighead Carp *Hypophthalmichthys nobilis,* and Silver Carp *H. molitrix*, range

expansion above barriers to fish movement in two tributaries of the Missouri River in eastern

South Dakota.

Methods: We collected water samples above and below two perceived barriers to fish

movement: a natural chain of waterfalls in the Big Sioux River and a spillway at the downstream

end of a manmade reservoir in the Vermillion River. We used filtration methods to collect

invasive carp eDNA from water samples and implemented qPCR techniques to quantify the

amount of eDNA in each sample. A total of 213 samples and 38 field blanks were collected over

two years (2021/2022).

Result: We detected carp eDNA below both barriers in both years, and above the barrier in the

Vermillion River in 2022. The probability of detecting eDNA below the barrier in the Big Sioux

River was 51% (95% Credible Interval (CrI): 2 to 97%) and 77% (95% CrI: 8 to 99%) in the

Vermillion River. The probability of detecting eDNA above both barriers was significantly

smaller: 1% (95% CrI: 0.02 to 24%) for both rivers.

Conclusion: The detection of positive samples above the spillway barrier in the Vermillion

River provides the first evidence that Bighead and Silver Carp may have expanded their range to

habitats upstream of their documented range in eastern South Dakota.

Impact Statement: This study demonstrates the utility of using eDNA sampling methods to

detect Bighead and Silver Carp in areas of both known and unknown invasive carp presence in

smaller tributary streams to the Missouri River.

Introduction

    Bighead Carp *Hypophthalmichthys nobilis,* and Silver Carp *H. molitrix*, were introduced

into the southern United States in the early 1970s as a means of biocontrol of algae in

aquaculture ponds and wastewater treatment facilities (Kolar et al. 2005). Soon after their

introduction, both species escaped and began establishing populations throughout the Mississippi

River basin and beyond (Nico et al. 2023a, 2023b). In Midwestern rivers, these species have

become a threat to many freshwater lotic systems since their arrival in the early to mid 1990s

(Kolar et al. 2005) due to their ability to outcompete native planktivorous fishes and alter

ecosystems (Freedman et al. 2012; Pendleton et al. 2017; Phelps et al. 2017). In the Missouri

River Basin, many barriers limit the upstream movement of both invasive and native fishes. The

barriers include dams on the mainstem Missouri and low-head dams and impoundments in the

James, Vermillion, and Big Sioux Rivers (Figure 1). Gavin's Point Dam is a major barrier that

prevents upstream movement of these carp into upstream sections of the Missouri River, but they

may freely swim between the mainstem Missouri and its tributaries in the stretch of river below

Gavin's Point Dam.

    Streams in eastern South Dakota are on the northwestern-most leading edge of the

Bighead and Silver Carp invasion (Hayer et al. 2014). Bighead and Silver Carp first officially

appeared in South Dakota in 1998 and in 2003 respectively (Kolar et al. 2005). In the spring of

2019, above-average snowfall followed by rapid melting and early spring rains caused major

flooding events throughout rivers in the Midwest (Bagwell and Peters 2019; Flanagan et al.

2020). In eastern South Dakota, this flooding impacted multiple tributaries to the Missouri River

(Francis and Melliger 2021), causing them to rise to record-high flood stages. The flooding

allowed potential pathways for the spread of invasive carp above two perceived barriers to

upstream fish movement: a chain of natural waterfalls in Sioux Falls in the Big Sioux River and

a man-made spillway at the downstream end of Lake Vermillion in the Vermillion River. In fact,

carp of an undetermined species were visually confirmed swimming over the spillway in the

Vermillion River (B. Schall, South Dakota Department of Game, Fish, and Parks, personal

communication) immediately after it was compromised during the flooding in 2019. The

presence of Bighead and Silver Carp above these barriers in the Big Sioux and Vermillion Rivers

was previously determined to be nonexistent, but the flooding created new uncertainty

surrounding the invasion front in southeastern South Dakota.

Environmental DNA (eDNA) water sampling has been used to detect Bighead and Silver

Carp in other systems and has especially targeted barriers to fish passage into the Great Lakes

ecosystems (Erickson et al. 2016; Erickson et al. 2017; Jerde et al. 2013; Song et al. 2017), but

there is a lack of its implementation in smaller rivers, especially in the Missouri River Basin.

eDNA is a beneficial tool to use because physically sampling an organism is not required,

therefore reducing operating costs and minimizing disturbance of the ecosystem, while still

providing valuable contemporary evidence for the presence of a species (Thomsen et al. 2012).

eDNA taken from water samples allows for the detection of organisms in waterbodies where

populations are in low or unknown quantities (Lodge et al. 2012). This is especially important

when considering invasive species in novel environments (Erickson et al. 2016). Positive eDNA

detections of invasive organisms in waterbodies where their status was previously unknown can

then trigger management actions to prevent the unchecked spread of the species.

The purpose of this study was to measure the presence and quantity of invasive carp

eDNA below and above two major barriers to fish movement in the Big Sioux and Vermillion

Rivers (a chain of natural waterfalls and a man-made spillway, respectively) to determine if

recent flooding events allowed for Bighead and Silver Carp movement into habitats above those

two barriers. Secondary aims were to evaluate the efficacy of our eDNA sampling regime by 1)

determining the number of samples needed to obtain a positive eDNA detections, 2) comparing

the effects of filtering method on the amount of eDNA in each sample, 3) comparing the effect of

two different extraction protocols on the overall amount of eDNA in each sample, and 4)

determining the overall probability of detecting Bighead and Silver Carp eDNA above and below

each barrier in both rivers. Data collected from this study is intended to help inform local and

basin-wide management efforts to control invasive carp populations in eastern South Dakota and

beyond.

Methods

Study organisms and sites

The Big Sioux River is a fifth-order stream that begins in Roberts County, South Dakota

and flows 674 river kilometers southward through South Dakota to its confluence with the

Missouri River on the South Dakota-Iowa border (Omernik 1987). It is the most populated river

basin in South Dakota (Dieterman and Berry 1999). There exists a major barrier to upstream fish

movement in Sioux Falls—a chain of natural waterfalls called Falls Park. The Vermillion River

is a fifth-order stream that flows 242 river kilometers from the confluence of the East and West

Fork Vermillion Rivers southward through South Dakota to its confluence with the Missouri

River on the South Dakota-Nebraska border (Braaten and Berry Jr 1997). A large spillway

(12.5m high) at the downstream end of Lake Vermillion in the East Fork Vermillion River acts

as a potential barrier to upstream fish movement (Hayer et al. 2014). Both the Big Sioux River

and Vermillion River experience intermittent patterns of flooding and drought, and flows are

highly dependent on the amount of precipitation, including snowmelt, and agricultural runoff

(Underhill, 1959).

Sampling methods

Water sampling took place over two field sampling seasons. In the first field season, at

each sample location, two water samples were collected using two, 2-Liter high-density

polyethylene (HDPE) bottles. In the second field season, one, 2L sample was taken at each

sample location. Prior to sampling, all bottles were sterilized with 20% bleach solution for ≥ 10

seconds and rinsed with distilled water (Coulter et al. 2019). Disposable nitrile gloves were worn

during sample collection and were replaced after each sampling location. The most downstream

location in each river was sampled first, and each successive sample was collected upstream of

the previous sample to prevent eDNA cross-contamination. All samples were collected while

standing on the shoreline to reduce the risk of boots or waders contaminating the water with

invasive carp eDNA. One negative control sample (one, 2-L bottle of distilled water) was

collected above and below each barrier. After sampling, all bottles were placed in coolers on ice

until filtration was carried out no more than 8 hours after sample collection.

The water samples were filtered through 1.5 μm glass microfiber filters (Cytiva Whatman

Binder-Free Glass Microfiber Filters; grade 934-AH circles; 47 mm outside diameter;

borosilicate glass) (Eichmiller et al. 2014) using a magnetic polyphenylsulfone filter funnel

(Eichmiller et al. 2014; Nukazawa et al. 2018; Coulter et al. 2019). All supplies were sterilized

with 20% bleach for ≥ 10 seconds, rinsed with distilled water, and dried with a new paper towel

between samples. Water samples were drawn through the filter until the pores were clogged and

water could no longer pass through the filter, and the volume of water that passed through was

recorded. Each filter was stored in a separate 95% ethanol in a 15 mL polypropylene Falcon tube

at -20°C until eDNA extraction could occur.

eDNA extraction

Prior to extraction, filters were cut in half, with one half returned to the 15 mL Falcon

tube in the freezer and the other half cut into three smaller pieces. The pieces were placed in 1.5

mL microcentrifuge tubes and allowed to air dry in a fume hood over a period of ≥ 4 hours until

all ethanol had evaporated. Tools were sterilized with 20% bleach for ≥ 10 seconds, rinsed with

distilled water, and then rinsed with nanopure water to prevent contamination between filters.

eDNA extraction was performed in a designated sequencing room at the University of South

Dakota and according to the Qiagen DNeasy Blood and Tissue Extraction protocol (Qiagen

2023) with one modification at the end step to centrifuge 100 μL of elution buffer through the

spin columns twice instead of 200 μL once (i.e., "Kristie" protocol (Schmidt et al. 2021)).

Extracted samples were stored in 1.5 mL or 2 mL microcentrifuge tubes at -20 °C.

Quantitative Polymerase Chain Reaction (qPCR) analysis

The real-time quantitative polymerase chain reaction (qPCR) analysis was conducted

using an Applied Biosystems™ QuantStudio™ 3 Real-Time PCR System thermocycler (Thermo

Fisher). The forward primer, reverse primer, and gBlock used in this study were developed by

Erickson et al. (2017) and targeted the mitochondrial gene *nd5*, where the primer sequence binds

to an area of the mitochondrial genome that is conserved between both Bighead and Silver Carp,

and the probe sequence binds to an area of the mitochondrial genome that is specific to either

species (Erickson et al. 2017). Each well had a volume of 20 μL, containing 7 μL molecular

grade water, 10 μL TaqMan™ Fast Advanced Master Mix (Thermo Fisher), 1 μL of 10

nanomole forward and reverse primer mix, 1 μL of 5 nanomole probe mix, and 1 μL of extracted

eDNA sample. A 313-base pair Megamer ssGene Fragment, or gBlock, containing DNA

sequences conserved between both Bighead and Silver Carp was used to create a standard

dilution series which served as the eDNA calibration curve for each qPCR run (Erickson et al.

2017). The 96-well plates contained four replicates of each of the following: eight field samples,

one field blank, one extraction blank and one positive control. Positive controls were extracted

with the DNeasy Qiagen Blood and Tissue Kit from Bighead and Silver Carp tissues. Each plate

also contained three spiked replicates of each sample (extracted sample plus 0.1 µL of the 10^3

dilution per well). Ten, no-template control wells (1 µL nanopure water) and two replicates of a

10^6 standard dilution series derived from the G-block megamer were included on each plate

(Table 1). qPCR cycling conditions followed the protocol in Erickson et al. (2017). We

considered each sample to have been positive if at least one sample replicate amplified during

qPCR, and negative if none of the replicates amplified.

Pilot study

First, to evaluate the number of water sample replicates needed to obtain a positive

detection in areas of known Bighead and Silver Carp presence, we took five, 2-L samples in

September 2022 at one location in the Big Sioux River (42.495167, -96.477236) and one

location in the Vermillion River (42.772914, -96.930374), where the USFWS was

contemporaneously performing their eDNA sampling and Dozer trawling for invasive carp.

Second, to determine if eDNA quantities varied when following the Qiagen extraction protocol

versus the modified extraction protocol (i.e., "Kristie" protocol (Schmidt et al. 2021)), we

extracted one half of each filter using the Qiagen protocol and the other half using the "Kristie"

protocol (Schmidt et al. 2021) and then analyzed the samples.

Statistical analysis

We evaluated whether filtering method (i.e., field vs lab) caused a difference in the

probability of eDNA detection. We used Program R (version 4.1.1; R Core Team 2022) and the

brms package (Bürkner 2017) to fit a Bayesian generalized linear mixed model with a Bernoulli

likelihood with a logit link with the following model and default priors:

176                          $E_i \sim \text{Bernoulli}(p_i),$

177                     $\text{logit}(p_i) = \alpha + \boldsymbol{\beta X} + \gamma_{\text{sample\_name}}$

Where E_i is the presence or absence of eDNA in sample i, p_i is the probability of detection, α is

the intercept, $\boldsymbol{\beta X}$ is a vector of 3 parameters representing two predictors ($\mathbf{X}$ = filter method,

species) and their interaction and $\gamma_{\text{sample\_name}}$ is the random intercept for sample name.

To evaluate the probability of detection as a function of species, river, and location in

respect to the barriers in each river, we again used a Bernoulli likelihood with a logit link using

the following equation:

184                          $E_i \sim \text{Bernoulli}(p_i),$

185                $\text{logit}(p_i) = \alpha + \boldsymbol{\beta X} + \gamma_{\text{sample\_name}} + \gamma_{\text{species}}$

186                          $\alpha \sim \text{Normal }(-10,1)$

187                          $\boldsymbol{\beta X} \sim \text{Normal }(0,1)$

188                          $\gamma \sim \text{Student-t}(3, 0, 2.5)$

Where E_i, p_i, and α are as defined above, $\boldsymbol{\beta X}$ is a vector of 7 parameters representing three

predictors (quantity, location, river) and their interactions, and $\gamma_{\text{sample\_name}}$ and γ_{species} are the

random intercepts for sample name and species, respectively.

Both models were run using a Hamiltonian Monte Carlo No-U-Turn sampler with 4

Markov chains, 2,000 iterations per chain with a warm-up of 1,000 iterations. We evaluated the

fit of the model by running posterior predictive checks and determined it properly converged if

Gelman-Rubin statistics were less than 1.1 (Brooks and Gelman 1998). We sampled from the

posterior and estimated the relationships for the model and predicted the mean with 95% credible

intervals. We ran a sensitivity analysis to assess the impact of the priors on our presence/absence

model by doubling the standard deviation values to Normal (-10,2) for the intercept, Student's t

(3, 0, 5) for the random intercepts, and Normal (0,2) for the beta values.

We also visualized how other factors may have affected the quantity of eDNA in our

samples using *ggplot2* (Wickham et al. 2016), such as the length of time between sample

collection and filtering, sample month, the volume of water filtered, the number of days between

filtering and extraction, and the number of days between extraction and qPCR.

Results

Water sampling

A total of 213 water samples and 38 field blanks were collected after two years of field

sampling. (98 water samples and 20 field blanks in 2021, 115 water samples and 18 field blanks

in 2022). In 2021, one-half of the water samples were filtered in the field using a 7.2-liter,

manual pump MityVac MIT7400 fluid evacuator (Figure S-1a). The other half of the samples

were transported back to the lab and filtered using a 1000 mL Büchner flask (Figure S-1b). In

2022, samples were exclusively filtered using the field filtering method. The average filtered

volume was 382 mL (range 100 to 1650 mL). Each year, samples were collected from sampling

sites in both rivers three separate times. In 2021, Big Sioux River samples were collected from

three below-barrier sites and six above-barrier sites in three sampling sessions. This was

expanded to five below-barrier sites and eleven above-barrier sites in 2022 (Figure S2). In 2021,

Vermillion River samples were taken at two below-barrier sites and seven above-barrier sites.

This was expanded to four below-barrier sites and fourteen above-barrier sites in 2022 (Figure

S3).

Invasive carp were observed swimming below the barriers at almost all sampling

sessions, except for the below-barrier Vermillion River samples in October and September 2021,

where only fish carcasses of various species, including Bighead and Silver Carp, were observed

in the water below the spillway. In 2022, we visually observed carp at all water sampling

sessions below both barriers.

qPCR results

Bighead and Silver Carp eDNA was detected below both barriers in 2021 and 2022 in

varying quantities (Table 3). In July 2022, Bighead and Silver Carp eDNA was detected in two

and three samples respectively above the Lake Vermillion spillway (Figure S4), but not in

previous or successive samples. The probability of detecting eDNA below the barrier in the Big

Sioux River was 51% (95% Credible Interval (CrI): 2 to 97%) and in the Vermillion River was

77% (95% CrI: 8 to 99%). The probability of detecting eDNA above the barriers was

significantly smaller: 1% (95% CrI: 0.02 to 22%) in the Big Sioux River and 1% (95% CrI: 0.02-

24%) in the Vermillion River. The probability that detection was higher below the barriers than

above the barriers was greater than 99% (Figure 2).

There was little difference between the mean quantity of eDNA obtained from field and

lab filtering methods, with less than one-hundredth of one percent difference between field and

lab filtered samples (0.003%, 95% CrI: 0.001 to 0.6%) (Figure 3). This was the basis for our

choice to only perform field filtration in the second sampling season. Through pilot study

sampling, we determined that one, 2L water sample at each sampling location was sufficient to

obtain a positive detection for both Bighead and Silver Carp in both rivers where eDNA was

present, with a 10% probability of false negative detection. The probability of false negatives

halved when more samples were taken at each location (i.e., 5% for two samples per location,

2.5% for three samples per location, etc.). The extraction protocol type (i.e.,"Qiagen" versus

"Kristie") appeared to have an impact on mean eDNA quantities, which reinforced our decision

to follow the modified extraction protocol described in the methods section above. Other factors

such as the length of time between sample collection and filtering, sample month, the volume of

water filtered, the number of days between filtering and extraction, and the number of days

between extraction and qPCR did not appear to have any strong effect on the amount of eDNA

detected with qPCR (Supplementary Figures 5-9).

Model performance

Both models performed well, and observed data was within the predicted range of the

posterior estimates. Trace plots indicated proper mixing of the Markov chains, all R-hat values

were < 1.001, and all parameter estimates had large effective sample sizes. The Bayesian R^2

value for the filter method model was 0.82 (95% credible interval (CrI): 0.73 to 0.89) and for the

presence/absence model was 0.76 (95% CrI: 0.67 to 0.84). Posterior predictive checks (Berkhof

et al. 2000) generally replicated the shape of the data for both models. The sensitivity analysis

showed model parameter estimates were not sensitive to the priors (Figure 4).

Discussion

This study was the first to use eDNA to detect Bighead and Silver Carp above barriers to

upstream movement in Missouri River tributaries in eastern South Dakota. We were able to

demonstrate the utility of the filtration method to quantify eDNA from water samples within

these two rivers. The filtration method used in this study has the potential to be more accessible

than other eDNA sampling methods. The manual pneumatic pump and magnetic filter funnel are

relatively inexpensive, easily portable, and can be deployed on-site and allow for immediate

filtration upon water sample collection. We determined that filtering water from one 2-L sample

was sufficient to detect eDNA at least 90% of the time when both species of carp are present.

The number of days between filtration and extraction, or the number of days between extraction

and qPCR analysis did not appear to affect the quantity of eDNA detected in each positive

sample.

The probability of detecting eDNA in our samples below the barriers was high (>54%)

and the probability above the barriers was low but greater than zero (<1%) in both rivers. We

obtained three positive eDNA detections above the spillway in the Vermillion River, within Lake

Vermillion in June 2022, but not in previous or successive samples. We are confident that

positive samples above the spillway were true positives because there was no evidence of eDNA

contamination in the field blanks or qPCR no-template controls. There have been no official

reports of invasive carp above the spillway, however it is possible that invasive carp have been

persisting in Lake Vermillion in relatively low numbers and that we simply did not sample in

areas where they were present during the other rounds of sampling. The presence of boats on

Lake Vermillion during our July 2022 sampling session could explain the positive detections

above the spillway barrier. Invasive carp DNA could have sloughed off the hull of a boat that

had previously been in carp-positive waters, resulting in positive detections. Another explanation

is that invasive carp initiate spawning movements in early summer when the hydrograph begins

to rise (Erickson et al. 2016). Because eDNA has short persistence times in freshwater

ecosystems (Eichmiller et al. 2014), it is possible that the invasive carp made an upstream

spawning run out of the lake and into the Vermillion River immediately prior to sampling. Both

could offer potential explanations for the positive samples above the barrier within the lake.

South Dakota is predicted to experience an increase in climate extremes, including

persistent severe droughts, at least a 15% increase in annual precipitation, and an increase in the

intensity and interval of severe flooding (Frankson et al. 2022). The effects of flooding have

been evident in the recent past during the flood of 2019, when Falls Park became inundated with

water and the spillway at Lake Vermillion became compromised, thus providing potential

pathways for invasive carp to pass barriers to fish movement. Future eDNA water sampling or

physical sampling methods (i.e., capture of live individuals via electrofishing) will be required to

confirm the presence of live invasive carp above the barriers in both rivers. Water sampling

could be expanded to other small tributaries to assess the presence of these fish in finer detail

throughout eastern South Dakota. It may also be necessary to eventually reconsider the

effectiveness of these barriers during flooding events.

Conclusion

This study provides evidence that eDNA is a useful technology to implement in smaller

tributary systems on the leading edge of the current invasive carp range. We determined that

filtering one, 2-L sample per site provided a true positive eDNA detection at least 90% of the

time. We established a baseline for fish presence above barriers in the Big Sioux and Vermillion

Rivers, however, we recommend continued monitoring to increase the level of certainty of fish

presence above the barriers in both rivers, especially in the Vermillion River. We recommend

future eDNA water sampling to further elucidate the status of these species above the barriers,

coupled with physical sampling efforts to gauge invasion presence and risk into the future.

Acknowledgements

We thank T Devine and J Larsen for their assistance with water sampling and processing,

and K Schmidt for demonstrating the qPCR protocol. Dr. A Coulter provided comments and

revisions to the manuscript. Funding for this study was provided South Dakota Game Fish and

Parks (Grant number F-16-R-1) and the United States Fish and Wildlife Service (Grant number

F20AP11067-00).

Conflict of Interest

The authors declare no conflicts of interest. The use of trade names does not constitute

endorsement of the product or firm being described.

Data Availability Statement

Data and R code used in this manuscript are available in the GitHub repository:

https://github.com/lindseylabrie/carp_eDNA

Ethics Statement

This work was conducted under an approved protocol from the University of South Dakota's

Institutional Animal Care and Use Committee (protocol # 10-08-20-23D).

Tables

Table 1: A general 96-well plate set up for detecting Bighead and Silver Carp eDNA. "NTC"

stands for "no template control."

	1	2	3	4	5	6	7	8	9	10	11	12
A	Sample 1	Sample 2	Sample 3	Sample 4	Sample 5	Sample 6	Sample 7	Sample 8	Field blank	Extraction blank	10^6	10^6
B	Sample 1	Sample 2	Sample 3	Sample 4	Sample 5	Sample 6	Sample 7	Sample 8	Field blank	Extraction blank	10^5	10^5
C	Sample 1	Sample 2	Sample 3	Sample 4	Sample 5	Sample 6	Sample 7	Sample 8	Field blank	Extraction blank	10^4	10^4
D	Sample 1	Sample 2	Sample 3	Sample 4	Sample 5	Sample 6	Sample 7	Sample 8	Field blank	Extraction blank	10^3	10^3
E	NTC	NTC	NTC	NTC	NTC	NTC	NTC	NTC	NTC	NTC	10^2	10^2
F	Spiked sample 1	Spiked sample 2	Spiked sample 3	Spiked sample 4	Spiked sample 5	Spiked sample 6	Spiked sample 7	Spiked sample 8	Spiked field blank	Spiked extraction blank	10^1	10^1
G	Spiked sample 1	Spiked sample 2	Spiked sample 3	Spiked sample 4	Spiked sample 5	Spiked sample 6	Spiked sample 7	Spiked sample 8	Spiked field blank	Spiked extraction blank	Positive control	Positive control
H	Spiked sample 1	Spiked sample 2	Spiked sample 3	Spiked sample 4	Spiked sample 5	Spiked sample 6	Spiked sample 7	Spiked sample 8	Spiked field blank	Spiked extraction blank	Positive control	Positive control

325   Table 2: Average eDNA quantification (copies/µL) for each river, above and below barriers, for

326   each species in both years.

2021		Bighead Carp	Silver Carp
Vermillion River	Above	0	0
	Below	555	0.009
Big Sioux River	Above	0	0
	Below	18.4	1.54

2022		Bighead Carp	Silver Carp
Vermillion River	Above	**0.302**	**0.00960**
	Below	24.1	1.53
Big Sioux River	Above	0	0
	Below	2.74	0.391

Figures

Figure 1: A map of barriers to upstream fish movement in the Missouri, James, Vermillion, and

Big Sioux Rivers. Dams on the mainstem Missouri River are shown as purple diamonds, poor

barriers to fish movement such as low-head dams and rock impoundments in the James River are

shown as green triangles (not all barriers are shown at the current map scale), and the putative

barriers in the Big Sioux and Vermillion Rivers are shown as red pentagons.

Figure 2: Posterior predictions for the probability of a positive sample for Bighead Carp (BHC)

and Silver Carp (SVC) in the Big Sioux and Vermillion Rivers above and below the major

barriers to upstream fish movement. The hollow, colored circles represent data points, with zero

(0) on the Y-axis indicating no eDNA detection and one (1) on the Y-axis indicating a positive

eDNA detection. The probability of a difference between above and below barrier samples was

greater than 99%.

Figure 3: Estimated posterior probability of detecting eDNA from Bighead Carp (BHC) and

Silver Carp (SVC) in both rivers (combined) for field and lab filtered samples. The colored

circles represent actual data points, with zero (0) on the y-axis indicating no eDNA detection and

one (1) on the y-axis indicating a positive eDNA detection. Note that the y-axis is on a log(10)

scale.

Figure 4: A comparison of the original model estimates (A) and the sensitivity analysis (B) for

which the standard deviation of each prior was doubled. The colored circles represent actual data

points, with zero (0) on the y-axis indicating no eDNA detection and one (1) on the y-axis

indicating a positive eDNA detection.

References

Bagwell R, Peters B. 2019. Analysis of the 2019 Midwest US Flooding using NASA data. Paper

presented at the AGU Fall Meeting Abstracts.

Berkhof J, van Mechelen I, Hoijtink H. 2000. Posterior predictive checks: Principles and

discussion. Comput. Stat. 15(3):337-354. doi:10.1007/s001800000038

Braaten PJ, Berry Jr CR. 1997. Fish associations with four habitat types in a South Dakota

prairie stream. J. Freshw. Ecol. 12(3):477-489.

Brooks S, Gelman A. 1998. General Methods for Monitoring Convergence of Iterative

Simulations. J. Comput. Graphi. Stat. 7:434-455. doi:10.1080/10618600.1998.10474787

Bürkner PC. 2017. brms : An R Package for Bayesian Multilevel Models Using Stan. J. Stat.

Softw. 80. doi:10.18637/jss.v080.i01

Coulter DP, Wang P, Coulter AA, Van Susteren GE, Eichmiller JJ, Garvey J E, Sorensen PW.

2019. Nonlinear relationship between Silver Carp density and their eDNA concentration

in a large river. PLoS One, 14(6):e0218823.

Dieterman D, Berry C. 1999. Fish community and water quality changes in the Big Sioux River.

Prairie Naturalist. 30:199-224.

Eichmiller JJ, Bajer PG, Sorensen PW. 2014. The relationship between the distribution of

common carp and their environmental DNA in a small lake. PLoS One. 9(11):e112611.

Erickson RA, Merkes CM, Jackson CA, Goforth RR, Amberg JJ. 2017. Seasonal trends in eDNA

detection and occupancy of bigheaded carps. J. Great Lakes Res. 43(4):762-770.

doi:https://doi.org/10.1016/j.jglr.2017.06.003

Erickson RA, Rees CB, Coulter AA, Merkes CM, McCalla SG, Touzinsky KF, Walleser L,

Goforth RR, Amberg JJ. 2016. Detecting the movement and spawning activity of

bigheaded carps with environmental DNA. Mol. Ecol. Resour. 16(4):957-965.

Flanagan PX, Mahmood R, Umphlett NA, Haacker E, Ray C, Sorensen W, Shulski M, Stiles CJ,

Pearson D, Fajman P. 2020. A hydrometeorological assessment of the historic 2019 flood

of Nebraska, Iowa, and South Dakota. Bull Am Meteorol Soc. 101(6):E817-E829.

Frankson R, Kunkel KE, Champion SM, Easterling DR, Umphlett NA, Stiles CJ. 2022 South

Dakota State Climate Summary 2022. NOAA Technical Report NESDIS 150-SD.

NOAA/NESDIS, Silver Spring, MD, 5 pp.

Francis B, Melliger J. 2021. 2019 Missouri River Basin Historic Flood Event. Journal of Dam

Safety. 18(1):20–28.

Freedman JA, Butler SE, Wahl DH. 2012. Impacts of Invasive Asian Carps on Native Food

Webs. Final project report to Illinois–Indiana Sea Grant. University of Illinois, Sullivan.

Hayer CA, Breeggemann J, Klumb R, Graeb B, Bertrand K. 2014. Population characteristics of

bighead and silver carp on the northwestern front of their North American invasion.

Aquat. Invasions. 9:289-303. doi:10.3391/ai.2014.9.3.05

Jerde CL, Chadderton WL, Mahon AR, Renshaw MA, Corush J, Budny ML, Mysorekar S,

Lodge DM. 2013. Detection of Asian carp DNA as part of a Great Lakes basin-wide

surveillance program. Can. J. Fish. Aquat. Sci. 70(4):522-526.

Kolar CS, Chapman DC, Courtenay Jr WR, Housel CM, Williams JD, Jennings DP. (2005).

Asian Carps of the Genus Hypophthalmichthys (Pisces, Cyprinidae) ― A Biological

Synopsis and Environmental Risk Assessment.

Lodge DM, Turner CR, Jerde CL, Barnes MA, Chadderton L, Egan SP, Feder JL, Mahon AR,

Pfrender ME. 2012. Conservation in a cup of water: estimating biodiversity and

population abundance from environmental DNA. Mol. ecol. 21(11):2555-2558.

Nico LG, Fuller P, Li J. 2023a. Hypophthalmichthys molitrix (Valenciennes in Cuvier and

Valenciennes, 1844). Nonindigenous Aquatic Species Database. Retrieved from

https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=549

Nico LG, Fuller P, Li J. 2023b. Hypophthalmichthys nobilis (Richardson, 1845). Nonindigenous

Aquatic Species Database. Retrieved from

https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=551

Nukazawa K, Hamasuna Y, Suzuki Y. 2018. Simulating the advection and degradation of the

environmental DNA of common carp along a river. Environ. Sci. Technol. 52(18):10562-

10570.

Omernik JM. 1987. Ecoregions of the conterminous United States. Ann Assoc Am Geogr. 77(1):

118-125.

Pendleton RM, Schwinghamer C, Solomon LE, Casper AF. 2017. Competition among river

planktivores: are native planktivores still fewer and skinnier in response to the Silver

Carp invasion? Environ. Biol. Fishes. 100:1213-1222.

Phelps QE, Tripp SJ, Bales KR, James D, Hrabik RA, Herzog DP. 2017. Incorporating basic and

applied approaches to evaluate the effects of invasive Asian Carp on native fishes: A

necessary first step for integrated pest management. PLoS One, 12(9):e0184081.

Qiagen, Inc. 2023. DNeasy Blood & Tissue Handbook. Retrieved from

https://www.qiagen.com/us/resources/resourcedetail?id=68f29296-5a9f-40fa-8b3d-

1c148d0b3030&lang=en

R Core Team. 2021. R: A Language and Environment for Statistical Computing. Vienna,

Austria. Retrieved from https://www.R-project.org/

Rees HC, Maddison BC, Middleditch DJ, Patmore JR, Gough KC. 2014. The detection of

aquatic animal species using environmental DNA–a review of eDNA as a survey tool in

ecology. J. Appl. Ecol. 51(5):1450-1459.

Schmidt KJ, Soluk DA, Maestas SEM, Britten HB. 2021. Persistence and accumulation of

environmental DNA from an endangered dragonfly. Sci. Rep. 11(1):18987.

Song JW, Small MJ, Casman EA. 2017. Making sense of the noise: The effect of hydrology on

silver carp eDNA detection in the Chicago area waterway system. Sci. Total Environ.

605-606, 713-720. doi:https://doi.org/10.1016/j.scitotenv.2017.06.255

Stan Development Team. 2023. RStan: the R interface to Stan. Retrieved from https://mc-

stan.org/

Thomsen PF, Kielgast JO, Iversen LL, Wiuf C, Rasmussen M, Gilbert MT, Orlando L,

Willerslev E. 2012. Monitoring endangered freshwater biodiversity using environmental

DNA. Mol. ecol. 21(11):2565-2573.

Underhill JC. 1959. Fishes of the Vermillion River, South Dakota. Paper presented at the Proc

SD Acad Sci.

USFWS (U.S. Fish and Wildlife Service). 2022. Quality Assurance Project Plan (QAPP) eDNA

monitoring of bighead and silver carps. U.S. Fish and Wildlife Service, Midwest Region.

Bloomington, Minnesota, USA. https://www.fws.gov/sites/default/files/documents/

eDNA-QAPP-2022-Whitney-Genetics-Lab.pdf