Cross-talk between tissues is critical for intergenerational acclimation to environmental
change

Sneha Suresh[1], Megan J Welch[2], Philip L Munday[2], Timothy Ravasi[3,4], Celia Schunter[1,5*]

[1]Swire Institute of Marine Science and School of Biological Sciences, The University of
Hong Kong, Hong Kong SAR

[2]Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook
University, Townsville, Australia

[3]Marine Climate Change Unit, Okinawa Institute of Science & Technology Graduate
University, Onna-son, Japan

[4]Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook
University, Townsville, Australia
14
15 [5]State Key Laboratory of Marine Pollution, City University of Hong Kong, Hong Kong,
16 Hong Kong SAR, China
17
18
19 *Corresponding author

20 Correspondence: celiaschunter@gmail.com
21

Abstract

An organism's reaction to environmental changes is mediated by coordinated responses of multiple tissues. Additionally, parental priming may increase offsprings' acclimation potential to changing environmental conditions. As the effects of human-induced climate change, such as ocean acidification (OA), continue to intensify it is critical to assess the acclimation potential of species at the whole organismal scale. For this we need to understand the cross-talk between tissues in regulating and responding to pH changes. Here by using a multi-tissue approach we determine the influence of 1) variation in parental behavioural tolerance and 2) parental environment, on molecular responses of their offspring in a coral reef fish. The gills and liver showed the highest transcriptional response to OA conditions in juvenile fish regardless of the parental environment, while the brain and liver showed the greatest signal of intergenerational acclimation. Key functional pathways that were altered in the brain upon within-generational CO_2 exposure were restored to control levels when parents were exposed to OA conditions. Furthermore, the expression of a new complement of genes involved in key functions was altered in the offspring only when parents were previously exposed to OA conditions. Therefore, previous parental conditioning to OA can reprogram tissue transcriptomic profiles of the offspring enabling them to better cope in an environment with elevated CO_2 levels. Overall, our results reveal tissue-specific transcriptional changes underlying intergenerational plastic responses to elevated CO_2 exposure and highlight the integration of these changes in promoting organismal acclimation to OA.

Keywords: Climate change, multi-tissue, intergeneration, acclimation, ocean acidification, transcriptomics, spiny damselfish.

Introduction

With the global climate continuously shifting to more extreme conditions organisms need to acclimate and/or adapt to the changing environments in order to survive. The oceans are becoming increasingly acidified as they absorb a major portion of anthropogenic CO_2 emissions (Pörtner et al., 2022) leading to ocean acidification (OA) which is reported to negatively impact the physiology and behavior of various marine organisms including fish (Heuer & Grosell, 2014; Strader et al., 2020). However, increasing evidence suggests that multi-generational exposure to elevated CO_2 conditions could influence the adaptive capacity of future generations to OA conditions (Nagelkerken et al., 2023). In fact, several studies have reported transgenerational acclimation in a number of fish species as well as some invertebrates to OA (Strader et al., 2020). Specifically, transgenerational exposure to elevated CO_2 conditions has been shown to facilitate acclimation of metabolism, growth, survival, neuronal plasticity and behavior in independent studies (Allan et al., 2014; Miller et al., 2012; Monroe et al., 2021; Munday, 2014; Schade et al., 2014; Schunter et al., 2016, 2018; Stiasny et al., 2018) however, we are still learning about the underlying molecular mechanisms of such acclimation processes.

Additionally, variation both within and across species in the biological responses to OA also exists due to differences in their evolutionary and environmental history. Studies examining the effect of elevated CO_2 on metabolism, growth, development, and reproduction in fish show variable results with some species being more affected than others (Heuer & Grosell, 2014). Variation in sensitivity to elevated CO_2 within a population could be crucial in long-term adaptation through selection of more tolerant individuals. Indeed, individual variation in behavioural tolerance to elevated CO_2 exposure has been reported to be heritable and hence could facilitate rapid selection of tolerant genotypes in the population (Lehmann et al., 2022; Welch & Munday, 2017). Such selection for CO_2 tolerance has been shown to occur

in nature, which could result in populations consisting of individuals with greater behavioural

tolerance to elevated CO_2 (Munday et al., 2013). Furthermore, inter-individual variation in

sensitivity to ocean acidification could have an epigenetic basis (Ryu et al., 2018; Turner, 2009)

and in fact several studies have reported the expression levels of genes involved in epigenetic

processes to be altered upon exposure to elevated CO_2 conditions (Huang et al., 2019; Schunter

et al., 2018). Transfer of epigenetic factors from parents to offspring (epigenetic inheritance)

could be one of the potential mechanisms of inter- and trans-generational acclimation and

eventual adaptation to OA.

79   Adaptive processes to environmental changes at the organismal level require

integrated activity of various tissues, with each tissue undergoing changes in its transcriptional

landscape resulting in the overall response of the organism. However, to date, research has

mainly focused on individual tissue functional changes in response to OA with less emphasis

on how these changes integrate to create a whole-body response. Several studies have

examined the effects of OA on brain and neurosensory systems since the discovery of impaired

behavioural responses in various fish species in elevated CO_2 conditions. The altered

behavioural responses have been linked to changes in the functioning of the GABAergic

signaling pathway (Schunter et al., 2019) and the circadian rhythm in the brain of fish exposed

to elevated CO_2 (Lee et al., 2021; Schunter et al., 2016; Williams et al., 2019). Previous studies

have also focused on the effects of OA on the gill transcriptome due to it being the primary

organ involved in acid-base regulation, immune defense, and stress response, and hence plays

a vital role in maintaining cellular homeostasis under conditions of CO_2 stress (De Souza et al.,

2014; Deigweiher et al., 2008, 2010). These processes are energetically expensive and indeed

changes in the aerobic metabolic scope (Crespel et al., 2019; Gräns et al., 2014; Pimentel et al.,

2014; Rummer et al., 2013) and expression levels of key metabolic genes (Frommel et al., 2020)

have been reported in fish exposed to elevated CO_2. Therefore, exposure to elevated CO_2

affects various aspects of fish physiology such as metabolism, cellular redox status, ion

transport and acid-base homeostasis, neurological functioning and behavior thereby exerting a

whole-body functional reprogramming (Grosell et al. 2019). Therefore, a systematic

transcriptomic analysis is needed to determine how the biological processes associated with

each tissue integrate together within the whole organism to drive adaptive responses to elevated

CO_2 environments.

In this study we conducted an intergenerational CO_2 exposure experiment and

performed systematic analysis of gene expression changes in response to elevated CO_2 across

three tissues, the brain, the gills, and the liver, in the spiny damselfish *Acanthochromis*

polyacanthus. While this species can be sensitive to increases in water temperature and CO_2

levels, they have the potential to acclimate to the changing environmental conditions across

multiple generations (Donelson et al., 2012; Schunter et al., 2016, 2018; Veilleux et al., 2015).

A. polyacanthus has been used as a model to study the impacts of climate change, and to

investigate the molecular basis of intergenerational plasticity to environmental changes, due to

its advantageous life-history traits for laboratory studies (Robertson, 1973), however past

studies have only focused on single tissues (Ryu et al., 2018; Schunter et al., 2016, 2018). Here,

by using a multi-tissue transcriptomic approach we aim to determine how dynamic cross-talk

between tissues maintains whole-body homeostasis under future ocean acidification conditions.

Additionally, we also assess how the acclimatory response of offspring mediated by

transcriptional reprogramming across multiple tissues is influenced by variation in parental

sensitivity to elevated CO_2 and parental environment. Through systemic characterization of the

effects of OA we aim to identify how the adaptive processes within each tissue integrate to

drive intergenerational acclimation to OA at the organismal level.

Methods

Sample collection, behavioural testing, and experimental design

Adult *Acanthochromis polyacanthus* were collected from the wild on the Great Barrier Reef, Australia (18°38'24.3"S, 146°29'31.8"E) and exposed to elevated CO_2 (754 ± 92 µatm) for seven days following which their behavioural sensitivity to conspecific chemical alarm cues (CAC) was tested using a two-chamber flume as described previously (Schunter et al., 2016). Briefly, the fish were classified as being behaviorally sensitive or tolerant to elevated CO_2 based on the amount of time spent in water containing the CAC. Sensitive individuals spent ≥ 70% time in CAC whereas tolerant individuals spent ≤ 30% time in CAC. Individuals of similar size displaying the same behavioural phenotype (sensitive or tolerant) were then grouped into breeding pairs and held in either control (414 ± 46 µatm) or elevated CO_2 conditions (754 ± 92 µatm) for three months prior to the breeding season. Offspring clutches from each breeding pair were placed into three different experimental treatments resulting in three combinations of parent-offspring conditions for each parental phenotype: (1) Control treatment – Parents and offspring held at control condition (414 ± 46 µatm); (2) Developmental treatment – Parents held at control condition and offspring exposed to elevated CO_2 (754 ± 92 µatm) immediately after hatching; and (3) Intergenerational treatment – Parents and offspring exposed to elevated CO_2 (754 ± 92 µatm). The offspring were held in their respective conditions until they were five months old after which nine fish from each parental phenotype, from each treatment condition (N = 27 from each parental phenotype; N = 54 total fish sampled) were euthanized and the brain, gills and liver were dissected, snap frozen in liquid nitrogen and stored at -80 ºC until further processing (Supplementary Figure S1).

RNA extraction, sequencing, and gene expression analyses

Total RNA was extracted from the fish brains, livers and gills using the AllPrep

DNA/RNA Mini kit from Qiagen following the manufacturer's instructions. RNA quality was

determined using nanodrop and Agilent Bioanalyzer and samples having an RNA integrity

value (RIN) $\geq$ 8 were sequenced using Illumina HiSeq 2500 to get paired-end reads of 100 bp

at Macrogen Inc., South Korea. A total of 1,614.25 $\pm$ 3.05, 2,367.03 $\pm$ 5.19, and 2,227.23 $\pm$

6.37 million raw paired-end reads were obtained from the 162 sequenced libraries from brain,

gills and liver, respectively which included nine control, nine developmental and nine

intergenerational samples for each parental phenotype for each tissue (Supplementary Table

S1). The quality of the raw reads were examined using FastQC (Andrews, 2010) v0.11.8 and

adapters and low quality sequences were trimmed using Trimmomatic (Bolger et al., 2014)

v0.39 (ILLUMINACLIP: adapters.fa:2:30:15:8:true; SLIDINGWINDOW:4:20; MINLEN:32).

Only those sequences $\geq$ 32 bp in length with both the forward and reverse reads retained after

trimming were used for further analysis. Potential contaminant sequences were identified using

kraken (Wood & Salzberg, 2014) v2.0.8-beta, with a confidence score of 0.3, using the bacteria,

fungi and virus RefSeq genomic libraries as reference and removed from further analyses. A

total of 1,510.51 $\pm$ 2.62, 2,254.16 $\pm$ 4.95, and 2,116.25 $\pm$ 6.04 million high-quality sequences

were retained after the filtering process (Supplementary Table S1). These sequences were

mapped to the *Acanthochromis polyacanthus* reference genome (unpublished) using HISAT2

(Kim et al., 2019) v2.1.0. On average, 84 $\pm$ 1.83%, 91.22 $\pm$ 0.66%, and 93.33 $\pm$ 0.81% reads

mapped to the reference genome from the brain, gills, and liver respectively (Supplementary

Table S1). Raw read counts per gene were obtained using featureCounts (Liao et al., 2014)

v2.0.0 (parameters: -B -J -M --fraction), assigning fractional counts to multi-mapped reads.

Exploring the gene expression patterns across the whole dataset (162 samples) using principal

component analysis (PCA) revealed a clear clustering of samples by tissues indicating that

tissues vary greatly in their gene expression patterns (Supplementary Figure S2). Subsequent

analysis of differences in gene expression levels was therefore carried out separately for each

tissue using the DESeq2 (Love et al., 2014) v1.32.0 package in R (R Core Team, 2021) v4.2.1.

Principal component analysis (PCA) using the regularized log transformed (rlog)

counts was done in R v4.2.1 to detect and remove outlier samples. A likelihood ratio test (LRT)

using a model comparison approach was then used to determine the effect of family line in

driving the gene expression patterns and to determine the best design formula for the final DE

analysis. First, significant differences in gene expression were measured by comparing a model

including treatment and family line against a reduced model without the family line factor

separately for each tissue. For a total of 924, 910, and 923 genes in the brain, gills, and liver

respectively, the model including family line better explained the observed differences in gene

expression compared to the reduced model excluding this factor (FDR corrected p-value < 0.05;

Supplementary Table S2). Pair-wise comparisons between the control, developmental and

intergenerational treatment was then caried out in DESeq2 (accounting for the family effect,

using Wald test) separately for each parental phenotype for each tissue to determine the effect

of parental environment and parental tolerance to CO_2 on the molecular responses of the

offspring to elevated CO_2. For each pair-wise comparison, the genes were considered to be

significantly differentially expressed (DE) if the False Discovery Rate (FDR) adjusted p-value

was less than 0.05, the absolute log 2-fold change in expression was greater than 0.3 and

baseMean was greater than 10. Functional enrichment analysis of the significant DE genes was

carried out in OmicsBox (https://www.biobam.com/omicsbox) v1.4.11 using Fisher's Exact

Test (FDR corrected p-value < 0.05) with the option of reducing to most specific GO terms to

reduce redundancy. The genes associated with the enriched GO terms were further categorized

into broader functional groups based on their functional description from the UniProt

knowledgebase (UniProtKB; https://www.uniprot.org/). All figures are made using ggplot in

R v4.2.1.

Results

Molecular processes affected by all elevated CO_2 treatments

To understand a general effects of elevated CO_2 exposure regardless of time of exposure

to elevated CO_2, we identified genes that were commonly differentially expressed (DE) in both

the developmental and intergenerational treatments compared to control (Figure 1(a)(i)), which

are considered the general "CO_2 response genes". Irrespective of the parental environment, the

liver showed the greatest transcriptional response to elevated CO_2 exposure in offspring of both

sensitive and tolerant parental behavioral phenotypes, with 63 and 986 differentially expressed

(DE) genes, respectively (Figure 1(b); Supplementary Table S4). In the offspring of tolerant

parental phenotype, the gills followed with 883 DE genes, while the brain had 44 DE genes

(Figure 1(b)). Conversely, in the offspring of sensitive parental phenotype, the brain displayed

a higher number of DE genes (37) compared to the gills (9). Overall, offspring of parents with

tolerant behavioural phenotype showed increased capacity for gene expression regulation

across all three tissues in response to elevated CO_2 exposure. However, the CO_2 response genes

showed high tissue specificity with no genes being commonly DE across the three tissues in

the sensitive parental phenotype and only 3 and 81 genes being shared between the brain and

gills, and liver and gills respectively in the tolerant parental phenotype (Supplementary Figure

S3(a)).

Figure 1: (a) Schematic graph representing the expression profile of (i) CO_2-response genes, (ii) genes showing a rescue pattern, and (iii) intergeneration-specific genes. (b) Number of differentially expressed genes across the three treatments in all the three tissues. SP indicates samples with a sensitive parental phenotype and TP indicates samples with a tolerant parental phenotype for each of the respective tissues. Note scale break in y-axis at 100 DE genes.

Functional enrichment analysis of the CO_2 response genes in offspring with sensitive parental phenotype revealed translation and amino-acid synthesis to be important in the brain, while no significantly enriched functions were found in the gills and liver (Figure 2; Supplementary Table S5). For the offspring of tolerant parental phenotype, biosynthetic processes was commonly enriched in the brain and gills (Figure 2). Few other functional pathways were commonly enriched between the gills and liver, including metabolism, translation, protein transport, immune and stress response (Figure 2; Supplementary Table S5). Interestingly, while genes associated with metabolic pathways such as glycolysis, TCA cycle

and mitochondrial electron transport chain were commonly differentially regulated in both the

gills and liver, pentose phosphate pathway gene showed upregulation exclusively in the gills.

Additionally, immune response and translation showed tissue-specific regulation with

downregulation in the liver but upregulation in the gills (Supplementary Table S5).

Furthermore, RNA processing and primary active transmembrane transport were enriched only

in the liver (Supplementary Table S5). This suggests intricate and tissue-specific regulation of

various functions in response to elevated CO_2 exposure.

Figure 2: Functions that are significantly enriched (FDR <0.05) among the DE genes involved in overall CO_2 response. SP indicates offspring of behaviorally sensitive parents and TP indicates offspring of behaviorally tolerant parents. The color of the circles represents the FDR corrected p-value and size of the circles represents the number of genes associated with the respective function.

Parental exposure to elevated CO_2 "rescues" developmental effects

In the developmental treatment (compared to control and intergeneration) a total of 883, 55, and 108 genes that were DE in offspring with sensitive parental phenotype and 458, 284, and 434 genes that were DE in offspring with tolerant parental phenotype in the brain, gills, and liver respectively, returned to control levels when parents were previously exposed to OA conditions suggesting cross-generation plasticity resulting in "rescue" of gene expression levels (Figure 1(a)(ii); Supplementary Table S6). Parental conditioning had the largest effect on brain gene expression followed by the liver (Figure 1(b)). Similar to the CO_2-response genes, there were very few genes commonly DE across tissues (Supplementary Figure S3(b)).

Genes exhibiting a rescue pattern in the brain of offspring of behaviorally sensitive parents showed the highest degree of specificity in functional regulation with pathways involved in cytoskeleton, apoptosis, synaptic signaling, sodium and calcium channel activity, transcription regulation, cellular transport, and small GTPase signaling being exclusively enriched only in this tissue (Figure 3; Supplementary Table S7). Gills were found to play a key role in regulating ion transport in offspring of both behaviorally sensitive and tolerant parents. Specifically, the expression of genes encoding sodium and potassium channel transporters were altered in the gills upon developmental exposure to elevated CO_2 levels but returned to control levels in intergenerationally exposed fish (Figure 3; Supplementary Table S6, S7). Additionally, purine ribonucleoside salvage pathway and DNA replication were identified as key functions among the "rescue" genes in the offspring of parents with sensitive phenotype in the gills and liver respectively (Figure 3; Supplementary Table S7). This suggests that the molecular response of the offspring is influenced by the parental environment with specific functional

263   pathways with parental conditioning to elevated CO_2 resulting in selective regulation of

264   specific functions in each tissue.

265
Figure 3: Functions that are significantly enriched (FDR <0.05) among the DE genes showing
a rescue pattern. SP indicates offspring of behaviorally sensitive parents and TP indicates
offspring of behaviorally tolerant parents. The color of the circles represents the FDR corrected
p-value and size of the circles represents the number of genes associated with the respective
function.

Intergenerational specific response to elevated CO_2

We found a large transcriptional response to elevated CO_2 that was only seen in the

intergenerationally exposed fish and not in fish with only developmental (within generation)

exposure to elevated CO_2. This indicates plasticity of the offspring transcriptome due to

parental conditioning to elevated CO_2 and was especially marked in offspring of tolerant

parents (Figure 1(b)). These are genes that were only differentially expressed in the

intergenerational treatment (compared to control and development) but were at control levels

in the developmental treatment (Figure 1(a)(iii)). Specifically, 383, 207, and 1,226 genes were

DE in the brain, gills, and liver respectively in offspring with tolerant parents, while offspring

with sensitive parents had 36, 7, and 8 DE genes in the brain, gills, and liver respectively that

were specific to the intergenerational treatment (Supplementary Table S8). Offspring of both

tolerant and sensitive parents showed high tissue specificity in the intergenerational specific

transcriptional signature to elevated CO_2 with only one and five genes being commonly DE

across all three tissues in the sensitive and tolerant parental phenotypes respectively

(Supplementary Figure S3(c)).

The intergeneration specific response was substantially higher in offspring of

behaviourally tolerant parents with various functions showing tissue-specific regulation. Stress

response and metabolic pathways were commonly differentially regulated in the brain and liver

(Figure 4), but the liver showed a much larger number of DE genes (~400) associated with

metabolism compared to the brain (12). Additionally, signal transduction pathway was

enriched only in the brain while several functions such as biosynthetic pathways, endopeptidase

activity, lipid binding/ transport, transcriptional regulation, proteolysis, and spindle

organization were exclusive to the liver (Figure 4, Supplementary Table S9). Interestingly,

molecular signatures indicating pH regulation by bicarbonate retention were identified only in

the intergenerationally treated fish with tolerant parental phenotype, indicated by the

downregulation of SLC4A1, CFTR, SLC12A2, and SLC9A3 in the gills and upregulation of

SLC4A4 in the brain (Supplementary Table S6). Therefore, offspring of parents with tolerant

behavioural phenotype showed increased molecular signatures of intergenerational plasticity

and this was especially evident in the liver. This change in transcriptional signature could

regulate the above mentioned "rescue" pattern and equip offspring to better cope with OA

conditions.

Figure 4: Functions that are significantly enriched (FDR <0.05) among the intergeneration specific DE genes in offspring of behaviorally tolerant parents. The color of the circles represents the FDR corrected p-value and size of the circles represents the number of genes associated with the respective function.

Parental variability in CO_2 sensitivity impacts the offspring transcriptome

Parental behavioural phenotype was found to have a substantial influence on the offspring transcriptional response. Across all three tissues, offspring with behaviorally tolerant parents when faced with elevated CO_2 had larger changes in gene expression levels (log2FC > 5; Figure 5) and a greater number of DE genes involved in the overall CO_2 response (common in developmental and intergenerational CO_2 exposure compared to control; Figure 1(b)). There were also very few genes involved in overall CO_2 response shared between the two parental phenotypes, specifically only six, three, and eleven common DE genes in the brain, gills, and liver respectively (Supplementary Figure S3(a)). When considering genes involved in intergenerational plastic responses, there were more differentially expressed genes in the offspring of tolerant parents compared to those of sensitive parents, except for genes showing a rescue pattern in the brain (Figure 1(b). Additionally, none of the DE genes involved in intergenerational plasticity were shared in the liver tissue between the two parental phenotypes while the brain and gills had a small proportion of common DE genes (specifically, 121 and 11 common DE genes showing a rescue pattern in the brain and gills respectively and only two and one common DE genes in the intergeneration specific response in the brain and gills respectively (Supplementary Figure S3(b, c)).

The difference in transcriptional response between the two parental phenotypes was especially pronounced when considering genes showing an intergeneration-specific signature with offspring of tolerant phenotype having more DE genes with a much higher magnitude of gene expression changes (log2FC >5; Figure 5). The liver showed the greatest signal of intergenerational response (Figure 4). Overall, while there were some common transcriptional responses between offspring with sensitive and tolerant parents, offspring of tolerant parents showed a much stronger transcriptional response to elevated CO_2 in general and also had a stronger signature of intergenerational plasticity.

335
336 **Figure 5:** Log2 fold change in expression of genes involved in (a) overall CO_2 response, (b)
337 rescue pattern, and (c) intergenerational specific response across all three tissues. SP indicates
338 samples with a sensitive parental phenotype and TP indicates samples with a tolerant parental
339 phenotype.
340

Discussion

Changes in the offspring transcriptional landscape in response to elevated CO_2 exposure exhibited tissue-specific signatures which were largely influenced by previous parental exposure to elevated environmental CO_2, but also by the parental behavioural phenotype. Specifically, we found that gills are critical in maintaining overall cellular homeostasis in elevated CO_2 conditions, by regulating pH disturbances and defending against stress-induced cellular damage and potential infections, and the brain and liver had the greatest signal of intergenerational plastic response. In fact, intergenerationally treated fish no longer showed molecular signatures of altered neural signaling in the brain that were seen in the developmental (within generation) treatment. Additionally, differential regulation of a new complement of metabolic genes exclusively in the brain and liver of offspring from parents with prior exposure to elevated CO_2 implies that parental conditioning enhances the offspring's capacity to regulate metabolic processes to meet potential shifts in metabolic demands arising from ocean acidification conditions. *A. polyacanthus* is known to have a highly plastic genome enabling it to respond and acclimate to environmental changes (Bernal et al., 2020; Kang et al., 2022) and our results show that this persists across generations potentially enabling this species to rapidly acclimate to the changing ocean environment.

Genes that are always differentially expressed in elevated CO_2 conditions regardless of the type of exposure are key genes in the general response to ocean acidification (OA). The gills and liver exhibited a higher transcriptional response in all elevated CO_2 treatments suggesting that these tissues play an important role in the overall response of the fish to elevated CO_2. In both the developmental and intergenerational CO_2 treatments, key functional pathways such as cellular metabolism, stress and immune response were differentially regulated in these tissues. Exposure to elevated CO_2 conditions is known to induce stress and immune response pathways (reviewed in Strader et al., 2020). Interestingly, in our study while stress response

genes were upregulated in both gills and liver, genes involved in immune response were

upregulated in the gills but downregulated in the liver. This indicates that both the gills and

liver work in synergy to mitigate potential cellular damage from oxidative stress and maintain

cellular redox balance. However, given that gills are one of the major surface tissues

continuously exposed to the external environment, and also serve as a first line of defense

against potential infections (Harper & Wolf, 2009; Hu et al., 2023), selective activation of

immune responses in this tissue alone could be a preventive measure against opportunistic

infections under CO_2 stress as has been previously observed under elevated CO_2 conditions

(De Souza et al., 2014; Machado et al., 2020). Similar differences in tissue-specific regulation

of specific metabolic pathways were observed between the gills and liver in all elevated CO_2

treatments. Specifically, genes associated with the tricarboxylic acid (TCA) cycle were

upregulated in both tissues, but pentose phosphate pathway was upregulated only in the gills.

Interestingly, although TCA cycle genes were upregulated in the liver, various genes part of

the cytochrome complex, an integral component of the electron transport chain, and genes

associated with glycolysis like, Fructose-bisphosphate aldolase B and enolase, were

downregulated in the liver. This suggests that there is potentially an overall shift in metabolic

priorities and adjustment of ATP production to potentially facilitate redirection of metabolic

carbon fluxes to meet cellular demands for various metabolites and cofactors needed for other

biological processes such as biosynthetic pathways and the cellular stress response machinery

(Gansemer et al., 2020; Rokitta et al., 2012; Walsh & Milligan, 1993). Overall, our results

indicate conserved pathways in the gills and liver that are always activated in response to

elevated CO_2 exposure irrespective of previous parental conditions.

Parental exposure to altered environmental conditions can pre-acclimate the offspring

transcriptome to these new conditions via intergenerational plasticity. In juvenile *A.*

polyacanthus, developmental exposure to elevated CO_2 resulted in altered expression of genes

involved in key functional pathways which were restored to control levels in the

intergenerationally treated fish. This "rescue pattern" was predominantly evident in the brain

of offspring with sensitive parental phenotype. Specifically, the expression of genes involved

in synaptic plasticity and calcium channel activity, initially altered in the developmental

treatment, were similar to control levels in the intergenerationally treated fish. Exposure to OA

conditions has been shown to affect neural plasticity and neurogenesis in some fish species,

which could result in changes in neural circuitry and signaling (Costa et al., 2022; Lai et al.,

2017). While neural plasticity could facilitate increased flexibility to environmental changes

(Ebbesson & Braithwaite, 2012), it could result in behavioural alterations that have been

observed in fish exposed to OA conditions (Schunter et al., 2019). Increased GABAergic

signaling is a commonly observed within-generation response to elevated CO_2 in *A.*

polyacanthus (Schunter et al., 2018) and a similar increase in neural signaling pathways was

found in the olfactory epithelium of European seabass (*D. labrax*) even after prolonged

transgenerational OA exposure for two generations (Cohen-Rengifo et al. 2022). The

restoration of OA induced changes in neural signaling processes with previous parental

exposure to elevated CO_2 could indicate elevated intergenerational plasticity of *A.*

polyacanthus. In fact, a comparison of six species living in wild CO_2 seeps with long-term CO_2

exposures revealed *A. polyacanthus* to have substantially larger brain transcriptional plasticity

compared to the other species (Kang et al. 2022). The 'rescue pattern' of synaptic plasticity

genes in intergenerationally treated fish was also accompanied by a similar pattern in the

regulatory processes, underlying this plasticity. These regulatory processes are driven by

various cell adhesion and cytoskeleton genes that play a role in synaptic plasticity (Gordon-

Weeks & Fournier, 2014; Zapara et al., 2000). Therefore, intergenerational CO_2 exposure

potentially restores the dynamic equilibrium of cytoskeletal proteins, thereby re-establishing

synaptic signaling processes to control levels.

Additionally, genes involved in transcriptional regulation such as transcription factors

and RNA-mediated gene silencers also showed a "rescue pattern" in the brain. Changes in the

external environment can trigger reprogramming of transcriptional networks resulting in

dynamic regulation of gene expression (Swift & Coruzzi, 2017) as also suggested in wild fish

populations naturally exposed to elevated CO_2 (Petit-Marty et al., 2021). Therefore, these

regulatory genes could play a key role in developmental plastic responses to elevated

environmental CO_2 levels, however, these are no longer needed in intergenerationally treated

fish revealing acclimation. In the gills, on the other hand, this 'rescue' pattern was seen for

genes encoding ion transporters in offspring of both behaviorally sensitive and tolerant parents.

Notably, potassium channels, known for their significance in pH regulation and association

with CO_2 chemoreception in the gills (Qin et al., 2010), displayed differential expression in the

developmental treatment, possibly triggered by hypercapnia. This, in turn, could prompt

downstream compensatory responses to maintain pH homeostasis (Suresh et al., 2023, Qin et

al., 2010). However, the restoration of expression levels of these genes to control levels in

intergenerationally treated fish suggests that intergenerational exposure may enhance and

facilitate improved pH regulatory mechanisms.

We also found a new complement of genes to be differentially expressed only when the

parents are exposed to the same elevated CO_2 condition as their offspring, which could further

facilitate acclimation of the offspring enabling them to better cope with an elevated CO_2

environment. Such intergenerational specific transcriptional signature was predominantly

observed in the brain and liver of offspring of behaviorally tolerant parents, notably in genes

associated with metabolism. Furthermore, genes involved in ion transport leading to

bicarbonate retention showed intergeneration-specific differential expression, predominantly

in the gills and to a lesser extent in the brain of fish with tolerant parental phenotype. Increasing

internal bicarbonate ion concentrations is a commonly observed compensatory response in fish

to buffer acid-base disturbance caused by exposure to elevated CO_2 levels (Heuer & Grosell,

2014). Differential expression of bicarbonate transporters in the intergenerationally treated fish

suggests that parental conditioning enables the offspring to more effectively buffer pH changes

(Brauner et al., 2019; Chen et al., 2009; Choi, 2012.; Esbaugh, 2017). Therefore,

intergenerational CO_2 exposure could have resulted in transcriptional rearrangements of ion

transporters in the offspring, especially in the brain and gills, to maintain pH homeostasis

primarily by bicarbonate retention.

Parental conditioning to elevated CO_2 also increased the offspring's capacity of energy

metabolism indicated by upregulation of genes involved in fatty-acid metabolism in the brain

and liver. Additionally, lipid synthesis was also upregulated in the liver. Changes in lipid

metabolism has been shown to facilitate acclimation to elevated CO_2 as well as temperature in

previous studies (Liu et al., 2013, Strader et al., 2020, Veilleux et al., 2015). Therefore,

upregulation of lipid metabolism in both the brain and liver in intergenerationally treated fish

combined with the observed redirection of metabolic carbon fluxes observed when considering

the overall CO_2 response, might indicate a similar switch from carbohydrate to lipid

metabolism as the primary metabolic pathway for ATP production in elevated CO_2 conditions,

although further research is needed to confirm this. Furthermore, genes involved in cellular

stress response were upregulated in the brain of intergenerationally treated fish with

behaviorally tolerant parents. Notably, the cellular stress response pathway was upregulated in

the gills and liver in both developmental and intergenerational CO_2 treatment, but in the brain

only in the intergenerational CO_2 treatment. Given that cellular stress response pathway is an

energy demanding process (Strader et al., 2020), the systemic upregulation of this function in

all three tissues only in the intergenerationally treated fish could be facilitated by the increased

capacity of energy metabolism.

We also found a parental behavioural phenotype to influence the acclimation potential

of offspring to elevated CO_2 conditions. Offspring of tolerant parental phenotype showed a

stronger transcriptional response to elevated CO_2 exposure with higher number of DE genes

and larger fold changes across all three tissues. Furthermore, offspring of tolerant parental

phenotype showed increased intergenerational plastic responses compared to offspring of

behaviorally sensitive parents, with the brain and liver showing the largest signal of

intergenerational effects. Parental behavioural phenotype has been shown to influence the

offspring brain transcriptional response to elevated CO_2 in *A. polyacanthus* (Monroe et al.,

2021), with behavioural tolerance being heritable (Welch & Munday, 2017). Selection

experiments have indicated a genetic basis for individual variation in OA induced responses in

a variety of animals (Langer et al. 2009; Parker et al., 2011; Pistevos et al. 2011; Sunday et al.

2011), including the behavioural phenotype to chemical alarm cues in elevated CO_2 in *A.*

polyacanthus (Lehmann et al. 2022). This intraspecific variation in organismal response is key

in driving future adaptive evolution. Our results suggest that offspring of parents with a tolerant

behavioural phenotype have an increased capacity for intergenerational plasticity and potential

acclimation to future ocean acidification conditions.

This study used a systematic approach considering parental behaviour, environment,

and multiple tissues to assess molecular responses to future ocean conditions. Gills and liver

work together to reduce stress-induced cellular damage in offspring of tolerant parents under

all CO_2 treatments, while in the brain cellular stress response is upregulated only in

intergenerationally treated fish. Therefore, parental conditioning to elevated CO_2 potentially

enhances offspring's stress response at a systemic level. Additionally, prior parental exposure

to OA mediates the reprogramming of the offspring transcriptome, impacting energy

metabolism in the brain and liver of tolerant-parent offspring and resulting in rescue of

altered synaptic signaling pathways, observed in the developmental treatment, in the brain of

offspring with behaviourally sensitive parents. Overall, our study reveals how

intergenerational plasticity is facilitated from a whole-organism perspective and illustrates how

transcriptional changes across multiple tissues integrate to drive the plastic responses of fish to

the changing ocean chemistry.

Author contributions

The experiment was designed and run by MJW and PLM. Molecular lab work was performed by CS and sequenced by TR. SS carried out the transcriptome expression analysis with input from CS. SS lead the writing of the manuscript with input from CS and all authors read, edited and approved the final manuscript.

Ethics

Sample collection was carried out following all institutional and national law guidelines. The experiment was completed under James Cook University ethics approval A1828.

Competing financial interests

All authors declare they have no competing interests.

Acknowledgements

CS and SS were supported through the HKU start-up to CS. PLM was supported by the ARC Centre of Excellence for Coral Reef Studies and TR was supported by Okinawa Institute of Science and Technology (OIST).

Data availability

The brain RNA-Seq raw sequences are deposited in NCBI under BioProject ID PRJNA311159. The gills and liver RNA-Seq raw sequences are deposited in NCBI under BioProject ID PRJNA989422 (https://dataview.ncbi.nlm.nih.gov/object/PRJNA989422?reviewer=q3n0q75hbbf2p1v85n0h e3uv86).

References

Allan, B. J. M., Miller, G. M., Mccormick, M. I., Domenici, P., & Munday, P. L. (2014). Parental effects improve escape performance of juvenile reef fish in a high-CO_2 world. *Proceedings of the Royal Society B: Biological Sciences*, 281(1777). https://doi.org/10.1098/RSPB.2013.2179

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

Bernal, M. A., Schunter, C., Lehmann, R., Lightfoot, D. J., Allan, B. J. M., Veilleux, H. D., Rummer, J. L., Munday, P. L., & Ravasi, T. (2020). Species-specific molecular responses of wild coral reef fishes during a marine heatwave. *Science Advances*, 6(12). https://doi.org/10.1126/sciadv.aay3423

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30(15), 2114–2120. https://doi.org/10.1093/BIOINFORMATICS/BTU170

Brauner, C. J., Shartau, R. B., Damsgaard, C., Esbaugh, A. J., Wilson, R. W., & Grosell, M. (2019). Acid-base physiology and CO_2 homeostasis: Regulation and compensation in response to elevated environmental CO_2. In *Fish Physiology* (Vol. 37, pp. 69–132). Academic Press. https://doi.org/10.1016/BS.FP.2019.08.003

Chen, J. H., Cai, Z., & Sheppard, D. N. (2009). Direct Sensing of Intracellular pH by the Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) Cl Channel*♦. *Journal of Biological Chemistry*, 284(51), 35495-35506. https://doi.org/10.1074/jbc.M109.072678

Choi, I. (2012). SLC4A transporters. In *Current topics in membranes* (Vol. 70, pp. 77-103). Academic Press.

Cohen-Rengifo, M., Danion, M., Gonzalez, AA. et al. The extensive transgenerational transcriptomic effects of ocean acidification on the olfactory epithelium of a marine fish are associated with a better viral resistance. *BMC Genomics* 23, 448 (2022). https://doi.org/10.1186/s12864-022-08647-w

Costa, R. A., Olvera, A., Power, D. M., & Velez, Z. (2022). Ocean acidification affects the expression of neuroplasticity and neuromodulation markers in seabream. *Biology Open*, 11(3). https://doi.org/10.1242/bio.059073

Crespel, A., Anttila, K., Lelièvre, P., Quazuguel, P., Le Bayon, N., Zambonino-Infante, J. L., Chabot, D., & Claireaux, G. (2019). Long-term effects of ocean acidification upon energetics and oxygen transport in the European sea bass (*Dicentrarchus labrax*, Linnaeus). *Marine Biology*, 166(9), 116. https://doi.org/10.1007/s00227-019-3562-9

De Souza, K. B., Jutfelt, F., Kling, P., Förlin, L., & Sturve, J. (2014). Effects of Increased CO_2 on Fish Gill and Plasma Proteome. *PLOS ONE*, 9(7), e102901. https://doi.org/10.1371/JOURNAL.PONE.0102901

Deigweiher, K., Hirse, T., Bock, C., Lucassen, M., & Pörtner, H. O. (2010). Hypercapnia induced shifts in gill energy budgets of Antarctic notothenioids. *Journal of Comparative Physiology B*, 180(3), 347–359. https://doi.org/10.1007/s00360-009-0413-x

Deigweiher, K., Koschnick, N., Pörtner, H. O., & Lucassen, M. (2008). Acclimation of ion regulatory capacities in gills of marine fish under environmental hypercapnia. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 295(5),

R1660–R1670. https://doi.org/10.1152/ajpregu.90403.2008

Donelson, J. M., Munday, P. L., McCormick, M. I., & Pitcher, C. R. (2012). Rapid transgenerational acclimation of a tropical reef fish to climate change. *Nature Climate Change*, 2(1), 30–32. https://doi.org/10.1038/nclimate1323

Ebbesson, L. O. E., & Braithwaite, V. A. (2012). Environmental effects on fish neural plasticity and cognition. *Journal of Fish Biology*, 81(7), 2151–2174. https://doi.org/10.1111/j.1095-8649.2012.03486.x

Esbaugh, A. J. (2017). Physiological implications of ocean acidification for marine fish: emerging patterns and new insights. *Journal of Comparative Physiology B*, 188(1), 1–13. https://doi.org/10.1007/S00360-017-1105-6

Frommel, A. Y., Hermann, B. T., Michael, K., Lucassen, M., Clemmesen, C., Hanel, R., & Reusch, T. B. H. (2020). Differential gene expression patterns related to lipid metabolism in response to ocean acidification in larvae and juveniles of Atlantic cod. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 247, 110740. https://doi.org/10.1016/j.cbpa.2020.110740

Gansemer, E. R., McCommis, K. S., Martino, M., King-McAlpin, A. Q., Potthoff, M. J., Finck, B. N., Taylor, E. B., & Rutkowski, D. T. (2020). NADPH and Glutathione Redox Link TCA Cycle Activity to Endoplasmic Reticulum Homeostasis. *IScience*, 23(5), 101116. https://doi.org/10.1016/J.ISCI.2020.101116

Gordon-Weeks, P. R., & Fournier, A. E. (2014). Neuronal cytoskeleton in synaptic plasticity and regeneration. *Journal of Neurochemistry*, 129(2), 206–212. https://doi.org/10.1111/JNC.12502

Gräns, A., Jutfelt, F., Sandblom, E., Jönsson, E., Wiklander, K., Seth, H., Olsson, C., Dupont, S., Ortega-Martinez, O., Einarsdottir, I., Björnsson, B. T., Sundell, K., & Axelsson, M. (2014). Aerobic scope fails to explain the detrimental effects on growth resulting from warming and elevated CO_2 in Atlantic halibut. *Journal of Experimental Biology*, 217(5), 711–717. https://doi.org/10.1242/jeb.096743

Grosell M., Munday P. L., Farrell A. P., Brauner C. J. (2019). Fish Physiology Volume 37: Carbon Dioxide (Amsterdam: Elsevier).

Harper, C., & Wolf, J. C. (2009). Morphologic Effects of the Stress Response in Fish. *ILAR Journal*, 50(4), 387–396. https://doi.org/10.1093/ILAR.50.4.387

Heuer, R. M., & Grosell, M. (2014). Physiological impacts of elevated carbon dioxide and ocean acidification on fish. *American Journal of Physiology - Regulatory Integrative and Comparative Physiology*, 307(9), R1061–R1084. https://doi.org/10.1152/ajpregu.00064.2014

Hu, X., Bai, J., Liu, R., & Lv, A. (2023). Comprehensive transcriptomics and proteomics analysis of *Carassius auratus* gills in response to *Aeromonas hydrophila*. *Fish and Shellfish Immunology Reports*, 4, 100077. https://doi.org/10.1016/J.FSIREP.2022.100077

Huang, R., Ding, J., Gao, K., De Carvalho, M. H. C., Tirichine, L., Bowler, C., & Lin, X. (2019). A potential role for epigenetic processes in the acclimation response to elevated pCO_2 in the model diatom phaeodactylum tricornutum. *Frontiers in Microbiology*, 9, 3342. https://doi.org/10.3389/fmicb.2018.03342

Kang, J., Nagelkerken, I., Rummer, J. L., Rodolfo-Metalpa, R., Munday, P. L., Ravasi, T., & Schunter, C. (2022). Rapid evolution fuels transcriptional plasticity to ocean acidification. *Global Change Biology*, 28(9), 3007-3022. https://doi.org/10.1111/GCB.16119

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* 37(8), 907-915. https://doi.org/10.1038/s41587-019-0201-4

Lai, F., Fagernes, C. E., Bernier, N. J., Miller, G. M., Munday, P. L., Jutfelt, F., & Nilsson, G. E. (2017). Responses of neurogenesis and neuroplasticity related genes to elevated CO_2 levels in the brain of three teleost species. *Biology Letters*, 13(8). https://doi.org/10.1098/rsbl.2017.0240

Langer, G., Nehrke, G., Probert, I., Ly, J., & Ziveri, P. (2009). Strain-specific responses of Emiliania huxleyi to changing seawater carbonate chemistry. *Biogeosciences*, 6(11), 2637-2646.

Lee, D. W., Song, J. A., Park, H.-S., & Choi, C. Y. (2021). Circadian Rhythm Disturbances Due to Exposure to Acidified Conditions and Different Photoperiods in Juvenile Olive Flounder (*Paralichthys olivaceus*). *Ocean Science Journal*, 56(2), 198–206. https://doi.org/10.1007/s12601-021-00018-y

Lehmann, R., Schunter, C., Welch, M. J., Arold, S. T., Nilsson, G. E., Tegner, J. N., Munday, P. L., & Ravasi, T. (2022). Genetic architecture of behavioural resilience to ocean acidification. *BioRxiv*, 2022.10.18.512656. https://doi.org/10.1101/2022.10.18.512656

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, 30(7), 923–930. https://doi.org/10.1093/BIOINFORMATICS/BTT656

Liu, S., Wang, X., Sun, F., Zhang, J., Feng, J., Liu, H., Rajendran, K. V., Sun, L., Zhang, Y., Jiang, Y., Peatman, E., Kaltenboeck, L., Kucuktas, H., & Liu, Z. (2013). RNA-Seq reveals expression signatures of genes involved in oxygen transport, protein synthesis, folding, and degradation in response to heat stress in catfish. *Physiological genomics*, 45(12), 462-476

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12), 1–21. https://doi.org/10.1186/s13059-014-0550-8

Machado, M., Arenas, F., Svendsen, J. C., Azeredo, R., Pfeifer, L. J., Wilson, J. M., & Costas, B. (2020). Effects of Water Acidification on Senegalese Sole Solea senegalensis Health Status and Metabolic Rate: Implications for Immune Responses and Energy Use. *Frontiers in Physiology*, 11, 26. https://doi.org/10.3389/FPHYS.2020.00026/BIBTEX

Miller, G. M., Watson, S. A., Donelson, J. M., McCormick, M. I., & Munday, P. L. (2012). Parental environment mediates impacts of increased carbon dioxide on a coral reef fish. *Nature Climate Change*, 2(12), 858-861. https://doi.org/10.1038/NCLIMATE1599

Monroe, A. A., Schunter, C., Welch, M. J., Munday, P. L., & Ravasi, T. (2021). Molecular basis of parental contributions to the behavioural tolerance of elevated pCO_2 in a coral reef fish. *Proceedings of the Royal Society B: Biological Sciences*, 288(1964). https://doi.org/10.1098/rspb.2021.1931

Munday, P. L. (2014). Transgenerational acclimation of fishes to climate change and ocean acidification. *F1000Prime Reports*, 6. https://doi.org/10.12703/P6-99

Munday, P. L., McCormick, M. I., Meekan, M., Dixson, D. L., Watson, S. A., Chivers, D. P., & Ferrari, M. C. O. (2013). Selective mortality associated with variation in CO_2 tolerance in a marine fish. *Ocean Acidification*, 1, 1–5. https://doi.org/10.2478/OAC-2012-0001

Nagelkerken, I., Allan, B. J., Booth, D. J., Donelson, J. M., Edgar, G. J., Ravasi, T., Rummer, J. L., Vergés, A., & Mellin, C. (2023). The effects of climate change on the ecology of fishes. *PLOS Climate*, 2(8), e0000258.

OmicsBox – Bioinformatics Made Easy, BioBam Bioinformatics, March 3, 2019, https://www.biobam.com/omicsbox

Parker, L. M., Ross, P. M., & O'Connor, W. A. (2011). Populations of the Sydney rock oyster, *Saccostrea glomerata*, vary in response to ocean acidification. *Marine Biology*, 158, 689-697.

Petit-Marty, N., Nagelkerken, I., Connell, S. D., & Schunter, C. (2021). Natural CO_2 seeps reveal adaptive potential to ocean acidification in fish. *Evolutionary Applications*, 14(7), 1794–1806. https://doi.org/10.1111/eva.13239

Pimentel, M., Pegado, M., Repolho, T., & Rosa, R. (2014). Impact of ocean acidification in the metabolism and swimming behavior of the dolphinfish (*Coryphaena hippurus*) early larvae. *Marine Biology*, 161(3), 725–729. https://doi.org/10.1007/s00227-013-2365-7

Pistevos, J. C., Calosi, P., Widdicombe, S., & Bishop, J. D. (2011). Will variation among genetic individuals influence species responses to global climate change?. *Oikos*, 120(5), 675-689

Pörtner, H. O., Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., & Weyer, N. M. (2019). The ocean and cryosphere in a changing climate. *IPCC special report on the ocean and cryosphere in a changing climate*, 1155.

Qin, Z., Lewis, J. E., & Perry, S. F. (2010). Zebrafish (*Danio rerio*) gill neuroepithelial cells are sensitive chemoreceptors for environmental CO_2. *The Journal of Physiology*, 588(5), 861-872. https://doi.org/10.1113/JPHYSIOL.2009.184739

R Core Team. (2021). *R:* A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/.

Robertson, D. R. (1973). Field Observations on the Reproductive Behaviour of a Pomacentrid Fish, *Acanthochromis polyacanthus*. *Zeitschrift Für Tierpsychologie*, 32(3), 319–324. https://doi.org/10.1111/j.1439-0310.1973.tb01108.x

Rokitta, S. D., John, U., & Rost, B. (2012). Ocean Acidification Affects Redox-Balance and Ion-Homeostasis in the Life-Cycle Stages of *Emiliania huxleyi*. *PLOS ONE*, 7(12), e52212. https://doi.org/10.1371/JOURNAL.PONE.0052212

Rummer, J. L., Stecyk, J. A. W., Couturier, C. S., Watson, S. A., Nilsson, G. E., & Munday, P. L. (2013). Elevated CO_2 enhances aerobic scope of a coral reef fish. *Conservation Physiology*, 1(1), cot023. https://doi.org/10.1093/conphys/cot023

Ryu, T., Veilleux, H. D., Donelson, J. M., Munday, P. L., & Ravasi, T. (2018). The epigenetic landscape of transgenerational acclimation to ocean warming. *Nature Climate*

Change, 8(6), 504–509. https://doi.org/10.1038/s41558-018-0159-0

Schade, F. M., Clemmesen, C., Wegner, M., & Wegener, A. (2014). Within-and transgenerational effects of ocean acidification on life history of marine three-spined stickleback (*Gasterosteus aculeatus*). *Marine Biology*, 161(7), 1667–1676. https://doi.org/10.1007/s00227-014-2450-6

Schunter, C., Ravasi, T., Munday, P. L., & Nilsson, G. E. (2019). Neural effects of elevated CO_2 in fish may be amplified by a vicious cycle. *Conservation Physiology*, 7(1). https://doi.org/10.1093/CONPHYS/COZ100

Schunter, C., Welch, M. J., Nilsson, G. E., Rummer, J. L., Munday, P. L., & Ravasi, T. (2018). An interplay between plasticity and parental phenotype determines impacts of ocean acidification on a reef fish. *Nature Ecology & Evolution*, 2(2), 334–342. https://doi.org/10.1038/s41559-017-0428-8

Schunter, C., Welch, M. J., Ryu, T., Zhang, H., Berumen, M. L., Nilsson, G. E., Munday, P. L., & Ravasi, T. (2016). Molecular signatures of transgenerational response to ocean acidification in a species of reef fish. *Nature Climate Change*, 6(11), 1014–1018. https://doi.org/10.1038/nclimate3087

Stiasny, M. H., Mittermayer, F. H., Göttler, G., Bridges, C. R., Falk-Petersen, I. B., Puvanendran, V., Mortensen, A., Reusch, T. B. H., & Clemmesen, C. (2018). Effects of parental acclimation and energy limitation in response to high CO_2 exposure in Atlantic cod. *Scientific Reports*, 8(1), 1–8. https://doi.org/10.1038/s41598-018-26711-y

Strader, M. E., Wong, J. M., & Hofmann, G. E. (2020). Ocean acidification promotes broad transcriptomic responses in marine metazoans: A literature survey. *Frontiers in Zoology*, 17(1), 1–23. https://doi.org/10.1186/s12983-020-0350-9

Sunday, J. M., Crim, R. N., Harley, C. D., & Hart, M. W. (2011). Quantifying rates of evolutionary adaptation in response to ocean acidification. *PloS one*, 6(8), e22881

Suresh, S., Mirasole, A., Ravasi, T., Vizzini, S., & Schunter, C. (2023). Brain transcriptome of gobies inhabiting natural CO_2 seeps reveal acclimation strategies to long-term acidification. *Evolutionary Applications*, 16(7), 1345-1358

Swift, J., & Coruzzi, G. M. (2017). A matter of time — How transient transcription factor interactions create dynamic gene regulatory networks. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms*, 1860(1), 75–83. https://doi.org/10.1016/j.bbagrm.2016.08.007

Turner, B. M. (2009). Epigenetic responses to environmental change and their evolutionary implications. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1534), 3403–3418. https://doi.org/10.1098/rstb.2009.0125

Veilleux, H. D., Ryu, T., Donelson, J. M., van Herwerden, L., Seridi, L., Ghosheh, Y., Berumen, M. L., Leggat, W., Ravasi, T., & Munday, P. L. (2015). Molecular processes of transgenerational acclimation to a warming ocean. *Nature Climate Change*, 5(12), 1074–1078. https://doi.org/10.1038/nclimate2724

Walsh, P. J., & Milligan, C. L. (1993). Roles of Buffering Capacity and Pentose Phosphate Pathway Activity in the Gas Gland of the Gulf Toadfish Opsanus Beta. *Journal of Experimental Biology*, 176(1), 311–316. https://doi.org/10.1242/JEB.176.1.311

Welch, M. J., & Munday, P. L. (2017). Heritability of behavioural tolerance to high CO_2 in a
coral reef fish is masked by nonadaptive phenotypic plasticity. *Evolutionary*
Applications, 10(7), 682–693. https://doi.org/10.1111/eva.12483

Williams, C. R., Dittman, A. H., McElhany, P., Busch, D. S., Maher, M. T., Bammler, T. K.,
MacDonald, J. W., & Gallagher, E. P. (2019). Elevated CO_2 impairs olfactory-mediated
neural and behavioral responses and gene expression in ocean-phase coho salmon
(*Oncorhynchus kisutch*). *Global Change Biology*, 25(3), 963–977.
https://doi.org/10.1111/gcb.14532

Wood, D. E., & Salzberg, S. L. (2014). Kraken: Ultrafast metagenomic sequence
classification using exact alignments. *Genome Biology*, 15(3).
https://doi.org/10.1186/gb-2014-15-3-r46

Zapara, T. A., Simonova, O. G., Zharkikh, A. A., & Ratushnyak, A. S. (2000). The Effects of
the Dynamic State of the Cytoskeleton on Neuronal Plasticity. *Translated from*
Rossiiskii Fiziologicheskii Zhurnal Imeni I. M. Sechenova, I/Ol, 30(3), 128–138.

742