Date: 09-November-2023

Lateral hypothalamic GABAergic neurons encode alcohol memories.

Isis Alonso-Lozares[1,2], Pelle Wilbers[1,2], Lina Asperl[1,2], Sem Teijsse[1,2], Charlotte van der Neut[1,2], Dustin Schetters[1,2], Yvar van Mourik[1,2], Allison J. McDonald[1,2], Tim Heistek[3], Huibert D. Mansvelder[3], Taco J. De Vries[1,2], Nathan J. Marchant[1,2]

1. Department of Anatomy & Neurosciences, Amsterdam Neuroscience, Amsterdam University Medical Centers, 1081 HZ, Amsterdam, The Netherlands.
2. Compulsivity Impulsivity and Attention, Amsterdam Neuroscience, Amsterdam, The Netherlands.
3. Department of Integrative Neurophysiology, Center for Neurogenomics and Cognitive Research, Amsterdam Neuroscience, Vrije Universiteit, Amsterdam, Netherlands

Corresponding Author: Nathan Marchant (n.marchant@amsterdamumc.nl)

<u>Number of figures:</u> 7 (+5 Suppl.)

<u>Word counts</u>

      Abstract: 246 words

      Main text: 8305 words

Abstract

In alcohol use disorder the alcohol memories persist during abstinence, and exposure to stimuli associated with alcohol use can lead to relapse. This highlights the importance of investigating the neural substrates underlying not only relapse, but also encoding and expression of alcohol memories. GABAergic neurons in the Lateral Hypothalamus (LH-GABA) have been shown to be critical for food-cue memories and motivation, however the extent to which this role extends to alcohol-cue memories and motivations remains unexplored. In this study, we aimed to describe how alcohol related memories are encoded and expressed in LH GABAergic neurons. Our first step was to monitor LH-GABA calcium transients during acquisition, extinction, and reinstatement of an alcohol-cue memory using fiber photometry. We trained the rats on a Pavlovian conditioning task where one conditioned stimulus (CS+) predicted alcohol (20% EtOH) and another conditioned stimulus (CS-) had no outcome. We then extinguished this association through non-reinforced presentations of the CS+ and CS-, finally in two different groups we measured relapse under non-primed and alcohol primed induced reinstatement. Our results show that initially both cues caused increased LH-GABA activity, and after learning only the alcohol-cue increased LH-GABA activity. After extinction this activity decreases, and we found no differences in LH-GABA activity during reinstatement in either group. Next, we used inhibited LH-GABA neurons with optogenetics to show that activity of these neurons is necessary for the formation of an alcohol-cue associations. These findings suggest that LH-GABA might be involved in attentional processes modulated by learning.

Introduction

Alcohol is one of the most widely available and frequently consumed substances in our society (Eurostat, 2022). While most individuals (80-90%) can consume alcohol in a moderate and controlled manner, a small percentage develop alcohol use disorder (AUD). In the European Union alone, this disorder affects approximately 23 million people (Carvalho et al., 2019). AUD is characterized by persistent alcohol use despite negative consequences and a high likelihood of relapse (Hunt et al., 1971). Individuals with AUD often go through multiple cycles of treatment, periods of abstinence, and relapse before achieving full recovery (Fleury et al., 2016).

Relapse can occur when an individual encounters cues and environments associated with previous alcohol use (O'Brien et al., 1992; Sliedrecht et al., 2019), and these experiences can cause craving (Suzuki et al., 2020) which leads to relapse (Vafaie & Kober, 2022). These strong positive associations between neutral stimuli (e.g., a bar, a specific location, certain individuals) and alcohol are formed in the initial stages of addiction development and share many similarities with behavioural patterns triggered by natural rewards like food (Everitt & Robbins, 2016). Like natural rewards, drugs create anticipation of positive outcomes and reinforce the individual's motivation to seek them. Consequently, pathological associative learning is a critical component of the addiction cycle, where the brain's mechanisms that typically promote survival-related behaviours are hijacked by a drug's more subjectively potent stimuli (Hyman, 2005). It is therefore crucial to better understand how these memories are formed in the brain.

The Lateral Hypothalamus (LH) has been considered a critical substrate of appetitive and consummatory behaviours since it was discovered that lesions cause hypophagia, while stimulation has the opposite effect (Hetherington & Ranson, 1940; Delgado & Anand, 1953). Others pointed at the LH having a general role in motivation (Valenstein et al., 1968a, 1968b, 1970), and more recent data suggests the role of LH extends beyond promoting feeding behaviour to orchestrating complex learning processes related to the encoding and maintenance of motivationally relevant cue-food associations (Cole et al., 2015; Jennings et al., 2015; Sharpe et al., 2017), as well as other learning processes (Kosse & Burdakov, 2019; Burdakov & Peleg-Raibstein, 2020). In relation to relapse, increased activity in LH is associated with context-induced reinstatement of alcohol seeking after both extinction and punishment (Marchant et al., 2009; 2014), as well as context-induced reinstatement of extinguished cocaine seeking (Hamlin et al., 2008).

Recent evidence suggests that the GABAergic subpopulation of LH (LH-GABA) are involved in associative learning processes and motivated behaviours. These neurons are functionally and anatomically distinct from the Orexin (Ox) and Melanin-Concentrating

Hormone (MCH) populations (Jennings et al., 2015; Karnani et al., 2013), and have reciprocal projections with Ventral Tegmental Area (VTA; Nieh et al., 2016) and Nucleus Accumbens Shell (NAcS; O'Connor et al., 2015), areas well known for their roles in reward learning. LH-GABA activity is associated with appetitive or consummatory behaviours (Jennings et al., 2015), and opto-inhibition of LH-GABA subpopulation prevents learning of cue-food associations (Sharpe, 2017).

Currently the role of LH-GABA in alcohol related learning is unknown. Here, we trained rats on an alcohol Pavlovian conditioning task and used fiber photometry calcium imaging (Nakai et al., 2001; Cui et al., 2014) to show that LH-GABA activity is initially high to cues when they are delivered, and across learning this activity is sustained to the alcohol-predictive cue but decreases to the non-predictive cue. We also show in extinction that activity in LH-GABA neurons track violations in expected outcomes, with sustained activity when alcohol is expected but not delivered. To investigate a causal role of LH-GABA neurons we used optogenetics (Mahn et al., 2018) to inhibit activity during conditioning and found that acquisition is decreased during cue presentation. These data highlight the importance of LH-GABA in both encoding and expression of alcohol associated memories.

Results

Exp. 1: Validation of the dual-virus approach to target GABAergic neurons in LH

To target the GABAergic subpopulation of LH neurons we used dual-virus approach involving combination of a GAD-Cre virus with different Cre-dependent viruses (Wakabayashi et al., 2019). To first validate this approach, we injected a cocktail of one AAV encoding Cre driven by the GAD promoter with another AAV encoding Cre-dependent eGFP. Immunofluorescence labelling MCH and OX-expressing neurons revealed minimal overlap between both OX-immunoreactive (Fig. 1B) and MCH-immunoreactive (Fig. 1C) neurons and GFP expression driven under the GAD promoter. Using RNA-scope to label GAD mRNA (Fig. 1A), we found high overlap between GAD+ and eGFP (Fig. 1D). These data show that the dual-virus approach was successful in confining expression of AAV-induced transgenes to GABAergic neurons.

To validate that blue light effectively inhibits the inhibitory opsin GtACR2 (Mahn et al., 2018) expressing neurons, we made whole cell recordings from fluorescent labelled neurons (Fig. 1D-E). Current injection into GtACR2 expression cells depolarized them above spike threshold, and activating GtACR2 with blue light significantly decreased the rate of current-induced action potentials (n=6).

Figure 1. *Validation of the dual-virus approach to target LH-GABA and optogenetic inhibition of LH-GABA neurons expressing GtACR2 using light.* Representative images of eGFP expression and co-labelling of OX (**A**), MCH (**B**) and GAD (**C**). Data are mean (+/- SEM; n = 4 for MCH and OX validation, n = 5 for GAD validation) % of eGFP+ only or eGFP+OX (**A**), eGFP+MCH (**B**), eGFP+GAD (**C**). (**D**) Example of light-evoked suppression of neuronal firing inGtACR2 expressing neurons. (**E**) Summary data plotting mean (+/- SEM; n = 6) spike frequency in the presence and absence of light stimulation. Blue bars indicate light delivery.

Exp. 2: Monitoring LH-GABA activity during alcohol conditioning, extinction, and primed reinstatement of alcohol-seeking

In experiment 2, we aimed to identify LH-GABA activity during alcohol conditioning, extinction, and primed reinstatement of alcohol seeking. We used fiber photometry to measure real-time population level LH-GABA neuron calcium (Ca2+) transients throughout the entire task. Figure 2A shows the experimental timeline, the expression of the Ca2+ sensor jGCaMP7f (Dana et al., 2019) in LH-GABA was achieved using a mixture of one AAV encoding Cre driven by the GAD promoter with another AAV encoding Cre-dependent jGCaMP7f under the hSyn1 promoter. Throughout the entire task we measured fluorescence via an optic fiber cannula implanted above LH (Fig. 2B).

Figure 2. *Monitoring LH-GABA activity using fiber photometry calcium imaging: Experimental design and histological validation.* (**A**) Outline of the experimental procedure (n =18; 2 males, 16 females). (**B**) Trial design for photometry experiments. (**C**) Representative image of jGCaMP7f expression and fiber implant in LH, and fiber placement and expression validation.

Learning across conditioning sessions

Behavioural data: We used repeated measures ANOVA with the within-subjects factors Session (First, Last) and CS Type (CS+, CS-) to compare alcohol seeking during the first and the last conditioning sessions (Fig. 3A and 3B). We found main effects of Session and CS Type, as well as a significant Session x CS Type interaction on time spent in the magazine both during (Fig. 3B; $F_{(1, 17)} = 133,7$, $p < 0.0001$) and after (Fig. 3C; $F_{(1, 17)} = 4747$, $p < 0.0001$) CS presentation. We found no difference between CS Type on time in the magazine during the pre-CS time period (Supp. Fig. 2). These data show that, throughout the course of the conditioning phase, the rats learned to discriminate between the alcohol-predictive and neutral cues, and alcohol seeking increased during the CS+ but not the CS-.

Photometry data: We made comparisons within a single session between the CS types (CS+ vs CS-), and across the different sessions within the same CS type (First vs. Last; Fig 3. D-G). In the first conditioning session (Fig. 3D), we found that LH-GABA activity was significantly higher for both CS types. Bootstrapping analysis revealed a significant increase from baseline for CS+ (0.31s to 11.76s) and CS- (0.31s to 10.94s), with no significant difference in activity between the two CS types (permutation tests). In the last conditioning session (Fig. 3E), bootstrapping analysis revealed a significant increase from baseline for CS+ (0.31s to 9.62s) and CS- (0.44s to 10.6s), however in contrast to the first session we found a significant difference in LH-GABA activity between CS+ and CS- (0.44 to 6.9s), with LH-GABA activity to the CS+ being significantly higher than to the CS-. These data suggest that acquisition of a cue-alcohol memory is associated with increased activity of LH-GABA neurons. To further investigate this, we performed additional analyses comparing CS+ or CS- activity between the first and last conditioning sessions. These analyses show that learning was associated with changes in activity to both the CS+ and CS-. We found that acquisition of the cue-alcohol memory was associated with both longer sustained activity to the CS+ (Fig. 3F; Permutation test: 2.14 to 7s), and decreased activity to the CS- (Fig. 3G; Permutation test: 0.5 to 1.57s). Overall, these data show that the activity of LH-GABA neurons is bidirectionally shaped by learning experience. We found both an increase in activity to a motivationally relevant (alcohol-predictive) cue, as well as decreased activity to a motivationally non-relevant cue (CS-).

Figure 3. *Monitoring LH-GABA during alcohol Pavlovian conditioning.* (**A**) Outline of experimental procedure (n = 2 males, n = 16 females). Mean ± standard error of the mean (SEM) time spent in the alcohol magazine during (**B**) and after (**C**) CS presentation comparing the first and last conditioning sessions. Ca2+ traces of LH-GABA activity centered around cue onset (-5s to +20s) comparing LH-GABA activity to CS+ and CS- on the first (**D**) and last (**E**) conditioning sessions. (CS+ Cond1: n = 169; CS- Cond1: n = 168; CS+ Cond19: n =170; CS- Cond19: n = 169). The same mean Ca2+ traces are plotted for comparisons across sessions between CS+ (**F**) and CS-(**G**). For all photometry traces, bars at bottom of graph indicate significant deviations from baseline (dF/F ≠ 0) determined via bootstrapped confidence intervals (99% CIs), or significant differences between the specific events (CS+ Cond1; CS- Cond1; CS+ Cond19; CS- Cond19) determined via permutation tests with alpha 0.01 for comparisons between CS type and session. Vertical dashed lines indicate CS onset and alcohol delivery for CS+, horizontal line indicates baseline (dF/F = 0).

Expression of the learned alcohol memory

Behavioural data: Repeated measures ANOVA comparing the first and the last conditioning sessions (Fig. 4B and 4C) revealed a significant Session x CS Type interaction on time spent in the magazine both during (Fig. 4B; $F_{(1, 17)} = 166,0$, $p < 0.0001$) and after (Fig. 4C; $F_{(1, 17)} = 1772$, $p < 0.0001$) CS presentation. Thus, we found that rats spent significantly more time in the alcohol port during CS+ than CS- in the absence of alcohol delivery, suggesting the learned memory was expressed without the alcohol outcome.

Photometry data: We made comparisons within a single session between the different CS types (CS+ vs CS-), and across the different sessions within the same CS type (First vs. Last; Fig 4. D-G). Here we plotted the same data from Fig. 3D into Fig. 4D, and results for the last conditioning session are described above. On the first extinction session (Fig. 4D), bootstrapping analysis revealed a significant increase from baseline for both CS+ (0.25s to 20s), and CS- (0.44s to 8.6s, 10s to 10.12s, and 10.37s to 10.81s). Comparison of activity between CS types revealed that activity to CS+ remains significantly higher compared to CS- both during cue presentation, and 10 seconds after (Fig. 4D; 0.31 to 20s). To determine if LH-GABA also track extinction of a previously learned alcohol-predictive cue, we next compared LH-GABA activity to either CS+ (Fig 4.F) or CS- (Fig. 4G) between the last conditioning and first extinction sessions. Interestingly, while we found no difference in LH-GABA activity to CS+ or CS- when the cues were delivered, we did observe a higher sustained activity in the first extinction session compared to the last conditioning session (9.4 to 12.9s; 16.8 to 18.42s). This sustained activity was during the period when alcohol is expected but not delivered. This observation suggests LH-GABA neuron activity is related to reward-prediction error, by increasing their activity when expectations of a reward are violated.

A Experimental phase

Alcohol seeking

B During CS presentation

C After CS presentation

LH-GABA activity

D Last conditioning session

E First extinction session

F Last conditioning vs first extinction CS+

G Last conditioning vs first extinction CS-

Figure 4. *Monitoring LH-GABA during alcohol Pavlovian conditioning and extinction.* (**A**) Outline of experimental procedure (n = 2 males, n = 16 females). Mean ± standard error of the mean (SEM) time spent in the alcohol magazine during (**B**) and after (**C**) CS presentation comparing the last conditioning and first extinction sessions. Ca2+ traces of LH-GABA activity centred around cue onset (-5s to +20s) comparing LH-GABA activity to CS+ and CS- on the last conditioning (**D**) and first extinction (**E**) sessions. (CS+ Cond19: n = 170; CS- Cond19: n = 169; CS+ EXT1: n = 61; CS- EXT1: n = 136). The same mean Ca2+ traces are plotted for comparisons across sessions between CS+ (**F**) and CS-(**G**). For all photometry traces, bars at bottom of graph indicate significant deviations from baseline (dF/F ≠ 0), or significant differences between the specific events (CS+ Cond19; CS- Cond19; CS+ EXT1; CS- EXT1), determined via bootstrapped confidence intervals (99% CIs), and permutation tests with alpha 0.01 for comparisons between CS type and session. Vertical dashed lines indicate CS onset and alcohol delivery for CS+, horizontal line indicates baseline (dF/F = 0).

Extinction of a learned alcohol memory

Behavioural data: Repeated measures ANOVA comparing the first and the last extinction sessions (Fig. 5B and 5C) revealed a significant Session x CS Type interaction on time spent in the magazine both during (Fig. 5B; $F_{(1, 17)} = 19,96$, $p < 0.001$) and after (Fig. 5C; $F_{(1, 17)} = 34,75$, $p < 0.0001$) CS presentation. Thus, we show that rats spent significantly more time in the alcohol port during and after CS+ onset on the first extinction session compared to CS-. This contrasts with the last extinction session, where we show that time spent in the alcohol port did not significantly differ between CS+ and CS-.

Photometry data: We made comparisons within a single session between the different CS types (CS+ vs CS-), and across the different sessions within the same CS type (First vs. Last; Fig 5. D-G). Here we have plotted the same data from Fig. 4D into Fig. 5D. Results for the first extinction session are described above. On the last extinction session, we found that LH-GABA has excitatory Ca^{2+} transients after CS onset (Fig. 5E). Bootstrapping analysis revealed a significant increase from baseline for CS+ (0.25s to 20s), and CS- (0.44s to 2.57s and 8.5s to 8.9s). The permutation tests comparing the activity between CS types for the last extinction session show that activity to the CS+ remained higher than CS- during the last extinction session (Fig. 5E; 0.44 to 1.64 s; 2.58 to 5.03 s).

To further investigate the different patterns of activity of LH-GABA during the first and last extinction sessions, we performed permutation tests of LH-GABA activity to CS+ (Fig. 5F) and CS- (Fig. 5G), across sessions. Permutation tests revealed significant differences between the first and last extinction sessions for CS+ (Fig. 5F; 1.13 to 5.47 s; 7.23 to 11.5 s), but not CS- (Fig. 5G). Overall, these data show that extinction of a cue-alcohol memory is associated with both a decrease in alcohol seeking, and decreased cue-evoked activity of LH-GABA neurons.

Figure 5. *Monitoring LH-GABA during extinction.* (**A**) Outline of experimental procedure (n = 2 males, n = 16 females). Mean ± standard error of the mean (SEM) time spent in the alcohol magazine during (**B**) and after (**C**) CS presentation comparing the first and last extinction sessions. Ca2+ traces of LH-GABA activity centred around cue onset (-5s to +20s) comparing LH-GABA activity to CS+ and CS- on the first (**D**) and last (**E**) extinction sessions. (CS+ EXT1: n = 161; CS- EXT1: n = 136; CS+ EXT7: n = 161; CS- EXT7: n = 136). The same mean Ca2+ traces are plotted for comparisons across sessions between CS+ (**F**) and CS- (**G**). For all photometry traces, bars at bottom of graph indicate significant deviations from baseline (dF/F ≠ 0), or significant differences between the specific events (CS+ EXT1; CS- EXT1; CS+ EXT7; CS- EXT7), determined via bootstrapped confidence intervals (99% CIs), and permutation tests with alpha 0.01 for comparisons between CS type and session. Vertical dashed lines indicate CS onset and alcohol delivery for CS+, horizontal line indicates baseline (dF/F = 0).

Alcohol-primed induced reinstatement of extinguished alcohol seeking.

We separated the animals into two groups: one group received primed 24 hour before the reinstatement with no priming on the reinstatement test (Supp. Fig. 4), and the other group received alcohol primed both 24 hours before and just prior to the reinstatement test (Chaudhri et al., 2010; LeCocq et al., 2018; Fig. 6). Only the second group showed reliable reinstatement to the CS+, and therefore we present these data here. Results for the first group are reported in the supplementary results (Supp. Fig. 4).

Behavioural data: Repeated measures ANOVA comparing the last extinction session and reinstatement (Fig. 6E and 6F) revealed a significant Session x CS Type interaction on time spent in the magazine during (Fig. 6E; $F_{(1, 12)} = 17,86$, $p < 0.01$) but not after (Fig. 6F; $F_{(1, 12)} = 4,057$; $p = 0.07$) CS presentation. Specifically, we found that rats spent significantly more time in the alcohol port during CS+ on the reinstatement test, compared to during CS- presentation and during CS+ on the last extinction session (multiple comparisons, bonferroni corrected). These data show that the alcohol-prime procedure induced reinstatement of extinguished alcohol seeking, which was specific to the previously conditioned CS+.

Photometry data: We made comparisons within a single session between the different CS types (CS+ vs CS-), and across the different sessions within the same CS type (First vs. Last; Fig 6. D-G). Here we have plotted the same data from Fig. 5D into Fig. 6D. Results for the last extinction session are described above. On the reinstatement session, we found that LH-GABA has excitatory Ca2+ transients after CS onset (Fig. 6E). Bootstrapping analysis revealed a significant increase from baseline for CS+ but not CS- on the last extinction session (0.38s to 9.24s), and a significant increase from baseline for both CS+ (0.38s to 9.87s) and CS- (0.57s to 6.10s; 7.48s to 9.87s) on the reinstatement test. Permutation tests revealed a significant difference in activity between CS+ and CS- for the last extinction (2.64 to 4.91 s) and reinstatement (5.59 to 7.67) sessions. Despite the significant increase in alcohol seeking to the CS+ in reinstatement, permutation tests revealed no significant differences in LH-GABA activity to CS+ between the last extinction session and reinstatement. The only significant differences we found were between CS- across sessions (2.58 to 3.33 s).

Figure 6. *Monitoring LH-GABA during alcohol-primed induced reinstatement of alcohol seeking.* (**A**) Outline of experimental procedure (n = 13 females). Mean ± standard error of the mean (SEM) time spent in the alcohol magazine during (**B**) and after (**C**) CS presentation comparing the last extinction and reinstatement test sessions. Ca2+ traces of LH-GABA activity centred around cue onset (-5s to +20s) comparing LH-GABA activity to CS+ and CS- on the last extinction (**D**) and reinstatement test (**E**) sessions. (CS+ EXT7: n = 105; CS- EXT7: n = 80; CS+ RT: n = 92; CS- RT: n = 73). The same mean Ca2+ traces are plotted for comparisons across sessions between CS+ (**F**) and CS- (**G**). For all photometry traces, bars at bottom of graph indicate significant deviations from baseline (dF/F ≠ 0), or significant differences between the specific events (CS+ EXT7; CS- EXT7; CS+ RT; CS- RT), determined via bootstrapped confidence intervals (99% CIs), and permutation tests with alpha 0.01 for comparisons between CS type and session. Vertical dashed lines indicate CS onset and alcohol delivery for CS+, horizontal line indicates baseline (dF/F = 0).

Exp 3. Optogenetic inhibition of LH-GABA impairs alcohol memory learning.

Here we aimed to determine whether activity in LH-GABA is necessary for acquisition of the cue-alcohol memory. Figures 7A and 7B show the experimental timeline. We used the dual-virus approach to confine expression of the inhibitory opsin GtACR2 (Mahn et al., 2018; Fig. 1D-E) or control GFP/mCherry to LH-GABA neurons. We then trained the animals on the conditioning phase while delivering light to LH during presentation of both cues. We then tested the expression of the alcohol memory on a single extinction session, where no light inhibition or alcohol was delivered.

Figure 7D shows alcohol seeking during (top; with light inactivation) and after (bottom) CS presentation throughout conditioning sessions. During CS presentation, we observed significant Session x CS Type ($F_{(3,297, 52,75)} = 6,234$, $p < 0.001$) and Session x Group ($F_{(10, 160)} = 3,569$, $p > 0.001$) interactions, but no Session x Group x CS Type interaction. Further analyses showed that the Session x Group interaction was only present in alcohol seeking to the CS+ ($F_{(10, 160)} = 2,078$, $p < 0.05$), but not CS-.

During the 10 seconds after CS presentation, we observed significant Session x Group ($F_{(10, 160)} = 2,053$, $p < 0.95$), Session x Type of CS ($F_{(10, 160)} = 19,01$, $p < 0.001$), and Session x Group x Type of CS ($F_{(10, 160)} = 3,151$, $p < 0.01$) interactions, but no Group x CS Type interaction. Further analyses showed that the Session x Group interaction was only present in alcohol seeking to the CS+ ($F_{(10, 160)} = 2,944$, $p < 0.01$), but not CS-. Overall, these data show that optogenetic inhibition of LH-GABA during conditioning prevented acquisition of the CS-alcohol memory, and there was no observed effect on behaviour to the CS- across sessions. In addition, we measured the alcohol remaining in the magazine at the end of each session and found that all alcohol was consumed. Thus, while opto-inhibition of LH-GABA decreased magazine entries during the CS presentation, the alcohol was consumed during the non-CS periods.

Figure 7E shows alcohol seeking in the extinction test during (top) and after (bottom) CS presentation. We observed no significant effects of Group or CS Type, or an interaction between them during CS presentation. After CS presentation, we did observe a significant effect of Group ($F_{(1, 16)} = 7,637$, $p < 0.05$), as well as CS Type ($F_{(1, 16)} = 8,959$, $p < 0.01$), but no Group x CS Type interaction. Post-hoc analysis revealed a significant difference in alcohol seeking between groups to the CS+ ($p < 0.001$), but not the CS-. These data show that the control animals showed expression of the learned cue-alcohol memory, while the opsin group did not.

A Experimental design

B Trial design

C Histological validation

During CS presentation

D Conditioning

E Extinction

After CS presentation

Conditioning

Extinction

Figure 7. *Inactivation of LH-GABA during conditioning.* (**A**) Outline of the experimental procedure (n =18 females). (**B**) Trial design for optogenetics experiments. (**C**) Representative image of GtACR2 expression and fiber implants in LH, and fiber placement and expression validation. (**D**) Mean ± standard error of the mean (SEM) time spent in the alcohol magazine during (top) and after (bottom) CS presentation across conditioning sessions. Blue overlay depicts when optogenetic inhibition was delivered. (**E**) Mean ± standard error of the mean (SEM) time spent in the alcohol magazine during (top) and after (bottom) CS presentation on the extinction test. *p < 0.05.

Discussion

In this study we describe the role of LH-GABA neurons in the acquisition and expression of a cue-alcohol memory. We first validated the dual-virus approach to confine Cre-dependent expression of fluorescent proteins, sensors, and opsins to LH-GABA neurons. We then used this approach with fibre photometry calcium imaging in male and female rats to show that LH-GABA neurons increase activity to cues prior to learning, and as learning progressed the cue-evoked activity adapts. We found that LH-GABA activity is maintained for the alcohol-associated cue (CS+) but is decreased to the cue with no consequence (CS-). We also found evidence for prediction-error signalling during extinction, where LH-GABA neurons showed sustained activity when alcohol was expected but not delivered. Finally, during reinstatement we found that LH-GABA neurons maintain higher activity to the CS+ compared to the CS-, but this activity was not significantly higher than the last extinction session. This suggests that LH-GABA activity may not be critical for primed-induced reinstatement. Next, we used optogenetics in female rats to show that inhibition of LH-GABA during conditioning impaired learning of cue-alcohol associations. This study demonstrates the importance of LH-GABA in memory processes related to alcohol seeking.

Methodological considerations

Several issues must be considered in the interpretation of these findings. The photometry and optogenetics experiments used slightly different conditioning protocol, where the alcohol was delivered during the cue in photometry but after the cue in optogenetics. We did this to avoid potential non-specific effects of LH-GABA inhibition on alcohol drinking. We found that alcohol delivery during the cue led to higher alcohol seeking during the cue period in experiment 2 compared to experiment 3. However, we also found that alcohol seeking during the post-cue period was not different between these experiments, providing evidence for learning in both training conditions.

In the reinstatement tests, we used two different procedures in two different cohorts of rats. In the first, the alcohol prime was given 24 hours prior to the test without alcohol primed during the reinstatement test (LeCocq et al., 2018). In the second the alcohol prime was given both 24 hours before and at the beginning of the reinstatement test (Lé et al., 1998). We found that the first group did not reliably reinstate alcohol seeking to the CS+, while the second group did. We therefore only analysed the photometry data from the second group.

The effect of sex and oestrous cycle has gained interest in recent years (Torres et al., 2014). In exp. 3 we used only female rats, and while we did not measure the cycle of these rats and cannot be sure if the phase of their cycle could have affected their behaviour, it has been suggested that there are no sex differences in operant behaviour related to drinking

(Priddy et al., 2017), and this might extend to Pavlovian behaviour. In experiment 2 we used both male and female rats and did not observe any significant differences in behaviour or LH-GABA activity between the sexes. Therefore, we argue that the effect of opto-inhibition of LH-GABA neurons on acquisition of a cue-alcohol memory likely extends to both male and female rats.

Finally, although our findings show a specific role of the GABAergic subpopulation of LH neurons in cue-alcohol reward learning, these are a heterogeneous population of neurons. LH-GABA neurons are comprised of populations which express different peptides (Mickelsen et al., 2017; Mickelsen et al., 2019), and of different output projections (Bonnavion et al., 2016). Therefore, in this study we have recorded and manipulated activity of the LH-GABA neuron sub-population which is comprised of various phenotypes and efferent targets. It will be of interest in future studies to determine which of these outputs are critical for alcohol learning. Given that LH-GABA neurons send dense projections to the VTA, it is likely that this pathway is critical for the acquisition of the cue-alcohol memory. Furthermore, we previously showed that LH-GABA projections to VTA regulate learning about food reward-paired cues (Sharpe et al., 2017) identifying this pathway as critical for learning. Given our observation of sustained activity of LH-GABA neurons when an expected alcohol reward is not delivered, we propose that this information may be conveyed to VTA to modulate the output of VTA dopamine neurons during extinction learning. It will be of interest in future studies to determine the contribution of LH-GABA input to VTA in the generation of this signal.

Cue-induced activity of LH-GABA neurons changes with acquisition of a cue-alcohol association, and signals omission of an expected alcohol reward.

Throughout all phases of the experiment, we found LH-GABA activity is increased to the presentation of a cue. This observation shows that LH-GABA activity changes in response to most changes in the environment. Interestingly there is precedent for a function such as this from historical lesion data, showing sensory neglect from LH lesions (Marshall et al., 1971; Marshall et al., 1974). However, an important finding of this study learning is associated with significant changes in the patterns of activity.

On the first conditioning session, we found LH-GABA activity increases to both cues which initially have no motivational relevance. By the last conditioning day, LH-GABA activity was significantly higher to CS+ compared to the CS-. We found that LH-GABA activity significantly decreased to the non-predictive cue and was sustained to the alcohol-predictive cue. This pattern of activity might reflect a role of LH-GABA in gating activity to motivationally significant stimuli, maintaining activity to behaviourally relevant stimuli, while decreasing

activity to irrelevant ones. This function may be consistent with observations that LH is important for attentional processes involved in learning (Wheeler et al., 2014). Consistent with this, on the first extinction session we found that activity to the alcohol-predictive cue remained significantly higher compared to the last conditioning session after the cue was turned off, when alcohol was expected. This sustained activity is consistent with a prediction error signal, which signals violation of an expected reward (Schultz, 1998; Den Ouden et al., 2012). We have previously proposed that LH-GABA projections to VTA are involved in the generation of the prediction error signal in VTA dopamine neurons (Sharpe et al., 2017). Furthermore, violations in expected outcomes also increase attentional processes (Holland & Schiffino, 2016), providing further evidence for a role in LH-GABA in attentional processes.

These processes may be orchestrated through excitatory projections from the basolateral amygdala (BLA) to the LH (Repucci & Petrovich, 2016), which are known to signal salient sensory-specific properties of stimuli based (Holland & Gallagher, 2003; Parkes & Westbrook, 2010, Hoang & Sharpe, 2021). Given the strong reciprocal projections of LH-GABA and VTA (Watabe-Uchida et al., 2012; Nieh et al., 2015, 2016), and the role this area is suggested to play in attention (Wheeler et al., 2014), a key component of associative learning (Pelley et al., 2016), and memory updating (Burdakov & Pele Raibstein, 2020), it is likely that LH-GABA play a role in increasing attention to encode new stimulus-outcome associations.

LH-GABA activity decreases after extinction of a cue-alcohol association and is not related to primed-induced reinstatement.

By the last extinction session, we found that LH-GABA activity to the CS+ is significantly decreased compared to the first extinction session. We propose that this further demonstrates a role of LH-GABA in acting as an information gating filter (Burdakov & Raibstein, 2020). In this model, LH-GABA activity is decreased to the CS+ because extinction has weakened the incentive value of this cue. Interestingly, even after alcohol seeking is fully extinguished, we still observe significant differences in LH-GABA activity between the CS+ and CS-. This suggests that LH-GABA activity to the formerly alcohol-predictive cue is not completely erased by extinction, which is consistent with the view that extinction is not memory erasure but rather creates a new memory which gates the expression of the originally learned memory (Bouton, 2014).

During primed-induced reinstatement, we found that LH-GABA activity to the CS+ was significantly higher than to the CS- for some periods of the cue presentation. However, activity to the CS+ was not increased relative to the last extinction session, suggesting that

LH-GABA activity is not specifically related to alcohol-primed induced relapse. One potential explanation for the lack of effect could be related to variation within the group, where some rats did not reinstate alcohol seeking. Another possibility is that after extinction training, memory expression is mediated through alternative brain regions. For example, in cocaine seeking studies, inactivation of the prefrontal cortex impacts reinstatement after extinction with no effect on cocaine seeking on the first day of extinction (Peters et al., 2009). Finally, inactivation of nucleus accumbens core, but not shell, decreases primed-induced reinstatement of extinguished operant alcohol seeking (Chaudhri et al., 2008). Interestingly the nucleus accumbens core does not project to LH while the shell does (Marchant et al., 2009; Marchant et al., 2014), providing anatomical justification for a potential lack of a role of LH-GABA in priming induced reinstatement of cued alcohol seeking. Finally, activity of the mu opioid receptor in ventral hippocampus is also necessary for primed-induced reinstatement to an extinguished alcohol-predictive CS (LeCocq et al., 2023). Thus, alcohol-primed induced reinstatement may depend on other brain regions than the GABAergic sub-population of LH.

This interpretation implies that the function of LH-GABA neurons is critical in the formation of cue-alcohol memories, and potentially in extinction learning, but not in the reinstatement of these memories. Given that LH has previously been implicated in reinstatement procedures (Marchant et al., 2012), it is possible that different neuronal sub-populations in LH, such as Ox, are involved in reinstatement. Finally, addictive behaviours are characterised by attentional biases for drug-related cues (Field & Cox, 2008; Anderson, 2017). Given the LH role in attentional processes this area could be involved during relapse by directing attention to the drug-related cues.

<u>Optogenetic inhibition of LH-GABA impairs learning of alcohol-predictive cues.</u>

In experiment 3 we show that optogenetic inhibition of LH-GABA neurons during cue presentation significantly impaired learning of the alcohol-predictive cue. This observation is consistent with the photometry observations during conditioning. These conclusions were strengthened during the extinction test without opto-inhibition of LH-GABA neurons, where only the control group showed alcohol seeking post CS+. This observation is comparable to our previous work in the transgenic GAD-Cre rat, showing that opto-inhibition of LH-GABA neurons impairs acquisition of a cue-food association (Sharpe et al., 2017), and thus extends the role of LH-GABA neurons in reward learning to addictive substances such as alcohol.

The LH harbours a wide range of different neuropeptide expressing neurons, such as Orexin/hypocretin (Ox/Hcrt; De Lecea et al., 1998), Melanin-Concentrating Hormone (MCH;

Bittencourt et al., 1992) Neurotensin, Galanin (Wheeler et al., 2014; Jennings et al., 2015), as well as putative γ-aminobutyric acid- (GABA) and Glutamate- releasing neurons (Bonnavion et al., 2016). These diverse neuronal sub-populations have been shown to host a variety of functions, including associated with alcohol seeking (Dayas et al., 2008; Duncan et al., 2006) and associative learning (Aitta-aho et al., 2016; Izawa et al., 2019). The contribution of LH-GABA to these functions may be consistent with the findings from these studies, indicating that the multiple different cellular phenotypes within LH contribute towards addictive behaviours (Marchant et al., 2012).

<u>Concluding remarks</u>

In this study we sought to investigate the role of LH-GABA in learning about stimuli which predict alcohol rewards. Our results show that activity in LH-GABA is necessary for the acquisition of alcohol-predictive cues. We also show that the activity of LH-GABA neurons is modulated through experience, based on changes in the motivational properties of stimuli across learning. Future studies are needed to determine the extent to which this activity is necessary for relapse, but our data indicate that other brain regions may be critical for alcohol-primed induced reinstatement of alcohol seeking. Overall, our findings highlight an important role for LH-GABA in a fundamental learning process and show that this system is a critical component of memory formation in alcohol addiction.

Materials and methods

Subjects

We used 40 adult wild-type Long-Evans rats (4 males and 36 females), from Janvier (France). All behavioural and surgical procedures were performed in the animal facility of the Vrije Universiteit, Amsterdam, The Netherlands. Rats were kept in temperature and humidity-controlled rooms, in an inverted 12/12 light/dark cycle, and we trained the rats during the dark cycle. Food and water were provided ad libitum. Animals were kept in cages of either 2 or 3 cage mates. All procedures were approved by the Instantie voor Dierenwelzijn of the Vrije Universiteit, Amsterdam, The Netherlands.

We did not make specific analysis to determine sample size prior to any experiments. Cohorts were balanced by experimental group prior to the experiment starting. We allocated the rats randomly to one of the groups within the optogenetics and chemogenetics experiments. We did not exclude any rats for reasons of behavioural validation. Rats in Experiments 2 and 3 that had expression or fiber placement not within LH, were excluded form analysis. Rats in Experiment 3 that had either unilateral expression, misplaced expression or a misplaced fiber placement were excluded from analysis.

Apparatus

All procedures were performed in standard Med Associates operant chambers with data collected through the MED-PC V program (Med Associates, Georgia, VT, USA). Each chamber had a grid floor and a large opening at the top to allow patch cords through. Each chamber was equipped with a white noise generator, and one or two light cues, as well as a magazine port where the alcohol was delivered (20% ethanol). The magazines were equipped with a beambreak device to measure total time spent in the magazine. Custom made Med-PC programs controlled all aspects of the task.

For fiber photometry, excitation and emission light was relayed to and from the animal via optical fiber patch cords (0.48 NA, 400-μm flat tip; Doric Lenses). Blue excitation light (490 or 470 nm LED [M490F2 or M470F2, Thorlabs]) was modulated at 211 Hz and passed through a 460–490 nm filter (Doric Lenses), while isosbestic light (405 nm LED [M405F1, Thorlabs]) was modulated at 531 Hz and passed through a filter cube (Doric Lenses). GCaMP7f fluorescence was passed through a 500- to 550-nm emission filter (Doric Lenses) onto the photoreceiver (Newport 2151). Light intensity at the tip of the fiber was measured before every training session and kept at 21 μW. A real-time processor (RZ5P, Tucker Davis Technologies) controlled excitation lights, demodulated fluorescence signals and received timestamps of behavioural events. Data were saved at 1017.25 Hz and analysed with custom-made Matlab scripts, available at: https://github.com/ialozares/AlcoholConditioning

For optogenetics experiments, light (470nm) was delivered to the animals via optical bifurcating fiber patch cords (0.48 NA, 200-µm flat tip; Doric Lenses) connected to the head of the animals. Light delivery began 500ms prior to CS onset until 500ms after CS presentation. Light intensity was kept at 5mW and measured at the beginning of each session for each operant chamber. Light delivery was controlled by custom Med-PC scripts.

Drugs

Deschloroclozapine (DCZ) (MedChem Express, 1977-07-7) was dissolved in 1-2% dimethyl sulfoxide (DMSO) in saline to a final volume of 0.1ml per kg. Alcohol solutions were prepared using 96% (v/v) ethanol diluted with tap water and were administered using normal bottles.

Surgeries

The day before and thirty minutes prior to surgery, we injected rats with the analgesic Rymadil® (5 mg/kg; Merial, Velserbroek, The Netherlands) Surgery was performed under isoflurane gas anesthesia (PCH; Haarlem). The surgical area on the skull was shaved & disinfected before surgery and the skin was locally anesthetized using a subcutaneous injection of 0.1 ml 10% lidocaine. Craniotomies above LH, VTA or NaCs were performed, followed by AAV injections using a micro-syringe connected to a micropump (UltraMicroPump 4, WPI surgical instruments), which was lowered into the brain until it reached the DV coordinates (see below for details). Additionally, for fibre photometry and optogenetics experiments, 6 screws were drilled on the skull to support the cement needed for fibre anchoring. Next, the optic fibres were implanted using the same stereotactic apparatus and bregma-relative coordinates as used for the AAV injection. The optic fiber implant(s) (when applicable) were then secured to the skull using dental cement (IV Tetric EvoFlow 2g A1, Henry Schein, Almere) and jewellers screws. Rymadil (5 mg/kg; s.c.) was administered for 2 days after the surgery. Rats were given one week of recovery following surgery.

AAV combinations

For all experiments, we injected a cocktail of one AAV expressing Cre under the GAD promotor, and another AAV encoding Cre-dependent jGCamP7f, GTACR2, or control GFP or mCherry. For unilateral inhibition experiments, we additionally injected an AAV-retro encoding flp in LH, and an AAV encoding FLP-hm4di or control mCherry.

<u>AAV injections for validation experiments</u>

1µl of AAV solution was injected bilaterally into LH (AP: + 2.3, ML: 2, DV: -8.4 to -8, two infusions) over 2.5 min, the needle was then left in place for 5 min, brought up to DV:-8, and then injected again for 2.5 min and left to diffuse for 5 min.

<u>AAV injections for fiber photometry</u>

1μl of AAV solution was injected unilaterally into LH (AP: + 2.3, ML: 2, DV: -8.4 to -8, two infusions) over 2.5 min, the needle was then left in place for 5 min, brought up to DV:-8, and then injected again for 2.5 min and left in place for 5 min. A 400μ optic fiber (Doric Lenses) was then implanted above LH.

AAV injections for optogenetics

0.5μl of AAV solution was injected unilaterally into LH (AP: + 2.3, ML: 2, DV: -8.4 to -8, two infusions) over 2.5 min, the needle was then left in place for 5 min, brought up to DV:-8, and then injected again for 2.5 min and left in place for 5 min. A 200μ optic fiber (Doric Lenses) was then implanted above LH.

AAV injections for chemogenetics

0.5μl of AAV solution was injected unilaterally into either NAcS(AP: + 1.7, ML: + 2.4, DV: -7.6) or VTA (AP: -5.4, ML: + 2.61, DV: -7.5) over 5min. The needle was then left in place for 5 min.

Viral vectors

Experiment	AAV	Provider & product code	Titer (gc/mL)
All	ssAAV-5/2-hGAD67-chI-iCre-SV40p(A)/ ssAAV-9/2-hGAD67-chI-iCre-SV40p(A)	UZH-vector core v197-9 & v197-5	5.6×10^{12}
Photometry	ssAAV-5/2-hSyn1-chI-dlox-jGCaMP7f(rev)-dlox-WPRE-SV40p(A)	UZH-vector core v319-5	2.4×10^{12}
Chemogenetics – retro-flp	scAAV-retro/2-hSyn1-FLPo-SV40p(A)	UZH-vector core V59-retro	7.2×10^{12}
Chemogenetics	ssAAV-5/2-hSyn1-dFRT-hM4D(Gi)_mCherry(rev)-dFRT-WPRE-hGHp(A)	UZH-vector core v190-5	5.2×10^{12}
Chemogenetics - Control	AAV-5/2-hSyn1-dFRT-mCherry(rev)-dFRT-WPRE-hGHp(A)x-WPRE-hGHp(A)	UZH-vector core v188-5	$6.8 \times 10E12$
Optogenetics	pAAV_hSyn1-SIO-stGtACR2-FusionRed	Addgene 105677	2.1×10^{13}
Optogenetics - Control	ssAAV-5/-hSyn1-dlox-mCherry(rev)-dlox-WPRE-hGHp(A)	UZH-vector core v133-5	6.1×10^{12}

Validation	ssAAV-5/2-hEF1α-dlox-EGFP(rev)-dlox-WPRE-bGHp(A)	UZH-vector core v217-5	4.2x10^12

Behavioural procedures

Intermittent access homecage alcohol

For all behavioural experiments, we first habituated rats to alcohol consumption by using an alcohol homecage procedure (Simms et al., 2008; Wise RA, 1973; Supp. Fig. 1). Rats had intermittent 24h access to a bottle containing 20% ethanol (EtOH), and a bottle containing water. Additionally, we used two control empty cages without animals to account for spillage. After 24h the bottles were weighed and swapped for a single water bottle for the following 24h. These two phases were repeated 12 times. The overall consumption of alcohol per rat after each session was measured in grams, by subtracting pre and post measurements for each cage and the average of the two control cages, accounting for 2-gram spillage and multiplying by the density of 20% ethanol (0.97).

During this phase, we habituated the rats to the optogenetics or photometry patch cords by tethering them to the patchcord and leaving them in their designated behavioural boxes for 30 minutes on the alcohol off days (Tuesday and Thursday). Alcohol magazines were removed from the boxes during habituation. For the chemogenetics inhibition experiment, we habituated the rats to an i.p. Injection by injecting approx. 0.1ml saline i.p. prior to 4 habituation sessions.

Alcohol Pavlovian conditioning task

Conditioning Phase: We placed the rats in the operant chambers described earlier and trained once a day during weekdays (Mon-Fri) for 45 minutes per group, per day. In these sessions, we trained the rats to associate a cue with alcohol delivery in the alcohol magazine (CS+ trials) and a different cue with no programmed consequences (CS- trials). Cues consisted of three different stimuli (white noise, light, or flickering light) counterbalanced across rats. In each session, rats received 8 presentations of each CS (16 trials total). Trials were further divided into three phases (pre-CS, during-CS, and post-CS), each lasting 10 seconds and resulting in a 30-second trial window. Alcohol (20% EtOH) was delivered for 6 seconds into the magazine for CS+ trials. Trials were followed by a variable inter-trial-interval (vITI 2mins). Drinking was assessed by measuring the time beam breaks in the magazines were broken for all trial and vITI phases. Alcohol leftover was measured after each training session to assess the amount animals were drinking. For all experiments, we gave every rat between 19-21 conditioning sessions.

Extinction phase: We placed the rats in the operant chambers and trained them once a day during weekdays for 45 minutes per day. Extinction sessions were procedurally identical to the conditioning sessions, with the exception that no alcohol was delivered after the CS+. Alcohol seeking was again assessed by measuring the time spent in the magazine during all CS trial and vITI phases.

Fluid exposure session: Rats received a single fluid exposure session after the last extinction session. In this session, alcohol was delivered in the alcohol magazine on the same time schedule as the conditioning phase, but the cues were not presented.

Alcohol-prime induced reinstatement test: Alcohol was delivered into the magazine for 6 seconds at the beginning of the test session. During the test session, both cues were presented in a similar manner to the conditioning and extinction sessions, but no alcohol was delivered. Alcohol seeking was measured as the time spent in the magazine during all CS and vITI phases.

Histological validation

Immunohistochemical labelling of MCH and Ox

After a waiting period of approximately 4 weeks to allow for virus expression, we deeply anesthetized rats with isoflurane and Euthasol® injection (i.p.) and transcardially perfused them with ~100 ml of normal saline followed by ~400 ml of 4% paraformaldehyde in 0.1M sodium phosphate (pH 7.4). The brains were removed and post-fixed for 2 h, and then 30% sucrose in 0.1M PBS for 48 h at 4°C. Brains were then frozen on dry ice, and coronal sections were cut (40 μm) using a Leica Microsystems cryostat and stored frozen in 0.1M PBS containing 30% sucrose at -20°C.

We selected a 1-in-4 series and first rinsed free-floating sections (3 x 10 minutes) before incubation in PBS containing 0.5% Triton-X and 10% Normal Donkey Serum (NDS) and incubated for at least 48 h at 4°C in primary antibody. Sections were then repeatedly washed with PBS and incubated for 2-4 h in PBS + 0.5% Triton-X with 2% NDS and secondary antibody. After another series of washes in PBS, slices were stained with DAPI (0.1 ug/ml) for 10 min prior to mounting onto gelatin-coated glass slides, air-drying and cover-slipping with Mowiol and DABCO.

Primary antibodies were guinea pig anti-orexin (1:250; Synapic Systens ART: 389004), and rabbit anti-MCH (1:500; Phoenix Biotech ART: G-070-47). Secondary antibodies were Alexa Fluor 488 anti-guinea pig (1:500; Molecular Probes ART: A11073) and IRDye 680LT anti-rabbit (1:500; Li-cor ART: 926-68023).

In-situ hybridization labelling of GAD-mRNA

We performed RNA in situ hybridization for glutamate decarboxylase1 (GAD1) and eGFP mRNA and according to "User Manual for Fresh Frozen Tissue" from RNAscope Multiplex Fluorescent Reagent Kit V2 assay (Advanced Cell Diagnostics). Specifically, freshly frozen brains were cryosectioned (20 μm) onto Superfrost Plus glass slides (Fisher Scientific) and stored at −80°C. Brain sections were fixed in 10% neutral buffered formalin for 15 min at 4°C, rinsed twice in distilled water, gradually dehydrated for 5 min each in 50%, 70% and twice 100% ethanol. The slides were then incubated with Hydrogen Peroxide for 10 min, and Protease IV for 30 min at RT. In between steps sections were rinsed twice in

1xPBS. After this, GAD1 (316401-C2) and eGFP (400281) specific target probes were applied to the sections and incubated at 40°C for 2 hr in the HybEZ oven (Advanced Cell Diagnostics). Sections were then treated with amplifier probes by applying AMP1 and AMP2 at 40°C for 30 min each, followed by AMP3 at 40°C for 15minfor 30 min. Signals for channel 1 (eGFP, Opal520), and channel 2 (GAD, Opal690) were developed by applying the appropriate HRP probes onto the tissue. Finally, the nuclei were stained using DAPI for 30 s to stain nuclei (blue color). In between steps sections were rinsed twice in 1x Wash Buffer (Advanced Cell Diagnostics).

<u>Histological validation after behavioural experiments</u>

After each behavioural experiment, we deeply anesthetized rats with isoflurane and Euthasol® injection (i.p.) and transcardially perfused them with ~100 ml of normal saline followed by ~400 ml of 4% paraformaldehyde in 0.1M sodium phosphate (pH 7.4). The brains were removed and post-fixed for 2 h, and then 30% sucrose in 0.1M PBS for 48 h at 4°C. Brains were then frozen on dry ice, and coronal sections were cut (40 μm) using a Leica Microsystems cryostat and stored frozen in 0.1M PBS containing 30% sucrose at -20°C.

We selected a 1-in-4 series and first rinsed free-floating sections (3 x 10 minutes) before washing in PBS. Following this, slices were stained with DAPI (0.1 ug/ml) for 10 min prior to mounting onto gelatin-coated glass slides, air-drying and cover-slipping with Mowiol and DABCO.

Image acquisition and neuronal quantification

We used a VectraPolaris slide scanner (VUmc imaging core) to image slides at 10x magnification. For validation of GAD-cre approach experiments (Exp. 1), images from Bregma -2.04 to Bregma -2.76 were scanned and imported into QuPath for analysis. For photometry and optogenetic experiments, images containing LH, from Bregma - 2.04 mm to -2.76 mm were identified and the boundary of expression and fiber placement for each rat was plotted onto the respective Paxinos and Watson atlas. For chemogenetic experiments images from Bregma 2.16 to Bregma 1.56 (NAcS), and Bregma -4.92 to Bregma -5.40 (VTA) were identified and the boundary of expression was plotted onto the respective Paxinos and Watson atlas.

LH was manually labelled using DAPI for identification of anatomical landmarks and boundaries. Each value was the result of an average of each count from 4-5 adjacent sections per rat (approx Bregma -2.04 mm to 2.76 mm). To identify GAD-, eGFP-, MCH-, Ox-, and redFusion-positive cells, we used the 'Cell detection' feature in QuPath, with an identical threshold applied across all sections. To identify GAD+eGFP, MCH+eGFP(or redFusion), and OX+eGFP(or redFusion), each region of interest was exported to ImageJ. The overlays representing the cells (GAD, eGFP, MCH, Ox, and redFusion) were then filled,

converted to a binary layer, and then multiplied using the ImageJ function 'Image calculator'. Double labelling is reported as a percentage of total neurons labelled with the AAV fluorescent reporter for that given region of interest.

Ex vivo slice physiology

Rats were anesthetized (5% isoflurane, i.p. injection of 0.1 ml/g pentobarbital) and perfused with ice-cold N-methyl-D-glucamin (NMDG) solution containing (in mM): 93 NMDG, 2.5 KCl, 1.2 NaH2PO4, 30 NaHCO3, 20 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), 25 glucose, 5 sodium ascorbate, 3 sodium pyruvate, 10 MgSO4, 0.5 CaCl2, at pH 7.3 adjusted with HCl. After removing the brain, 250 μm thick brain slices were cut in ice-cold NMDG solution using a Leica VT1200 vibratome. The slices were transferred to a holding chamber with carbogenated artificial cerebrospinal fluid (aCSF; in mM: 125 NaCl, 3 KCl, 1.2 NaH2PO4, 1 MgSO4, 2 CaCl2, 26 NaHCO3, and 10 glucose). Slices were stored at room temperature for at least 1 hour before transferring them to the recording chamber.

During recordings slices were kept at 34°C in aCSF. Whole cell recordings were made using borosilicate glass patch-pipettes (3–5 MΩ) were used with a K-gluconate-based internal solution containing (in mM): 135 K-gluconate, 4 NaCl, 2 MgATP, 10 phosphocreatine, 0.3 GTP (sodium salt), 0.2 EGTA, 10 HEPES at a pH of 7.3 using a Multiclamp 700B amplifier (Molecular Devices) and pClamp software (Molecular Devices). Optogenetic stimulation was done at 470 nm using a DC4100 4-channel LED-driver (Thorlabs, Newton, NJ) as light source.

Experimental design

Exp 2.a: Monitoring LH-GABA neurons during alcohol pavlovian conditioning (n = 4 female, n = 2 male)

Fig 2A shows the experimental outline. Alcohol delivery coincided with presentation of the CS+, beginning 4 seconds after CS+ onset. All phases were carried out as described in the behavioural procedure.

Exp 2.b: Unilateral chemogenetic inhibition and monitoring neurons during alcohol Pavlovian conditioning (n = 12, female)

Supp. Fig. 4 shows the experimental outline. In this experiment we delivered 0.1 mL of 20% alcohol into the magazine. Alcohol delivery coincided with presentation of the CS+, beginning 4 seconds after CS+ onset. All phases were carried out as described in the behavioural procedure. We used chemogenetics to chemogenetically inactivated projections from VTA or NAcS to LH on the days 1 or 2, and 17 or 18 (counterbalanced d-clz or sal) of the conditioning phases, and on the first extinction day. For tests during other conditioning phase, we injected both GFP and hM4D9i expressing rats with d-clz (0.1 mg/kg injection i.p.) or saline (0.1 mg/kg injection i.p), counterbalanced across the two test days. For the first extinction day test, we injected all animals with d-clz (0.1 mg/kg) before the start of the

session. We found very sparse or off target DREADDs expression (Supp. Fig. 3), leading to sample sizes not big enough to make statistical comparisons. Moreover, we found no d-CLZ effects on LH-GABA activity or behaviour (Supp. Fig. 4) and given the behavioural task was identical, we combined behavioural and neural data from animals in experiments 2a and 2b.

Exp 3: Inhibiting LH-GABA neurons during encoding of alcohol-predictive cues (n = 18, female)

Fig 7A shows the experimental outline. During the presentation of the CS, light (470 nm) was delivered into the LH, thereby inhibiting the GABAergic neurons. Light delivery occurred via a bifurcating patch-cord (Doric Lenses Inc) connected to the optic fibres on the head of the rat. Light delivery occurred only during the conditioning phase, and alcohol was delivered 0.5 seconds after CS+ offset. This was done to avoid potential effects of LH-GABA inhibition on drinking. After conditioning, we tested all rats in a single extinction session. During this session, cues were presented in the same manner as during conditioning, but there was no light delivery during the cue presentation, and no alcohol delivered after CS+ offset.

Statistics

Behavioural data were analysed using GraphPad Prism 9.1.0. Phases were analysed separately. Dependent variables were the averaged time spent in the magazine during CS+ and during CS-, or post CS+ and post CS- per session across phases (Conditioning, Extinction, Reinstatement).

Experiment 2 - Behaviour

We used a repeated measures analysis of variance (ANOVA), with type of CS (CS+, CS-) and session (Across Conditioning sessions, Last Conditioning vs. First Extinction; Across Extinction sessions; Last Extinction vs Reinstatement) as the within-subject factors.

Experiment 2 - Photometry

Recorded signals were first down sampled by a factor of 64, giving a final sampling rate of 15.89 Hz. The 405 nm isosbestic signal was fit to the 490 nm calcium-dependent signal using a first-order polynomial regression. A normalized, motion-artifact-corrected ΔF/F was then calculated as follows: ΔF/F = (490 nm signal − fitted 405 nm signal)/fitted 405 nm signal. The resulting ΔF/F was then detrended via a 90-s moving average, and low-pass filtered at 3 Hz. ΔF/F from 10 s before CS (baseline) to 20 s after CS onset were collated. These traces were then baseline corrected and converted into z-scores by subtracting the mean baseline activity during the first 5 s of the baseline and dividing by the standard deviation of those 5 s.

Traces were grouped by type of CS (CS+ or CS-) and by session (First conditioning, last conditioning, first extinction, last extinction, reinstatement test). Two analysis approaches were used, bootstrapping and permutation tests, the rationale for each is

described in detail in Jean-Richard-Dit-Bressel et al., 2020 and Garreh et al., 2022. The output of every statistical test we conducted is available in the raw data files.

Bootstrapping was used to determine whether calcium activity type of CS was significantly different from baseline ($\Delta F/F = 0$). A distribution of bootstrapped means were obtained by randomly sampling from traces with replacement (n traces for that response type; 5000 iterations). A 95% confidence interval was obtained from the 2.5th and 97.5th percentiles of the bootstrap distribution, which was then expanded by a factor of sqrt (n/(n − 1)) to account for narrowness bias (Jean-Richard-Dit-Bressel et al., 2020).

Permutation tests were used to assess significant differences in calcium activity between CS+ and CS- across the different phases of the task. Observed differences between CS type or session were compared against a distribution of 1000 random permutations (difference between randomly regrouped traces) to obtain a p value per time point. Alpha of 0.05 was Bonferroni corrected based on the number of comparison conditions, resulting in alpha of 0.01. For both bootstrap and permutation tests, only periods that were continuously significant for at least 0.25 s were identified as significant (Jean-Richard-Dit-Bressel et al., 2020).

Experiment 3 - Behaviour

We used a 3-way ANOVA to determine the effects of group (control vs. experimental), type of CS (CS+, CS-), and session (Across Conditioning sessions, Last Conditioning vs. First Extinction) and the interaction between them.

Funding and Disclosure

The work was supported by an NWO VIDI grant (016.Vidi.188.022)

Acknowledgements

The authors gratefully acknowledge the VUmc Histology Imaging Unit for their support & assistance in whole-slide imaging. The authors declare no conflict of interest.

References

Aitta-Aho, T., Pappa, E., Burdakov, D., & Apergis-Schoute, J. (2016). Cellular activation of hypothalamic hypocretin/orexin neurons facilitates short-term spatial memory in mice. *Neurobiology of learning and memory*, *136*, 183-188.

Alonso-Lozares, I. 2023. AlcoholConditioning. GitHub. https://github.com/ialozares/AlcoholConditioning

Anderson, B. A. (2016). What is Abnormal About Addiction-Related Attentional Biases? *Drug and Alcohol Dependence*, *167*, 8–14. https://doi.org/10.1016/j.drugalcdep.2016.08.002

Bittencourt, J. C., Presse, F., Arias, C., Peto, C., Vaughan, J., Nahon, J. L., ... & Sawchenko, P. E. (1992). The melanin-concentrating hormone system of the rat brain: an immuno-and hybridization histochemical characterization. *Journal of Comparative Neurology*, *319*(2), 218-245.

Bonnavion, P., Mickelsen, L.E., Fujita, A., de Lecea, L. and Jackson, A.C. (2016), Hubs and spokes of the lateral hypothalamus: Cell types, circuits and behaviour. J Physiol, 594: 6443-6462. Https://doi.org/10.1113/JP271946—Google

Bouton, M. E. (2004a). Context and behavioral processes in extinction. *Learning & Memory (Cold Spring Harbor, N.Y.)*, *11*(5), 485–494. https://doi.org/10.1101/lm.78804

Bouton, M. E. (2004b). Context and Behavioral Processes in Extinction. *Learning & Memory*, *11*(5), 485–494. https://doi.org/10.1101/lm.78804

Burdakov, D., & Peleg-Raibstein, D. (2020). The hypothalamus as a primary coordinator of memory updating. *Physiology & Behavior*, *223*. https://doi.org/10.1016/j.physbeh.2020.112988

Carvalho, A. F., Heilig, M., Perez, A., Probst, C., & Rehm, J. (2019). Alcohol use disorders. *The Lancet*, *394*(10200), 781–792. https://doi.org/10.1016/S0140-6736(19)31775-1

Chaudhri, N., Sahuque, L. L., Cone, J. J., & Janak, P. H. (2008). Reinstated ethanol-seeking in rats is modulated by environmental context and requires the nucleus accumbens core. *The European Journal of Neuroscience*, *28*(11), 2288–2298. https://doi.org/10.1111/j.1460-9568.2008.06517.x

Cole, S., Hobin, M. P., & Petrovich, G. D. (2015). Appetitive associative learning recruits a distinct network with cortical, striatal, and hypothalamic regions. *Neuroscience*, *0*, 187–202. https://doi.org/10.1016/j.neuroscience.2014.11.026

Cox, V. C., Kakolewski, J. W., & Valenstein, E. S. (1968). Effects of Ventromedial Hypothalamic Damage in Hypophysectomized Rats. *Journal of Comparative and Physiological Psychology*, *65*(1), 145–148. https://doi.org/10.1037/h0020853

Cui, G., Jun, S. B., Jin, X., Luo, G., Pham, M. D., Lovinger, D. M., Vogel, S. S., & Costa, R. M. (2014). Deep brain optical measurements of cell type–specific neural activity in behaving mice. *Nature Protocols*, *9*(6), Article 6. https://doi.org/10.1038/nprot.2014.080

Dana, H., Sun, Y., Mohar, B., Hulse, B. K., Kerlin, A. M., Hasseman, J. P., ... & Kim, D. S. (2019). High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. *Nature methods*, *16*(7), 649-657.

Dayas, C. V., McGranahan, T. M., Martin-Fardon, R., & Weiss, F. (2008). Stimuli Linked to Ethanol Availability Activate Hypothalamic CART and Orexin Neurons in a Reinstatement Model of Relapse. *Biological Psychiatry*, *63*(2), 152–157. https://doi.org/10.1016/j.biopsych.2007.02.002

De Lecea, L., Kilduff, T. S., Peyron, C., Gao, X. B., Foye, P. E., Danielson, P. E., ... & Sutcliffe, J. (1998). The hypocretins: hypothalamus-specific peptides with neuroexcitatory activity. *Proceedings of the National Academy of Sciences*, *95*(1), 322-327.

Delgado, J. M. R., & Anand, B. K. (1953). Increase of food intake induced by electrical stimulation of the lateral hypothalamus. *The American Journal of Physiology*, *172*(1), 162–168. https://doi.org/10.1152/ajplegacy.1952.172.1.162

Den Ouden, H., Kok, P., & De Lange, F. (2012). How Prediction Errors Shape Perception, Attention, and Motivation. *Frontiers in Psychology*, *3*. https://www.frontiersin.org/articles/10.3389/fpsyg.2012.00548

DiFazio, L. E., Fanselow, M., & Sharpe, M. J. (2022). The effect of stress and reward on encoding future fear memories. *Behavioural Brain Research*, *417*, 113587. https://doi.org/10.1016/j.bbr.2021.113587

Duncan, E. A., Rider, T. R., Jandacek, R. J., Clegg, D. J., Benoit, S. C., Tso, P., & Woods, S. C. (2006). The regulation of alcohol intake by melanin-concentrating hormone in rats. *Pharmacology, Biochemistry, and Behavior*, *85*(4), 10.1016/j.pbb.2006.11.004. https://doi.org/10.1016/j.pbb.2006.11.004

Everitt, B. J., & Robbins, T. W. (2016). Drug Addiction: Updating Actions to Habits to Compulsions Ten Years On. *Annual Review of Psychology*, *67*, 23–50. https://doi.org/10.1146/annurev-psych-122414-033457

Field, M., & Cox, W. M. (2008). Attentional bias in addictive behaviors: A review of its development, causes, and consequences. *Drug and Alcohol Dependence*, *97*(1–2), 1–20. https://doi.org/10.1016/j.drugalcdep.2008.03.030

Fleury, M.-J., Djouini, A., Huỳnh, C., Tremblay, J., Ferland, F., Ménard, J.-M., & Belleville, G. (2016). Remission from substance use disorders: A systematic review and meta-analysis. *Drug and Alcohol Dependence*, *168*, 293–306. https://doi.org/10.1016/j.drugalcdep.2016.08.625

Hamlin, A. S., Clemens, K. J., & McNally, G. P. (2008). Renewal of extinguished cocaine-seeking. Neuroscience, 151(3), 659-670.

Hetherington, A. W., & Ranson, S. W. (1940). Hypothalamic lesions and adiposity in the rat. *The Anatomical Record*, *78*(2), 149–172. https://doi.org/10.1002/ar.1090780203

Hoang, I. B., & Sharpe, M. J. (2021). The basolateral amygdala and lateral hypothalamus bias learning towards motivationally significant events. *Current Opinion in Behavioral Sciences*, *41*, 92-97.

Holland, P. C., & Schiffino, F. L. (2016). Mini-review: Prediction errors, attention and associative learning. *Neurobiology of Learning and Memory*, *131*, 207–215. https://doi.org/10.1016/j.nlm.2016.02.014

Hunt, W. A., Barnett, L. W., & Branch, L. G. (1971). Relapse rates in addiction programs. *Journal of Clinical Psychology*, *27*(4), 455–456. https://doi.org/10.1002/1097-4679(197110)27:4<455::aid-jclp2270270412>3.0.co;2-r

Hyman, S. E. (2005). Addiction: A disease of learning and memory. *The American Journal of Psychiatry*, *162*(8), 1414–1422. https://doi.org/10.1176/appi.ajp.162.8.1414

Izawa, S., Chowdhury, S., Miyazaki, T., Mukai, Y., Ono, D., Inoue, R., ... & Yamanaka, A. (2019). REM sleep–active MCH neurons are involved in forgetting hippocampus-dependent memories. *Science*, *365*(6459), 1308-1313.

Janak, P. H., & Chaudhri, N. (2010). The Potent Effect of Environmental Context on Relapse to Alcohol-Seeking After Extinction. *The Open Addiction Journal*, *3*, 76–87. https://doi.org/10.2174/1874941001003010076

Jennings, J. H., Ung, R. L., Resendez, S. L., Stamatakis, A. M., Taylor, J. G., Huang, J., ... & Stuber, G. D. (2015). Visualizing hypothalamic network dynamics for appetitive and consummatory behaviors. *Cell*, *160*(3), 516-527.

Karnani, M. M., Szabó, G., Erdélyi, F., & Burdakov, D. (2013). Lateral hypothalamic GAD65 neurons are spontaneously firing and distinct from orexin- and melanin-concentrating hormone neurons. *The Journal of Physiology*, *591*(Pt 4), 933–953. https://doi.org/10.1113/jphysiol.2012.243493

Key figures on Europe – 2022 edition. (n.d.). Retrieved September 19, 2023, from https://ec.europa.eu/eurostat/web/products-key-figures/-/ks-ei-22-001

Kosse, C., & Burdakov, D. (2019). Natural hypothalamic circuit dynamics underlying object memorization. *Nature communications*, *10*(1), 2505.

Le Pelley, M. E., Mitchell, C. J., Beesley, T., George, D. N., & Wills, A. J. (2016). Attention and associative learning in humans: An integrative review. *Psychological bulletin*, 142(10), 1111.

Lê, A. D., Quan, B., Juzytch, W., Fletcher, P. J., Joharchi, N., & Shaham, Y. (1998). Reinstatement of alcohol-seeking by priming injections of alcohol and exposure to stress in rats. *Psychopharmacology*, *135*(2), 169–174. https://doi.org/10.1007/s002130050498

LeCocq, M. R., Chander, P., & Chaudhri, N. (2023). Blocking μ-opioid receptors attenuates reinstatement of responding to an alcohol-predictive conditioned stimulus through actions in the ventral hippocampus. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *48*(10), 1484–1491. https://doi.org/10.1038/s41386-023-01640-1

LeCocq, M. R., Lahlou, S., Chahine, M., Padillo, L. N., & Chaudhri, N. (2018). Modeling Relapse to Pavlovian Alcohol-Seeking in Rats Using Reinstatement and Spontaneous Recovery Paradigms. *Alcoholism, Clinical and Experimental Research*, *42*(9), 1795–1806. https://doi.org/10.1111/acer.13825

Mahn, M., Gibor, L., Patil, P., Cohen-Kashi Malina, K., Oring, S., Printz, Y., Levy, R., Lampl, I., & Yizhar, O. (2018). High-efficiency optogenetic silencing with soma-targeted anion-conducting channelrhodopsins. *Nature Communications*, *9*(1), Article 1. https://doi.org/10.1038/s41467-018-06511-8

Marchant, N. J., Hamlin, A. S., & McNally, G. P. (2009). Lateral hypothalamus is required for context-induced reinstatement of extinguished reward seeking. *Journal of Neuroscience*, *29*(5), 1331-1342.

Marchant, N. J., Millan, E. Z., & McNally, G. P. (2012). The hypothalamus and the neurobiology of drug seeking. *Cellular and Molecular Life Sciences*, *69*, 581-597.

Marchant, N. J., Rabei, R., Kaganovsky, K., Caprioli, D., Bossert, J. M., Bonci, A., & Shaham, Y. (2014). A Critical Role of Lateral Hypothalamus in Context-Induced Relapse to Alcohol Seeking after Punishment-Imposed Abstinence. *The Journal of Neuroscience*, *34*(22), 7447–7457. https://doi.org/10.1523/JNEUROSCI.0256-14.2014

Marshall, J. F., Turner, B. H., & Teitelbaum, P. (1971). Sensory neglect produced by lateral hypothalamic damage. *Science*, *174*(4008), 523-525.

Marshall, J. F., & Teitelbaum, P. (1974a). Further analysis of sensory inattention following lateral hypothalamic damage in rats. *Journal of Comparative and Physiological Psychology*, *86*(3), 375–395. https://doi.org/10.1037/h0035941

Marshall, J. F., & Teitelbaum, P. (1974b). Further analysis of sensory inattention following lateral hypothalamic damage in rats. *Journal of Comparative and Physiological Psychology*, *86*(3), 375–395. https://doi.org/10.1037/h0035941

Mickelsen, L. E., Kolling, F. W., Chimileski, B. R., Fujita, A., Norris, C., Chen, K., ... & Jackson, A. C. (2017). Neurochemical heterogeneity among lateral hypothalamic hypocretin/orexin and melanin-concentrating hormone neurons identified through single-cell gene expression analysis. *eneuro*, *4*(5).

Mickelsen, L. E., Bolisetty, M., Chimileski, B. R., Fujita, A., Beltrami, E. J., Costanzo, J. T., ... & Jackson, A. C. (2019). Single-cell transcriptomic analysis of the lateral hypothalamic area reveals molecularly distinct populations of inhibitory and excitatory neurons. *Nature neuroscience*, *22*(4), 642-656.

Nakai, J., Ohkura, M., & Imoto, K. (2001). A high signal-to-noise Ca(2+) probe composed of a single green fluorescent protein. *Nature Biotechnology*, *19*(2), 137–141. https://doi.org/10.1038/84397

Nieh, E. H., Matthews, G. A., Allsop, S. A., Presbrey, K. N., Leppla, C. A., Wichmann, R., ... & Tye, K. M. (2015). Decoding neural circuits that control compulsive sucrose seeking. Cell, 160(3), 528-541.

Nieh, E. H., Vander Weele, C. M., Matthews, G. A., Presbrey, K. N., Wichmann, R., Leppla, C. A., Izadmehr, E. M., & Tye, K. M. (2016). Inhibitory Input from the Lateral Hypothalamus to the Ventral Tegmental Area Disinhibits Dopamine Neurons and Promotes Behavioral Activation. *Neuron*, *90*(6), 1286–1298. https://doi.org/10.1016/j.neuron.2016.04.035

O'Brien, C. P., Childress, A. R., McLellan, A. T., & Ehrman, R. (1992). Classical conditioning in drug-dependent humans. *Annals of the New York Academy of Sciences*, *654*, 400–415. https://doi.org/10.1111/j.1749-6632.1992.tb25984.x

Parkes, S. L., & Westbrook, R. F. (2010). The Basolateral Amygdala Is Critical for the Acquisition and Extinction of Associations between a Neutral Stimulus and a Learned Danger Signal But Not between Two Neutral Stimuli. *Journal of Neuroscience*, *30*(38), 12608–12618. https://doi.org/10.1523/JNEUROSCI.2949-10.2010

Peters, J., LaLumiere, R. T., & Kalivas, P. W. (2008). Infralimbic prefrontal cortex is responsible for inhibiting cocaine seeking in extinguished rats. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *28*(23), 6046–6053. https://doi.org/10.1523/JNEUROSCI.1045-08.2008

Peters, J., Kalivas, P. W., & Quirk, G. J. (2009). Extinction circuits for fear and addiction overlap in prefrontal cortex. *Learning & Memory*, *16*(5), 279–288. https://doi.org/10.1101/lm.1041309

Pickens, C. L., Saddoris, M. P., Setlow, B., Gallagher, M., Holland, P. C., & Schoenbaum, G. (2003). Different roles for orbitofrontal cortex and basolateral amygdala in a reinforcer devaluation task. *Journal of Neuroscience*, *23*(35), 11078-11084.

Priddy, F. H., Lewis, D. J. M., Gelderblom, H. C., Hassanin, H., Streatfield, C., LaBranche, C., Hare, J., Cox, J. H., Dally, L., Bendel, D., Montefiori, D., Sayeed, E., Ackland, J., Gilmour, J., Schnepp, B. C., Wright, J. F., & Johnson, P. (2019). Adeno-associated virus vectored immunoprophylaxis to prevent HIV in healthy adults: A phase 1 randomised

controlled trial. *The Lancet. HIV*, *6*(4), e230–e239. https://doi.org/10.1016/S2352-3018(19)30003-7

Reppucci, C. J., & Petrovich, G. D. (2016). Organization of connections between the amygdala, medial prefrontal cortex, and lateral hypothalamus: A single and double retrograde tracing study in rats. *Brain Structure & Function*, *221*(6), 2937–2962. https://doi.org/10.1007/s00429-015-1081-0

Rooke, S. E., Hine, D. W., & Thorsteinsson, E. B. (2008). Implicit cognition and substance use: A meta-analysis. *Addictive Behaviors*, *33*(10), 1314–1328. https://doi.org/10.1016/j.addbeh.2008.06.009

Schultz, W. (1998). Predictive reward signal of dopamine neurons. *Journal of Neurophysiology*, *80*(1), 1–27. https://doi.org/10.1152/jn.1998.80.1.1

Sharpe MJ, Marchant NJ, Whitaker LR, Richie CT, Zhang YJ, Campbell EJ, Koivula PP, Necarsulmer JC, Mejias-Aponte C, Morales M, Pickel J, Smith JC, Niv Y, Shaham Y, Harvey BK, Schoenbaum G. Lateral Hypothalamic GABAergic Neurons Encode Reward Predictions that Are Relayed to the Ventral Tegmental Area to Regulate Learning. Curr Biol. 2017 Jul 24;27(14):2089-2100.e5. doi: 10.1016/j.cub.2017.06.024. Epub 2017 Jul 6. PMID: 28690111; PMCID: PMC5564224.

Sharpe, M.J., Batchelor, H.M., Mueller, L.E. et al. Past experience shapes the neural circuits recruited for future learning. Nat Neurosci 24, 391–400 (2021). https://doi.org/10.1038/s41593-020-00791-4

Sliedrecht, W., de Waart, R., Witkiewitz, K., & Roozen, H. G. (2019). Alcohol use disorder relapse factors: A systematic review. *Psychiatry Research*, *278*, 97–115. https://doi.org/10.1016/j.psychres.2019.05.038

Suzuki, S., Mell, M. M., O'Malley, S. S., Krystal, J. H., Anticevic, A., & Kober, H. (2020). Regulation of Craving and Negative Emotion in Alcohol Use Disorder. *Biological Psychiatry. Cognitive Neuroscience and Neuroimaging*, *5*(2), 239–250. https://doi.org/10.1016/j.bpsc.2019.10.005

Torres, O. V., Walker, E. M., Beas, B. S., & O'Dell, L. E. (2014). FEMALE RATS DISPLAY ENHANCED REWARDING EFFECTS OF ETHANOL THAT ARE HORMONE DEPENDENT. *Alcoholism, Clinical and Experimental Research*, *38*(1), 108–115. https://doi.org/10.1111/acer.12213

Vafaie, N., & Kober, H. (2022). Association of Drug Cues and Craving With Drug Use and Relapse: A Systematic Review and Meta-analysis. *JAMA Psychiatry*, *79*(7), 641–650. https://doi.org/10.1001/jamapsychiatry.2022.1240

Valenstein, E. S., Cox, V. C., & Kakolewski, J. W. (1968). Modification of motivated behavior elicited by electrical stimulation of the hypothalamus. *Science*, *159*(3819), 1119-1121.

Valenstein, E. S., Cox, V. C., & Kakolewski, J. W. (1970). Reexamination of the role of the hypothalamus in motivation. *Psychological Review*, *77*(1), 16–31. https://doi.org/10.1037/h0028581

Wakabayashi, K. T., Feja, M., Baindur, A. N., Bruno, M. J., Bhimani, R. V., Park, J., Hausknecht, K., Shen, R.-Y., Haj-Dahmane, S., & Bass, C. E. (2019). Chemogenetic activation of ventral tegmental area GABA neurons, but not mesoaccumbal GABA terminals, disrupts responding to reward-predictive cues. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *44*(2), 372–380. https://doi.org/10.1038/s41386-018-0097-6

Wakabayashi, Y., Miyatsuka, T., Miura, M., Himuro, M., Taguchi, T., Iida, H., Nishida, Y., Fujitani, Y., & Watada, H. (2022). STAT3 suppression and β-cell ablation enhance α-to-β reprogramming mediated by Pdx1. *Scientific Reports*, *12*(1), Article 1. https://doi.org/10.1038/s41598-022-25941-5

Watabe-Uchida, M., Zhu, L., Ogawa, S. K., Vamanrao, A., & Uchida, N. (2012). Whole-brain mapping of direct inputs to midbrain dopamine neurons. *Neuron*, *74*(5), 858–873. https://doi.org/10.1016/j.neuron.2012.03.017

Wheeler, D. S., Wan, S., Miller, A., Angeli, N., Adileh, B., Hu, W., & Holland, P. C. (2014). Role of lateral hypothalamus in two aspects of attention in associative learning. *The European Journal of Neuroscience*, *40*(2), 2359–2377. https://doi.org/10.1111/ejn.12592