Title
Shared and diverging neural dynamics underlying false and veridical perception

Short title
Neural dynamics underlying false and veridical percepts

Joost Haarsma[1], Dorottya Hetenyi[1], Peter Kok[1]
[1]Wellcome Centre for Human Neuroimaging, UCL Queen Square Institute of Neurology, University College London, London, UK.

Corresponding author
Joost Haarsma, j.haarsma@ucl.ac.uk

Pages
Manuscript: 32 pages

Figures
Manuscript: 6 Figures

Words
Abstract: 248
Introduction: 736
Discussion: 1902

Acknowledgements
This work was supported by a Wellcome/Royal Society Sir Henry Dale Fellowship [218535/Z/19/Z] and a European Research Council (ERC) Starting Grant [948548] to P.K. The Wellcome Centre for Human Neuroimaging is supported by core funding from the Wellcome Trust [203147/Z/16/Z].

Conflict of interests
The authors declare no conflicts of interests.

Abstract

We often mistake visual noise for meaningful images, which sometimes appear to be as convincing as veridical percepts. This suggests that there is considerable overlap between the mechanisms that underlie false and veridical perception. Yet, false percepts must arise at least in part from internally generated signals. Here, we apply multivariate analyses to human MEG data to study the overlap between veridical and false perception across two discrete stages of perceptual inference: discrimination of content (what did I see) and detection (did I see something?). To this end, participants performed a visual discrimination task requiring them to indicate the orientation of a noisy grating, as well as their confidence in having seen a grating. Importantly, on 50% of trials no gratings were presented (noise-only trials). On a subset of these noise-only trials, participants reported seeing a grating with high confidence, dubbed here false percepts. We found that a sensory signal reflecting the content of these false percepts was present both before and after stimulus onset. Uniquely, high confidence false, but not veridical, percepts were associated with increased pre-stimulus high alpha/low beta [11-14Hz] power, potentially reflecting enhanced reliance on top-down signalling on false percept trials. Later on, a shared neural code reflecting confidence in stimulus presence emerged for both false and veridical percepts, as revealed by cross-decoding. These findings suggest that false percepts arise through sensory-like signals reflecting both content and detection signals, similar to veridical percepts, with an increase in pre-stimulus alpha/beta power uniquely contributing to false percepts.

Significance statement

The neural mechanisms underlying false percepts are likely different from those that underlie veridical perception, as the former are generated endogenously, whereas the latter are the result of an external stimulus. Yet, false percepts often get confused for veridical perception, suggesting a converging mechanism. This study explores the extent to which the mechanisms diverge and converge. We found that both high confidence false and veridical percepts were accompanied by content-specific stimulus-like orientation signals, as well as a shared signal reflecting perceptual confidence. In contrast, we found that false, but not veridical, percepts were preceded by increased high alpha/low beta [11-14 Hz] power, possibly reflecting a reliance on endogenous signals.

Introduction

When we try to make sense of our noisy visual surroundings, we sometimes mistake our own internally generated signals for externally caused sensations. For example, a cat owner may have such a strong expectation to see their pet that they misperceive a blurry flash in their visual periphery for it. Despite these false perceptual experiences relying on internally generated signals, they sometimes appear to be just as real as veridical percepts, suggesting at least partially overlapping neural mechanisms between veridical and false perception. However, the extent to which the underlying neural mechanisms are similar remains unclear.

How does the brain make inferences about the world? Perceptual inference has two main aspects: what is the content of the experience, and how sure am I that I perceived it? When a silhouette appears in the dark, we need to infer what it is (a tree, a person?) as well as make a decision on how likely it is that I perceived this object. Indeed, some recent empirical and theoretical work has highlighted the importance of separating these distinct and discrete stages of inference (Mazor et al., 2020a,b; Fleming et al., 2020; Dijkstra et al., 2023). The neural signals underlying false percepts may resemble those of veridical perception in both these aspects, i.e., there are signals reflecting the content of false percepts as well as signals reflecting confidence in having perceived said object. Alternatively, false percepts may not rely on sensory content signals, and only share signals reflecting confidence in having perceived something, rather than nothing. This important distinction also pertains to one of the most influential models of hallucinations, namely the predictive coding account of hallucinations. The predictive coding model states that hallucinations arise through increased influence of prior expectations (Powers et al., 2017; Kafadar et al., 2020; Haarsma et al., 2020; Benrimoh et al., 2023; Schmack et al., 2021). However, it remains unclear whether these priors act on the content level of stimulus discrimination, on the level of stimulus detection, or both. Therefore, characterising at which stages of perceptual inference false percepts arise is of crucial importance to further our understanding of hallucinations, as well as subjective perception more generally.

While false and veridical percepts may share considerable overlap, their neural origins must be separate in some sense, as they rely on different neural inputs by definition – the former being generated internally, and the latter externally. Pre-stimulus oscillatory dynamics are a candidate mechanism for a point of divergence. Previous studies have consistently linked a decrease in alpha power to increased hit rates by inducing a more liberal detection threshold, while not affecting sensitivity, possibly through increasing neuronal excitability (Achim et al., 2013; Benwell et al., 2017; Boncompte et al., 2016; Brüers & VanRullen, 2018; Chaumon & Busch, 2014; Ergenoglu et al., 2004; Iemi et al., 2017; Iemi & Busch, 2018; Mathewson et al., 2014; Samaha et al., 2017, 2020; Van Dijk et al., 2008). Yet, evidence linking decreased alpha to increased false percepts remains mixed (Keil et al., 2014; Lange et al., 2013). In contrast, a second body of work has linked increased alpha and beta to top-down cognitive processes believed to contribute to false percepts. For example, increased beta power plays a role in interpreting ambiguous stimuli in the language (Iversen et al., 2009) and visual domains (Hipp et al., 2011; Okazaki et al., 2008), as well as mental imagery (Bartsch et al., 2015; Villena-González et al., 2018), and perceptual predictions (Auksztulewicz et al., 2017; Bastos et al., 2012, 2020; Engel et al., 2001; Fujioka et al., 2012; Sherman et al., 2016; Turner et al., 2023). Thus, if top-down mechanisms play a role in driving false percepts, we might find enhanced power in the alpha/beta frequency bands to be related to false percepts.

Here, we used multivariate decoding techniques in combination with MEG to study the
overlap between false and veridical perception at the content and detection-level, as well as
to study a potentially unique role for neural oscillations in driving false percepts. To preview,
false percepts resembled veridical perception to the extent that both were associated with
stimulus-like content-specific signals reflecting the perceived gratings, as well as a shared
signal reflecting confidence in stimulus presence. Uniquely however, false, but not veridical
percepts were associated with an increase in pre-stimulus high alpha/low beta (11-14Hz)
power, possibly reflecting endogenous feedback signals. Together, these findings show that
false percepts arise from sensory-like neural mechanisms, similar to veridical perception.

Results

{**Fig. 1 | Experimental design and behavioural findings. a**, An auditory cue was followed by either a low contrast grating embedded in noise (50% of trials), or a noise patch (50%). Participants indicated which orientation they saw and how confident they were that a grating was presented. **b**, One sound predicted the appearance of a 45°, or clockwise, oriented grating, whilst the other predicted a 135°, or anti-clockwise, orientated grating. Auditory cues were 100% valid on grating-present trials. **c,** Participants were more accurate on high confidence trials. **d,** Upon debriefing participants believed that only 18% of trials contained noise, compared to the real number of 50%. **e,** Participants were driven by the cue on noise-only trials when they were aware of the cues meaning. **f,** On noise-only trials, participants reported high confidence in having seen a}

Participants experienced false percepts that were independent of perceptual expectation cues
Participants performed a grating discrimination task under noisy conditions, while on 50% of the trials noise-only patches were presented. Participants indicated both the perceived orientation as well as their degree of confidence in having perceived a grating (regardless of orientation). Participants accurately identified the grating orientation on grating-present trials more often than expected by chance (mean accuracy=0.66, SD=.12, T{24}=6.5, *p*<.001). Furthermore, they were more accurate when they were confident that they had seen a grating (i.e., higher than average confidence across trials) than when they were not (high: mean=.70, SD=.13; low: mean=.59, SD=.11; paired t-test) (T{24}=6.76, *p*<.001) (Fig. 1c), demonstrating that they were able to perform the task and used the confidence ratings in a meaningful way. It is worth repeating that here participants reported their confidence in having seen a grating, rather than confidence in their orientation report, and thus effectively gave a perceptual awareness response (Sandberg & Overgaard, 2015). Participants were slightly more confident on grating-present trials (mean confidence =2.29, SD=.64, on a scale of 1-4) than noise-only trials (mean=2.25, SD=.64) (T{24}=2.28, *p*=.031). Upon debriefing, all participants but one underestimated the frequency of noise-only trials, believing on average that .18 (SD=.16) of trials contained just noise, while the true proportion was .50 (Fig. 1d). Strikingly, participants reported perceiving a grating with high confidence (3 out of 4 or higher) on 35% of noise-only trials (Fig. 1f). The perceptual expectation cues significantly biased which orientation participants perceived on noise-only trials (.55 false percepts congruent with the cue, chance level is .50, T{24}=2.53, *p* =.018). This effect was driven by the individuals who became aware of the meaning of the cues (N = 7 out of 25; Fig. 1e), potentially reflecting a response bias. Indeed, those aware of the cue had significantly stronger effects of the cue (T{24}=4.14, *p*<.001). High confidence false percepts were not more affected by the cues than low confidence percepts, i.e. guesses (T{23}= 1.18, *p*=.24). In sum, these results indicate that participants regularly had false grating percepts, but that these were hardly, if at all, affected by the predictive cues. This is highly consistent with a previous study using the same experimental paradigm (Haarsma et al., 2023).

{**Fig. 2 | Decoding of perceived orientation.** A linear discriminator analyses was used to decode orientation-specific activity from a functional localiser, which was then generalised to the main experiment. The orientation of a clearly presented grating could be decoded from the functional localiser with strong clusters in the temporal generalisation matrix **(a)** and the diagonal **(b). b,** The peak of the localiser window that served as the training timepoint for the generalisation to the main experiment is indicated in shaded red. **c,** Orientation decoding was primarily driven the occipital lobe. **d,** Generalising from the localiser to the main experiment revealed a clear signal reflecting the perceived orientation across all trials. Training on the peak of the localiser decoding (100-110ms) and separating by stimulus presence revealed significant clusters on noise-only trials **(e)** and stimulus present trials **(f).}**

Orientation-specific activity reflecting the perceived stimulus was present both pre- and post-stimulus on high confidence false percept trials

In order to test the hypothesis that false percepts arise from sensory-like processes we performed a linear discriminant analysis (LDA) to decode the orientation of high contrast, task-irrelevant gratings presented in a separate localiser block in order to then test for the presence of these signals during the main experiment. First, we confirmed that the orientation of these high contrast gratings could be decoded from the MEG signal (Fig. 2a). Cluster-based statistics on the diagonal of the temporal generalisation matrix revealed significant decoding from 55ms-270ms post-stimulus ($p<.001$, Cohen's d= 1.87), peaking at 105ms (Fig. 2b). Further, source localisation revealed that the occipital lobe contributed primarily to the decoding of the grating orientation (Fig. 2c). We then generalised from the localiser to the main experiment to see whether sensory signals contributed to high confidence false percepts. Here, we trained the decoder on a 10ms window centred on the peak of the localiser decoding (100-110ms). We subtracted orientation-specific stimulus traces on low confidence trials from high confidence ones, for both grating present and noise-only trials. We focussed on the pre- and post-stimulus time-windows separately (-500 to 0ms and 0 to 1000ms, respectively). First, across all trials we found a clear orientation-specific neural signal for high vs. low confidence trials during the post-stimulus time-window, from 180ms-855ms ($p<.001$, Cohen's d= 0.87) (Fig. 2d). Analysing grating-present and noise-only trials separately revealed orientation-specific signals for grating-present trials from 505ms-610ms and 865ms-940ms ($p=.018$ & $p=.043$, Cohen's d= 0.75 & 0.71) (Fig. 2f), and critically also on noise-only trials from 390ms-510ms ($p=.014$, Cohen's d= 0.68) (Fig. 2e). Further, during the pre-stimulus time window there was significant orientation-specific activity on noise-only trials from -55ms to -5ms & -205ms to -150ms ($p=.047$ & $p=.050$, Cohen's d= 0.64 & 0.52) (Supplementary Fig.1). Note, because only noise patches with low (2%) signal energy for all orientations were selected to be included in the present experiment, this orientation-specific activity could not have arisen from the noise-patch itself. No pre-stimulus orientation-specific signals were found on grating present trials. Taken together, false percepts were associated with orientation-specific sensory-like signals in both the pre-stimulus and post-stimulus time windows.

*{*__Fig. 3 | Decoding of perceptual confidence.__ A linear discriminator analyses was used to identify a perceptual confidence signal. __a,__ A significant confidence signal was found on all trials after stimulus onset. Analysing noise-only __(b)__ and grating-present __(c)__ trials separately revealed similar results. Cross-decoding from noise-only to grating-present trials __(d),__ and vice versa __(e),__ demonstrated that there was a shared neural signal on grating-present and noise-only trials. __f,__ When balancing low and high confidence trial counts decoding was still successful, demonstrating that the effect was not confounded by imbalanced trial counts. __g,__ Percentage absolute signal change of the high confidence condition compared to the low confidence condition in source localisation. Here the combined, noise-only, and grating-present trials in the 1000ms to 2000ms time window are presented separately. Across all three conditions, parietal and frontal cortices contributed most strongly to the linear discrimination analyses. S = onset of stimulus, Or = onset orientation response cue, Cr = onset of confidence response cue.}*

A shared neural code reflecting confidence in stimulus presence emerged 250ms post-stimulus

We next decoded confidence in stimulus presence on both grating-present and noise-only trials using an LDA (Mostert et al., 2015). Rather than training on a separate localiser (during which participants did not give confidence responses), here we used cross-validation to train and test within the main experiment blocks, in order to isolate a signal reflecting participants' confidence in having seen a grating (regardless of its content). Across both grating-present

and noise-only trials, a confidence signal emerged 265ms post stimulus and was sustained
throughout the post-stimulus window (265ms-2000ms, $p<.001$, Cohen's d= 1.6) (Fig. 3a). This
signal was present on both noise-only (410ms-1280, 1280-2000ms, $p<.001$, Cohen's d= 1.1,
1.2) (Fig. 3b) and grating-present trials (425ms-670, 700-2000ms, $p<.001$, Cohen's d= 0.83,
1.26) (Fig. 3c). To test whether the neural representation of perceptual confidence in veridical
and false percepts was the same, we trained on perceptual confidence on noise-only trials,
and decoded the confidence on grating-present trials, and vice versa. Cluster-based analyses
revealed a significant effect in the post-stimulus time window, demonstrating that the
confidence signal on noise-only trials generalised to grating-present trials (235ms-385ms,
415ms-470ms, 480ms-2000ms, $p<.001$, Cohen's d= 0.85, 0.76, 1.38) (Fig. 3d) and vice versa
(225ms-295ms, 320ms-385ms, 415ms-875ms, 885ms-1000ms, $p=.008$, .015, .001, .003,
Cohen's d= 0.93, 0.81, 1.26, 1.19) (Fig. 3e).
To ensure that the decoding results were not confounded by unequal numbers of low and
high confidence trials, we repeated the analyses drawing a random number of trials from the
overrepresented condition equal to the number of trials in the underrepresented condition.
In some individuals this led to very few trials to perform the decoding analyses on, leading to
unreliable results. We therefore removed participants with fewer than 50 trials to train the
decoder on. This control analysis confirmed significant decoding of confidence post-stimulus
(all trials: 260ms-2000ms ; Fig. 3f).
For visualisation purposes, we reconstructed the cortical sources of the confidence signal by
performing source localisation separately on the magnetic fields underlying high and low
confidence perceptual reports and computing a signal difference map. This source
reconstruction was conducted on the 1000-2000ms time window, where decoding was the
strongest. The source map was overlaid on a 3D cortical surface, revealing parietal and frontal
regions for both noise-only and grating-present trials (Fig. 3g).

Within experiment cross-validated decoding of perceived orientation
Note that the confidence decoding analysis involved cross-validated training and testing
within the main experiment, whereas for the orientation decoding analysis we trained on a
separate localiser and generalised to the experiment in order to provide a strong test of the
sensory nature of the orientation signals. For completeness' sake, and to aid comparison with
confidence decoding, we also applied a within experiment cross-validation approach to
decoding the reported orientations (Supplementary Fig. 2). Broadly, this confirmed our
finding that subjective percepts were reflected by orientation-specific neural signals that
were modulated by confidence (Supplementary Fig. 2d). Note, these signals cannot reflect
motor responses, as the response mapping was randomised. When decoding participants
motor responses, a significant cluster was present exclusively in the response time-window
(cluster-based analyses on diagonal, all trials: 1015-2000ms, $p<.0001$, Cohen's D=2.43), which
was not modulated by confidence ($p>.1$).

{**Fig. 4 | Frequency analyses. a,** Scalp topographies of differences in pre-stimulus beta power for high minus
low confidence false percepts. **b,** Difference in average time x frequency between high and low confidence
trials on noise-only trials.. **c,** Difference in average time x frequency between high and low confidence trials on
grating-present trials. **d,** T-map of the difference in source localisation for the 13Hz frequency band +- 1.5Hz
between high and low confidence response on noise-only trials. A network of parietal and occipital regions was
active prior to high confidence false percepts}

False percepts were preceded by an increase in pre-stimulus high alpha/low beta power
To explore whether pre-stimulus neural oscillations played a role in driving false percepts, we
conducted a frequency analyses on the pre-stimulus time window. Specifically, we tested
whether changes in power in the alpha and beta bands preceded high confidence false
percepts, that is, trials on which participants indicated high confidence in having seen a
grating in the absence of one. We tested for an interaction using a two-way repeated
measures ANOVA with confidence and stimulus presence as two-level factors within the -
750ms to 0ms time window (i.e., the time window from the auditory cue to the onset of the
stimulus). There was a significant interaction in the beta band (12-20Hz) (-650ms to -50ms,
p=.0039), but not in the alpha band (8-12Hz, -650ms to 0ms, p=.0939). Post-hoc analyses
revealed that this interaction was driven by an increase in beta power preceding high
confidence false percepts (-650ms to 0ms, p=.0108, cluster-based permutation-test) (Fig. 4a).
We conducted post-hoc analyses to estimate the exact frequencies that drove these effects
by repeating the analyses in steps of 1Hz. This revealed the interaction between stimulus
presence and confidence started in the high alpha band (11Hz, -700ms to -300ms, p=.0499;
12Hz, -700ms to 0ms, p=.002) and continued into the beta band (13Hz, -700ms to 0ms,
p=.0080; 14Hz, -650ms to -50ms, p=.030), as did the increase in beta power preceding high
confidence false percepts (11Hz, -700ms to 0ms, p=.0039; 12Hz, -700ms to 0ms, p=.0059;
13Hz, -750ms to -150ms, p=.0079; 14Hz, -700ms to -400ms, p=.049).
Thus, the increase in power preceding high confidence false percepts was in the high alpha,
low beta band (Fig. 4b). In contrast, no effects were found comparing high and low confidence
reports on grating-present trials for either alpha or beta power, p>.5, Fig. 4c). No effects were
found in the post-stimulus time-window for either false or veridical percepts (p>.5). In sum,
an increase in high alpha/low beta band power preceded high confidence false percepts, but
not high confidence veridical percepts.
To further explore the temporal profile of this effect, we extended the time window to -
2000ms to 0ms. This control analysis revealed that the interaction between stimulus presence
and confidence on beta power started after the onset of the auditory cue (-650ms to -50ms,
p=.018).
We performed source localisation analyses on the difference in beta power prior to high and
low confidence false percepts in the -750 to 0ms pre-stimulus time window centred on 13Hz
(+- 1.5 Hz). This revealed that the increase in beta power arose from a network including
parietal and occipital regions (Fig. 4d).
304
ERF analyses
Finally, we performed ERF analyses to explore whether confidence in stimulus presence was
reflected in the amplitude of the ERFs on grating-present and noise-only trials. For these
purposes we contrasted high and low confidence trials, and computed cluster-based analyses
on these differences. There were no effects of confidence on ERF amplitude on the sensor
level on either noise-only or grating-present trials (p>.5).
311
Discussion
The present study investigated whether the neural mechanisms underlying false and veridical
perception are shared. Specifically, we tested at which stage of perceptual inference
commonalities between veridical and false perception emerge. During a perceptual
discrimination task, participants indicated whether they perceived a grating with a specific
orientation. On a subset of noise-only trials, participants reported seeing gratings with high

confidence, which we dubbed high confidence false percepts. We found that on these false
percept trials a sensory-like signal reflecting the perceived orientation emerged both prior to
and following the onset of the noise stimulus. The same was true for veridical trials, albeit
only post-stimulus. Further, a signal reflecting confidence in having seen a grating emerged
around 250ms on both false percept and veridical trials, originating from a frontal/parietal
network. Cross-decoding between false and veridical trials showed that the neural patterns
underlying this confidence signal was shared. Finally, uniquely to high confidence false
percept trials, there was an increase in pre-stimulus high alpha/low beta power (11-14Hz),
arising from a parietal-occipital network. Together, these findings show that the neural signals
that underlie false and veridical percepts are largely shared both on the level of sensory-like
content signals and in terms of neural signals reflecting confidence in stimulus presence, with
an additional unique role for pre-stimulus oscillations on false percept trials.

The present study distinguished two aspects of perceptual inference: content discrimination
and detection. Recent theoretical work has highlighted the importance of these different
stages of inference, and how they are conceptually different (Mazor et al., 2020; Fleming et
al., 2020; Dijkstra et al., 2023). When applied to false percepts, this conceptual distinction
suggests that these experiences could in principle arise in different ways. That, is a false
percept could occur due to sensory-like activity (a sensory signal resembling a particular
stimulus causes the agent to report a false percept), or through the neural mechanisms that
are concerned with reporting the presence of a stimulus (I saw something rather than
nothing). Here we find evidence suggesting that false percepts arise from low-level sensory-
like signals reflecting the content of these false percepts. To our knowledge, this is the first
study to decode the content of false percepts from the MEG signal and show that they
resemble sensory signals. Two fMRI studies have reported content-specific signals related to
false percepts. First, Pajani et al. (2015) linked spontaneous pre-stimulus orientation-specific
activity to falsely reported orientations in noise. Second, a previous layer-specific fMRI study
from our group used the same experimental paradigm as the present study (Haarsma et al.,
2023). Notably, the behavioural findings across these two studies were highly consistent, with
both reporting a limited effect of cued orientations on perception, which was only significant
in those who became aware of the meaning of the auditory cues. Regardless, in both studies
participants reported high confidence false percepts on noise-only trials in almost identical
proportions (35% and 36% in the present and previous study, respectively). In the layer-
specific imaging study, we found that the content of false percepts was reflected in the middle
layers of the early visual cortex, suggesting a feedforward signal driving the content of false
percepts. Together with the findings of the present study, we speculate that on noise-only
trials, enhanced alpha/beta power and stimulus-specific activity fluctuations in the early
visual cortex work in concert to generate high confidence false percepts. In line with this,
increased alpha/beta power preceding false percepts source localised to a parieto-occipital
network, making it ideally situated to interact with orientation-specific signals in the early
visual cortex, potentially enhancing them through gain modulation. Indeed, previous studies
have linked interactions between these areas to the modulation of sensory activity by priors
(Rahnev et al., 2011). The nature of this potential interaction between alpha/beta oscillations
and stimulus-specific activity fluctuations requires much further investigation.

Beyond a sensory signal reflecting the content of the reported false percepts, we also found
a signal reflecting the confidence in stimulus presence, which emerged 250-300ms post-

stimulus. Cross-decoding between grating-present and noise-only trials reveals that this neural signal was shared between false and veridical percepts. One might argue that this signal reflects motor preparation, potentially due to a facilitated motor response on high confidence trials. However, this is unlikely, because participants were required to provide an orientation response before their confidence response. Because the response-mapping for the orientation response was randomised, a motor response could not be prepared, precluding the possibility for the presence of a motor preparation signal. Instead, the uncovered neural signal is more likely to track confidence in the presence of a stimulus. Previous studies using variations of perceptual discrimination and detection tasks have reported neural correlates of perceptual confidence signals originating from the dorsolateral and (ventral) medial prefrontal cortex (Bang et al., 2020; Bang & Fleming, 2018; Gherman & Philiastides, 2018; Lau & Passingham, 2006; Shekhar & Rahnev, 2018; Yeon et al., 2020). Note that confidence here reflected the degree to which one was certain a stimulus was present, and is therefore strongly linked to stimulus visibility, which previous studies have shown to be encoded in frontal regions (Mazor et al., 2022). A likely neuromodulatory system that contributes to this signal is the dopamine system, which extensively innervates frontal regions (Björklund & Dunnett, 2007). Indeed, a series of studies have shown that dopamine may play an important role in modulating perceptual confidence, with dopaminergic agonists increasing confidence in perceptual detection (Lou et al., 2011). In rats, causally manipulating dopamine modulates confidence in false alarms (Schmack et al., 2021). Further, in primates, subjective stimulus intensity is reflected by the dopamine signal (De Lafuente & Romo, 2011). Most relevantly, a dopaminergic perceptual confidence signal in the caudate emerges around 300ms post-stimulus during decision-making (Lak et al., 2017), aligning with the emergence of the confidence signal in our cluster-based analyses. However, it should be noted that cluster-based analyses preclude exact inferences about the latency of signals (Maris & Oostenveld, 2007; Sassenhagen & Draschkow, 2019).

A number of influential studies have reported that false percepts are driven by overly strong priors (Schmack et al., 2021; Powers et al., 2017; Kafadar et al., 2020; Haarsma et al., 2020). While the false percepts in the present study were not induced by the perceptual expectations that we tried to elicit using predictive auditory cues, we do not think this should be taken as evidence against the predictive coding account of hallucinations. Rather, it highlights a possible additional mechanism through which prior expectations might induce false percepts. How might this work? In theory, perceptual priors could act on either level of stimulus content, or the level of stimulus presence. Indeed, both types of priors could lead to an increased tendency to report false alarms. For example, a cue signalling the onset of a sensory stimulus could increase false alarms by injecting sensory-like signals in the visual system. This could subsequently be mistaken for an external stimulus, as it passes a reality threshold, leading to the experience of a false percept. Indeed, recent work has suggested that imagination might mix with sensory signals, which under certain circumstances can lead to these signals to be mistaken for externally caused signals (Dijkstra et al., 2023). Alternatively, a prior that signals the likely presence of a stimulus, regardless of its content, could drive false percepts by increasing the likelihood of reporting a stimulus being present (Powers et al., 2017; Kafadar et al., 2020; Benrimoh et al., 2023). A study conducted in mice suggests that the dopamine system could be a candidate neurotransmitter system for acting as a prior on stimulus presence. Here, high confidence false percepts were related to, and could be induced by, increasing pre-stimulus dopamine levels in the caudate (Schmack et al.,

2021). Indeed, previous work has linked dopamine in the caudate to perceptual confidence,
thus increasing dopamine in this region pre-stimulus could de facto resemble a prior on
stimulus presence (Lak et al., 2017). Alternatively, dopamine could mediate content-specific
priors downstream. Regardless, it is clear that false percepts could arise in various ways.
Ultimately, future studies are needed to study whether hallucination proneness is associated
with an increased reliance on priors or presence, or priors on content, or both.
418
What could the role of increased alpha/beta power be in driving high confidence false
percepts? As discussed above, in stimulus detection paradigms, it is well known that a
decrease in alpha and beta power is related to increased hit rates (Achim et al., 2013; Benwell
et al., 2017; Boncompte et al., 2016; Brüers & VanRullen, 2018; Chaumon & Busch, 2014;
Ergenoglu et al., 2004; Mathewson et al., 2014; Samaha et al., 2017, 2020; Van Dijk et al.,
2008). However, some studies have found the opposite for false percepts, where it is an
increase in beta power, rather than a decrease, that drives false percepts (Keil et al., 2014;
Poorganji et al., 2023). This suggests that the mechanism underlying perceptual thresholds
for external stimuli (increased excitability) might be different from those for internally driven
false percepts. One possible mechanism is suggested by previous studies which have linked
high alpha/beta power to top-down endogenous signals. A number of studies have
substantiated this, linking high alpha/beta power – often arising from the parietal cortex – to
attention, imagination, illusory perception, and prediction (Arnal & Giraud, 2012;
Auksztulewicz et al., 2017; Bartsch et al., 2015; Bastos et al., 2020; Buschman & Miller, 2007,
2009; De Lange et al., 2013; Engel & Fries, 2010; Fujioka et al., 2012; Hipp et al., 2011; Iversen
et al., 2009; Okazaki et al., 2008; Pesaran et al., 2008; Raposo et al., 2023; Sherman et al.,
2016; Villena-González et al., 2018). Moreover, previous fMRI studies have reported lower
BOLD activity prior to false percepts (Hesselmann et al., 2010; Pajani et al., 2015). Because
alpha/beta power and BOLD activity are inversely correlated (Scheeringa et al., 2011, 2016),
speculatively, increased alpha/beta power might have played a role in these fMRI studies as
well. However, this does not answer the question of what mechanism underlies fluctuations
in beta power in the present study. Speculatively, they might reflect beliefs about stimulus
presence. That is, while the specifically cued orientations (e.g., a low tone predicts a left-tilted
grating) did not influence which orientation participants reported on high confidence false
percepts trials, there was likely still a strong expectation of *a grating* (of either orientation)
being present on each trial, as the participants were not explicitly told that there would be
noise-only trials. This was confirmed by participants grossly underestimating the proportion
of noise-only trials upon debriefing. As previous studies have linked an increase in beta power
to stimulus predictions (Auksztulewicz et al., 2017; Bastos et al., 2020; Fujioka et al., 2012;
Turner et al., 2023), we speculate here that beta power may reflect endogenous trial-by-trial
fluctuations in stimulus expectation, which may have contributed to the high confidence false
percepts in this study. Additionally, top-down mechanisms like imagination or choice history
could have added to these effects (Dijkstra & Fleming, 2023; Urai et al., 2019).
452
In conclusion, the present study revealed that the neural signals underlying false and veridical
perception are largely shared, both at the level of content-specific sensory signals and
confidence in stimulus presence. Unique to false percepts, increased high alpha/low beta
power arising from a parietal occipital network specifically preceded high confidence false
percepts, but not veridical perception. Thereby, the current study sheds light on how false

percepts arise from basic sensory mechanisms causing internal sensory signals to be mistaken

for reality.

Methods
Ethics statement
This study was approved by the University College London Research Ethics Committee
(R13061/RE002) and conducted according to the principles of the Declaration of Helsinki. All
participants gave written informed consent prior to participation and received monetary
compensation (£7.50 an hour for behavioural training, £10 an hour for MEG).
468
469 *Participants*
470 Twenty-five healthy human volunteers with normal or corrected-to-normal vision
471 participated in the MEG experiment. Two participants were excluded due to missing trigger
472 data. The final sample consisted of 23 participants (22 female; age 25 ± 4 years; mean ± SD).
473
Stimuli
Grayscale luminance-defined sinusoidal grating stimuli were generated using MATLAB
(MathWorks, Natick, Massachusetts, United States of America, RRID:SCR_001622) and the
Psychophysics Toolbox (Brainard, 1997). During the behavioural session, the stimuli were
presented on a PC (1024 × 768 screen resolution, 60-Hz refresh rate). During the MEG
recording session, stimuli were projected onto a screen in front of the participant (1920 ×
1200 screen resolution, 60-Hz refresh rate). On grating-present trials (50%), auditory cues
were followed by a grating after a 750ms delay (0.5-cpd spatial frequency, 33ms duration),
displayed in an annulus (outer diameter: 10° of visual angle, inner diameter: 1°, contrast
decreasing linearly to 0 over 0.7° at the inner and outer edges), surrounding a fixation bull's
eye (0.7° diameter). These stimuli were combined with one of 4 noise patches, which resulted
in a 4% contrast grating embedded in 20% contrast noise during the MEG session. On noise-
only trials, one of the 4 noise patches was presented on its own. Noise patches were created
by smoothing pixel-by-pixel Gaussian noise with a Gaussian smoothing filter, ensuring that
the spatial frequency of the noise matched the gratings (Wyart et al., 2012). This was done to
ensure that the noise patches and gratings had similar low-level properties, increasing the
likelihood of false percepts (Pajani et al., 2015). To avoid including noise patches which
contained grating-like orientation signals by chance, 1000 noise patches were processed
through a bank of Gabor filters with varying preferred orientations. Only noise patches with
low (2%) signal energy for all orientations were selected to be included in the present
experiment. The resulting four noise patches were used for all trials throughout the
experiment, in a counterbalanced manner, ensuring that reported false percepts could only
be triggered by internal mechanisms (Haarsma et al., 2023; Pajani et al., 2015). During the
practice session on the first day, the contrast of the gratings was initially high (80%), gradually
decreasing to 4% towards the end of the practice. The central fixation bull's-eye was present
throughout the trial, as well as during the intertrial interval (ITI; randomly varied between
1000 and 1200ms).
501
Experimental procedure
Participant were required to perform a visual perceptual discrimination task. Trials consisted
of an auditory expectation cue, followed by a grating stimulus embedded in noise on 50% of
trials (750ms stimulus onset asynchrony (SOA) between cue and grating). The auditory cue
(high or a low tone) predicted the orientation of the grating stimulus (45° or 135°) on grating-
present trials. On these grating-present trials, a grating with the orientation predicted by the
auditory cue was presented embedded in noise, while on noise-only trials (50%) only a noise

patch was presented. The stimulus was presented for 33ms. After the stimulus disappeared,
the orientation response prompt appeared, consisting of a left and right pointing arrow on
either side of the fixation dot (location was counterbalanced). Participants were required to
select the arrow corresponding to their answer (left arrow for anti-clockwise, or 135°, right
arrow for clockwise, or 45°; 1s response window) through a button press with their right hand,
using either a button box in the MEG, or a keyboard during behavioural training. Subsequently
the letters "CONF?" appeared on the screen probing participants to indicate their confidence
that they had seen a grating (1 = I did not see a grating, 2 = I may have seen a grating, 3 = I
probably saw a grating, 4 = I am sure I saw a grating), using 1 of 4 buttons with their left hand
(1.25s response window). It is worth highlighting here that confidence here reflected
participants' belief that a grating was present, not their confidence in their orientation report.
It therefore reflects a perceptual awareness scale response (Sandberg & Overgaard, 2015).
On the first day of testing, participants took part in a behavioural practice session. The
practice consisted of an instruction phase with 7 blocks of 16 trials where the task was made
progressively more difficult, whilst verbal and written instructions were provided. During
these practice runs, the auditory cues predicted the orientation of the grating stimulus with
100% validity (45° or 135°; no noise-only trials). After the completion of the instructions, the
participants completed 4 runs of 128 trials each, separated into 2 blocks of 64 trials each. In
the first 2 runs the expectation cues were 100% valid, to ensure participants learnt the
association, whilst in the final 2 runs the cues were 75% valid (i.e., the grating had an
unexpected orientation on 25% of trials), to test whether participants might have adopted a
response bias. Grating contrast decreased over the 4 runs, specifically the contrast levels were
7.5, 6, 5, and 4%, while the contrast of the noise patches remained constant at 20%. No noise-
only trials were presented on day 1. On the second day, participants performed the same task
during the MEG recording. During this session, 8 runs were completed, each consisting of 64
trials. This time the grating contrast was fixed at 4% on grating-present trials, and on 50% of
the trials the gratings were omitted and only noise patches were presented, resulting in noise-
only trials (Fig. 1a). On grating-present trials the cues always predicted the orientation of the
grating with 100% validity (Fig. 1b). On noise-only trials the cue was by definition invalid, since
no grating was presented. After each run a localiser run followed where gratings oriented
either 45° or 135° were presented while participants performed a distracting fixation dimming
task. The purpose of this localiser run was to uncover an orientation-specific MEG signal,
which will not feature in the present paper as we were not able to uncover a significant
orientation-specific signal evoked by the noisy stimuli presented in the current experiment.
This analysis did reveal orientation-specific activity during the response window, which will
be reported in detail elsewhere. Each run lasted ~8 minutes, totalling ~64 minutes.
545
Pre-processing of MEG data
MEG was recorded continuously at 600 samples/second, using a whole-head 273 channel
axial gradiometer system (CTF Omega, VSM MedTech), while participants sat upright. A
photodiode was used to measure the onset of the visual stimuli through the presentation of
a small white square in the bottom-right corner of the screen on both localiser trials as well
as main experiment trials. This was done to ensure that the trials were aligned exactly to
stimulus presentation. Note that the white square was not visible to participants, as it was
covered by the electrode. The first experimental run (out of 8) for each subject was removed,
leaving 7 runs for analyses. Trials were segmented 3000ms pre-stimulus and 3000ms post-
stimulus. Movement and eye-blink artefacts were manually selected and inspected before

being rejected from the data. Independent component analyses were applied to the complete
dataset to identify components that reflect eye-blinks as well as cardiac related signals, which
were manually inspected and removed from the data for each subject.
559
ERF analyses
ERF analyses were conducted for exploratory reasons. We tested for differences in ERF
amplitude for high and low confidence trials. Cluster-based analyses where then conducted
on the sensor-level using Monte-Carlo permutation tests (N=10000), at a significance
threshold of $p<.05$ for the initial threshold for determining a significant difference in ERF as
well as for determining significant clusters. ERFs were computed separately for high and low
confidence trials, for both grating-present and noise-only trials.
567
Frequency analyses
Frequency power was estimated across all MEG channels from 2000ms pre-stimulus to
2000ms post-stimulus in steps of 50ms, for the frequencies 2Hz to 30Hz with steps of 1Hz,
using Morlet wavelets (width=7). To test for pre-stimulus changes in alpha and beta power,
we used cluster-based-permutation tests with 10000 iterations, at a significance threshold of
$p < 0.05$, while averaging over the alpha band (8-12Hz) or the beta band (12-20Hz). For pre-
stimulus effects we tested the time window from the onset of the cue to the onset of the
stimulus, i.e. -750ms to 0ms. For post-stimulus effects the time window of interest ranged
from 0ms to 1000ms. We conducted additional exploratory analyses that included the full
pre-stimulus time-window -2000ms and full post-stimulus time-window. Note that these time
windows include the onset of the auditory stimulus (-750ms) and the orientation response
cues (1000ms). We furthermore repeated the analyses for each frequency between 8 and 20
Hz with steps of 1Hz to estimate which specific frequencies drove the effects.
581
Decoding analyses
We decoded both stimulus content (grating orientation) as well as participants' confidence in
having seen a stimulus from the MEG signal using a two-class linear discriminant analysis
(LDA) (Mostert et al. 2015). For content decoding, we were interested in testing whether
subjectively perceived orientations were accompanied by sensory-like orientation signals, on
both false percept and veridical trials. To obtain a measure of a sensory orientation signal, we
trained a decoder on MEG data from separate localiser blocks to distinguish the orientation
of clearly presented (but task-irrelevant) gratings. Localiser blocks were presented at the end
of each run (of which there were 8), which each contained 80 trials of clearly presented
gratings. Once trained, we applied this decoder to the main experiment, to test for
orientation-specific neural activity on false percept and veridical trials. To train the decoder,
the 273 MEG channels were used as features. To confirm the ability of the decoder to retrieve
orientation signals, we also trained and tested it within the localiser data using a leave-one
out procedure, where all localiser blocks except one served as the training data to decode the
remaining block. This procedure was repeated for all blocks. Time points were averaged
across a moving time window containing 17 samples ((17/600) = 28.3ms), with steps of 3
samples ((3/600) = 5ms). The covariance matrix was taken into account in the decoder as
recommended in previous studies to address correlations between neighbouring MEG
sensors (Brouwer & Heeger, 2009; Mostert et al., 2015). For details on the implementation of
the decoder, see Mostert et al. 2015.

For the purpose of decoding confidence in having seen a stimulus, we used a cross-validated
decoding procedure similar to the approach used to decode orientations within the localiser
blocks. Specifically, using a leave-one-block-out procedure, we decoded confidence in having
seen a grating during the main experiment, by training the LDA on binarized confidence
response, such that the confidence responses reflected higher or lower than average
confidence. All other parameters of the LDA were identical to the orientation decoding
analysis.
To test for sensory-like orientation-specific activity during the main experiment we trained on
the group-level peak of the localiser decoding and averaged over a 10ms time window
(100ms-110ms), and generalised to the main experiment. Subsequently, the evidence traces
for trials where a left-tilted grating was reported was subtracted from right-tilted traces,
resulting in relative evidence in favour of the reported stimulus. Subsequently, low
confidence trials were subtracted from high confidence trials to show activity specific for
orientations that the participant indicated to have seen. Then, cluster-based permutation
tests were conducted on the time courses of these signals. In the first step of the permutation
test, clusters were defined by adjacent points that crossed a significance threshold of p <.05.
The number of permutations for the temporal generalisation matrix was limited to 1,000 due
to the large sample space. In contrast, when performing cluster-based analyses on a single
time course, as when we generalised from the localiser to the main experiment, or the
diagonal of the temporal generalisation matrix for the confidence decoding 10,000
permutations were used. A cluster in the true data was considered significant if p<.05 based
on the null distribution generated by the permutations. The time window for the confidence
decoding was initially -1000ms to 2000ms relative to stimulus onset to include the time
window where the orientation responses were given (1000-2000ms). For the analyses
pertaining to finding content specific signals, we focussed on pre- and post-stimulus time
windows separately, and excluded the time window where the orientation was reported (i.e.,
1000ms onwards). We performed a number of follow-up analyses for the confidence
decoding. These follow-up analyses involved cross-decoding between noise-only and grating-
present trials, to explore whether they shared a neural code for perceptual confidence.
Furthermore, we performed control analyses where we balanced the number of trials in the
high and low perceptual confidence conditions by subsampling to the condition with the
fewest trials, to ensure that the effects were not driven by an overrepresentation of a single
condition.
635
Source localisation
To visualise the source of the confidence decoding we performed source localisation analyses.
We did not collect individual anatomical MRI scans for our subjects, instead, a template MRI
and default head and source models as present in the Fieldtrip toolbox
(www.fieldtriptoolbox.org) were used (see below) (Oostenveld et al., 2011). Previous studies
have demonstrated that little anatomical specificity is lost using a group-based template
approach (Holliday et al., 2003).
The spatial pattern that underlies classification in a linear discrimination analysis is driven by
the difference in magnetic fields between the two conditions on which the decoding is based.
Therefore, one can visualise the source of a decoder by estimating the sources of the two
different conditions, and compute the difference (Haufe et al., 2014). For our purposes we
computed the absolute difference within the 1000ms to 2000ms time window (where
decoding was strongest), divided by the source map of the low confidence condition, thereby

estimating percentage signal change. Taking the absolute difference will visualise which
source signals are involved in perceptual confidence, without making assumptions about the
sign of the signal.
We also performed source localisation on the frequency analyses. Here, there was a direct
translation from the univariate effects to the underlying source map, as they seek to refute
the same null-hypothesis. We therefore computed cluster-based statistics on the -750ms to
0ms time window for the difference in 13Hz (+- 1.5Hz) power between high and low
confidence false percept conditions in source space. The same parameters were used as on
the sensor level (10000 permutations, cluster-defining threshold: p=.05, cluster-level
threshold: p=.05).
For both the frequency and decoding source localisation, we used the default forward and
source models from the Fieldtrip toolbox which were then warped to participants' specific
fiducials based on the MEG sensors. The spatial filter was computed for the time windows of
interest in the averaged data, which was subsequently applied separately to the two
conditions of interest (high and low confidence trials). Source localisation was then computed
for the two conditions of interests. For the confidence decoding a percentage absolute signal
change was computed in source space, whereas for the frequency analyses t-maps were
visualised for significant clusters.
667
Behavioural analyses
We tested participants' accuracy scores, as well as the relationship between accuracy and
confidence in stimulus presence. We further tested whether participants were significantly
biased by the auditory cues on noise-only trials, and performed post-hoc tests to see whether
these effects were driven by cue awareness. All tests were conducted within-subject and two-
sided. Statistics were conducted in JASP (JASP team, 2023).
674
675

Figures

Fig. 1 | Experimental design and behavioural findings. a, An auditory cue was followed by either a low contrast grating embedded in noise (50% of trials), or a noise patch (50%). Participants indicated which orientation they saw and how confident they were that a grating was presented. **b**, One sound predicted the appearance of a 45°, or clockwise, oriented grating, whilst the other predicted a 135°, or anti-clockwise, orientated grating. Auditory cues were 100% valid on grating-present trials. **c,** Participants were more accurate on high confidence trials. **d,** Upon debriefing participants believed that only 18% of trials contained noise, compared to the real number of 50%. **e,** Participants were driven by the cue on noise-only trials when they were aware of the cues meaning. **f,** On noise-only trials, participants reported high confidence in having seen a

687   grating on 35% of trials.

688
Fig. 2 | Decoding of perceived orientation. A linear discriminator analyses was used to decode orientation-
specific activity from a functional localiser, which was then generalised to the main experiment. The
orientation of a clearly presented grating could be decoded from the functional localiser with strong clusters in
the temporal generalisation matrix **(a)** and the diagonal **(b). b,** The peak of the localiser window that served as
the training timepoint for the generalisation to the main experiment is indicated in shaded red. **c,** Orientation
decoding was primarily driven the occipital lobe. **d,** Generalising from the localiser to the main experiment
revealed a clear signal reflecting the perceived orientation across all trials. Training on the peak of the localiser
decoding (100-110ms) and separating by stimulus presence revealed significant clusters on noise-only trials **(e)**
and stimulus present trials **(f).**

698
Fig. 3 | Decoding of perceptual confidence. A linear discriminator analyses was used to identify a perceptual
confidence signal. **a,** A significant confidence signal was found on all trials after stimulus onset. Analysing

noise-only **(b)** and grating-present **(c)** trials separately revealed similar results. Cross-decoding from noise-only
to grating-present trials **(d),** and vice versa **(e),** demonstrated that there was a shared neural signal on grating-
present and noise-only trials. **f,** When balancing low and high confidence trial counts decoding was still
successful, demonstrating that the effect was not confounded by imbalanced trial counts. **g,** Percentage
absolute signal change of the high confidence condition compared to the low confidence condition in source
localisation. Here the combined, noise-only, and grating-present trials in the 1000ms to 2000ms time window
are presented separately. Across all three conditions, parietal and frontal cortices contributed most strongly to
the linear discrimination analyses. S = onset of stimulus, Or = onset orientation response cue, Cr = onset of
confidence response cue.

710
Fig. 4 | Frequency analyses. a, Scalp topographies of differences in pre-stimulus beta power for high minus
low confidence false percepts. **b,** Difference in average time x frequency between high and low confidence
trials on noise-only trials.. **c,** Difference in average time x frequency between high and low confidence trials on
grating-present trials. **d,** T-map of the difference in source localisation for the 13Hz frequency band +- 1.5Hz
between high and low confidence response on noise-only trials. A network of parietal and occipital regions was
active prior to high confidence false percepts

717
718
719 **Supplementary Fig. 1 | Additional figures for orientation and stimulus-specific decoding of perceived**
720 **gratings. a,** Subtracting low from high confidence false percepts, a clear signal is present reflecting the
721 orientation of perceived gratings on all trials. **b,** On false percept trials there was additional orientation and
722 stimulus-specific activity reflecting false percepts in the pre-stimulus time window.

723
Supplementary Fig. 2 | Within decoding of reported orientation. a, As a comparison to the within confidence
decoding we used the same approach to decode the reported orientations during the main experiment. Taking
all trials together, we find a signal reflecting the reported percept emerged from 120ms to 315ms post-
stimulus (*p*=.026, Cohen's d=1.02), and from 325ms until the end of the decoding time window of 2000ms
(*p*<.001, Cohen's d=1.13). **b,** This signal was present on both noise-only (295-630ms, 750-1750ms & 1840-
1960ms *p*=.0018, *p*<.001 & *p*=.038, Cohen's d= 1.03, .89 & .65 respectively). **c,** and grating-present trials
(210ms-2000ms, *p*<.001, Cohen's d=1.43). **d,** This signal was modulated by participants confidence in stimulus
presence, such that decoding was stronger for high confidence trials at 485-705ms, which overlaps with the
timewindow for sensory-like signals that underlie false percepts (*p*=.0019, Cohen's d = .77).
733

References

Achim, A., Bouchard, J., & Braun, C. M. J. (2013). EEG amplitude spectra before near

threshold visual presentations differentially predict detection/omission and short–

long reaction time outcomes. *International Journal of Psychophysiology*, *89*(1), 88–

98. https://doi.org/10.1016/j.ijpsycho.2013.05.016

Aitken, F., Menelaou, G., Warrington, O., Koolschijn, R. S., Corbin, N., Callaghan, M. F., &

Kok, P. (2020). Prior expectations evoke stimulus-specific activity in the deep layers

of the primary visual cortex. *PLOS Biology*, *18*(12), e3001023.

https://doi.org/10.1371/journal.pbio.3001023

Albers, A. M., Kok, P., Toni, I., Dijkerman, H. C., & de Lange, F. P. (2013). Shared

Representations for Working Memory and Mental Imagery in Early Visual Cortex.

Current Biology, *23*(15), 1427–1431. https://doi.org/10.1016/j.cub.2013.05.065

Arnal, L. H., & Giraud, A.-L. (2012). Cortical oscillations and sensory predictions. *Trends in

Cognitive Sciences*, *16*(7), 390–398. https://doi.org/10.1016/j.tics.2012.05.003

Auksztulewicz, R., Friston, K. J., & Nobre, A. C. (2017). Task relevance modulates the

behavioural and neural effects of sensory predictions. *PLOS Biology*, *15*(12),

e2003143. https://doi.org/10.1371/journal.pbio.2003143

Bang, D., Ershadmanesh, S., Nili, H., & Fleming, S. M. (2020). Private–public mappings in

human prefrontal cortex. *eLife*, *9*, e56477. https://doi.org/10.7554/eLife.56477

Bang, D., & Fleming, S. M. (2018). Distinct encoding of decision confidence in human medial

prefrontal cortex. *Proceedings of the National Academy of Sciences*, *115*(23), 6082–

6087. https://doi.org/10.1073/pnas.1800795115

Bartsch, F., Hamuni, G., Miskovic, V., Lang, P. J., & Keil, A. (2015). Oscillatory brain activity in

the alpha range is modulated by the content of word-prompted mental imagery:

759   Alpha oscillations during mental imagery. *Psychophysiology*, *52*(6), 727–735.

760   https://doi.org/10.1111/psyp.12405

761 Bastos, A. M., Lundqvist, M., Waite, A. S., Kopell, N., & Miller, E. K. (2020). Layer and rhythm

762   specificity for predictive routing. *Proceedings of the National Academy of Sciences*,

763   *117*(49), 31459–31469. https://doi.org/10.1073/pnas.2014868117

764 Bastos, A. M., Usrey, W. M., Adams, R. A., Mangun, G. R., Fries, P., & Friston, K. J. (2012).

765   Canonical Microcircuits for Predictive Coding. *Neuron*, *76*(4), 695–711.

766   https://doi.org/10.1016/j.neuron.2012.10.038

767 Benwell, C. S. Y., Tagliabue, C. F., Veniero, D., Cecere, R., Savazzi, S., & Thut, G. (2017).

768   Prestimulus EEG Power Predicts Conscious Awareness But Not Objective Visual

769   Performance. *Eneuro*, *4*(6), ENEURO.0182-17.2017.

770   https://doi.org/10.1523/ENEURO.0182-17.2017

771 Björklund, A., & Dunnett, S. B. (2007). Dopamine neuron systems in the brain: An update.

772   *Trends in Neurosciences*, *30*(5), 194–202. https://doi.org/10.1016/j.tins.2007.03.006

773 Boncompte, G., Villena-González, M., Cosmelli, D., & López, V. (2016). Spontaneous Alpha

774   Power Lateralization Predicts Detection Performance in an Un-Cued Signal Detection

775   Task. *PLOS ONE*, *11*(8), e0160347. https://doi.org/10.1371/journal.pone.0160347

776 Brouwer, G. J., & Heeger, D. J. (2009). Decoding and Reconstructing Color from Responses in

777   Human Visual Cortex. *The Journal of Neuroscience*, *29*(44), 13992–14003.

778   https://doi.org/10.1523/JNEUROSCI.3577-09.2009

779 Brüers, S., & VanRullen, R. (2018). Alpha Power Modulates Perception Independently of

780   Endogenous Factors. *Frontiers in Neuroscience*, *12*, 279.

781   https://doi.org/10.3389/fnins.2018.00279

Buschman, T. J., & Miller, E. K. (2007). Top-Down Versus Bottom-Up Control of Attention in

the Prefrontal and Posterior Parietal Cortices. *Science*, *315*(5820), 1860–1862.

https://doi.org/10.1126/science.1138071

Buschman, T. J., & Miller, E. K. (2009). Serial, Covert Shifts of Attention during Visual Search

Are Reflected by the Frontal Eye Fields and Correlated with Population Oscillations.

Neuron, *63*(3), 386–396. https://doi.org/10.1016/j.neuron.2009.06.020

Chaumon, M., & Busch, N. A. (2014). Prestimulus Neural Oscillations Inhibit Visual

Perception via Modulation of Response Gain. *Journal of Cognitive Neuroscience*,

26(11), 2514–2529. https://doi.org/10.1162/jocn_a_00653

Collerton, D., Barnes, J., Diederich, N. J., Dudley, R., Ffytche, D., Friston, K., Goetz, C. G.,

Goldman, J. G., Jardri, R., Kulisevsky, J., Lewis, S. J. G., Nara, S., O'Callaghan, C.,

Onofrj, M., Pagonabarraga, J., Parr, T., Shine, J. M., Stebbins, G., Taylor, J.-P., … Weil,

R. S. (2023). Understanding visual hallucinations: A new synthesis. *Neuroscience &*

Biobehavioral Reviews, *150*, 105208.

https://doi.org/10.1016/j.neubiorev.2023.105208

De Lafuente, V., & Romo, R. (2011). Dopamine neurons code subjective sensory experience

and uncertainty of perceptual decisions. *Proceedings of the National Academy of*

Sciences, *108*(49), 19767–19771. https://doi.org/10.1073/pnas.1117636108

De Lange, F. P., Rahnev, D. A., Donner, T. H., & Lau, H. (2013). Prestimulus Oscillatory

Activity over Motor Cortex Reflects Perceptual Expectations. *The Journal of*

Neuroscience, *33*(4), 1400–1410. https://doi.org/10.1523/JNEUROSCI.1094-12.2013

Dijkstra, N., Bosch, S. E., & Van Gerven, M. A. J. (2017). Vividness of Visual Imagery Depends

on the Neural Overlap with Perception in Visual Areas. *The Journal of Neuroscience*,

37(5), 1367–1373. https://doi.org/10.1523/JNEUROSCI.3022-16.2016

Dijkstra, N., & Fleming, S. M. (2023). Subjective signal strength distinguishes reality from

imagination. *Nature Communications*, *14*(1), 1627. https://doi.org/10.1038/s41467-

023-37322-1

Dijkstra, N., Mostert, P., Lange, F. P. D., Bosch, S., & Van Gerven, M. A. (2018). Differential

temporal dynamics during visual imagery and perception. *eLife*, *7*, e33904.

https://doi.org/10.7554/eLife.33904

Engel, A. K., & Fries, P. (2010). Beta-band oscillations—Signalling the status quo? *Current*

Opinion in Neurobiology, *20*(2), 156–165.

https://doi.org/10.1016/j.conb.2010.02.015

Engel, A. K., Fries, P., & Singer, W. (2001). Dynamic predictions: Oscillations and synchrony

in top–down processing. *Nature Reviews Neuroscience*, *2*(10), 704–716.

https://doi.org/10.1038/35094565

Ergenoglu, T., Demiralp, T., Bayraktaroglu, Z., Ergen, M., Beydagi, H., & Uresin, Y. (2004).

Alpha rhythm of the EEG modulates visual detection performance in humans.

Cognitive Brain Research, *20*(3), 376–383.

https://doi.org/10.1016/j.cogbrainres.2004.03.009

Fujioka, T., Trainor, L. J., Large, E. W., & Ross, B. (2012). Internalized Timing of Isochronous

Sounds Is Represented in Neuromagnetic Beta Oscillations. *The Journal of*

Neuroscience, *32*(5), 1791–1802. https://doi.org/10.1523/JNEUROSCI.4107-11.2012

Gherman, S., & Philiastides, M. G. (2018). Human VMPFC encodes early signatures of

confidence in perceptual decisions. *eLife*, *7*, e38293.

https://doi.org/10.7554/eLife.38293

Haarsma, J., Deveci, N., Corbin, N., Callaghan, M. F., & Kok, P. (2022). *Perceptual*

expectations and false percepts generate stimulus-specific activity in distinct layers of

the early visual cortex [Preprint]. Neuroscience.

https://doi.org/10.1101/2022.04.13.488155

Haarsma, J., Deveci, N., Corbin, N., Callaghan, M. F., & Kok, P. (2023). Expectation cues and

false percepts generate stimulus-specific activity in distinct layers of the early visual

cortex Laminar profile of visual false percepts. *The Journal of Neuroscience*, JN-RM-

0998-23. https://doi.org/10.1523/JNEUROSCI.0998-23.2023

Harrison, S. A., & Tong, F. (2009). Decoding reveals the contents of visual working memory

in early visual areas. *Nature*, *458*(7238), 632–635.

https://doi.org/10.1038/nature07832

Haufe, S., Meinecke, F., Görgen, K., Dähne, S., Haynes, J.-D., Blankertz, B., & Bießmann, F.

(2014). On the interpretation of weight vectors of linear models in multivariate

neuroimaging. *NeuroImage*, *87*, 96–110.

https://doi.org/10.1016/j.neuroimage.2013.10.067

Hesselmann, G., Sadaghiani, S., Friston, K. J., & Kleinschmidt, A. (2010). Predictive Coding or

Evidence Accumulation? False Inference and Neuronal Fluctuations. *PLoS ONE*, *5*(3),

e9926. https://doi.org/10.1371/journal.pone.0009926

Hipp, J. F., Engel, A. K., & Siegel, M. (2011). Oscillatory Synchronization in Large-Scale

Cortical Networks Predicts Perception. *Neuron*, *69*(2), 387–396.

https://doi.org/10.1016/j.neuron.2010.12.027

Holliday, I. E., Barnes, G. R., Hillebrand, A., & Singh, K. D. (2003). Accuracy and applications

of group MEG studies using cortical source locations estimated from participants'

scalp surfaces. *Human Brain Mapping*, *20*(3), 142–147.

https://doi.org/10.1002/hbm.10133

Iemi, L., & Busch, N. A. (2018). Moment-to-Moment Fluctuations in Neuronal Excitability

Bias Subjective Perception Rather than Strategic Decision-Making. *Eneuro*, *5*(3),

ENEURO.0430-17.2018. https://doi.org/10.1523/ENEURO.0430-17.2018

Iemi, L., Chaumon, M., Crouzet, S. M., & Busch, N. A. (2017). Spontaneous Neural

Oscillations Bias Perception by Modulating Baseline Excitability. *The Journal of*

Neuroscience, *37*(4), 807–819. https://doi.org/10.1523/JNEUROSCI.1432-16.2016

Iversen, J. R., Repp, B. H., & Patel, A. D. (2009). Top-Down Control of Rhythm Perception

Modulates Early Auditory Responses. *Annals of the New York Academy of Sciences*,

1169(1), 58–73. https://doi.org/10.1111/j.1749-6632.2009.04579.x

Keil, J., Müller, N., Hartmann, T., & Weisz, N. (2014). Prestimulus Beta Power and Phase

Synchrony Influence the Sound-Induced Flash Illusion. *Cerebral Cortex*, *24*(5), 1278–

1288. https://doi.org/10.1093/cercor/bhs409

Kok, P., Failing, M. F., & de Lange, F. P. (2014). Prior Expectations Evoke Stimulus Templates

in the Primary Visual Cortex. *Journal of Cognitive Neuroscience*, *26*(7), 1546–1554.

https://doi.org/10.1162/jocn_a_00562

Kok, P., Mostert, P., & De Lange, F. P. (2017). Prior expectations induce prestimulus sensory

templates. *Proceedings of the National Academy of Sciences*, *114*(39), 10473–10478.

https://doi.org/10.1073/pnas.1705652114

Lak, A., Nomoto, K., Keramati, M., Sakagami, M., & Kepecs, A. (2017). Midbrain Dopamine

Neurons Signal Belief in Choice Accuracy during a Perceptual Decision. *Current*

Biology, *27*(6), 821–832. https://doi.org/10.1016/j.cub.2017.02.026

Lange, J., Oostenveld, R., & Fries, P. (2013). Reduced Occipital Alpha Power Indexes

Enhanced Excitability Rather than Improved Visual Perception. *The Journal of*

Neuroscience, *33*(7), 3212–3220. https://doi.org/10.1523/JNEUROSCI.3755-12.2013

Lau, H. C., & Passingham, R. E. (2006). Relative blindsight in normal observers and the neural

correlate of visual consciousness. *Proceedings of the National Academy of Sciences*,

103(49), 18763–18768. https://doi.org/10.1073/pnas.0607716103

Lawrence, S. J. D., Van Mourik, T., Kok, P., Koopmans, P. J., Norris, D. G., & De Lange, F. P.

(2018). Laminar Organization of Working Memory Signals in Human Visual Cortex.

Current Biology, *28*(21), 3435-3440.e4. https://doi.org/10.1016/j.cub.2018.08.043

Lou, H. C., Skewes, J. C., Thomsen, K. R., Overgaard, M., Lau, H. C., Mouridsen, K., &

Roepstorff, A. (2011). Dopaminergic stimulation enhances confidence and accuracy

in seeing rapidly presented words. *Journal of Vision*, *11*(2), 15–15.

https://doi.org/10.1167/11.2.15

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data.

Journal of Neuroscience Methods, *164*(1), 177–190.

https://doi.org/10.1016/j.jneumeth.2007.03.024

Mathewson, K. E., Beck, D. M., Ro, T., Maclin, E. L., Low, K. A., Fabiani, M., & Gratton, G.

(2014). Dynamics of Alpha Control: Preparatory Suppression of Posterior Alpha

Oscillations by Frontal Modulators Revealed with Combined EEG and Event-related

Optical Signal. *Journal of Cognitive Neuroscience*, *26*(10), 2400–2415.

https://doi.org/10.1162/jocn_a_00637

Mazor, M., Dijkstra, N., & Fleming, S. M. (2022). Dissociating the Neural Correlates of

Subjective Visibility from Those of Decision Confidence. *The Journal of Neuroscience*,

42(12), 2562–2569. https://doi.org/10.1523/JNEUROSCI.1220-21.2022

Mostert, P., Kok, P., & De Lange, F. P. (2015). Dissociating sensory from decision processes in

human perceptual decision making. *Scientific Reports*, *5*(1), 18253.

https://doi.org/10.1038/srep18253

Okazaki, M., Kaneko, Y., Yumoto, M., & Arima, K. (2008). Perceptual change in response to a

bistable picture increases neuromagnetic beta-band activities. *Neuroscience*

Research, *61*(3), 319–328. https://doi.org/10.1016/j.neures.2008.03.010

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open Source

Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data.

Computational Intelligence and Neuroscience, *2011*, 1–9.

https://doi.org/10.1155/2011/156869

Pajani, A., Kok, P., Kouider, S., & de Lange, F. P. (2015). Spontaneous Activity Patterns in

Primary Visual Cortex Predispose to Visual Hallucinations. *Journal of Neuroscience*,

35(37), 12947–12953. https://doi.org/10.1523/JNEUROSCI.1520-15.2015

Peelen, M. V., & Kastner, S. (2011). A neural basis for real-world visual search in human

occipitotemporal cortex. *Proceedings of the National Academy of Sciences*, *108*(29),

12125–12130. https://doi.org/10.1073/pnas.1101042108

Pesaran, B., Nelson, M. J., & Andersen, R. A. (2008). Free choice activates a decision circuit

between frontal and parietal cortex. *Nature*, *453*(7193), 406–409.

https://doi.org/10.1038/nature06849

Poorganji, M., Zomorrodi, R., Zrenner, C., Bansal, A., Hawco, C., Hill, A. T., Hadas, I., Rajji, T.

K., Chen, R., Zrenner, B., Voineskos, D., Blumberger, D. M., & Daskalakis, Z. J. (2023).

Pre-Stimulus Power but Not Phase Predicts Prefrontal Cortical Excitability in TMS-

EEG. *Biosensors*, *13*(2), 220. https://doi.org/10.3390/bios13020220

Rahnev, D., Lau, H., & De Lange, F. P. (2011). Prior Expectation Modulates the Interaction

between Sensory and Prefrontal Regions in the Human Brain. *Journal of*

Neuroscience, *31*(29), 10741–10748. https://doi.org/10.1523/JNEUROSCI.1478-

11.2011

Raposo, I., Szczepanski, S. M., Haaland, K., Endestad, T., Solbakk, A.-K., Knight, R. T., &

Helfrich, R. F. (2023). Periodic attention deficits after frontoparietal lesions provide

causal evidence for rhythmic attentional sampling. *Current Biology*,

S0960982223013143. https://doi.org/10.1016/j.cub.2023.09.065

Samaha, J., Iemi, L., Haegens, S., & Busch, N. A. (2020). Spontaneous Brain Oscillations and

Perceptual Decision-Making. *Trends in Cognitive Sciences*, *24*(8), 639–653.

https://doi.org/10.1016/j.tics.2020.05.004

Samaha, J., Iemi, L., & Postle, B. R. (2017). Prestimulus alpha-band power biases visual

discrimination confidence, but not accuracy. *Consciousness and Cognition*, *54*, 47–

55. https://doi.org/10.1016/j.concog.2017.02.005

Sánchez-Fuenzalida, N., Van Gaal, S., Fleming, S. M., Haaf, J. M., & Fahrenfort, J. J. (2023).

Predictions and rewards affect decision-making but not subjective experience.

Proceedings of the National Academy of Sciences, *120*(44), e2220749120.

https://doi.org/10.1073/pnas.2220749120

Sandberg, K., & Overgaard, M. (2015). Using the perceptual awareness scale (PAS).

Behavioral Methods in Consciousness Research, 181–196.

Sassenhagen, J., & Draschkow, D. (2019). Cluster-based permutation tests of MEG/EEG data

do not establish significance of effect latency or location. *Psychophysiology*, *56*(6),

e13335. https://doi.org/10.1111/psyp.13335

Scheeringa, R., Fries, P., Petersson, K.-M., Oostenveld, R., Grothe, I., Norris, D. G., Hagoort,

P., & Bastiaansen, M. C. M. (2011). Neuronal Dynamics Underlying High- and Low-

Frequency EEG Oscillations Contribute Independently to the Human BOLD Signal.

Neuron, *69*(3), 572–583. https://doi.org/10.1016/j.neuron.2010.11.044

Scheeringa, R., Koopmans, P. J., Van Mourik, T., Jensen, O., & Norris, D. G. (2016). The

relationship between oscillatory EEG activity and the laminar-specific BOLD signal.

Proceedings of the National Academy of Sciences, *113*(24), 6761–6766.

https://doi.org/10.1073/pnas.1522577113

Schmack, K., Bosc, M., Ott, T., Sturgill, J. F., & Kepecs, A. (2021). Striatal dopamine mediates

hallucination-like perception in mice. *Science*, *372*(6537), eabf4740.

https://doi.org/10.1126/science.abf4740

Shekhar, M., & Rahnev, D. (2018). Distinguishing the Roles of Dorsolateral and Anterior PFC

in Visual Metacognition. *The Journal of Neuroscience*, *38*(22), 5078–5087.

https://doi.org/10.1523/JNEUROSCI.3484-17.2018

Sherman, M. T., Kanai, R., Seth, A. K., & VanRullen, R. (2016). Rhythmic Influence of Top–

Down Perceptual Priors in the Phase of Prestimulus Occipital Alpha Oscillations.

Journal of Cognitive Neuroscience, *28*(9), 1318–1330.

https://doi.org/10.1162/jocn_a_00973

Stokes, M., Thompson, R., Cusack, R., & Duncan, J. (2009). Top-Down Activation of Shape-

Specific Population Codes in Visual Cortex during Mental Imagery. *The Journal of*

Neuroscience, *29*(5), 1565–1572. https://doi.org/10.1523/JNEUROSCI.4657-08.2009

Stokes, M., Thompson, R., Nobre, A. C., & Duncan, J. (n.d.). *Shape-specific preparatory*

activity mediates attention to targets in human visual cortex.

Turner, W., Blom, T., & Hogendoorn, H. (2023). Visual Information Is Predictively Encoded in

Occipital Alpha/Low-Beta Oscillations. *The Journal of Neuroscience*, *43*(30), 5537–

5545. https://doi.org/10.1523/JNEUROSCI.0135-23.2023

Urai, A. E., De Gee, J. W., Tsetsos, K., & Donner, T. H. (2019). Choice history biases

subsequent evidence accumulation. *eLife*, *8*, e46331.

https://doi.org/10.7554/eLife.46331

Van Dijk, H., Schoffelen, J.-M., Oostenveld, R., & Jensen, O. (2008). Prestimulus Oscillatory

Activity in the Alpha Band Predicts Visual Discrimination Ability. *The Journal of*

Neuroscience, *28*(8), 1816–1823. https://doi.org/10.1523/JNEUROSCI.1853-07.2008

Villena-González, M., Palacios-García, I., Rodríguez, E., & López, V. (2018). Beta Oscillations

Distinguish Between Two Forms of Mental Imagery While Gamma and Theta Activity

Reflects Auditory Attention. *Frontiers in Human Neuroscience*, *12*, 389.

https://doi.org/10.3389/fnhum.2018.00389

Wyart, V., Nobre, A. C., & Summerfield, C. (2012). Dissociable prior influences of signal

probability and relevance on visual contrast sensitivity. *Proceedings of the National*

Academy of Sciences, *109*(9), 3593–3598. https://doi.org/10.1073/pnas.1120118109

Yeon, J., Shekhar, M., & Rahnev, D. (2020). Overlapping and unique neural circuits are

activated during perceptual decision making and confidence. *Scientific Reports*,

10(1), 20761. https://doi.org/10.1038/s41598-020-77820-6

986