Machine learning approaches identify immunologic signatures of total and intact HIV DNA during long-term antiretroviral therapy

Lesia Semenova[1], Yingfan Wang[1], Shane Falcinelli[2,3], Nancie Archin[2,3], Alicia D Cooper-Volkheimer[5], David M Margolis[2,3,4], Nilu Goonetilleke[3,4], David M Murdoch[5], Cynthia D Rudin[1*], and Edward P Browne[2,3,4*]

[1]Department of Computer Science, Duke University
[2]Department of Medicine, UNC Chapel Hill
[3]UNC HIV Cure center UNC Chapel Hill
[4]Department of Microbiology and Immunology, UNC Chapel Hill
[5]Department of Medicine, Duke University
[*]to whom correspondence should be addressed:
cynthia@cs.duke.edu (919) 660-6555
epbrowne@email.unc.edu (919) 843-1058

Abstract: Antiretroviral therapy (ART) halts HIV replication; however, cellular / immue cell viral reservoirs persist despite ART. Understanding the interplay between the HIV reservoir, immune perturbations, and HIV-specific immune responses on ART may yield insights into HIV persistence. A cross-sectional study of peripheral blood samples from 115 people with HIV (PWH) on long-term ART was conducted. High-dimensional immunophenotyping, quantification of HIV-specific T cell responses, and the intact proviral DNA assay (IPDA) were performed. Total and intact HIV DNA was positively correlated with T cell activation and exhaustion. Years of ART and select bifunctional HIV-specific CD4 T cell responses were negatively correlated with the percentage of intact proviruses. A Leave-One-Covariate-Out (LOCO) inference approach identified specific HIV reservoir and clinical-demographic parameters that were particularly important in predicting select immunophenotypes. Dimension reduction revealed two main clusters of PWH with distinct reservoirs. Additionally, machine learning approaches identified specific combinations of immune and clinical-demographic parameters that predicted with approximately 70% accuracy whether a given participant had qualitatively high or low levels of total or intact HIV DNA. The techniques described here may be useful for assessing global patterns within the increasingly high-dimensional data used in HIV reservoir and other studies of complex biology.

Keywords.

HIV reservoir, subgroup analysis

Introduction.

The development of ART has vastly improved morbidity and mortality associated with HIV infection. However, a long-lived proviral reservoir precludes a cure of infection (1–4). Furthermore, despite ART,

PWH are still at increased risk of morbidity and mortality related to sequelae of persistent immune activation. This persistent immune activation is thought to be multifactorial: a product of residual immune pertubation from untreated HIV infection, persistent proviral expression during ART, and specific clinical risk factors for immune activation that are enriched amongst PWH. Understanding the interplay between the HIV reservoir and persistent immune activation during long-term ART is important to inform strategies for a cure.

While a number of cell types have been proposed as contributing to the HIV reservoir, the largest and most well-characterized reservoir is within CD4 T cells. Molecular analyses of residual HIV proviruses in CD4 T cells during ART have shown that the reservoir is comprised of a heterogeneous mixture of full-length replication-competent proviruses and a larger set of defective proviruses containing internal deletions or APOBEC-mediated hypermutation(s) (5–7). Infected cells can undergo clonal expansion and contraction over time, potentially in response to homeostatic cues or antigen-driven activation (8–10). Studies of SIV-infected non-human primates and HIV-infected humans show that a pool of latently-infected cells forms very early during infection (3, 11). However, studies suggest the reservoir is enriched in viral variants that were circulating at the time of ART initiation, indicating that initiation of ART may promote the entry of recently infected cells into the reservoir (12, 13).

HIV infected cells are rare in peripheral blood, making measuring and characterizing the reservoir challenging. Additionally, the size of the reservoir is highly variable between individuals, ranging over multiple logs in scale. During therapy, a subset of the reservoir remains transcriptionally active, expressing viral RNA and, in some cases, viral protein, leading to persistent viremia or "blips" during therapy (14). The factors that regulate reservoir size and transcriptional activity are only beginning to become clear. Similarly, the impact of the latent reservoir on the immune system of PWH is not fully understood. Over time on therapy, intact proviruses become preferentially depleted from transcriptionally active genomic regions and enriched in heterochromatic regions, suggesting that ongoing viral expression promotes infected cell depletion, either through a cytopathic effect from viral proteins or from immune killing of infected cells with residual expression (15).

In addition to the presence of a latent HIV reservoir, PWH on therapy exhibit a persistent level of T cell immune activation and other altered phenotypes compared to uninfected individuals, involving cells of both the innate and adaptive immune systems. Specifically, HIV infection is associated with increased levels of inflammatory biomarkers and increased levels of activation for immune cells (16–19). PWH also exhibit a depleted naïve T cell compartment (20) and increased expression of activation/exhaustion and senescence markers such as PD-1 and KLRG1 (21–24). These alterations may contribute to increased levels of disease and death in PWH. The molecular mechanisms underlying persistent immune activation and dysfunction for PWH on ART are unclear, but it is possible that ongoing viral gene expression and reverse transcription generates ligands for pattern recognition receptors in the innate immune sys-

tem (25). Viral protein expression during therapy likely maintains antigen-specific cells of the adaptive immune system. Given that viral protein expression may be relatively rare during therapy, another important potential cause of persistent immune activation during ART is persistent sequelae of the massive disruptions to T cell homeostasis and lymphoid structures that occur during untreated HIV infection.

It is possible that the HIV reservoir and the host immune system interact in multiple complex ways during therapy. The dynamics of how the HIV reservoir affects the immune system and vice versa are important to characterize and may lead to novel ideas to promote reservoir depletion or to inhibit immune dysfunction that may be related to ongoing HIV expression in persistently infected cells. A number of previous studies have examined the relationship between reservoir size and the immune system (26–45). Consistent major findings from these studies include enrichment of the HIV reservoir in CD4 T cells expressing activation and exhaustion markers, positive correlation of the reservoir with central memory T cells (Tcm) and transitional memory (Ttm) T cell subsets, and inverse correlation of the HIV reservoir with the CD4 nadir and the CD4/CD8 T cell ratio. One important limitation of most of these studies is that they were not able to distinguish intact proviruses from defective proviruses, making the relationship between the immune system and the intact reservoir unclear. Recent studies indicate enrichment of the intact HIV reservoir in cells expressing VLA-4, but no correlation with classic T cell activation markers (26, 37). Additional assessment of immune correlates of the intact HIV reservoir in additional cohorts is needed. Furthermore, most of these studies examined a limited number of surface markers and may have missed associations that would be revealed by a higher-resolution examination of immunophenotypes. Finally, there is a recent, growing appreciation for the profound influence of ART initiation timing and length of treatment on the size and composition of the HIV reservoir that must be accounted for in assessing persistent immune activation in PWH on therapy (46, 47).

In this study, we use a mixture of traditional statistical approaches and novel data science methods to assess the relationship between the immune system and the HIV reservoir across a cohort of 115 PWH. Collectively, this study highlights the utility of machine learning approaches to identify otherwise imperceptible global patterns in high-parameter studies of the HIV reservoir. Additionally, these approaches corroborate recent findings in the HIV persistence field regarding selective intact proviral decay and immune dynamics and highlight the complex immunologic signatures of HIV latency.

Results.

Cohort description, reservoir quantification, and immunophenotyping.
115 participants with HIV infection for at least one year and who had been receiving antiretroviral therapy (ART) for at least 0.9 years (median 9 years, IQR 5.2-16.6) were studied **(Table 1)**. The study cohort was 77 percent male and median participant age was 45.

The Intact Proviral DNA assay (IPDA) was performed on isolated total CD4 T cells (48). The IPDA

enables estimation of the frequency of total proviruses, intact proviruses, and the percent intact of total proviruses for each participant. Immunophenotyping was performed using 25-color spectral flow cytometry on peripheral blood mononuclear cells (PBMCs) from each study participant. A representative flow gating is shown in **Figure S1**. Briefly, live lymphocytes were first defined as T cells (CD3+/CD56-), NK cells (CD3-/CD56+), or NKT cells (CD3+/CD56+). T cells were then further subdivided as CD4+ or CD8+ T cells, and then as naïve T cells (Tn, CD45RA+/CCR7+), central memory T cells (Tcm, CD45RA-/CCR7+), effector memory cells (Tem, CD45RA-/CCR7-) and terminally differentiated effector cells (Teff, CD45RA+/CCR7-). Expression of various surface markers within each of these subsets was then examined (CD38, HLA-DR, PD-1, KLRG1, CD127, CD27, NKG2A). In addition to measuring the baseline abundance of surface proteins, we also stimulated PBMCs from each participant with a pool of HIV-derived peptides and measured intracellular cytokines TNFα, IL-2, and IFNγ as well as the surface expression of the degranulation marker CD107a following background subtraction of the signal from unstimulated cells. Frequencies of CD4 or CD8 T cells that were single- or double-positive for TNFα, IL-2, IFNγ, or CD107 were used for analysis.

These analyses resulted in the determination of a total of 144 parameters for the cohort of n = 115 PWH. These 144 parameters include 133 immunophenotypic cell frequencies, 3 HIV reservoir parameters (frequency of intact HIV DNA, frequency of total HIV DNA, and percentage intact of total HIV DNA), and 8 clinical-demographic parameters. Clinical-demographic variables (**Table 1**) included age, biological sex, years of ART, estimated years of HIV infection prior to ART (here, we use categorical variables: years before ART=NA, years before ART< 1, years before ART≥ 1), CD4 nadir, most recent CD4 T cell count, and race (Caucasian, African-American, other). The median and interquartile range (IQR) for all immune cells population variables are presented in **Table S2**. No significant differences in the reservoir based on biological sex or race were observed (**Table S1**).

Correlation of total, intact, and percent intact HIV DNA with immunophenotypes.

For the initial analysis, we performed Spearman correlations of the clinical-demographic and immunophenotypic data with HIV reservoir metrics. 69 host variables correlated (unadjusted p < 0.05) with one or more characteristics of the HIV reservoir (**Table S3**). Following Benjamini-Hochberg p-value adjustment for multiple comparisons using a false discovery rate correction at 5%, 31 variables were found to be correlated with total HIV DNA, 3 with intact HIV DNA, and 4 with the percentage of intact HIV DNA (**Table 2**). Overall, this analysis indicated that the HIV reservoir size is significantly associated with the abundance of several immunophenotype frequencies and clinical-demographic variables. Of note, the correlations observed were generally weak to moderate in magnitude (the mean of the absolute values of Spearman correlation coefficients was 0.31 for total HIV DNA, 0.33 for intact HIV DNA, and 0.35 for percent intact HIV DNA).

With regard to clinical-demographic variables, CD4 nadir (Spearman $r = -0.32$) was negatively correlated with total HIV DNA, whereas age ($r = 0.31$) and years of ART ($r = 0.31$) were positively correlated with total reservoir frequencies (**Table 2**). In contrast, years of ART ($r = -0.45$) was negatively correlated with the percentage of intact proviruses (**Table 2**). Given the growing appreciation in the field for the role of ART initiation and timing on the HIV reservoir and immune recovery, we examined this association further (**Figure 1**). Correlation plots of total, intact, and percentage intact HIV DNA versus years of ART were visualized **(Figure 1A-C)**. As expected, the plots showed no significant correlation for intact HIV DNA versus years of ART (**Figure 1B**), while total reservoir size was positively correlated with the time of ART (**Figure 1A**, Spearman $r = 0.31$). Interestingly, when we examined a plot of percent intact proviruses versus time on therapy (**Figure 1C**), we observed a biphasic decay pattern. Indeed, when we fitted piece-wise linear functions with allowances for up to 2 breaks, the best-fit model had decay slope changes at 1.6 and 9 years of ART (R^2 score of piece-wise linear model with two breaks was 44.18%, while linear model without any breaks – 16.3%). In PWH within approximately 0-6 years of ART initiation, a significant proportion of the participants exhibited high fractions of intact proviruses (50-90%). After approximately 6 years of ART, however, the majority of participants had reservoirs for which intact proviruses represented a minor fraction of the overall reservoir. Of note, several immune features also demonstrated statistically significant correlations with years of ART, including CD8 T cell expression of CD38, CD8 Tcm frequencies, and CD107a+IL-2+IFNγ-TNFα- CD8 T cells (**Figures S2, S3, S4, Table S4**).

Examination of immunophenotypic parameter correlations with HIV DNA metrics demonstrated several notable correlations (**Table 2, Figures S2, S3, S4**). As previously reported, CD4 T cell frequencies were negatively correlated with total HIV DNA frequencies ($r = -0.36$), whereas CD8 T cells were positively correlated ($r = 0.41$). This relationship held true for intact HIV DNA frequencies but not for the percentage of intact HIV DNA (**Table 2**). Frequencies of CD4 and CD8 T naïve cells, as well as their expression of CD38, were significantly negatively correlated with total HIV DNA frequency. By contrast, CD4 and CD8 Tcm and Tem cells were weakly positively correlated with total HIV DNA. The direction of these correlations was concordant for intact HIV DNA frequencies, albeit without statistical significance. Markers of T cell activation and exhaustion were weakly but significantly positively correlated with total HIV DNA, including NKG2A+ CD4 T, PD-1+ CD4 T, HLA-DR+ CD4 T, PD-1+ CD8 T, and PD-1+/CCR7+ CD8 T frequencies, among others (**Table 2, Figures S2, S3, S4**). Again, the direction of these correlations were conserved for intact HIV DNA frequencies, albeit without statistical significance. For both intact and total HIV DNA, an inverse correlation with the CD4/CD8 T cell ratio was observed (**Figure S5**).

In general, the magnitude of correlation was stronger for total HIV DNA compared to intact HIV DNA (**Table 2**). This pattern may reflect a larger influence of total HIV DNA on immunophenotype of the

host. A notable exception to this pattern was CD127+ CD4 T cell frequencies, which were significantly negatively correlated with intact but not total HIV DNA. For percentage of intact HIV DNA, negative correlations were observed with the frequencies of 2 subsets of bifunctional HIV peptide-stimulated CD4 T cells (IL-2+TNFα+ and IFNγ+IL-2+), as well as years on ART. Changes in immune markers in this study relative to years of ART were also indicative of progressive immune recovery (**Figure S2-S4 and Table S4**). Specifically, years of ART negatively correlated with CD38 expression on CD8 T cells and positively correlated with the frequency of CD8 Tcm cells as well as CD107+IL-2+IFNγ-TNFα- HIV-specific CD8 T cells (**Table S4**). There are likely many more immune parameters that change in concert with ART as recently reported (47); however, the noise associated with cross-sectional immunophenotype data likely precludes their identification in this study.

Leave-One-Covariate-Out (LOCO) inference identifies specific HIV reservoir and clinical-demographic parameters important for the prediction of several immunophenotypes.

Given the evidence that reservoir and immune recovery dynamics likely occur in concert, we next used a variable importance approach - Leave-One-Covariate-Out (LOCO) inference analysis, to account for potential confounding variables and more carefully assess the relative importance of a given variable to an immunophenotype. The LOCO inference approach is demonstrated in **Figure 2A**. First, a least-squares regression linear model that predicts the dependent variable is fitted on a set of independent variables, and an R^2 value is generated (**Figure 2A Step 1**). Next, one independent variable is excluded from the model, the model is refitted, and an adjusted R^2 value is determined (**Figure 2A Step 2**). Finally, the change in R^2 scores, $\triangle R^2$, is calculated (**Figure 2A Step 3**). This process is then repeated for all independent variables in the model, and the $\triangle R^2$ values are plotted in a heatmap for the variables of interest (**Figure 2B-D**).

We applied the LOCO inference analysis to all 133 immunophenotype parameters (dependent variables). Three separate analyses were conducted for a set of independent variables consisting of age, sex, years of ART, CD4 nadir, recent CD4 account, years of HIV infection prior to ART (=NA, <1, ≥ 1), and 1 of 3 HIV reservoir metrics (total HIV DNA, intact HIV DNA, or percentage intact HIV DNA). Multicollinearity analysis confirmed that independent variables from this analysis were not correlated to a degree that would interfere with model generation (**Table S5**). Each analysis produced a heatmap of 133 dependent immunophenotype variables, with visualization of the variables for each immunophenotype (**Figure S6, Tables S6-S8**). For each of these analyses, we displayed the top 10 variables with the largest $\triangle R^2$ value for LOCO inference incorporating either total (**Figure 2B**), intact (**Figure 2C**), or percentage intact (**Figure 2D**) HIV DNA. In general, the least-squares linear regression models were at best weakly explanatory (R^2 values in range [-8.38%, 31.69%]) of the global variability in immunophenotypes (**Tables S6-S8**, **Figure 2B-D**, **Figure S7**), however, the LOCO analysis is still valuable for understanding

which clinical/demographic variables and HIV characteristics are important to the immunophenotypes.

Across all 3 analyses, age emerged as an important covariate of several immunophenotypes related to T cell subsets, particularly the naïve and central memory compartments, as well as KLRG-1, a marker of immune senescence (**Figure 2 and Figure S6**). This is consistent with well-described age-related changes in T cell subset populations (49, 50). Notably, biological sex was a key variable for model prediction of NK cell frequency and NK cell NKG2A expression. Years of ART was important for predicting the frequencies of HLA-DR+ CD8 T cell subsets including Tcm, Tn, and Tem, but more so for the model incorporating intact, rather than total, HIV DNA (**Figure 2B and 2C**). This suggests that both years on ART and total HIV DNA are important explanatory variables for the variance in CD8 T cell activation levels. Total HIV DNA was also noted to be an important predictor of PD-1+ CD4 T naïve cells, HIV-specific IL-2+TNFα-IFNγ-CD107-CD4 T cells, and CD4/CD8 T cell frequencies, among others (**Figure 2B, Figure S6**). Total HIV DNA and, to a lesser extent, intact HIV DNA were important predictors of CD4 T effector memory HLA-DR+CD38+ expression (**Figure 2B and 2C**). In general, total and intact HIV DNA were identified to be predictive of similar immune features (**Figure S6, Tables S6-S8)**; however, total HIV DNA generally had greater magnitude $\triangle R^2$ than intact HIV DNA. The percentage of intact HIV DNA did not appear to contribute greatly to the predictive power of the model for the variance in the immunophenotypes examined (**Figure S6C, Tables S10, S11**), most likely due to its relatively higher correlation with years of ART (**Figures 1C**). Nonetheless, the LOCO inference findings suggest a complex biological concert of immunologic changes that occur over years of life and suppressive ART, and which may be influenced by the HIV reservoir.

Receiver operator characteristic curve analysis demonstrates distinct immune parameters associated with intact and total reservoir frequencies.

The complex relationship between the HIV reservoir and the immune system led us to hypothesize that machine learning (ML) approaches that combine several parameters into models of the HIV reservoir could be a useful approach to understand this interaction. Before this could be attempted, we decided to first identify a defined set of the most valuable variables from which ML models could be built. To achieve this, we first binarized the 3 reservoir metrics (total, intact, %intact) into high (above median) or low (below median) reservoir groups and generated receiver operator (ROC) curves for the 144 clinical-demographic and immunophenotype parameters (**Figure 3**). The area under the curve (AUC) of the ROC curve indicates the ability of the feature (immune parameter or clinical/demographic information) to correctly identifying a participant as having qualitatively low or high total HIV DNA (**Figure 3A**), intact HIV DNA (**Figure 3B**), or percentage intact HIV DNA (**Figure 3C**). A model that randomly guesses high or low reservoir has an AUC of 0.5. In this analysis, 57 host variables had an AUC higher than 0.6 for one or more of total HIV DNA, intact HIV DNA, or percent intact DNA (**Table S9**). The most effective

individual immune markers for classification of high versus low total HIV DNA included %NKG2A+ CD4 T (AUC = 0.70), %PD-1+ Tn CD4 T (AUC = 0.68), and %CD38+/HLA-DR- CD8 T (AUC = 0.68). In contrast, the most effective markers for classifying based on the frequency of intact HIV DNA included %CD127+ CD4 T (AUC = 0.71), %CD8 T (AUC = 0.66), and %Tn CD8 T (AUC = 0.65). Finally, for percentage intact HIV DNA, years of ART (AUC = 0.72), %CD107-IFNγ-IL-2+TNFα+ CD4 T (AUC = 0.65) and %KLRG1+/PD1+ CD4 T (AUC = 0.65) were the most effective (**Figure 3**).

Similar to the findings in the Spearman correlations (**Table 2**) and LOCO inference analysis (**Figure 2**), there were more variables that had AUC values > 0.6 for total reservoir size (**Table S9**): 44 variables above this threshold for the total reservoir, but only 19 variables for the intact reservoir and 23 variables for percent intact. Overall, this approach allowed us to derive a ranked list of the most predictive immune parameters for each aspect of the HIV reservoir, and these highly ranked features were thus used for subsequent dimension reduction and ML modeling.

Dimension reduction reveals two clusters of PWH with distinct HIV reservoirs.

To examine the overall structure of dataset we employed an unsupervised dimension reduction (DR) machine learning approach (PaCMAP (51)) using the 10 immune cell features with the highest AUC values for classifying participants based on total, intact, or percentage intact HIV DNA (**Figure 3, Table S9**). Interestingly, while no clear clustering was observed when we use intact or percentage intact HIV DNA associated features, we observed two distinct clusters of PWH (cluster 1 and cluster 2, **Figure 4A**) when using total HIV DNA associated features. These clusters were of roughly equal size with 50 participants in cluster 1 and 65 in cluster 2. Projecting total reservoir size (above or below the median) onto the clustering plot, we observe a strong distinction between the two clusters in terms of total HIV DNA reservoir characteristics, (**Figure 4B,C**). Analysis of quantitative reservoir frequencies across clusters (**Figure 4C-E**) demonstrated that cluster 1 is characterized by a smaller absolute intact reservoir frequency, but greater percentage of intact proviruses (**Figure 4C-E**). In contrast, cluster 2 is defined by a larger absolute intact reservoir frequency, but lower percentage of intact proviruses (**Figure 4C-E**).

To gain insight into the immune cell features that distinguish these two clusters, we generated ROC curves for the 133 immune parameters based on their ability to identify membership in cluster 1 versus cluster 2 for each participant (**Figure 4F**). Many features that distinguished the clusters overlapped with the features selected for the clustering analysis, including expression patterns of CD38, HLA-DR, KLRG-1 and PD-1 on CD4 and CD8 T cells and CD4 T naïve cell frequencies. However, some novel features emerged, including CD4 KLRG1-CD27+ frequency, CD8 KLRG1-PD1- frequency, and CD8 Tn frequency (**Figure 4F and Figure S8**). Cluster 2, which had a higher total and intact absolute reservoir size but lower percentage of intact proviruses, was associated with a lower fraction of naïve CD4 and CD8 T cells. Additionally, cluster 2 had generally higher expression of immune exhaustion markers, in-

cluding KLRG1 and PD-1 (**Figure S8**). These data indicate that cluster membership is highly associated with the overall level of immune activation and exhaustion for the individual. We have previous shown that cannabis use reduces the overall level of immune activation and inflammation in PWH, despite having minimal impact on the size of the HOV reservoir (52). Interestingly, a subset of participants in this study had clinical data regarding cannabis (CB) use. CB users were enriched in cluster 1, which had lower levels of immune exhaustion markers (**Figure S8**). Notably, with regard to other clinical variables, cluster 2 had an older median age, longer time on ART, lower CD4 nadir, and lower current CD4 T cell count (**Figure S8**).

Overall, these data demonstrate that, within the cohort of PWH on long-term ART, there were two major clusters of participants with distinct reservoir characteristics and immunophenotypes. These two clusters may reflect the results of a complex concert of age-related immune changes, immune recovery following years of ART, and decay of the intact HIV reservoir, among other possibilities.

Decision tree visualization of PWH with respect to reservoir characteristics.
Since the interaction of the immune system and the HIV reservoir is multifactorial, we hypothesized that models that consider multiple parameters simultaneously could more accurately describe the overall dataset, and provide insights regarding the biology of the HIV reservoir and the host immune system. To accomplish this, we employed a decision tree approach to visualize combinations of variables that classify participants as having high (above median) or low (below median) reservoir size. As compared to the dimension reduction technique, decision tree visualization is an interpretable supervised approach and does not require post-hoc analysis.

We first selected 35 variables with the highest ROC AUC values for either total, intact, or percentage intact HIV DNA to be considered for model generation (**Figure 4**). Using these parameters, we fitted Generalized and Scalable Optimal Sparse Decision Trees (GOSDT) (53) to the data in order to classify high versus low total HIV DNA, intact HIV DNA, or percentage intact HIV DNA. We required the trees to achieve at least 80% accuracy for classifying PWH in the cohort, as well as have at least five PWH in each leaf. Since these trees are based on the entire dataset, these models are thus descriptive rather than predictive.

For the total HIV DNA decision tree, only four immune variables were required to accurately describe high versus low HIV DNA status (**Figure 5A, Figure S9**): CD8 T cell frequency, CD4 nadir, %CD38+HLA-DR- CD8 T cells, and %NKG2A+ CD4 T cells. The tree divided the cohort into five subgroups (leaves), among which three have high total reservoir size and two have low total reservoir size. Comparing the labels provided by GOSDT model with the actual data, the tree achieved 83.5% accuracy (i.e. misclassifying 19 PWH among the overall cohort of 115 PWH). Notably, when we combined all samples from "high total reservoir" leaves and all samples from "low total reservoir" leaves, we observed

a significant difference in the actual median total reservoir size for these two groups (266/M for low total and 1288.5/M for high total, Mann Whitney U test p-value is 3.56e-13, **Figure 5B**).

We then repeated this approach for classifying the cohort participants with respect to intact reservoir size (**Figure 5C**, **Figure S10**). Despite the overall lower correlation between the immune cell phenotypes and the intact reservoir size, this tree nevertheless achieved 82.6% accuracy. The tree had six leaves (three with low intact reservoir, three with high intact reservoir) and relied on %Tn CD8 T cells, %CD107-IFNγ+IL-2+TNFα- CD4 T cells, %CD127+ CD4 T cells, %CD38-HLA-DR+ Tcm CD8 T cells, and CD4 Nadir. When we combined the high intact leaves together and the low intact leaves together, we observed a significant difference in median intact reservoir frequency between the groups (29.5/M for low intact and 101/M for high intact, Mann Whitney U test p-value is 9.59e-08, **Figure 5D**).

Finally, a third iteration of GOSDT generation was performed, classifying the cohort participants with respect to percentage intact of total proviruses (**Figure 5E**, **Figure S11**). This tree achieved 82.4% accuracy and relied on years of ART treatment, %KLRG1+CD27- CD4 T cells, %CD127+ CD4 T cells, and recent clinical CD4 count. This tree has two leaves with low percent intact reservoir and three leaves with high percent intact reservoir. We observed a significant difference in the actual percent intact values between the model generated high percent intact and low percent intact leaves (median 5.7% for low percent intact leaves and median 15.3% for high percent intact leaves: Mann Whitney U test, p-value is 5.5e-10, **Figure 5F**).

Collectively, these decision trees highlight combinations of immune parameters that can accurately describe qualitatively high versus low total, intact, and percentage intact HIV DNA in a cohort of n = 115 PWH. This visualization serves as a basis for mechanistic hypotheses about the interactions of the immune system and HIV reservoir during long-term ART.

Machine learning algorithms identify immune feature combinations that predict high versus low reservoir metrics with ~70% test accuracy.

Although clustering and decision tree analysis permit visualization and understanding of global structures within a dataset, we were curious if combinations of immune and clinical-demographic parameters could actually accurately predict (rather than only visualize) whether a given participant had a high (above median) or low (below median) values with respect to total HIV DNA, intact HIV DNA, or percent intact HIV DNA. We considered five machine learning algorithms including Logistic Regression with L2 regularization (LR), CART, Support Vector Machines with RBF kernel (SVM), Random Forest (RF), and Gradient Boosted Trees (GBT). For reservoir size characteristics, we measured accuracy of the models over 10 random splits of the data into training and test sets (**Figure 6A and Table S10)**. Overall, we found that for total reservoir size Logistic Regression achieved highest mean classification accuracy (69.31%) in test data. The accuracy of these models is likely limited by the sample size (n=115) and noise in the

data, as evidenced by lower R^2 score of machine learning models for non-binarized HIV characteristics (**Figure S12**). To examine the contribution of individual immune features to model performance, we examined logistic regression coefficients for each immune cell variable in the model. Since Logistic Regression coefficients are associated with expected change in log odds (based on $\log_e$), we can think about the coefficient β for variable X in the following way: increasing variable X by one unit multiplies the odds of high reservoir size (probability that the reservoir size is high divided by the probability that the reservoir size is low) by e^β.

In **Figure 6B** we visualize the logistic regression model for one data split among ten we considered for total reservoir size. For this split we observe that higher values of %NKG2A+ CD4 T, %PD-1+ Tn CD4 T, and %Tcm CD8 T are associated with a increased probability of total reservoir size being high. On the other hand, an increase in %Tn CD4 T decreases the odds of high reservoir size. The model visualized in **Figure 6B** achieved 75.86% training and 75% test accuracy.

Similarly, logistic regression models performed well for predicting intact reservoir size compared to other methods (average 65.17% test accuracy, see **Figure 6C**). In **Figure 6D** we visualize the Logistic Regression model for one fixed data split. In this model we observed that higher values of %CD107-IFNγ+IL-2+TNFα- CD4 T and %CD127+ CD4 T, and % CD4 T are associated with lower probability of intact reservoir size being high, while higher values of %KLRG1+CD27+ CD8 T are associated with increased probability of intact reservoir size being high. For analysis of percentage intact HIV DNA, we display training and test accuracy values in **Figure 6E**. For the visualized model for one data split (**Figure 6F**), an increase in %CD127+ CD4 T, years of ART, %KLRG1+PD-1+ CD4 T, and %Tem CD4 T leads to a lower probability of percent intact HIV DNA being high.

Overall, these analyses demonstrate that we can use machine learning tools to construct models that can predict with approximately 70 percent accuracy whether a given PWH has qualitatively low or high total, intact or percentage of HIV DNA. The coefficients identified in each of these models, including NKG2A and CD127 expression on CD4 T cells, identify specific immunologic nodes that serve as a basis for the generation of hypotheses about the interplay between total versus intact HIV DNA and the host the immune system.

Discussion.

In this cross-sectional study of peripheral blood samples from a well-characterized clinical cohort of 115 PWH on long-term ART, we used IPDA and high parameter flow cytometry to better understand the associations between the HIV reservoir and a broad range of immunophenotypes related to T cell differentiation, activation, exhaustion, senescence, HIV-specific T cell responses, as well as NK cell phenotypes. Using data science approaches, specific HIV reservoir features (total, intact, percentage intact HIV DNA) and clinical-demographic variables (age, sex, CD4 nadir, recent CD4 count, years on

ART, years of HIV infection prior to initiating ART) were identified that were particularly important for predicting specific immunophenotypes in PWH on long-term ART. Distinct host immune correlates of intact versus total HIV DNA were also identified. Further, machine learning approaches enabled identification of combinations of immune parameters that accurately classified and predicted whether a given PWH is likely to have qualitatively high or low total, intact, or percentage intact HIV DNA. Collectively, this study provides novel insights into HIV reservoir biology and notably also demonstrates the utility of specific data science and machine learning approaches for HIV reservoir studies. Development and application of these analytic approaches are important to enable biological insights from high-parameter studies of the HIV reservoir and other complex biology, particularly with the advent of high-dimensional techniques such as CYTOF and scRNAseq.

Previous studies have sought to identify immune and other correlates of the total HIV DNA reservoir (26–45). These studies have identified several activation/exhaustion markers that correlate with HIV reservoir frequency, including T cell expression of HLA-DR, CD38, PD-1, CTLA-4, and LAG-3, the CD4/CD8 ratio, and NK cell KLRG1 expression. However, not all studies have found associations of total HIV DNA and immune subsets, particularly for CD4 T cell HLA-DR and CD38 expression(40, 54–56) during long-term ART. The correlative results from this present study corroborate many of these studies, and provide additional insights. A number of correlations between total HIV DNA and immune activation and exhaustion were identified. There were weak to moderate **positive** correlations observed between **total HIV DNA and CD4 T cell expression** of NKG2A, PD-1, KLRG1, **CD27**, HLA-DR as well as CD8 T cell expression of PD-1, CCR7, and CD38. **Total HIV DNA was inversely** correlated with CD8 T cell expression of CD38, **CD4 T cell expression of CD27**, the frequency of naïve CD4 and CD8 T cells, and the CD4/CD8 ratio.

Interestingly, when we examined immune correlates of intact HIV DNA measured by the IPDA, many of the same relationships as described above were noted, but with decreased magnitude and lacking statistical significance, with the exception of the inverse correlation observed for CD4/CD8 T cell ratio and intact HIV DNA. This is consistent with a recent study using the IPDA that did not find associations of intact HIV DNA with CD4 T cell HLA-DR and CD38 co-expression (57), as well as a study using near-full-length sequencing of single cells that found only slight enrichment of HLA-DR or PD-1 in cells with genetically intact, inducible proviruses (37).

Uniquely, however, CD127 expression on CD4 T cells was significantly inversely associated with intact reservoir frequency. CD127 is the alpha-chain IL-7 receptor and is expressed on long-lived memory T cells that are depleted during untreated HIV infection. Interestingly CD127+ cells have recently been found to be highly susceptible to latent HIV infection in tissues (58), and IL-7 is associated with slower natural reservoir decay (27). Furthermore, IL-7 signaling is known to drive expression of VLA-4, which was recently reported to highly enrich for genetically intact, inducible proviruses (37). This obser-

vational finding that intact but not total HIV DNA inversely correlates with CD127 expression on CD4 T cells requires further investigation. Finally, this initial immune correlate analysis demonstrated that the percentage of intact HIV DNA was negatively correlated with years of ART as well as and HIV-specific CD4 T cell responses. These findings are consistent with known selective decay of intact proviruses over years of ART therapy (57, 59, 60).

Dimension reduction machine learning approaches identified two robust clusters of PWH when using total HIV DNA reservoir-associated immune cell frequencies (**Figure 4A**), but not for intact or percentage intact HIV DNA (**Figure 4B and 4C**). This pattern is likely indicative of a more profound association of total HIV DNA with host immunophenotype relative to intact HIV DNA. This is certainly plausible giving the growing appreciation for the immunogenicity and high frequency of defective proviruses within the HIV reservoir (61–64). These clusters exhibited distinct reservoir characteristics, with one cluster being enriched with participants with a larger total reservoir size, a lower percentage of intact viruses, a lower frequency of naïve T cells and elevated expression of activation and exhaustion markers, while the other contained PWH with smaller total reservoirs, higher percentage of intact viruses, higher frequency of naïve T cells and lower expression of activation and exhaustion markers. The basis for the two distinct clusters is unclear, but the cluster membership was notably associated with age and time on therapy. The existence of two distinct clusters of PWH with different immune features and reservoir characteristics could have important implications for HIV cure strategies – these two groups may respond differently to a given approach, and cluster membership may need to be considered to optimize a given strategy. It is tempting to speculate that these two clusters represent differing severity of the initial immune insult of untreated HIV infection, and possibly dichotomous degrees of immune recovery following initiation of ART. These two clusters are also interesting in the context of a previous study examining reservoir characteristics which proposed two major types of viral reservoir within PWH. One type with smaller reservoirs that were enriched in Tcm cells and associated with lower levels of Ki67+ cells and immune activation, and another type with larger reservoirs that were enriched in Ttm cells and associated with higher levels of Ki67+ cells and immune activation (27). It is possible that these two proposed types of reservoirs represent the two clusters we observe using immune markers and dimension reduction, although we were unable to examine the frequency of HIV proviruses specifically within Tcm and Ttm cells in this study.

In addition to unsupervised dimension reduction approaches, we also found that we were able to construct simple, interpretable decision trees that could describe our cohort of PWH with more than 80% accuracy with respect to the size of the reservoir. Furthermore, when we constructed decision trees to classify PWH separately based on total reservoir, intact reservoir and percentage intact, distinct sets of features were important (though some were the same) for each of these models. The decision tree analyses highlight complex immune parameter combinations that serve as a basis for unique hypotheses

about HIV reservoir biology. For instance, lower levels of HIV-specific CD4 T cell responses and lower levels of CD8 Tcm HLA-DR expression were useful for identifying individuals with higher frequencies of intact HIV DNA. This suggests that individuals with higher levels of intact proviruses may have lower levels of HIV-specific immune responses and memory CD8 T cell activation that could facilitate clearance of intact proviruses. Notably, the same relationships were not observed for the total HIV DNA decision tree, highlighting the likely more potent immunogenicity of intact proviruses.Another striking example of machine learning approaches identifying potentially insightful reservoir biology is the percentage intact HIV DNA decision tree. As expected, years of ART was critical to predict the percentage of intact proviruses, but CD4 T cell CD127 expression, CD4 T cell KLRG expression and recent CD4 count also played a role. Notably, following approximately 7 years of ART, lower CD4 T cell CD127 expression or low recent CD4 T cell count were useful to classify individuals with higher percentages of intact proviruses. We hypothesize that this finding suggests incomplete immune recovery (low CD4 T cell CD127 expression and low CD4 T cell count) is associated with a larger percentage of intact proviruses, perhaps because of a lack of dilution with *de novo* naïve CD4 T cells over many years of ART.

Logistic regression models that we trained to classify high versus low total reservoir, intact reservoir, and percentage intact were able to predict out-of-sample with ∼70% accuracy on average. Each of these models relied on five features only and were simple enough for us to visualize and assess how important each variable is for every characteristic of reservoir size. These models corroborated the potential biological relationships identified in the decision tree analysis, and confirmed that these variables could actually be used to predict out-of-sample whether total, intact or percentage intact HIV DNA was qualitatively high or low.

Our findings should be considered in the light of some inherent limitations and caveats. This study is cross-sectional in nature and is primarily observational, so caution should be used interpreting findings associated with time on therapy. As with any observational study, confounding bias may influence our results, particularly for correlative analyses. Another limitation to this study is that our analyses are limited to peripheral blood, and may not represent HIV reservoirs and immune cells located in tissues known to harbor latent HIV such as gut, spleen and brain.

Nevertheless, our study does have some strengths compared to previous work in this area, specifically a highly detailed flow panel, allowing us to quantify numerous immune subsets with a high degree of resolution. Also, the use of the intact proviral DNA assay allows us to separately examine the association of the immune system with intact and total viral DNA. It is however, important to note that IPDA does not measure all proviral sequences in PWH and is likely failing to identify some sequences due to proviral polymorphisms or dual deletions in the two IPDA amplicons. Also, a minor fraction of IPDA+ proviruses are actually defective due to the limited specificity of using two short amplicons to select for full-length intact proviruses (48, 65).

Overall, these findings suggest a complex concert of immune recovery, HIV reservoir dynamics, and intrinsic host factors (age, biological sex) that shapes host immunophenotype, even after years of ART. Mechanistic work will be needed to fully dissect the dynamic relationship between the immune system and the reservoir. Nevertheless, these findings support a model in which ongoing interaction between the HIV reservoir and the host immune cells continue to drive an association, albeit a relatively minor one, with persistent CD4 T cell activation during long-term ART. Notably, both intact and defective proviruses appear to contribute to this immune signature of HIV latency during long-term ART.

This study identifies several areas for future investigation. The question of whether the HIV reservoir directly drives persistent immune activation could be addressed directly by specifically suppressing viral transcription during therapy and examining the impact on the immune system. Future clinical trials with effective latency reversal and clearance regimens could include measurement of HIV-specific T cell responses, as this analysis suggests that lower levels may be associated with reduced intact HIV DNA clearance.

There is also growing appreciation for additional factors that correlate with the HIV reservoir. Specifically, the size of the reservoir has been linked to the abundance of specific bacteria species in the intestinal microbiome (66). Also, genetic studies have indicated the existence of single nucleotide polymorphisms (SNPs) that are associated with reservoir size, including SNPs in MX1 and IRF7 (67–69). These, along with future studies exploring the interplay of the host immune system and HIV reservoir during long-term might leverage new high-dimensional technologies such as scRNAseq and mass cytometry. The machine learning approaches described and applied in this study may be particularly useful for gaining biological insights into these high parameter datasets, particularly using analysis of global patterns in the data that may be otherwise imperceptible to humans.

Methods.

Cohort and sample collection.

115 people with HIV (PWH) were recruited from two clinical sites. 66 PWH were recruited at Duke University medical center and 49 PWH at UNC Chapel Hill. Peripheral blood was collected and peripheral blood mononuclear cells (PBMCs) were obtained by Ficoll separation, then frozen in 90% fetal bovine serum 10% dimethylsulfoxide (DMSO) (52, 59, 70, 71).

Flow cytometry.

Cell preparation and staining followed previously described methods (72). All antibodies were titrated to optimize signal-to-noise ratio on PBMCs prior to use, assuming a 50µL staining volume. Cell viability was examined using Zombie-NIR Fixable Viability Dye (0.4µL per 50µL staining volume; Biolegend). Antibodies used for surface staining were as follows: KLRG1-BV421 (SA231A2 clone, Biolegend,

CD45RA-PacBlue (H100 clone, BL), CD8-BV570 (RPA-T8 clone, Biolegend), CD127-BV605 (A019D5 clone, Biolegend), CD56-BV650 (5.1H11 clone, Biolegend), CCR7-BV711 (G043H7 clone, Biolegend), CD27-BV750 (O323 clone, Biolegend), PD1-VioBright515 (REA165 clone, Miltenyi), NKG2A-PE-Vio615 (REA110 clone, Miltenyi), CD16-PerCP-Cy5.5 (33G8 clone, Biolegend), CD38-PCPeF710HB7 clone, TF), CD14-SparkNIR685 (63D3 clone, Biolegend), CD19-SparkNIR685 (HIB19 clone, Biolegend), HLA-DR-APC-F750 (L243 clone, Biolegend). Antibodies used for intracellular staining were as follows: CD3-BV480 (UCHT1 clone, BD), CD4-PerCP (L200 clone, BD), IFN-g-PE-Cy7 (4S.B3 clone, Biolegend), IL-2-APC (MQ1-17H12 clone, Biolegend), TNFα-AF700 (Mab11 clone, Biolegend). Samples were analyzed using a Cytek Aurora spectral flow cytometer. Flow cytometry analysis was performed in FlowJo v10.8 software.

Intact Proviral DNA Assay (IPDA).

Cryopreserved samples of PBMCs from each study participant were viably thawed. A portion was used for immunophenotyping as described above, and the remainder were subjected to total CD4 T cell negative selection with the StemCell Technologies EasySep[TM] Human CD4+ T Cell Enrichment Kit (Cat# 19052). CD4 T cell DNA was extracted using the QIAamp DNA Mini Kit and quantified on a NanoDrop 1000 (Thermo Fisher Scientific). IPDA was performed as originally described (48), with a validated PCR annealing temperature modification to increase signal to noise ratio (59). Gating for positive droplets was set using negative (DNA elution buffer and HIV-seronegative CD4 T-cell DNA), and positive (Integrated DNA Technologies gblock amplicon) positive control wells processed in parallel. DNA shearing index values were similar to those reported previously (median, 0.33; interquartile range [IQR], 0.31–0.34) (59, 70, 71). A median of 1.04 x 106 (Q1 8.71 x 105, Q3 1.16 x x 106) cell equivalents were assessed for each donor.

Statistics.

Variable data analysis.

To create binarized labels that represent reservoir characteristics, we split the data at the median, which is 553/M for total reservoir size, 53/M for intact reservoir size, and 8.64 percent for percent intact. For every variable (immune cell frequency, demographics, and clinical information), we computed Spearman correlation and an Area Under the Curve (AUC) value. To compute the AUC value, we first created a Receiver Operating Characteristic (ROC) curve by plotting the true positive rate versus the true negative rate for every cell subset frequency. AUC is then computed as the area under the ROC curve using the trapezoidal rule.

To assess the importance of clinical, demographic information and HIV reservoir characteristics in

describing the immune markers, we performed a leave-one-covariate-out (LOCO) analysis. We trained a linear regression model (M_1) with intercept based on all variables. We removed each variable at a time and trained a new model without this variable. Then we computed the difference in R^2 score before and after dropping the variable. Variance Inflation Factors (VIFs) for age, sex, years of ART, CD4 nadir, recent CD4 account, and years of HIV infection prior to HIV infection were less than 2, indicating an acceptable level of correlation between these independent variables (73). However, higher VIF values were observed for intact and total HIV DNA, therefore separate LOCO inference analyses were conducted in order to avoid artificial fluctuations in model fit due to multicollinearity.

Data visualization.

We used PaCMAP (51) to reduce the dimensionality of the dataset to a 2-dimensional space. Due to the relatively small sample size of our dataset (n=115), we set the number of neighbors to 5. Additionally, we used GOSDT (53) which computes sparse optimal trees to identify patterns in the dataset. For all the visualization trees we set the depth budget parameter to 4 and the regularization parameter to 0.02.

Training and generalization.

Our training procedure for classification is described in **Algorithm 1** in Supplemental Materials, which is a standard machine learning pipeline with an added search for the number of variables. For the full hyperparameters list see **Table S8**. We normalized continuous variables to have values between 0 and 1. For logistics regression models, we set the regularization parameter of $\ell - 2$ regularization based on cross-validation. The values are 10 for total reservoir size and percent intact and 100 for intact reservoir size.

Study approval.

Written informed consent was obtained for all study participants. The study design was reviewed and approved by IRB for both Duke University and UNC Chapel Hill.

Data availability.

Deidentified datasets will be made publically available upon manuscript publication.

Author contributions.

Designed the study – EPB, DMM, CDR. Conducted experiments – SDF, ADVC. Analyzed the data – LS, YW, SDF. Wrote the manuscript: All authors.

Conflict of interest statement.

The authors have declared that no conflict of interest exists.

Acknowledgements.

This work was supported by the following grants from the National Institutes of Health: NIAID R01 AI143381 (EPB), NIAID UM1 AI164567 (DM Margolis), NIDA R61 DA047023 (EPB), NIDA R01 DA054994 (CDR), NIAID F30 AI145588 (SDF).

References

1. Finzi D, *et al.* Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy. *Science.* 1997;278(5341):1295–1300.

2. Chun TW, *et al.* Presence of an inducible HIV-1 latent reservoir during highly active antiretroviral therapy. *Proc Natl Acad Sci U S A.* 1997;94(24):13193–13197.

3. Chun TW, *et al.* Early establishment of a pool of latently infected, resting CD4(+) T cells during primary HIV-1 infection. *Proc Natl Acad Sci U S A.* 1998;95(15):8869–8873.

4. Wong JK, *et al.* Recovery of replication-competent HIV despite prolonged suppression of plasma viremia. *Science.* 1997;278(5341):1291–1295.

5. Ho YC, *et al.* Replication-competent noninduced proviruses in the latent reservoir increase barrier to HIV-1 cure. *Cell.* 2013;155(3):540–551.

6. Imamichi H, *et al.* Defective HIV-1 proviruses produce novel protein-coding RNA species in HIV-infected patients on combination antiretroviral therapy. *Proc Natl Acad Sci U S A.* 2016;113(31):8783–8788.

7. Bruner KM, *et al.* Defective proviruses rapidly accumulate during acute HIV-1 infection. *Nat Med.* 2016;22(9):1043–1049.

8. Maldarelli F, *et al.* HIV latency. Specific HIV integration sites are linked to clonal expansion and persistence of infected cells. *Science.* 2014;345(6193):179–183.

9. Wang Z, *et al.* Expanded cellular clones carrying replication-competent HIV-1 persist, wax, and wane. *Proc Natl Acad Sci U S A.* 2018;115(11):E2575-E2584.

10. Collora JA, *et al.* Single-cell multiomics reveals persistence of HIV-1 in expanded cytotoxic T cell clones. *Immunity.* 2022;55(6):1013–1031.

11. Whitney JB, *et al.* Rapid seeding of the viral reservoir prior to SIV viraemia in rhesus monkeys. *Nature.* 2014;512(7512):74–77.

12. Abrahams MR, *et al.* The replication-competent HIV-1 latent reservoir is primarily established near the time of therapy initiation. *Sci Transl Med.* 2019;11(513).

13. Martin GE, *et al.* Levels of Human Immunodeficiency Virus DNA Are Determined Before ART Initiation and Linked to CD8 T-Cell Activation and Memory Expansion. *J Infect Dis.* 2020;221(7):1135–1145.

14. Suzuki K, *et al.* HIV-1 viral blips are associated with repeated and increasingly high levels of cell-associated HIV-1 RNA transcriptional activity. *AIDS.* 2021;35(13):2095–2103.

15. Lian X, *et al.* Signatures of immune selection in intact and defective proviruses distinguish HIV-1 elite controllers. *Sci Transl Med.* 2021;13(624):eabl4097.

16. Hunt PW, *et al.* T cell activation is associated with lower CD4+ T cell gains in human immunodeficiency virus-infected patients with sustained viral suppression during antiretroviral therapy. *J Infect Dis.* 2003;187(10):1534–1543.

17. Bastard JP, *et al.* Circulating interleukin-6 levels correlate with residual HIV viraemia and markers of immune dysfunction in treatment-controlled HIV-infected patients. *Antivir Ther.* 2012;17(5):915–919.

18. Kottilil S, *et al.* Innate immune dysfunction in HIV infection: effect of HIV envelope-NK cell interactions. *J Immunol.* 2006;176(2):1107–1114.

19. Lichtfuss GF, *et al.* Biomarkers of immune dysfunction following combination antiretroviral therapy for HIV infection. *Biomark Med.* 2011;5(2):171–186.

20. Prescott LM. Loss of naive T cells may play major role in weakened immune response. *J Int Assoc Physicians AIDS Care.* 1995;1(2):32.

21. Day CL, *et al.* PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and disease progression. *Nature.* 2006;443(7109):350–354.

22. Ibegbu CC, *et al.* Expression of killer cell lectin-like receptor G1 on antigen-specific human CD8+ T lymphocytes during active, latent, and resolved infection and its relation with CD57. *J Immunol.* 2005;174(10):6088–6094.

23. Tavenier J, *et al.* Immunosenescence of the CD8(+) T cell compartment is associated with HIV-infection, but only weakly reflects age-related processes of adipose tissue, metabolism, and muscle in antiretroviral therapy-treated HIV-infected patients and controls. *BMC Immunol.* 2015;16:72.

24. Wang S, *et al.* An atlas of immune cell exhaustion in HIV-infected individuals revealed by single-cell transcriptomics. *Emerg Microbes Infect.* 2020;9(1):2333–2347.

25. Alter G, *et al.* Single-stranded RNA derived from HIV-1 serves as a potent activator of NK cells. *J Immunol.* 2007;178(12):7658–7666.

26. Horsburgh BA, *et al.* High levels of genetically intact HIV in HLA-DR+ memory T cells indicates their value for reservoir studies. *AIDS.* 2020;34(5):659–668.

27. Chomont N, *et al.* HIV reservoir size and persistence are driven by T cell survival and homeostatic proliferation. *Nat Med.* 2009;15(8):893–900.

28. Cockerham LR, *et al.* CD4+ and CD8+ T cell activation are associated with HIV DNA in resting CD4+ T cells. *PLoS One.* 2014;9(10):e110731.

29. Hatano H, *et al.* Cell-based measures of viral persistence are associated with immune activation and programmed cell death protein 1 (PD-1)-expressing CD4+ T cells. *J Infect Dis.* 2013;208(1):50–56.

30. Ruggiero A, *et al.* During Stably Suppressive Antiretroviral Therapy Integrated HIV-1 DNA Load in Peripheral Blood is Associated with the Frequency of CD8 Cells Expressing HLA-DR/DP/DQ. *EBioMedicine.* 2015;2(9):1153–1159.

31. Lee E, *et al.* Memory CD4 + T-cells expressing HLA-DR contribute to HIV persistence during pro-longed antiretroviral therapy. *Front. Microbiol..* 2019;10:2214.

32. Huang Y, *et al.* Immune activation and exhaustion marker expression on T-cell subsets in ART-treated adolescents and young adults with perinatal HIV-1 infection as correlates of viral persistence. *Front. Immunol..* 2023;14:1007626.

33. Banga R, *et al.* PD-1(+) and follicular helper T cells are responsible for persistent HIV-1 transcription in treated aviremic individuals. *Nat. Med..* 2016;22(7):754–761.

34. Fromentin R, *et al.* PD-1 blockade potentiates HIV latency reversal ex vivo in CD4+ T cells from ART-suppressed individuals. *Nat. Commun..* 2019;10(1):814.

35. Fromentin R, *et al.* CD4+ T cells expressing PD-1, TIGIT and LAG-3 contribute to HIV persistence during ART. *PLoS Pathog..* 2016;12(7):e1005761.

36. Pardons M, *et al.* Single-cell characterization and quantification of translation-competent viral reser-voirs in treated and untreated HIV infection. *PLoS Pathog..* 2019;15(2):e1007619.

37. Dufour C, *et al.* Phenotypic characterization of single CD4+ T cells harboring genetically intact and inducible HIV genomes. *Nat. Commun..* 2023;14(1):1115.

38. Zhu T, Cao W, Li T. HIV DNA positively correlates with HLA-DR+CD8+ T lymphocytes over 8-year suppressive antiretroviral therapy. *AIDS.* 2023;37(8):1335–1337.

39. Wu VH, *et al.* Profound phenotypic and epigenetic heterogeneity of the HIV-1-infected CD4+ T cell reservoir. *Nat. Immunol..* 2023;24(2):359–370.

40. Gálvez C, *et al.* Atlas of the HIV-1 reservoir in peripheral CD4 T cells of individuals on successful antiretroviral therapy. *MBio.* 2021;12(6):e0307821.

41. Bernal S, *et al.* Impact of obefazimod on viral persistence, inflammation, and immune activation in people with HIV on suppressive antiretroviral therapy. *J. Infect. Dis..* 2023.

42. Astorga-Gamaza A, *et al.* KLRG1 expression on natural killer cells is associated with HIV persistence, and its targeting promotes the reduction of the viral reservoir. *Cell Rep. Med..* 2023;4(10):101202.

43. Fisher K, *et al.* Unequal distribution of genetically-intact HIV-1 proviruses in cells expressing the immune checkpoint markers PD-1 and/or CTLA-4. *Front. Immunol..* 2023;14:1064346.

44. Dubé M, *et al.* Spontaneous HIV expression during suppressive ART is associated with the magnitude and function of HIV-specific CD4+ and CD8+ T cells. *Cell Host Microbe.* 2023;31(9):1507–1522.e5.

45. Takata H, *et al.* An active HIV reservoir during ART is associated with maintenance of HIV-specific CD8+ T cell magnitude and short-lived differentiation status. *Cell Host Microbe.* 2023;31(9):1494–1506.e4.

46. Siliciano JD, Siliciano RF. In vivo dynamics of the latent reservoir for HIV-1: New insights and implications for cure. *Annu. Rev. Pathol..* 2022;17(1):271–294.

47. Sponaugle A, *et al.* Dominant CD4+ T cell receptors remain stable throughout antiretroviral therapy-mediated immune restoration in people with HIV. *Cell Rep. Med..* 2023(101268):101268.

48. Bruner KM, *et al.* A quantitative approach for measuring the reservoir of latent HIV-1 proviruses. *Nature.* 2019;566(7742):120–125.

49. Thome JJC, *et al.* Spatial map of human T cell compartmentalization and maintenance over decades of life. *Cell.* 2014;159(4):814–828.

50. Lee KA, *et al.* Immune senescence, immunosenescence and aging. *Front. Aging.* 2022;3:900028.

51. Wang Y, *et al.* Understanding How Dimension Reduction Tools Work: An Empirical Approach to Deciphering t-SNE, UMAP, TriMap, and PaCMAP for Data Visualization. *Journal of Machine Learning Research.* 2021;22(201):1-73.

52. Falcinelli SD, *et al.* Impact of cannabis use on immune cell populations and the viral reservoir in people with HIV on suppressive antiretroviral therapy. *The Journal of Infectious Diseases.* 2023.

53. Lin J, *et al.* Generalized and scalable optimal sparse decision trees. in *International Conference on Machine Learning*:6150–6160PMLR 2020.

54. Poizot-Martin I, *et al.* Lack of correlation between the size of HIV proviral DNA reservoir and the level of immune activation in HIV-infected patients with a sustained undetectable HIV viral load for 10 years. *J Clin Virol.* 2013;57(4):351–355.

55. Gandhi RT, *et al.* Levels of HIV-1 persistence on antiretroviral therapy are not associated with markers of inflammation or activation. *PLoS Pathog.* 2017;13(4):e1006285.

56. Besson GJ, *et al.* HIV-1 DNA decay dynamics in blood during more than a decade of suppressive antiretroviral therapy. *Clin. Infect. Dis..* 2014;59(9):1312–1321.

57. Gandhi RT, *et al.* Selective Decay of Intact HIV-1 Proviral DNA on Antiretroviral Therapy. *J Infect Dis.* 2021;223(2):225–233.

58. Hsiao F, *et al.* Tissue memory CD4+ T cells expressing IL-7 receptor-alpha (CD127) preferentially support latent HIV-1 infection. *PLoS Pathog.* 2020;16(4):e1008450.

59. Falcinelli SD, *et al.* Longitudinal Dynamics of Intact HIV Proviral DNA and Outgrowth Virus Frequencies in a Cohort of Individuals Receiving Antiretroviral Therapy. *J Infect Dis.* 2021;224(1):92–100.

60. Peluso MJ, *et al.* Differential decay of intact and defective proviral DNA in HIV-1-infected individuals on suppressive antiretroviral therapy. *JCI Insight.* 2020;5(4).

61. Tumpach C, *et al.* Adaptation of droplet digital PCR-based HIV transcription profiling to digital PCR and association of HIV transcription and total or intact HIV DNA. *Viruses.* 2023;15(7).

62. Singh K, *et al.* Long-term persistence of transcriptionally active 'defective' HIV-1 proviruses: implications for persistent immune activation during antiretroviral therapy. *AIDS.* 2023;37(14):2119–2130.

63. Ishizaka A, *et al.* Short intracellular HIV-1 transcripts as biomarkers of residual immune activation in patients on antiretroviral therapy. *J. Virol..* 2016;90(12):5665–5676.

64. Martin HA, *et al.* New assay reveals vast excess of defective over intact HIV-1 transcripts in antiretroviral therapy-suppressed individuals. *J. Virol..* 2022;96(24):e0160522.

65. Gaebler C, *et al.* Sequence evaluation and comparative analysis of novel assays for intact proviral HIV-1 DNA. *J. Virol..* 2021;95(6).

66. Borgognone A, *et al.* Gut microbiome signatures linked to HIV-1 reservoir size and viremia control. *Microbiome.* 2022;10(1):59.

67. Siegel DA, *et al.* Host variation in type I interferon signaling genes (MX1), C-C chemokine receptor type 5 gene, and major histocompatibility complex class I alleles in treated HIV+ noncontrollers predict viral reservoir size. *AIDS.* 2023;37(3):477–488.

68. Thorball CW, *et al.* Host Genomics of the HIV-1 Reservoir Size and Its Decay Rate During Suppressive Antiretroviral Treatment. *J Acquir Immune Defic Syndr.* 2020;85(4):517–524.

69. Zhang Z, *et al.* IRF7 and RNH1 are modifying factors of HIV-1 reservoirs: a genome-wide association analysis. *BMC Med.* 2021;19(1):282.

70. Falcinelli SD, *et al.* Impact of biological sex on immune activation and frequency of the latent HIV reservoir during suppressive antiretroviral therapy. *J. Infect. Dis..* 2020;222(11):1843–1852.

71. Gay CL, *et al.* Stable latent HIV infection and low-level viremia despite treatment with the broadly neutralizing antibody VRC07-523LS and the latency reversal agent vorinostat. *J. Infect. Dis..* 2022;225(5):856–861.

72. Healy ZR, Murdoch DM. OMIP-036: Co-inhibitory receptor (immune checkpoint) expression analysis in human T cell subsets. *Cytometry A.* 2016;89(10):889–892.

73. James G, *et al. An introduction to statistical learning*;112. Springer 2013.

Tables and Figures.

	Percentage (count)	Median	[Q1, Q3]	[Min, Max]
Age		45	[37, 53]	[23,65]
Sex (% male)	76.52% (88)			
Race				
Black	60% (69)			
White	37.39% (43)			
Other	2.61% (3)			
Years of HIV		11	[7, 19.85]	[1, 33.6]
Years before ART< 1	55.65% (64)			
Years before ART≥ 1	38.26% (44)			
Years before ART=NA	6.09% (7)			
Years of ART		9	[5.23, 16.63]	[0.9,33.5]
Recent CD4 count		799	[624.5, 962]	[319, 1970]
CD4 Nadir		313.5	[163.25, 463.25]	[2, 1080]

Table 1: *Participant demographic and clinical characteristics.*
For demographics and clinical information, we report percentage for categorical variables, medians and [Q1, Q3] for real-value variables. ART is antiretroviral therapy. CD4 counts reported in cells/mm^3. Years of HIV has 1 missing value, Years of ART has 7, and CD4 Nadir has 3; consequently, these missing values are not included in median and quantiles computations. Years before ART means years of HIV infection before ART initiation.

Figure 1: *Duration of treatment and the HIV reservoir.*
Scatterplots for years of ART vs HIV reservoir size total reservoir size **(A)**, intact reservoir size **(B)** and percent intact **(C)** are shown. Each dot represents an individual study participant. Correlation coefficients and corresponding p-values are shown for each plot. Patients that have missing values of years of ART were not included in the plot. For percent intact, piece-wise linear function with two breaks is fitted. For total reservoir size the linear function is fitted.

Variable	Total	Intact	Percent intact
%CD8 T	**0.4052**	**0.3562**	0.0068
%CD38+/HLA-DR- CD4 T	**-0.3891**	-0.1098	0.2664
%KLRG1-/PD-1- CD4 T	**-0.3808**	-0.1289	0.2334
%Tn CD4 T	**-0.3802**	-0.2031	0.1714
%NKG2A+ CD4 T	**0.3618**	0.2904	0.0179
%PD-1-/CCR7+ CD4 T	**-0.3590**	-0.1082	0.2283
%CD4 T	**-0.3564**	**-0.3195**	-0.0079
%Tcm CD8 T	**0.3466**	0.1752	-0.1814
%CD38+ CD4 T	**-0.3366**	-0.0611	0.2829
%PD-1-/CCR7- CD4 T	**0.3300**	0.1824	-0.0938
%CD38+/HLA-DR- CD8 T	**-0.3267**	-0.0837	0.2636
%PD-1+ CD4 T	**0.3222**	0.0664	-0.2470
Age	**0.3172**	0.1669	-0.1471
CD4 Nadir	**-0.3164**	-0.1512	0.1927
%PD-1+ Tn CD4 T	**0.3119**	0.1246	-0.0962
%Tn CD8 T	**-0.3028**	-0.2697	0.0154
Years of ART	**0.3062**	-0.0688	**-0.4523**
%PD-1+/CCR7+ CD8 T	**0.2926**	0.1361	-0.1254
%CD38-/HLA-DR+ CD4 T	**0.2849**	0.0920	-0.1500
%KLRG1-/PD-1- CD8 T	**-0.2757**	-0.2182	0.0162
%PD-1+ Tn CD8 T	**0.2738**	0.1411	-0.0705
%CD38+/HLA-DR- Tn CD8 T	**-0.2676**	-0.0610	0.2119
%PD-1+/CCR7+ CD4 T	**0.2665**	0.0859	-0.1673
%KLRG1+/CD27+ CD8 T	**0.2606**	0.2212	0.0319
%CD27+ CD4 T	**-0.2602**	-0.0531	0.1767
%KLRG1-/CD27+ CD4 T	**-0.2575**	-0.0464	0.2171
%HLA-DR+ CD4 T	**0.2565**	0.1110	-0.0970
%PD-1+ CD8 T	**0.2541**	0.1969	0.0035
%CD38+ CD8 T	**-0.2418**	0.0109	**0.3114**
%CD38-/HLA-DR+ Tn CD8 T	**0.2404**	0.0738	-0.1346
%Tem CD4 T	**0.2402**	0.0888	-0.1669
%CD127+ CD4 T	-0.1298	**-0.3160**	-0.2539
%CD107-IFNγ-IL-2+TNFα+ CD4 T	0.0568	-0.1775	**-0.3223**
%CD107-IFNγ+IL-2+TNFα- CD4 T	0.0285	-0.2455	**-0.3265**

Table 2: *PWH features correlate with HIV reservoir characteristics.*
The abundance of 144 immune cell populations were determined by flow cytometry and the HIV reservoir was quantified by intact proviral DNA assay for a cohort of 115 people with HIV (PWH). Each abundance and clinical and demographic variable was correlated with total HIV reservoir size, intact reservoir size and the percentage of intact proviruses. Spearman correlation coefficients (**bold**) are shown for 36 variables that had significant p-values (< 0.05) after Benjamini–Hochberg correction for multiple comparisons. Each feature/subset is ranked by the absolute value of the correlation coefficient for the total reservoir size. For years of ART, we compute correlation based on 108 participants, excluding participants with missing years of ART values.

Figure 2: *Leave-one-covariate-out (LOCO) analysis for clinical-demographic features and reservoir characteristics while predicting immunophenotypes.*
A. Explanation of leave-one-covariate-out (LOCO) analysis based on example of %Tn CD8 T for clinical-demographic features and reservoir characteristics while predicting immunophenotypes. Analysis was performed for all 133 immunophenotypes considered in the study. The top 10 biggest drops in adjusted R^2 scores are reported for models that use total reservoir size (**B.**), intact reservoir size (**C.**), or percent intact (**D.**) as features in addition to clinical and demographic information. Participants with missing years of ART values are excluded from this analysis. The missing value of the CD4 nadir for one participant is imputed. In **Figure S6**, the LOCO analysis results are visualized for all 133 immunophenotypes.

Figure 3: *Receiver operating characteristic (ROC) curves identify PWH parameters that can classify reservoir characteristics.*
For total reservoir (**A**), intact reservoir (**B**), and percent intact (**C**), ROC curves are plotted for all 144 immune markers, demographics, and clinical variables (shown in gray). Axes represent the true positive rate (TPR) and the false positive rate (FPR) for each variable for classifying study participants into low (below median) versus high (above median) reservoir size. ROC curves for ten variables with the highest AUC values are shown in color for every HIV reservoir characteristic. Striped black lines represent the ROC curves of a random model. For years of ART ROC curves, we exclude participants with missing years of ART values.

Figure 4: *Dimension reduction reveals two major clusters of PWH with distinct immune systems and reservoirs.*
A. PaCMAP was applied to the data using the ten immune cell features with the highest AUC values for classifying participants based on total reservoir size, and two clusters (cluster 1 and cluster 2) are identified. **B**. Same as A but data points are color-coded by total reservoir size (high=purple, low=pink). Total reservoir size (**C.**), intact reservoir size (**D.**), and percent intact (**E.**) is shown for participants within each cluster. **F.** Key immune cell features that distinguish cluster 1 from cluster 2 are identified by visualizing the features with the highest AUC values with respect to classifying cohort participants based on cluster membership. Axes represent the true positive rate (TPR) and the false positive rate (FPR) for each variable. Immune markers and clinical-demographic features are shown for each cluster in Figure S8.

Figure 5: *Decision tree visualization of the association of immune cell subsets with reservoir characteristics.*

(A, C, E) Host variables (immune cell frequencies, demographic, and clinical information) were used to visualize the PWH dataset using the optimal sparse decision trees algorithm GOSDT. The overall set of PWH was classified as likely having high (above median, orange "leaves") or low (below median, blue "leaves") total reservoir size **(A)**, intact reservoir size **(C)**, and percent intact **(E)**. In each leaf, "med" denotes the median HIV characteristic of PWH, N is the number of PWH in the leaf, and MN is the number of misslabeled PWH. **(B, D, F)** PWH in model leaves associated with high (orange) or low (blue) reservoir size characteristics were aggregated and a Mann-Whitney U test was performed to determine statistical significance between the actual total reservoir size of the "high" and "low" groups for total reservoir size **(B)**, intact reservoir size **(D)**, and percent intact **(F)**. For the percent intact tree we exclude participants with missing values of years of ART. For total and intact reservoir size, missing values of years of ART were imputed, however, since the trees do not use this variable, imputations do not influence results. Visualization trees are explained with sets of rules in Supplemental Materials: **Figure S9** for total reservoir size, **Figure S10** for intact reservoir size, and **Figure S11** for percent intact.

Figure 6: *Predicting HIV reservoir characteristics with machine learning.*
Average training and test accuracies over ten training and test data splits for Random Forest (RF), Gradient Boosted Trees (GBT), Support Vector Machines with RBF kernel (SVM), Logistic Regression (LR), and CART models for total reservoir size (**A**), intact reservoir size (**C**), and percent intact (**E**) are shown. For one split of training and test sets, logistic regression models are visualized for total reservoir (**B**), intact reservoir (**D**), and percent intact (**F**). On the y-axis we show variables used by the model, while the x-axis displays coefficient values for individual variables used by models. For percent intact models we exclude participants with missing values of years of ART. For total and intact reservoir size, missing values of years of ART were imputed. The missing value of the CD4 nadir for one participant was imputed as well using the MICE algorithm.

Machine learning approaches identify immunologic signatures of total and intact HIV DNA during long-term antiretroviral therapy.

Supplementary Methods.

CD4/CD8 ratio regression analysis

To examine the connection between CD4/CD8 and (CD127+ CD4)/CD8 ratio and HIV characteristics (**Figure S5**), we first computed the natural logarithm (log$_e$) of the intact, total and percent intact as the outcome and normalized outcome of each immune feature (meaning that our values of ratio and outcome are between 0 and 1). We removed outliers using the DBSCAN algorithm (1), which is an unsupervised clustering algorithm that groups data points based on density into a single cluster. For DBSCAN, we set the parameter that determines radius of a circle around each point (that is used to compute density) as 0.15 and minimum number of samples in each cluster as 10. Then we fitted the remaining datapoints using Linear Regression.

Regression machine learning models to predict reservoir size from immune cell frequencies.

To predict reservoir size based on immune cell frequencies, clinical and demographics information (**Figure S12**), similar to classification, we sought to find the set of model parameters and types of models that would be able to fit the dataset without overfitting. We used several approaches: Linear Regression (LR), Ridge Regression (RR), Kernel Regression with RBF kernel (KR), Decision Tree Regressor (DT), Random Forest (RF), and Gradient Boosted Trees (GBT). We followed the procedure described in Algorithm 1, optimized the R^2 score, returned the mean and standard deviation of the R^2 score of training and test sets, and chose features based on the absolute value of the correlation coefficient. We normalized variables to have values between 0 and 1. We then fitted the models to the normalized natural logarithm (log$_e$) of total, intact reservoir size and percent intact and then further normalization makes outcomes to be between 0 and 1. For total and intact reservoir size we visualized the Ridge Regression model, which performed the best based on mean test R^2 score (**Figure S12 D, E, G, H**). We set the regularization parameter to 1, as this value was best on average according to cross-validation for total and intact reservoir size. For percent intact, we visualized Linear Regression based on 4 variables (**Figure S12 F, I**) as it achieved the second highest R^2 score after Kernel Ridge based on 25 variables.

References

1. Ester M, *et al.* A density-based algorithm for discovering clusters in large spatial databases with noise. in *2nd International Conference on Knowledge Discovery and Data Mining (KDD-96)*;96:226–231 1996.

Figure S1: *Representative flow cytometry gating is shown for one sample from the 115-person cohort.*

Group	Size	Intact reservoir	Total reservoir	% intact
Male	88	63.5 [18.75 146.25]	681 [198 1433.75]	8.73 [4.9 20.58]
Female	27	41 [19, 92.5]	489 [317.5, 637]	8.64 [4.86, 15.56]
p-value		0.4521	0.1937	0.9451
Black	69	43 [19, 113]	502 [181, 1246]	9.77 [4.64, 16.4]
White	43	69 [26 191.5]	605 [289.5 1415.5]	8 [5.31 20.81]
Other	3	67 [41, 520]	1055 [720, 1165,]	6.39 [5.20, 41.37]
p-value		0.5079	0.4916	0.9914

Table S1: *Intact, Total reservoir size and % intact for demographic subgroups.*
We report medians and [Q1, Q3] for total reservoir size, intact reservoir size, and percent intact. Mann-Whitney U test was used to compute p-values for gender and Kruskal-Wallis H-test for race.

Immune marker	Med	[Q1, Q3]	Immune marker	Med	[Q1, Q3]
%CD4 T	52.90	[43.85, 59.60]	%CD27+ CD4 T	86.20	[79.15, 90.10]
%CD38+ CD4 T	46.60	[39.65, 54.30]	%CD127+ CD4 T	97.10	[95.20, 98.30]
%HLA-DR+ CD4 T	12.30	[9.76, 16.40]	%KLRG1+ CD4 T	17.20	[10.85, 26.25]
%NKG2A+ CD4 T	0.10	[0.06, 0.20]	%PD-1+ CD4 T	45.40	[39.70, 54.25]
%Tcm CD4 T	49.90	[42.30, 56.65]	%Tn CD4 T	30.40	[22.10, 39.90]
%Teff CD4 T	1.73	[0.89, 3.50]	%Tem CD4 T	15.70	[10.95, 19.70]
%CD38-/HLA-DR+ CD4 T	8.16	[6.22, 11.00]	%CD38+/HLA-DR+ CD4 T	4.37	[3.16, 5.94]
%CD38+/HLA-DR- CD4 T	40.20	[32.90, 47.75]	%KLRG1-/CD27+ CD4 T	74.40	[67.30, 83.05]
%KLRG1+/CD27+ CD4 T	6.70	[3.67, 11.65]	%KLRG1+/CD27- CD4 T	7.68	[4.54, 13.40]
%KLRG1-/CD27- CD4 T	6.01	[4.43, 8.04]	%KLRG1-/PD-1+ CD4 T	32.80	[27.00, 37.95]
%KLRG1+/PD-1+ CD4 T	12.60	[7.46, 17.30]	%KLRG1+/PD-1- CD4 T	4.79	[3.12, 8.63]
%KLRG1-/PD-1- CD4 T	49.30	[38.35, 55.90]	%PD-1-/CCR7+ CD4 T	50.90	[40.50, 56.35]
%PD-1+/CCR7+ CD4 T	29.70	[25.00, 35.05]	%PD-1+/CCR7- CD4 T	14.60	[10.40, 18.90]
%PD-1-/CCR7- CD4 T	3.77	[2.56, 5.25]	%CD8 T	40.30	[33.35, 47.45]
%CD27+ CD8 T	65.40	[54.05, 75.30]	%CD38+ CD8 T	35.70	[26.05, 45.45]
%CD127+ CD8 T	99.20	[98.40, 99.60]	%HLA-DR+ CD8 T	25.70	[18.75, 36.05]
%KLRG1+ CD8 T	62.10	[47.40, 73.40]	%PD-1+ CD8 T	37.50	[26.10, 47.50]
%Tn CD8 T	30.40	[21.60, 41.50]	%Teff CD8 T	26.90	[20.35, 36.70]
%Tem CD8 T	19.00	[13.00, 27.15]	%CD38-/HLA-DR+ CD8 T	16.10	[10.55, 22.75]
%CD38+/HLA-DR+ CD8 T	8.43	[5.86, 13.45]	%CD38+/HLA-DR- CD8 T	19.60	[14.45, 31.15]
%KLRG1-/CD27+ CD8 T	35.00	[23.45, 48.80]	%KLRG1+/CD27+ CD8 T	25.90	[19.35, 33.45]
%KLRG1+/CD27- CD8 T	30.30	[21.45, 43.40]	%KLRG1-/CD27- CD8 T	2.42	[1.58, 4.57]
%KLRG1-/PD-1+ CD8 T	7.40	[5.17, 10.95]	%KLRG1+/PD-1+ CD8 T	27.80	[18.35, 36.75]
%KLRG1+/PD-1- CD8 T	30.20	[23.35, 38.25]	%Tcm CD8 T	14.60	[10.30, 22.00]
%KLRG1-/PD-1- CD8 T	27.70	[17.75, 43.70]	%PD-1-/CCR7+ CD8 T	30.70	[22.45, 42.95]
%PD-1+/CCR7+ CD8 T	13.30	[10.10, 16.70]	%PD-1+/CCR7- CD8 T	22.00	[14.85, 27.50]
%PD-1-/CCR7- CD8 T	28.30	[18.20, 36.40]	%NK	6.37	[3.98, 10.50]
%NKG2A+ NK	31.60	[21.40, 49.85]	%NKG2A-/CD16+ NK	52.20	[34.25, 64.90]
%NKG2A+/CD16+ NK	16.50	[10.21, 24.35]	%NKG2A+/CD16- NK	13.10	[8.26, 23.75]
%NKG2A-/CD16- NK	13.80	[9.26, 19.15]	%NKT	4.33	[2.33, 6.56]
%NKG2A+ NKT	9.31	[4.79, 19.70]	%NKG2A-/CD16+ NKT	6.31	[1.66, 12.95]
%NKG2A+/CD16+ NKT	0.43	[0.14, 1.23]	%NKG2A+/CD16- NKT	8.48	[4.57, 17.65]
%NKG2A-/CD16- NKT	77.40	[65.85, 89.35]	%CD38-/HLA-DR+ Tcm CD4 T	9.93	[7.52, 12.25]
%CD38+/HLA-DR+ Tcm CD4 T	4.71	[3.13, 6.30]	%CD38+/HLA-DR- Tcm CD4 T	22.90	[17.00, 28.45]
%CD38-/HLA-DR- Tcm CD4 T	61.80	[55.20, 66.75]	%PD-1+ Tcm CD4 T	58.90	[52.55, 65.80]
%CD38-/HLA-DR+ Tn CD4 T	0.66	[0.30, 1.12]	%CD38+/HLA-DR+ Tn CD4 T	7.29	[5.21, 9.38]
%CD38+/HLA-DR- Tn CD4 T	81.30	[76.20, 87.55]	%CD38-/HLA-DR- Tn CD4 T	8.52	[5.80, 14.20]
%PD-1+ Tn CD4 T	7.26	[4.40, 12.65]	%CD38-/HLA-DR+ Teff CD4 T	17.50	[11.20, 25.75]
%CD38+/HLA-DR+ Teff CD4 T	19.00	[10.75, 28.15]	%CD38+/HLA-DR- Teff CD4 T	16.00	[11.25, 27.15]
%CD38-/HLA-DR- Teff CD4 T	38.50	[27.40, 50.50]	%PD-1+ Teff CD4 T	56.40	[39.30, 72.25]
%CD38-/HLA-DR+ Tem CD4 T	27.40	[22.45, 33.80]	%CD38+/HLA-DR+ Tem CD4 T	7.31	[5.31, 10.10]
%CD38+/HLA-DR- Tem CD4 T	4.77	[3.49, 6.96]	%CD38-/HLA-DR- Tem CD4 T	57.00	[50.75, 65.80]
%PD-1+ Tem CD4 T	82.20	[76.00, 86.55]	%CD38-/HLA-DR+ Tcm CD8 T	22.10	[16.70, 28.20]
%CD38+/HLA-DR+ Tcm CD8 T	7.11	[4.23, 12.10]	%CD38+/HLA-DR- Tcm CD8 T	5.28	[3.71, 7.18]
%CD38-/HLA-DR- Tcm CD8 T	62.70	[51.65, 72.10]	%PD-1+ Tcm CD8 T	63.60	[53.80, 71.10]
%CD38-/HLA-DR+ Tn CD8 T	6.02	[3.29, 9.82]	%CD38+/HLA-DR+ Tn CD8 T	3.68	[2.42, 5.36]
%CD38+/HLA-DR- Tn CD8 T	45.40	[31.70, 57.25]	%CD38-/HLA-DR- Tn CD8 T	43.00	[31.75, 54.45]
%PD-1+ Tn CD8 T	15.00	[9.96, 24.35]	%CD38-/HLA-DR+ Teff CD8 T	27.80	[21.55, 37.25]
%CD38+/HLA-DR+ Teff CD8 T	16.60	[11.35, 25.95]	%CD38-/HLA-DR- Teff CD8 T	12.40	[7.18, 18.45]
%CD38-/HLA-DR- Teff CD8 T	39.20	[26.30, 47.20]	%PD-1+ Teff CD8 T	33.90	[24.10, 41.85]
%CD38-/HLA-DR+ Tem CD8 T	29.70	[21.55, 36.05]	%CD38+/HLA-DR+ Tem CD8 T	13.80	[9.31, 21.10]
%CD38+/HLA-DR- Tem CD8 T	5.31	[3.27, 9.88]	%CD38-/HLA-DR- Tem CD8 T	46.30	[35.30, 56.80]
%PD-1+ Tem CD8 T	62.40	[49.70, 71.80]	%IL-2+ CD8 T	0.00	[-0.03, 0.03]
%IL-2+ CD4 T	0.02	[0.01, 0.04]	%TNFα+ CD4 T	0.04	[0.01, 0.06]
%TNFα+ CD8 T	0.04	[0.01, 0.12]	%IFNγ+ CD4 T	0.02	[0.00, 0.04]
%IFNγ+ CD8 T	0.05	[0.01, 0.21]	%CD107+ CD4 T	0.02	[0.01, 0.04]
%CD107+ CD8 T	0.23	[0.12, 0.55]	%CD107+IFNγ+IL-2-TNFα- CD4 T	0.00	[0.00, 0.00]
%CD107+IFNγ-IL-2+TNF- CD4 T	0.00	[0.00, 0.00]	%CD107+IFNγ-IL-2-TNFα+ CD4 T	0.00	[0.00, 0.00]
%CD107+IFNγ-IL-2-TNFα- CD4 T	0.01	[0.01, 0.02]	%CD107-IFNγ+IL-2+TNFα- CD4 T	0.00	[0.00, 0.00]
%CD107-IFNγ+IL-2-TNFα+ CD4 T	0.00	[-0.01, 0.01]	%CD107-IFNγ+IL-2-TNFα- CD4 T	0.00	[-0.01, 0.00]
%CD107-IFNγ-IL-2+TNFα+ CD4 T	0.01	[0.01, 0.02]	%CD107-IFNγ-IL-2+TNFα- CD4 T	0.00	[-0.01, 0.00]
%CD107-IFNγ-IL-2-TNFα+ CD4 T	0.01	[0.00, 0.02]	%CD107+IFNγ+IL-2-TNFα- CD8T	0.03	[0.01, 0.07]
%CD107+IFNγ-IL-2+TNF- CD8 T	0.00	[0.00, 0.00]	%CD107+IFNγ-IL-2-TNFα+ CD8 T	0.01	[0.00, 0.03]
%CD107+IFNγ-IL-2-TNFα- CD8 T	0.14	[0.08, 0.30]	%CD107-IFNγ+IL-2+TNFα- CD8 T	0.00	[0.00, 0.00]
%CD107-IFNγ+IL-2-TNFα+ CD8 T	0.00	[0.00, 0.01]	%CD107-IFNγ+IL-2-TNFα- CD8 T	0.00	[-0.01, 0.01]
%CD107-IFNγ-IL-2+TNFα+ CD8 T	0.00	[0.00, 0.00]	%CD107-IFNγ-IL-2+TNFα- CD8 T	-0.01	[-0.04, 0.00]
%CD107-IFNγ-IL-2-TNFα+ CD8 T	0.00	[0.00, 0.00]			

Table S2: *Immune subsets characteristics*. We report medians, 25% and 75% percentiles for every immune variable used in the study.

Variable	Total	Intact	% intact	Variable	Total	Intact	% intact
%CD8 T	**0.4052**	**0.3562**	0.0068	%CD38+/HLA-DR- Tcm CD8 T	-0.2169	-0.0182	0.1970
%CD38+/HLA-DR- CD4 T	**-0.3891**	-0.1098	0.2664	%CD107-IFNγ+IL-2-TNFα- CD4 T	-0.2154	-0.1633	-0.0599
%KLRG1-/PD-1- CD4 T	**-0.3808**	-0.1289	0.2334	%KLRG1-/CD27+ CD8 T	-0.2112	-0.1894	-0.0021
%Tn CD4 T	**-0.3802**	-0.2031	0.1714	%KLRG1+/PD-1+ CD8 T	0.2089	0.1922	0.0473
%NKG2A+ CD4 T	**0.3618**	0.2904	0.0179	%CD38+/HLA-DR- Tn CD4 T	-0.2066	0.0140	0.1909
%PD-1-/CCR7+ CD4 T	**-0.3590**	-0.1082	0.2283	%PD-1+/CCR7- CD4 T	0.1971	0.0410	-0.1338
%CD4 T	**-0.3564**	**-0.3195**	-0.0079	%CD38-/HLA-DR+ CD8 T	0.1947	0.0572	-0.0860
%Tcm CD8 T	**0.3466**	0.1752	-0.1814	%KLRG1+ CD8 T	0.1887	0.1709	0.0048
%CD38+ CD4 T	**-0.3366**	-0.0611	0.2829	%KLRG1-/PD-1+ CD4 T	0.1877	0.0818	-0.0963
%PD-1-/CCR7- CD4 T	**0.3300**	0.1824	-0.0938	%NKG2A-/CD16+ NK	0.1855	0.0433	-0.1222
%CD38+/HLA-DR- CD8 T	**-0.3267**	-0.0837	0.2636	%KLRG1+/PD-1- CD4 T	0.1849	0.0616	-0.1388
%PD-1+ CD4 T	**0.3222**	0.0664	-0.2470	%CD38+/HLA-DR+ Tem CD4 T	0.1798	0.2668	0.1961
Age	**0.3172**	0.1669	-0.1471	%KLRG1+ CD4 T	0.1771	0.0002	-0.2034
CD4 Nadir	**-0.3164**	-0.1512	0.1927	%KLRG1+/PD-1+ CD4 T	0.1650	-0.0392	-0.2355
%PD-1+ Tn CD4 T	**0.3119**	0.1246	-0.0962	Years before ART < 1	-0.1645	0.0527	0.2513
%Tn CD8 T	**-0.3028**	-0.2697	0.0154	% NKT	0.1616	-0.0223	-0.1942
Years of ART	**0.3062**	-0.0688	**-0.4523**	%CD38-/HLA-DR+ Tn CD4 T	0.1612	-0.0675	-0.2160
%PD-1+/CCR7+ CD8 T	**0.2926**	0.1361	-0.1254	%PD-1+/CCR7- CD8 T	0.1352	0.1899	0.1057
%CD38-/HLA-DR+ CD4 T	**0.2849**	0.0920	-0.1500	%CD38-/HLA-DR- Tn CD8 T	0.1349	-0.0452	-0.2308
%KLRG1-/PD-1- CD8 T	**-0.2757**	-0.2182	0.0162	%CD38-/HLA-DR- Tcm CD8 T	0.1302	-0.0048	-0.2000
%PD-1+ Tn CD8 T	**0.2738**	0.1411	-0.0705	Recent CD4 count	-0.1302	-0.1944	-0.0970
%CD38+/HLA-DR- Tn CD8 T	**-0.2676**	-0.0610	0.2119	%CD127+ CD4 T	-0.1298	**-0.3160**	-0.2539
%PD-1+/CCR7+ CD4 T	**0.2665**	0.0859	-0.1673	Years before ART = NA	0.1183	-0.0312	-0.1950
%KLRG1+/CD27+ CD8 T	**0.2606**	0.2212	0.0319	%CD127+ CD8 T	0.1136	-0.1217	-0.2458
%CD27+ CD4 T	**-0.2602**	-0.0531	0.1767	%CD38+/HLA-DR+ Tcm CD8 T	-0.0920	0.0701	0.2340
%KLRG1-/CD27+ CD4 T	**-0.2575**	-0.0464	0.2171	%CD107-IFNγ-IL-2+TNFα+ CD8 T	0.0909	-0.0499	-0.2270
%HLA-DR+ CD4 T	**0.2565**	0.1110	-0.0970	%CD38+/HLA-DR- Teff CD8 T	-0.0833	0.1328	0.2462
%PD-1+ CD8 T	**0.2541**	0.1969	0.0035	%CD38+/HLA-DR- Tem CD8 T	-0.0660	0.1177	0.1958
%CD38+ CD8 T	**-0.2418**	0.0109	**0.3114**	%CD107-IFNγ-IL-2+TNFα+ CD4 T	0.0568	-0.1775	**-0.3223**
%CD38-/HLA-DR+ Tn CD8 T	**0.2404**	0.0738	-0.1346	%IL-2+ CD4 T	-0.0515	-0.1936	-0.1868
%Tem CD4 T	**0.2402**	0.0888	-0.1669	%CD38+/HLA-DR+ Tem CD8 T	0.0510	0.1694	0.2107
%PD-1-/CCR7+ CD8 T	-0.2321	-0.2173	-0.0237	%CD38+/HLA-DR+ CD8 T	0.0461	0.1660	0.1865
%CD38-/HLA-DR- Tem CD4 T	-0.2302	-0.2365	-0.1349	%CD107-IFNγ+IL-2+TNFα- CD4 T	0.0285	-0.2455	**-0.3265**
%KLRG1+/CD27- CD4 T	0.2271	0.0207	-0.1935	%CD38+/HLA-DR- Tem CD4 T	-0.0134	0.1551	0.2013
%Tcm CD4 T	0.2262	0.1745	-0.0492				

Table S3: *Host features correlate with HIV reservoir characteristics.*
The abundance of 119 immune cell populations was determined by flow cytometry and the HIV reservoir was quantified by intact proviral DNA assay for a cohort of 115 people with HIV (PWH). Each abundance and clinical and demographic variable was correlated with total HIV reservoir size, intact reservoir size, and the percentage of intact proviruses. Spearman correlation coefficients are shown for 69 variables that were correlated (p<0.05) with one or more reservoir characteristics. Each feature/subset is ranked by the absolute value of the correlation coefficient for the total reservoir size. Variables that had significant p-values (<0.05) after Benjamini–Hochberg correction for multiple comparisons (144 comparisons in our case) with one of the characteristics are shown in bold. For years of ART, correlations are computed based on 108 participants, excluding participants with missing years of ART values. For CD4 Nadir, correlations are computed based on 114 patients, excluding participants with the missing CD4 Nadir value.

Figure S2: *Abundance of immune cell subsets correlates with HIV reservoir. Continuation in Figures S3, S4.*
Scatterplots for selected examples of immune cell subsets versus HIV reservoir size or percent intact are shown. Additionally, scatterplots of immune cell subsets versus time on therapy are displayed. Each dot represents an individual study participant. Correlation coefficients and corresponding p-values are shown for each plot. For time plots, participants that have missing values of years of ART were not included in the plot. The piece-wise linear function with one break is fitted if a break is between 1 and 20 years, otherwise, a linear function is fitted.

Figure S3: *Abundance of immune cell subsets correlates with HIV reservoir. Continuation in Figures S2, S4.*
Scatterplots for selected examples of immune cell subsets versus HIV reservoir size or percent intact are shown. Additionally, scatterplots of immune cell subsets versus time on therapy are displayed. Each dot represents an individual study participant. Correlation coefficients and corresponding p-values are shown for each plot. For time plots, participants that have missing values of years of ART were not included in the plot. The piece-wise linear function with one break is fitted if a break is between 1 and 20 years, otherwise, a linear function is fitted.

Figure S4: *Abundance of immune cell subsets correlates with HIV reservoir. Continuation in Figures S2, S3.*
Scatterplots for selected examples of immune cell subsets versus HIV reservoir size or percent intact are shown. Additionally, scatterplots of immune cell subsets versus time on therapy are displayed. Each dot represents an individual study participant. Correlation coefficients and corresponding p-values are shown for each plot. For time plots, participants that have missing values of years of ART were not included in the plot. The piece-wise linear function with one break is fitted if a break is between 1 and 20 years, otherwise, a linear function is fitted.

Figure S5: *CD4/CD8 and (%CD127+ CD4T)/CD8 ratios correlate with total and intact reservoir size.*
Normalized (values transformed to be between 0 and 1) CD4/CD8 (**A**) and (%CD127+ CD4T)/CD8 (**B**)
ratios are shown on the x-axis and the normalized natural logarithm (log$_e$) of the total reservoir, intact
reservoir and percent intact on the y-axis. Spearman correlation was computed between ratios and HIV
reservoir characteristics. Outliers (red data points) were removed with the DBSCAN clustering algorithm
and a linear regression model was fitted (black line) to the remaining data points. Spearman correlations,
R^2 scores and mean squared error after outlier removal are displayed.

Variable	Spearman correlation coefficient, All participants	Spearman correlation coefficient, Participants with years of ART< 10 years
Age	**0.4550**	0.2629
%CD38+/HLA-DR- CD8 T	**-0.4132**	-0.3788
%CD38+/HLA-DR- Tn CD8 T	**-0.4035**	-0.3075
%CD38+ CD8 T	**-0.3770**	-0.3025
%Tcm CD8 T	**0.3684**	0.1595
%CD38-/HLA-DR- Tn CD8 T	**0.3580**	0.2222
%CD38+/HLA-DR- Tcm CD8 T	**-0.3138**	-0.2666
CD4 Nadir	**-0.3078**	0.1265
%CD107+IFNγ-IL-2+TNF- CD8 T	**0.2903**	0.1728

Table S4: *PWH immune features correlate with years of ART.*
The abundance of 119 immune cell populations was determined by flow cytometry and the HIV reservoir was quantified by intact proviral DNA assay for a cohort of 115 people with HIV (PWH). Each abundance and clinical and demographic variable was correlated with years of therapy for all participants (middle column)and for participants who had been on therapy for less than 10 years (right column). Spearman correlation coefficients are shown for all variables that were correlated (p$<$0.05) after Benjamini-Hochberg correction with years of treatment for all participants or for participants that were on ART for less than 10 years. Variables that had significant p-value ($<$0.05) after correction for multiple comparisons are shown in bold. Correlation is computed based on 108 PWH for all participants and on 60 PWH for PWH who were on therapy for less than 10 years). Computations exclude PWH with missing years of ART values. For CD4 Nadir, correlations are computed excluding the participant with missing CD4 Nadir value.

Total model		Intact model		% intact model	
Variable	**VIF value**	**Variable**	**VIF value**	**Variable**	**VIF value**
Total	1.2489	Intact	1.0505	% intact	1.3273
Age	1.4585	Age	1.4534	Age	1.4605
Sex	1.1335	Sex	1.1071	Sex	1.1165
Years of ART	1.5010	Years of ART	1.4474	Years of ART	1.6326
nadir CD4	1.3141	nadir CD4	1.2810	nadir CD4	1.3753
Recent CD4 count	1.2992	Recent CD4 count	1.3108	Recent CD4 count	1.2885
Years before ART	1.1296	Years before ART	1.1049	Years before ART	1.1565

Table S5: *Multicolinearity analysis for variables used in models to predict immunophenotypes.*
Variance Inflation Factor (VIF) for every variable is computed for models that fit immunophenotypes as a target and are based on clinical, demographic information and HIV characteristics (total, intact reservoir size or percent intact) as features. Note that here "Years before ART" is a continuous variable that we used for VIF computations only. In our other analysis we categorize this variable and use "Years before ART=NA", "Years before ART< 1", and "Years before ART≥ 1" instead.

Figure S6: *Leave-one-covariate-out (LOCO) analysis for clinical-demographic features and reservoir characteristics while predicting immunophenotypes.*

Analysis that is described in **Figure 2** was performed for all 133 immunophenotypes considered in the study. Drops in adjusted R^2 scores are reported for models that use total (**A.**), intact reservoir size (**B.**), or percent intact (**C.**) as features in addition to clinical and demographic information such as age, biological sex, years of treatment, CD4 nadir, recent CD4 count, and years of HIV before treatment ($=$NA, < 1, ≥ 1). On the x-axis, we show features that were dropped from the model. On the y-axis, we display immunophenotypes, which are targets (outcomes) for the linear regression models. In **Tables S6-S8** we show the actual values of drops in adjusted R^2 score.

Figure S7: *Coefficient visualization for linear regression models that predict immunophenotypes based on clinical, demographic information and HIV characteristics.* **(A., B., C.)** LOCO analysis from **Figure 3** for total-based (**A.**), intact-based (**B.**) and percent intact-based (**C.**) models. The drops in adjusted R^2 scores are shown after removing a feature and training a new model without it. **(D., E., F.)** Coefficient visualization for models that include clinical and demographic information such as age, biological sex, years of treatment, CD4 nadir, recent CD4 count, and years of HIV before treatment = NA, years of HIV before treatment < 1, years of HIV before treatment ≥ 1 and total reservoirs size (**D.**) or intact reservoir size (**E.**), or percent intact (**F.**). No features are dropped from these models, they are "Include all" models from tables **Tables S6-S8**. On the x-axis, the feature is shown, and on the y-axis the target (immunophenotypes from (A., B., C.)). The heatmap displays the coefficient in front of that variable in the model (if the model is %CD4 T= β_1Total + β_2Age + β_3Sex + ..., then $\beta_1, \beta_2, \beta_3, ...$ are visualized), where positive coefficients are shown in red and negative in blue.

	Adj. R^2	$\triangle R^2$, difference in adjustable R^2 after dropping a covariate						
Model Immunophenotype	Include all	Total	Age	Sex	Years of ART	CD4 nadir	Recent CD4 count	Years before ART
%CD4 T	0.2736	0.0468	0.0134	0.0492	-0.0001	0.0089	0.1290	-0.0175
%CD27+ CD4 T	0.0380	0.0110	-0.0087	-0.0016	-0.0010	-0.0055	0.0142	0.0000
%CD38+ CD4 T	0.0737	0.0026	0.0101	0.0038	-0.0089	0.0352	0.0092	-0.0239
%CD127+ CD4 T	-0.0026	-0.0048	0.0282	-0.0081	-0.0098	0.0008	0.0100	-0.0292
%HLA-DR+ CD4 T	0.1701	0.0889	-0.0029	0.0138	-0.0070	-0.0028	0.0133	-0.0092
%KLRG1+ CD4 T	0.0485	-0.0016	0.0190	-0.0090	-0.0083	0.0003	0.0135	0.0109
%NKG2A+ CD4 T	0.1179	0.0844	0.0607	-0.0064	-0.0080	-0.0006	-0.0058	-0.0181
%PD-1+ CD4 T	0.1156	0.0355	-0.0075	0.0042	-0.0030	0.0045	0.0113	-0.0262
%Tcm CD4 T	0.0377	-0.0097	0.0440	-0.0046	-0.0086	-0.0019	-0.0090	-0.0083
%Tn CD4 T	0.1139	0.0061	0.0317	0.0059	-0.0070	-0.0065	0.0237	-0.0263
%Teff CD4 T	-0.0409	0.0017	-0.0099	-0.0088	-0.0097	-0.0014	-0.0082	-0.0155
%Tem CD4 T	0.0480	0.0082	-0.0096	-0.0052	-0.0085	-0.0096	0.0388	-0.0100
%CD38-/HLA-DR+ CD4 T	0.1339	0.0629	-0.0087	0.0217	0.0019	0.0101	-0.0080	0.0136
%CD38+/HLA-DR+ CD4 T	0.1334	0.0680	0.0074	-0.0021	-0.0067	-0.0086	0.0470	-0.0256
%CD38+/HLA-DR- CD4 T	0.1220	0.0318	0.0009	0.0125	-0.0072	0.0308	-0.0061	-0.0218
%KLRG1-/CD27+ CD4 T	0.0619	-0.0089	-0.0002	-0.0070	-0.0059	0.0032	0.0261	-0.0007
%KLRG1+/CD27+ CD4 T	0.0321	0.0064	0.0186	-0.0087	-0.0092	0.0388	-0.0028	-0.0270
%KLRG1+/CD27- CD4 T	0.0242	-0.0095	-0.0065	-0.0039	-0.0039	-0.0021	0.0054	0.0261
%KLRG1-/CD27- CD4 T	0.1745	0.1169	0.0496	-0.0049	-0.0034	-0.0062	0.0063	-0.0080
%KLRG1-/PD-1+ CD4 T	0.1058	0.1003	-0.0078	-0.0085	-0.0090	-0.0037	-0.0056	0.0044
%KLRG1+/PD-1+ CD4 T	0.0367	-0.0002	-0.0018	0.0062	-0.0009	-0.0049	0.0052	0.0179
%KLRG1+/PD-1- CD4 T	0.0358	-0.0073	0.0366	0.0027	-0.0075	0.0015	0.0098	-0.0161
%KLRG1-/PD-1- CD4 T	0.1655	0.0168	0.0129	-0.0067	-0.0065	0.0164	0.0315	-0.0199
%PD-1-/CCR7+ CD4 T	0.1199	0.0433	-0.0055	-0.0020	-0.0058	-0.0022	0.0214	-0.0246
%PD-1+/CCR7+ CD4 T	0.0887	0.0168	0.0027	-0.0074	-0.0048	0.0201	-0.0086	-0.0096
%PD-1+/CCR7- CD4 T	0.0388	0.0078	-0.0064	0.0039	-0.0081	-0.0096	0.0207	-0.0064
%PD-1-/CCR7- CD4 T	-0.0002	0.0114	-0.0018	-0.0051	-0.0077	-0.0041	0.0196	-0.0184
%CD8 T	0.2684	0.0620	-0.0067	0.0436	-0.0058	0.0081	0.1129	-0.0166
%CD27+ CD8 T	0.0422	0.0001	0.0378	0.0058	-0.0083	-0.0096	0.0263	-0.0132
%CD38+ CD8 T	0.2734	-0.0062	0.0978	0.0130	-0.0061	0.0360	0.0749	-0.0009
%CD127+ CD8 T	-0.0299	-0.0073	-0.0071	-0.0073	-0.0062	0.0067	0.0050	-0.0233
%HLA-DR+ CD8 T	0.0243	0.0044	0.0073	-0.0098	-0.0098	-0.0060	0.0323	-0.0205
%KLRG1+ CD8 T	0.1687	-0.0079	0.1416	0.0240	-0.0059	-0.0059	0.0062	-0.0198
%PD-1+ CD8 T	0.1710	0.0247	0.0736	0.0218	-0.0084	-0.0084	0.0031	-0.0246
%Tn CD8 T	0.2632	-0.0062	0.1920	0.0185	-0.0063	-0.0073	0.0187	-0.0125
%Teff CD8 T	-0.0257	-0.0096	0.0010	0.0112	0.0084	-0.0103	-0.0078	-0.0217
%Tem CD8 T	0.0213	-0.0090	0.0309	-0.0077	-0.0086	-0.0093	0.0378	-0.0290
%CD38-/HLA-DR+ CD8 T	0.0331	-0.0049	0.0293	-0.0082	-0.0098	-0.0096	-0.0040	0.0017
%CD38+/HLA-DR+ CD8 T	0.0296	0.0060	-0.0096	-0.0074	-0.0097	0.0037	0.0588	-0.0282
%CD38+/HLA-DR- CD8 T	0.3169	-0.0049	0.1369	0.0338	-0.0047	0.0193	0.0159	0.0020
%KLRG1-/CD27+ CD8 T	0.1670	-0.0082	0.1265	0.0265	-0.0049	-0.0057	0.0140	-0.0188
%KLRG1+/CD27+ CD8 T	0.1028	0.0045	0.0804	0.0073	-0.0069	0.0018	-0.0081	-0.0242
%KLRG1+/CD27- CD8 T	0.0470	0.0015	0.0533	0.0063	-0.0089	-0.0094	0.0171	-0.0145
%KLRG1-/CD27- CD8 T	-0.0028	-0.0084	0.0158	-0.0101	-0.0077	-0.0101	0.0131	-0.0291
%KLRG1-/PD-1+ CD8 T	-0.0150	0.0455	-0.0102	-0.0090	-0.0094	-0.0101	-0.0089	-0.0300
%KLRG1+/PD-1+ CD8 T	0.1595	0.0029	0.0870	0.0328	-0.0084	-0.0084	0.0090	-0.0250
%KLRG1+/PD-1- CD8 T	0.0067	0.0080	0.0526	-0.0066	-0.0060	-0.0039	-0.0081	-0.0200
%Tcm CD8 T	0.2447	0.0105	0.1236	-0.0076	0.0033	-0.0074	-0.0045	-0.0221
%KLRG1-/PD-1- CD8 T	0.2288	-0.0061	0.1628	0.0252	-0.0057	-0.0045	0.0061	-0.0178
%PD-1-/CCR7+ CD8 T	0.1778	-0.0047	0.1121	0.0379	-0.0013	-0.0080	0.0121	-0.0170
%PD-1+/CCR7+ CD8 T	0.1880	0.0498	0.0535	0.0050	-0.0072	-0.0002	0.0109	-0.0241

%PD-1+/CCR7- CD8 T	0.0975	-0.0030	0.0368	0.0150	-0.0088	-0.0050	0.0485	-0.0268
%PD-1-/CCR7- CD8 T	-0.0254	0.0025	0.0114	-0.0014	0.0021	-0.0097	-0.0053	-0.0173
% NK	0.1177	-0.0074	-0.0082	0.1481	-0.0064	0.0031	-0.0088	-0.0238
%NKG2A+ NK	0.1098	0.0031	-0.0038	0.1538	-0.0080	-0.0083	0.0152	-0.0260
%NKG2A-/CD16+ NK	0.0988	0.0283	-0.0057	0.1088	-0.0082	-0.0022	0.0301	-0.0260
%NKG2A+/CD16+ NK	0.0279	-0.0087	0.0148	0.0549	-0.0097	-0.0091	-0.0098	-0.0244
%NKG2A+/CD16- NK	0.1178	0.0103	-0.0081	0.1412	-0.0055	-0.0047	0.0472	-0.0255
%NKG2A-/CD16- NK	0.0096	0.0301	-0.0094	-0.0017	-0.0100	0.0069	0.0027	-0.0292
% NKT	-0.0626	0.0056	-0.0107	-0.0072	-0.0107	-0.0101	-0.0075	-0.0281
%NKG2A+ NKT	-0.0114	-0.0044	-0.0058	-0.0038	0.0221	-0.0097	-0.0100	-0.0297
%NKG2A-/CD16+ NKT	0.0330	0.0675	0.0011	0.0084	-0.0078	-0.0066	-0.0064	-0.0262
%NKG2A+/CD16+ NKT	0.0376	0.0126	-0.0064	0.0308	0.0122	-0.0091	0.0180	0.0009
%NKG2A+/CD16- NKT	-0.0187	-0.0071	-0.0065	-0.0075	0.0177	-0.0095	-0.0086	-0.0302
%NKG2A-/CD16- NKT	0.0727	0.0556	0.0068	0.0160	0.0233	-0.0090	-0.0085	-0.0272
%CD38-/HLA-DR+ Tcm CD4 T	-0.0016	0.0137	-0.0070	0.0125	0.0021	0.0015	-0.0099	-0.0226
%CD38+/HLA-DR+ Tcm CD4 T	0.0870	-0.0012	-0.0027	0.0020	-0.0044	-0.0014	0.0908	-0.0271
%CD38+/HLA-DR- Tcm CD4 T	0.0316	0.0129	-0.0075	-0.0098	-0.0098	0.0380	0.0497	-0.0284
%CD38-/HLA-DR- Tcm CD4 T	0.0806	-0.0036	-0.0092	-0.0032	-0.0089	0.0321	0.0910	-0.0271
%PD-1+ Tcm CD4 T	0.0049	0.0269	-0.0066	-0.0089	-0.0076	0.0040	-0.0064	-0.0293
%CD38-/HLA-DR+ Tn CD4 T	-0.0197	0.0203	-0.0052	0.0052	-0.0053	-0.0051	-0.0061	-0.0221
%CD38+/HLA-DR+ Tn CD4 T	-0.0368	-0.0054	-0.0063	-0.0093	-0.0085	-0.0104	0.0244	-0.0308
%CD38+/HLA-DR- Tn CD4 T	0.0344	0.0239	-0.0040	0.0226	-0.0082	0.0071	0.0320	-0.0258
%CD38-/HLA-DR- Tn CD4 T	0.0720	0.0123	-0.0074	0.0316	-0.0083	0.0109	0.0876	-0.0249
%PD-1+ Tn CD4 T	0.0716	0.1019	0.0038	-0.0009	-0.0086	0.0034	-0.0028	-0.0270
%CD38-/HLA-DR+ Teff CD4 T	-0.0065	-0.0067	-0.0059	-0.0085	-0.0102	0.0007	0.0249	-0.0108
%CD38+/HLA-DR+ Teff CD4 T	0.1348	-0.0048	0.0205	-0.0087	-0.0018	0.0066	0.0932	-0.0257
%CD38+/HLA-DR- Teff CD4 T	0.0156	-0.0099	-0.0010	-0.0085	-0.0090	-0.0085	0.0174	0.0028
%CD38-/HLA-DR- Teff CD4 T	0.0877	-0.0009	0.0141	-0.0058	-0.0076	0.0085	0.0461	-0.0257
%PD-1+ Teff CD4 T	0.0268	-0.0035	-0.0092	0.0498	0.0067	-0.0081	0.0084	-0.0249
%CD38-/HLA-DR+ Tem CD4 T	0.0114	0.0169	-0.0100	-0.0056	0.0088	0.0071	0.0179	0.0067
%CD38+/HLA-DR+ Tem CD4 T	0.1559	0.1513	-0.0022	-0.0064	-0.0085	-0.0065	0.0146	-0.0189
%CD38+/HLA-DR- Tem CD4 T	-0.0035	-0.0087	-0.0066	-0.0083	-0.0100	0.0356	0.0185	0.0023
%CD38-/HLA-DR- Tem CD4 T	0.0725	0.1179	-0.0086	-0.0060	0.0002	-0.0093	-0.0086	-0.0268
%PD-1+ Tem CD4 T	-0.0838	-0.0107	-0.0064	-0.0088	-0.0086	-0.0109	-0.0109	-0.0322
%CD38-/HLA-DR+ Tcm CD8 T	-0.0646	-0.0104	-0.0040	-0.0093	-0.0094	-0.0102	-0.0065	-0.0312
%CD38+/HLA-DR+ Tcm CD8 T	0.0716	0.0128	0.0132	-0.0025	-0.0084	0.0097	0.0452	-0.0191
%CD38+/HLA-DR- Tcm CD8 T	0.1146	-0.0048	0.0351	0.0031	-0.0089	0.0492	0.0305	-0.0240
%CD38-/HLA-DR- Tcm CD8 T	0.0990	0.0020	0.0311	-0.0054	-0.0091	0.0169	0.0519	-0.0228
%PD-1+ Tcm CD8 T	-0.0074	-0.0054	-0.0101	0.0092	-0.0083	-0.0097	0.0143	-0.0298
%CD38-/HLA-DR+ Tn CD8 T	0.0757	0.0441	0.0202	-0.0054	-0.0075	-0.0016	-0.0086	-0.0263
%CD38+/HLA-DR+ Tn CD8 T	0.0107	0.0204	-0.0025	-0.0086	-0.0038	-0.0003	0.0269	-0.0289
%CD38+/HLA-DR- Tn CD8 T	0.2479	0.0085	0.0579	0.0522	-0.0054	0.0212	0.0615	-0.0204
%CD38-/HLA-DR- Tn CD8 T	0.2043	-0.0079	0.0452	0.0408	-0.0063	0.0228	0.1133	-0.0220
%PD-1+ Tn CD8 T	0.0969	0.0493	0.0285	0.0173	-0.0075	-0.0071	-0.0007	-0.0268
%CD38-/HLA-DR+ Teff CD8 T	0.0310	-0.0098	-0.0094	0.0063	0.0167	-0.0074	-0.0096	0.0070
%CD38+/HLA-DR+ Teff CD8 T	-0.0086	-0.0049	-0.0090	-0.0013	-0.0102	0.0020	0.0498	-0.0242
%CD38+/HLA-DR- Teff CD8 T	0.0839	-0.0010	-0.0070	-0.0093	0.0088	0.0193	0.0295	0.0118
%CD38-/HLA-DR- Teff CD8 T	0.0358	0.0043	-0.0080	0.0203	-0.0078	0.0148	0.0838	-0.0274
%PD-1+ Teff CD8 T	-0.0511	-0.0075	-0.0074	-0.0100	-0.0105	-0.0084	0.0097	-0.0258
%CD38-/HLA-DR+ Tem CD8 T	0.0108	-0.0096	-0.0054	0.0059	-0.0005	-0.0098	-0.0094	0.0091
%CD38+/HLA-DR+ Tem CD8 T	0.0070	0.0186	-0.0011	-0.0100	-0.0077	-0.0039	0.0327	-0.0291
%CD38+/HLA-DR- Tem CD8 T	0.0319	-0.0062	-0.0097	-0.0093	0.0051	0.0000	0.0249	-0.0006
%CD38-/HLA-DR- Tem CD8 T	-0.0057	0.0094	-0.0095	-0.0024	-0.0064	0.0024	0.0370	-0.0258
%PD-1+ Tem CD8 T	-0.0133	0.0220	-0.0102	0.0102	-0.0101	-0.0064	-0.0086	-0.0295
%IL-2+ CD8 T	-0.0439	0.0087	-0.0098	-0.0063	0.0132	-0.0104	-0.0104	-0.0308
%IL-2+ CD4 T	0.0483	0.0491	-0.0086	0.0088	0.0147	-0.0001	-0.0094	-0.0186

%TNFα+ CD4 T	-0.0031	0.0052	-0.0076	0.0128	0.0288	-0.0097	0.0019	-0.0167
%TNFα+ CD8 T	-0.0252	0.0112	-0.0053	-0.0069	0.0160	-0.0040	0.0107	-0.0279
%IFNγ+ CD4 T	-0.0513	-0.0015	-0.0027	-0.0023	-0.0070	-0.0106	-0.0078	-0.0262
%IFNγ+ CD8 T	-0.0600	0.0007	-0.0107	-0.0092	-0.0023	-0.0098	0.0054	-0.0280
%CD107+ CD4 T	-0.0542	-0.0099	-0.0012	-0.0105	-0.0024	-0.0105	0.0059	-0.0254
%CD107+ CD8 T	-0.0545	0.0004	-0.0087	-0.0054	0.0002	-0.0085	0.0083	-0.0313
%CD107+IFNγ+IL-2-TNFα- CD4 T	-0.0576	-0.0102	-0.0006	-0.0101	-0.0088	-0.0103	-0.0015	-0.0265
%CD107+IFNγ-IL-2+TNF- CD4 T	-0.0490	-0.0099	-0.0076	-0.0106	-0.0101	-0.0074	0.0022	-0.0153
%CD107+IFNγ-IL-2-TNFα+ CD4 T	-0.0466	-0.0093	-0.0103	-0.0086	0.0083	0.0106	-0.0040	-0.0173
%CD107+IFNγ-IL-2-TNFα- CD4 T	-0.0137	-0.0050	0.0086	-0.0075	0.0088	-0.0091	0.0318	-0.0301
%CD107-IFNγ+IL-2+TNFα- CD4 T	-0.0040	0.0168	-0.0056	-0.0025	-0.0100	-0.0035	0.0298	-0.0261
%CD107-IFNγ+IL-2+TNFα+ CD4 T	-0.0234	-0.0103	0.0026	0.0378	-0.0054	-0.0033	-0.0015	-0.0299
%CD107-IFNγ+IL-2-TNFα- CD4 T	0.0006	-0.0021	-0.0039	0.0131	-0.0065	-0.0097	0.0029	-0.0284
%CD107-IFNγ-IL-2+TNFα+ CD4 T	0.2042	0.0364	-0.0048	0.0333	0.1008	0.0116	0.0002	-0.0139
%CD107-IFNγ-IL-2+TNFα- CD4 T	0.0720	0.0747	-0.0027	0.0070	-0.0076	0.0093	-0.0050	-0.0231
%CD107-IFNγ-IL-2-TNFα+ CD4 T	-0.0021	-0.0081	-0.0092	0.0042	0.0499	-0.0029	-0.0009	-0.0131
%CD107+IFNγ+IL-2-TNFα- CD8T	-0.0685	-0.0066	-0.0097	-0.0100	-0.0100	-0.0108	0.0024	-0.0294
%CD107+IFNγ-IL-2+TNF- CD8 T	-0.0142	-0.0096	0.0038	-0.0099	0.0149	-0.0030	-0.0096	-0.0274
%CD107+IFNγ-IL-2-TNFα+ CD8 T	0.0041	0.0087	0.0157	0.0009	0.0023	0.0117	0.0044	-0.0275
%CD107+IFNγ-IL-2-TNFα- CD8 T	-0.0068	-0.0030	-0.0018	0.0046	0.0056	-0.0084	0.0104	-0.0138
%CD107-IFNγ+IL-2+TNFα- CD8 T	-0.0072	0.0281	0.0011	-0.0055	0.0104	-0.0102	-0.0033	-0.0259
%CD107-IFNγ+IL-2-TNFα+ CD8 T	0.0455	0.0175	0.0032	-0.0091	0.0554	-0.0058	0.0280	-0.0277
%CD107-IFNγ+IL-2-TNFα- CD8 T	-0.0587	0.0025	-0.0064	-0.0088	-0.0040	-0.0080	-0.0035	-0.0312
%CD107-IFNγ-IL-2+TNFα+ CD8 T	0.0349	-0.0097	-0.0085	0.0080	0.0832	0.0008	0.0209	-0.0232
%CD107-IFNγ-IL-2+TNFα- CD8 T	-0.0553	-0.0028	-0.0048	-0.0085	-0.0102	-0.0073	-0.0048	-0.0233
%CD107-IFNγ-IL-2-TNFα+ CD8 T	0.0152	0.0000	-0.0036	-0.0099	0.0142	-0.0099	-0.0099	-0.0230

Table S6: *Adjusted R^2 scores and differences in adjusted R^2 for LOCO analysis for the model that contains total reservoir size.*
The first column contains all immunophenotypes and is a target for the regression model. The second column contains adjusted R^2 of the linear regression model with features such as total reservoir size, age, biological sex, years of treatment, CD4 nadir, recent CD4 count, and years of HIV before treatment (=NA, < 1, ≥ 1). The next seven columns show differences in adjusted R^2 score (ΔR^2) after removing a specific feature. Participants with missing years of ART values are excluded from this analysis. The missing value of the CD4 nadir for one participant is imputed.

Immunophenotype \ Model	Adj. R^2 Include all	ΔR^2, difference in adjustable R^2 after dropping a covariate						
		Intact	Age	Sex	Years of ART	CD4 nadir	Recent CD4 count	Years before ART
%CD4 T	0.2554	0.0286	0.0096	0.0385	0.0067	0.0223	0.1265	-0.0124
%CD27+ CD4 T	0.0188	-0.0082	-0.0096	0.0024	0.0492	-0.0082	0.0174	0.0053
%CD38+ CD4 T	0.0631	-0.0081	0.0157	0.0087	0.0114	0.0428	0.0039	-0.0232
%CD127+ CD4 T	0.0792	0.0770	0.0227	-0.0076	-0.0071	-0.0026	0.0013	-0.0247
%HLA-DR+ CD4 T	0.1372	0.0560	-0.0056	0.0254	0.0664	0.0094	0.0123	0.0038
%KLRG1+ CD4 T	0.0626	0.0125	0.0200	-0.0091	-0.0040	-0.0030	0.0174	0.0044
%NKG2A+ CD4 T	0.0996	0.0661	0.0529	-0.0023	-0.0090	0.0137	-0.0066	-0.0077
%PD-1+ CD4 T	0.0790	-0.0011	-0.0059	0.0113	0.0031	0.0143	0.0143	-0.0260
%Tcm CD4 T	0.0377	-0.0097	0.0446	-0.0045	-0.0071	-0.0016	-0.0090	-0.0083
%Tn CD4 T	0.1002	-0.0076	0.0375	0.0105	-0.0024	-0.0041	0.0267	-0.0266
%Teff CD4 T	-0.0523	-0.0097	-0.0105	-0.0073	0.0613	-0.0045	-0.0074	-0.0126
%Tem CD4 T	0.0321	-0.0077	-0.0097	-0.0025	-0.0034	-0.0093	0.0425	-0.0060
%CD38-/HLA-DR+ CD4 T	0.0863	0.0153	-0.0090	0.0345	0.1054	0.0257	-0.0079	0.0279
%CD38+/HLA-DR+ CD4 T	0.1421	0.0767	0.0051	0.0032	0.0107	-0.0069	0.0410	-0.0220
%CD38+/HLA-DR- CD4 T	0.0848	-0.0054	0.0055	0.0216	0.0129	0.0461	-0.0079	-0.0193
%KLRG1-/CD27+ CD4 T	0.0652	-0.0056	0.0018	-0.0062	0.0044	0.0038	0.0308	-0.0012
%KLRG1+/CD27+ CD4 T	0.0329	0.0072	0.0172	-0.0077	0.0014	0.0293	-0.0020	-0.0283
%KLRG1+/CD27- CD4 T	0.0293	-0.0044	-0.0052	-0.0026	0.0294	-0.0018	0.0091	0.0247
%KLRG1-/CD27- CD4 T	0.1403	0.0827	0.0405	0.0008	0.0212	0.0044	0.0049	-0.0200
%KLRG1-/PD-1+ CD4 T	0.0671	0.0616	-0.0092	-0.0059	-0.0069	0.0092	-0.0061	-0.0117
%KLRG1+/PD-1+ CD4 T	0.0603	0.0233	-0.0008	0.0055	-0.0016	-0.0072	0.0097	0.0098
%KLRG1+/PD-1- CD4 T	0.0368	-0.0063	0.0365	0.0040	-0.0023	-0.0003	0.0105	-0.0179
%KLRG1-/PD-1- CD4 T	0.1424	-0.0062	0.0183	-0.0045	-0.0011	0.0259	0.0359	-0.0177
%PD-1-/CCR7+ CD4 T	0.0762	-0.0004	-0.0028	0.0037	0.0088	0.0056	0.0256	-0.0229
%PD-1+/CCR7+ CD4 T	0.0692	-0.0027	0.0060	-0.0053	-0.0007	0.0311	-0.0084	-0.0152
%PD-1+/CCR7- CD4 T	0.0232	-0.0078	-0.0080	0.0085	-0.0060	-0.0097	0.0236	-0.0019
%PD-1-/CCR7- CD4 T	-0.0200	-0.0083	0.0013	-0.0077	0.1314	-0.0074	0.0232	-0.0151
%CD8 T	0.2344	0.0280	-0.0077	0.0314	-0.0002	0.0227	0.1134	-0.0108
%CD27+ CD8 T	0.0367	-0.0054	0.0350	0.0033	0.0270	-0.0089	0.0252	-0.0165
%CD38+ CD8 T	0.2875	0.0080	0.1036	0.0140	0.0350	0.0328	0.0655	-0.0037
%CD127+ CD8 T	0.0920	0.1146	-0.0081	-0.0071	-0.0068	0.0027	-0.0038	-0.0135
%HLA-DR+ CD8 T	0.0447	0.0248	0.0074	-0.0095	0.2107	-0.0083	0.0262	-0.0148
%KLRG1+ CD8 T	0.1682	-0.0084	0.1403	0.0233	0.0582	-0.0061	0.0052	-0.0200
%PD-1+ CD8 T	0.1654	0.0191	0.0798	0.0294	-0.0079	-0.0072	0.0021	-0.0242
%Tn CD8 T	0.2650	-0.0044	0.1937	0.0199	0.0383	-0.0074	0.0171	-0.0110
%Teff CD8 T	-0.0253	-0.0092	0.0010	0.0110	0.1180	-0.0103	-0.0077	-0.0226
%Tem CD8 T	0.0212	-0.0092	0.0287	-0.0082	0.0077	-0.0091	0.0339	-0.0291
%CD38-/HLA-DR+ CD8 T	0.0333	-0.0047	0.0308	-0.0087	0.3354	-0.0090	-0.0045	0.0060
%CD38+/HLA-DR+ CD8 T	0.0689	0.0454	-0.0093	-0.0063	0.0033	-0.0017	0.0476	-0.0276
%CD38+/HLA-DR- CD8 T	0.3152	-0.0066	0.1417	0.0371	0.0833	0.0225	0.0148	0.0037
%KLRG1-/CD27+ CD8 T	0.1668	-0.0084	0.1259	0.0263	0.0623	-0.0059	0.0130	-0.0189
%KLRG1+/CD27+ CD8 T	0.0995	0.0012	0.0851	0.0110	0.0077	0.0072	-0.0081	-0.0258
%KLRG1+/CD27- CD8 T	0.0397	-0.0057	0.0496	0.0034	0.0376	-0.0086	0.0157	-0.0178
%KLRG1-/CD27- CD8 T	-0.0046	-0.0101	0.0141	-0.0100	-0.0059	-0.0100	0.0146	-0.0290
%KLRG1-/PD-1+ CD8 T	-0.0334	0.0271	-0.0100	-0.0103	-0.0016	-0.0070	-0.0090	-0.0301
%KLRG1+/PD-1+ CD8 T	0.1596	0.0030	0.0908	0.0382	-0.0078	-0.0083	0.0078	-0.0246
%KLRG1+/PD-1- CD8 T	-0.0024	-0.0010	0.0484	-0.0082	0.1103	-0.0071	-0.0084	-0.0236
%Tcm CD8 T	0.2393	0.0051	0.1306	-0.0075	0.0086	-0.0060	-0.0045	-0.0211
%KLRG1-/PD-1- CD8 T	0.2306	-0.0042	0.1645	0.0270	0.0661	-0.0033	0.0050	-0.0165
%PD-1-/CCR7+ CD8 T	0.1797	-0.0028	0.1141	0.0412	0.0532	-0.0073	0.0107	-0.0150
%PD-1+/CCR7+ CD8 T	0.1674	0.0292	0.0615	0.0119	0.0043	0.0101	0.0107	-0.0238

%PD-1+/CCR7- CD8 T	0.0996	-0.0009	0.0382	0.0178	-0.0090	-0.0067	0.0457	-0.0265
%PD-1-/CCR7- CD8 T	-0.0320	-0.0040	0.0092	-0.0036	0.1052	-0.0104	-0.0057	-0.0209
% NK	0.1168	-0.0083	-0.0081	0.1551	0.0143	0.0050	-0.0089	-0.0244
%NKG2A+ NK	0.1078	0.0012	-0.0046	0.1667	-0.0084	-0.0068	0.0157	-0.0265
%NKG2A-/CD16+ NK	0.0934	0.0229	-0.0068	0.1255	-0.0088	0.0056	0.0314	-0.0269
%NKG2A+/CD16+ NK	0.0269	-0.0098	0.0134	0.0590	-0.0072	-0.0093	-0.0098	-0.0248
%NKG2A+/CD16- NK	0.1241	0.0165	-0.0078	0.1544	-0.0070	-0.0001	0.0506	-0.0235
%NKG2A-/CD16- NK	0.0047	0.0252	-0.0088	-0.0049	-0.0049	0.0190	0.0037	-0.0293
% NKT	-0.0729	-0.0047	-0.0108	-0.0089	0.0939	-0.0108	-0.0072	-0.0261
%NKG2A+ NKT	-0.0172	-0.0102	-0.0070	-0.0018	0.0327	-0.0102	-0.0098	-0.0295
%NKG2A-/CD16+ NKT	0.0782	0.1127	0.0006	0.0163	-0.0081	0.0038	-0.0030	-0.0274
%NKG2A+/CD16+ NKT	0.0364	0.0113	-0.0071	0.0380	0.0077	-0.0067	0.0192	0.0099
%NKG2A+/CD16- NKT	-0.0217	-0.0102	-0.0075	-0.0064	0.0280	-0.0099	-0.0078	-0.0303
%NKG2A-/CD16- NKT	0.0696	0.0526	0.0042	0.0254	0.0334	-0.0045	-0.0081	-0.0271
%CD38-/HLA-DR+ Tcm CD4 T	-0.0205	-0.0052	-0.0086	0.0192	0.0696	0.0082	-0.0103	-0.0195
%CD38+/HLA-DR+ Tcm CD4 T	0.1029	0.0148	-0.0021	0.0036	0.0155	-0.0043	0.0816	-0.0260
%CD38+/HLA-DR- Tcm CD4 T	0.0100	-0.0086	-0.0091	-0.0094	0.0074	0.0505	0.0427	-0.0281
%CD38-/HLA-DR- Tcm CD4 T	0.0756	-0.0087	-0.0093	-0.0050	-0.0079	0.0374	0.0815	-0.0274
%PD-1+ Tcm CD4 T	-0.0213	0.0007	-0.0085	-0.0102	0.0004	0.0134	-0.0057	-0.0286
%CD38-/HLA-DR+ Tn CD4 T	-0.0474	-0.0074	-0.0077	0.0118	-0.0060	-0.0001	-0.0079	-0.0189
%CD38+/HLA-DR+ Tn CD4 T	-0.0399	-0.0085	-0.0079	-0.0101	0.0504	-0.0105	0.0303	-0.0309
%CD38+/HLA-DR- Tn CD4 T	0.0091	-0.0015	-0.0064	0.0321	0.0154	0.0168	0.0280	-0.0235
%CD38-/HLA-DR- Tn CD4 T	0.0639	0.0042	-0.0082	0.0393	-0.0055	0.0200	0.0867	-0.0224
%PD-1+ Tn CD4 T	0.0080	0.0383	-0.0018	0.0075	-0.0004	0.0206	-0.0045	-0.0290
%CD38-/HLA-DR+ Teff CD4 T	-0.0100	-0.0102	-0.0047	-0.0077	0.0565	0.0029	0.0215	-0.0095
%CD38+/HLA-DR+ Teff CD4 T	0.1612	0.0217	0.0239	-0.0084	-0.0012	0.0117	0.0800	-0.0246
%CD38+/HLA-DR- Teff CD4 T	0.0158	-0.0097	-0.0014	-0.0083	0.0059	-0.0085	0.0183	0.0030
%CD38-/HLA-DR- Teff CD4 T	0.0930	0.0044	0.0140	-0.0047	-0.0030	0.0144	0.0423	-0.0265
%PD-1+ Teff CD4 T	0.0257	-0.0046	-0.0094	0.0470	0.0441	-0.0065	0.0078	-0.0231
%CD38-/HLA-DR+ Tem CD4 T	0.0118	0.0172	-0.0100	-0.0031	0.1069	0.0171	0.0196	0.0177
%CD38+/HLA-DR+ Tem CD4 T	0.2064	0.2019	-0.0034	-0.0010	0.0019	-0.0068	0.0084	-0.0235
%CD38+/HLA-DR- Tem CD4 T	-0.0038	-0.0089	-0.0055	-0.0089	0.0036	0.0347	0.0219	0.0020
%CD38-/HLA-DR- Tem CD4 T	0.0797	0.1251	-0.0091	-0.0005	0.0572	-0.0053	-0.0091	-0.0198
%PD-1+ Tem CD4 T	-0.0818	-0.0087	-0.0054	-0.0092	0.0617	-0.0109	-0.0109	-0.0321
%CD38-/HLA-DR+ Tcm CD8 T	-0.0571	-0.0028	-0.0029	-0.0094	0.2539	-0.0104	-0.0079	-0.0306
%CD38+/HLA-DR+ Tcm CD8 T	0.1217	0.0629	0.0154	-0.0004	-0.0060	0.0022	0.0343	-0.0226
%CD38+/HLA-DR- Tcm CD8 T	0.1203	0.0009	0.0354	0.0045	0.0623	0.0576	0.0334	-0.0224
%CD38-/HLA-DR- Tcm CD8 T	0.1396	0.0426	0.0348	-0.0045	0.0433	0.0101	0.0408	-0.0242
%PD-1+ Tcm CD8 T	-0.0013	0.0007	-0.0101	0.0111	0.0058	-0.0101	0.0117	-0.0297
%CD38-/HLA-DR+ Tn CD8 T	0.0290	-0.0026	0.0287	-0.0009	0.2810	0.0067	-0.0080	-0.0252
%CD38+/HLA-DR+ Tn CD8 T	0.0510	0.0608	-0.0018	-0.0071	0.0047	-0.0058	0.0194	-0.0280
%CD38+/HLA-DR- Tn CD8 T	0.2360	-0.0034	0.0644	0.0615	0.0139	0.0300	0.0578	-0.0191
%CD38-/HLA-DR- Tn CD8 T	0.2065	-0.0057	0.0471	0.0422	-0.0021	0.0221	0.1077	-0.0222
%PD-1+ Tn CD8 T	0.0641	0.0165	0.0359	0.0277	0.0336	-0.0014	-0.0020	-0.0265
%CD38-/HLA-DR+ Teff CD8 T	0.0347	-0.0061	-0.0091	0.0057	0.1491	-0.0075	-0.0097	0.0058
%CD38+/HLA-DR+ Teff CD8 T	0.0169	0.0205	-0.0082	-0.0012	-0.0079	-0.0012	0.0405	-0.0260
%CD38+/HLA-DR- Teff CD8 T	0.0882	0.0033	-0.0072	-0.0091	0.0581	0.0140	0.0267	0.0058
%CD38-/HLA-DR- Teff CD8 T	0.0463	0.0148	-0.0080	0.0177	0.0235	0.0082	0.0768	-0.0282
%PD-1+ Teff CD8 T	-0.0533	-0.0096	-0.0069	-0.0096	0.0301	-0.0092	0.0100	-0.0269
%CD38-/HLA-DR+ Tem CD8 T	0.0105	-0.0098	-0.0056	0.0068	0.3076	-0.0096	-0.0094	0.0082
%CD38+/HLA-DR+ Tem CD8 T	0.0590	0.0705	0.0002	-0.0094	-0.0055	-0.0078	0.0234	-0.0277
%CD38+/HLA-DR- Tem CD8 T	0.0284	-0.0097	-0.0098	-0.0088	0.0755	-0.0017	0.0269	-0.0026
%CD38-/HLA-DR- Tem CD8 T	0.0203	0.0354	-0.0092	-0.0037	0.1168	-0.0031	0.0298	-0.0206
%PD-1+ Tem CD8 T	-0.0176	0.0177	-0.0101	0.0163	-0.0079	-0.0095	-0.0089	-0.0302
%IL-2+ CD8 T	-0.0183	0.0343	-0.0095	-0.0046	0.0105	-0.0099	-0.0094	-0.0285
%IL-2+ CD4 T	0.0432	0.0440	-0.0092	0.0165	0.0134	-0.0070	-0.0093	-0.0095

%TNFα+ CD4 T	-0.0125	-0.0042	-0.0066	0.0179	0.0284	-0.0084	0.0010	-0.0128
%TNFα+ CD8 T	-0.0420	-0.0056	-0.0072	-0.0045	0.0116	0.0008	0.0082	-0.0262
%IFNγ+ CD4 T	-0.0538	-0.0040	-0.0017	-0.0001	-0.0082	-0.0103	-0.0078	-0.0240
%IFNγ+ CD8 T	-0.0694	-0.0086	-0.0107	-0.0079	-0.0047	-0.0084	0.0037	-0.0265
%CD107+ CD4 T	-0.0540	-0.0096	-0.0010	-0.0104	-0.0018	-0.0103	0.0062	-0.0246
%CD107+ CD8 T	-0.0654	-0.0105	-0.0098	-0.0030	0.0016	-0.0067	0.0051	-0.0316
%CD107+IFNγ+IL-2-TNFα- CD4 T	-0.0579	-0.0106	-0.0001	-0.0102	-0.0091	-0.0104	-0.0018	-0.0261
%CD107+IFNγ-IL-2+TNF- CD4 T	-0.0403	-0.0012	-0.0068	-0.0105	-0.0081	-0.0081	-0.0004	-0.0126
%CD107+IFNγ-IL-2-TNFα+ CD4 T	-0.0379	-0.0005	-0.0103	-0.0085	0.0372	0.0141	-0.0020	-0.0144
%CD107+IFNγ-IL-2-TNFα- CD4 T	-0.0180	-0.0093	0.0068	-0.0084	0.0252	-0.0098	0.0331	-0.0302
%CD107-IFNγ+IL-2+TNFα- CD4 T	0.0165	0.0373	-0.0056	-0.0044	-0.0078	-0.0077	0.0241	-0.0208
%CD107-IFNγ+IL-2+TNFα+ CD4 T	-0.0227	-0.0097	0.0035	0.0397	-0.0035	-0.0032	-0.0024	-0.0299
%CD107-IFNγ+IL-2-TNFα- CD4 T	-0.0039	-0.0066	-0.0028	0.0169	0.0074	-0.0101	0.0031	-0.0293
%CD107-IFNγ-IL-2+TNFα+ CD4 T	0.1869	0.0191	-0.0063	0.0441	0.0921	0.0022	-0.0001	-0.0072
%CD107-IFNγ-IL-2+TNFα- CD4 T	0.0574	0.0602	-0.0050	0.0159	-0.0094	-0.0028	-0.0059	-0.0154
%CD107-IFNγ-IL-2-TNFα+ CD4 T	0.0014	-0.0047	-0.0081	0.0073	0.0609	-0.0020	-0.0040	-0.0132
%CD107+IFNγ+IL-2-TNFα- CD8T	-0.0726	-0.0107	-0.0092	-0.0094	-0.0098	-0.0106	0.0009	-0.0288
%CD107+IFNγ-IL-2+TNF- CD8 T	-0.0055	-0.0008	0.0028	-0.0097	0.0795	-0.0041	-0.0100	-0.0280
%CD107+IFNγ-IL-2-TNFα+ CD8 T	-0.0106	-0.0059	0.0119	0.0050	0.0045	0.0197	0.0025	-0.0292
%CD107+IFNγ-IL-2-TNFα- CD8 T	-0.0117	-0.0079	-0.0045	0.0090	0.0244	-0.0073	0.0051	-0.0150
%CD107-IFNγ+IL-2+TNFα- CD8 T	-0.0254	0.0099	-0.0016	-0.0019	0.0120	-0.0091	-0.0039	-0.0221
%CD107-IFNγ+IL-2-TNFα+ CD8 T	0.0199	-0.0081	-0.0009	-0.0078	0.0461	-0.0018	0.0221	-0.0272
%CD107-IFNγ+IL-2-TNFα- CD8 T	-0.0597	0.0014	-0.0056	-0.0076	-0.0005	-0.0098	-0.0031	-0.0303
%CD107-IFNγ-IL-2+TNFα+ CD8 T	0.0353	-0.0092	-0.0086	0.0084	0.1003	0.0015	0.0215	-0.0228
%CD107-IFNγ-IL-2+TNFα- CD8 T	0.0040	0.0565	-0.0059	-0.0079	-0.0092	-0.0089	-0.0080	-0.0256
%CD107-IFNγ-IL-2-TNFα+ CD8 T	0.0241	0.0090	-0.0034	-0.0099	0.0384	-0.0091	-0.0098	-0.0193

Table S7: *Adjusted R^2 scores and differences in adjusted R^2 for LOCO analysis for the model that contains intact reservoir size.*
The first column contains all immunophenotypes and is a target for the regression model. The second column contains adjusted R^2 of the linear regression model with features such as intact reservoir size, age, biological sex, years of treatment, CD4 nadir, recent CD4 count, and years of HIV before treatment (=NA, < 1, ≥ 1). The next seven columns show differences in adjusted R^2 score ($\triangle R^2$) after removing a specific feature. Participants with missing years of ART values are excluded from this analysis. The missing value of the CD4 nadir for one participant is imputed.

Immunophenotype / Model	Adj. R^2 Include all	ΔR^2, difference in adjustable R^2 after dropping a covariate						
		% intact	Age	Sex	Years of ART	CD4 nadir	Recent CD4 count	Years before ART
%CD4 T	0.2226	-0.0042	0.0132	0.0312	0.0119	0.0227	0.1479	-0.0137
%CD27+ CD4 T	0.0449	0.0179	0.0034	-0.0003	-0.0057	-0.0022	0.0263	-0.0087
%CD38+ CD4 T	0.1254	0.0542	0.0327	0.0028	-0.0073	0.0152	0.0009	-0.0258
%CD127+ CD4 T	0.0887	0.0866	0.0244	-0.0016	-0.0008	-0.0091	0.0057	-0.0170
%HLA-DR+ CD4 T	0.0718	-0.0094	0.0091	0.0298	-0.0091	0.0048	0.0259	-0.0053
%KLRG1+ CD4 T	0.0530	0.0029	0.0190	-0.0096	-0.0094	-0.0066	0.0135	-0.0026
%NKG2A+ CD4 T	0.0256	-0.0079	0.0420	0.0004	-0.0084	0.0116	-0.0015	-0.0132
%PD-1+ CD4 T	0.0995	0.0193	0.0011	0.0082	-0.0057	0.0008	0.0254	-0.0267
%Tcm CD4 T	0.0385	-0.0089	0.0446	-0.0041	-0.0080	-0.0008	-0.0091	-0.0110
%Tn CD4 T	0.1099	0.0021	0.0460	0.0083	-0.0081	-0.0075	0.0339	-0.0262
%Teff CD4 T	-0.0402	0.0024	0.0060	-0.0082	-0.0085	0.0009	-0.0051	-0.0198
%Tem CD4 T	0.0517	0.0119	-0.0093	-0.0042	-0.0095	-0.0092	0.0540	-0.0159
%CD38-/HLA-DR+ CD4 T	0.0656	-0.0054	0.0203	0.0350	-0.0040	0.0152	-0.0047	0.0136
%CD38+/HLA-DR+ CD4 T	0.0604	-0.0050	0.0039	0.0077	0.0030	-0.0073	0.0614	-0.0252
%CD38+/HLA-DR- CD4 T	0.1324	0.0422	0.0192	0.0157	-0.0088	0.0182	-0.0087	-0.0248
%KLRG1-/CD27+ CD4 T	0.0823	0.0115	0.0035	-0.0077	-0.0090	-0.0036	0.0332	-0.0103
%KLRG1+/CD27+ CD4 T	0.0160	-0.0098	0.0225	-0.0071	-0.0075	0.0271	-0.0052	-0.0285
%KLRG1+/CD27- CD4 T	0.0484	0.0147	-0.0115	-0.0054	-0.0088	0.0059	0.0104	0.0099
%KLRG1-/CD27- CD4 T	0.0525	-0.0052	0.0246	0.0019	0.0020	-0.0031	0.0205	-0.0163
%KLRG1-/PD-1+ CD4 T	-0.0010	-0.0064	-0.0084	-0.0055	-0.0081	0.0008	0.0014	-0.0054
%KLRG1+/PD-1+ CD4 T	0.0500	0.0130	-0.0001	-0.0001	-0.0089	-0.0094	0.0059	-0.0001
%KLRG1+/PD-1- CD4 T	0.0341	-0.0090	0.0356	0.0053	-0.0059	-0.0018	0.0089	-0.0189
%KLRG1-/PD-1- CD4 T	0.1663	0.0176	0.0259	-0.0058	-0.0081	0.0096	0.0475	-0.0227
%PD-1-/CCR7+ CD4 T	0.1004	0.0238	0.0085	0.0011	-0.0078	-0.0048	0.0400	-0.0264
%PD-1+/CCR7+ CD4 T	0.0682	-0.0037	0.0148	-0.0057	-0.0048	0.0198	-0.0067	-0.0103
%PD-1+/CCR7- CD4 T	0.0464	0.0153	-0.0087	0.0053	-0.0095	-0.0086	0.0333	-0.0136
%PD-1-/CCR7- CD4 T	-0.0080	0.0037	0.0638	-0.0066	-0.0067	-0.0030	0.0311	-0.0218
%CD8 T	0.1989	-0.0074	0.0053	0.0258	0.0003	0.0200	0.1360	-0.0139
%CD27+ CD8 T	0.0323	-0.0098	0.0504	0.0023	-0.0070	-0.0091	0.0218	-0.0162
%CD38+ CD8 T	0.2852	0.0056	0.1183	0.0090	-0.0072	0.0209	0.0719	-0.0088
%CD127+ CD8 T	0.0451	0.0677	-0.0077	-0.0012	0.0156	-0.0076	0.0014	-0.0052
%HLA-DR+ CD8 T	0.0203	0.0004	0.0698	-0.0088	-0.0067	-0.0097	0.0349	-0.0137
%KLRG1+ CD8 T	0.1735	-0.0031	0.2042	0.0258	-0.0075	-0.0041	0.0041	-0.0177
%PD-1+ CD8 T	0.1426	-0.0037	0.0881	0.0354	-0.0056	-0.0066	0.0067	-0.0246
%Tn CD8 T	0.2655	-0.0038	0.2266	0.0227	-0.0074	-0.0073	0.0186	-0.0094
%Teff CD8 T	-0.0265	-0.0104	0.0353	0.0102	0.0087	-0.0103	-0.0081	-0.0227
%Tem CD8 T	0.0267	-0.0037	0.0259	-0.0073	-0.0069	-0.0098	0.0334	-0.0285
%CD38-/HLA-DR+ CD8 T	0.0282	-0.0098	0.1017	-0.0090	-0.0097	-0.0092	-0.0027	0.0029
%CD38+/HLA-DR+ CD8 T	0.0398	0.0163	-0.0048	-0.0026	-0.0032	-0.0070	0.0598	-0.0280
%CD38+/HLA-DR- CD8 T	0.3152	-0.0066	0.1514	0.0365	-0.0047	0.0187	0.0139	0.0010
%KLRG1-/CD27+ CD8 T	0.1707	-0.0045	0.1261	0.0285	-0.0068	-0.0042	0.0119	-0.0168
%KLRG1+/CD27+ CD8 T	0.1003	0.0020	0.0883	0.0156	-0.0090	0.0125	-0.0084	-0.0265
%KLRG1+/CD27- CD8 T	0.0358	-0.0096	0.0687	0.0028	-0.0082	-0.0090	0.0126	-0.0170
%KLRG1-/CD27- CD8 T	-0.0016	-0.0072	0.0139	-0.0101	-0.0068	-0.0101	0.0166	-0.0295
%KLRG1-/PD-1+ CD8 T	-0.0699	-0.0094	-0.0089	-0.0108	-0.0076	-0.0093	-0.0108	-0.0318
%KLRG1+/PD-1+ CD8 T	0.1564	-0.0002	0.0930	0.0444	-0.0070	-0.0074	0.0104	-0.0240
%KLRG1+/PD-1- CD8 T	-0.0114	-0.0100	0.0916	-0.0087	-0.0039	-0.0066	-0.0096	-0.0226
%Tcm CD8 T	0.2265	-0.0076	0.1372	-0.0073	0.0086	-0.0063	-0.0062	-0.0219
%KLRG1-/PD-1- CD8 T	0.2315	-0.0034	0.1974	0.0305	-0.0075	-0.0014	0.0060	-0.0150
%PD-1-/CCR7+ CD8 T	0.1771	-0.0054	0.1078	0.0450	-0.0054	-0.0066	0.0130	-0.0141
%PD-1+/CCR7+ CD8 T	0.1294	-0.0088	0.0710	0.0147	-0.0034	0.0068	0.0029	-0.0256

%PD-1+/CCR7- CD8 T	0.0997	-0.0008	0.0392	0.0224	-0.0085	-0.0084	0.0496	-0.0259
%PD-1-/CCR7- CD8 T	-0.0384	-0.0105	0.0979	-0.0045	0.0060	-0.0104	-0.0072	-0.0207
% NK	0.1179	-0.0072	-0.0041	0.1584	-0.0043	0.0064	-0.0088	-0.0250
%NKG2A+ NK	0.1063	-0.0003	-0.0064	0.1625	-0.0074	-0.0087	0.0087	-0.0255
%NKG2A-/CD16+ NK	0.0750	0.0044	-0.0076	0.1232	-0.0082	-0.0031	0.0171	-0.0262
%NKG2A+/CD16+ NK	0.0673	0.0307	0.0094	0.0490	-0.0074	-0.0039	-0.0089	-0.0161
%NKG2A+/CD16- NK	0.0997	-0.0079	-0.0091	0.1630	-0.0086	-0.0004	0.0407	-0.0246
%NKG2A-/CD16- NK	-0.0267	-0.0062	-0.0068	-0.0058	-0.0103	0.0091	-0.0038	-0.0304
% NKT	-0.0500	0.0182	0.0037	-0.0074	-0.0094	-0.0081	-0.0025	-0.0303
%NKG2A+ NKT	-0.0161	-0.0091	-0.0089	-0.0012	0.0122	-0.0103	-0.0099	-0.0291
%NKG2A-/CD16+ NKT	-0.0444	-0.0098	-0.0058	0.0204	-0.0105	-0.0026	-0.0101	-0.0301
%NKG2A+/CD16+ NKT	0.0173	-0.0078	-0.0090	0.0390	0.0075	-0.0087	0.0107	-0.0001
%NKG2A+/CD16- NKT	-0.0199	-0.0084	-0.0040	-0.0060	0.0088	-0.0102	-0.0083	-0.0302
%NKG2A-/CD16- NKT	0.0071	-0.0099	0.0149	0.0302	0.0063	-0.0065	-0.0100	-0.0294
%CD38-/HLA-DR+ Tcm CD4 T	-0.0222	-0.0069	0.0026	0.0184	-0.0006	0.0024	-0.0102	-0.0235
%CD38+/HLA-DR+ Tcm CD4 T	0.0961	0.0079	0.0032	0.0087	0.0061	-0.0077	0.0910	-0.0248
%CD38+/HLA-DR- Tcm CD4 T	0.0397	0.0211	-0.0036	-0.0096	-0.0085	0.0260	0.0338	-0.0285
%CD38-/HLA-DR- Tcm CD4 T	0.1107	0.0264	-0.0090	-0.0017	-0.0082	0.0165	0.0752	-0.0241
%PD-1+ Tcm CD4 T	-0.0226	-0.0006	-0.0113	-0.0101	-0.0079	0.0036	-0.0013	-0.0303
%CD38-/HLA-DR+ Tn CD4 T	-0.0349	0.0051	-0.0077	0.0091	-0.0044	-0.0065	-0.0095	-0.0250
%CD38+/HLA-DR+ Tn CD4 T	-0.0329	-0.0015	0.0079	-0.0094	-0.0098	-0.0098	0.0318	-0.0302
%CD38+/HLA-DR- Tn CD4 T	0.0167	0.0061	-0.0060	0.0289	-0.0091	0.0046	0.0188	-0.0276
%CD38-/HLA-DR- Tn CD4 T	0.0565	-0.0032	-0.0095	0.0383	-0.0085	0.0100	0.0723	-0.0260
%PD-1+ Tn CD4 T	-0.0284	0.0018	0.0027	0.0074	-0.0104	0.0067	-0.0096	-0.0302
%CD38-/HLA-DR+ Teff CD4 T	-0.0059	-0.0061	0.0037	-0.0083	-0.0101	-0.0014	0.0202	-0.0141
%CD38+/HLA-DR+ Teff CD4 T	0.1788	0.0392	0.0236	-0.0067	0.0170	0.0271	0.0863	-0.0214
%CD38+/HLA-DR- Teff CD4 T	0.0228	-0.0026	0.0015	-0.0076	-0.0099	-0.0097	0.0154	-0.0051
%CD38-/HLA-DR- Teff CD4 T	0.1131	0.0245	0.0151	-0.0007	0.0035	0.0279	0.0441	-0.0262
%PD-1+ Teff CD4 T	0.0225	-0.0078	-0.0117	0.0427	0.0133	-0.0056	0.0102	-0.0227
%CD38-/HLA-DR+ Tem CD4 T	-0.0070	-0.0016	0.0190	-0.0001	-0.0045	0.0210	0.0139	0.0191
%CD38+/HLA-DR+ Tem CD4 T	0.0170	0.0124	-0.0084	0.0052	0.0044	-0.0069	0.0261	-0.0292
%CD38+/HLA-DR- Tem CD4 T	-0.0037	-0.0089	-0.0044	-0.0085	-0.0099	0.0350	0.0216	0.0025
%CD38-/HLA-DR- Tem CD4 T	-0.0366	0.0088	-0.0010	0.0057	-0.0104	-0.0035	-0.0068	-0.0226
%PD-1+ Tem CD4 T	-0.0835	-0.0104	0.0200	-0.0088	-0.0083	-0.0109	-0.0109	-0.0322
%CD38-/HLA-DR+ Tcm CD8 T	-0.0639	-0.0097	0.0108	-0.0088	-0.0103	-0.0106	-0.0066	-0.0308
%CD38+/HLA-DR+ Tcm CD8 T	0.0602	0.0014	0.0102	0.0041	-0.0018	-0.0023	0.0498	-0.0251
%CD38+/HLA-DR- Tcm CD8 T	0.1118	-0.0076	0.0459	0.0047	-0.0089	0.0462	0.0262	-0.0244
%CD38-/HLA-DR- Tcm CD8 T	0.0982	0.0012	0.0406	-0.0015	-0.0065	0.0040	0.0543	-0.0260
%PD-1+ Tcm CD8 T	-0.0062	-0.0041	-0.0099	0.0149	-0.0101	-0.0101	0.0149	-0.0297
%CD38-/HLA-DR+ Tn CD8 T	0.0228	-0.0088	0.1109	-0.0007	-0.0097	0.0026	-0.0062	-0.0269
%CD38+/HLA-DR+ Tn CD8 T	-0.0163	-0.0065	-0.0072	-0.0053	0.0054	-0.0074	0.0336	-0.0301
%CD38+/HLA-DR- Tn CD8 T	0.2321	-0.0073	0.0922	0.0618	-0.0041	0.0242	0.0525	-0.0204
%CD38-/HLA-DR- Tn CD8 T	0.2051	-0.0071	0.0461	0.0394	-0.0072	0.0181	0.1133	-0.0226
%PD-1+ Tn CD8 T	0.0382	-0.0094	0.0731	0.0299	-0.0097	-0.0041	-0.0062	-0.0282
%CD38-/HLA-DR+ Teff CD8 T	0.0392	-0.0016	0.0246	0.0088	0.0071	-0.0091	-0.0097	-0.0014
%CD38+/HLA-DR+ Teff CD8 T	0.0224	0.0260	-0.0088	-0.0056	-0.0044	-0.0077	0.0463	-0.0284
%CD38+/HLA-DR- Teff CD8 T	0.0856	0.0007	0.0069	-0.0086	-0.0023	0.0063	0.0307	-0.0009
%CD38-/HLA-DR- Teff CD8 T	0.0398	0.0083	0.0023	0.0108	0.0013	0.0002	0.0860	-0.0285
%PD-1+ Teff CD8 T	-0.0542	-0.0106	-0.0087	-0.0094	-0.0102	-0.0094	0.0117	-0.0270
%CD38-/HLA-DR+ Tem CD8 T	0.0120	-0.0084	0.0383	0.0079	-0.0039	-0.0092	-0.0094	0.0033
%CD38+/HLA-DR+ Tem CD8 T	-0.0015	0.0101	-0.0028	-0.0087	0.0042	-0.0099	0.0364	-0.0290
%CD38+/HLA-DR- Tem CD8 T	0.0298	-0.0083	0.0117	-0.0085	-0.0007	-0.0038	0.0266	-0.0064
%CD38-/HLA-DR- Tem CD8 T	-0.0161	-0.0010	0.0106	-0.0069	0.0025	-0.0063	0.0411	-0.0208
%PD-1+ Tem CD8 T	-0.0432	-0.0079	0.0021	0.0207	-0.0093	-0.0101	-0.0070	-0.0310
%IL-2+ CD8 T	-0.0578	-0.0052	-0.0060	-0.0044	0.0113	-0.0106	-0.0105	-0.0313
%IL-2+ CD4 T	0.0123	0.0131	-0.0085	0.0257	-0.0062	-0.0096	-0.0100	-0.0060

%TNFα+ CD4 T	-0.0130	-0.0047	-0.0009	0.0217	0.0111	-0.0069	0.0000	-0.0105
%TNFα+ CD8 T	-0.0252	0.0112	-0.0087	-0.0059	0.0212	-0.0067	0.0023	-0.0296
%IFNγ+ CD4 T	-0.0553	-0.0055	0.0010	0.0023	-0.0103	-0.0095	-0.0084	-0.0225
%IFNγ+ CD8 T	-0.0582	0.0026	-0.0101	-0.0087	0.0006	-0.0105	-0.0002	-0.0297
%CD107+ CD4 T	-0.0548	-0.0104	0.0009	-0.0103	-0.0045	-0.0102	0.0055	-0.0248
%CD107+ CD8 T	-0.0552	-0.0004	-0.0100	-0.0044	0.0024	-0.0095	0.0025	-0.0311
%CD107+IFNγ+IL-2-TNFα- CD4 T	-0.0580	-0.0106	0.0016	-0.0102	-0.0090	-0.0104	-0.0021	-0.0267
%CD107+IFNγ-IL-2+TNF- CD4 T	-0.0451	-0.0061	-0.0066	-0.0105	-0.0105	-0.0095	0.0017	-0.0113
%CD107+IFNγ-IL-2-TNFα+ CD4 T	-0.0452	-0.0078	0.0180	-0.0075	0.0016	0.0154	-0.0039	-0.0141
%CD107+IFNγ-IL-2-TNFα- CD4 T	-0.0182	-0.0095	0.0031	-0.0083	0.0085	-0.0094	0.0371	-0.0302
%CD107-IFNγ+IL-2+TNFα- CD4 T	0.0166	0.0374	-0.0015	-0.0082	0.0016	-0.0099	0.0304	-0.0138
%CD107-IFNγ+IL-2+TNFα+ CD4 T	-0.0195	-0.0065	0.0019	0.0415	-0.0080	-0.0005	-0.0003	-0.0302
%CD107-IFNγ+IL-2-TNFα- CD4 T	-0.0074	-0.0101	-0.0035	0.0180	-0.0045	-0.0102	0.0055	-0.0293
%CD107-IFNγ-IL-2+TNFα+ CD4 T	0.1690	0.0012	0.0046	0.0524	0.0551	-0.0021	-0.0030	-0.0057
%CD107-IFNγ-IL-2+TNFα- CD4 T	-0.0122	-0.0094	-0.0078	0.0208	-0.0099	-0.0034	-0.0001	-0.0205
%CD107-IFNγ-IL-2-TNFα+ CD4 T	0.0022	-0.0039	0.0023	0.0043	0.0541	-0.0054	-0.0032	-0.0170
%CD107+IFNγ+IL-2-TNFα- CD8T	-0.0655	-0.0036	-0.0084	-0.0098	-0.0090	-0.0107	-0.0010	-0.0305
%CD107+IFNγ-IL-2+TNF- CD8 T	0.0195	0.0241	0.0090	-0.0099	0.0363	-0.0090	-0.0098	-0.0291
%CD107+IFNγ-IL-2-TNFα+ CD8 T	-0.0090	-0.0043	0.0087	0.0039	0.0015	0.0105	-0.0010	-0.0275
%CD107+IFNγ-IL-2-TNFα- CD8 T	-0.0099	-0.0060	-0.0011	0.0065	0.0059	-0.0088	0.0059	-0.0128
%CD107-IFNγ+IL-2+TNFα- CD8 T	-0.0208	0.0145	-0.0045	-0.0033	0.0135	-0.0101	-0.0085	-0.0284
%CD107-IFNγ+IL-2-TNFα+ CD8 T	0.0673	0.0392	-0.0044	-0.0087	0.0729	-0.0087	0.0138	-0.0276
%CD107-IFNγ+IL-2-TNFα- CD8 T	-0.0690	-0.0078	-0.0020	-0.0076	-0.0050	-0.0085	-0.0066	-0.0316
%CD107-IFNγ-IL-2+TNFα+ CD8 T	0.0361	-0.0085	-0.0086	0.0095	0.0697	0.0025	0.0219	-0.0220
%CD107-IFNγ-IL-2+TNFα- CD8 T	-0.0596	-0.0071	-0.0001	-0.0063	-0.0105	-0.0099	-0.0039	-0.0271
%CD107-IFNγ-IL-2-TNFα+ CD8 T	0.0474	0.0323	0.0009	-0.0088	0.0437	-0.0040	-0.0096	-0.0106

Table S8: *Adjusted R^2 scores and differences in adjusted R^2 for LOCO analysis for the model that contains percent intact.*
The first column contains all immunophenotypes and is a target for the regression model. The second column contains adjusted R^2 of the linear regression model with features such as percent intact, age, biological sex, years of treatment, CD4 nadir, recent CD4 count, and years of HIV before treatment (=NA, < 1, ≥ 1). The next seven columns show differences in adjusted R^2 score (ΔR^2) after removing a specific feature. Participants with missing years of ART values are excluded from this analysis. The missing value of the CD4 nadir for one participant is imputed.

Variable	Bin Total	Bin Intact	Bin % intact	Variable	Bin Total	Bin Intact	Bin % intact
%NKG2A+ CD4 T	**0.6953**	0.6084	0.5416	%PD-1-/CCR7+ CD8 T	0.6207	0.6134	0.5115
%PD-1+ Tn CD4 T	**0.6827**	0.5749	0.5313	%CD38+ CD8 T	0.6142	0.5425	0.6222
%CD38+/HLA-DR- CD8 T	**0.6777**	0.5797	0.6084	%CD38-/HLA-DR- Teff CD4 T	0.6122	0.5587	0.5017
%Tn CD4 T	**0.6763**	0.5933	0.5665	%CD38-/HLA-DR+ Tn CD8 T	0.6122	0.5420	0.5933
%KLRG1-/PD-1- CD4 T	**0.6745**	0.5789	0.6224	%KLRG1-/CD27+ CD8 T	0.6096	0.6101	0.5265
%Tcm CD8 T	**0.6729**	0.6205	0.5541	%CD38+/HLA-DR- Tn CD4 T	0.6078	0.5050	0.5531
%CD38+/HLA-DR- CD4 T	**0.6700**	0.5507	0.5889	%CD107-IFNγ+IL-2-TNFα- CD8 T	0.6065	0.5712	0.5133
%CD8 T	**0.6629**	0.6547	0.5484	%CD38+/HLA-DR+ Tem CD4 T	0.6057	0.5829	0.5941
Age	**0.6627**	0.6264	0.5476	%CD38+/HLA-DR- Tcm CD8 T	0.6039	0.5150	0.5750
%PD-1+/CCR7+ CD8 T	**0.6608**	0.5907	0.5328	%CD38-/HLA-DR+ Tcm CD8 T	0.6030	0.6053	0.5015
%PD-1+ CD8 T	**0.6562**	0.6207	0.5070	%KLRG1-/PD-1+ CD4 T	0.6019	0.5346	0.5290
%PD-1+ Tn CD8 T	**0.6553**	0.5880	0.5597	%NKG2A-/CD16+ NK	0.6012	0.5257	0.5848
%Tn CD8 T	**0.6550**	0.6485	0.5138	%KLRG1+ CD8 T	0.5954	0.6021	0.5251
%PD-1-/CCR7+ CD4 T	**0.6537**	0.5473	0.6178	%KLRG1-/CD27+ CD4 T	0.5945	0.5322	0.6205
%PD-1+/CCR7+ CD4 T	0.6475	0.5656	0.5740	%CD27+ CD4 T	0.5776	0.5018	0.6021
%Tcm CD4 T	0.6452	0.5998	0.5104	%KLRG1+/CD27- CD4 T	0.5723	0.5017	0.6181
%KLRG1-/PD-1- CD8 T	0.6440	0.6308	0.5234	%PD-1+ Tcm CD4 T	0.5705	0.5005	0.6158
%PD-1+ CD4 T	0.6434	0.5286	0.6295	%PD-1+ Tem CD4 T	0.5688	0.6124	0.5489
%CD4 T	0.6429	0.6338	0.5322	%KLRG1+ CD4 T	0.5649	0.5243	0.6236
%CD107-IFNγ+IL-2-TNFα- CD4 T	0.6420	0.5670	0.5185	Recent CD4 count	0.5620	0.6320	0.6104
%CD38+ CD4 T	0.6414	0.5299	0.5932	%KLRG1+/PD-1+ CD4 T	0.5602	0.5100	0.6450
%CD127+ CD4 T	0.6367	**0.7051**	0.6323	%Tem CD4 T	0.5576	0.5230	0.6145
%CD38+/HLA-DR- Tn CD8 T	0.6364	0.5558	0.5724	%IL-2+ CD4 T	0.5416	0.6022	0.5687
%PD-1-/CCR7- CD4 T	0.6340	0.6010	0.5473	%CD107-IFNγ+IL-2+TNFα- CD4 T	0.5342	0.6308	0.6248
%KLRG1+/PD-1+ CD8 T	0.6335	0.6098	0.5020	%CD38+/HLA-DR- Teff CD8 T	0.5306	0.5613	0.6276
CD4 Nadir	0.6303	0.6110	0.5563	Years before ART < 1	0.5299	0.5222	0.6092
Years of ART	0.6296	0.5069	**0.7171**	%CD127+ CD8 T	0.5195	0.6051	0.6048
%CD38-/HLA-DR- Tem CD4 T	0.6249	0.5513	0.5867	%CD107-IFNγ-IL-2+TNFα+ CD4 T	0.5003	0.6000	0.6493
%KLRG1+/CD27+ CD8 T	0.6224	0.6473	0.5153				

Table S9: *Host features classify PWH with respect to HIV reservoir characteristics.*
The abundance of each of the 144 variables was analyzed for performance in classifying each PWH within the cohort as having reservoir size above or below the median using a receiver operating characteristics (ROC) curve. The area under the curve (AUC) is reported. The table includes variables that had an AUC value larger than 0.6 for one of the reservoir size characteristics. We highlight variables that had an AUC value higher than 0.65 in **bold**. Variables are ranked by AUC for the total reservoir size. For years of ART, AUC values are computed based on 108 PWH, excluding participants with missing years of ART values. For CD4 Nadir, AUC values are computed based on 114 PWH, excluding one participant] with the missing CD4 Nadir value.

Figure S8: *Dimension reduction supplemental figures.*
A. Participant age is shown within each cluster. **B.** Participant years of ART are shown within each cluster. **C.** Participant CD4 nadir is shown within each cluster. **D.** Participant CD4 count is shown within each cluster. **E-O**: Participant immune features of interest are shown, where plots of immune features with similar names are placed nearby. **P.**: Clusters with data points color-coded by intact reservoir size (high=purple, low=pink). **Q.**: Clusters with data points color-coded by percent intact reservoir size (high=purple, low=pink). **R.** Relative proportions of cannabis (CB) users and non-users (non-CB) are shown for each cluster. **S.** Total reservoir sizes for non-users and CB users are shown. **T.** The ages of study participants for non-users and CB users are shown.

Total reservoir size

To read the tree, consider each leaf and follow the path from the root to this leaf.
All features that are encountered on the path will construct a rule.

Figure S9: *The total reservoir size visualization tree from Figure 6 is explained with a set of rules.*
For every leaf, the path that leads to this leaf is described. The histogram of the variable and split value for every node used in the tree is shown.

Figure S10: *The intact reservoir size visualization tree from Figure 6 is explained with a set of rules.* For every leaf, the path that leads to this leaf is described. The histogram of the variable and split value for every node used in the tree is shown.

Percent intact

To read the tree, consider each leaf and follow the path from the root to this leaf.
All features that are encountered on the path will construct a rule.

Figure S11: *The percent intact visualization tree from Figure 6 is explained with a set of rules.*
For every leaf, the path that leads to this leaf is described. The histogram of the variable and split value for every node used in the tree is shown.

Figure S12: *Using machine learning to predict reservoir size.*
Average training and test R^2 scores over different training and test data splits for Linear Regression (LR), Ridge Regression (RR), Kernel Regression with RBF kernel (KR), Decision Tree Regressor (DT), Random Forest (RF), and Gradient Boosted Trees (GBT) models are shown for predicting total reservoir size (**A**), intact reservoir size (**B**), and percentage intact (**C**). For one split of training and test sets, ridge regression performance is shown for total reservoir size (**D**), intact reservoir (**E**), and linear regression for percentage intact (**F**). On the x-axis, the actual values of HIV reservoir characteristics are shown, while on the y-axis the outputs of models are shown for training data (blue dots) and test data (red dots). For the same training and test split, ridge regression model feature coefficients are visualized for total reservoir (**G**), intact reservoir (**H**), and linear regression model feature coefficients are visualized for percent intact (**I**). On the y-axis we show variables used by the model, while on the x-axis coefficient values for linear models based on these variables. For percent intact models we exclude participants with missing values of years of ART. For total and intact reservoir size, missing values of years of ART were imputed. The missing value of the CD4 nadir for one participant was imputed as well using the MICE algorithm.

Algorithm 1 Training procedure for classification (regression)

    Input: Outcome, algorithm, set of hyperparameters from Table S10
    Output: Mean and standard deviation of train and test accuracies (R^2 scores)

1: Compute product of hyperparameters for grid search, let H be one hyperparameter setting
2: **for** k in the set of top AUC (correlation) variables to consider **do**
3:     Take the first k variables from the set of variables
4:     **for** test seed in the array of random seeds for train/test splits **do**
5:         Sort data according to test seed
6:         Split data into test and train in proportion 25% to 75%
7:         **for** every hyperparameters settings H **do**
8:             **for** validation seed in the array of random seeds for train/valid splits **do**
9:                 Sort data according to validation seed
10:                 Split data into validation and train for validation in proportion 25% to 75%
11:                 Train algorithm with hyperparameter setting H on train for the validation set
12:                 Compute accuracy (R^2 score) on validation set
13:             **end for**
14:             Compute mean validation accuracy (R^2 score) on all validation sets for hyperparameter setting H
15:         **end for**
16:         Find hyperparameter setting (H_m) with the highest mean validation accuracy
17:         Train algorithm with hyperparameter setting H_m on train data
18:         Compute train and test accuracy (R^2 score)
19:     **end for**
20:     Compute accuracy (R^2 score) mean and std over different train and test sorting
21: **end for**
22: Among mean validation accuracies (R^2 scores) for different k find the maximum
23: Return train and test accuracies (R^2 scores) mean and standard deviation that corresponds to the number of variables k with the highest mean validation accuracy (R^2 score)

General	
Random seeds for train/test splits	0..10
Random seeds for train/valid splits	0..20
Set of the top AUC variables to consider	2..9,10,15,20,25,30,40,50
Set of the top absolute value of correlation coefficient to consider	2..9,10,15,20,25,30,40, all
Logistic Regression (classifier)	
Penalty	L2
Regularization parameter	0.00001, 0.0001, 0.001, 0.01, 0.1, 1, 10, 100, 1000, 10000
SVM with RBF kernel (classifier)	
Regularization parameter	0.00001, 0.0001, 0.001, 0.01, 0.1, 1, 10, 100, 1000, 10000
Ridge Regression (regressor)	
Regularization parameter	0.00001, 0.0001, 0.001, 0.01, 0.1, 1, 10, 100, 1000, 10000
Kernel Ridge Regression (regressor)	
Regularization parameter	0.00001, 0.0001, 0.001, 0.01, 0.1, 1, 10, 100, 1000, 10000
Kernel parameter	0.00001, 0.0001, 0.001, 0.01, 0.1, 1, 10, 100, 1000, 10000
CART (classifier) and **Decition Tree** (regressor)	
Maximum depth	2..7, 10
Minimum number of samples in the leaf	4, 6, 8, 10
Random Forest and Gradient Boosted Trees (classifier and regressor)	
Maximum depth	2..7, 10
Minimum number of samples in the leaf	4, 6, 8, 10
Maximum number of estimators	2, 5, 10, 20, 50, 100, 200

Table S10: *Ranges of hyperparameters values that we used to perform grid search for classification and regression.*