Title: Building a barrier: investigating the functional role and evolution of induced trichome production

Authors: Nia M Johnson[1,2] and Regina S Baucom[1]
[1]Ecology and Evolutionary Biology Department, 4034 Biological Sciences Building, University of Michigan, Ann Arbor, MI 48109 [2]Negaunee Institute for Plant Conservation Science and Action, Chicago Botanic Garden, 1000 Lake Cook Rd., Glencoe, IL, 60022, USA,

Corresponding author: nmjo@umich.edu

ORCIDs:
NMJ- orcid.org/0000-0002-3110-991X
RSB- orcid.org/0000-0001-7960-498X

Acknowledgments
The authors thank Michael Palmer and Jeremy Moghtader at the Matthaei Botanical Gardens for assistance with experiment logistics and herbicide applications. Many thanks to Kaira Liggett, Shantrell Tremmell, and Alanna Miyashiro who helped with field maintenance and with capturing pictures for herbivory estimates. Raj Gautam and Jordan Herman who helped collect and count seeds. We also thank Gregg Sobocinski at the University of Michigan Imaging Core for assistance with capturing trichome images.

Funding
This research was supported by a Rackham Research Grant, Howard Hughes Medical Institute Grant (Gilliam Fellows grant to both NMJ and RSB), and the Nancy Williams Walls Award for Field Research.

Conflict of Interest
The authors declare no conflict of interest.

Author contributions
N.M.J and R.S.B. contributed to the experimental design, writing, and editing of this manuscript.

Data Availability Statement
The data underlying this article are available in Github, at
https://github.com/niajohnson1/HerbicideInducedTrichomes

Abstract

Functional traits are increasingly recognized as fundamental determinants for mediating
ecological resilience. As global change accelerates, phenotypic plasticity in functional traits may
provide a modality for fine-tuning responses to atypical or stressful environments. While various
studies have explored induced plant trichomes as a response to biotic stress, emerging evidence
suggests that trichome induction may also function as a broader tactic for managing a range of
environmental stressors, including abiotic chemical stress. Using the annual invasive weed
velvetleaf (*Abutilon theophrasti*), we examine whether herbicide exposure induces changes in
trichome traits and whether these responses are associated with herbicide resistance and fitness
outcomes. We quantified changes in trichome density and trichome type proportions across
control and glyphosate (active ingredient in "Roundup") treated plants. We identified positive
correlations between induced total trichome density and herbicide resistance, as well as induced
branched trichomes and herbicide resistance. Selection analysis further revealed positive linear
selection acting upon induced trichome density in the presence of herbicide, as well as
correlative selection favoring induced trichome density and intermediate plant growth. Overall,
our study indicates that plastic defenses like trichomes may contribute to plant performance
under novel abiotic stress and highlights key constraints shaping the evolution of stress-
responsiveness.

Keywords: functional traits, abiotic stressor, chemical stress, induced plant defense, trichome
production, phenotypic plasticity

Introduction

Understanding the relationship between functional traits and environmental shifts is a growing area of both evolutionary and ecological research given that global change is intensifying the frequency and severity of stressors (Aubin et al 2016, Ahrens et al 2019, Madsen-Hepp et al 2023). Phenotypic plasticity, the ability of a genotype to express various phenotypes as a response to environmental heterogeneity (Sultan and Stearns 2005), is a proposed mechanism by which plant species may modulate functional traits and persist amidst rapid environmental change (Valladares et al 2007). As abiotic challenges such as temperature extremes, altered precipitation regimes, and chemical exposures escalate, investigating how plastic functional trait responses are deployed across environments can reveal important insights into processes influencing population resilience.

Among functional traits, plant trichomes—epidermal outgrowths of various types—are widely recognized for their protective roles in responding to both biotic and abiotic stress. Although often studied as a defense against herbivory (Doss et al 1987, Haberlandt 1914, Agren and Schemske 1993, Westerbergh and Nyberg 1995, Dalin and Bjorkman 2003), trichomes also respond to abiotic factors such as ultraviolet radiation (Tattini et al 2000, Yamasaki et al 2007), drought (Boughalleb and Hajlaoui 2011), elevated CO_2 (Karowe and Gribb 2011), and temperature (Liakopoulos et al., 2006). These responses suggest that trichome production may be a general stress-responsive trait, adjusting to diverse environmental cues. However, the degree to which different trichome types respond plastically to novel stressors, such as herbicide exposure, is still poorly understood.

As chemical agents applied across agricultural (Gianessi 2013), urban (Meftaul et al 2020), and restoration landscapes (Humphries et al 2020), herbicides represent a particularly relevant and underexplored abiotic stressor. There is sufficient evidence that herbicide drifts into surrounding ecosystems (Kleijn and Snoeijing 1997, Marrs et al 1989, De Snoo and Van der Poll 1999, Egan and Mortensen 2012, Hwang et al 2022), unintentionally exposing non-target plants to chemical stress (Egan & Mortensen, 2012, Silva et al 2016). Recent uptakes in herbicide use (Benbrook 2016) also highlight the crucial need to understand its impacts on population and community dynamics. Though some studies have examined herbicide effects on plant growth and herbivory (Dewar et al., 2000; Wu et al., 2001), the extent to which herbicide exposure induces plastic changes in plant defense traits such as trichomes—and whether these traits play a role in herbicide resistance and plant fitness—remains unclear.

Here, we explore the extent to which trichome traits respond plastically to herbicide exposure and examine the potential functional significance and evolutionary trajectory of such responses. Using the model weed species *Abutilon theophrasti* (hereafter velvetleaf), a species with diverse trichome types and a history of herbicide exposure in agricultural settings, we assess whether

trichome traits are induced under herbicide stress, and whether such plastic changes correlate with resistance and/or fitness outcomes. There are four types of trichomes on the surface of velvetleaf, including two multicellular types that are glandular (peltate and capitate) and two unicellular types that are non-glandular (single and branched) (Sterling and Putnam 1987). While we know that glandular trichomes have the ability to synthesize and secrete flavonoids as chemical defense (Sterling et al 1987), limited work has explored the defensive role for the non-glandular types in this species. Further, velvetleaf has evolved reduced susceptibility to herbicides routinely used in weed management, including atrazine (Anderson and Gronwald 1991) and glyphosate (Hartzler and Battles 2001), but the mechanisms underlying this reduced susceptibility are unknown. Thus, for both of these reasons – *i.e.*, variation in trichome types and reduced herbicide susceptibility – velvetleaf provides a unique study system with which to address questions about stress-induced functional trait responses.

In past work, we found positive correlations between constitutively produced branched trichomes and glyphosate resistance, indicating that this trichome type may serve as an herbicide resistance trait in velvetleaf (Johnson and Baucom 2024). We further showed positive selection on branched trichomes in the presence of the herbicide, indicating that herbicide exposure could lead to adaptive evolution of this trichome type. We also observed that trichome phenotypes exhibited high plasticity between growth room and field conditions, and reasoned that, similar to trichome induction when plants are damaged by herbivory or abiotic factors, damage from herbicide may induce trichomes in surviving plants. In this work, we investigate trichome plasticity in *Abutilon theophrasti* and examine potential impacts of trichome induction on plant fitness and performance.

We address the following specific questions: Does the density or proportion of different trichome types (branched, single, capitate, peltate) vary when exposed to herbicide in field conditions, such that trichome traits are induced in *Abutilon theophrasti* exposed to herbicide? Is there a relationship between induced trichome traits and herbicide resistance, suggesting that trichome induction may be a general feature of more resistant plants? Further, we wanted to assess the evolutionary trajectory of induced trichome defense. Thus, we asked: Is there genetic variation associated with induced trichome traits, and do plants that display trichome plasticity also exhibit higher fitness in the presence of herbicide, indicating a benefit of trichome induction? Are there fitness costs associated with trichome plasticity, and/or tradeoffs between this form of defense and growth or reproduction? The presence of genetic variation in induced trichome traits would provide evidence that induced trichome traits have the potential to evolve, whereas a benefit or cost of trichome plasticity would provide evidence for increases or brakes on this form of plant defense.

Materials and Methods

Study organism

Velvetleaf is an invasive annual native to Asia and is frequently found in and around corn, cotton, and soybean fields globally (Warwick and Black 1985). First introduced to Pennsylvania and Virginia in the late 17th century, velvetleaf was originally cultivated as a commercial fiber crop (Spencer 1984). Despite its commercial success in other regions, the fiber industry in the US was not able to use velvetleaf effectively, leading to the lack of management and rapid growth in and around agricultural fields. Today, this annual species has become one of the most detrimental weed species between 32º and 45º N latitude, particularly in southwestern Canada and the United States (Warwick and Black 1985).

Velvetleaf is a highly self-pollinating, hexaploid species with 2n = 42 chromosomes, and with high levels of fixed heterozygosity as a result of polyploidy (Warwick and Black 1985). Seed production ranges from 700 to 17,000 seeds per individual (Winter 1960; Chandler and Dale 1974; Hartgerink and Bazzaz 1984; Warwick and Black 1985), and seed dormancy can last up to 50 years (Warwick and Black 1988). Studies on velvetleaf accessions in North America reveal phenotypic variation that correlates with latitudinal environmental heterogeneity. For instance, southern populations found in North America produced fewer, larger seeds with lower rates of seed dormancy when compared to northern populations (Warwick and Black 1985). This species has also been shown to exhibit resistance or decreased susceptibility to several herbicide classes including triazine (Ritter 1986, Anderson and Gronwald 1991) and glyphosate (Hartzler and Battles 2001).

In addition to resistance to herbicides, velvetleaf also has traits that reduce the potential for herbivory. These traits include glandular trichomes on the plant surface, which serve as chemical defense structures via the synthesis and storage of flavonoids such as anthocyanin, quercetin, kaempferol, and myricetin (Morris and Wang 2013). The glandular morphs are multicellular structures that include peltate trichomes, which contain 4 - 5 cells, and capitate trichomes, which contain 12 - 15 cells (Sterling 1987, Fig 1). Velvetleaf also has non-glandular trichomes, which serve as physical defense structures by shielding the plant body. The non-glandular morphs are unicellular structures that include single trichomes, which grow perpendicular to the plant surface, and branched trichomes, which grow 4 - 8 arms (Sterling 1987, Fig 1).

Field Experiment

We conducted a field experiment to measure induced trichomes in field conditions and in response to herbicide application. Seeds were sampled from individuals found in populations located near agricultural farms in Dexter, MI (see SFig 1 of Johnson and Baucom, 2024) and were planted and selfed in growth room conditions to generate replicate, selfed seed for the field experiment (Johnson and Baucom 2024). We planted scarified velvetleaf seeds in a randomized block design comprising two treatments (herbicide and control). Three replicate individuals per

maternal line were planted in each treatment/block combination for a total of 720 seeds (40
maternal lines x 2 treatments x 3 blocks x 3 replicates). When plants reached an average height
of 11 cm, roughly 5 weeks after germination, we applied glyphosate at 550g/ha ai to the
herbicide treatments, as our previous study revealed genetic variation for reduced susceptibility
at this application dose (Johnson and Baucom 2024). We recorded height measurements to
estimate growth rate two weeks after planting and once again when we collected fitness
estimates, approximately 12 weeks after planting.

As an annual species, velvetleaf seed count indicates the success for which individual genetic
material may be passed on to subsequent generations, thus we elected to use seed count as an
estimate of fitness. Plants that germinated but did not survive herbicide exposure were given a
fitness of zero. To estimate herbicide damage in the field, two weeks after herbicide application
we recorded the number of yellowing leaves and divided that by the total number of leaves per
individual. Because velvetleaf begins to shed its leaves during seed maturation, we choose this
metric to estimate herbicide damage rather than biomass so that fitness data could also be
collected. Herbicide resistance was defined operationally as the inverse of herbicide damage (1
minus the proportion of yellowing leaves).

Due to low germination rates and mortality following herbicide application in the field, there
were 25 maternal lines remaining for which induced trichomes could be estimated (i.e.
represented in both treatments). Once plants began to flower, approximately 10 weeks after
germination, we collected one leaf from 2-3 randomly chosen plants per maternal line in each
treatment for trichome imaging (n = 130), ensuring that we sampled leaves that were newly
formed post- herbicide exposure. Along with measuring total trichome density, we calculated the
proportion of each trichome type by dividing the absolute number of each trichome structure by
the total trichome density recorded in each $3mm^2$ segment. We estimated the weekly growth rate
as the difference in height measured 2 weeks after germination and 12 weeks after germination
divided by 10.

Data Analysis – Phenotypic plasticity in trichome production
We conducted all statistical analyses in R (version 3.4.2, R Development Core Team). We ran a
series of univariate models to assess the potential for plasticity in trichome production in
different environments. For each model, the response variable was one of the observed trichome
traits (density and the proportion of each type) and the fixed effect was the environment (*i.e.*
control vs herbicide). We performed subsequent post-hoc tests using Tukey to detect differences
between environments.

Genetic variation underlying induced trichome traits

We tested for genetic variation underlying induced trichome traits to determine if trichome
induction would be expected to respond to selection and evolve. To do so, we fit the following
model for each trait:
222
223 $$y = u + \text{treatment} + (1|\text{ maternal line}) + \text{treatment} \times (1|\text{ maternal line}) + e$$
224
where each respective trichome traits (total density, and the proportion of branched, single,
peltate, and capitate) was the response variable, u is the intercept or mean of the model,
treatment was a fixed effect, maternal line was a random effect, maternal line by treatment
interaction was a random effect, and e is the error term. Our specific effect of interest was the
interaction of maternal line with treatment, since such an interaction would indicate that trichome
traits of maternal lines responded differently to the control and herbicide environments and thus
implicate adaptive plasticity. As such, the remaining analysis was performed on trichome traits
with significant maternal by treatment effects. For each trait, we first removed the effects of
block by performing a regression of the trait with the block effect as the sole predictor variable,
and then used the residuals in each subsequent analysis of variance. We determined the
significance of predictor variables using the stats package (R Core Team 2022) to generate *F*-
statistics for the fixed effects. We used the lmerTest package (Kuznetsova et al 2017) to perform
a log-likelihood ratio test for each random effect.
238
Herbicide resistance and induced trichomes
We calculated induced trichomes in field settings as the mean difference between damaged (*i.e.*
herbicide treatment) and control (*i.e.* control treatment) states for each maternal line in the field
experiment. As such, this measurement produced a positive number when trichomes increased
(*i.e.*, induced resistance) in response to herbicide and a negative number when trichomes
decreased (*i.e.*, induced susceptibility).
245
246 We elected to perform the remaining analysis on induced total density, and induced changes in
247 the proportion branched and capitate trichomes because these three traits exhibited significant
248 genetic variation (see Results). We estimated Pearson correlation coefficients to determine if
249 induction of these traits had a significant relationship with herbicide resistance and tested the
250 significance of the correlation coefficient using the stats package.
251
Genotypic selection for induced trichomes
We performed genotypic selection analysis to determine if trichome induction in response to
herbicide exposure is under selection and to assess if there was a cost when compared to control
conditions. We estimated differentials to measure selection on induced trichome density,
proportion branched and capitate trichomes in the presence and absence of herbicide. Because
the proportion of branched trichomes and capitate trichomes are not independent of total
trichome density, we estimated selection differentials rather than gradients. We estimated

selection differentials (S) by performing univariate regressions of relative fitness in each trait
separately which captures total (indirect and direct) selection acting upon each trait. Relative
fitness was calculated as the final seed count per individual divided by mean seed count for each
treatment.
263
Inducible defenses are hypothesized to come at the expense of plant growth as a form of cost
(Herms and Mattson 1992), thus, we explored if there was correlative selection acting on the
interaction between growth rate and induced trichomes. We performed non-linear selection
analyses (γ) containing trichome traits and growth rate as linear terms, quadratic terms, and the
cross-product of the focal trichome trait and growth rate. We assessed the potential of quadratic
or correlative selection by doubling the quadratic regression coefficients. Significant quadratic
terms would indicate non-linear (disruptive or stabilizing) selection acting upon growth rate or
induced trichome traits, whereas significant correlative selection would indicate selection
favoring a combination of the focal traits.
273
274
275 **Results**
276
Trichome production in *Abutilon theophrasti* is highly plastic, with significant changes in
trichome density produced according to environment. When comparing between the control
(non-herbicide) environment and herbicide environment in the field, plants exposed to herbicide
generally exhibited fewer total trichomes (*i.e.* reduced trichome density: Fig 2A). We found no
evidence for changes in the proportions of the four trichome trichome types between control and
herbicide environments (Fig 2B-E).
283
However, induced trichome density varied by maternal line when plants were exposed to
herbicide (STable 1). We identified genetic variation in induction of the proportion of branched
trichomes (maternal line by treatment effect: $\chi^2 = 3.92$, p = 0.048, Table 1, Fig 3), and the
proportion of capitate trichomes (maternal line by treatment effect: $\chi^2 = 5.64$, p = 0.018, Table 1,
Fig 3). Of the 25 total maternal lines, 9 increased the proportion of branched trichomes (36%),
19 increased the proportion of capitate trichomes (76%), and 6 increased the total trichome
density (24%) after herbicide treatment. We also found significant maternal line by treatment
interaction for total trichome density ($\chi^2 = 10.05$, p = 0.002, Table 1, Fig 3).
292
Herbicide resistance and induced trichomes
We investigated if there was a significant relationship between induced trichomes and herbicide
resistance (1 – proportion herbicide damage). We found that lines that exhibited a greater
induction of branched trichomes and a greater induction of total trichome density exhibited
higher levels of herbicide resistance (r = 0.44, p = 0.035, Fig 4A; r = 0.43, p = 0.040, Fig 4C).

There was no evidence of a relationship between the proportion of capitate trichomes and
herbicide resistance (r = -0.37, p = 0.084, Fig 4B).

Genotypic selection for induced trichomes
In the absence of herbicide, we found no evidence for selection acting upon any induced trait
(branched: S = -0.03, p = 0.976; capitate: S = -0.39, p = 0.698, total density: S = -0.01, p = 0.99;
STable 2). In the presence of herbicide, however, we found negative selection for induced
capitate trichomes (S = -2.13, p = 0.045, Fig 4, STable 2) and positive selection for induced
trichome density (S = 2.56, p = 0.018, Fig 4, STable 2). We found no evidence of selection on
induced branched trichomes (S = 0.010, p = 0.925, Fig 4, STable 2) in the presence of herbicide.

In the absence of herbicide, we identified significant correlative selection acting branched
trichomes (γ = 6.13, p = 0.006, STable 3), in those genotypes with the highest growth rate and a
greater ability to alter the proportion of branched and capitate trichomes exhibited highest
fitness. However, there was no evidence of correlative selection between growth rate and
induced total density in this environment (γ = -0.87, p = 0.669, STable 3).

In contrast, in the presence of herbicide, there was evidence for correlative selection between
growth rate and induced total density (γ = 5.71, p = 0.010, STable 3), in that genotypes with the
highest fitness exhibited intermediate growth and high levels of induced total density (Fig 6.
There was no evidence of correlative selection acting upon growth rate and either induced
branched (γ = -0.77, p = 0.070, STable 3) or induced capitate (γ = -2.58, p = 0.091, STable 3)
trichomes in the presence of herbicide.

Discussion

The goal of this study was to evaluate phenotypic plasticity in trichomes as a response to
chemical stress from a common incrementally sprayed herbicide (glyphosate). Plasticity in
defense traits such as trichomes may evolve in response to spatial and temporal variability in
environmental stressors, allowing plants to balance the costs and benefits of defense, and are
particularly favored when the presence of the stressor is unpredictable (Agrawal and Karban
1999; Karban and Baldwin 2007). We explored the cost and benefits of induced trichomes,
which are apparent in the absence and presence of a damaging agent, respectively, reflecting a
tradeoff between defense and reproduction (Rhoades 1979, Zangerl and Bazzaz 1992, Gavrilets
and Scheiner, 1993).

In the broadest sense, plastic responses cause real-time modifications of resource allocation and
can result in a range of defense strategies. For example, plants can induce phytochemicals
(Baldwin and Schmelz 1994, Textor and Gershenzon 2009) and increase physical defenses such
as thorns and trichomes (Young 1987, Milewski et al 1991, Gonzales et al 2008). Of the research

on trichomes, ample work has examined environmental factors that induce trichome production to include: herbivory (Baur et al 1991, Agrawal 1999, Rautio et al 2002, Dalin and Bjorkman 2003), pathogens (Zhang et al 2020), water availability (Nagata et al 1999, Bosu and Wagner 2014, Vanhoutte et al 2017), UV radiation (Höglund and Larsson 2005), elevated CO_2 (Karowe and Grubb 2011) and temperature (Ehleringer 1982, Perez-Estrada 2000, Shibuya et al 2016). However, few studies have identified the degree to which different trichome types respond plastically to novel stressors, acting as a multi-functionality stress-responsive trait (Sack and Buckley 2020). Here, we discuss our results conducted on velvetleaf (*Abutilon theophrasti*) as it relates to induced trichome production in response to herbicide exposure.

Induced trichome production as a response to herbicide
Most plants have developed the ability to perceive and respond to external stimuli (Braam, 2005), and recent studies indicate that plant trichomes can directly sense external forces in anticipation of threats (Peiffer et al., 2009; Matsumura et al., 2022). Trichomes have been hypothesized to contribute to herbicide resistance (Devine et al., 1992; Baucom, 2019), but limited studies have empirically tested this idea. While we previously demonstrated that the proportion of branched trichomes is positively correlated with herbicide resistance in this species (Johnson and Baucom, 2024), suggesting a potential role as a constitutive resistance trait, the possibility that trichome plasticity also functions as an induced defense had not yet been explored.

In the present study, we show that trichomes are inducible in response to herbicide exposure, although the direction of the response varied. Across all plants, there was a general decrease in total trichome density on newly formed leaves following herbicide treatment, with no significant change in the proportion of trichome types. There were, however, significant maternal line effects in which half of maternal lines in this study had an increase in the proportion of capitate trichomes, our finding differs from previous work in M. guttatus which revealed that damage induced an increase in capitate trichome production. One important caveat is that we did not assess trichomes on the leaves that were directly exposed to the herbicide, which may exhibit different induction patterns. Despite an overall decrease in trichome density, however, we observed substantial variation among maternal lines. Specifically, 76% of maternal lines exhibited an increase in proportion of capitate trichomes, which aligns with previous work in *M. guttatus* which revealed that herbivory damage induced an increase in capitate trichome production (Holeski 2007). We also found that 24% of maternal lines exhibited an increase in total trichome density following herbicide exposure, and we detected a significant maternal line × treatment interaction for both traits. Moreover, those maternal lines with higher herbicide resistance showed stronger trichome induction overall—including increases in both total trichome density and the proportion of branched trichomes.

While we do not know the mechanistic basis of this positive correlation between herbicide
resistance and induced trichomes, we hypothesize three potential reasons for this relationship.
First, it is possible that genotypes that are more generally defended are likewise able to induce
trichomes when damaged. In this scenario, which we imagine as agnostic to the particular
damaging agent, the plant experiences stress that then triggers increased trichome production in
genotypes that are better defended in a broad range of ways. In other words, individuals that are
more herbicide-resistant also appear better able to mount a defense response, consistent with a
generalizable inducible defense mechanism. Second, damage from an abiotic source, and
specifically a chemical stressor like an herbicide, may trigger trichome production, potentially
enhancing the plant's ability to sequester chemicals within these structures, a process
increasingly documented across diverse species (reviewed in Li et al 2022). Given that trichome
vacuoles can occupy up to 95% of the total trichome volume (Gutiérrez-Alcalá et al., 2000), an
increase in branched trichomes may be especially advantageous, as this type often possesses the
highest vacuolar capacity (Calvert et al., 1985). Third, genes that influence trichome
development in *A. thaliana* have known pleiotropic effects that impact other physiological
functions (Bird and Gray 2003, Kirik et al 2004), thus it is possible that the induced trichomes
we observed in velvetleaf could be linked to a response not measured in our experiment. While
uncovering the exact mechanism underlying this relationship was outside of the scope of this
study, further research is needed to determine if the induction in plant trichomes in more resistant
genotypes is due to being more defended in general, having the ability to sequester chemicals for
detoxification purposes, a pleiotropic relationship to an unknown response that we did not
capture, or a combination of any of these explanations.
399
Costs and benefits of trichome plasticity
The adaptive potential of damage-induced changes to trichome production are dependent on the
benefits and costs of the phenotypic change (Dalin et al 2008). Positive correlations between
induced increases in trichome traits and herbicide resistance suggests that plants would benefit
from the induction of trichomes as long as the fitness benefits of this induction outweigh the
costs of induction. Our linear selection analysis revealed that in the presence of herbicide there
was positive selection acting upon the induced increase of total trichome density, but no
evidence of selection acting upon the induced increase of proportion branched trichomes. This
indicates that there are fitness benefits associated with the induced increase of total trichome
density.
410
In the event of significant fitness cost, the pattern of selection would reveal fitness peaks in the
absence of herbicide that correspond with the absence of the resistance character in question (*i.e.*
induced total trichomes or induced increase of proportion branched trichomes) (Mauricio 1998).
While many models assume that fitness costs increase linearly with allocation towards defense
(Rhoades 1979, Simms and Rausher 1987), our results revealed no evidence of selection acting
upon any of the induced trichome traits in the absence of herbicide, suggesting a lack of fitness

costs. This finding aligns with several past studies in which costs for induced resistance had not been detected (Brown 1988, Simms 1992, Karban 1993, Mole 1994), but it also conflicts with other work that has identified costs of induced defenses (Tally et al 1999, Heil et al 2000, Cipollini 2002). Some explanations for the lack of costs may be that costs are too small, or that the costs may only be detectable in highly competitive environments (Simms 1992, Karban 1993). Here, we find that for induced trichome density, the net pattern of selection is positive, such that the benefits associated with trichome induction outweigh any cost.

Other constraints on trichome plasticity

Plastic responses are expected to evolve when environmental cues provide a predictable indicator of appropriate adaptive phenotypes that exhibit genetic variation (Padilla and Adolf 1996; Reed et al. 2010). Thus, one constraint on the evolution of plastic defense responses would be the lack of genetic variation underlying such phenotypes. In this work we identified significant maternal line variation for induced total trichome density, as well as induced changes in the proportion of branched and capitate trichomes, indicating that these plastic trichome traits have the potential to evolve. While past work from other researchers has identified genetic variation in density, glandular, and non-glandular trichome types independently (Agrawal et al 2002, Holeski 2007, Ogran et al 2020), our study is one of few designed to detect genetic variation for induced trichome production for multiple trichome traits.

Past work tells us that in some cases induced plant defense can directly hamper reproductive output (Agrawal 1999, Redman 2001), and in other cases can directly hamper plant growth (Siemens 2002, Hermosa et al 2013). As such, we explored the possibility that allocation to induced defenses could come at a cost to plant growth. Following optimal defense theory, we predicted that in the presence of herbicide, selection should favor genotypes that allocated away from growth and toward induced defense, whereas in the absence of herbicide selection should favor genotypes that exhibited high plant growth and low levels of induced defense. We found different responses according to the type of trichome phenotype we considered. In the presence of herbicide, for total trichome density, we identified correlative selection favoring induced increases in the total number of trichomes and intermediate growth rates, indicating that induced trichomes come at the expense of increased growth. In contrast, in the absence of herbicide, we found no evidence of correlative selection acting upon induced total trichome density and growth rate, suggesting that plant growth is not limited in the absence of damage when trichome production is not induced.

In comparison, while we did not identify correlative selection acting on growth rate and either the proportion of branched and capitate trichomes in the presence of herbicide, in the absence of herbicide we found evidence of correlative selection in favor of both high growth rate and high inducibility for these trichome traits. This indicates that genotypes with the highest growth rate and high plastic ability to alter the proportion of trichome types have the highest fitness in the

absence of herbicide. Overall, these results suggest that in the presence of herbicide, genotypes
with intermediate growth and high plasticity in trichome production, but not plasticity in the
proportion of trichome types, exhibit the highest fitness. These selection results align with past
studies that indicate inducibility is negatively associated with species growth (Jacobsen 2022).
Thus, while we did not find costs on induced trichomes in terms of fitness, the pattern of
correlative selection in the presence of herbicide indicates a cost in terms of plant growth.
463
464 *Conclusion*
465
Over the past four decades, much has been learned about the importance of trichomes in
modulating leaf boundary dynamics, and recent developments suggest that we may be
underestimating the eco-evolutionary significance of these plant appendages. As chemical stress
is one of the main global drivers of environmental change, functional trait plasticity may play a
key role in the persistence of plants in novel conditions. The current study examines induced
trichome production as a reflection of the cost-benefit ratio associated with herbicide damage.
Our results provide evidence of a fitness benefit for induced trichome production in the presence
of herbicide, demonstrating that induced responses can serve a stress responsive function against
herbicide. It also provides evidence of evolutionary constraints in terms of correlative selection
acting upon induced trichome production and reduced plant growth, indicating limiting factors
that may impede, but not prevent, adaptive trichome plasticity. Future research should explore
the multi-functionality of various trichome types and the potential for transgenerational trichome
plasticity in ecosystems impacted by chemical stressors.
479

References

Agrawal, A.A., 1998. Induced responses to herbivory and increased plant performance. Science,
279(5354), pp.1201-1202.

Agrawal, A.A., 1999. Induced responses to herbivory in wild radish: effects on several
herbivores and plant fitness. Ecology, 80(5), pp.1713-1723.

Agrawal, A.A., Conner, J.K., Johnson, M.T. and Wallsgrove, R., 2002. Ecological genetics of an
induced plant defense against herbivores: additive genetic variance and costs of
phenotypic plasticity. Evolution, 56(11), pp.2206-2213.

Agrawal, A. A., Hastings, A. P., Johnson, M. T. J., Maron, J. L. , and Salminen, J. P. 2012.
"Insect Herbivores Drive Real-Time Ecological and Evolutionary Change in Plant
Populations." Science 338: 113–6

Agrawal, A. A., and R. Karban. 1999. Why induced defenses may be favored over constitutive
strategies in plants, p. 45-61. In: The ecology and evolution of inducible defenses. R.
Tollrian and C. D. Harvell (eds.). Princeton University Press, Princeton.

Agren, J. and Schemske, D.W., 1993. The cost of defense against herbivores: an experimental
study of trichome production in Brassica rapa. The American Naturalist, 141(2), pp.338-
350.

Anderson, M.P. and Gronwald, J.W., 1991. Atrazine resistance in a velvetleaf (Abutilon
theophrasti) biotype due to enhanced glutathione S-transferase activity. Plant physiology,
96(1), pp.104-109.

Baur, R., Binder, S. and Benz, G., 1991. Nonglandular leaf trichomes as short-term inducible
defense of the grey alder, Alnus incana (L.), against the chrysomelid beetle, Agelastica
alni L. Oecologia, pp.219-226.

Baldwin, I.T. and Schmelz, E.A., 1994. Constraints on an induced defense: the role of leaf area.
Oecologia, 97, pp.424-430

Baldwin, I.T., 1998. Jasmonate-induced responses are costly but benefit plants under attack in
native populations. Proceedings of the National Academy of Sciences, 95(14), pp.8113-
8118.

Barto, E.K. and Cipollini, D., 2005. Testing the optimal defense theory and the growth-
differentiation balance hypothesis in Arabidopsis thaliana. Oecologia, 146, pp.169-178.

Baucom, R.S., 2019. Evolutionary and ecological insights from herbicide-resistant weeds: what
have we learned about plant adaptation, and what is left to uncover?. New Phytologist,
223(1), pp.68-82.

Benbrook, C.M., 2016. Trends in glyphosate herbicide use in the United States and globally.
Environmental Sciences Europe, 28(1), pp.1-15.

Bidart-Bouzat, M.G., Mithen, R. and Berenbaum, M.R., 2005. Elevated CO_2 influences
herbivory-induced defense responses of Arabidopsis thaliana. Oecologia, 145, pp.415-
424.

Bird, S.M. and Gray, J.E., 2003. Signals from the cuticle affect epidermal cell differentiation.
New Phytologist, 157(1), pp.9-23.

Bode, R. F., and Kessler, A. 2012. "Herbivore Pressure on Goldenrod (Solidago altissima L.,
Asteraceae): Its Effects on Herbivore Resistance and Vegetative Reproduction." Journal
of Ecology 100: 795–801.

Boughalleb, F. and Hajlaoui, H., 2011. Physiological and anatomical changes induced by
drought in two olive cultivars (cv Zalmati and Chemlali). Acta Physiologiae Plantarum,
33, pp.53-65.

Braam, J., 2005. In touch: plant responses to mechanical stimuli. New Phytologist, 165(2),
pp.373-389.

Brown, D.G., 1988. The cost of plant defense: an experimental analysis with inducible proteinase
inhibitors in tomato. Oecologia, 76, pp.467-470.

Calvert, H.E., Pence, M.K. and Peters, G.A., 1985. Ultrastructural ontogeny of leaf cavity
trichomes in Azolla implies a functional role in metabolite exchange. Protoplasma,
129(1), pp.10-27.

Chandler, J.M. and Dale, J.E., 1974. Comparative growth of four malvaceous species.
Comparative growth of four malvaceous species., pp.116-117.

Cipollini, D.F., 2002. Does competition magnify the fitness costs of induced responses in
Arabidopsis thaliana? A manipulative approach. Oecologia, 131, pp.514-520

Coverdale, Tyler C., and Agrawal, A.A. 2022. " Experimental Insect Suppression Causes Loss of
    Induced, But Not Constitutive, Resistance in Solanum carolinense." Ecology 103(11):
    e3786.

Dalin, P., Ågren, J., Björkman, C., Huttunen, P. and Kärkkäinen, K., 2008. Leaf trichome
    formation and plant resistance to herbivory. Induced plant resistance to herbivory, pp.89-
    105.

Dalin, P. and Björkman, C., 2003. Adult beetle grazing induces willow trichome defence against
    subsequent larval feeding. Oecologia, pp.112-118

De Snoo, G.R. and Van der Poll, R.J., 1999. Effect of herbicide drift on adjacent boundary
    vegetation. Agriculture, ecosystems & environment, 73(1), pp.1-6.

Devine, M., Duke, S.O. and Fedtke, C., 1992. Physiology of herbicide action. PTR Prentice Hall.

Dewar, A.M., Haylock, L.A., Bean, K.M. and May, M.J., 2000. Delayed control of weeds in
    glyphosate-tolerant sugar beet and the consequences on aphid infestation and yield. Pest
    Management Science: formerly Pesticide Science, 56(4), pp.345-350.

Doss, R.P., Shanks JR, C.H., Chamberlain, J.D. and Garth, J.K., 1987. Role of leaf hairs in
    resistance of a clone of beach strawberry, Fragaria chiloensis, to feeding by adult black
    vine weevil, Otiorhynchus sulcatus (Coleoptera: Curculionidae). Environmental
    Entomology, 16(3), pp.764-768.

Egan, J.F. and Mortensen, D.A., 2012. Quantifying vapor drift of dicamba herbicides applied to
    soybean. Environmental toxicology and chemistry

Ehleringer, J., 1982. The influence of water stress and temperature on leaf pubescence
    development in Encelia farinosa. American Journal of Botany, 69(5), pp.670-675.

Gavrilets, S. and Scheiner, S.M., 1993. The genetics of phenotypic plasticity. VI. Theoretical
    predictions for directional selection. Journal of evolutionary biology, 6(1), pp.49-68.

Gibert, J.P., Allen, R.L., Hruska III, R.J. and DeLong, J.P., 2017. The ecological consequences
    of environmentally induced phenotypic changes. Ecology Letters, 20(8), pp.997-1003.

Gonzales, W.L., Negritto, M.A., Suarez, L.H. and Gianoli, E., 2008. Induction of glandular and
non-glandular trichomes by damage in leaves of Madia sativa under contrasting water
regimes. Acta oecologica, 33(1), pp.128-132.

Grether, G.F., 2005. Environmental change, phenotypic plasticity, and genetic compensation.
The American Naturalist, 166(4), pp.E115-E123.

Gutiérrez-Alcalá, G., Gotor, C., Meyer, A.J., Fricker, M., Vega, J.M. and Romero, L.C., 2000.
Glutathione biosynthesis in Arabidopsis trichome cells. Proceedings of the National
Academy of Sciences, 97(20), pp.11108-11113.

Haberlandt, G., 1914. Physiological plant anatomy. Macmillan and Company, limited.

Hahn, P. G., and Maron, J. L. 2016. "A Framework for Predicting Intraspecific Variation in Plant
Defense." Trends in Ecology and Evolution 31: 646–56.

Hamann, E., Blevins, C., Franks, S.J., Jameel, M.I. and Anderson, J.T., 2021. Climate change
alters plant–herbivore interactions. New Phytologist, 229(4), pp.1894-1910.

Hartgerink, A.P. and Bazzaz, F.A., 1984. Seedling-scale environmental heterogeneity influences
individual fitness and population structure. Ecology, 65(1), pp.198-206.

Hartzler, R.G. and Battles, B.A., 2001. Reduced fitness of velvetleaf (Abutilon theophrasti)
surviving glyphosate. Weed Technology, 15(3), pp.492-496.

Hwang, J.I., Norsworthy, J.K., Houston, M.M., Piveta, L.B., Priess, G.L., Zaccaro-Gruener,
M.L., Barber, L.T. and Butts, T.R., 2022. Large-scale evaluation of physical drift and
volatility of 2, 4-D choline in cotton: a four-year field study. Pest Management Science,
78(8), pp.3337-3344.

Heil, M., Hilpert, A., Kaiser, W. and Linsenmair, K.E., 2000. Reduced growth and seed set
following chemical induction of pathogen defence: does systemic acquired resistance
(SAR) incur allocation costs?. Journal of Ecology, 88(4), pp.645-654.

Hermosa, R., Rubio, M.B., Cardoza, R.E., Nicolás, C., Monte, E. and Gutiérrez, S., 2013. The
contribution of Trichoderma to balancing the costs of plant growth and defense. Int.
Microbiol, 16(2), pp.69-80.

Herms, D.A. and Mattson, W.J., 1992. The dilemma of plants: to grow or defend. The quarterly
review of biology, 67(3), pp.283-335.

Höglund, S. and Larsson, S., 2005. Abiotic induction of susceptibility in insect-resistant willow.
Entomologia Experimentalis et Applicata, 115(1), pp.89-96.

Holeski, L.M., 2007. Within and between generation phenotypic plasticity in trichome density of
Mimulus guttatus. Journal of evolutionary biology, 20(6), pp.2092-2100.

Jacobsen, D.J., 2022. Growth rate and life history shape plant resistance to herbivores. American
Journal of Botany, 109(7), pp.1074-1084.

Johnson, N.M. and Baucom, R.S., 2024. The double life of trichomes: understanding their dual
role in herbivory and herbicide resistance. Evolution, 78(6), pp.1121-1132.

Karban, R., 1993. Costs and benefits of induced resistance and plant density for a native shrub,
Gossypium thurberi. Ecology, 74(1), pp.9-19.

Karban, R., Agrawal, A.A., Thaler, J.S. and Adler, L.S., 1999. Induced plant responses and
information content about risk of herbivory. Trends in Ecology & Evolution, 14(11),
pp.443-447.

Karban, R. and Baldwin, I.T., 2007. Induced responses to herbivory. University of Chicago
Press.

Karowe, D.N. and Grubb, C., 2011. Elevated CO 2 increases constitutive phenolics and
trichomes, but decreases inducibility of phenolics in Brassica rapa (Brassicaceae). Journal
of chemical ecology, 37, pp.1332-1340.

Kleijn, D. and Snoeijing, G.I.J., 1997. Field boundary vegetation and the effects of agrochemical
drift: botanical change caused by low levels of herbicide and fertilizer. Journal of Applied
Ecology, pp.1413-1425.

Kirik, V., Simon, M., Huelskamp, M. and Schiefelbein, J., 2004. The ENHANCER OF TRY
AND CPC1 gene acts redundantly with TRIPTYCHON and CAPRICE in trichome and
root hair cell patterning in Arabidopsis. Developmental biology, 268(2), pp.506-513.

Kuznetsova, A., Brockhoff, P.B. and Christensen, R.H., 2017. lmerTest package: tests in linear
mixed effects models. Journal of statistical software, 82, pp.1-26.

Li, C., Mo, Y., Wang, N., Xing, L., Qu, Y., Chen, Y., ... & Kopittke, P. M. (2023). The
overlooked functions of trichomes: Water absorption and metal detoxication. Plant, Cell
& Environment, 46(3), 669-687.

Marrs, R.H., Williams, C.T., Frost, A.J. and Plant, R.A., 1989. Assessment of the effects of herbicide spray drift on a range of plant species of conservation interest. Environmental Pollution, 59(1), pp.71-86.

Matsumura, M., Nomoto, M., Itaya, T., Aratani, Y., Iwamoto, M., Matsuura, T., Hayashi, Y., Mori, T., Skelly, M.J., Yamamoto, Y.Y. and Kinoshita, T., 2022. Mechanosensory trichome cells evoke a mechanical stimuli–induced immune response in Arabidopsis thaliana. Nature communications, 13(1), p.1216.

Mauricio, R., 1998. Costs of resistance to natural enemies in field populations of the annual plant Arabidopsis thaliana. The American Naturalist, 151(1), pp.20-28.

McKey, D., 1974. Adaptive patterns in alkaloid physiology. The American Naturalist, 108(961), pp.305-320.

Milewski, A.V., Young, T.P. and Madden, D., 1991. Thorns as induced defenses: experimental evidence. Oecologia, pp.70-75.

Mole, S., 1994. Trade-offs and constraints in plant-herbivore defense theory: a life-history perspective. Oikos, pp.3-12

Morris, J.B. and Wang, M.L., 2013. Anthocyanin indexes, quercetin, kaempferol and myricetin concentration in leaves and fruit of Abutilon theophrasti Medik. genetic resources. Plant Genetic Resources, 11(1), pp.87-89.

Ogran, A., Conner, J., Agrawal, A.A. and Barazani, O., 2020. Evolution of phenotypic plasticity: Genetic differentiation and additive genetic variation for induced plant defence in wild arugula Eruca sativa. Journal of evolutionary biology, 33(2), pp.237-246

Parmesan, C., 2006. Ecological and evolutionary responses to recent climate change. Annu. Rev. Ecol. Evol. Syst., 37, pp.637-669.

Padilla, D.K. and Adolph, S.C., 1996. Plastic inducible morphologies are not always adaptive: the importance of time delays in a stochastic environment. Evolutionary Ecology, 10, pp.105-117.

Peiffer, M., Tooker, J.F., Luthe, D.S. and Felton, G.W., 2009. Plants on early alert: glandular trichomes as sensors for insect herbivores. New Phytologist, 184(3), pp.644-656.

Pérez-Estrada, L.B., Cano-Santana, Z. and Oyama, K., 2000. Variation in leaf trichomes of
Wigandia urens: environmental factors and physiological consequences. Tree Physiology,
20(9), pp.629-632.

Pimm, S.L., 2009. Climate disruption and biodiversity. Current biology, 19(14), pp.R595-R601.

Poelman, E.H. and Kessler, A., 2016. Keystone herbivores and the evolution of plant defenses.
Trends in plant science, 21(6), pp.477-485.

R Core Team, 2022. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Rautio, P., Markkola, A., Martel, J., Tuomi, J., Härmä, E., Kuikka, K., Siitonen, A., Riesco, I.L.
and Roitto, M., 2002. Developmental plasticity in birch leaves: defoliation causes a shift
from glandular to nonglandular trichomes. Oikos, 98(3), pp.437-446.

Redman, A.M., Cipollini, D.F. and Schultz, J.C., 2001. Fitness costs of jasmonic acid-induced
defense in tomato, Lycopersicon esculentum. Oecologia, 126, pp.380-385.

Reed, T.E., Wanless, S., Harris, M.P., Frederiksen, M., Kruuk, L.E. and Cunningham, E.J., 2006.
Responding to environmental change: plastic responses vary little in a synchronous
breeder. Proceedings of the Royal Society B: Biological Sciences, 273(1602), pp.2713-
2719.

Rhoades, D.F., 1979. Evolution of plant chemical defenses against herbivores. Herbivores-Their
Interaction with Secondary Plant Metabolites, pp.3-48.

Ritter, R.L., 1986. Triazine resistant velvetleaf and giant foxtail control in no-tillage corn.
Triazine resistant velvetleaf and giant foxtail control in no-tillage corn., pp.50-52.

Shibuya, T., Itagaki, K., Ueyama, S., Hirai, N. and Endo, R., 2016. Atmospheric humidity
influences oviposition rate of Tetranychus urticae (Acari: Tetranychidae) through
morphological responses of host Cucumis sativus leaves. Journal of economic
entomology, 109(1), pp.255-258.

Siemens, D.H., Garner, S.H., Mitchell-Olds, T. and Callaway, R.M., 2002. Cost of defense in the
context of plant competition: Brassica rapa may grow and defend. Ecology, 83(2),
pp.505-517.

Simms, E.L., 1992. to Herbivory. Plant resistance to herbivores and pathogens: ecology, evolution, and genetics, p.392.

Simms, E.L. and Rausher, M.D., 1987. Costs and benefits of plant resistance to herbivory. The American Naturalist, 130(4), pp.570-581.

Spencer, N.R., 1984. Velvetleaf, Abutilon theophrasti (Malvaceae), history and economic impact in the United States. Economic Botany, 38(4), pp.407-416.

Sterling, T.M., Houtz, R.L. and Putnam, A.R., 1987. Phytotoxic exudates from velvetleaf (Abutilon theophrasti) glandular trichomes. American Journal of Botany, 74(4), pp.543-550.

Sterling, T.M. and Putnam, A.R., 1987. Possible role of glandular trichome exudates in interference by velvetleaf (Abutilon theophrasti). Weed Science, 35(3), pp.308-314.

Sultan, S.E. and Stearns, S.C., 2005. Environmentally contingent variation: phenotypic plasticity and norms of reaction. Variation, pp.303-332.

Tally, A., Oostendorp, M., Lawton, K., Staub, T. and Bassi, B., 1999. Commercial development of elicitors of induced resistance to pathogens. Induced plant defenses against pathogens and herbivores: biochemistry, ecology, and agriculture., pp.357-369.

Tattini, M., Gravano, E., Pinelli, P., Mulinacci, N. and Romani, A., 2000. Flavonoids accumulate in leaves and glandular trichomes of Phillyrea latifolia exposed to excess solar radiation. The New Phytologist, 148(1), pp.69-77

Textor, S. and Gershenzon, J., 2009. Herbivore induction of the glucosinolate–myrosinase defense system: major trends, biochemical bases and ecological significance. Phytochemistry Reviews, 8, pp.149-170.

Tollrian, R. and Harvell, C.D. eds., 1999. The ecology and evolution of inducible defenses. Princeton university press.

Valladares, F., Gianoli, E. and Gómez, J.M., 2007. Ecological limits to plant phenotypic plasticity. New phytologist, 176(4), pp.749-763.

Vanhoutte, B., Schenkels, L., Ceusters, J. and De Proft, M.P., 2017. Water and nutrient uptake in Vriesea cultivars: trichomes vs. roots. Environmental and Experimental Botany, 136, pp.21-30

Warwick, S.I. and Black, L.D., 1985. Genecological variation in recently established populations of Abutilon theophrasti (velvetleaf). Canadian Journal of Botany, 64(8), pp.1632-1643.

Warwick, S.I. and Black, L.D., 1988. THE BIOLOGY OF CANADIAN WEEDS.: 90. Abutilon theophrasti. Canadian Journal of Plant Science, 68(4), pp.1069-1085.

Westerbergh, A. and Nyberg, A.B., 1995. Selective grazing of hairless Silene dioica plants by land gastropods. Oikos, pp.289-298.

Winter, D.M., 1960. The development of the seed of Abutilon theophrasti I. Ovule and embryo. American journal of botany, 47(1), pp.8-14.

Wu, J.C., Xu, J.X., Liu, J.L., Yuan, S.Z., Cheng, J.A. and Heong, K.L., 2001. Effects of herbicides on rice resistance and on multiplication and feeding of brown planthopper (BPH), Nilaparvata lugens (Stål)(Homoptera: Delphacidae). International Journal of Pest Management, 47(2), pp.153-159.

Yamasaki, S., Noguchi, N. and Mimaki, K., 2007. Continuous UV-B irradiation induces morphological changes and the accumulation of polyphenolic compounds on the surface of cucumber cotyledons. Journal of Radiation Research, 48(6), pp.443-454.

Young, T.P. and Okello, B.D., 1998. Relaxation of an induced defense after exclusion of herbivores: spines on Acacia drepanolobium. Oecologia, 115, pp.508-513

Zangerl, A.R. and Bazzaz, F.A., 1992. Defense Allocation. Plant resistance to herbivores and pathogens: ecology, evolution, and genetics, p.363.

Figures

Figure 1. Trichome structures present on the adaxial surface of *Abutilon theophrasti* leaves using

confocal imaging (x10). A) Branched trichomes are unicellular structures with 4-8 appendages.

B) Single trichomes are unicellular unbranched structures. C) Peltate trichomes are multicellular

glandular structures made up of 4-5 cells. D) Capitate trichomes are multicellular glandular

structures made up of 12-15 cells. Density is calculated as the total number of all trichomes

captured in the image.

844
845
846 Figure 2. Treatment effects observed on (**a**) trichome density (F = 15.81, p = 0.02) (**b**) the
847 proportion of single trichomes (F = 0.21, p = 0.65), (**c**) the proportion of capitate trichomes (F =
848 1.11, p = 0.29), (**d**) the proportion of peltate trichomes (F = 0.20, p = 0.65), and (**e**) the
849 proportion of branched trichomes (F = 0.87, p = 0.35) in *Abutilon theophrasti*.

857   Figure 3. Illustration of genetic variation for induced trichome phenotypes in *Abutilon*

858   *theophrasti* maternal lines grown in field conditions in either the control (non-herbicide

859   environment) or herbicide present environment. Shown are the significant maternal line by

860   treatment effects for the proportion branched trichomes ($\chi^2 = 3.92$, p = 0.048), the proportion

861   capitate trichomes ($\chi^2 = 5.64$, p = 0.018), and total trichome density ($\chi^2 = 10.05$, p = 0.002).

Figure 4. Relationships between herbicide resistance and (**a**) induced proportion branched and (**b**) induced proportion capitate and (**c**) induced total trichome density for *Abutilon theophrasti*. Shown are Pearson's correlation coefficients; significant correlations are indicated in bold. Data points represent maternal line means. Induced trichome traits were measured as trichomes damaged – trichome control, and herbicide resistance was measured as 1 – proportion of leaf yellowing.

Figure 5. The relationship between relative fitness and (**a**) induced proportion branched ($S =$

0.10, p = 0.925) and (**b**) induced proportion capitate ($S =$ -2.13, p = 0.045) and (**c**) induced total

trichome density ($S =$ 2.56, p = 0.018) in *Abutilon theophrasti*. Solid lines represent selection

differentials in the presence of herbicide. Data points represent maternal line means; induced

trichome traits were measured as damaged – control states, and herbicide resistance was

measured as 1 – proportion of leaf yellowing.

880
881 Figure 6. Fitness surface for correlative selection acting upon *Abutilon theophrasti* induced total
882 trichome density and growth rate in the presence of herbicide ($\gamma = 5.71$, p = 0.010), tested using
883 non-linear selection gradients (multivariate selection). Relative fitness is depicted by the color
884 gradient; yellow is the highest fitness, blue is the lowest fitness, and green is intermediate.

885 Tables
886

	Maternal Line x Treatment Effects		
	$\chi 2$	df	p - values
Proportion Branched	**3.92**	**1**	**0.048**
Proportion Single	3.26	1	0.071
Proportion Capitate	**5.64**	**1**	**0.018**
Proportion Peltate	0.71	1	0.399
Total Density	**10.05**	**1**	**0.002**

888 Table 1. Results from the test for genetic variation in *Abutilon theophrasti* showing the effects of
889 maternal line by treatment on trichome traits (proportion branched, single, capitate, peltate, and
890 total density) captured in the field. Significant effects are indicated in boldface. The full
891 ANOVAs are presented in STable 1.

892   Supplementary Figures and Tables
893

894
895   SFigure 1. Locations of *Abutilon theophrasti* populations sampled and used for this study.

	Treatment (df =1)		Maternal Line (df =1)		Maternal Line x Treatment (df = 1)	
Trait	F	p-value	χ^2	p-value	χ^2	p-value
Proportion Branched	1.56	0.221	0.34	0.056	3.92*	0.048*
Proportion Capitate	1.94	0.168	0.00	1.000	5.64*	0.018*
Proportion Single	0.35	0.561	0.23	0.639	3.26	0.071
Proportion Peltate	0.44	0.512	0.08	0.767	0.71	0.399
Total Density	3.22	0.079	0.00	1.000	10.05**	0.002**

897
898   STable 1. Influence of herbicide treatment and maternal line on trichome traits (proportion
899   branched, capitate, single, peltate, and total density) when observed in the field. Significant
900   effects are indicated with asterisks: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Phenotypic Selection (*S*) Differentials						
	Control			Herbicide		
Trait	*S*	SE	p-value	*S*	SE	p-value
Induced Branched	-0.22	0.15	0.826	0.10	0.47	0.925
Induced Capitate	-0.38	0.04	0.707	-2.13*	0.11*	0.045*
Total Induced Density	0.50	0.09	0.625	2.56*	0.37*	0.018*

902
903
904   STable 2. Total selection (*S*) on *Abutilon theophrasti* induced trichome traits (proportion
905   branched, proportion capitate, and total density) in the absence and presence of herbicide. Shown
906   are selection differential values, standard errors, and p-values for traits in each treatment.
907   Significant effects are indicated with asterisks: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Trait	Non-linear Selection (γ) Gradients					
	Control			Herbicide		
	γ	SE	p-value	γ	SE	p-value
Induced Branched x Growth Rate	6.13**	0.37**	0.006**	-0.77	2.08	0.070
Induced Capitate x Growth Rate	3.97	1.08	0.062	-3.58	5.33	0.091
Induced Density x Growth Rate	-0.87	0.01	0.669	5.71*	0.05*	0.010*

910 STable 3. Correlative selection (γ) on *Abutilon theophrasti* induced trichome traits (proportion
911 branched, proportion capitate, and total density) and growth rate in the absence and presence of
912 herbicide. Shown are selection differential values, standard errors, and p-values for traits in each
913 treatment. Significant effects are indicated with asterisks: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.
914