A Modified BPaL Regimen for Tuberculosis Treatment replaces Linezolid with Inhaled Spectinamides

Malik Zohaib Ali [1,2,6], Taru S. Dutt [1,2], Amy MacNeill [2], Amanda Walz [1,2], Johnathan Patterson[1,2], Camron Pearce [1,2,6], Ha Lam[1,2], Marcela Henao-Tamayo[2], Richard E. Lee [3], Jiuyu Liu [3], Gregory T. Robertson [1,2], Anthony J. Hickey [4], Bernd Meibohm [5], Mercedes Gonzalez-Juarrero [1,2]

[1]Mycobacteria Research Laboratories, Colorado State University, 80523, USA

[2] Microbiology, Immunology and Pathology, Colorado State University, 80523, USA

[3] Department of Chemical Biology and Therapeutics, St. Jude Children's Research Hospital, 38105, USA

[4]Technology Advancement and Commercialization, RTI International, 27709, USA

[5] Department of Pharmaceutical Sciences, University of Tennessee Health Science Center, 38163, USA

[6] Program in Cell & Molecular Biology, Colorado State University, 80523, USA

Keywords: mycobacterium tuberculosis, inhalation, spectinamides , BPaL, linezolid, MDR-TB, XDR-TB, therapy, multidrug regimen , C3HeB/FeJ mice , Balb/c mice , 1599

Corresponding Author Information: Mercedes.gonzalez-juarrero@colostate.edu

Abstract

The Nix-TB clinical trial evaluated a new 6-month regimen containing three-oral-drugs; bedaquiline (B), pretomanid (Pa) and linezolid (L) (BPaL regimen) for treatment of tuberculosis (TB). This regimen achieved remarkable results as almost 90% of the multidrug resistant (MDR) or extensively drug resistant (XDR) TB participants were cured but many patients also developed severe adverse effects (AEs). The AEs were associated with the long-term administration of the protein synthesis inhibitor linezolid. Spectinamide 1599 (S) is also a protein synthesis inhibitor of *Mycobacterium tuberculosis* with an excellent safety profile but which lacks oral bioavailability. Here we hypothesize that inhaled spectinamide 1599, combined with BPa ––BPaS regimen––has similar efficacy to that of BPaL regimen while simultaneously avoiding the L-associated AEs. The BPaL and BPaS regimens were compared in the Balb/c and C3HeB/FeJ murine chronic TB efficacy models. After 4-weeks of treatment, both regimens promoted equivalent bactericidal effect in both TB murine models. However, treatment with BPaL resulted in significant weight loss and the complete blood count suggested development of anemia. These effects were not similarly observed in mice treated with BPaS. BPaL treatment also decreased myeloid to erythroid ratio and increased concentration of proinflammatory cytokines in bone marrow compared to mice receiving BPaS regimen. During therapy both regimens improved the lung lesion burden, reduced neutrophil and cytotoxic T cells counts while increased the number of B and helper and regulatory T cells. These combined data suggest that inhaled spectinamide 1599 combined with BPa is an effective TB regimen that avoids L-associated AEs.

IMPORTANCE Tuberculosis (TB) is an airborne infectious disease that spreads via aerosols containing *Mycobacterium tuberculosis* (Mtb), the causative agent of TB. TB can be cured by administration of 3-4 drugs for 6-9 months but there are limited treatment options for patients infected with multidrug (MDR) and extensively resistant (XDR) strains of Mtb. BPaL is a new all-oral combination of drugs consisting of Bedaquiline (B), Pretomanid (Pa) and Linezolid (L). This regimen was able to cure ~90% of MDR and XDR TB patients in clinical trials but many patients developed severe adverse effects (AEs) associated to the long-term administration of linezolid. We evaluated a new regimen in which Linezolid in the BPaL regimen was replaced with inhaled spectinamide 1599. In the current study, we demonstrate that 4-weeks of treatment with inhaled spectinamide 1599 in combination with Bedaquiline and Pretomanid has equivalent efficacy to the BPaL drug combination and avoids the L-associated-AEs.

Introduction

Tuberculosis (TB) remains one of the leading causes of death initiated by an infectious agent. In 2021, the World Health Organization reported 10.6 million new TB cases worldwide and among those, 450,000 cases were also diagnosed as multidrug resistant (MDR) or extensively drug resistant (XDR) TB (1). Treatment of MDR- and XDR-TB patients is lengthy and is often poorly tolerated due to significant associated side effects (2).

Recently, a 6-month novel treatment regimen of three-oral-drugs: bedaquiline (B), pretomanid (Pa) and linezolid (L) referred to as the BPaL regimen was approved. Preclinical studies demonstrated the better efficacy of BPaL regimen for drug sensitive TB compared to the standard TB chemotherapy (1,3,4) and thereafter, the BPaL regimen was tested in the Nix-TB clinical trial conducted in South Africa (5). This trial enrolled patients with XDR-TB and treatment-intolerant or non-responsive MDR-TB, including HIV positive patients with a CD4 count of 50 or higher. The results were remarkable as 95 out of 107 patients were cured though many patients had a high rate of treatment-associated adverse effects (AEs).

The long-term administration of linezolid (an oxazolidinone antibiotic) was likely the causative agent resulting in bone marrow myelosuppression (48%), peripheral neuropathy, optic neuritis (81%) and anemia (37%) in patients treated with the BPaL regimen (5). Subsequently, the ZeNix trial adjusted the BPaL regimen to a linezolid dose of 600mg. This trial also had remarkable results; it cured 84-91% of patients (9 to 26 weeks of therapy, respectively) and resulted in fewer AEs than those observed in the Nix-TB trial (6). Apart from AEs, several studies have also raised awareness of high doses of

linezolid leading to development of linezolid-resistant *Mycobacterium tuberculosis* (Mtb)

(7–9). At present, the TB-drug development field is working to modify the BPaL regimen

to maintain or improve its efficacy while diminishing treatment-associated AEs (10).

Spectinomycin, an aminocyclitol antibiotic, is a broad spectrum antibiotic used

mainly for the treatment of *Neisseria gonorrhoeae* (11). It inhibits bacterial protein

synthesis (12) and has a very high safety profile with no known ototoxicity and

nephrotoxicity (13,14). The efficacy of spectinomycin against Mtb is very poor but its

structural modification led to the development of a new series of semisynthetic analogs

called spectinamides (12,15,16). Spectinamides bind selectively with the bacterial 30S

ribosomes and importantly, unlike linezolid, they do not bind to the mitochondrial 30S in

mammalians cells. The latter represents a great advantage for reduced potential for side

effects, such as ototoxicity and myeloid suppression that are commonly associated with

other protein synthesis inhibitors such as amikacin and linezolid, respectively (17,18).

Spectinamides' potent anti-tubercular activity is attributed to its ability to evade drug efflux

by Rv1258c major facilitator superfamily transporter present on the surface of Mtb

(19,20). Spectinamides have shown excellent activity against MDR- and XDR-Mtb strains

(16,21). However, their poor oral availability have limited their usage to injectable forms.

One of the lead spectinamides, 1599, has demonstrated promising results *in vitro*

and *in vivo* and was shown to lack cross-resistance with existing anti-TB drugs (16,19–

23). 1599, delivered subcutaneously, proved an effective partner agent when combined

with rifampin and pyrazinamide and also with bedaquiline, pretomanid or moxifloxacin in

TB mouse efficacy models of increasing complexity (21).

One of the limitations of using 1599 as injectable is a potential risk for poor patient

compliance (24) and direct administration of aerosolized antibiotics to the lungs has been

studied for decades as an alternative to systemic drug administration via injection.

Aerosolized administration of 1599 has been tested in preclinical *in vivo* studies using

liquid formulation of the drug. These studies have shown that inhaled 1599, used in

monotherapy or in combination with pyrazinamide, is efficacious and well tolerated in

murine TB efficacy models (25,26). A comparative study assessing biodistribution of the

drug in relation to the administration route demonstrated that 1599 showed 48 times

higher exposure in mouse lungs via inhalation compared to equivalent dosages

administered by subcutaneous injection; the latter may explain the increased efficacy of

this drug following intrapulmonary aerosol (22,26). Moreover, 1599 was shown to be

amenable to dry powder formulation and delivery, suggesting a pathway to a more

patient-friendly delivery system (27). Therefore, in this study, we hypothesized that

combining BPa with inhaled S will maintain equivalent efficacy to the BPaL regimen while

avoiding the accompanying toxicities that occurred with long-term BPaL administration to

human MDR/XDR-TB patients.

Results

Linezolid and spectinamide 1599 show similar efficacy in monotherapy

To compare the efficacy of L or S in monotherapy, Mtb-infected C3HeB/FeJ (n=7)

and Balb/c (n=4-6) mice received L (administered 5/7 days per week orally at 100 mg/Kg)

or S (administered 3/7 days per week on alternate days via intrapulmonary aerosol

delivery at 100 mg/Kg and 50 mg/Kg, respectively) for 4 weeks. At the end of treatment,

the animals were sacrificed, and the CFU in lungs and spleen was enumerated (Figure

1A-D). Treatment of C3HeB/FeJ mice with L (7.30 ± 0.45 $\log_{10}$) or S (6.90 ± 0.48 $\log_{10}$)

for 4 weeks resulted in an average of 0.51 and 0.91 $\log_{10}$ CFU reduction in the lungs

respectively, compared to untreated (UnRx) control (7.81 ± 0.36 $\log_{10}$) (Figure 1A). This

difference failed to achieve statistical significance owing to the larger standard deviation

associated with heterogenous advanced lung pathology observed in this model

(26,28,29). In contrast, L (4.49 ± 0.10 $\log_{10}$) treated C3HeB/FeJ mice had significantly

lower spleen bacterial burden compared to UnRx (5.43 ± 0.27 $\log_{10}$), with no significant

difference between the L and S (4.94 ± 0.11 $\log_{10}$) treatment arm (Figure 1C). In Balb/c

mice, L or S treatment was found to promote significant reduction in lung bacterial burden

(0.83 and 0.77 $\log_{10}$ respectively) compared to UnRx or the vehicle only controls (Figure

1B). There was no significant difference in lung bacterial burden after L or S treatment in

Balb/c mice and there was no change in spleen bacterial burden compared to UnRx

control (Figure 1D).

Combination therapy with BPaL or BPaS have similar efficacy

We further tested and compared the efficacy of L or S when used in combination

therapy with BPa. Mtb-infected C3HeB/FeJ mice were treated with either BPaL (B=

25mpk; Pa= 100 mg/Kg and L= 100 mg/Kg all administered 5/7 a week via oral gavage)

or BPaS (BPa as in BPaL and S= 100 mg/Kg administered 3/7 a week on alternate days

via intrapulmonary aerosol delivery) for 4 weeks. The comparative analysis from

combined data of three independent studies is shown in Figure 1E&G (data from

individual studies are shown in Figures S1-3). Two of three studies contained an extra

group of BPa treated mice as a reference control. Compared to UnRx (7.48 ± 0.12 $\log_{10}$)

control, mice in the BPa (5.49 ± 0.42 $\log_{10}$), BPaL (4.76 ± 0.38 $\log_{10}$) and BPaS (4.98 ±

0.35 $\log_{10}$) treatment groups had significantly reduced the lung bacterial burden by 1.99,

2.72, and 2.50 $\log_{10}$ respectively (Figure 1E). Although a higher CFU reduction was

observed in the lungs of C3HeB/FeJ mice treated with BPaL or BPaS, these differences

failed to achieve statistical significance compared to BPa backbone regimen. All three

regimens proved highly effective at reducing spleen bacterial burden in C3HeB/FeJ mice,

with most mice returning no CFU within the limit of detection (LOD) employed herein

(Figure 1G).

The effect of BPaL and BPaS (S= 50 mg/Kg) combination therapy on the bacterial

burden in lungs and spleen of Mtb-infected Balb/c mice was determined at the end of 4-

weeks of treatment. Figure 1F&H show the combined lung and spleen CFU data from two

independent studies (data for each study is shown in Figures S7-8). The combined result

demonstrated that compared to UnRx (5.24 ± 0.17 $\log_{10}$) control, mice in the BPaL (1.29

± 0.13 $\log_{10}$) and BPaS (1.44 ± 0.17 $\log_{10}$) treatment groups returned significantly fewer

CFU in the lungs, with most mice returning no CFU within the LOD employed herein

(Figure 1F). As in C3HeB/FeJ TB model, no significant difference was observed in the

lung CFU of Balb/c mice treated with either the BPaL or BPaS regimen. BPaL and BPaS

therapy reduced Balb/c spleen bacterial burden to below the LOD of the assay with no

CFU recovered for any treated mice (Figure 1H). In summary, these results support our

hypothesis and demonstrate that both BPaL and BPaS multidrug regimens show

equivalent bactericidal effects in C3HeB/FeJ and Balb/c chronic TB efficacy models.

Monitoring of adverse effects

Five approaches were employed to monitor treatment-associated AEs in mice in this study including 1) changes in the live body weight of mice; 2) lung histopathology and lesion scoring 3) evaluation of complete blood count (CBC), 4) clinical pathology to study myelosuppression in bone marrow and 5) changes in content of immune cells in the lungs, spleen, bone marrow and blood.

BPaL therapy decreases the live body weight of mice

No significant difference in the live body weight among the treatment groups in either C3HeB/FeJ or Balb/c mice was observed following 4-weeks of treatment with S or L alone (Figure 2A&B). On the other hand, C3HeB/FeJ or Balb/c mice treated with BPa, BPaL or BPaS showed marginal loses of live body weight, ranging from 2.37-5.13% and only mice receiving the BPaL regimen, when compared to the UnRx control, reached statistically significant loses in live body weight by the end of treatment (Figure 2C&D).

BPaL and BPaS therapy result in significant lower lung lesion burden

The lesions in Mtb-infected UnRx C3HeB/FeJ mice (Figure 2E) showed a spectrum of diverse granuloma types ranging from aggregations of macrophages and lymphocytes to highly organized granulomas with collagen encapsulation and a region of central caseous necrosis that resembles to those found in some human patients (28,29). By comparison, lung lesions of Mtb-infected Balb/c mice (Figure 2G) consisted of granulomas with a very homogeneous structure (28) formed also by aggregations of macrophages and lymphocytes without a necrotic core. Mice treated for 4 weeks with

the BPaL or BPaS regimen presented with significant reduction in the number and size

of TB granulomas in both C3HeB/FeJ and Balb/c TB models compared to their

respective UnRx control, with no significant difference in lesion burden score between

the combination drug treatment groups (Figure 2F&H).

Association of L with altered blood profile and mild anemia in mice

Complete blood count (CBC) profile was performed on Mtb-infected C3HeB/FeJ

mice at the end of 4-weeks of treatment to quantify treatment-associated hematological

changes. The results obtained from mice treated with L or S alone are summarized in

Figure 3A. Compared to UnRx control and S treated mice, L treatment significantly

increased the red blood cell distribution width standard deviation (RDWs), while both L

and S treatment were associated with a significant decrease in the mean corpuscular

hemoglobin concentration (MCHC) compared to UnRx control (Figure 3A).

The effect of combination therapy with the BPaL or BPaS regimen on CBC profile

was analyzed at 1-, 2- and 4-weeks of treatment (Figure 3B). None of the 20 parameters

of CBC changed during the first 2 weeks of treatment. However, L-containing BPaL

regimen was again associated with a significant increase in the RDWs, and lower

hemoglobin (HGB) compared to UnRx control after 4 weeks. This effect was not observed

in mice treated with BPa or BPaS (Figure 3B). However, as in monotherapy, there was a

trend towards lower overall MCHC in mice treated with either BPaL or BPaS. The mean

platelet volume (MPV) was marginally higher at 4 weeks in mice treated with BPa

compared to BPaL or BPaS, albeit not significantly different from UnRx control (Figure

3B). Given that no difference in HGB or RDWs was observed between UnRx control and

their comparator BPa and BPaS regimens, we concluded that the significant HGB

decrease and RDWs increase (often observed during development of anemia) were

associated with inclusion of L in BPaL regimen.

Spectinamide 1599 recovers the altered ratio of myeloid to erythroid cells in bone

marrow

To further evaluate if L was associated with myelosuppressive effect, we

performed hematopathology analysis on bone marrow from Mtb-infected C3HeB/FeJ and

Balb/c mice at the end of treatment. For C3HeB/FeJ mice, the number of myeloid (M) and

erythroid (E) precursor cells were calculated from H&E stained sections and their myeloid

to erythroid ratio (M:E) was determined by counting 300 bone marrow cells in 5 different

regions (Figure 3C). The BPa and BPaL treatment significantly decreased myeloid cells

while increasing the proportion of erythroid cells (Figure 3D). Hence, the corresponding

ratio in bone marrow of animals treated with BPaL or BPa was lower compared to UnRx

control. Importantly, the BPaS treatment did not show any difference in the content of

myeloid or erythroid cells when compared to UnRx control. In Mtb-infected Balb/c mice,

the number of myeloid and other cell types were counted, however, no significant

difference was found among the control and treatment groups (data not shown).

BPaL therapy increases proinflammatory cytokine response in bone marrow

A comparative analysis for concentration of cytokines and chemokines in the bone

marrow, plasma, and lung samples from Mtb-infected C3HeB/FeJ mice treated with BPaL

or BPaS regimen was also conducted. The bone marrow samples demonstrated a

significant difference between the BPaL and BPaS groups, with appreciably higher level of pro-inflammatory cytokines and chemokines (IL-1β, IL-12p70, IL-23, TNFα, GROα (CXCL1), MP-2α (CXCL2), IP-10 (CXCL10), MP-1α (CCL3), RANTES (CCL5), MCP-3 (CCL7) and Eotaxin (CCL11)) in the BPaL- compared to BPaS-treated mice (Figure 4B). The plasma and lung samples, however, had similar cytokine and chemokine contents between the treatment groups except for MCP-3 (CCL7) in plasma which was significantly higher in BPaS compared to BPaL group (Figure S11). We also performed a correlation analysis of bone marrow cytokine and chemokine content with lung CFU obtained from treatment (BPaL or BPaS) and UnRx groups (Figure 4C). The analysis suggested that compared to UnRx control, there was a strong correlation between the pronounced reduction of lung bacterial burden and profound reduction in bone marrow cytokine and chemokine contents observed in mice from BPaL or BPaS groups. Similar correlations were found between lung CFU and content of cytokines and chemokines in lung (Figure S12) and plasma (Figure S13).

BPaL and BPaS therapies reduce inflammation-associated cells

We further assessed the environment of immune cells using flow cytometry in bone marrow, lungs, and blood from each group of Mtb-infected C3HeB/FeJ mice (Figure 5). The bone marrow (Figure 5A) results revealed that as compared to UnRx control, there was a significant decrease in the percentage of inflammatory myeloid phenotypes (CD45+CD3-CD11b+CD11c-Ly6C+CCR2+, CD45+CD3-CD11b+CD11c-Ly6C+CCR2+MHC-II+ and CD45+CD3-D14+CCR2+) in response to therapy with either BPa, BPaL, or BPaS. In contrast, neutrophils (CD45+CD3-CD11b+CD11c-

Ly6C+Ly6G^{high}), precursor T cells (CD45+CD3+) and B cells (CD45+CD3-CD19+B220-)

were significantly increased in either BPa, BPaL, or BPaS treatment groups compared

to UnRx control in bone marrow. This reduced inflammatory response in treatment

groups is also consistent in blood shown by significantly reduced inflammatory myeloid

cells (CD45+CD3-CD11b+CD11c-Ly6C+CCR2+) (Figure 5B). Interestingly, the

response to therapy in the lungs (Figure 5C) was manifested by a significant increase of

CD3+CD4+ T helper cells and B-1 cells (CD3-CD19+) and a reverse trend for

CD3+CD8+ and γδ-T cells (CD3+CD8+γδTCR+).

Furthermore, we also assessed changes in the distribution of immune cells in the

lungs from Mtb-infected C3HeB/FeJ mice in response to therapy using multiplex

fluorescent immunohistochemistry (mfIHC). A 7-color composite image for cell markers

(B220, CD4, CD8, Foxp3, F4/80 and Ly6G) along with DAPI is shown in Figure 6A,

while their single-color staining is shown in Figure 6B. Figure 6A shows a typical

necrotic TB granuloma comprised of central necrosis, peripheral rim and lung

parenchyma. The analysis of mfIHC images revealed that the BPaL and BPaS

treatments significantly and dramatically lowered the number of neutrophils (count

based on Ly6G+) compared to UnRx control, however, F4/80+ cells were observed

significantly higher in BPaS compared to UnRx control (Figure 6C, S14). Interestingly,

the spearman's correlation plot (Figure 6D) shows that a significant decrease in

neutrophils was positively correlated with the corresponding increase in most of the

other immune cells.

Discussion

292
We used two preclinical chronic TB murine efficacy models (30–32) to investigate

BPa, BPaL and BPaS, for efficacy and any associated AEs during the course of 4-

weeks of treatment. Based on the diversity of outcomes observed during human TB

disease (22) and as no single animal model recapitulates the wide spectrum of human

TB pathology (33), we chose the Balb/c and C3HeB/FeJ murine TB models. The Balb/c

chronic TB model is representative of a long-term Mtb chronic infection that develops

homogenous lung granulomatous lesions restraining the bacilli within intracellular

compartments (34) of macrophages and foamy macrophages. In contrast, low-dose

aerosol Mtb infection of C3HeB/FeJ mice also results in a chronic infection, but their

lungs exhibit a heterogenous spectrum of lesions including granulomas similar to those

seen in Balb/c chronic TB model in addition to caseous necrotic lesions surrounded by a

fibrotic rim (28,29,35). The caseum of these necrotic granulomas creates a hypoxic

environment and contains abundant extracellular bacilli (28,29,36) in a similar fashion to

necrotic granulomas found in some human TB patients. It is believed that the fibrotic,

necrotic and hypoxic environment of these granulomas creates barriers to drug

penetration, alters bacterial phenotype, and all together challenge therapeutic outcomes

(30–32,37). Therefore, to understand the implications of drug efficacy and drug

associated AEs in scenario without (Balb/c) and with (C3HeB/FeJ) necrotic granulomas,

both murine TB efficacy models were employed.

Overall, our antimicrobial data are in accordance with the granuloma spectrum of

both mouse models discussed above where a more robust reduction in lung bacterial

burden was observed in the absence (Balb/c) versus the presence (C3HeB/FeJ) of

necrotic lesions. Both multidrug regimens as BPaL or BPaS significantly reduced lung

bacterial burden by 3.8-4.0 $\log_{10}$ and 2.5-2.7 $\log_{10}$ in Balb/c and C3HeB/FeJ TB models,

respectively (Figures 1E-H). We therefore conclude that both regimens promote similar

bactericidal effects in murine models lacking, or featuring, advanced pulmonary

pathology. Furthermore, the potent antimicrobial effect of the BPaL and BPaS regimens

also resulted in improvement of the pathological outcome during chronic infection with

Mtb but only the L-containing BPaL regimen, not BPaS, was associated with a

significant decrease in the live body weight at week 4 in Mtb-infected Balb/c and

C3HeB/FeJ mice (Figure 2C&D). The extent of weight loss is an important preclinical

and clinical parameter in TB patients because it determines the severity of disease

progression (38), and it is also an indicator of *in vivo* drug efficacy (39). The bactericidal

effects of S (as monotherapy) (40) and BPaL (41) in mice observed in these studies is

in line with previous reports and similar inverse relationship between body weight and L

exposure was recently reported in human patients (42).

Among L-associated hematological side effects, the incidence of anemia is

reported up to 62.5% in MDR and XDR-TB patients (43–45) and the onset of this effect

can occur 2 weeks to 2 months post L administration (46). In human TB patients treated

with anti-TB drugs (47–49), increased RDWs and lower HGB are associated with anemia

and these parameters serve as markers of disease prognosis. Using a CBC profile of 20

peripheral blood parameters, our preclinical study failed to detect any difference between

L or S 4-weeks monotherapy in terms of total red blood cells, white blood cells, platelets,

or hemoglobin (HGB) concentration compared to UnRx control (data not shown).

However, L treatment alone increased the RDWs and both L and S decreased the MCHC

(Figure 3A). Likewise, none of the 20-blood parameters evaluated showed any change
after the first 2 weeks of therapy with BPa, BPaL, or BPaS (Figure 3B). However, by 4
weeks, a significant drop in HGB and an increase in the RDWs was apparent for the BPaL
group (Figure 3B). This is interpreted to mean that mild hematological effects observed
in mice treated for 4 weeks with L or BPaL are dependent on the number of L-doses
administered and are thus, time-dependent. Overall, the onset of these hematological
effects when testing an L-containing regimen (BPaL) is in agreement with previous
studies (46,50) in human patients.

The mechanism of L-induced toxicity is attributed to its binding with the host
mitochondrial ribosomes leading to mitochondrial toxicities (17). The latter results in
activation of Nlrp3 inflammasome (51) and subsequently results in L-mediated bone
marrow myelosuppression (52); a phenomenon consistent with the hematologic
anomalies seen in patients treated with L for extended time periods. Because
spectinamides do not bind to mitochondrial ribosomes there is reduced potential for
similar side effects (16,18). To conclusively test this hypothesis, we performed bone
marrow histopathology to quantify the myeloid to erythroid ratio (M:E), a parameter that
provides information about the relative proportions of myeloid lineage (granulocytes,
monocytes, and their precursors) to erythroid lineage (53). The BPa and BPaL regimens
altered M:E in the C3HeB/FeJ TB model by suppressing myeloid and inducing erythroid
lineages (Figure 3C&D) whereas no such difference was observed in mice treated with
BPaS compared to untreated control. L was previously shown to impact M:E ratio,
although an opposite trend was observed in those studies, which employed 12-days of

L administration and a different strain of otherwise healthy mice (51). Time course

studies using a single consistent assay method are needed to resolve this discrepancy.

Elevation of interleukin 1β (IL-1β) levels and activation of Nlrp3 inflammasome

have been previously linked to myelosuppression (52). A 26-plexed immunoassay on

bone marrow samples from Mtb-infected C3HeB/FeJ mice revealed that most of the

proinflammatory cytokines and chemokines including IL-1β, IL-12p70 and TNF-α were

present at significantly higher concentrations in animals treated with BPaL compared to

BPaS (Figure 4A&B). The presence of elevated IL-1β was previously reported during

monotherapy with L (51,54). Studies from UnRx Mtb-infected C3HeB/FeJ mice also

revealed highly elevated levels of cytokines and chemokines (Figure 4C). As expected,

the positive therapeutic effect of BPaL and BPaS (as seen by reduction in bacterial and

lesion burden of lungs) also correlated with decreased level of proinflammatory

cytokines in bone marrow, lung, and plasma (Figures 4C, S12 and S13 respectively).

Similar to changes observed for cytokine profile, the flow cytometry data of bone

marrow, blood, and lungs along with quantification of mfIHC lung image analysis

revealed a significant change in the percentage of myeloid and lymphoid phenotypes

during treatment compared to UnRx control (Figures 5 and 6). Most notably, the

therapeutic effect of the BPa, BPaL and BPaS treatments reduced inflammatory

myeloid cells expressing CCR2 in blood and bone marrow and reduction in cytotoxic T

cells (CD3+CD8+) and γδT cells (CD3+CD8+γδTCR+) in lungs. In addition, BPa, BPaL

and BPaS therapy significantly increased the influx of helper T cells (CD3+CD4+),

regulatory T cells (Foxp3+) and B cells (CD3-CD19+) in lungs. These results suggest

that the combination therapy promotes the immune system's equilibrium by reducing

inflammation and enhancing adaptive immune responses.

An additional striking finding from this study was a strong concordance between

the decrease in lung and spleen bacterial burden and a corresponding decrease in the

number of cells expressing the neutrophil associated marker (Ly6G) (Figure 6C&D). An

implication of this finding is that the favorable treatment outcomes may promote a

corresponding decline in Ly6G neutrophil population as suggested before (55). Among

all immune cell phenotypes studied, no differences were found between BPaL and

BPaS treated animals and only the F4/80 +cells in the BPaS (but not in the BPaL)

treated animals showed significant increase when compared to the UnRx control.

The TB drug development field is working towards developing shorter and safer

therapies with a common goal of developing new multidrug regimens of low pill burden

that are accessible to patients, of short duration (ideally 2-3 months) and consist of 3-4

drugs of novel mode-of-action with proven efficacy, safety, and limited toxicity. Here we

present initial results that are in line with these goals. Further studies with inhaled

spectinamide 1599 and BPa are therefore warranted to determine if 2-3 months of

treatment with BPaS, further reduces the bacterial burden and leads to sterilizing

activity without associated AEs.

Materials and Methods

Female C3HeB/FeJ and Balb/c mice at 6-8 weeks of age were purchased from

the Jackson Laboratories. All protocols and use of these animals were approved by the

Institutional Animal Care and Use Committee (IACUC) at CSU. Animals were infected

with a low dose aerosol infection of Mtb (Erdman strain; ATCC 35801) using an

inhalation exposure system (Glas-Col, Terre Haute, IN) calibrated to deliver ~50-100

colony forming units (CFU) to the lungs (22). Clinical observations (e.g., inactivity, rough

fur, hunched posture, increased respiratory rate or effort) were monitored daily and their

live body weights were taken weekly.

Drug preparation and treatment

Bedaquiline fumarate (B) and linezolid (L) were obtained from LKT laboratories,

pretomanid (Pa) from ChemShuttle and 1599 (S: dihydrochloride) was provided by Dr.

Lee at St. Jude Children's Research Hospital. B was administered at 25 mg/kg. Pa and

L were administered at 100 mg/kg each and S was delivered by inhalation in liquid form

at 50 and 100 mg/kg to Balb/c and C3HeB/FeJ mice, respectively. The drugs were

formulated in weekly batches according to the live body weight of the animals, aliquoted

for single daily dosing, and stored at 4 °C in the dark.

The drugs were prepared as reported previously (21). Drug treatment was

started 4 weeks post-aerosol infection for Balb/c and at 8-9 weeks post-aerosol infection

for C3HeB/FeJ mice to allow time for lung pathology to fully develop. All drugs were

administered once daily for 5 days/week for 4 weeks by oral (gavage) administration

except S which was administered 3 days/week by intrapulmonary aerosol delivery using

the Penn Century microsprayer as reported previously (22). B was administered in the

morning, Pa one hour after B and, L and S at least 4 hours after Pa.

Necropsy

After 4-weeks of treatment, C3HeB/FeJ and Balb/c mice were humanely

euthanized by CO_2 narcosis. Blood, lungs, spleen, femur and tibia bones were collected

from each mouse for further processing and analysis.

Assessment of Efficacy

The efficacy of the treatment was assessed by determining changes in bacterial

burden [measured as colony forming units (CFU)] in the lungs and spleen of animals at

necropsy. The lungs and spleen were homogenized and prepared as reported

previously (22). The lung homogenates were plated onto 7H11 agar plates

supplemented with 0.4% activated charcoal to reduce the carryover effect of drugs and

incubated at 37 °C for 6-8 weeks before the final CFU count. The remaining lung

homogenate was centrifuged, and the supernatant was collected and stored at -80 °C

for evaluation of cytokines and chemokines.

Histopathology and lesion scoring

The lungs were fixed in 4% paraformaldehyde (PFA) for 48 hours then

embedded in paraffin for histopathology. Sections from formalin fixed and paraffin

embedded (FFPE) tissues were cut at 5 µm, stained with hematoxylin and eosin (H&E)

and scanned at 40X magnification using multispectral automated PhenoImager (Akoya

Biosciences) for histopathological evaluation. The extent of lung lesion burden was

quantified in blinded digital images using an open-source QuPath software for image

analysis as described previously (56).

The sternum and one femur were fixed in 4% PFA and processed for histology.

To evaluate the myelosuppressive effect of the drugs, bone sections were cut at 5 μm

and stained with H&E. The number of myeloid and erythroid cells from 5 different

regions of the bone were blinded and then counted by a veterinary clinical pathologist.

Bone marrow collection

Briefly, a 0.6 mL sterile Eppendorf tube punctured at the bottom with the help of a

26-gauge needle was inserted into a 1.5 mL sterile Eppendorf tube. One end of the

epiphysis of the long bones was cut open to expose the bone marrow and placed down

into the small Eppendorf tube-system. The tubes were centrifuged at 10,000 x g for 15

seconds and the marrow was collected from the base of the large Eppendorf tube. The

bone marrow was resuspended in PBS and centrifuged again. Thereafter, the

supernatant was collected and stored at -80 °C for evaluation of cytokines and

chemokines while the bone marrow cells were saved in 4% PFA and freezing media for

further use in clinical pathology analysis and flow cytometry, respectively.

Processing of blood

For complete blood count (CBC) analysis of C3HeB/FeJ animals during the

treatment, blood was collected in K2-EDTA tubes via submandibular vein puncture as

described previously (57). The blood was immediately analyzed in VETSCAN® HM5

hematology analyzer (Zoetis).

At the time of necropsy, whole blood was collected via cardiac puncture in K2-

EDTA containing tubes. After adding an equal volume of PBS, the samples were

centrifuged at 800 × g for 10 min at 25 °C with the brake off (deceleration = 0). The top

plasma layer was collected and stored at -80 °C for evaluation of their cytokine's

content. The buffy coat was collected, washed and the erythrocytes were lysed using

Miltenyi RBC lysis buffer (Miltenyi, CA). The cells were washed and resuspended in 500

µL of complete DMEM media and prepared for flow cytometry analysis.

Cytokine quantification

Multiplex immunoassay was performed using a Luminex bead-based multiplex

ELISA kit (ProcartaPlex Mouse Cytokine & Chemokine Panel 1 26plex, reference #

EPXR260-26088-901, Invitrogen). Each sample was normalized to the total protein

concentration determined by Bicinchoninic acid (BCA) assay (Thermo Fisher). The BCA

and Luminex assay were performed according to the manufacturer's instructions and

the final stained samples were fixed with 4% PFA prior to acquisition. Sample data were

acquired on a MAGPIX instrument running xPONENT 4.3 software (Luminex Corp.).

Heatmaps were generated using R pheatmap package. Correlation analysis of cytokine

contents in bone marrow, plasma and lungs with the lung bacterial burden was

performed using the corrplot package in R.

Immune cell populations analysis

Single cell suspension of bone marrow, blood, and lungs from C3HeB/FeJ mice

was prepared as described previously (56). Cells counting, viability staining and cell

staining (Table S1) was performed accordingly (56). Samples were acquired using

Cytek Aurora^{TM} 4-Laser spectral flow cytometer where 100,000 events were recorded.

Data were analyzed in FlowJo software (BD Biosciences) using manual gating (58).

Multiplex fluorescent immunohistochemistry

Five μm sections of FFPE lung tissues were stained for multiplex fluorescent

immunohistochemistry (mfIHC) by the Imaging Core at the University of Colorado,

Anschutz Medical Campus, Denver. The mfIHC was performed for a panel of 6-color

antibodies + DAPI using Opal-plex Tyramide Signal Amplification (TSA) technique using

a Leica Bond III autostainer. The details of antibodies and Opal fluorophores used are

given in Table S2. Each antibody was optimized using Opal 3-Plex Anti-Rb Detection Kit

(Akoya Biosciences Inc. cat# NEL830001KT) and stained with automated LabSat^{TM}

Research (Lunaphore Technologies SA, Epredia). Slides were scanned using

multispectral automated PhenoImager^{TM} (Akoya Biosciences) and analyzed for several

immune cell populations using the inForm$^®$ tissue Finder (Version 2.4.8) and

Phenochart^{TM} (Version 1.0.12) software (Akoya Biosciences).

Statistical analysis. Bacterial burden data were expressed as CFU which were Log$_{10}$-

transformed and analyzed using GraphPad Prism version 9.5.1 (GraphPad software, La

Jolla, CA). The statistical analysis was performed using a Tukey's multiple comparison

test as part of either one-way or two-way ANOVA and mixed-model effect where

necessary. The correlation analysis was performed using the spearman's correlation

test. Flow cytometry and mfIHC data were graphed in R studio and statistical evaluation

was performed using stats package in R.

Figure 1. Bactericidal effect of BPaL and BPaS in TB mouse models after 4-weeks of treatment

Balb/c and C3HeB/FeJ female mice were chronically infected with a low dose aerosol infection of Mtb Erdman strain to deliver $\sim$75 and $\sim$100 bacilli respectively. Post-infection, Balb/c and C3HeB/FeJ mice were rested for 4 and 8-9 weeks respectively until they were randomly assigned to the study groups. The mice were treated with monotherapy of linezolid or spectinamide 1599 or combination therapy of BPaL, BPa or BPaS for 4 weeks. Bedaquiline (B), pretomanid (Pa) and linezolid (L) were administered at 25, 100 and 100 mg/kg respectively by oral gavage for 5/7 a week while spectinamide 1599 at 50 and 100 mg/kg in Balb/c and C3HeB/FeJ TB models respectively for 3/7 a week on the alternate days via intrapulmonary aerosol delivery. On the third day of the last treatment, the mice were euthanized, and their lungs and spleen were collected. The organs were homogenized, serially diluted and plated on 7H11 agar with 4% charcoal (to avoid drug carry-over effect) to determine bacterial burden in the form of colony forming units (CFU) in each sample. CFU were enumerated after 4-6 weeks of incubation at 37 °C and expressed as $\log_{10}$.C3HeB/FeJ and Balb/c TB models showing efficacy of monotherapy (A-D) and combination therapy (E-H). The C3HeB/FeJ graphs (E&G) represent the pooled data from three independent studies (n=3-8 each, Figures S1-3) and two of the three studies contained BPa as a reference control. The Balb/c graphs (F&H) represent the pooled data from two independent studies (n=5 each, Figures S7-8).

Statistical significance was calculated using one-way ANOVA with Tukey's multiple comparison test, $p < 0.05$ was considered significant and ** = $p<0.001$, *** = $p<0.0001$,

539   **** = P<0.0001. UnRx = untreated, L = linezolid, S = spectinamide 1599, LOD: limit of

540   detection.

Figure 2. Effect of therapy on the live body weight and lung histopathology of TB mouse models after 4-weeks of treatment

The live body weight of animals in Figure 1 treated with monotherapy (A&B) or multidrug therapy (C&D) was recorded on a weekly basis. The average weight and SEM of each group is shown. At the end of therapy, the mice were euthanized, and their lungs were collected and processed for histopathology and lesion scoring (E-H). FFPE sections were cut at 5 µm, stained with hematoxylin and eosin (H&E) and imaged at 40x (E&G). Lesion scores (F&H) were calculated as the proportion of infected area over the total lung area per animal. Statistical significance was calculated using mixed-effect model (body weight) and one-way ANOVA (lesion score) with Tukey's multiple comparison test, and $p < 0.05$ was considered significant and ** = p<0.001, *** = P<0.0001. UnRx = untreated, L =

553   linezolid,   S   =   spectinamide   1599

Figure 3. Complete blood count profiling and bone marrow histopathology of Mtb infected C3HeB/FeJ TB model during 4-weeks of therapy

During therapy of mice in Figure 1, the blood of live animals was collected at 1- and 2-weeks via the puncture of submandibular vein. At the end of 4-weeks therapy, the blood was collected by cardiac puncture. Blood was collected in K2-EDTA (purple top) tube and immediately analyzed for the complete blood count in VETSCAN® HM5 hematology analyzer (Zoetis). (A) monotherapy, (B) multidrug therapy. The MCHC (mean corpuscular hemoglobin concentration) and RDWs (red blood cell distribution width-standard deviation) along with the HGB (hemoglobin concentration) and MPV (mean platelet volume) are shown. A horizontal dotted line indicates the lower end of the reference interval for C3HeB/FeJ mice.

The sternum, femur and tibias bones from each mouse were collected, fixed in 4% PFA, and processed for histology (C&D). Sections were cut at 5 µm, stained with hematoxylin and eosin (H&E) and imaged at 40x. (C) Representative photos of bone marrow sections showing myeloid (black arrows) and erythroid (white arrows) cells in bone marrow of untreated (UnRx) and treatment (BPa, BPaS and BPaS) groups. (D) The number of myeloid (M) and erythroid (E) among a total of 300 cells in 5 different regions were counted for each group and M:E was calculated. Statistical significance was calculated using one-way ANOVA with Tukey's test for multiple comparisons. $p < 0.05$ was considered significant and ** = p<0.001, *** = P<0.0001.

Figure 4. Bone marrow cytokine and chemokine profiling in Mtb infected

C3HeB/FeJ TB model after 4-weeks of treatment

Femur bones from selected studies in mice in Figure 1 were collected to harvest the bone marrow. The bone marrow was resuspended in PBS, centrifuged and the supernatant was collected for the evaluation of cytokine's content. BPaL and BPaS therapy showed profile of 26 cytokines and chemokines in bone marrow and the data were converted to Z score and represented as a heatmap (A) and graphically (B). (C) Spearman's correlation analysis of bone marrow cytokines and chemokines with the lung bacterial burden (CFU).

Statistical significance was calculated using the t test. $p < 0.05$ was considered significant and ** = p<0.001*** = p<0.0001, **** = P<0.00001.

Figure 5. Immune cell populations in the bone marrow, lung and blood of

C3HeB/FeJ TB model after 4-weeks of treatment

The bone marrow, lung and blood from selected studies from Figure 1 were evaluated by

flow cytometry. The samples were processed for a panel of 17-color antibodies and the

data were analyzed by FlowJo software using manual gating strategy. The myeloid and

lymphoid phenotypes present in the untreated (UnRx) and treatment (BPa, BPaL or

BPaS) groups are shown. Statistical significance was calculated using one-way ANOVA

with Tukey's test for multiple comparisons. $p < 0.05$ was considered significant and ** =

p<0.001*** = p<0.0001, **** = P<0.00001.

Figure 6. Immune cell populations in the lungs of Mtb infected C3HeB/FeJ TB model after 4-weeks of treatment

Selected mice from those shown in Figure 1 were processed for multiplex fluorescent immunohistochemistry (mfIHC). The mfIHC was performed for a panel of 6-color antibodies + DAPI using Opal-plex Tyramide Signal Amplification (TSA). Slides were scanned using multispectral automated PhenoImager™ (Akoya Biosciences) and analyzed for different immune cell populations using the inForm® tissue Finder and Phenochart software (Akoya Biosciences). (A) The lung mfIHC full composite image displays B220, CD4, CD8, F4/80, FoxP3 and Ly6G markers along with DAPI staining for nuclei in the TB granuloma. The central and peripheral regions of a TB granuloma and the parenchyma of lung are also shown. (B) Single color composite image of individual markers with DAPI showing distribution of each immune cell population in the TB granuloma. (C) Cell populations (%) of several immune cells per total number of phenotypes calculated in untreated (UnRx) and treatment (BPaL and BPaS) groups based on a panel of 6-color antibodies + DAPI. (D) Spearman's correlation matrix for several immune cell populations (B220, CD4, CD8, F4/80, FoxP3, Ly6G) showing all relationships. A coefficient with a value of either +1 (blue), 0 (white), or -1 (red) indicates a perfect association, no association, and a perfect negative association of ranks, respectively. Numbers indicate the correlation coefficient.

Acknowledgments: This research was supported by the National Institute of Allergy and
Infectious Diseases and the Office of the Director of the National Institutes of Health (grants
R01AI120670, R01AI090810 and 1S10OD030263 respectively). Dr. Malik Zohaib Ali was a
Fulbright Foreign Student Program grantee from Pakistan. Dr. Charles Daley from the Division of
Mycobacterial and Respiratory Infections, National Jewish Health, Denver, Colorado 80206, USA
advised on current TB drug treatments. The content is solely the responsibility of the authors and
does not necessarily represent the official views of the National Institutes of Health. We
acknowledge the staff of the Laboratory Animal Resources at Colorado State University for
providing animal care.

References

1.   Campbell JR, Nsengiyumva P, Chiang LY, Jamieson F, Khadawardi H, Mah HKH, et al. Costs of Tuberculosis at 3 Treatment Centers, Canada, 2010–2016. Emerg Infect Dis. 2022;28(9):1814–23.

2.   Zhang MW, Zhou L, Zhang Y, Chen B, Peng Y, Wang F, et al. Treatment outcomes of patients with multidrug and extensively drug-resistant tuberculosis in Zhejiang, China. Eur J Med Res. 2021;26(1):1–10.

3.   Williams K, Minkowski A, Amoabeng O, Peloquin CA, Taylor D, Andries K, et al. Sterilizing activities of novel combinations lacking first- and second-line drugs in a murine model of tuberculosis. Antimicrob Agents Chemother. 2012;56(6):3114–20.

4.   Tasneen R, Betoudji F, Tyagi S, Li SY, Williams K, Converse PJ, et al. Contribution of oxazolidinones to the efficacy of novel regimens containing bedaquiline and pretomanid in a mouse model of tuberculosis. Antimicrob Agents Chemother. 2016;60(1):270–7.

5.   Conradie F, Diacon AH, Ngubane N, Howell P, Everitt D, Crook AM, et al. Treatment of Highly Drug-Resistant Pulmonary Tuberculosis. N Engl J Med. 2020;382(10):893–902.

6.   Conradie F, Bagdasaryan TR, Borisov S, Howell P, Mikiashvili L, Ngubane N, et al. Bedaquiline–Pretomanid–Linezolid Regimens for Drug-Resistant Tuberculosis.

N Engl J Med. 2022;387(9):810–23.

7.   TB-Alliance. Update on Pretomanid , in Combination with Bedaquiline and

Linezolid (the BPaL Regimen) in Drug-Resistant TB. 2022.

8.   Jackson A. Foresight. In: Drugs and the Future: Brain Science, Addiction and

Society. 2006. p. 7–10.

9.   Richter E, Rüsch-Gerdes S, Hillemann D. First linezolid-resistant clinical isolates

of Mycobacterium tuberculosis. Antimicrob Agents Chemother. 2007;51(4):1534–

6.

10.   Li SY, Converse PJ, Betoudji F, Lee J, Mdluli K, Upton A, et al. Next-Generation

Diarylquinolines Improve Sterilizing Activity of Regimens with Pretomanid and the

Novel Oxazolidinone TBI-223 in a Mouse Tuberculosis Model. Antimicrob Agents

Chemother. 2023;67(4):1–9.

11.   Holloway WJ. Spectinomycin. Med Clin North Am. 1982;66(1).

12.   Kanchugl, S. and Selmer M. Structural Recognition of Spectinomycin by

Resistance Enzyme ANT(9) from Enterococcus faecalis. Antimicrob Agents

Chemother. 2020;64(June):e00371-20.

13.   Owusu E, Amartey BT, Afutu E, Boafo N. Aminoglycoside Therapy for

Tuberculosis: Evidence for Ototoxicity among Tuberculosis Patients in Ghana.

Diseases. 2022;10(1):10.

14.   De Jager R, Van Altena R. Hearing loss and nephrotoxicity in long-term

aminoglycoside treatment in patients with tuberculosis. Int J Tuberc Lung Dis.

2002;6(7):622–7.

15.   Temrikar ZH, Kodidela S, Kumar S, Liu J, Robertson GT, Lee RE, et al.

Characterization of spectinamide 1599 efficacy against different mycobacterial

phenotypes. Tuberculosis [Internet]. 2023;140(April):102342. Available from:

https://doi.org/10.1016/j.tube.2023.102342

16.   Richard E. Lee, Julian G. Hurdle, Jiuyu Liu, David F. Bruhn, Tanja Matt, Michael

S. Scherman, Pavan K Vaddady, Zhong Zheng, Jianjun Qi, Rashid Akbergenov,

Sourav Das, Dora B. Madhura, Chetan Rathi, Ashit Trivedi, Cristina Villellas,#,

Robin. B. Lee, Rakes and AJL. Spectinamides: A New Class of Semisynthetic

Anti-Tuberculosis Agents that Overcome Native Drug Efflux. Nat Med.

2017;20(2):152–8.

17.   De Vriese AS, Van Coster R, Smet J, Seneca S, Lovering A, Van Haute LL, et al.

Linezolid-induced inhibition of mitochondrial protein synthesis. Clin Infect Dis.

2006;42(8):1111–7.

18.   Modongo C, Pasipanodya JG, Zetola NM, Williams SM, Sirugo G, Gumboc T.

Amikacin concentrations predictive of ototoxicity in multidrug-resistant

tuberculosis patients. Antimicrob Agents Chemother. 2015;59(10):6337–43.

19.   Liu J, Bruhn DF, Lee RB, Zheng Z, Janusic T, Scherbakov D, et al. Structure-

activity relationships of spectinamide antituberculosis agents: A dissection of

ribosomal inhibition and native efflux avoidance contributions. ACS Infect Dis.

2017;3(1):72–88.

20.   Bruhn DF, Scherman MS, Liu J, Scherbakov D, Meibohm B, Böttger EC, et al. In

vitro and in vivo evaluation of synergism between anti-tubercular spectinamides

and non-classical tuberculosis antibiotics. Sci Rep. 2015;5(May):1–10.

21.   Robertson GT, Scherman MS, Bruhn DF, Liu J, Hastings C, McNeil MR, et al.

Spectinamides are effective partner agents for the treatment of tuberculosis in

multiple mouse infection models. J Antimicrob Chemother. 2017;72(3):770–7.

22.   Gonzalez-Juarrero M, Lukka PB, Wagh S, Walz A, Arab J, Pearce C, et al.

Preclinical Evaluation of Inhalational Spectinamide-1599 Therapy against

Tuberculosis. ACS Infect Dis. 2021;7(10):2850–63.

23.   Wagh S, Rathi C, Lukka PB, Parmar K, Temrikar Z, Liu J, et al. Model-based

exposure-response assessment for spectinamide 1810 in a mouse model of

tuberculosis. Antimicrob Agents Chemother. 2021;65(11):1–15.

24.   Nirmal A, Sznajder K, Patil R, Shaikh B. Preference of inhalants over

pills/injections among pulmonary tuberculosis patients in Western India: A cross-

sectional study. J Clin Tuberc Other Mycobact Dis. 2021;23.

25.   Matthieu B, Matthieu J, Nicolas G, Patrice G, Srine M, William C, et al.

Comparison of intrapulmonary and systemic pharmacokinetics of colistin

methanesulfonate (CMS) and colistin after aerosol delivery and intravenous

administration of CMS in critically ill patients. Antimicrob Agents Chemother.

2014;58(12):7331–9.

26.   Rathi C, Lukka PB, Wagh S, Lee RE, Lenaerts AJ, Braunstein M, et al.

783   Comparative pharmacokinetics of spectinamide 1599 after subcutaneous and

784   intrapulmonary aerosol administration in mice. Tuberculosis. 2019;114(December

785   2018):119–22.

786   27.   Hickey AJ, Misra A, Fourie PB. Dry powder antibiotic aerosol product

787   development: Inhaled therapy for tuberculosis. J Pharm Sci [Internet].

788   2013;102(11):3900–7. Available from: http://dx.doi.org/10.1002/jps.23705

789   28.   Irwin SM, Driver E, Lyon E, Schrupp C, Ryan G, Gonzalez-Juarrero M, et al.

790   Presence of multiple lesion types with vastly different microenvironments in

791   C3HeB/FeJ mice following aerosol infection with Mycobacterium tuberculosis.

792   DMM Dis Model Mech. 2015;8(6):591–602.

793   29.   Driver ER, Ryan GJ, Hoff DR, Irwin SM, Basaraba RJ, Kramnik I, et al. Evaluation

794   of a mouse model of necrotic granuloma formation using C3HeB/FeJ mice for

795   testing of drugs against Mycobacterium tuberculosis. Antimicrob Agents

796   Chemother. 2012;56(6):3181–95.

797   30.   Harper J, Skerry C, Davis SL, Tasneen R, Weir M, Kramnik I, et al. Mouse model

798   of necrotic tuberculosis granulomas develops hypoxic lesions. J Infect Dis.

799   2012;205(4):595–602.

800   31.   Irwin SM, Gruppo V, Brooks E, Gilliland J, Scherman M, Reichlen MJ, et al.

801   Limited activity of clofazimine as a single drug in a mouse model of tuberculosis

802   exhibiting caseous necrotic granulomas. Antimicrob Agents Chemother.

2014;58(7):4026–34.

32.   S Aly, K Wagner, C Keller, S Malm, A Malzan, S Brandau, Bange E. Oxygen

status of lung granulomas in Mycobacterium tuberculosis-infected mice. J Pathol.

2006;210:298–305.

33.   Singh AK, Gupta UD. Animal models of tuberculosis: Lesson learnt. Indian J Med

Res [Internet]. 2018;147:456–63. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/23144490

34.   De Groote MA, Gilliland JC, Wells CL, Brooks EJ, Woolhiser LK, Gruppo V, et al.

Comparative studies evaluating mouse models used for efficacy testing of

experimental drugs against Mycobacterium tuberculosis. Antimicrob Agents

Chemother. 2011;55(3):1237–47.

35.   Dpre, Robertson GT, Ramey ME, Massoudi LM, Carter CL, Zimmerman M, et al.

Comparative Analysis of Pharmacodynamics in the C3HeB / FeJ. Antimicrob

Agents Chemother aac.asm.org. 2021;65 Issue(July):e00583-21.

36.   Lanoix JP, Lenaerts AJ, Nuermberger EL. Heterogeneous disease progression

and treatment response in a C3HeB/FeJ mouse model of tuberculosis. DMM Dis

Model Mech. 2015;8(6):603–10.

37.   Walter ND, Ernest JP, Dide-Agossou C, Bauman AA, Ramey ME, Rossmassler K,

et al. Lung microenvironments harbor Mycobacterium tuberculosis phenotypes

with distinct treatment responses . Antimicrob Agents Chemother. 2023;67(9):1–

15.

38. Van Crevel R, Karyadi E, Netea MG, Verhoef H, Nelwan RHH, West CE, et al.

Decreased plasma leptin concentrations in tuberculosis patients are associated

with wasting and inflammation. J Clin Endocrinol Metab. 2002;87(2):758–63.

39. Nikonenko B V., Samala R, Einck L, Nacy CA. Rapid, simple in vivo screen for

new drugs active against Mycobacterium tuberculosis. Antimicrob Agents

Chemother. 2004;48(12):4550–5.

40. Pstragowski, M. Zbrzezna, M. and Bujalska-Zadrozny M. Advances in

pharmacotherapy of Tuberculosis [Internet]. Vol. 74, Acta Poloniae - Drug

Research. 2017. p. 3–11. Available from:

https://www.embase.com/search/results?subaction=viewrecord&id=L604925228&

from=export

41. Nuermberger EL, Martínez-Martínez MS, Sanz O, Urones B, Esquivias J, Soni H,

et al. GSK2556286 Is a Novel Antitubercular Drug Candidate Effective In Vivo

with the Potential To Shorten Tuberculosis Treatment. Vol. 66, Antimicrobial

Agents and Chemotherapy. 2022.

42. Wasserman S, Denti P, Brust JCM, Abdelwahab M, Hlungulu S, Wiesner L, et al.

Linezolid pharmacokinetics in South African patients with drug-resistant

tuberculosis and a high prevalence of HIV coinfection. Antimicrob Agents

Chemother. 2019;63(3).

43. Abebe G, , Zegeye Bonsa WK. Early Safety and Efficacy of Linezolid-Based

Combination Therapy among Patients with Drug-Resistant Tuberculosis in

North-western Nigeria. Int J Mycobacteriology. 2021;10:129–35.

44. Palomino JC, Martin A. Drug resistance mechanisms in Mycobacterium
tuberculosis. Antibiotics. 2014;3(3):317–40.

45. Sotgiu G, Centis R, D'Ambrosio L, Alffenaar JWC, Anger HA, Caminero JA, et al.
Efficacy, safety and tolerability of linezolid containing regimens in treating MDR-
TB and XDR-TB: Systematic review and meta-analysis. Eur Respir J.
2012;40(6):1430–42.

46. Tang S, Yao L, Hao X, Zhang X, Liu G, Liu X, et al. Efficacy, safety and tolerability
of linezolid for the treatment of XDR-TB: A study in China. Eur Respir J [Internet].
2015;45(1):161–70. Available from: http://dx.doi.org/10.1183/09031936.00035114

47. Nezamodini ZS, Rezvani Z, Kian K. Electronic Physician (ISSN : 2008-5842).
Electron Physician. 2017;9(January):3592–7.

48. Kassa E, Enawgaw B, Gelaw A, Gelaw B. Effect of anti-tuberculosis drugs on
hematological profiles of tuberculosis patients attending at University of Gondar
Hospital, Northwest Ethiopia. BMC Hematol. 2016;16(1):1–11.

49. Luo M, Liu M, Wu X, Wu Y, Yang H, Qin L, et al. Impact of anemia on prognosis
in tuberculosis patients. Ann Transl Med. 2022;10(6):329–329.

50. Gerson SL, Kaplan SL, Bruss JB, Le V, Arellano FM, Hafkin B, et al. Hematologic
effects of linezolid: Summary of clinical experience. Antimicrob Agents
Chemother. 2002;46(8):2723–6.

51.   Iyer SS, Iyer SS, He Q, He Q, Janczy JR, Janczy JR, et al. Mitochondrial

Cardiolipin Is Required for Nlrp3 Inflammasome Activation. Immunity [Internet].

2013;39(2):311–23. Available from:

http://linkinghub.elsevier.com/retrieve/pii/S1074761313003294%5Cnpapers3://pu

blication/doi/10.1016/j.immuni.2013.08.001

52.   Winchell CG, Mishra BB, Phuah JY, Saqib M, Nelson SJ, Maiello P, et al.

Evaluation of IL-1 Blockade as an Adjunct to Linezolid Therapy for Tuberculosis in

Mice and Macaques. Front Immunol. 2020;11(May):1–18.

53.   Elmore SA. Enhanced Histopathology of the Bone Marrow Susan. Toxicol Pathol

[Internet]. 2006;34(5):666–86. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624763/pdf/nihms412728.pdf

54.   Jasenosky LD, Scriba TJ, Hanekom WA, Goldfeld AE. T cells and adaptive

immunity to Mycobacterium tuberculosis in humans. Immunol Rev.

2015;264(1):74–87.

55.   Lovewell RR, Baer C, Mishra BB, Smith CM, Christopher M. Granulocytes act as

a niche for Mycobacterium tuberculosis growth. Mucosal Immunol.

2021;14(1):229–41.

56.   Dutt TS, Karger BR, Fox A, Youssef N, Dadhwal R, Ali MZ, et al. Mucosal

exposure to non-tuberculous mycobacteria elicits B cell-mediated immunity

against pulmonary tuberculosis. Cell Rep [Internet]. 2022;41(11):111783.

Available from: https://doi.org/10.1016/j.celrep.2022.111783

57.   Golde WT, Gollobin P, Rodriguez LL. A rapid, simple, and humane method for

887   submandibular bleeding of mice using a lancet. Lab Anim (NY). 2005;34(9):39–

888   43.

889   58.   Fox A, Dutt TS, Karger B, Rojas M, Obregón-Henao A, Anderson GB, et al. Cyto-

890   Feature Engineering: A Pipeline for Flow Cytometry Analysis to Uncover Immune

891   Populations and Associations with Disease. Sci Rep. 2020;10(1):1–12.

Supporting information

Figure S1 (C3HeB/FeJ Study 1). Bacterial burden (CFU) in Mtb infected C3HeB/FeJ mice treated with BPaL and BPaS regimen for 4 weeks

Figure S2 (C3HeB/FeJ Study 2). Bacterial burden (CFU) in Mtb infected C3HeB/FeJ mice treated with BPa, BPaL and BPaS regimen for 4 weeks

Figure S3 (C3HeB/FeJ Study 3). Bacterial burden (CFU) in Mtb infected C3HeB/FeJ mice treated with BPa, BPaL and BPaS regimen for 4 weeks

Figure S4 (C3HeB/FeJ Study 1). Change in the average live body weight of Mtb infected C3HeB/FeJ mice during drug treatment

Figure S5 (C3HeB/FeJ Study 2). Change in the average live body weight of Mtb infected C3HeB/FeJ mice during drug treatment

Figure S6 (C3HeB/FeJ Study 3). Change in the average live body weight of Mtb infected C3HeB/FeJ mice during drug treatment

Figure S7 (Balb/c Study 1). Bacterial burden (CFU) in Mtb infected Balb/c mice treated with BPaL and BPaS regimen for 4 weeks

Figure S8 (Balb/c Study 2). Bacterial burden (CFU) in Mtb infected Balb/c mice treated with BPaL and BPaS regimen for 4 weeks

Figure S9 (Balb/c Study 1). Change in the average live body weight of Mtb infected Balb/c mice during drug treatment

Figure S10 (Balb/c Study 2). Change in the average live body weight of Mtb infected Balb/c mice during

drug treatment

Figure S11. Change in the cytokines and chemokines profile in Mtb infected C3HeB/FeJ mice during drug

treatment

Figure S12. Spearman's correlation analysis between bacterial burden (CFU) and cytokine and

chemokine profile in the lungs of Mtb infected C3HeB/FeJ mice treated with BPaL and BPaS regimen

Figure S13. Spearman's correlation analysis between lung bacterial burden (CFU) and plasma cytokine

and chemokine profile in Mtb infected C3HeB/FeJ mice treated with BPaL and BPaS regimen

Figure S14. Immune cell populations in the lungs of Mtb infected C3HeB/FeJ TB model after 4 weeks

therapy

Table S1. **Reagents**

Table S2. Antibodies