Astyanax mexicanus surface and cavefish chromosome-scale assemblies for trait variation discovery

Wesley C. Warren[1,2], Edward S. Rice[1], Maggs X[1], Emma Roback[3], Alex Keene[4], Fergal Smith[5], Denye Ogeh[5], Leanne Haggerty[5], Rachel A. Carroll[1], Suzanne McGaugh[3], Nicolas Rohner[6,7]

[1]Department of Animal Sciences, Department of Surgery, University of Missouri, Bond Life Sciences Center, Columbia, MO

[2]Institute for Data Science and Informatics, University of Missouri, Columbia, MO

[3]Department of Ecology, Evolution, and Behavior, University of Minnesota, Saint Paul, MN

[4]Department of Biology, Texas AM University, College Station, TX

[5]European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, Hinxton, Cambridge CB10 1SD, UK

[6]Stowers Institute for Medical Research, Kansas City, MO

[7]Department of Molecular and Integrative Physiology, KU Medical Center, Kansas City, KS

Authors for correspondence: Wesley C. Warren (warrenwc@missouri.edu); Nicolas Rohner (nro@stowers.org)

Keywords: *Astyanax mexicanus*, cavefish, chromosome assembly

Abstract

The ability of organisms to adapt to sudden extreme environmental changes produces

some of the most drastic examples of rapid phenotypic evolution. The Mexican Tetra,

Astyanax mexicanus, is abundant in the surface waters of northeastern Mexico, but

repeated colonizations of cave environments have resulted in the independent evolution

of troglomorphic phenotypes in several populations. Here, we present three chromosome-

scale assemblies of this species, for one surface and two cave populations, enabling the

first whole-genome comparisons between independently evolved cave populations to

evaluate the genetic basis for the evolution of adaptation to the cave environment. Our

assemblies represent the highest quality of sequence completeness with predicted protein-

coding and non-coding gene metrics far surpassing prior resources and, to our

knowledge, all long-read assembled teleost genomes, including zebrafish. Whole genome

synteny alignments show highly conserved gene order among cave forms in contrast to a

higher number of chromosomal rearrangements when compared to other phylogenetically

close or distant teleost species. By phylogenetically assessing gene orthology across

distant branches of amniotes, we discover gene orthogroups unique to *A. mexicanus.*

When compared to a representative surface fish genome, we find a rich amount of

structural sequence diversity, defined here as the number and size of insertions and

deletions as well as expanding and contracting repeats across cave forms. These new

more complete genomic resources ensure higher trait resolution for comparative,

functional, developmental, and genetic studies of drastic trait differences within a species.

Introduction

Natural trait alterations in the Mexican tetra cavefish *Astyanax mexicanus*, some with

extreme phenotypic consequences, have offered insight into how genetic adaptation

works (ROHNER 2018) (MCGAUGH *et al.* 2020). This species exists in two distinct forms:

a traditional river-dweller and a blind, depigmented cave-dweller. Many studies have

highlighted the use of the *A. mexicanus* surface fish to cave-life as an adaptation model to

investigate the unique molecular mechanisms underlying various disease traits, such as

obesity, sleep disorders, diabetes, heart regeneration, and many others, which has

elevated its importance as an evolutionary model and led to its rapidly expanding use

(DUBOUE *et al.* 2011; ASPIRAS *et al.* 2015) (OJHA AND WATVE 2018) (STOCKDALE *et al.*

2018) (BILANDŽIJA 2019). Furthermore, we envision its wider use with the expansion of

A. mexicanus genetic resources by taking advantage of a relatively short evolutionary

time of transition, ~150,000 years from surface to cave and clearly demarcated

phenotypic differences (HERMAN *et al.* 2018).

High-quality and nearly complete chromosomes of many species are becoming

more readily available, but challenges remain for genomes with difficult structural

features, for example, the highly repetitive zebrafish genome (CHERNYAVSKAYA *et al.*

2022). Further research has refined best practices across broad phyla to resolve

difficulties in building multiple genome assemblies with near gap-free representation

(JARVIS *et al.* 2022). To date, few aquatic species have reached these higher levels of

contiguity and representation, but recently developed long-read hybrid methods improve

de novo assembly (RAUTIAINEN *et al.* 2023). Many recently published aquatic genomes,

although significantly more complete in sequence representation, have relied on error-

prone long-read reads and their correction with short-reads (MOORE *et al.* 2023)

(ROBERTS *et al.* 2023). This new generation of genome assemblies has significantly

advanced our knowledge of historically underrepresented sequences, in particular sex

chromosomes (IMARAZENE *et al.* 2021) (DU *et al.* 2022).

In cavefish, a recent long-read based assembly improved identification of

candidate genes underlying QTLs, discovered fixed or variable deletions in each cave

form, and guided gene editing experiments (WARREN *et al.* 2021). However, this

assembly of a surface-dwelling individual suffered from a high level of gene

fragmentation due to the use of high-error long reads. An individual from the Pachón

cave-dwelling population was recently assembled with high levels of contiguity and

facilitated the discovery of novel sex chromosome origins (IMARAZENE *et al.* 2021).

However, given the high phenotypic divergence of the various populations, references of

multiple populations are necessary to best perform comparative genomics between

surface and cave morphs. To this end, we present here three highly continuous

chromosome-scale assemblies of individuals from the Río Choy surface population, as

well as the Molino and Tinaja cave populations to complement Pachón and for the first

time present more complete references that span demographically unique and

independent cave populations. The generation of these resources in *A. mexicanus* will

support the community's use of this aquatic model of human disease, particularly those

wishing to compare trait diversity in the well-studied zebrafish model.

Methods

Sequencing and assembly. The *A. mexicanus* DNA samples were obtained from female

fishes of surface, Molino, and Tinaja origins (Fig. 1A) reared at the Stowers Institute

aquatic facility under IACUC approved protocol (Protocol ID: 2021-126). High

molecular weight DNA was isolated using the salting out method described in the 10X

genomics demonstrated protocol (10X Genomics; Pleasanton, CA) from muscle tissue to

generate single molecular real time (SMRT) sequences using HiFi sequence base calling

output mode from the Sequel II instrument (Pacific Biosciences) according to the

manufacturer's protocols. From all SMRT sequences, total HiFi coverage ranged from

27-46x using a genome size estimate of 1.4Gb with average read lengths across all

samples of 21kb (Suppl. Table 1).

We assembled these reads into contigs using hifiasm v0.13-r308 with default

options (CHENG *et al.* 2021). To look for haplotigs, we aligned CLR (Circular Long

Reads) reads to the contigs using minimap v2.20 with arguments "-ax map-pb". We ran

purge_haplotigs v1.1.1 (GUAN *et al.* 2020) on all three assemblies; the Molino and Tinaja

assemblies had detectable signals of haplotigs but surface did not, so we removed

haplotigs from Molino and Tinaja using coverage cutoffs (100, 197, 250) and (30, 193,

250), respectively. Haplotigs represent assembled sequences that are essentially a copy of

the other haplotype and thus are removed to avoid redundancy in genome representation.

To perform scaffolding a sibling of the reference fish was used to generate and sequence

a HiC library developed with the Proximo Hi-C kit (Phase Genomics: Seattle, WA)

according to the manufacturers protocol. We next assembled the contigs into scaffolds

using a custom pipeline (see Code Availability) that aligns the Hi-C reads to the contigs

with bwa mem v0.7.17 (LI AND DURBIN 2009), postprocesses the alignments, and finally

scaffolds with SALSA v2.2 (GHURYE *et al.* 2019) and Juicebox (version 1.11.08)

(DURAND *et al.* 2016), as previously described (GHURYE AND POP 2019). We curated the

scaffolds using a combination of synteny and manual examination of the Hi-C heatmaps

(ROBINSON *et al.* 2018; RHIE *et al.* 2021; WARRENLAB. 2022). Agptools was used to

finalize chromosome assignments to be consistent with chromosome nomenclature of the

surface assembled genome reported in Warren et al (WARREN *et al.* 2021).

Chromosomes were numbered by aligned synteny to assembled chromosomes of the

previously assembled surface form of *A. mexicanus* (WARREN *et al.* 2021). The

completeness of gene representation was assessed using BUSCO v4.1.2

(Actinopterygii_odb10) (SIMAO *et al.* 2015).

Whole genome interspecies and intraspecies analysis. To better understand shared gene

organization with other teleost experimental models and across cave morphs, we first

built and visualized syntenic orthology of surface fish chromosomes (AstMex3) to widely

used fish experimental models including zebrafish (GRCz11), medaka (ASM223467v1),

and platyfish (X_maculatus-5.0) chromosomes using the Genespace R package (LOVELL

et al. 2022). Using the parse annotation function of Genespace we generated the gene bed

file from the gff files of each species. After files were parsed, formatted correctly and all

headers matched in the protein and gene bed file, we initiated Genespace as described in

Lovell (LOVELL *et al.* 2022). We repeated this same Genespace workflow per morph

compared to our surface fish assembly to evaluate possible assembly errors or natural

structural variation (SV).

Gene annotation. Gene annotation of all of our assemblies, in addition to the previously

assembled Pachón (IMARAZENE *et al.* 2021), was carried out by the use of the

standardized Ensembl workflows (AKEN *et al.* 2016). However, only the surface genome

described in this study was also annotated with the NCBI pipeline (PRUITT *et al.* 2014).

Numerous RNAseq data sets that exist in the NCBI sequence archive for surface and

cavefish from different tissue sources were used to improve the accuracy of protein-

coding and non-coding gene model builds. A full accounting of all gene annotations was

generated using AGAT v1.0.0 (J. 2020). We used standard parsing tools to compare lists

of gene symbols in all four annotations to one another. Here, we assumed homology of

matching gene symbols and ignored duplicates.

Gene orthology. To compare gene orthology with other vertebrates, we ran OrthoFinder

version 2.5.4 (EMMS AND KELLY 2019) on fifteen species total (

Table 2). To mitigate the effect of multiple transcripts per gene, we used primary

transcripts (the longest version) only (Suppl. Table 2), per OrthoFinder recommendations

(EMMS AND KELLY 2019). MAFFT was used for multiple sequence alignment and

Fasttree for gene tree inference in the OrthoFinder analysis. We ran a secondary analysis

for comparison using IQtree for gene tree inference. We generated orthogroups across all

species which included protein sequences from primary transcripts of *A. mexicanus*

surface genome, zebrafish (*Danio rerio*), Japanese medaka (*Oryzias latipes*), platyfish

(*Xiphophorus maculatus*), spotted gar (*Lepisosteus oculatus*), eastern brown snake

(*Pseudonaja textilis*), common wall lizard (*Podarcis muralis*), chicken (*Gallus gallus*),

human (*Homo sapiens*), chimpanzee (*Pan troglodytes*), mouse (*Mus musculus*), rat

(*Rattus norvegicus*), gray short-tailed opossum (*Monodelphis domestica*), western clawed

frog (*Xenopus tropicalis*), and a tunicate, *Ciona intestinalis,* as the outgroup. These taxa

were chosen 1) because they all have available Ensembl annotations, thus easily

compatible with Orthofinder algorithms, and 2) establish an even phylogenetic sampling

across amniotes and anamniotes. In addition to the four teleosts, we include the spotted

gar because this species originated prior to the teleost whole genome duplication, making

it an important bridge for ortholog predictions between teleosts and other vertebrates

(BRAASCH *et al.* 2016). We collated orthogroups that contain *A. mexicanus* sequences

lacking gene symbols, so that gene symbol annotation can be inferred from that of other

members in the orthogroup.

Structural variation analysis. To initially estimate the distribution and number of SVs

present in these independently derived cave morph genomes we used Assemblytics 1.2.1

(NATTESTAD AND SCHATZ 2016), that estimates tandem or repeat sequence contractions

and expansions, as well as deletions or insertions when compared to the surface genome

(NATTESTAD AND SCHATZ 2016). This approach can reflect uniqueness to a cave morph

or similarity across morphs when compared to the surface genome. The nucmer

algorithm of MUMmer release 4.x (KURTZ *et al.* 2004) was run to align assemblies at

their contig level to minimize false positives (NATTESTAD AND SCHATZ 2016). The

resulting delta file was input into the Assemblytics browser and run using default
parameters: a unique contig sequence anchoring size of 10,000 bp, and the variants
classified by type and size ranging from 50-500 bp and 500-10,000 bp for plot
visualization.
182
Results and Discussion
Chromosome-scale assembly of the surface and cavefish forms. We generated reference
genomes from single lab-reared *A. mexicanus* surface or cave morphs that are descended
from varying Mexico localities, representing independent origins of the cave form from
Tinaja and Molino populations as well as a surface Río Choy population. We sequenced
and assembled each genome using SMRT CCS (circular consensus sequencing) with a
genome coverage depth of 27.4 to 46.2 (RUAN AND LI 2020) (Suppl. Table 1). The final
de novo assemblies resulted in ungapped assembly length ranges of 1.36-1.41 Gb, total
contigs of 123-292, and N50 contig lengths of 12-47Mb (Table 1). Using on average 200
million 150bp Hi-C reads for chromosome-scale scaffolding resulted in the generation of
25 total chromosomes for the surface and each cave morph. Due to the exceptional
contiguity of the surface assembly (47 Mb N50 contig length), we established the
chromosome structure of this assembly first, then investigated discrepancies of the other
cavefish assemblies using multiple pairwise alignments, including the recently published
long-read assembly of the Pachón morph (IMARAZENE *et al.* 2021). Few order or
orientation errors were found and corrected; however, three chromosomes with complete
orientation differences in all cave forms must be further investigated for possible curation
fixes (Suppl. Fig. 1). The surface genome contiguity surpassed all our cave morphs and,
to our knowledge, all long-read assembled teleost genomes to date found in the NCBI
assembly archive, including the zebrafish (fDanRer4.1). Across all assemblies, only 3.7
to 7.4% of sequences could not be properly assigned to chromosomes. These new
assemblies contribute 90 Mb in new sequence on average and show a substantial
reduction (25-fold) in assembly gaps, when evaluated against the prior version of the
surface fish genome (A. mexicanus-2.0; Table 1). Our surface assembly exhibits a high
level of contiguity and completeness, despite that the total interspersed repeats for the
surface genome was 45.8%, which is only slightly lower than zebrafish's 48.4% with a

similar estimated genome size of 1.4 Gb.

Structural differences. One question we wished to address was: is gene order highly

conserved among surface and cave morph chromosomes despite their demographic origin

differences (Fig. 1A) and up to 190,000 generations of cave morph divergence from their

surface ancestor (MORAN R.L. 2022). We find no major chromosomal discrepancies in

chromosome order when performing pairwise comparisons of the Pachon, Tinaja, or

Molino cave morphs to the representative surface fish ancestor (Suppl. Fig. 1). Also, of

interest was: are there examples of unexpected teleost interspecies conservation in

chromosome gene order that can aid future studies aimed at understanding cave morph

standing genetic variation. The alignments of all 25 *A. mexicanus* chromosomes to three

distantly related teleost genomes (zebrafish, platyfish, and medaka) using aligned genes

are expected to display patterns of organizational divergence dependent on their

phylogenetic relationships. Overall, 56% of *A. mexicanus* chromosomes display complex

genomic rearrangements relative to all three distantly related teleost species (Fig. 1B). In

one example, the largest *A. mexicanus* chr1 (134 Mb), aligns with four zebrafish

chromosomes: 5, 16, 20, and 24 (Fig. 1C), suggesting multiple fissions or fusions

occurred throughout teleost genome evolution. In contrast, there were some

chromosomes that are nearly syntenic for gene order, such as *A. mexicanus* chr14 versus

zebrafish chr17 (Fig. 1C). A separate pairwise alignment of the surface fish and zebrafish

genomes (Danio rerio GRCz11) support these findings of variable synteny (Suppl. Fig.

2). These teleost interspecies alignments confirm earlier studies, and highlight the

complex trajectory in the evolution of the teleost genome (VOLFF 2005) (BRAASCH *et al.*

2016).

Gene annotation. We first assessed the sequence completeness of the assemblies using

benchmarking universal single-copy ortholog (BUSCO) (SIMAO *et al.* 2015) scores and

found on average 96.8% of BUSCOs present in their complete and unfragmented form,

2.4% missing, and 1.3% duplicated (Suppl. Table 3). Protein-coding genes from all four

genomes were predicted using Ensembl (AKEN *et al.* 2016) with the average being

26,974. The surface genome was also annotated using the NCBI workflow, with small

differences in the total protein-coding genes count (Suppl. Table 4). In contrast, large

differences were seen in the surface genome pseudogene annotation, 1,376 versus 183,

when comparing the NCBI and Ensembl output (Suppl. Table 4). Interestingly, a small

increase in Ensembl predicted pseudogenes in cave morphs compared to the surface is

observed (Suppl. Table 4) apart from Pachon. The correct annotation of pseudogenes

"non-functional" genes across species is a persistent problem, and especially deserves

further attention in cave species genomes given its importance in understanding

troglomorphic genetic adaptation (HARRISON 2021). While all these newly annotated *A.*

mexicanus genomes are improved over prior assemblies (WARREN *et al.* 2021), estimates

of non-coding genes were substantially improved (a 36-fold increase; Suppl. Table 4),

owing mostly to improved assembly contiguity and accuracy. In total, all these surface

and cave morph gene sets show Molino, Tinaja, and Surface have similar numbers of

mRNAs, exons, CDSs, and total CDS lengths, but in most comparisons, the increased

surface genome contiguity resulted in improved gene annotation, supporting the use of

AstMex3_surface as the standard reference for future *A. mexicanus* computational

experiments (Suppl. Table 5).

In pairwise comparisons, the surface genome had 1,041, 978, and 1,054 gene

symbols not found in Pachon, Molino, and Tinaja annotations, respectively (Suppl. Table

6). Similarly, each cave population had over 800 gene symbols not found in the surface

genome.

Gene Orthology. We targeted a specific span of model species across the vertebrate

phylogeny to classify potential orthologs (Fig. 2A). By analyzing all detected gene

ortholog relationships, we were able to better understand the utility of *A. mexicanus* for

comparative inference across aquatic models and other vertebrates (Fig. 2B). In the

OrthoFinder analysis, 95% of all genes were assigned to 20,612 orthogroups (Suppl.

Table 7; Fig. 2C). Fifty percent of genes assigned to orthogroups are in orthogroups of 19

genes or more, and over 25% of orthogroups contained all species (n = 5,285) (Fig. 2D),

of which 20.8% are single copy orthogroups (n = 1,097) (Suppl. Table 7; Fig. 2E). An

additional 14.4% of orthogroups (n = 2,988) contained at least one copy from every

vertebrate species and zero copies from the outgroup (Fig. 2B). All vertebrates had >96%

of their genes assigned to orthogroups (Fig. 2C; Suppl. Table 8). The outgroup, *C.*

intestinalis, had 66.8% of genes assigned to orthogroups (Suppl. Table 8). For *A.*

mexicanus, 96.3% of genes were assigned to orthogroups (Suppl. Table 8). We found

more one to one orthologs of *A. mexicanus* with humans than for zebrafish with humans

(n = 6,341 and n = 5,118, respectively; Fig. 2E). We identified 378 orthogroups between

A. mexicanus and humans that lack zebrafish orthologs (Fig. 2E). Overall, more

orthogroups contain genes from *A. mexicanus* and humans (n = 11,444) than zebrafish

and humans (n = 11,272). Addressing specifically orthologous relationships, 16,582

human genes (70.5%) have an ortholog with *A. mexicanus*, whereas 16,221 human genes

(68.9%) have an ortholog with *Danio rerio*. These findings increase the breadth of

orthologs available in an alternative fish model for comparative trait dissection. Zebrafish

have more many-to-one and many-to-many orthologs with humans and more species-

specific orthogroups than *A. mexicanus* (n = 245 and n = 157, respectively; Fig. 2E). In

general, the availability of higher-quality genome assemblies is revealing teleost whole

genome duplication to be a pervasive source of genetic variation across these taxa

(ALBALAT AND CANESTRO 2016), and understanding the main evolutionary forces

impacting *A. mexicanus* gene losses or gains is beyond the scope of this study (ADRIAN-

KALCHHAUSER *et al.* 2020). In identifying orthologs, we also estimated the number of

duplications across the species tree, both at the tips and internal nodes which could

represent differential genome fractionation across lineages stemming from the teleost

whole genome duplication or lineage-specific duplications. At the base of teleosts, our

analysis identified 2,451 duplications. In the tip branches of teleosts, we identified 2,647,

2,825, 4,812, and 8,509 duplicates for *X. maculatus*, *O. latipes*, *A. mexicanus*, and *D.*

rerio, respectively. We suspect that teleost whole genome duplication, annotation bias,

phylogenetic sampling, and multiple evolutionary events or forces underpin these results.

Five orthogroups (OG0000022, OG0000023, OG0000045, OG0000050, and

OG0000086) each have over 100 genes from *A. mexicanus*. These uniquely large copy

numbers suggest that these genes experienced unique evolutionary pressures in *A.*

mexicanus. The orthogroup with the largest number of *A. mexicanus* gene copies is

OG0000022 (n = 215). This group and the group with the second largest number of *A.*

mexicanus genes, OG0000023 (n = 205), are dominated by *A. mexicanus* zinc finger

proteins. Additional studies using independent tests of gene family expansions and

contractions, such as CAFE (Mendes et al. 2021), will be needed to truly resolve

duplication events prior to explorations of their functional roles in the *A. mexicanus* cave

adaptation. We provide gene symbols for human and *A. mexicanus* orthologs as a

resource for comparative studies (Suppl. Table 9). These *A. mexicanus* protein-coding

gene resources further efforts to improve genome annotation for *A. mexicanus*, offering

another aquatic model beyond zebrafish for human comparative studies. Future research

to identify gene duplicates, pseudogenization events, and nonsynonymous protein coding

changes unique to *A. mexicanus* within a framework of surface and independent cave

morphs will address hypotheses about how this species was adaptable to extreme

environmental change.

Structural variation. Cave morphs display many phenotypic differences from their

surface ancestor that motivated us to evaluate genome evolution on a finer scale beyond

chromosomal gene order evaluated herein and prior studies that have focused on protein-

coding gene changes (WARREN *et al.* 2021) (MORAN R.L. 2022). Discovery of SVs

among cave morphs compared to surface has only been estimated using short reads to a

less contiguous reference genome (WARREN *et al.* 2021). To evaluate moderately-sized

SVs (50 to 10,000 bp) among these cave morphs compared to the surface genome, we

estimated the presence of deletions, insertions, and contractions and expansions of

repeats using Assemblytics (NATTESTAD AND SCHATZ 2016). The number of insertions

and deletions shows comparable total bases affected across these cave morphs relative to

surface regardless of their sequence length distribution (500-10,000 bp Fig. 3;50-500 bp

Suppl. Fig. 3; Suppl. Table 10). The average total size of detected cave morph deletions

and insertions was 19.4 MB (1.42% of the genome size) when aligned to the surface

genome (Suppl. Table 10). SV divergences unique to each cave morph were also evident.

For example, the Tinaja morph had the highest total sequence size for all insertions (12.9

MB), followed by Pachón (12.8 Mb) and Molino (10.8 Mb) (Suppl. Table 10). Similarly

for all deletions, Tinaja was highest (7.54 MB), then Pachón (7.10 Mb), and Molino

(7.08Mb) (Suppl. Table 10). Interestingly, total inserted sequence by size distribution

changes cave morph order with Pachón (9.9 Mb) having the largest amount for the 500 to

10,000 bp range followed by Tinaja (9.7 Mb) (Suppl. Table 10). When evaluating the
cave morph repeat landscape, we find an even larger percentage of the genome impacted
relative to insertions and deletions, with the average repeat expansion or contraction
being 3.05 and 2.91% of the total genome, respectively (Suppl. Table 10). These broadly
classified SV and repeat results highlight their differences vary by category and cave
morph origin, which may be the result of numerous factors, including assembly
completeness and accuracy, mixed haplotype assembly architecture, and the diversified
origins of each cave morph reference (HERMAN *et al.* 2018).

Study conclusions. The surface-to-cave genomic transitions that occurred in *A. mexicanus*
establish a unique model for the study of natural polygenic trait adaptation. Here, we
provide initial evidence that these more complete genomes will substantially advance our
capability to resolve these signatures of genetic adaption. The availability of nearly
complete genome copies of a surface and the independently evolved cave morphs will
drive future reevaluations of all types of segregating single nucleotide variants and SVs
in a pangenome-dependent manner (SIREN *et al.* 2021).

Acknowledgments. We thank the Stowers aquatics group for fish care and help with
shipments of fish samples. AK, SEM, NR are supported by NIH 1R01GM127872-01.
This work was also supported by NIH R24OD011198 to WCW, AK, NR. NSF EDGE
Award 1923372 to NR, SEM, and NSF IOS-1933428 to SEM, NR. Some computation
for this work was performed on the high-performance computing infrastructure provided
by Research Computing Support Services and in part by the National Science Foundation
under grant number CNS-1429294 at the University of Missouri, Columbia MO. The
Minnesota Supercomputing Institute (MSI) at the University of Minnesota provided
resources that contributed to the research results reported within this paper.

Data availability. All raw and processed data for this study are available by querying

NCBI BioProject accession numbers PRJNA807270, PRJNA819394, and

PRJNA819399. In addition, each assembly is available under GenBank numbers

GCA_023375975.1, GCA_023375835.1, and GCA_023375845.1. The full availability of

orthofinder results are attainable on figshare.

Code availability. Scripts used for this study are available at the following GitHub
repositories:
https://github.com/esrice/hic-pipeline
https://github.com/WarrenLab/purge-haplotigs-nf
https://github.com/WarrenLab/agptools

Literature cited

Adrian-Kalchhauser, I., A. Blomberg, T. Larsson, Z. Musilova, C. R. Peart *et al.*, 2020
The round goby genome provides insights into mechanisms that may
facilitate biological invasions. BMC Biol 18**:** 11.
Aken, B. L., S. Ayling, D. Barrell, L. Clarke, V. Curwen *et al.*, 2016 The Ensembl gene
annotation system. Database 2016.
Albalat, R., and C. Canestro, 2016 Evolution by gene loss. Nat Rev Genet 17**:** 379-391.
Aspiras, A. C., N. Rohner, B. Martineau, R. L. Borowsky and C. J. Tabin, 2015
Melanocortin 4 receptor mutations contribute to the adaptation of cavefish
to nutrient-poor conditions. Proc Natl Acad Sci U S A 112**:** 9668-9673.
Bilandžija, H., Hollifield, B., Steck, M., Meng, G., Ng, M., Koch, A.D., Gračan, R.,Ćetković,
H., Porter, M.L., Renner, K.J., Jeffery, W.R., 2019 Phenotypic plasticity as an
important mechanism of cave colonization and adaptation in Astyanax
cavefish. bioRxiv.
Braasch, I., A. R. Gehrke, J. J. Smith, K. Kawasaki, T. Manousaki *et al.*, 2016 The
spotted gar genome illuminates vertebrate evolution and facilitates human-
teleost comparisons. Nat Genet 48**:** 427-437.
Cheng, H., G. T. Concepcion, X. Feng, H. Zhang and H. Li, 2021 Haplotype-resolved de
novo assembly using phased assembly graphs with hifiasm. Nat Methods 18**:**
170-175.
Chernyavskaya, Y., X. Zhang, J. Liu and J. Blackburn, 2022 Long-read sequencing of
the zebrafish genome reorganizes genomic architecture. BMC Genomics 23**:**
116.
Du, K., M. Pippel, S. Kneitz, R. Feron, I. da Cruz *et al.*, 2022 Genome biology of the
darkedged splitfin, Girardinichthys multiradiatus, and the evolution of sex
chromosomes and placentation. Genome Res 32**:** 583-594.
Duboue, E. R., A. C. Keene and R. L. Borowsky, 2011 Evolutionary convergence on
sleep loss in cavefish populations. Curr Biol 21**:** 671-676.
Durand, N. C., J. T. Robinson, M. S. Shamim, I. Machol, J. P. Mesirov *et al.*, 2016
Juicebox Provides a Visualization System for Hi-C Contact Maps with
Unlimited Zoom. Cell Syst 3**:** 99-101.
Emms, D. M., and S. Kelly, 2019 OrthoFinder: phylogenetic orthology inference for
comparative genomics. Genome Biol 20**:** 238.
Ghurye, J., and M. Pop, 2019 Modern technologies and algorithms for scaffolding
assembled genomes. PLoS Comput Biol 15**:** e1006994.
Ghurye, J., A. Rhie, B. P. Walenz, A. Schmitt, S. Selvaraj *et al.*, 2019 Integrating Hi-C
links with assembly graphs for chromosome-scale assembly. PLoS Comput
Biol 15**:** e1007273.
Guan, D., S. A. McCarthy, J. Wood, K. Howe, Y. Wang *et al.*, 2020 Identifying and
removing haplotypic duplication in primary genome assemblies.
Bioinformatics 36**:** 2896-2898.
Harrison, P. M., 2021 Computational Methods for Pseudogene Annotation Based on
Sequence Homology. Methods Mol Biol 2324**:** 35-48.

Herman, A., Y. Brandvain, J. Weagley, W. R. Jeffery, A. C. Keene *et al.*, 2018 The role of
gene flow in rapid and repeated evolution of cave-related traits in Mexican
tetra, Astyanax mexicanus. Mol Ecol 27**:** 4397-4416.
Imarazene, B., K. Du, S. Beille, E. Jouanno, R. Feron *et al.*, 2021 A supernumerary "B-
sex" chromosome drives male sex determination in the Pachon cavefish,
Astyanax mexicanus. Curr Biol 31**:** 4800-4809 e4809.
J., D., 2020 AGAT: Another Gff Analysis Toolkit to handle annotations in any
GTF/GFF format., pp. in *Zenodo*.
Jarvis, E. D., G. Formenti, A. Rhie, A. Guarracino, C. Yang *et al.*, 2022 Semi-automated
assembly of high-quality diploid human reference genomes. Nature 611**:** 519-
531.
Kurtz, S., A. Phillippy, A. L. Delcher, M. Smoot, M. Shumway *et al.*, 2004 Versatile and
open software for comparing large genomes. Genome Biol 5**:** R12.
Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-
Wheeler transform. Bioinformatics 25**:** 1754-1760.
Lovell, J. T., A. Sreedasyam, M. E. Schranz, M. Wilson, J. W. Carlson *et al.*, 2022
GENESPACE tracks regions of interest and gene copy number variation
across multiple genomes. Elife 11.
McGaugh, S. E., J. E. Kowalko, E. Duboue, P. Lewis, T. A. Franz-Odendaal *et al.*, 2020
Dark world rises: The emergence of cavefish as a model for the study of
evolution, development, behavior, and disease. J Exp Zool B Mol Dev Evol
334**:** 397-404.
Moore, B., M. Herrera, E. Gairin, C. Li, S. Miura *et al.*, 2023 The chromosome-scale
genome assembly of the yellowtail clownfish Amphiprion clarkii provides
insights into the melanic pigmentation of anemonefish. G3 (Bethesda) 13.
Moran R.L., R., E.J., Ornelas-García, C.P., Gross, J.B., Donny, A., Wiese, J., Keene, A.C.,
Kowalko, J.E., Rohner, N., McGaugh, S.E., 2022 Selection-driven trait loss in
independently evolved cavefish populations. bioRxiv.
Nattestad, M., and M. C. Schatz, 2016 Assemblytics: a web analytics tool for the
detection of variants from an assembly. Bioinformatics 32**:** 3021-3023.
Ojha, A., and M. Watve, 2018 Blind fish: An eye opener. Evol Med Public Health 2018**:**
186-189.
Pruitt, K. D., G. R. Brown, S. M. Hiatt, F. Thibaud-Nissen, A. Astashyn *et al.*, 2014
RefSeq: an update on mammalian reference sequences. Nucleic Acids Res 42**:**
D756-763.
Rautiainen, M., S. Nurk, B. P. Walenz, G. A. Logsdon, D. Porubsky *et al.*, 2023
Telomere-to-telomere assembly of diploid chromosomes with Verkko. Nat
Biotechnol.
Rhie, A., S. A. McCarthy, O. Fedrigo, J. Damas, G. Formenti *et al.*, 2021 Towards
complete and error-free genome assemblies of all vertebrate species. Nature
592**:** 737-746.
Roberts, M. B., D. T. Schultz, R. Gatins, M. Escalona and G. Bernardi, 2023
Chromosome-level genome of the three-spot damselfish, Dascyllus
trimaculatus. G3 (Bethesda) 13.

Robinson, J. T., D. Turner, N. C. Durand, H. Thorvaldsdottir, J. P. Mesirov *et al.*, 2018
Juicebox.js Provides a Cloud-Based Visualization System for Hi-C Data. Cell
Syst 6**:** 256-258 e251.
Rohner, N., 2018 Cavefish as an evolutionary mutant model system for human
disease. Dev Biol 441**:** 355-357.
Ruan, J., and H. Li, 2020 Fast and accurate long-read assembly with wtdbg2. Nat
Methods 17**:** 155-158.
Simao, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva and E. M. Zdobnov,
2015 BUSCO: assessing genome assembly and annotation completeness with
single-copy orthologs. Bioinformatics 31**:** 3210-3212.
Siren, J., J. Monlong, X. Chang, A. M. Novak, J. M. Eizenga *et al.*, 2021 Pangenomics
enables genotyping of known structural variants in 5202 diverse genomes.
Science 374**:** abg8871.
Stockdale, W. T., M. E. Lemieux, A. C. Killen, J. Zhao, Z. Hu *et al.*, 2018 Heart
Regeneration in the Mexican Cavefish. Cell Rep 25**:** 1997-2007 e1997.
Volff, J. N., 2005 Genome evolution and biodiversity in teleost fish. Heredity (Edinb)
94**:** 280-294.
Warren, W. C., T. E. Boggs, R. Borowsky, B. M. Carlson, E. Ferrufino *et al.*, 2021 A
chromosome-level genome of Astyanax mexicanus surface fish for comparing
population-specific genetic differences contributing to trait evolution. Nat
Commun 12**:** 1447.
Warrenlab., 2022 Nextflow workflow for purging haplotigs from a genome assembly.

Author notes

Competing interests. The authors declare no competing interests.

Figure legends

Figure 1. Summary of interspecies chromosomal synteny. A. Geographical representation
of the independent cave morph populations and their physical appearance used in this
study. Image courtesy of Alex Keene. B. Gene ortholog synteny of *A. mexicanus* against
three teleost species. C. Higher resolution image of *A. mexicanus* (surface genome)
aligned to the same three teleost species in B.

Figure 2. Summary of OrthoFinder analysis of vertebrate species *Astyanax mexicanus*,
Danio rerio, *Oryzias latipes*, *Xiphophorus maculatus*, *Lepisosteus oculatus*, *Pseudonaja*
textilis, *Podarcis muralis*, *Gallus gallus*, *Homo sapiens*, *Pan troglodytes*, *Mus musculus*,
Rattus norvegicus, *Monodelphis domestica*, *Xenopus tropicalis*, and *Ciona*
intestinalis. Bar charts describe data for each species, aligned to the matching species in
the tree. A. Phylogenetic tree built with all species using shared gene orthologs. B.
Number of orthogroups each classified by type per species. C. Percentage of genes by
orthogroups by species. D. Number of species-specific orthogroups per species. E.
Ortholog multiplicities for all species.
505

507 Figure 3. Genomic structural variation among *A. mexicanus* cave morph assemblies when
508 compared to surface. Cave morph specific distribution by counts for A. Molino, B.
509 Tinaja, and C. Pachón for the size distribution of 500 to 10,000 bp.
510
511

512 Table 1. Representative genome assembly metrics for sequenced *A. mexicanus* genomes.

Common name	Assembled version	Total size Mb	Total contigs	Contig N50 length Mb	Unplaced Mb
Surface	Astyanax mexicanus-2.0	1,291	3,030	1.7	394
Río Choy Surface	AstMex3_surface	1,373	123	47	51
Molino	AstMex3_Molino	1,361	252	12	101
Tinaja	AstMex3_Tinaja	1,410	292	26	78
Pachón	AMEX_1.1	1,378	529	14.8	29

A. Species Tree	B. Orthogroup Types	C. % Genes in Orthogroups	D. Species-Specific Orthogroups	E. Multiplicity

Homo sapiens

Species (A):
Astyanax mexicanus
Danio rerio
Oryzias latipes
Xiphophorus maculatus
Lepisosteus oculatus
Pseudonaja textilis
Podarcis muralis
Gallus gallus
Homo sapiens
Pan troglodytes
Mus musculus
Rattus norvegicus
Monodelphis domestica
Xenopus tropicalis
Ciona intestinalis

B. Number of Ogs — 0, 4000, 8000, 12000

C. % — 0, 50, 100

D. Number of Ogs — 0, 200, 400

E. % — 0, 50, 100

Legend (B):
- 1:1 OGs with all species present
- other OGs with all species present
- vertebrates only 1:1 OGs
- other OGs with all vertebrates present
- other OGs containing species

Legend (E):
- 1 : 1
- 1 : many
- many : 1
- many : many

A. Molino   B. Tinaja   C. Pachon

Insertions
Deletions
Repeat expansion
Repeat contraction

Counts

Variant size (bp)