1

Evolution of a small phage protein overcomes antiphage defense in *Enterococcus*

faecalis

Nathan P. Bullen[2,3,*], Cydney N. Johnson[1,*], Shelby E. Andersen[1], Garima Arya[1], Sonia R. Marotta[2,3], Yan-Jiun Lee[4], Peter R. Weigele[4], John C. Whitney[2,3,#], Breck A. Duerkop[1,#]

*Authors contributed equally to this work

[1]Department of Immunology and Microbiology, University of Colorado School – Anschutz Medical Campus, School of Medicine, Aurora, CO, USA, 80045

[2]Department of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, ON, Canada, L8S 4L8

[3]Michael DeGroote Institute for Infectious Disease Research, McMaster University, Hamilton, ON, Canada

[4]Research Department, New England Biolabs, Ipswich, MA, USA, 01938

[#]Correspondence: Breck A. Duerkop, breck.duerkop@cuanschutz.edu or John C. Whitney, jwhitney@mcmaster.ca

Key words: Bacteriophages, *Enterococcus faecalis*, plasmid, antiphage, type IV restriction enzyme

ABSTRACT

The prevalence of multidrug resistant (MDR) bacterial infections continues to rise as the development of new antibiotics needed to combat these infections remains stagnant. MDR enterococci, which are a common cause of hospital-acquired infections, are emerging as one of the major contributors to this crisis. A potential therapeutic approach for combating MDR enterococci is bacteriophage (phage) therapy, which entails the use of lytic viruses to infect and kill pathogenic bacteria. While phages that lyse some strains of MDR enterococci have been identified, other strains display high levels of phage resistance and the mechanisms underlying this resistance are poorly defined. Here, we use a CRISPR interference (CRISPRi) screen to identify a genetic locus found on a mobilizable plasmid from vancomycin-resistant *Enterococcus faecalis* involved in phage resistance. This locus encodes a putative serine recombinase followed by a Type IV restriction enzyme (TIV-RE) and we show that this enzyme is sufficient to restrict the replication of lytic phage phi47 in *E. faecalis*. We further find that phi47 can evolve to overcome restriction by acquiring a missense mutation in a novel TIV-RE inhibitor protein encoded by many enterococcal phages. We show that this inhibitor, which we have named type IV restriction inhibiting factor A (*tifA*), directly binds to and inactivates diverse TIV-REs. Overall, our findings significantly advance our understanding of phage defense in drug-resistant *E. faecalis* and provide mechanistic insight into how phages can evolve to overcome antiphage defense systems.

INTRODUCTION

Typically members of the gut microbiota of humans and other animals, the enterococci

are Gram-positive bacteria that can cause life-threatening disease following antibiotic-mediated

dysbiosis[1,2]. During antibiotic treatment, multidrug-resistant (MDR) *Enterococcus faecalis* can

outgrow antibiotic-sensitive members of the microbiota and this increases the likelihood that

MDR *E. faecalis* will breach the intestinal barrier and enter the bloodstream leading to

infection[1,2]. These MDR infections are difficult to treat due to limited antibiotic options and

consequently, bacteriophages or phages (viruses that infect bacteria) are increasingly being

considered as a therapeutic option as the frequency of MDR infections continues to rise[3-5].

Phage therapy offers advantages over traditional antibiotic therapy because phages exhibit

strain-level specificity, which reduces off-target bacterial killing, they are self-limiting following

the exhaustion of their bacterial host, and their abundance in nature implies that a phage

specific to a target bacterium can usually be identified[6-10].

As is the case with the emergence of antimicrobial resistance, exposure of bacteria to

phages also imposes a strong selective pressure that leads to the evolution or acquisition of

mechanisms to resist infection. Recent studies have shown that bacteria have evolved diverse

antiphage defense systems to prevent successful phage infection[11-13]. These systems are often

concentrated in bacterial genomes within so-called "defense-islands", which are characterized

by the clustering of genes encoding two or more defense systems and are typically carried on

mobile genetic elements, including integrative and conjugative elements, plasmids,

transposons, and temperate phages[14-16]. This capacity to mobilize and spread between bacterial

species represents a robust pan-genomic immune system that contributes to the rapid evolution

of resistance to phage infection and constitutes a significant challenge to the effective

application of phage therapy.

Antiphage defenses utilize a variety of mechanisms to overcome phage infection. These

include restriction-modification systems in which host proteins epigenetically label nucleobases

and/or the phosphate backbone to distinguish host DNA from phage DNA, RNA-based adaptive

nucleic acid targeting systems such as CRISPR-Cas[16-21], abortive infection systems and other

phage interference systems that stop phage replication and spread[22-26], chemical defenses that

restrict phage infection[27], and phage receptor alterations that abrogate the ability of phages to

interact with the bacterial cell surface[28-32]. In turn, phages have evolved counter-defenses to

antiphage defense systems including enzymes that modify tail proteins for bacterial cell

recognition, DNA modification to avoid restriction systems, and phage-encoded proteins that

interfere with antiphage defenses[33-37]. Compared to bacterial antiphage defense systems, the

breadth of phage-encoded proteins that support counter-defenses and the mechanisms that

drive their evolution are poorly understood. Thus, prior to phages being widely used as

therapeutics for MDR bacterial infections, a detailed understanding of the mechanisms by which

bacteria subvert phage infection and, conversely, how phages overcome these mechanisms is

needed.

In this study, we sought to understand why some MDR strains of *E. faecalis* are

susceptible to infection by virulent enterococcal phages whereas others exhibit resistance[38]. By

conducting a CRISPR interference (CRISPRi) screen on one such resistant strain, we pinpoint a

mechanism of resistance to a two-gene locus residing on a pheromone responsive plasmid

(PRP) that inhibits infection of the enterococcal phage phi47. Further analysis revealed that

resistance to phi47 infection is solely conferred by the activity of one of these genes, EF_B0059,

encoding a putative type IV restriction enzyme (TIV-RE). We subsequently isolated phages that

are resistant to restriction by EF_B0059 and discovered that the *orf65* gene of phi47 encodes a

small protein that can evolve to not only bind and inactivate EF_B0059, but also other highly

divergent type IV restriction enzymes. From this, we posit that *orf65* encodes a rapidly evolving

and broadly acting anti-TIV-RE protein. Overall, our data demonstrate that *E. faecalis* utilizes a

mobilizable antiphage system to resist phage infection, and that phages escape this restriction

through a coevolving small open reading frame found in enterococcal and related phages.

RESULTS

The pheromone responsive plasmid pTEF2 is sufficient to inhibit phage infection

In our attempts to understand the mechanisms underlying phage-host interactions in

MDR enterococci, we previously observed that the lytic phage phi47 can infect *E. faecalis* strain

SF28073 but not the closely related strain V583[38]. phi47 readily adsorbs to both SF28073 and

V583, indicating that this resistance cannot be attributed to differences in the cell surface of

these strains preventing phage binding[38]. From this, we hypothesized that *E. faecalis* V583 may

encode one or more antiphage defense systems that restrict the replication of phi47.

A comparison of the *E. faecalis* SF28073 and V583 genomes revealed that they are very

similar (nearly 90% sequence identity across shared sequences) (Fig. S1A). However,

sequence comparison of the three natural plasmids carried by these strains showed that the

plasmids of V583 (pTEF1, pTEF2, pTEF3) differ significantly from that of SF28073 (pSF1,

pSF2, pSF3) (Fig S1B). Based on these observations, we hypothesized that one or more

plasmid(s) of V583 may be responsible for resistance to phi47 infection. To explore this idea, we

infected *E. faecalis* V19, a plasmid-cured derivative of V583, with phi47[39,40]. Unlike V583, V19

supports robust replication of phi47 indicating that one or more of the three V583 plasmids

confers resistance to phi47 (Fig. 1A). Of these three plasmids, pTEF1 and pTEF2 are PRPs that

mobilize through conjugation upon sensing a peptide pheromone signal produced by plasmid-

free cells[41]. In contrast to pTEF1 and pTEF2, pTEF3 is a comparatively small, non-mobilizable

plasmid[42]. Given that plasmid mobilization has been previously demonstrated to disseminate

antiphage defenses[43-45], we began our investigation with pTEF1 and pTEF2.

To determine if an *E. faecalis* V583 mobilizable plasmid mediates resistance to phi47,

we conjugated pTEF1 and pTEF2 into the phi47 susceptible *E. faecalis* V19 strain. Subsequent

infections with phi47 showed pTEF1 had no effect on the ability of phi47 to replicate in V19

whereas the carriage of pTEF2 completely restricted phi47 replication (Fig. 1A). This pTEF2-

mediated resistance to phi47 is independent of the *E. faecalis* strain background because

mobilization of pTEF2 into *E. faecalis* JH2-2, which shares 87% of its genome at a nucleotide

identity of 99% or greater with V583[46], was sufficient to prevent replication of phi47 (Fig. 1A). To

confirm that no infectious phages were being released from pTEF2-containing cells, we also

measured total infectious phi47 particle production during infection (Fig. 1B). While the strains

V19 and SF28073 support the production of infectious phi47 particles, V19 pTEF2 grown in the

presence of phi47 did not show an increase in recoverable phages over time. This data

demonstrates that pTEF2 alone confers resistance to phi47.

Some antiphage defense systems are predicted to function through a mechanism

termed abortive infection whereby the activation of the antiphage system stops the spread of

virus at the expense of the viability of the bacterial host[22,23,47-49]. Such population-level

resistance can be measured by infecting bacteria with phage at high <u>m</u>ultiplicities <u>o</u>f <u>i</u>nfection

(MOI) because this results in the complete collapse of the bacterial culture. In contrast, some

defense systems confer individual-level resistance to phage, often by directly targeting phage

replication processes during early infection. In these instances, bacterial growth upon phage

infection is comparable at low and high MOIs. To determine if phi47 resistance is due to

abortive infection, we measured the optical density (OD_{600}) of *E. faecalis* cultures during

infection by phi47 at an MOI of 10 (Fig. S2A). In doing so, we observed that V19 pTEF2 cells

continued to grow during phi47 infection at a similar rate to that of uninfected V19 pTEF2. Thus,

pTEF2 antiphage defense is not mediated by the death or inability of the bacteria to grow, but

instead provides direct immunity to phi47. Additionally, we confirmed that phi47 adsorbs equally

to the cell surface of both V19 and V19 pTEF2 compared to V583 (Fig. S2B). Taken together,

our data suggests that the pheromone responsive plasmid pTEF2 encodes an antiphage

defense system that prevents phage replication in actively growing cells.

A GmrSD-like protein encoded on pTEF2 is responsible for phage restriction

We next determined which gene(s) on pTEF2 were responsible for the restriction of

phi47. Of the 62 open reading frames (ORFs) encoded by pTEF2, approximately half are

predicted to be involved in the maintenance and conjugation of the plasmid. Furthermore, more

than half of the genes on pTEF2 share 95-97% sequence identity to another *E. faecalis* PRP,

pCF10, which, unlike pTEF2, does not confer resistance to phi47 when conjugated into V19

(Fig. 2A and 2B). Comparing the sequences of pTEF2 and pCF10 revealed that pTEF2 encodes

a unique 15 kb region encompassing 18 ORFs not present in pCF10 (EF_B0048 to EF_B0065).

We reasoned that the gene(s) involved in phi47 restriction were therefore likely encoded by one

or more these 18 ORFs.

To test this prediction, we performed a CRISPR interference (CRISPRi) gene

knockdown screen in *E. faecalis* V19 pTEF2. CRISPRi uses a catalytically inactive Cas9

(dCas9) that binds to target DNA sequences when directed by complementary small guide

RNAs (sgRNAs), resulting in transcriptional repression of target genes[50]. We used plasmids

expressing dCas9 and sgRNAs that targeted each of the 18 ORFs unique to pTEF2 and

transformed these into V19 pTEF2 cells to selectively knock down target gene expression. The

relative viability of the resulting transformants was then measured in the presence or absence of

phi47 (Fig. 2C). Strikingly, under these experimental conditions, the sgRNA-dependent

repression of either EF_B0058 or EF_B0059 in the presence of phi47 resulted in a significant

decrease in cell viability when compared to V19 pTEF2. V19 transformed with an sgRNA

targeting *gfp* (sgRNA-GFP) was used as a non-specific control. To validate the findings of our

screen, we generated single gene knock outs of either EF_B0058 or EF_B0059 in both V583

and V19 carrying pTEF2 (Fig. 2D and 2E). Subsequent infections with phi47 in these strains

revealed that the deletion of EF_B0058 did not restore the ability of phi47 to replicate in V583 or

V19 whereas deletion of EF_B0059 did. This result indicates that EF_B0059 alone confers

resistance to phi47 infection. A closer examination of the genetic context of EF_B0058 and

EF_B0059 found that these genes are encoded in tandem and are separated by a small

intergenic region (54 base pairs), suggesting that they may be encoded on the same mRNA

(Fig. 2F). To test this prediction, we performed reverse-transcription PCR to generate cDNA

from total RNA purified from *E. faecalis* V19 with and without pTEF2. PCR reactions using this

cDNA as a template and primers that span the intergenic region between EF_B0059 and

EF_B0058 revealed that EF_B0058 and EF_B0059 exist on the same mRNA (Fig. 2G). From

this finding, we concluded that the sgRNA targeting EF_B0058 likely also disrupts the

expression of downstream EF_B0059, resulting in a reduced capacity to restrict phi47.

Bioinformatic analyses, including the generation of a high-confidence Alphafold2 model

of EF_B0059, predict that this antiphage protein belongs to the GmrSD-like type IV modification-

dependent restriction enzyme family (Fig. S3A and S3B). GmrSD family proteins are defined by

N-terminal DUF262 and C-terminal DUF1524 domains connected by a small α-helical domain of

variable length (Fig. S3A)[51]. DUF262 domains function as NTPases and contain a conserved

(I/V)DGQQR motif that forms a nucleotide binding pocket whereas DUF1524 domains are

structurally similar to HNH endonucleases and contain a conserved EHxxP catalytic motif found

in members of the broad His-Me finger endonuclease superfamily[51,52]. EF_B0059 possesses an

intact DUF262 VDGQQR motif but has a glutamate in place of the conserved proline in its

DUF1524 EHxxP motif (Fig. S3A). The precise molecular details describing how these two

domains coordinate DNA substrate recognition and cleavage are not yet understood, however,

NTP hydrolysis by the N-terminal DUF262 has been shown to drive conformational changes

that activate DUF1524's nuclease activity resulting in DNA cleavage[51,53]. Moreover, these

enzymes have been shown to target diverse types of modified DNA, in particular DNA

containing modified cytosine bases including 5-methylcytosine[54,55]. Given this precedent, we

expressed EF_B0059 in *E. coli* XL1-Blue, a K-12 derivative strain that contains 5-

methylcytosine and N6-methyladenosine, under the assumption that overexpression of this

putative TIV-RE would target the DNA of an evolutionarily distant bacterial species. In line with

our hypothesis, we found that the heterologous expression of EF_B0059 is extremely toxic to *E.*

coli XL1-Blue (Fig. 2H and 2I). To test our assumption that EF_B0059 recognizes modified DNA

for restriction, we repeated these intoxication experiments using the *E. coli* strain ER2796,

which lacks the three active methyltransferases typically found in K-12 derivative strains,

namely the Dam, Dcm, and *Eco*KI methylases. Surprisingly, we found that this strain still

succumbs to intoxication by EF_B0059 (Fig. S4A). These data, combined with sequence

analyses suggesting that the phi47 genome does not encode DNA modifying enzymes, indicate

that EF_B0059 may not require modified phage DNA for restriction. To explore this possibility,

we next examined the chemical structure of phi47 DNA. To this end, we isolated and digested

DNA from phi47 virions and analyzed the resultant nucleotides by high-performance liquid

chromatography coupled with mass spectrometry (Fig. S4B). Comparison of the phi47 DNA

digestion products to that of nucleotide standards revealed that phi47 DNA is comprised of

unmodified nucleotides. We also considered the possibility that EF_B0059 may act on phage

DNA modifications conferred by the host cell from which the phage is propagated. To test this,

we infected *E. faecalis* strains SF28073 and JH2-2 with phi47 and then measured the ability of

the output phages to replicate in *E. faecalis* JH2-2 + pTEF2 (Fig. S4C). Regardless of the host

strain used to propagate phi47, EF_B0059 prevented phage replication. When taken together

with our biochemical data, this observation strongly suggests that phi47 DNA does not modify

its DNA and that restriction of phage replication by EF_B0059 occurs through a modification-

independent mechanism. Importantly, we found that EF_B0059 function is dependent on both

its NTPase and nuclease activities because inactivating mutations to the conserved catalytic

sites of either domain were non-toxic to *E. coli* XL1-Blue (Fig. 2H and 2I). Thus, EF_B0059

restriction presumably arises through its enzymatic activity acting on the phage genome.

However, the mechanism by which EF_B0059 inhibits phi47 replication and how it discriminates

between phage and host DNA remain to be determined.

pTEF2-mediated resistance is overcome by a phage-encoded protein

Because phages are entirely dependent on their host for replication, they are under

strong selective pressure to evolve ways to evade host defense systems. Thus, we next asked if

phi47 can evolve to overcome EF_B0059-dependent restriction. To determine this, three phi47

single-plaque isolates were independently raised and purified to a titer of $\sim10^{12}$ PFU/mL. 1×10^9

PFU of each isolate was plated on *E. faecalis* V19 pTEF2 and two resistant plaques were

isolated for each replicate. Subsequent infections with the phages isolated from each of these

plaques confirmed that they replicate equally on V19 and V19 pTEF2 (Fig. S5A). Intriguingly,

whole genome sequencing of these phages found that they all harbor a single point mutation in

the small open reading frame, *orf65*, that results in a nonsynonymous mutation at amino acid 32

from leucine to phenylalanine (Fig. 3A, Table S1). Mutations in phage genomes can arise to

prevent DNA targeting defense systems from recognizing phage DNA substrate[56,57]. However, it

is unlikely that any single change in DNA sequence would confer resistance given that

restriction enzymes frequently target multiple sites within a phage genome[54,55]. We reasoned

that the gene product of the mutated *orf65* may instead be responsible for evading EF_B0059

activity. To test this hypothesis, we infected *E. faecalis* V19 pTEF2 cells ectopically expressing

either the wildtype or mutated Orf65 (Orf65^{L32F}) with phi47 (Fig. 3B). These results confirmed

that Orf65^{L32F}, but not the wild-type protein, confers resistance to EF_B0059-dependent

restriction.

To understand how this phage protein prevents restriction by EF_B0059 we next used

Alphafold2 to generate a high confidence model of Orf65[L32F] (Fig. 3C and S5B)[58-60]. Orf65 is

predicted to be a small protein consisting of two globular domains connected by a short linker.

Structural alignments to proteins of known structure identified minimal overall similarity to

characterized proteins, although the N-terminal domain within which the L32F mutation is

localized bears some structural resemblance to SH3 domains of several eukaryotic proteins with

the top scoring hit being the SH3 domain of myosin IB from *Entamoeba histolytica* (rmsd 2.3 Å,

9% sequence identity). In bacteria, SH3-like domains bind diverse ligands and molecules in a

variety of unrelated proteins and thus this similarity provides minimal functional insight into how

Orf65 confers resistance to EF_B0059[61]. However, in examining the predicted structure of

Orf65, we also noted the presence of an unusually high number of acidic residues (21.3%) that

are found on the surface of the protein resulting in a large electronegative surface potential (Fig.

3D). This high degree of surface electronegativity is reminiscent of the anti-restriction protein

Ocr of phage T7. Ocr is a negatively charged protein (29% acidic amino acids) whose structure

mimics B-DNA, which facilitates its binding and inactivation of Type I restriction enzymes by

competing with their DNA substrates[62,63]. Although the predicted structure of Orf65[L32F] does not

resemble Ocr or mimic the structure of B-DNA, its similar electronegative surface properties led

us to hypothesize that it may function as a direct inhibitor of EF_B0059.

If our hypothesis is correct, we would expect Orf65[L32F] to inhibit the activity of EF_B0059

in the absence of any additional phi47 or *E. faecalis* encoded factors. To test this, we

heterologously co-expressed EF_B0059 with either wildtype Orf65 or the Orf65[L32F] variant in *E.*

coli XL1-Blue. Consistent with our phi47 infection data in *E. faecalis*, expression of the Orf65[L32F]

variant but not the wildtype protein restored *E. coli* growth when co-expressed with EF_B0059

(Fig. 3E and 3F). Next, to determine if the observed Orf65[L32F] inactivation is dependent on a

physical interaction with EF_B0059, we immunoprecipitated FLAG-tagged EF_B0059 from cells

co-expressing either VSV-G-tagged Orf65 or VSV-G-tagged Orf65[L32F] (Fig. 3G). Western

blotting of the precipitated fractions showed that wild-type EF_B0059 co-precipitates with a

small amount of Orf65, which our co-expression data indicate is insufficient to alleviate the toxic

effects of EF_B0059 on *E. coli* (Fig. 3E). However, we found that Orf65[L32F] was substantially

more enriched than the wildtype protein upon immunoprecipitation suggesting that the inhibitory

properties of this variant are due to enhanced binding to EF_B0059. Importantly, in the absence

of EF_B0059, neither Orf65 nor Orf65[L32F] precipitate with α-FLAG conjugated agarose,

confirming that the observed interactions are specific (Fig. S5C). Overall, these data suggest

that while the wildtype Orf65 interacts with EF_B0059, the L32F mutation likely increases its

binding affinity. Interestingly, mutation of leucine 32 to alanine or to other non-phenylalanine

aromatic residues does not result in Orf65 variants capable of binding and inactivating

EF_B0059, suggesting that the presence of phenylalanine at this position specifically stabilizes

the Orf65[L32F]-EF_B0059 complex (Fig. S4E and S5D). Taken together, our data indicate that

phi47 encodes a small, negatively charged protein that can evolve to inhibit the activity of

EF_B0059. Furthermore, protein-protein interaction studies demonstrate that this inhibition is

mediated by the direct binding of Orf65 to EF_B0059 and that this interaction is strongly

influenced by a phenylalanine substitution at position 32 of Orf65.

Orf65 represents a family of phage-encoded proteins that inhibit type IV restriction
enzymes.

The isolation of phage mutants that can infect phi47-resistant *E. faecalis* pTEF2

revealed that phi47 can evolve to overcome the restriction activity of EF_B0059. It achieves this

through a single point mutation in the small protein Orf65, which promotes the binding and

inactivation of EF_B0059. While wildtype Orf65 also interacts with EF_B0059, it does so with

qualitatively lower affinity and does not display the same inhibitory effect as Orf65[L32F].

Intriguingly, sequence comparisons between Orf65 homologues found in the genomes of

enterococcal phages shows that phenylalanine naturally occurs at position 32 in the *E. faecalis*

phage SDS2, suggesting that phages related to phi47 may have previously evolved to

overcome restriction by EF_B0059 and/or homologs of this TIV-RE (Fig. 4A). This observation

suggests that the L32F substitution we isolated may be representative of an ongoing

evolutionary arms race between phages that encode *orf65* and the hosts that use EF_B0059

and related proteins to restrict their growth. To probe this experimentally, we first asked if Orf65

from phage SDS2, which contains 84% sequence identity to Orf65 from phi47, can inactivate

EF_B0059. However, owing to factors unrelated to the activity of EF_B0059, we found that

phage SDS2 was unable to infect the phi47-susceptible strain V19 or the phi47-resistant strain

V583. Further, we found that over-expression of the SDS2 Orf65 homolog is toxic to *E. coli*

thereby precluding us from testing if it can inactivate EF_B0059 in this heterologous expression

system (Fig. S6A). However, if EF_B0059 and Orf65 are co-evolving together as an effector-

inhibitor pair, we reasoned that the wildtype Orf65 protein is likely already effective at inhibiting

an EF_B0059-related restriction enzyme without requiring any amino acid substitutions. To test

this hypothesis, we searched a database of human intestinal bacterial isolate genomes using

the EF_B0059 sequence to identify candidate homologues[64]. As expected, genes from

enterococci and related bacteria were enriched in this context. From this search, we identified

two predicted TIV-REs including GC1503 from *Clostridium sporogenes,* and GC874 from

Leuconostoc mesenteroides. These homologues share 57% sequence similarity to one another

and approximately 40% sequence similarity to EF_B0059. Despite these relatively low

sequence similarities, comparison of the Alphafold2 predicted structures of these proteins with

that of EF_B0059 suggests that all three enzymes are structurally very similar (Fig. S7).

Heterologous expression of these proteins in *E. coli* demonstrated that like EF_B0059,

GC1503 is toxic to *E. coli*, whereas GC874 is not, suggesting GC1503 functions more similarly

to EF_B0059 (Fig. S6B). Using this same assay, we asked if the co-expression of GC1503 with

either Orf65 or Orf65[L32F] restores *E. coli* growth. Remarkably, and in contrast to EF_B0059, we

found that both wildtype and the L32F mutant of Orf65 were able to restore *E. coli* viability (Fig.

4B and S6C). Moreover, co-immunoprecipitation experiments using FLAG-tagged GC1503

revealed that wildtype Orf65 forms a complex with GC1503 that is comparable in abundance to

that of the L32F mutant containing complex (Fig. 4C). These data indicate that although position

32 of Orf65 from phi47 is essential to stabilize its interaction with EF_B0059, it has no effect on

its ability to inhibit GC1503. Indeed, each of the L32 variants (Trp, Tyr, and Ala) that we

originally tested with EF_B0059 and were found to be non-protective are still protective against

GC1503 activity (Fig. S6D). Therefore, the molecular mechanism of Orf65 binding to EF_B0059

homologous proteins appears to arise from multiple protein-protein contacts whose significance

at the individual amino acid level differs depending on the TIV-RE in question.

We next asked if the inhibitory activity of Orf65 is generalizable to more divergent TIV-

REs. To this end, we tested if the expression of Orf65 or its L32F variant interferes with the

activity of BrxU, a recently characterized TIV-RE from the Gram-negative bacterium *Escherichia*

fergusonii, which shares only 26% sequence similarity with EF_B0059[55]. Remarkably,

expression of either Orf65 or Orf65[L32F] interfered with the ability of BrxU to restrict the infection

of *E. coli* by a waste-water isolated coliphage with wildtype Orf65 exhibiting a greater inhibitory

effect than Orf65[L32F] (Fig. 4D). In accordance with these data, co-immunoprecipitation

experiments with FLAG-tagged BrxU found that both wildtype and Orf65[L32F] interact with BrxU.

However, in contrast to EF_B0059, the L32F mutation had a negative effect on the capacity of

Orf65 to bind BrxU (Fig. 4E).

Together, our data support the conclusion that Orf65 of phi47 encodes a novel anti-TIV-

RE factor. The results of our *E. coli* toxicity data in combination with our phage evolution

experiments suggest that these phage proteins retain the ability to bind and inactivate diverse

TIV-REs, but can evolve to become efficient inhibitors upon selection by a particular TIV-RE.

Due to this newly defined role in overcoming anti-phage defense, we propose to name Orf65

type IV restriction inhibiting factor A (TifA).

DISCUSSION

The need for alternative therapies to combat MDR bacterial infections has renewed

interest in understanding how bacteria defend against phage infection. The majority of this

research has prioritized the discovery and characterization of antiphage defense systems in

model bacteria such as *E. coli* and *Bacillus subtilis,* with minimal focus on systems found in

medically relevant and less-genetically tractable bacteria[20,21,25,65,66]. Moving forward there is a

need to prioritize the characterization of such systems in bacteria that are responsible for MDR

infections that will ultimately be the targets of phage therapy. In this study, we identify a

previously unknown TIV-RE encoded on the enterococcal plasmid pTEF2 that limits phi47

phage replication. phi47 can overcome this restriction via a single point mutation of leucine to

phenylalanine (L32F) in the previously uncharacterized protein Orf65, which herein has been

renamed TifA. This mutation results in a gain-of-function inhibitory protein-protein interaction

between TifA and EF_B0059. Intriguingly, EF_B0059 inhibition strictly depends on a

phenylalanine at position 32 and is lost when other aromatic amino acids are substituted in this

position. However, wildtype TifA binds and neutralizes diverse TIV-REs, suggesting that TifA

represents a broadly active inhibitor of TIV-REs that undergoes adaptive evolution.

pTEF2 belongs to a family of plasmids known as pheromone responsive plasmids

(PRPs) that can be mobilized among enterococcal species. PRPs frequently carry genes

required for antimicrobial resistance and pathogenicity and thus play a crucial role in virulence[67].

Here, we show these plasmids also contribute to the ongoing battle between enterococci and

their phage parasites by encoding dedicated antiphage defense systems. Considering the

pervasiveness of phage predation within complex microbial ecosystems such as the mammalian

gut combined with the proficient PRP-mediated genetic transfer observed in these contexts, we

speculate that PRPs contribute significantly to the ability of the enterococci to counter phage

threats, potentially serving both as a reservoir for and mechanism to mobilize antiphage defense

systems[68,69].

We further discovered that within pTEF2, EF_B0059 is found in an operon with

EF_B0058, which is predicted to encode a serine recombinase. Such recombinases have well-

established roles as mediators of genotypic diversity and population-level fitness in bacteria[70].

Recent studies indicate that recombinases are frequently found in phage defense islands, and

although their direct contribution to the mobilization of antiphage defenses remains

unconfirmed, they are presumed to aid in the diversification of defense islands via homologous

recombination[14]. Indeed, so-called defense islands are commonly used to discover novel

antiphage systems, however, the mechanisms that drive their genetic variation and mobility are

poorly understood. It will be of interest in future studies to probe if and how such recombinases

contribute to the diversification of antiphage defense systems in the enterococci and more

broadly across bacterial taxa.

We found that EF_B0059 provides *E. faecalis* resistance to infection by the lytic phage

phi47. The precise mechanism by which EF_B0059 prevents phi47 replication is unknown,

however, we found that expression of EF_B0059 is toxic to *E. coli* and that this toxicity is

abrogated upon mutation of conserved active site residues in either the N-terminal DUF262 or

C-terminal DUF1524 domains. These data suggest that EF_B0059 functions similarly to other

enzymes that harbor these domains such as the TIV-REs GmrSD and BrxU, which target and

degrade DNA methylated at cytosine bases. Interestingly, however, we also found that

EF_B0059 remains toxic to an *E. coli* strain that is unable to methylate its DNA. In line with this

finding, a sequence analysis of the phi47 genome failed to identify genes with homology to

known DNA modifying enzymes, suggesting phi47 does not modify its DNA. This notion is

further reinforced by the mutations identified in our isolation of escape phages that evade

EF_B0059 restriction. If phi47 encoded DNA modifying enzymes that sensitize it to restriction by

EF_B0059, we would expect to isolate phage mutants for any such genes. The lack of

identification of such genes in the phi47 genome in addition to the LC-MS analysis of phi47 DNA

showing no obvious modifications, suggests that phi47 does not directly modify its DNA.

Together, these observations indicate that EF_B0059 targets unmodified DNA, distinguishing it

from previously characterized TIV-REs. This raises the obvious question of how EF_B0059

differentiates between host and phage DNA. One possible explanation is that EF_B0059

functions more akin to classical restriction-modification systems in which functionally and

genetically paired modification systems modify the host genome at specific sites that block

cleavage by their cognate restriction endonuclease. This explanation seems unlikely given that

the 18 genes we identified as being unique to pTEF2 compared to EF_B0059-deficient pCF10,

are not predicted to encode proteins with the capacity to modify DNA. A more plausible

explanation is that EF_B0059 recognizes a feature of the phage genome that is unrelated to a

specific base modification, such as a unique sequence motif or higher-order DNA structure. For

example, It has been previously reported that Wadjet systems cleave closed-circular DNA[71].

Moreover, EF_B0059 activity could be stimulated by an unknown signal that is present during

phage infection, perhaps related to the entry of the phage genome into the cell or the early

stages of its replication. Future studies examining how phages lacking a TifA homolog are able

to escape EF_B0059-dependent restriction may answer some of these questions and shed light

on how this novel defense system discriminates between self and non-self DNA.

It is important to note that protein-based antirestriction mechanisms, including those that

target TIV-REs have been described previously. Indeed, the discovery of GmrSD in *E. coli* as a

modification-dependent restriction enzyme that prevents T4 phage replication also coincided

with the characterization of the T4 phage internal protein I (IPI) as an inhibitor of GmrSD. IPI is

packaged into the T4 capsid and upon infection is released into the host cell where it protects

the phage DNA from degradation by binding to and inhibiting the enzymatic activity of

GmsrSD[53,72]. Our data indicate that like IPI, TifA inhibits EF_B0059 through a direct binding

mechanism. However, IPI and TifA share neither sequence nor structural homology indicating

that these TIV-RE inhibitors evolved independently. Intriguingly, the independent emergence of

counter-defense proteins like IPI and TifA, which target the same family of defense systems,

might provide insight into why phage genomes are replete with hypothetical genes that lack

homology to other genes beyond those found in closely related phages. It is possible that many

of these encode species-specific counter defense proteins that have independently evolved to

counteract related bacterial defense mechanisms. Our discovery of TifA adds to a growing list of

recent phage-bacteria coevolution studies showing that many of these small cryptic phage

proteins function specifically to counter antiphage defense[73,74].

Our data show that the wild-type TifA can bind and disrupt the activity of GC1503 and

BrxU, whereas a single missense mutation provides it with the capacity to inhibit EF_B0059.

Given their divergent sequences, these findings raise the question as to how TifA is able to

interact with and inhibit such diverse TIV-REs. We suspect that an explanation may lie in unique

biochemical characteristics of TifA. Specifically, our structural modeling indicates that TifA has a

highly electronegative surface that likely facilitates its binding to the electropositive regions of

TIV-REs that are involved in DNA binding. Although the details underpinning DNA recognition in

TIV-REs are incomplete, recent biochemical and structural studies of BrxU demonstrated that its

DNase activity is linked to its homodimerization, which results in the formation of a large

electropositive cleft that the authors posit is well suited for DNA binding. In the case of TifA,

adaptive mutations that enhance its binding to a particular TIV-RE would be selected for as

improved inhibitory activity would result in successful phage infections. Such a mechanism of

inhibition would explain why we observed a weak interaction between EF_B0059 and wild-type

phi47 TifA that is enhanced by the L32F mutation identified in the phi47 escape mutants.

446   Competitive inhibition models have been established for several antirestriction proteins

targeting other types of restriction enzymes. For example, the T7 phage protein Ocr

competitively inhibits multiple diverse Type I restriction enzymes by mimicking the structure and

chemical properties of B-form DNA[63]. As we suspect is the case for TifA, selection for point

mutations in Ocr results in a more efficient inhibitor of specific type I restriction enzymes[62].

Although our data suggest that Ocr and TifA may be functionally analogous in their capacity to

inhibit multiple restriction enzymes, the predicted structure of TifA does not appear to explicitly

mimic the charge distribution of DNA in any obvious way and thus we refrain from classifying

TifA as a bona fide DNA mimic. Indeed, further biochemical and structural investigations are

required to understand how TifA inhibits TIV-REs. Nevertheless, our data suggest that like DNA-

mimic antirestriction proteins, TifA likely provides phi47 and related phages with a robust

antiphage defense strategy, allowing these phages to adapt to and inhibit diverse TIV-REs.

<u>METHODS</u>

Bacterial strains, bacteriophages, and growth conditions.

All bacterial and bacteriophage strains used in this study can be found in Table S2. *E.*

faecalis strains were grown with aeration in Brain Heart Infusion (BHI, Becton Dickinson) broth

or on BHI agar at 37°C. To maintain pLZ12A, pMSP3545, pGCP123 plasmids cultures were

supplemented with 15 µg/mL chloramphenicol, 20 µg/ml erythromycin, and 500 µg/mL

kanamycin, respectively. For induction of the CRISPRi plasmids, cells were grown with 50

ng/mL nisin (MilliporeSigma). Phage plaque assays were performed using Todd-Hewitt broth

(THB, Becton Dickinson) agar supplemented with 10 mM $MgSO_4$ as described elsewhere[29]. To

generate strains of *E. faecalis* carrying pTEF2 and its derivatives, we performed pheromone

responsive plasmid matings. V19 pTEF2 and derivatives were selected with 7.5 µg/mL

vancomycin and 200 µg/mL spectinomycin. JH2-2 pTEF2 and derivatives were selected with 50

µg/mL rifampicin and 200 µg/mL spectinomycin. The construction of EF_B0058 and EF_B0059

mutant strains of *E. faecalis* followed a previously published protocol for allelic exchange[75].

Primers used to generate deletion constructs that were cloned into pLT06 can be found in Table

S2. *Escherichia coli* and *Escherichia fergusonii* were grown in Lennox lysogeny broth (LB,

Fisher Scientific) with aeration or on LB agar at 37°C. When appropriate, cultures were

supplemented with 15 µg/mL chloramphenicol, 20 µg/ml erythromycin, 50 µg/mL kanamycin,

200 µg/mL trimethoprim, 15 µg/mL gentamicin, 0.5 mM IPTG, and 0.2% w/v L-rhamnose.

Clostridium sporogenes and *Leuconostoc mesenteroides* were grown on BHI broth and agar

supplemented with 0.5 g/L L-cysteine, 10 mg/L hemin, and 1 mg/L vitamin K at 37°C in an

anerobic chamber (5% CO_2, 5% H_2, 90% N_2; Shel Labs).

Plasmid construction

A detailed list of the primers and plasmids used in this study can be found in Table S2. For DNA

amplification by PCR*, E. faecalis, E fergusonii, L. mesenteroides, C. sporogenes,* and phi47

genomic DNA was purified using a ZymoBIOMICS DNA Miniprep Kit (Zymo Research) or a

previously published protocol for phenol/chloroform extraction[29]. PCR reactions used to create

DNA fragments for allelic exchange were performed using KOD Hot Start DNA Polymerase

(EMD Millipore) and cloned into pLT06[75]. phi47 *orf65* (*tifA*) and *orf65*-L32F were PCR amplified

using Q5 master mix (New England Biolabs) and cloned into pLZ12A, a derivative of the shuttle

vector pLZ12 carrying the *bacA* promoter (41). CRISPRi sgRNA pGCP123 plasmids were

designed and cloned following previously published protocols[50]. Plasmids were transformed into

E. faecalis by electroporation. Briefly, *E. faecalis* was grown in 100 ml of BHI broth to an OD_{600}

of 0.3. The cells were washed once with 10 ml of ice-cold 10% glycerol and resuspended in 500

µl of lysozyme solution (30 µg/ml lysozyme, 20% sucrose, 10 mM EDTA, 50 mM NaCl and 10

mM Tris-HCl, pH 8.0) and incubated at 37ºC for 20 min. The cells were washed four times with

500 µl of ice-cold electroporation buffer (500 mM sucrose and 10% glycerol). Cell pellets were

resuspended in 250 µl of ice-cold electroporation buffer and 0.5 – 1.0 µg of plasmid DNA was

electroporated into 50 µl of the *E. faecalis* suspension using a 0.2 mm cuvette at 2.5k V, 200 Ω,

and 25 µF. Plasmids used in *E. coli* include pSCrhaB2-CV, a low-copy (20-30 copies per cell),

tightly regulated expression vector containing a rhamnose-inducible promotor used for the

expression of EF_B0059 and other TIV-REs and pPSV39-CV, a high-copy expression vector

containing a *lac*UV5 promoter used for the expression of TifA[76-78]. All expression plasmids were

constructed using standard restriction enzyme-based cloning procedures. Reagents used for

cloning, including restriction enzymes, T4 DNA ligase, and Phusion DNA polymerase were

obtained from New England Biolabs. Plasmids were transformed into chemically competent *E.*

coli by heat-shock and transformants were subsequently isolated by selection with the

appropriate antibiotics.

Genomic comparisons of *E. faecalis* strains

The chromosomes of *E. faecalis* SF28073 (NZ_CP060804.1) and V583 (NC_004668.1) were

obtained from NCBI and aligned using the MAUVE multiple genome alignment tool[79]. To

compare plasmid sequences between *E. faecalis* strains, sequences of the plasmids pTEF1

(NC_004669.1), pTEF2 (NC_004670.1), pTEF3 (NC_004671.1), pSF1 (NZ_CP060801.1), pSF2

(NZ_CP060802.1), and pSF3 (NZ_CP060803.1) were obtained from NCBI and aligned using

the clinker alignment tool[80].

Pheromone responsive plasmid mating

For pheromone responsive plasmid conjugation, 150 µL of overnight *E. faecalis* cultures were subcultured into 5 mL of BHI broth and grown for 90 minutes with aeration at 37°C. Donor and recipient bacteria were mixed at a 1:1 ratio in a volume of 1 mL and applied to a 0.45-micron filter. The filter was removed from the housing and placed on a BHI agar plate with the bacteria side of the filter facing up. These plates were incubated overnight at 37°C. The following day, the filter was removed from the plate and placed in 5 mL of sterile PBS and vortexed to dislodge the bacteria from the filter. Serially diluted bacteria were plated on BHI agar containing antibiotics as described above to select for recipient cells carrying the plasmid and were grown at 37°C for no more than 48 hrs. Potential transconjugants were screened via colony PCR to confirm the presence of the plasmid. To generate *E. faecalis* V19 pTEF2 and JH2-2 pTEF2 the donor was *E. faecalis* OG1RF pTEF2-specR[40]. To generate *E. faecalis* V19 pTEF1, *E. faecalis* OG1SSp pTEF1 was used as the donor[40].

Growth curves

For broth culture infections of *E. faecalis* with phi47, 15 mL of THB broth was inoculated with *E. faecalis* and grown to an OD_{600} of 0.3. Cultures were infected with phi47 at a multiplicity of infection of either 1 or 10, as indicted, and supplemented with 10 mM $MgSO_4$. Cultures were incubated at room temperature for 5 minutes, then moved to 37°C with aeration. Phage production was quantified by removing 100 µL of the culture, adding 33.3 µL of chloroform, and quantifying the number of phages using plaque assays[38]. For the *E. coli* growth curves, overnight cultures containing the indicated plasmids were diluted 1:100 into LB broth supplemented with 200 µg/mL trimethoprim and 15 µg/mL gentamicin. For each culture, 200 µL was seeded into a 96-well plate in triplicate. Growth was measured at 15-minute intervals until

an OD_{600} of 0.4 was reached, upon which protein expression was induced by the addition of

either 0.2% w/v L-rhamnose or 0.2% w/v L-rhamnose and 0.5 mM IPTG. For the infection of *E.*

coli cells expressing BrxU and Orf65 with coliphage, overnight cultures were diluted 1:100 in LB

and grown to an OD_{600} of 0.5. Expression of Orf65 and/or BrxU was induced by the addition 0.5

mM IPTG and 0.2% w/v L-rhamnose, respectively. After 30 minutes, 500 µL of *E. coli* was

mixed with 5 mL of semi-solid LB (0.75% w/v agar) supplemented with 0.5 mM IPTG and/or

0.2% L-rhamnose and poured onto an 1.5% w/v LB agar plate. The agar overlay was allowed to

dry and 5 µL of serially diluted coliphages were spotted on top. After overnight incubation at 37

°C, the plaques were quantified and compared between inducing conditions.

Reverse transcription-PCR

cDNA was generated from 1 µg of RNA using qScript cDNA SuperMix (QuantaBio) and the

cDNA was purified using a QIAquick PCR Purification Kit (Qiagen). 2 µL of cDNA was used in a

PCR reaction with primers specific to EF_B0058 and EF_B0059 (Table S2). The resulting PCR

products were run on a 1% w/v agarose gel to visualize PCR bands.

Phage adsorption assay

E. faecalis cultures grown in BHI were pelleted at 3220 x *g* for 10 minutes and resuspended to

10^8 CFU/mL in SM-plus buffer (100 mM NaCl, 50 mM Tris-HCl, 8 mM $MgSO_4$, 5 mM $CaCl_2$

[pH 7.4]). Phages were added at an MOI of 0.1 and incubated at room-temperature without

agitation for 10 minutes. The bacteria-phage suspensions were centrifuged at 24,000 x *g* for 1

minute and the supernatant was collected to determine the number of unbound phages using a

plaque assay. SM-plus buffer supplemented with phage only was used as a negative control.

Western blotting

To analyze the expression of EF_B0059, its active site mutants, and Orf65, 5 mL cultures of *E.*

coli containing the indicated plasmids were grown to an OD_{600} of 0.5, upon which protein

expression was induced by the addition of 0.2% w/v L-rhamnose and 0.5 mM IPTG. The cells

were pelleted by centrifugation at 7,800 x *g* after 30 minutes of protein expression. Cell pellets

were resuspended in 50 μL of PBS and mixed 1:1 with Laemmli buffer and boiled at 95°C for 10

minutes. Samples were centrifuged at 16,000 x *g* and 5 μL was loaded onto a 10% SDS PAGE

gel. The gel was run for 15 minutes at 95 V, followed by 40 minutes at 195 V, and total protein

was wet transferred to a nitrocellulose membrane using a mini Trans-Blot electrophoretic

system (Bio-Rad). After the transfer, the blot was blocked in TBS-T supplemented with 5% w/v

non-fat milk, shaking gently at room temperature for 1 hour. The blot was washed three times

with TBS-T and primary antibody - either anti-VSV-G (MilliporeSigma, V4888) or anti-FLAG

(MilliporeSigma, F7425) - was added to the blot at a 1:5000 dilution followed by an overnight

incubation shaking gently at 4°C. The blot was washed three times with TBS-T, followed by a 1-

hour incubation with secondary antibody (anti-rabbit HRP conjugate). After three more washes

with TBS-T, the blot was developed by the addition of Clarity Max ECL substrate (Bio-Rad) and

chemiluminescence was visualized using a ChemiDoc Touch Imaging System (Bio-Rad).

CRISPRi screening

Using previously described CRISPRi constructs for *E. faecalis*[50], each gene unique to pTEF2

was knocked down. In brief, pMSP3545-dCas9$_{Str}$ was leveraged to express the dCas9 to inhibit

gene expression, while pGCP123 carried each of the guide RNAs (sgRNAs) to facilitate

targeted inhibition. sgRNA oligonucleotides were designed against pTEF2 open reading frames

and targeted the non-template strand. sgRNA oligonucleotides were ordered from Twist

Bioscience. All sgRNA oligonucleotides used can be found in Table S2. *E. faecalis* V19 pTEF2

with CRISPRi and sgRNA plasmids were cultured overnight in BHI broth containing 200 µg/mL

kanamycin, 50 µg/mL erythromycin, and 50 ng/mL nisin. Overnight cultures were diluted to an

OD_{600} of 0.1 in SM-plus buffer. 6 µL of 10-fold serial dilutions were spotted on THB agar plates

supplemented with 10mM $MgSO_4$, 200 µg/mL kanamycin, 50 µg/mL erythromycin, and 50

ng/mL nisin, with and without $1x10^7$ PFU/mL of phi47 embedded in the agar. Plates were

incubated at 37°C for 48 hours and bacteria were enumerated. Relative viability was calculated

by dividing the CFU of cells grown in the presence of phi47 by the CFU grown in the absence of

phi47.

LC-MS analysis of phi47 DNA

DNA isolated from phage phi47 was digested into nucleosides before being subjected to liquid

chromatography-mass spectrometry (LC-MS) analysis. Approximately 1-2 µg phi47 DNA was

treated with Nucleoside Digestion Mix (New England Biolabs) following the manufacturer's

protocol at 37°C for >2h. The resulting mixture of nucleosides were filtered through a hydrophilic

PTFE 0.2 µm centrifugal filter, and the filtrate was subjected to the reverse phase LC-MS for

nucleoside separation and MS detection. LC-MS was performed on an Agilent 1290 Infinity II

UHPLC-MS system equipped with a G7117 Diode Array Detector and a LC/MSD XT G6135

Single Quadrupole Mass Detector. Chromatography performed with the instrument was on a

Waters XSelect HSS T3 C18 column (4.6 × 150 mm, 3 µm particle size) operated at a flow rate

of 0.5 mL/min with a binary gradient mobile phase consisting of 10 mM ammonium acetate (pH

4.5) and methanol. The course of chromatography was monitored at 260 nm. MS was operated

in both positive (+ESI) and negative (-ESI) electrospray ionization modes. MS was performed

with a capillary voltage of 2500 V at both modes, a fragmentation voltage of 70 V, and a mass

range of m/z 100 to 1000. Agilent ChemStation software was used for the primary LC-MS data

processing.

Structural modeling and comparisons

Protein structures were modelled using AlphaFold2 within Colabfold (V.1.5.2) that is part of

Google Collaboratory. Predictions were run using the default parameters[60]. The resulting

structures were visualized, and electrostatic surface potential was mapped using ChimeraX[81].

Structural comparisons and RMSD calculations were performed using the DALI webserver[82].

Identification of bacteria encoding GC1503 and GC874

To identify homologs of EF_B0059, we searched a local database containing the Surette lab

whole genome collection of diverse human bacterial phyla[64]. For this, the amino acid sequence

of EF_B0059 was searched against this database using tblastn in BLAST v2.13.0+, with default

settings and the '-t 6' option for generating tabular results[83]. The taxonomy of the genomes was

identified using GTDB-Tk[2] v202.0, and the results were merged with the BLAST hits in R

version 4.0.3[84].

Isolation of *E. coli* bacteriophages from wastewater

Bacteriophages were isolated from wastewater as previously described with minor alterations[55].

In brief, 5 mL of wastewater was centrifuged at 9,000 x *g* and then filtered through a 0.22 µM

filter. The cleared wastewater was mixed with 5 mL of LB and inoculated with 500 µL of an

overnight *E. coli* culture. After a 24-hour incubation shaking at 37°C, 2 mL aliquots were

removed from cultures that showed evidence of phage propagation (complete cell lysis), and

centrifuged at 12,000 x g. The supernatant was filtered through a 0.22 µM filter to remove any

remaining bacterial cells. The filtered supernatant was serially diluted in SM-plus buffer and 10

µL was added to 500 µL of an overnight $E.\ coli$ culture. This phage-bacterial mixture was then

added to 5 mL of semi-solid LB (0.75% w/v agar), mixed gently, and overlayed onto an LB agar

plate. After overnight incubation, the resulting plaques were isolated. Agar plugs of individual

plaques were used to inoculate cultures of $E.\ coli$ grown to mid-log phase as determined by

OD_{600} measurements. Upon lysis of the $E.\ coli$ cultures, the resulting phages were purified as

described above. The above infection protocol was repeated until a uniform plaque morphology

was obtained. Using this approach, we isolated 2 phages from 8 unique wastewater samples for

a total of 16 isolated coliphages. Subsequent plaque assays found that the replication of six of

these phages were restricted by BrxU to varying degrees. From these six, we used the phage

that displayed the largest degree of replication inhibition, which we term JCW01, to assay if co-

expression of $Orf65/Orf65^{L32F}$ interfered with BrxU-dependent restriction.

Sequencing and genomic analysis of pTEF2 escape phi47 isolates

phi47 genomic DNA was sequenced using the Illumina NextSeq 2000 platform to 300 Mbp

depth at SeqCenter (Pittsburgh, PA). The resulting reads were mapped to the phi47 genome

(NCBI accession ON086985.1) and SNPs were identified using CLC Genomics Workbench v20

(Qiagen) with a minimum coverage of 15, a minimum frequency of 30%, and ploidy of 0.

Co-immunoprecipitation assays

To probe the interaction between TifA and a given TIV-RE, 50 mL cultures of $E.\ coli$ were grown

to an OD_{600} of 0.5 upon which expression of FLAG-tagged TIV-RE and VSV-G-tagged Orf65

were induced with the addition 0.2% w/v L-rhamnose and 0.5 mM IPTG, respectively. After 30

minutes the cells were pelleted and resuspended in 2 mL of lysis buffer (150 mM NaCl, 20 mM

Tris-HCl pH 7.5) and lysed by sonication. The cell lysates were cleared by centrifugation at

16,000 x g for 30 minutes at 4 °C. Cleared cell lysates were incubated with 50 µL of anti-FLAG

agarose beads (Genscript, L00425) for 2 hours at 4 °C using a tube rotator. The beads were

washed three times with 10 mL of lysis buffer and bound protein was eluted by boiling in 50 µL

of Laemmli buffer. 5 µL of the elution was subjected to SDS-PAGE and western blotting as

described above using anti-FLAG and anti-VSV-G as the primary antibodies as described

above.

DATA AVAILABILITY

DNA sequencing reads can be found at the European Nucleotide Archive under project

accession PRJEB68205.

FIGURE LEGENDS

Figure 1. **The pheromone responsive plasmid pTEF2 is sufficient to inhibit phi47**

replication. (A) Capacity of phage phi47 to form plaques measured by $\log_{10}$ PFU/mL on the

indicated *E. faecalis* strains. **(B)** PFU/mL produced upon phi47 infection of indicated *E. faecalis*

strains over 80 minutes. All strains were infected with an equivalent MOI of 1.0.

Figure 2. *EF_B0059* **encodes a putative type IV restriction enzyme that prevents infection**

by phi47. (A) Sequence alignments between the pheromone responsive plasmids pTEF2 and

pCF10. Genes highlighted in yellow indicate a sequence identify greater than 95%. **(B)** Capacity

of phage phi47 to form plaques measured by PFU/mL on the indicated *E. faecalis* strains. **(C)**

Relative viability of *E. faecalis* V19 pTEF2 cells upon transcription knock-down of the indicated

pTEF2 encoded genes of interested using CRISPR interference. Relative viability was

determined by comparing the growth of each strain in the presence and absence of phi47.

Statistical significance determined by ANOVA with multiple corrections, *p=0.0004. **(D)** and **(E)**

Outcome of phi47 infections measured by PFU/mL in *E. faecalis* strains V583 (D) and V19 (E) in

which the genes encoding EF_B0058 or EF_B0059 on pTEF2 are deleted. **(F)** Schematic of the

EF_B0058 and EF_B0059 genes. **(G)** Reverse transcription-PCR analysis of the EF_B0058 and

EF_B0059 mRNA transcript in *E. faecalis* V19 with or without pTEF2. Primer sets used to probe

the structure of the EF_B0058-EF_B0059 transcriptional operon are shown in red (EF_B0058

specific), green (intergenic) and blue (EF_B0059 specific). **(H)** Growth curves of *E. coli*

expressing EF_B0059, the indicated active site mutants, or empty vector. Arrow indicates when

protein expression was induced. **(I)** Western blot of VSV-G-tagged EF_B0059 in *E. coli* strains

from (H), 30 minutes post-induction. A non-specific band from the PAGE of these samples

stained with Coomassie Brilliant Blue is used as a loading control (load). Data are

representative of three independent experiments.

Figure 3. A mutation in phage phi47 protein Orf65 prevents EF_B0059-dependent

restriction of phi47. (A) Gene schematic of the genomic loci of *orf65* in the phi47 genome. **(B)**

Capacity of phi47 to form plaques measured by PFU/mL on *E. faecalis* strain V19 or V19

carrying pTEF2 and co-expressing Orf65 or Orf65^{L32F}. **(C)** Ribbon representation of the

predicted structure of Orf65 as generated using Alphafold2. The structure is colored according

to the predicted local distance difference test (pLDDT). The distribution of the acidic residues

throughout the structure is also shown. **(D)** The predicted amino acid surface electrostatic

potential of Orf65. **(E)** Growth curves of *E. coli* expressing EF_B0059 and either Orf65 or

Orf65^{L32F}, or empty vector. The arrow indicates when protein expression was induced. **(F)**

Western blot of VSV-G-tagged EF_B0059 and Orf65/Orf65^{L32F} in *E. coli* strains from (E) 30

minutes post-induction. A non-specific band from the PAGE of these samples stained with

Coomassie Brilliant Blue is used as a loading control (load). Data are representative of three

independent experiments. **(G)** Western blot analysis of the Co-immunoprecipitation of VSV-G-

tagged Orf65 or Orf65^{L32F} with FLAG-tagged EF_B0059. Tagged proteins were co-expressed in

E. coli for 30 minutes upon which cells were lysed and FLAG-EF_B0059 was precipitated using

anti-FLAG antibody conjugated to agarose beads. Data are representative of three independent

experiments.

Figure 4. Wild-type Orf65 inactivates distantly related type IV restriction enzymes. (A)

Multiple sequence alignment of residues 25-42 from Orf65 homologs identified in the genomes

of 9 closely related enterococcal phages. The residues are colored according to their percent

similarity at that site. **(B)** Growth curves of *E. coli* expressing GC1503 and either Orf65 or

Orf65^{L32F}, or empty vector. Arrow indicates when protein expression was induced. **(C)** Western

blot analysis of the Co-immunoprecipitation of VSV-G-tagged Orf65 or Orf65^{L32F} with FLAG-

tagged GC1503. Tagged proteins were co-expressed in *E. coli* for 30 minutes upon which cells

were lysed and FLAG-tagged EF_B0059 was precipitated using anti-FLAG antibody conjugated

to agarose beads. Data are representative of three independent experiments. **(D)** Outcome of

coliphage infection measured by PFU/mL in *E. coli* strain XL-1 blue co-expressing BrxU and

either Orf65 or Orf65$^{©F}$. **(E)** Outcome of Co-immunoprecipitation as in (C) but with FLAG-tagged

BrxU.

Table S1. SNPs in Phi47 isolates before and after selection by pTEF2. Three isolates of Phi47

were independently grown to high titer and subjected to selection by pTEF2. Phi47.1, Phi47.2,

and Phi47.3 represent the three original stocks of Phi47 prior to selection. pTEF2R 1 and

pTEF2R 2 denote each of the two pTEF2-resistant isolates sequenced for each original stock.

Orf65 (TifA) mutations are bolded.

Table S2. Bacterial/bacteriophage strains, plasmids, and oligonucleotides used in this

study.

Figure S1. Genomic comparisons of *E. faecalis* strains V583 and SF28073. (A) Whole

genome alignments of alignments of *E. faecalis* V583 and SF28073 chromosomes **(B)**

Sequence alignments between plasmids from *E. faecalis* V583 (pTEF1, pTEF2, pTEF3) and *E.*

faecalis SF28073 (pSF1, pSF2, pSF3).

Figure S2. pTEF2 prevents assembly of intracellular phage particles in an abortive

infection-independent manner. (A) Growth curves of the indicated *E. faecalis* strains infected

with Phi47 at an MOI of 10 The growth *E. faecalis* strain V19 containing pTEF2 was measured

with and without (uninfected) the addition of phi47. (**B**) Percent of phi47 adsorbed to *E. faecalis*

strains V583, V19, V19 pTEF2, and OG1RF (negative control).

Figure S3. The predicted structure of EF_B0059 resembles GmrSD-like type IV restriction

enzymes. (A) Ribbon representation of the predicted structure of EF_B0059 generated using

Alphafold2. The predicted local distance difference tests (pLDDT) per position is mapped onto

the model. Purple corresponds to a more confident prediction at that position. The N-terminal

DUF262 and C-terminal DUF1524 domains, and the residues that define each active site are

indicated. **(B)** The predicted alignment error (Å) of all residues against all residues for the model

in (A). Lower error (blue) corresponds to a well-defined relative position for a given residue pair.

Figure S4. EF_B0059 likely targets unmodified phage DNA. (A) Growth curves of *E. coli*

strains XL1-blue and ER2796 expressing EF_B0059 or empty vector (EV). Arrow indicates

when inducer was added to the cultures. Data represent means ± SD, n= 3. **(B)** Chromatograms

of digested phi47 DNA, deoxyribonucleotide (dNs) standards, or ribonucleotide (rNs) standards.

The *m/z*, measured, and expected masses for each peak in the phi47 trace is indicated below

(C). Capacity of phage phi47 to form plaques measured by PFU/mL on *E. faecalis* strain JH2-2

or JH2-2 + pTEF2 following passage through the indicated *E. faecalis* strains.

Figure S5. The L32F mutation in Orf65 is essential for overcoming restriction by

EF_B0059. (A) Outcome of either wildtype of evolved phi47 infections measured by PFU/mL of

the *E. faecalis* strain V19 with or without pTEF2. **(B)** The predicted alignment error (Å) of all

residues against all residues for the structural model in (Fig. 3C). Lower error (blue)

corresponds to a well-defined relative position for a given residue pair. **(C)** Anti-FLAG

immunoprecipitation of VSV-G tagged Orf65 or Orf65^{L32F} expressed in the absence of

EF_B0059-FLAG, detected by Western blot. **(D)** Growth curves of *E. coli* co-expressing

EF_B0059 and Orf65 or the indicated Orf65 mutant. **(E)** Anti-FLAG immunoprecipitation of VSV-

G tagged Orf65, or the indicated L32 mutant.

Figure S6. GC1503, but not GC874, is toxic to *E. coli.* (A) Growth curves of *E. coli*

expressing the indicated Orf65 variants. Orf65^{SDS2} refers to the Orf65 homolog from the

enterococcus phage SDS2. **(B)** Growth curves of *E. coli* expressing either EF_B0059, GC1503,

GC874, or an empty vector control. Arrow indicates when protein expression was induced. **(C)**

Western blot of VSV-G-tagged EF_B0059 and Orf65 or Orf65^{L32F} in *E. coli* strains 30 minutes

post-induction. A non-specific band from the PAGE of these samples stained with Coomassie

Brilliant Blue is used as a loading control (load). Data are representative of three independent

experiments. **(D)** Growth curves of *E. coli* co-expressing GC1503 and the indicated ORf65

mutant. Arrow indicates when protein expression was induced.

Figure S7. GC1503 and GC874 are predicted to be structurally similar to EF_B0059. (A)

Structural alignment of the predicted GC1503 (grey) and GC874 (red) models with that of

EF_B0059 (blue). Domains corresponding to the N-terminal DUF262 and C-terminal DUF1524

are highlighted, along with the sequences corresponding to the conserved catalytic resides for

each structure in these domains. The alignment statistics, including the sequence identify and

Cα root mean squared deviations (rmsd) are also indicated. **(B)** Structural model of GC874

(upper) coloured according to the per position predicted local distance difference test (pLDDT)

and the predicted alignment error (Å) plot. **(C)** Structural model of GC1503 (upper) coloured

according to the per position predicted local distance difference test (pLDDT) and the predicted

alignment error (Å) plot.

REFERENCES

798   1.   Donskey CJ, Chowdhry TK, Hecker MT, Hoyen CK, Hanrahan JA, Hujer AM, Hutton-
799   Thomas RA, Whalen CC, Bonomo RA, Rice LB. Effect of antibiotic therapy on the
800   density of vancomycin-resistant enterococci in the stool of colonized patients. *N Engl J*
801   *Med.* **343**, 1925-1932 (2000). doi:10.1056/NEJM200012283432604

2.   Ubeda C, Bucci V, Caballero S, Djukovic A, Toussaint NC, Equinda M, Lipuma L, Ling L, Gobourne A, No D, Taur Y, Jenq RR, van den Brink MRM, Xavier JB, Pamer EG. Intestinal microbiota containing *Barnesiella* species cures vancomycin-resistant *Enterococcus faecium* colonization. *Infect Immun*. **81**, 965-973 (2013). doi:10.1128/IAI.01197-12

3.   Suh GA, Lodise TP, Tamma PD, Knisely JM, Alexander J, Aslam S, Barton KD, Bizzell E, Totten KMC, Campbell JL, Chan BK, Cunningham SA, Goodman KE, Greenwood-Quaintance KE, Harris AD, Hesse S, Maresso A, Nussenblatt V, Pride D, Rybak MJ, Sund Z, van Duin D, Van Tyne D, Patel R. Considerations for the use of phage therapy in clinical practice. *Antimicrob Agents Chemother*. **66**, e0207121 (2022). doi:10.1128/aac.02071-21

4.   Strathdee SA, Hatfull GF, Mutalik VK, Schooley RT. Phage therapy: From biological mechanisms to future directions. *Cell*. **186**, 17-31 (2023). doi:10.1016/j.cell.2022.11.017

5.   Kortright KE, Chan BK, Koff JL, Turner PE. Phage therapy: A renewed approach to combat antibiotic-resistant bacteria. *Cell Host Microbe*. **25**, 219-232 (2019). doi:10.1016/j.chom.2019.01.014

6.   Sulakvelidze A, Alavidze Z, Morris JG. Bacteriophage therapy. *Antimicrob Agents Chemother*. **45**, 649-659 (2001). doi:10.1128/AAC.45.3.649-659.2001

7.   Langdon A, Crook N, Dantas G. The effects of antibiotics on the microbiome throughout development and alternative approaches for therapeutic modulation. *Genome Med*. **8**, 39 (2016). doi:10.1186/s13073-016-0294-z

8.   Moelling K, Broecker F, Willy C. A wake-up call: We need phage therapy now. *Viruses*. **10**, 688 (2018). doi:10.3390/v10120688

9.   Brives C, Pourraz J. Phage therapy as a potential solution in the fight against AMR: obstacles and possible futures. *Palgrave Commun*. **6**, 1-11 (2020). doi:10.1057/s41599-020-0478-4

10. Cook R, Brown N, Redgwell T, Rihtman B, Barnes M, Clokie M, Stekel DJ, Hobman J, Jones MA, Millard A. INfrastructure for a PHAge REference Database: Identification of

large-scale biases in the current collection of cultured phage genomes. *Phage (New*
Rochelle). 2, 214-223 (2021). doi:10.1089/phage.2021.0007

11. Ofir G, Sorek R. Contemporary phage biology: From classic models to new insights.
Cell. **172**, 1260-1270 (2018). doi:10.1016/j.cell.2017.10.045

12. Hampton HG, Watson BNJ, Fineran PC. The arms race between bacteria and their
phage foes. *Nature*. **577**, 327-336 (2020). doi:10.1038/s41586-019-1894-8

13. Duncan-Lowey B, Kranzusch PJ. CBASS phage defense and evolution of antiviral
nucleotide signaling. *Curr Opin Immunol*. **74**, 156-163 (2022).
doi:10.1016/j.coi.2022.01.002

14. Hochhauser D, Millman A, Sorek R. The defense island repertoire of the *Escherichia coli*
pan-genome. *PLoS Genet*. **19**, e1010694 (2023). doi:10.1371/journal.pgen.1010694

15. Makarova KS, Wolf YI, Snir S, Koonin EV. Defense islands in bacterial and archaeal
genomes and prediction of novel defense systems. *J Bacteriol*. **193**, 6039-6056 (2011).
doi:10.1128/jb.05535-11

16. Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, Amitai G, Sorek R.
Systematic discovery of antiphage defense systems in the microbial pangenome.
Science. **359**, (2018). doi:10.1126/science.aar4120

17. Brouns SJ, Jore MM, Lundgren M, Westra ER, Slijkhuis RJ, Snijders AP, Dickman MJ,
Makarova KS, Koonin EV, van der Oost J. Small CRISPR RNAs guide antiviral defense
in prokaryotes. *Science*. **321**, 960-964 (2008). doi:10.1126/science.1159689

18. Mojica FJ, Diez-Villasenor C, Garcia-Martinez J, Soria E. Intervening sequences of
regularly spaced prokaryotic repeats derive from foreign genetic elements. *J Mol Evol*.
60, 174-182 (2005). doi:10.1007/s00239-004-0046-3

19. Tock MR, Dryden DT. The biology of restriction and anti-restriction. *Curr Opin Microbiol*.
8, 466-472 (2005). doi:10.1016/j.mib.2005.06.003

20. Xiong X, Wu G, Wei Y, Liu L, Zhang Y, Su R, Jiang X, Li M, Gao H, Tian X, Zhang Y, Hu
L, Chen S, Tang Y, Jiang S, Huang R, Li Z, Wang Y, Deng Z, Wang J, Dedon PC, Chen
S, Wang L. SspABCD-SspE is a phosphorothioation-sensing bacterial defence system
with broad anti-phage activities. *Nat Microbiol.* **5**, 917-928 (2020). doi:10.1038/s41564-
020-0700-6

21. Goldfarb T, Sberro H, Weinstock E, Cohen O, Doron S, Charpak-Amikam Y, Afik S, Ofir
G, Sorek R. BREX is a novel phage resistance system widespread in microbial
genomes. *EMBO J.* **34**, 169-183 (2015). doi:10.15252/embj.201489455

22. Pecota DC, Wood TK. Exclusion of T4 phage by the *hok/sok* killer locus from plasmid
R1. *J Bacteriol.* **178**, 2044-2050 (1996). doi:10.1128/jb.178.7.2044-2050.1996

23. Fineran PC, Blower TR, Foulds IJ, Humphreys DP, Lilley KS, Salmond GP. The phage
abortive infection system, ToxIN, functions as a protein-RNA toxin-antitoxin pair. *Proc
Natl Acad Sci U S A.* **106**, 894-899 (2009). doi:10.1073/pnas.0808832106

24. Dy RL, Przybilski R, Semeijn K, Salmond GP, Fineran PC. A widespread bacteriophage
abortive infection system functions through a Type IV toxin-antitoxin mechanism. *Nucleic
Acids Res.* **42**, 4590-4605 (2014). doi:10.1093/nar/gkt1419

25. Cohen D, Melamed S, Millman A, Shulman G, Oppenheimer-Shaanan Y, Kacen A,
Doron S, Amitai G, Sorek R. Cyclic GMP-AMP signalling protects bacteria against viral
infection. *Nature.* **574**, 691-695 (2019). doi:10.1038/s41586-019-1605-5

26. Millman A, Bernheim A, Stokar-Avihail A, Fedorenko T, Voichek M, Leavitt A,
Oppenheimer-Shaanan Y, Sorek R. Bacterial retrons function in anti-phage defense.
Cell. **183**, 1551-1561 e1512 (2020). doi:10.1016/j.cell.2020.09.065

27. Kronheim S, Daniel-Ivad M, Duan Z, Hwang S, Wong AI, Mantel I, Nodwell JR, Maxwell
KL. A chemical defence against phage infection. *Nature.* **564**, 283-286 (2018).
doi:10.1038/s41586-018-0767-x

28. Burmeister AR, Fortier A, Roush C, Lessing AJ, Bender RG, Barahman R, Grant R,
Chan BK, Turner PE. Pleiotropy complicates a trade-off between phage resistance and
antibiotic resistance. *Proc Natl Acad Sci U S A.* **117**, 11207-11216 (2020).
doi:10.1073/pnas.1919888117

29. Duerkop BA, Huo W, Bhardwaj P, Palmer KL, Hooper LV. Molecular basis for lytic bacteriophage resistance in enterococci. *mBio*. **7**, e01304-01316 (2016). doi:10.1128/mBio.01304-16

30. Gordillo Altamirano F, Forsyth JH, Patwa R, Kostoulias X, Trim M, Subedi D, Archer SK, Morris FC, Oliveira C, Kielty L, Korneev D, O'Bryan MK, Lithgow TJ, Peleg AY, Barr JJ. Bacteriophage-resistant *Acinetobacter baumannii* are resensitized to antimicrobials. *Nat Microbiol*. **6**, 157-161 (2021). doi:10.1038/s41564-020-00830-7

31. Eugster MR, Morax LS, Huls VJ, Huwiler SG, Leclercq A, Lecuit M, Loessner MJ. Bacteriophage predation promotes serovar diversification in *Listeria monocytogenes*. *Mol Microbiol*. **97**, 33-46 (2015). doi:10.1111/mmi.13009

32. Canfield GS, Chatterjee A, Espinosa J, Mangalea MR, Sheriff EK, Keidan M, McBride SW, McCollister BD, Hang HC, Duerkop BA. Lytic bacteriophages facilitate antibiotic sensitization of *Enterococcus faecium*. *Antimicrob Agents Chemother*. **65**, e00143-00121, AAC.00143-00121 (2021). doi:10.1128/AAC.00143-21

33. Paul BG, Burstein D, Castelle CJ, Handa S, Arambula D, Czornyj E, Thomas BC, Ghosh P, Miller JF, Banfield JF, Valentine DL. Retroelement-guided protein diversification abounds in vast lineages of bacteria and archaea. *Nat Microbiol*. **2**, 17045 (2017). doi:10.1038/nmicrobiol.2017.45

34. Rusinov IS, Ershova AS, Karyagina AS, Spirin SA, Alexeevski AV. Avoidance of recognition sites of restriction-modification systems is a widespread but not universal anti-restriction strategy of prokaryotic viruses. *BMC Genomics*. **19**, 885 (2018). doi:10.1186/s12864-018-5324-3

35. Golovenko D, Manakova E, Tamulaitiene G, Grazulis S, Siksnys V. Structural mechanisms for the 5'-CCWGG sequence recognition by the N- and C-terminal domains of EcoRII. *Nucleic Acids Res*. **37**, 6613-6624 (2009). doi:10.1093/nar/gkp699

36. Blower TR, Evans TJ, Przybilski R, Fineran PC, Salmond GP. Viral evasion of a bacterial suicide system by RNA-based molecular mimicry enables infectious altruism. *PLoS Genet*. **8**, e1003023 (2012). doi:10.1371/journal.pgen.1003023

37. Samson JE, Belanger M, Moineau S. Effect of the abortive infection mechanism and
type III toxin/antitoxin system AbiQ on the lytic cycle of *Lactococcus lactis* phages. *J*
Bacteriol. **195**, 3947-3956 (2013). doi:10.1128/JB.00296-13

38. Chatterjee A, Johnson CN, Luong P, Hullahalli K, McBride SW, Schubert AM, Palmer
KL, Carlson PE, Duerkop BA. Bacteriophage resistance alters antibiotic-ediated
intestinal expansion of enterococci. *Infect Immun.* **87**, e00085-00019 (2019).
doi:10.1128/IAI.00085-19

39. Furlan S, Matos RC, Kennedy SP, Doublet B, Serror P, Rigottier-Gois L. Fitness
restoration of a genetically tractable *Enterococcus faecalis* V583 derivative to study
decoration-related phenotypes of the enterococcal polysaccharide antigen. *mSphere.* **4**,
(2019). doi:10.1128/mSphere.00310-19

40. Manson JM, Hancock LE, Gilmore MS. Mechanism of chromosomal transfer of
Enterococcus faecalis pathogenicity island, capsule, antimicrobial resistance, and other
traits. *Proc Natl Acad Sci U S A.* **107**, 12269-12274 (2010).
doi:10.1073/pnas.1000139107

41. Dunny GM. Enterococcal sex pheromones: signaling, social behavior, and evolution.
Annu Rev Genet. **47**, 457-482 (2013). doi:10.1146/annurev-genet-111212-133449

42. Paulsen IT, Banerjei L, Myers GSA, Nelson KE, Seshadri R, Read TD, Fouts DE, Eisen
JA, Gill SR, Heidelberg JF, Tettelin H, Dodson RJ, Umayam L, Brinkac L, Beanan M,
Daugherty S, DeBoy RT, Durkin S, Kolonay J, Madupu R, Nelson W, Vamathevan J,
Tran B, Upton J, Hansen T, Shetty J, Khouri H, Utterback T, Radune D, Ketchum KA,
Dougherty BA, Fraser CM. Role of mobile DNA in the evolution of vancomycin-resistant
Enterococcus faecalis. *Science.* **299**, 2071-2074 (2003). doi:10.1126/science.1080613

43. Jeltsch A, Pingoud A. Horizontal gene transfer contributes to the wide distribution and
evolution of type II restriction-modification systems. *J Mol Evol.* **42**, 91-96 (1996).
doi:10.1007/BF02198833

44. Harms A, Brodersen DE, Mitarai N, Gerdes K. Toxins, targets, and triggers: An overview
of toxin-antitoxin biology. *Mol Cell.* **70**, 768-784 (2018). doi:10.1016/j.molcel.2018.01.003

45. Rocha EPC, Bikard D. Microbial defenses against mobile genetic elements and viruses: Who defends whom from what? *PLoS Biol.* **20**, e3001514 (2022). doi:10.1371/journal.pbio.3001514

46. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* **215**, 403-410 (1990). doi:10.1016/S0022-2836(05)80360-2

47. Fernandez-Garcia L, Wood TK. Phage-defense systems are unlikely to cause cell suicide. *Viruses.* **15**, (2023). doi:10.3390/v15091795

48. Lopatina A, Tal N, Sorek R. Abortive infection: Bacterial suicide as an antiviral immune strategy. *Annu Rev Virol.* **7**, 371-384 (2020). doi:10.1146/annurev-virology-011620-040628

49. Aframian N, Eldar A. Abortive infection antiphage defense systems: separating mechanism and phenotype. *Trends Microbiol.* **31**, 1003-1012 (2023). doi:10.1016/j.tim.2023.05.002

50. Afonina I, Ong J, Chua J, Lu T, Kline KA. Multiplex CRISPRi system enables the study of stage-specific biofilm genetic requirements in *Enterococcus faecalis*. *mBio.* **11**, e01101-01120 (2020). doi:10.1128/mBio.01101-20

51. Machnicka MA, Kaminska KH, Dunin-Horkawicz S, Bujnicki JM. Phylogenomics and sequence-structure-function relationships in the GmrSD family of Type IV restriction enzymes. *BMC Bioinformatics.* **16**, 336 (2015). doi:10.1186/s12859-015-0773-z

52. Jablonska J, Matelska D, Steczkiewicz K, Ginalski K. Systematic classification of the His-Me finger superfamily. *Nucleic Acids Res.* **45**, 11479-11494 (2017). doi:10.1093/nar/gkx924

53. Bair CL, Black LW. A type IV modification dependent restriction nuclease that targets glucosylated hydroxymethyl ytosine modified DNAs. *J Mol Biol.* **366**, 768-778 (2007). doi:10.1016/j.jmb.2006.11.051

54. He X, Hull V, Thomas JA, Fu X, Gidwani S, Gupta YK, Black LW, Xu SY. Expression and purification of a single-chain Type IV restriction enzyme Eco94GmrSD and

1020 determination of its substrate preference. *Sci Rep.* **5**, 9747 (2015).
1021 doi:10.1038/srep09747

1022

1023 55. Picton DM, Luyten YA, Morgan RD, Nelson A, Smith DL, Dryden DTF, Hinton JCD,
1024 Blower TR. The phage defence island of a multidrug resistant plasmid uses both BREX
1025 and type IV restriction for complementary protection from viruses. *Nucleic Acids Res.* **49**,
1026 11257-11273 (2021). doi:10.1093/nar/gkab906

1027

1028 56. Stern A, Sorek R. The phage-host arms race: shaping the evolution of microbes.
1029 *Bioessays.* **33**, 43-51 (2011). doi:10.1002/bies.201000071

1030

1031 57. Rocha EP, Danchin A, Viari A. Evolutionary role of restriction/modification systems as
1032 revealed by comparative genome analysis. *Genome Res.* **11**, 946-958 (2001).
1033 doi:10.1101/gr.gr-1531rr

1034

58. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe
O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper J,
Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N, Kohli P, Kleywegt
G, Birney E, Hassabis D, Velankar S. AlphaFold protein structure database: Massively
expanding the structural coverage of protein-sequence space with high-accuracy
models. *Nucleic Acids Res.* **50**, D439-D444 (2021). doi:10.1093/nar/gkab1061

1041

59. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool
K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie
A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D,
Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver
D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. Highly accurate protein
structure prediction with AlphaFold. *Nature.* **596**, 583-589 (2021). doi:10.1038/s41586-
021-03819-2

1049

1050 60. Mirdita M, Schutze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold:
1051 making protein folding accessible to all. *Nat Methods.* **19**, 679-682 (2022).
1052 doi:10.1038/s41592-022-01488-1

1053

1054 61. Kamitori S, Yoshida H. Structure-function relationship of bacterial SH3 domains. In: *SH*
1055 *Domains* (ed Kurochkina N). Springer, Cham (2015). doi:10.1007/978-3-319-20098-9_4

1056

1057　62.　Roberts GA, Stephanou AS, Kanwar N, Dawson A, Cooper LP, Chen K, Nutley M,
1058　　　　Cooper A, Blakely GW, Dryden DTF. Exploring the DNA mimicry of the Ocr protein of
1059　　　　phage T7. *Nucleic Acids Res.* **40**, 8129-8143 (2012). doi:10.1093/nar/gks516

1060

1061　63.　Walkinshaw MD, Taylor P, Sturrock SS, Atanasiu C, Berge T, Henderson RM,
1062　　　　Edwardson JM, Dryden DTF. Structure of Ocr from bacteriophage T7, a protein that
1063　　　　mimics B-form DNA. *Mol Cell.* **9**, 187-194 (2002). doi:10.1016/s1097-2765(02)00435-5

1064

1065　64.　Derakhshani H, Bernier SP, Marko VA, Surette MG. Completion of draft bacterial
1066　　　　genomes by long-read sequencing of synthetic genomic pools. *BMC Genomics.* **21**, 519
1067　　　　(2020). doi:10.1186/s12864-020-06910-6

1068

1069　65.　Xu S-Y, Corvaglia AR, Chan S-H, Zheng Y, Linder P. A type IV modification-dependent
1070　　　　restriction enzyme SauUSI from *Staphylococcus aureus* subsp. aureus USA300. *Nucleic
1071　　　　Acids Res.* **39**, 5597-5610 (2011). doi:10.1093/nar/gkr098

1072

1073　66.　Ofir G, Melamed S, Sberro H, Mukamel Z, Silverman S, Yaakov G, Doron S, Sorek R.
1074　　　　DISARM is a widespread bacterial defence system with broad anti-phage activities. *Nat
1075　　　　Microbiol.* **3**, 90-98 (2018). doi:10.1038/s41564-017-0051-0

1076

1077　67.　Sterling AJ, Snelling WJ, Naughton PJ, Ternan NG, Dooley JSG. Competent but
1078　　　　complex communication: The phenomena of pheromone-responsive plasmids. *PLoS
1079　　　　Pathog.* **16**, e1008310 (2020). doi:10.1371/journal.ppat.1008310

1080

1081　68.　Licht TR, Laugesen D, Jensen LB, Jacobsen BL. Transfer of the pheromone-inducible
1082　　　　plasmid pCF10 among *Enterococcus faecalis* microorganisms colonizing the intestine of
1083　　　　mini-pigs. *Appl Environ Microbiol.* **68**, 187-193 (2002). doi:10.1128/AEM.68.1.187-
1084　　　　193.2002

1085

1086　69.　Huycke MM, Gilmore MS, Jett BD, Booth JL. Transfer of pheromone-inducible plasmids
1087　　　　between *Enterococcus faecalis* in the Syrian hamster gastrointestinal tract. *J Infect Dis.*
1088　　　　**166**, 1188-1191 (1992). doi:10.1093/infdis/166.5.1188

1089

1090　70.　Rice PA. Serine resolvases. *Microbiol Spectr.* **3**, MDNA3-0045-2014 (2015).
1091　　　　doi:10.1128/microbiolspec.MDNA3-0045-2014

1092

1093   71.   Deep A, Gu Y, Gao YQ, Ego KM, Herzik MA, Jr., Zhou H, Corbett KD. The SMC-family
1094   Wadjet complex protects bacteria from plasmid transformation by recognition and
1095   cleavage of closed-circular DNA. *Mol Cell.* **82**, 4145-4159 e4147 (2022).
1096   doi:10.1016/j.molcel.2022.09.008

1097

1098   72.   Jamet A, Touchon M, Ribeiro-Goncalves B, Carrico JA, Charbit A, Nassif X, Ramirez M,
1099   Rocha EPC. A widespread family of polymorphic toxins encoded by temperate phages.
1100   *BMC Biol.* **15**, 75 (2017). doi:10.1186/s12915-017-0415-1

1101

1102   73.   Millman A, Melamed S, Leavitt A, Doron S, Bernheim A, Hor J, Garb J, Bechon N,
1103   Brandis A, Lopatina A, Ofir G, Hochhauser D, Stokar-Avihail A, Tal N, Sharir S, Voichek
1104   M, Erez Z, Ferrer JLM, Dar D, Kacen A, Amitai G, Sorek R. An expanded arsenal of
1105   immune systems that protect bacteria from phages. *Cell Host Microbe.* **30**, 1556-1569
1106   e1555 (2022). doi:10.1016/j.chom.2022.09.017

1107

1108   74.   LeRoux M, Srikant S, Teodoro GIC, Zhang T, Littlehale ML, Doron S, Badiee M, Leung
1109   AKL, Sorek R, Laub MT. The DarTG toxin-antitoxin system provides phage defence by
1110   ADP-ribosylating viral DNA. *Nat Microbiol.* **7**, 1028-1040 (2022). doi:10.1038/s41564-
1111   022-01153-5

1112

1113   75.   Thurlow LR, Thomas VC, Hancock LE. Capsular polysaccharide production in
1114   *Enterococcus faecalis* and contribution of CpsF to capsule serospecificity. *J Bacteriol.*
1115   **191**, 6203-6210 (2009). doi:10.1128/jb.00592-09

1116

1117   76.   Lefebre MD, Valvano MA. Construction and evaluation of plasmid vectors optimized for
1118   constitutive and regulated gene expression in *Burkholderia cepacia* complex isolates.
1119   *Appl Environ Microbiol.* **68**, 5956-5964 (2002). doi:10.1128/AEM.68.12.5956-5964.2002

1120

1121   77.   Silverman JM, Agnello DM, Zheng H, Andrews BT, Li M, Catalano CE, Gonen T,
1122   Mougous JD. Haemolysin coregulated protein is an exported receptor and chaperone of
1123   type VI secretion substrates. *Mol Cell.* **51**, 584-593 (2013).
1124   doi:10.1016/j.molcel.2013.07.025

1125

1126   78.   Cardona ST, Valvano MA. An expression vector containing a rhamnose-inducible
1127   promoter provides tightly regulated gene expression in *Burkholderia cenocepacia.*
1128   *Plasmid.* **54**, 219-228 (2005). doi:10.1016/j.plasmid.2005.03.004

1129

1130 79. Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene
1131   gain, loss and rearrangement. *PLoS One*. **5**, e11147 (2010).
1132   doi:10.1371/journal.pone.0011147

1133

1134 80. Gilchrist CLM, Chooi Y-H. clinker & clustermap.js: automatic generation of gene cluster
1135   comparison figures. *Bioinformatics*. **37**, 2473-2475 (2021).
1136   doi:10.1093/bioinformatics/btab007

1137

1138 81. Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE.
1139   UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci*.
1140   **27**, 14-25 (2018). doi:10.1002/pro.3235

1141

1142 82. Holm L. Using Dali for protein structure comparison. *Methods Mol Biol*. **2112**, 29-42
1143   (2020). doi:10.1007/978-1-0716-0270-6_3

1144

1145 83. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
1146   BLAST+: architecture and applications. *BMC Bioinformatics*. **10**, 421 (2009).
1147   doi:10.1186/1471-2105-10-421

1148

1149 84. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk v2: memory friendly
1150   classification with the genome taxonomy database. *Bioinformatics*. **38**, 5315-5316
1151   (2022). doi:10.1093/bioinformatics/btac672

1152

ACKNOWLEDGEMENTS

We would like to thank Dr. Michael Gilmore for providing *E. faecalis* OG1RF pTEF2-specR and

E. faecalis OG1SSp pTEF1, Dr. Katrine Whiteson for sharing phage SDS2, Dr. Kimberly Kline

for providing CRISPRi genetic tools, Dr. James Samuelson and Dr. Elisabeth Raleigh for

supplying the *E. coli* strain ER2796, and Dr. Michael Surette for sharing *Clostridium sporogenes*

and *Leuconostoc mesenteroides* strains. We also thank Dr. Lindsey Marmont for her

constructive feedback on the manuscript.

1160

AUTHOR CONTRIBUTIONS

B.D., J.W., C. J., N.B., and S. A. conceptualized the project. C.J. and N.B. performed the

majority of experiments. S. A., G. A., S. M., and Y-J. L. assisted with experiments. P. W.

provided valuable insight for data analysis. C. J. and N. B. wrote the manuscript. B. D., J. W.,

and S. A. edited the manuscript. B. D., J. W., C. J., S. A., and G. A. acquired funding that

supported the study.

1167

1168 **COMPETING INTERESTS**

1169 The authors declare no competing interests.

1170

FUNDING

This work was support by National Institutes of Health grants R01AI141479 (B.A.D.),

T32AI052066 (C.N.J.), and F31AI157050 (C.N.J.), the National Science Foundation Graduate

Research Fellowship Program (S.E.A.), American Heart Association Postdoctoral Fellowship

24POST1183502 (G. A.), the Burroughs Welcome Fund Investigators in the Pathogenesis of

Infectious Disease Program (J.C.W.), and a Canada Research Chair in Molecular Microbiology

(J.C.W.).

A phi47 *orf64* *orf65* *orf66* *orf67*

G → A
L32F

200 bp

B

V19 + pTEF2

$\log_{10}$ pfu mL^{-1}

V19   −   + Orf65   + Orf65^{L32F}

ND   ND

C

F32

SH3-domain

N

C

Alphafold2 confidence
low — high

D

Charge
negative — positive

E

- Vector Control
- + EV
- + Orf65
- + Orf65^{L32F}

EF_B0059

OD$_{600}$

Time (hours)

F

EF_B0059

Control   + EV   + Orf65   + Orf65^{L32F}

75
50

EF_B0059

Orf65

Load

pScrhaB2::EF_B0059-VSV-G
pPSV39::Orf65-VSV-G

G

pPSV39::Orf65-VSV-G

Orf65^{L32F} Orf65

Input
Orf65
EF_B0059

Elute
Orf65
EF_B0059

pScrhaB2::EF_B0059-FLAG