The T Cell Receptor β Chain Repertoire of Tumor Infiltrating Lymphocytes Improves

Neoantigen Prediction and Prioritization

Thi Mong Quynh Pham[1], Thanh Nhan Nguyen[1], Bui Que Tran Nguyen[1], Thi Phuong Diem

Tran[1], Nguyen My Diem Pham[1], Hoang Thien Phuc Nguyen[1], Thi Kim Cuong Ho[1], Dinh Viet

Linh Nguyen[1], Huu Thinh Nguyen[2], Duc Huy Tran[2], Thanh Sang Tran[2], Truong-Vinh Ngoc

Pham[2], Minh-Triet Le[2], Thi Tuong Vy Nguyen[1], Minh-Duy Phan[1], Hoa Giang[1], Hoai-Nghia

Nguyen[1], Le Son Tran[1]

1. Medical Genetics Institute, Ho Chi Minh City, Vietnam

2. University Medical Center Ho Chi Minh City, Ho Chi Minh City, Vietnam

* Correspondence:

Le Son Tran (leson1808@gmail.com), Hoai-Nghia Nguyen (nhnghia81@gmail.com)

Keywords: neoantigen, colorectal cancer (CRC), TCR sequencing (TCRseq), tumor variant

calling, neoantigen identification workflow, neoantigen prioritization, cancer immunotherapy.

ABSTRACT

In the realm of cancer immunotherapy, the meticulous selection of neoantigens plays a

fundamental role in enhancing personalized treatments. Traditionally, this selection process has

heavily relied on predicting the binding of peptides to human leukocyte antigens (pHLA).

Nevertheless, this approach often overlooks the dynamic interaction between tumor cells and the

immune system. In response to this limitation, we have developed an innovative prediction

algorithm rooted in machine learning, integrating T cell receptor β chain (TCRβ) profiling data

from colorectal cancer (CRC) patients for a more precise neoantigen prioritization. TCRβ

sequencing was conducted to profile the TCR repertoire of tumor-infiltrating lymphocytes (TILs)

from 28 CRC patients. The data unveiled both intra-tumor and inter-patient heterogeneity in the

TCRβ repertoires of CRC patients, likely resulting from the stochastic utilization of V and J

segments in response to neoantigens. Our novel combined model integrates pHLA binding

information with pHLA-TCR binding to prioritize neoantigens, resulting in heightened

specificity and sensitivity compared to models using individual features alone. The efficacy of

our proposed model was corroborated through ELISpot assays on long peptides, performed on
four CRC patients. These assays demonstrated that neoantigen candidates prioritized by our
combined model outperformed predictions made by the established tool NetMHCpan. This
comprehensive assessment underscores the significance of integrating pHLA binding with
pHLA-TCR binding analysis for more effective immunotherapeutic strategies.

INTRODUCTION

In metastatic CRC patients, immunotherapy has fulfilled promise in improving survival rate
(1). Immune checkpoint inhibitors (ICIs), which block negative regulatory pathways in T-cell
activation, have been approved by the US Food and Drug Administration (FDA) for the
treatment of deficient mismatch repair (dMMR) or high microsatellite instability (MSI-H) CRC
patients (2-4). However, there is an urgent need for alternative immunotherapy strategies for
metastatic CRC patients, as patients with proficient mismatch repair (pMMR) or microsatellite
stability (MSS) have not shown significant responses to immune checkpoint inhibitors (2, 5).

Neoantigens (neopeptides) have emerged as potential targets for personalized cancer
immunotherapy, including CRC (6, 7). Neoantigens are peptides that result from somatic
mutations and can be displayed by class I human leukocyte antigen (HLA-I) molecules on the
surface of cancer cells, thereby activating immune-mediated tumor killing (8). Recent studies
have demonstrated that the presence of neoantigens is associated with better responses to
immune checkpoint inhibitor (ICI) therapy in CRC patients (9, 10). A high neoantigen burden
has been linked to improved overall survival and progression-free survival in patients with
various solid tumors, including CRC (9, 10). Therefore, neoantigen-based immunotherapies are
considered to have significant potential for improving treatment outcomes for CRC patients.

The successful development of neoantigen-based therapies hinges upon the identification of
neoantigens that exhibit a strong binding affinity to their respective HLA-I molecules and
demonstrate high immunogenicity. Initially, DNA sequencing of tumor tissues and paired
Peripheral Blood Mononuclear Cells identifies cancer-associated genomic mutations. RNA
sequencing then determines the patient's HLA-I allele profile and the gene expression levels of
mutated genes. *In silico* tools are then applied to analyze the tumor somatic variant, HLA-I allele,
and gene expression data, predicting the binding affinity of neoantigens to the patient's HLA-I
alleles and their potential to activate T cell responses (11-13). This standard workflow has been
successful in identifying clinically relevant neoantigens in various malignancies (12, 14).

However, despite its achievements, the current approach's impact on patient outcomes

remains limited due to the scarcity of mutations in cancer patients that lead to effective

immunogenic neoantigens. This limitation arises primarily from the fact that the selection of

neoantigen candidates by these workflows relies solely on neoantigen-HLA-I binding affinity as

the criteria for immunogenicity prediction (15). While HLA-I binding is indeed a crucial factor

for neoantigen presentation, it does not fully account for T cell receptor (TCR) recognition and

interaction. The recognition of the neoantigen-HLA complex through TCR is of paramount

importance for T cell activation and eliciting an immune response (16, 17). Various factors, such

as the specific TCR repertoire, TCR clonality, and the structural characteristics of the TCR-

peptide-HLA complex, profoundly influence TCR recognition. Unfortunately, these critical

aspects are not entirely captured by HLA-I binding prediction alone.

The TCR is a critical component of the adaptive immune system, responsible for recognizing

specific antigens presented by antigen-presenting cells (APCs). This membrane-bound

heterodimer protein complex is expressed on the surface of T cells and exists in two distinct

forms: TCRα/TCRβ for αβ T cells and TCRγ/TCRδ for γδ T cells, both intricately associated

with invariant CD3 chain molecules (18, 19). TCR diversity arises from the recombination of

variable (V), diversity (D), and joining (J) genes at the TCRβ and TCRδ loci, along with VJ

recombination at the TCRα and TCRγ loci (20), giving rise to a broad array of unique TCRs

collectively known as the TCR repertoire. The complementarity-determining region 3 (CDR3),

situated at the junction of V, D, and J gene segments (21), plays a pivotal role in antigen

recognition, with the unique combination of CDR3 sequences contributing significantly to the

specificity and diversity of the TCR repertoire. With advancements in technology, TCR

sequencing has become a powerful technique used to characterize the diversity and composition

of TCR repertoires. Valuable information directly obtained through TCR sequencing, including

TCR clonotype diversity, V(D)J gene usage, repertoire size, clonal expansion, and repertoire

changes, has provided invaluable insights into immune responses, antigen recognition, and the

development of targeted immunotherapies.

Sequencing the T cell receptors (TCRs) of TILs or lymphocytes found in peripheral blood

provides crucial insights into the T-cell repertoire and their responses against neoantigens

associated with tumors (22-24). This information holds paramount importance in identifying

TCRs that specifically target these neoantigens. Additionally, it proves to be invaluable in

assessing the immunogenic potential of predicted neoantigens. Unfortunately, the current
bioinformatic workflows used in neoantigen selection and prioritization do not incorporate TCR
sequencing data. Therefore, we speculate that integrating TCR sequencing data into the
assessment of immunogenicity of predicted neoantigens holds promise for unveiling effective
neoantigens which can induce immune response, consequently, advancing the development of
personalized immunotherapies for cancer.

To achieve this goal, we first performed TCR sequencing on frozen samples collected from
CRC patients to profile TCRβ repertoire of tumor filtrating T cells. We next exploited the
peptide-HLA (pHLA) and pHLA-TCR binding affinity of peptides with known immunogenicity
status from six public databases including 10X Genomics (25-28), McPAS (29), PRIME (30),
VDJdb (31), IEDB (32) and TBAdb (33) to develop a predictive algorithm. Subsequently, we
employed this algorithm to predict and rank neoantigens by using the TCRβ, HLA types and
mutation profiles identified from patients' frozen tumor tissues. Finally, we experimentally
verified the efficacy of our model by measuring the immunogenicity of the ranked peptides by an
ELISpot assay.

MATERIALS AND METHODS

Tumor biopsy and peripheral blood collection

Twenty-eight patients diagnosed with CRC were enrolled in this study at the University
Medical Center in Ho Chi Minh City between June 2022 and May 2023. CRC confirmation was
based on abnormal colonoscopies and histopathological analysis. The stages of CRC were
determined according to guidelines provided by the American Joint Committee on Cancer and
the International Union for Cancer Control (the eighth version) (34, 35). Prior to participation, all
patients provided written informed consent for tumor and whole blood sample collection.
Relevant clinical data, including demographics, cancer stages, and pathology information, were
extracted from the medical records of the University Medical Center. Detailed information
regarding the clinical factors of the patients can be found in **Supplementary Table 1**. This study
was approved by the Ethics Committee of the University of Medicine and Pharmacy in Ho Chi
Minh City, Vietnam. For four patients, ten mL of peripheral blood was collected before surgery
and stored in Heparin tubes prior isolation of PBMCs.

Targeted DNA, RNA and TCRβ sequencing

The DNA/RNA samples were isolated using either the AllPrep DNA/RNA Mini Kit or the AllPrep DNA/RNA/miRNA Universal Kit (Qiagen, Germany) as per the manufacturer's protocol. In addition, matched genomic DNA from the white blood cells (WBC) of individuals was also extracted from the buffy coat using the GeneJET Whole Blood Genomic DNA Purification Mini kit (ThermoFisher, MA, USA), following the manufacturer's instructions. Genomic DNA samples from the patients's paired tumor tissues and WBCs were used to prepare DNA libraries for DNA sequencing with the ThruPLEX Tag-seq Kit (Takara Bio, USA). The libraries were then pooled and hybridized with pre-designed probes for 95 targeted genes (Integrated DNA Technologies, USA). This gene panel encompasses commonly mutated genes in CRC tumors, as reported in the Catalog of Somatic Mutations in Cancer (COSMIC) database. The DNA libraries were then subjected to massive parallel sequencing on the DNBSEQ G400 sequencer (MGI, Shenzhen, China) for paired-end reads of 2x100 bp with a sequencing depth of 10X.

Isolated total RNA was processed using the NEBNext Poly(A) mRNA Magnetic Isolation Module (New England Biolabs, MA, USA) to isolate intact poly(A)+ RNA following the manufacturer's instructions. RNA libraries were prepared using the NEBNext Ultra Directional RNA Library Prep Kit for Illumina (New England Biolabs). These libraries were subsequently sequenced for paired-end reads of 2x100 bp on an MGI system at a sequencing depth of 50X.

For TCRβ library construction, mRNA was utilized with the SMARTer Human TCR a/b Profiling Kit v2 (Takara, USA). The resulting libraries were sequenced for paired-end reads of 2x250 bp on an Illumina system at a sequencing depth of 7.5X.

Microsatellite instability status

Microsatellite instability (MSI) status was determined in both tumor DNA and corresponding DNA from blood, serving as the control, using the MSI Analysis System, version 1.2 (Promega, US). This analysis identified alterations in 5 mononucleotide repeat markers: BAT-25, BAT-26, NR-21, NR-24, and MONO-27. The results were classified as microsatellite stability (MSS) in cases where either none or only one marker was unstable. Samples showing alterations in two or more markers were categorized as microsatellite instability-high (MSI-H).

Variant calling from DNAseq and RNAseq data

The approach of integrating RNA-seq into calling variants was performed as described in our previous publication (36). Briefly, we used Dragen (Illumina) (37) in tumor-normal mode to detect somatic mutations from DNAseq data. The default filtering thresholds of Dragen were applied for the detection of single nucleotide polymorphisms (SNPs) and insertions/deletions (indels). SNPs were further filtered using the dbSNP and 1000 Genome datasets. Germline mutations in tumor tissues were identified by comparison with matched WBC DNA samples. Mutations within immunoglobulin and HLA genes were excluded due to alignment difficulties in these highly polymorphic regions, necessitating specialized analysis tools. Additionally, synonymous mutations were removed from downstream analysis. For analysis, we included somatic mutations that exceeded a minimum threshold of $\geq 2\%$ variant allele frequency (VAF) in DNA extracted from fresh frozen tissues.

Sequencing reads underwent trimming using Trimmomatic (38) and were then aligned to the human reference genome using STAR (version 2.6.0c) (39). Prior to alignment, raw sequencing reads were subjected to quality checks using FastQC version 0.11.9 (40). VarScan 2 (41), , which accepts both DNA and RNAseq data, was employed to identify mutations in paired tumor and WBC samples across 95 cancer-associated genes. This analysis was carried out in tumor-normal mode. Four filtering steps were applied: (i) only calls with a PASS status were utilized, (ii) population SNPs overlapping with a panel of normal samples from the 1000 Genome dataset were excluded, (iii) somatic mutations included for analysis met a minimum threshold of $\geq 10\times$ read depth and $\geq 2\%$ variant allele frequency (VAF) in RNA extracted from FF tissue, and (iv) synonymous mutations and those related to HLA were removed from downstream analysis.

The resulting BAM files were sorted, indexed using Samtools version 1.10 (42), and had PCR duplicates removed using Picard tools version 2.25.6 (43). Somatic variants were manually reviewed using Integrative Genomics Viewer (v2.8.2). The VCF files generated by Dragen (for DNAseq) and VarScan 2 (for RNAseq) were subsequently annotated using the Ensembl Variant Effect Predictor (VEP version 105) (44) to extract information about the potential effects of variants on the phenotypic outcome.

CDR3β calling from TCRseq data

To define the CDR3β from TCRseq, we utilized Cogent NGS Immune Profiler Software v1.0
(45). Before calling clonotypes, the raw sequencing reads underwent filtering based on the
following criteria: (i) allowing only one mismatch while splitting reads by matching read
sequences to different receptor chains, (ii) excluding reads shorter than 30 bp and reads
ambiguously matched to multiple receptor chains, (iii) excluding reads that failed correction
when linker-based correction was enabled, (iv) excluding reads that failed the abundance check
during sequencing error correction, and (v) excluding molecular identifier groups (MIGs) with
fewer than 3 unique molecular identifier (UMI) reads. The filtered MIGs were then assembled
and aligned to the V(D)J reference to define the TCR clonotype.

Shannon index and clonality

We utilized two indices to characterize T-cell diversity and expansion: the Shannon entropy
index and the clonality index.
Shannon entropy:

$$-\sum_{i=1}^{n} p_i \log_e (p_i)$$

193   Clonality:

$$1 - \frac{\sum_{i=1}^{n} p_i \log_e (p_i)}{\log_e (n)}$$

These indices consider both the number of T-cell clone types 'n' and the frequency 'p_i' of each
clone. Here, 'p_i' represents the proportion of the i-th clone within the TCR library containing n
clones.

***In silico* prediction of peptide-HLA binding and peptide-HLA-TCR binding**

Class I HLA alleles (HLA-A/B/C) with two-digit resolution were identified from patient
tumor RNAseq data using the OptiType tool (**Supplementary Table 2)** (46, 47). The annotated
VCF files were analyzed using pVACseq, a tool in pVACtools (v1.5.9) (11, 48, 49). We used the
default settings, except for disabling the coverage and MAF filters. We used all peptide-HLA-I
binding algorithms that were implemented in pVACseq to predict 8 to 11-mer epitopes binding
to HLA-I (A, B, or C) for downstream analysis. Mutants with lower binding affinity than

wildtype peptides were prioritized using a two-step ranking process. The minimum binding
affinity score was calculated from the distribution of scores among the peptides derived from
each mutation, and priority was given to mutations with the lowest binding affinity scores.
Moreover, in pVACseq, we collected peptides with binding affinity scores lower than wildtype
from multiple tools, calculated minimum binding affinity scores for each unique peptide, and
incorporated this data into a combined machine learning model.

We used the peptide-HLA-TCR binding algorithm implemented in pMTNet (50) to predict
peptide binding to HLA and TCR with default settings. These scores represented the predicted
likelihood of peptides being immunogenic. A ranking value for immunogenicity was assigned to
each unique peptide by determining the minimum TCR ranking of its immunogenicity score.

Construction of a combined machine learning model

To identify immunogenic peptides, we conducted a thorough search across multiple
databases, including 10X Genomics (25-28), McPAS (29), VDJdb (31), IEDB (32), and TBAdb
(33). For the creation of non-immunogenic pHLA-TCR complex, we assembled a negative
dataset using the PRIME tool (30). This dataset was generated by associating each pHLA with
10 randomly generated TCRs sourced from the following databases: 10X Genomics, McPAS,
VDJdb, IEDB, and TBAdb. To objectively train and evaluate the model, we separated the dataset
mentioned above into two data subsets: discovery dataset (70%) and validation dataset (30%).
These subsets are mutually exclusive and do not overlap (**Supplementary Figure 1**). pHLA-
TCR complex in the discovery dataset, whether labeled as immunogenic or non-immunogenic,
were used for model training to classify whether a peptide is positive or not. We examined three
machine learning algorithms - Logistic Regression (LR), Random Forest (RF), and Extreme
Gradient Boosting (XGB) - for each feature type (pHLA binding, pHLA-TCR binding), as well
as for combined features. Feature selection was tested using a k-fold cross-validation approach
on the discovery dataset with 'k' set to 10-fold. This process splits the discovery dataset into 10
equal-sized folds, iteratively using 9 folds for training and 1 fold for validation. Model
performance was evaluated using the 'roc_auc' (Receiver Operating Characteristic Area Under
the Curve) metric, which measures the model's ability to distinguish between positive and
negative peptides. The average of these scores provides a robust estimate of the model's
performance and generalizability. The model with the highest 'roc_auc' average score, XGB,

was chosen for all features. The model cut-off was set based on a threshold specificity of >90%

or >95% to achieve high specificity for selecting peptides. This combined model's performance

was evaluated on the independent validation dataset of immunogenic and non-immunogenic

pHLA-TCR complex to assess its ability to classify positive peptides from negative peptides. In

our GitHub repository, we have included links to the GitHub repositories for NetMHCpan and

pMTNet (https://github.com/QuynhPham1220/Combined-model).

Ranking coverage score calculation

The approach of compare ranking between two algorithms was performed as described in a

previous publication (51). Briefly, the rank coverage score was based on the ranking value

calculation given by the formula:

$$\text{Rank Coverage Score} = \frac{\sum_{n \in negative} rank\,(n)}{T \times num\,(n)} \times coverage(n) - \frac{\sum_{n \in positive} rank\,(p)}{T \times num\,(p)} \times coverage(p)$$

$$coverage\,(k) = \frac{max\,rank(k)}{T} \quad k \in (n, p)$$

Where, *T* presented the total neoepitope number identified and *p* and *n* presented the positive and

negative peptides, respectively, that were experimentally validated in vitro.

If positive peptides have smaller rank values than negative peptides, this will result in a high

rank coverage score, indicates the better ranking result.

Isolation, culture, and stimulation of PBMCs with long peptides

Peripheral blood samples were collected from eight patients prior to surgery using BD

Vacutainer Heparin Tubes (BD Biosciences, NJ, USA). Peripheral blood mononuclear cells

(PBMCs) were isolated through gradient centrifugation using Lymphoprep (STEMCELL

Technologies) within 4 hours of blood collection. The PBMCs were then resuspended in a

solution of FBS (10%) and DMSO (7-10x10^6 cells/mL) for cryopreservation in liquid nitrogen.

Frozen PBMCs were thawed in AIM-V media (Gibco, Thermo Scientific, MA, USA)

supplemented with 10% FBS (Cytiva, USA) and DNase I (Stemcell Technology, Canada) (1

mg/mL) solution. A total of 10^5 PBMCs were allowed to rest in a 96-round bottom well-plate

containing AIM V media supplemented with 10% FBS, 10 mM HEPES, and 50 mM β-

mercaptoethanol overnight before stimulation with synthesized long peptides at a concentration

of 5 mM in a humidified incubator at 37°C with 5% CO2. PBMCs were further stimulated with

GM-CSF (2000 IU/mL, Gibco, MT, USA) and IL-4 (1000 IU/mL, Invitrogen, MA, USA) for 24

hours. Following this initial stimulation, LPS (100 ng/mL, SigmaAldrich, MA, USA) and IFN-γ

(10 ng/mL, Gibco, MT, USA) were added to the PBMCs along with the peptides for an

additional 12 hours. On the following day, IL-7, IL-15, and IL-21 (each at a concentration of 10

ng/mL) (Peprotech, NJ, USA) were added to the PBMC culture. The restimulation process

involved exposing the peptides to fresh media containing IL-7, IL-15, and IL-21 every 3 days,

for a total of 3 times. On day 12, PBMCs were restimulated with peptides and cultured in media

without cytokines. ELISpot assays were performed on stimulated PBMCs on day 13.

ELISpot assay on PBMCs stimulated with long peptides

Cultured T cells were transferred to an enzyme-linked immunospot (ELISpot) plate (Mabtech,

Sweden) and incubated for 20 hours at 37°C. Peripheral blood mononuclear cells (PBMCs)

cultured with DMSO were used as a negative control group, while PBMCs stimulated with anti-

CD3 were used as a positive control group. The ELISpot assay was performed on treated PBMCs

using the ELISpot Pro: Human IFN-g (ALP) kit (Mabtech, Sweden), following the

manufacturer's protocol. Developed spots on the ELISpot plate were then counted using an

ELISpot reader (Mabtech, Sweden). Reactivity was determined by measuring the fold increase in

the number of spots in PBMCs treated with mutant peptides compared to those treated with wild-

type peptides. A fold change of two was selected as the cutoff for positivity, indicating a

significant increase in reactivity (52).

Statistical analysis

In this study, t-tests were used to compare the TCR clones, clonality, and Shannon-index

among two groups of four categories (Microsatellite status, stage, gender, tumor location). Chi-

square test was used to compare the proportions of immununogenic and non-immunogenic

peptides. All statistical analyses were performed using R (version 4.3.0) with common data

analysis packages, including ggplot2 and pROC. The 95% confidence interval (95% CI) was

presented in brackets next to a value as appropriate.

RESULTS

The workflow of identifying neoantigens by combining both HLA and TCR binding characteristics

The identification of neoantigens has traditionally heavily relied on peptide-HLA (pHLA) binding prediction while often neglecting the significance of pHLA-TCR interactions (50). In this study, we introduce a novel workflow that integrates both pHLA and pHLA-TCR interactions to enhance the precision of neoantigen identification and prioritization (**Figure 1**). In the first step (**Figure 1A**), we conducted RNA and DNA sequencing on matched tumor tissues and peripheral blood mononuclear cells (PBMCs) collected from 28 CRC patients to detect cancer-associated nonsynonymous mutations and determine HLA types, as detailed in our previous work (36). Additionally, T-cell receptor sequencing (TCR-Seq) was performed to profile the TCRβ repertoire of TILs. In the second step (**Figure 1B**), we used the pVACseq and pMTNet tools to predict the binding affinities of both pHLA and pHLA-TCR interactions. To exploit the information of both pHLA and pHLA-TCR binding for selecting and ranking neoantigen candidates, we constructed a machine learning model by using immunogenic and non-immunogenic peptide information sourced from five publicly available databases (**Figure 1C**). Finally, we designed long peptide (LP) candidates encompassing the selected neoantigens and experimentally assessed their immunogenicity using peripheral blood mononuclear cells (PBMCs) obtained from the same patients (**Figure 1D**).

Heterogenous tumor infiltrating TCRβ profiles in colorectal cancer patients

Autologous TILs have exhibited varying reactivity levels to neoantigens, suggesting the potential of TIL-based recognition to improve the identification of immunogenic neoantigens (53). Furthermore, characterizing T cell receptors (TCRs) can complement efforts to predict immunogenicity. To explore this, we initiated our study by characterizing the TIL repertoire in a cohort of 28 colorectal cancer patients. This characterization involved sequencing the complementarity-determining region 3 (CDR3) of T-cell receptor beta (TCRβ), renowned for its remarkable diversity within the TCR gene. Our sequencing analysis yielded an average of 2,992,949 productive TCR reads per sample, with a range between 256,035 and 10,888,726 (**Supplementary Table 3**), following correction for duplications, sequencing errors, and exclusive use of uniquely barcoded reads mapped to TCRβ-CD3 sequences from the

ImMunoGeneTics (IMGT) databases. Across the 28 patients, we observed variable numbers of TCRβ-CDR3 clonotypes, ranging from 433 to 27,749 (**Figure 2A & Supplementary Table 3,** 28 patients were arranged in ascending order), indicating the intra-tumor heterogeneity of TCR clonotypes. Of these clonotypes, 59.5% exhibited a single uniquely barcoded read mapped to the TCRβ-CDR3 sequences (depicted in yellow in **Figure 2A**), while 40.5% had at least two TCRβ-CDR3 reads confidently identified (**Figure 2A**). As observed previously (54), the length distribution of CDR3 was approximately normally distributed (median 14, range 4-43, **Supplementary Figure 2A & Supplementary Table 4**). Subsequently, we explored potential associations between TCR diversity, as quantified by the Shannon index, and patient characteristics. In accordance with prior investigations, we identified an inverse relationship between the number of TCR clones and the Shannon index (**Figure 2B**). Based on our current sequencing depth, we have observed that many of our samples (14 out of 28) have reached sufficient saturation (**Figure 2C**) as their diversity of clonotypes was saturated. Additionally, the TCR clones selected in our studies are unique molecular identifier (UMI)-collapsed reads, with each representing at least three raw reads sharing the same UMI. Consequently, despite some samples having low sequencing depth for TCRβ sequencing, likely due to the diversity of TIL infiltration between patients, our TCRβ profiling analysis is robust and reliable. However, we did not detect any significant correlations between TCR clonality or diversity and clinical variables, including microsatellite status, tumor staging, patient gender and tumor location (**Supplementary Figure 4**). In contrast, MSI-H tumors, which are known to be rich in neoantigens (55, 56), displayed a significantly lower number of TCRβ clonotypes compared to MSS tumors (2181 versus 5330, p=0.057, **Supplementary Figure 3 & 4A**), aligning with previous research (57).

After filtering out TCR clones with read counts below 1, we obtained a total of 74,590 TCR clones. Subsequently, we conducted a comprehensive assessment of TCRβ repertoire similarity by calculating the proportion of overlapping TCRβ-CDR3 clones across the 28 patients. The preeminent majority, constituting 95.1% of all identified TCRβ-CDR3 clones, comprised unique clonotypes found in one patient, while the remaining 4.9% were recurrently observed in at least two patients (**Supplementary Figure 2B**). This observation underscores the substantial inter-patient heterogeneity in TCR profiles. The TCRβ repertoire is generated by the random rearrangement of variable (V), diversity (D) and joining (J) segments. We conducted a search

specifically targeting the D segment by scoring the similarities of sub-sequences within the
junction sequences against the reference sequences of D segments. Consistent with a previous
study, we were unable to unambiguously assign the D segments to any defined D region usage
due to the truncation of this region (58). In contrast to the diversified D segments, V and J
sequments displayed high recurrent rates across 28 patients (**Supplementary Figure 2B**). As
anticipated, we identified 59 distinct V segments (**Supplementary Figure 2C**) and 13 distinct J
segments (**Supplementary Figure 2D**), collectively sharing 185,627 clones across the 28 tumor
tissue samples. This underscores the conservation of these segments (**Supplementary Figure 2C**
& D). Conversely, we observed a varied combination of V and J segments, which significantly
contributes to the heterogeneity of the TIL TCRβ repertoire (**Supplementary Figure 2E**).
Collectively, our data elucidate the presence of both intra-tumor and inter-patient heterogeneity
in the TCRβ repertoires of CRC patients, likely stemming from the stochastic utilization of V
and J segments in response to neoantigens.

pHLA and pHLA-TCR interactions are two complement determinants of neoantigen
immunogenicity

While *in silico* tools predicting HLA-peptide binding affinity have traditionally played a
pivotal role in determining neoantigen immunogenicity (59), the evaluation of TCR-peptide
binding for screening immunogenic neoantigens remains understudied. In light of this, our study
aimed to assess the contributions of both pHLA and pHLA-TCR binding affinity in predicting
immunogenic neoantigens. To accomplish this, we gathered HLA and TCRβ sequences from
established datasets containing immunogenic and non-immunogenic pHLA-TCR complexes
(**Supplementary Table 5**). Subsequently, we employed NetMHCpan (13) and pMTNet (50)
tools to predict pHLA and pHLA-TCR binding, respectively.

For comparative purposes, we generated plots depicting predicted percentile rank values,
with lower percentile ranks signifying stronger binding affinity. As anticipated, our analysis
revealed a significantly higher prevalence of peptides with robust HLA binding (percentile rank
< 2%) among immunogenic peptides in contrast to their non-immunogenic counterparts (**Figure**
3A & B, p< 0.00001). Similarly, immunogenic peptides exhibited a greater proportion of
peptides with percentile ranks indicating pHLA-TCR binding of < 2% compared to non-
immunogenic peptides (**Figure 3C & D**, p=0.086). As recommended by NetMHCpan and
pMTNet, we considered peptide candidates with predicted percentile ranks below 2% as binders.

Utilizing this predefined threshold, both pHLA and pHLA-TCR binding affinity exhibited

comparable positive predictive values (PPV), at 68.5% (**Figure 3B)** and 64.3% (**Figure 3D)**,

respectively. This substantiates the significance of both pHLA and pHLA-TCR binding in

determining the immunogenicity of peptides. When we simultaneously applied the cutoff values

for both HLA-peptide and TCR-peptide percentile ranks (Q1 group, **Figure 3E**), the PPV

increased to 76.9% from 68.5% and 64.3% for individual binding features (**Figure 3F**). This

underscores the rationale for combining these two criteria to enhance the accuracy of neoantigen

prediction. Notably, pHLA-TCR binding displayed a remarkably lower sensitivity but higher

specificity in comparison to pHLA binding (**Figure 3G**), implying their potential as a

complementary criteria for the selection of immunogenic peptides.

Combination of peptide-HLA and peptide-HLA-TCR interactions improves neoantigen
prediction

The combination of peptide-HLA and peptide-HLA-TCR binding ranking with a fixed cutoff

values of 2% for each feature resulted in high specificity but low sensitivity (**Figure 3G**). To

optimize the precision of immugenic neoantigens, we examined three distinct machine learning

classifiers, namely Random Forest (RF), Logistic Regression (LR), and Extreme Gradient

Boosting (XGB) classifiers. We first partitioned the pHLA binding and pHLA-TCR binding

ranks for immunogenic and non-immunogenic peptides from publicly available databases into

separate discovery and validation datasets. These datasets were used to develop and validate

machine learning algorithms, as illustrated in **Figure 4A**. We next assessed the performance of

the three examined algorithms, employing a 10-fold cross-validation strategy. Among these

algorithms, XGB demonstrated the highest performance, achieving an area under the receiver

operating characteristic curve (AUC) of 0.82 in the training dataset and 0.84 in the validation

dataset (**Supplementary Figure 5**). Consequently, the model combining pHLA and pHLA-TCR

binding, referred to as the 'combined model' was chosen for further validation.

Our findings revealed that the combined model outperformed methods relying on pHLA-

TCR or pHLA binding feature separately. The combined model yielded sensitivity AUC values

of 0.82 (95% CI 0.81-0.84) and 0.84 (95% CI 0.82-0.86) for discovery and validation cohorts,

whereas the pHLA-TCR feature alone achieved AUC values of 0.69 (95% CI 0.66-0.71) and

0.74 (95% CI 0.71-0.77). Meanwhile, the pHLA feature resulted in AUC values of 0.76 (95% CI

0.75-0.78) and 0.74 (95% CI 0.72-0.77) (**Figure 4B**). In order to address the elevated false

positive rates associated with current prediction tools for neoantigens, we set the specificity at high thresholds of >95% and >99%. We observed that the combined model achieved a greater sensitivity (39.7% versus 5.9% and 19.7% at > 95% specificity; 47.1% versus 3.5% and 18.8% at > 99% specificity, **Figure 4C**), Negative Predictive Value (NPV, 87.7% versus 82% and 84.2% at > 95% specificity; 88.9% versus 81.5% and 83.9% at > 99% specificity, **Figure 4D**), and Positive Predictive Value (PPV, 65.9% versus 22.2% and 47.1% at > 95% specificity; 62.3% versus 11.8% and 39.4% at > 99% specificity, **Figure 4E**) compared to the single feature methods.

The accurate prioritization of neoantigen candidates with high immunogenicity holds the potential to streamline the validation process, reducing both costs and time expenditures. Consequently, we proceeded to evaluate the combined model's capacity to prioritize neoantigens by computing the ranking coverage score, which considers the accuracy in ranking immunogenic peptides versus non-immunogenic peptides (51). In the validation phase, the combined model exhibited superior rank coverage scores in comparison to the individual feature-based methods. The combined model attained a ranking coverage score of 0.37, while the single-feature methods, pHLA-TCR and pHLA, yielded scores of -0.26 and 0.25, respectively (**Figure 4F**). These findings underscore the notion that the incorporation of pHLA and pHLA-TCR binding criteria can enhance the accuracy of prediction and prioritization of immunogenic neoantigens.

Experimental validation of the pHLA-TCR and pHLA combined approach in selecting neoantigen candidates in patients with CRC

To experimentally validate the efficacy of our combined approach, we conducted a comparative analysis of the percentage of confirmed immunogenic neoantigen candidates and the ranking coverage scores between the conventional NetMHCpan method, which relies solely on pHLA binding percentiles, and our combined approach (**Figure 5A**). Due to the limited availability of patient blood samples, we were only able to perform validation on eight patients who possessed sufficient quantity of PBMCs. For each patient, we chose the top three neoantigen candidates predicted by each method. Among the neoantigen candidates, three were found to be common between the two methods, while twenty neoantigen candidates were uniquely identified by either NetMHCpan or the combined model (**Figure 5B**), bringing the total number of candidates to 44. Subsequently, we synthesized 44 long peptides (LPs) covering these 44 selected neoantigen candidates, along with 44 LPs corresponding to the wild-type sequences.

The LPs were utilized in an *ex vivo* ELISpot assay to measure the release of interferon-gamma

(IFNγ) from the four patients' PBMCs. If the stimulation with a mutant LP resulted in a > 2-fold

increase in IFNγ spots compared to its corresponding wildtype LP, the respective neoantigen

candidate was classified as an immunogenic neoantigen.

Out of the 44 selected LPs, we confirmed the immunogenicity of three LPs

(RNF213_W719S, MAP3K1_L1202F and TRRAP_T148TEL) identified by the NetMHCpan

method and seven LPs (NCOR2_R1963H, TRRAP_F1568S, DICER1_P1592L,

KMT2C_K3848T, BRAF_EV275-276-, SMAD4_G230R and PTPN13_D184A) selected by the

combined method (**Figure 5C & 5D**). Notably, six patients exhibited at least one immunogenic

peptide identified by the combined method, whereas none of the LPs in five patients, PT07,

PT18, PT21, PT24 and PT25, chosen via the NetMHCpan method were validated as

immunogenic (**Figure 5C & 5D**). To further assess the ranking accuracy of these two methods,

we calculated the ranking coverage scores. In agreement with our *in silico* analyses, we observed

higher rank coverage scores for the combined method in five out of the eight patients, resulting

in an overall score of 0.04 (**Figure 5E & Supplementary Figure 6**). In contrast, the

NetMHCpan method exhibited a lower rank coverage score of -0.37 (**Figure 5E &**

Supplementary Figure 6). To further evaluate our model, we gathered additional public data

and assessed its effectiveness in comparison to other models. We utilized immunogenic peptides

from databases such as NEPdb (60), NeoPeptide (61), dbPepneo (62), Tantigen (63), and

TSNAdb (64), ensuring there was no overlap with the datasets used for training and validation.

For non-immunogenic peptides, we used data from 10X Genomics Chromium Single Cell

Immune Profiling (25-28). The findings indicate that the combined model from pMTNet and

NetMHCpan outperforms NetTCR tool (65) (**Supplementary Table 6**). These outcomes

conclusively demonstrate the ability of the combined approach to enhance the prediction and

ranking of immunogenic neoantigens in cancer patients.

DISCUSSION

The selection of neoantigens plays a pivotal role in enhancing the efficacy of personalized

treatments in cancer immunotherapy. Historically, neoantigen selection has predominantly

hinged upon the prediction of peptide-human leukocyte antigen (pHLA) binding. However, the

limitations of this approach have become increasingly evident (66), as it often neglects the

dynamic interplay between tumor cells and the immune system. The findings of our study shed

new light on this critical aspect of neoantigen selection. Our research highlights the potential of integrating pHLA binding prediction with the assessment of pHLA-Tumor-Infiltrating Lymphocyte T Cell Receptor (TIL TCR) binding (**Figure 1**). By encompassing the interaction between neoantigens and the tumor microenvironment, we have demonstrated a substantial enhancement in the accuracy of neoantigen selection and prioritization. Our findings underscore the significance of comprehensive neoantigen assessment in harnessing the full potential of immunotherapeutic strategies in the fight against cancer.

Our investigation into the TIL TCRβ repertoire across 28 CRC patients has unveiled a complex tapestry of intra-tumor and inter-patient heterogeneity (**Figure 2A, Supplementary Figure 2B**). One underlying cause of this diversity is the random rearrangement of variable (V) and joining (J) segments, which contributes to the distinct TCRβ sequences observed (**Supplementary Figure 2C, D &E**). This remarkable diversity in the TCRβ profile, as elucidated in our study, presents a potential link to the heterogeneity of cancer mutations and neoantigens within CRC. The intricate relationship between the TCRβ repertoire and the genetic alterations within the tumor microenvironment underscores the need for a more comprehensive understanding of this dynamic interplay to develop more precise immunotherapeutic strategies (67). Furthermore, our research revealed a compelling observation regarding patients with MSI-H, who are known to exhibit high mutation and neoantigen burdens (68, 69). Intriguingly, MSI-H patients with strong immune reactivity to neoantigens was shown to display a lower number of TCR clonotypes and reduced diversity, as characterized by the Shannon index, compared to CRC patients with Microsatellite Stable (MSS) (**Supplementary Figure 4A**). This stark contrast in TCRβ diversity hints at the potential enrichment of neoantigen-reactive TCR clonotypes in MSI-H patients. These findings provide valuable insights into the immune responses to different subtypes of CRC and may guide the development of more tailored immunotherapies for patients with differing mutational landscapes.

By utilizing proven immunogenic and non-immunogenic peptides from public databases, we have demonstrated that predicting the strength of pHLA-TCR binding can be a crucial criterion for selecting immunogenic candidates (**Figure 3**). For the prediction of pHLA-TCR binding, we employed the well-established tool pMTnet (50). This choice was driven by its proven efficacy, based on its consistently high performance across different validated datasets (50). Although it exhibited lower sensitivity and positive predictive value (PPV) when compared to pHLA binding

alone, pHLA-TCR binding strength exhibited notably superior specificity (**Figure 3G**). This

implies that incorporating pHLA-TCR binding strength as a selection criterion would result in a

reduced false positive rate, a crucial factor in the context of neoantigen selection. What makes

our study particularly compelling is the convergence of these observations. The integration of

both criteria, specifically, pHLA binding and pHLA-TCR binding strength, emerged as a

strategy that not only enhanced the PPV but also fortified the rationale for combining these two

features in the neoantigen selection process.

The integration of both pHLA binding and pHLA-TCR binding strength features in our

approach exhibited superior performance in neoantigen selection and prioritization when

compared to the single-feature method (**Figure 4**). This observation aligns with the findings of

previous studies, which consistently indicate that combining multiple criteria enhances the

accuracy and efficacy of neoantigen identification (51, 70). Furthermore, our experimental

validation confirmed the robustness of our approach by consistently demonstrating performance

consistent with our analysis on publicly available data (**Figure 5 & Supplementary Table 6**).

When examining the percentile ranks of positive LPs predicted by netMHCpan from those

predicted by our combined model, the results further underscored the strength of our approach

(**Figure 5E**). This alignment between experimental validation and computational analysis

enhances the reliability and applicability of our neoantigen selection method. Interestingly, the

four identified neoantigen candidates with confirmed immunogenicity by the combined approach

have not been previously reported in public neoantigen databases, indicating that they could

serve as novel targets for neoantigen based therapies.

However, several limitations must be acknowledged. Firstly, the relatively small sample size

employed for validation raises the potential for ranking score bias. Additionally, while both

TCRα and TCRβ regions play essential roles in engaging peptide-bound HLA complexes, our

study focused solely on TCRβ sequences to predict pHLA-TCR binding strength. Future

investigations should include TCRα sequences to provide a more comprehensive analysis.

Although the PBMC and TILs were previously shown to be congruent in neoantigen reactivity

(71), it is important to recognize that differences in the contribution of TIL and PBMC TCR

repertoires in neoantigen selection may introduce variability in the selection and validation of

neoantigen candidates, and future studies are warranted to address the consistency in neoantigen

reactivity between these two sources of T cells. Moreover, to improve the accuracy and

effectiveness of the machine learning model in predicting and ranking neoantigens, we have developed an in-house tool called EpiTCR. This tool will utilize immunogenic assays, such as ELISpot and single-cell sequencing, for validation.

In summary, our study delves into the diversity and variation within the TCRβ repertoire of TILs in the tumor tissues of colorectal cancer patients. Through our research, we have introduced a novel approach for the identification and prioritization of neoantigen candidates. This approach combines the assessment of peptide-human leukocyte antigen (pHLA) binding and peptide-human leukocyte antigen-T cell receptor (pHLA-TCR) binding. Our findings underscore the significance of considering pHLA-TCR binding interactions as part of the process for selecting neoantigen candidates. This is a crucial step in the development of personalized immunotherapy strategies. By combining these two factors, we can more accurately identify the neoantigens that hold promise for effective immunotherapy, ultimately improving the prospects for tailored and effective cancer treatment.

Author contributions

Pham Thi Mong Quynh, Nguyen Thanh Nhan, Nguyen Bui Que Tran, Nguyen Hoang Thien Phuc, Tran Thi Phuong Diem, Pham Nguyen My Diem, Ho Thi Kim Cuong, and Nguyen Dinh Viet Linh were engaged in formal analysis, data curation, and methodology development for the manuscript draft. Huu Thinh Nguyen provided conceptualization and patient consultation. Duc Huy Tran, Tran Thanh Sang, Truong-Vinh Ngoc Pham, and Minh-Triet Le were responsible for patient recruitment and tissue histological analysis. Nguyen Thi Tuong Vy contributed to formal analysis and data curation. Minh-Duy Phan supported in conceptualization and manuscript editing. Hoa Giang contributed to conceptualization and methodology. Hoai Nghia Nguyen provided supervision and conceptualization guidance. Le Son Tran contributed to conceptualization and manuscript writing.

Acknowledgments

The authors thank all participants who agreed to take part in this study. We thank Dr. Kien Nguyen for proofreading our manuscript.

Financial & competing interests' disclosure

This research was funded by a NexCalibur Therapeutic grant (NC01). The authors including Le Son Tran and Minh Duy Phan hold the equity in NexCalibur Therapeutic.

FIGURE LEGEND

Figure 1. A novel workflow based on machine learning that integrates TCRβ sequencing data for the identification and ranking of CRC neoantigens.

(A) Tumor biopsies and peripheral blood from CRC patients were subjected to targeted DNA-seq, RNA-seq, and TCR-seq. (B) The prediction of peptide-HLA binding and peptide-HLA-TCR binding by indicated tools using the DNA-seq, RNA-seq, and TCR-seq data was performed. (C) Machine learning models were subsequently constructed based on the analysis of the peptide-HLA binding and peptide-HLA-TCR binding features to distinguish immunogenic antigens from non-immunogenic peptides. The immunogenicity of predicted neoantigen candidates prioritized by the model was validated by ELISpot to evaluate the effectiveness of this approach.

Figure 2. Tumor-infiltrating TCRβ profiles in 28 colorectal cancer patients.

(A) A bar plot depicting the distribution of TCR clonotypes among 28 CRC patients, categorized into two groups: those with a unique read count and those with read counts greater than or equal to 2 for each TCR clonotype. (B) The scatter plot illustrates the relationship between Shannon-index and the number of TCR clones. (C) The rarefaction plot shows the variable between sample size and diversity among 28 CRC samples.

Figure 3. Peptide-TCR and peptide-HLA interactions are two complementary determinants of neoantigen immunogenicity.

(A) The histogram displays the HLA percentile distribution of immunogenic antigens (red bar) and non-immunogenic peptides (grey bar). (B) The percentage of immunogenic antigens (red bar) and non-immunogenic peptides (grey bar) is compared between two groups based on HLA percentile: <2% and >= 2% (Chi-square test, p< 0.00001). (C) The histogram displays the TCR ranking distribution of immunogenic antigens (red bar) and non-immunogenic peptides (grey bar). (D) The percentage of immunogenic antigens (red bar) and non-immunogenic peptides (grey bar) is compared between two groups based on TCR ranking: <2% and >= 2% (Chi-square

test, p=0.086). (E) The scatter plot illustrates the relationship between the HLA percentile

distribution and TCR ranking of immunogenic antigens (red bar) and non-immunogenic peptides

(grey bar). (F) The percentage of immunogenic antigens (red bar) and non-immunogenic

peptides (grey bar) is analyzed in four distinct groups based on cutoffs of HLA percentile and

TCR ranking. (G) The bar plot illustrates the sensitivity and specificity of three neoantigen

prioritization approaches: based on neoantigen-HLA binding affinity alone (yellow bar),

neoantigen-TCR binding ranking alone (blue bar), and the combined method using both features

(red bar).

Figure 4. The combined model demonstrates improved sensitivity and specificity for

neoantigen prioritization.

(A) The workflow for constructing the model. (B) The ROC curves demonstrate the performance

of both the combined model and individual models in both the discovery and validation cohorts.

The bar graphs illustrate the sensitivity (C), negative predictive value (NPV) (D), and positive

predictive value (PPV) (E) at specificity levels of at least 95% or 99% for both the combined and

individual models in both the discovery and validation cohorts. (F) Ranking coverage scores for

the specified models in either the discovery or validation cohorts.

Figure 5. Validation of neoantigens identified *in silico* from the novel workflow through

ELISpot assays conducted on four CRC patients.

(A) A schematic diagram illustrates the procedural steps of neoantigen prioritization and the

ELISpot assay. (B) The count of neoantigens identified from each pipeline. (C) The fold change

in IFN-γ spots, relative to the wildtype peptides, is shown for 21 long peptides. Note: Only the

mutants that result in a positive value in ELISpot are depicted, along with their corresponding

amino acid changes and their associated rankings. (D) ELISpot assays on six long peptides

resulting in at least a 2-fold change in IFN-γ spots. (E) The bar graphs display ranking of

validated long peptides identified from the NetMHCpan tool (blue bar) or the combined method

(red bar) for individual patients and all patients.

Supplementary Figure 1. Dataset construction workflow.

Supplementary Figure 2. Quality control metrics for Tumor-Infiltrating Lymphocyte (TIL) TCRβ analysis.

(A) Distribution of CDR3β lengths in total TCR clones. (B) The pie chart displays the recurrence rates of TCR clones, V segments, and J segments when the read count of TCR clones exceeds 01. The graph illustrates the uniqueness of TCR clones and the shared presence of both V and J segments. (C) The heatmap depicts the Z-scored read counts of V segments or (D) J segments across 28 samples. Some V and J segments were found to be dominant in all samples. (E) The chord diagram illustrates the rearrangement of V and J segments, revealing random V and J combinations, with a few combinations exhibiting high frequencies.

Supplementary Figure 3. Rarefaction between MSI and MSS samples

The rarefaction plot illustrates the sample size and diversity of samples in two groups: MSI and MSS.

Supplementary Figure 4. Association between TIL TCRβ profiles and patients' characteristics.

The bar plot and dot plot compare TCR clones, Shannon-index, and clonality between MSI-H and MSS (A, B, C), stage II and III (D, E, F), female and male gender (G, H, I), and distal and proximal tumor locations (K, L, M).

Supplementary Figure 5. The performance of three machine learning models with three different algorithms is evaluated using receiver operating characteristic (ROC) curves.

The curves depict the performance of the combined model in the discovery cohort (A) and the validation cohort (B)

Supplementary Figure 6. The rank coverage score of the combined model compared to NetMHCpan.

The bar graphs display rank coverage scores of validated long peptides identified by the NetMHCpan tool (blue bars) and the combined method (red bars) for individual patients and all patients collectively.

Supplementary Table 1. Clinical Characteristics of 28 CRC patients

Supplementary Table 2. Comparison of HLA calling accuracy between Acras-HLA and Optitype .

Supplementary Table 3. Quality metrics of TCRβ sequencing

Supplementary Table 4. Percentage of unique and expanded TCR clones from 28 CRC patients

Supplementary Table 5. List of immunogenic peptides and non-immunogenic peptides from public databases

Supplementary Table 6. Performance of combined model in an independent dataset.

REFERENCES

1. K. Ganesh *et al.*, Immunotherapy in colorectal cancer: rationale, challenges and potential. *Nat Rev Gastroenterol Hepatol* **16**, 361-375 (2019).
2. J. C. Dudley, M. T. Lin, D. T. Le, J. R. Eshleman, Microsatellite Instability as a Biomarker for PD-1 Blockade. *Clin Cancer Res* **22**, 813-820 (2016).
3. D. T. Le *et al.*, PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med* **372**, 2509-2520 (2015).
4. M. J. Overman *et al.*, Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study. *Lancet Oncol* **18**, 1182-1191 (2017).
5. D. T. Le *et al.*, Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* **357**, 409-413 (2017).
6. Y. Yu *et al.*, Neoantigen-reactive T cells exhibit effective anti-tumor activity against colorectal cancer. *Hum Vaccin Immunother* **18**, 1-11 (2022).
7. V. M. Kim *et al.*, Neoantigen-based EpiGVAX vaccine initiates antitumor immunity in colorectal cancer. *JCI Insight* **5** (2020).
8. E. Blass, P. A. Ott, Advances in the development of personalized neoantigen-based therapeutic cancer vaccines. *Nat Rev Clin Oncol* **18**, 215-229 (2021).
9. D. Miao *et al.*, Genomic correlates of response to immune checkpoint therapies in clear cell renal cell carcinoma. *Science* **359**, 801-806 (2018).
10. M. Yarchoan, A. Hopkins, E. M. Jaffee, Tumor Mutational Burden and Response Rate to PD-1 Inhibition. *N Engl J Med* **377**, 2500-2501 (2017).
11. J. Hundal *et al.*, pVACtools: A Computational Toolkit to Identify and Visualize Cancer Neoantigens. *Cancer Immunol Res* **8**, 409-420 (2020).
12. Z. S. Chheda *et al.*, Novel and shared neoantigen derived from histone 3 variant H3.3K27M mutation for glioma T cell therapy. *J Exp Med* **215**, 141-157 (2018).
13. B. Reynisson, B. Alvarez, S. Paul, B. Peters, M. Nielsen, NetMHCpan-4.1 and NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent

motif deconvolution and integration of MS MHC eluted ligand data. *Nucleic Acids Res*
48, W449-W454 (2020).

14.   T. N. Schumacher, R. D. Schreiber, Neoantigens in cancer immunotherapy. *Science* **348**,
69-74 (2015).

15.   I. Chen, M. Y. Chen, S. P. Goedegebuure, W. E. Gillanders, Challenges targeting cancer
neoantigens in 2021: a systematic literature review. *Expert Rev Vaccines* **20**, 827-837
(2021).

16.   C. Szeto, C. A. Lobos, A. T. Nguyen, S. Gras, TCR Recognition of Peptide-MHC-I: Rule
Makers and Breakers. *Int J Mol Sci* **22** (2020).

17.   S. Guerder, R. A. Flavell, T-cell activation. Two for T. *Curr Biol* **5**, 866-868 (1995).

18.   M. S. Kuhns, H. B. Badgandi, Piecing together the family portrait of TCR-CD3
complexes. *Immunol Rev* **250**, 120-143 (2012).

19.   J. P. Rast *et al.*, alpha, beta, gamma, and delta T cell antigen receptor genes arose early in
vertebrate phylogeny. *Immunity* **6**, 1-11 (1997).

20.   E. Rosati *et al.*, Overview of methodologies for T-cell receptor repertoire analysis. *BMC*
Biotechnol **17**, 61 (2017).

21.   K. W. Wucherpfennig, E. Gagnon, M. J. Call, E. S. Huseby, M. E. Call, Structural
biology of the T-cell receptor: insights into receptor assembly, ligand recognition, and
initiation of signaling. *Cold Spring Harb Perspect Biol* **2**, a005140 (2010).

22.   N. Porciello, O. Franzese, L. D'Ambrosio, B. Palermo, P. Nistico, T-cell repertoire
diversity: friend or foe for protective antitumor response? *J Exp Clin Cancer Res* **41**, 356
(2022).

23.   Y. C. Lu *et al.*, Direct identification of neoantigen-specific TCRs from tumor specimens
by high-throughput single-cell sequencing. *J Immunother Cancer* **9** (2021).

24.   L. Mazzotti *et al.*, T-Cell Receptor Repertoire Sequencing and Its Applications: Focus on
Infectious Diseases and Cancer. *Int J Mol Sci* **23** (2022).

25.   https://www.10xgenomics.com/resources/datasets/cd-8-plus-t-cells-of-healthy-donor-1-1-
standard-3-0-2.

26.   https://www.10xgenomics.com/resources/datasets/cd-8-plus-t-cells-of-healthy-donor-2-1-
standard-3-0-2.

27.   https://www.10xgenomics.com/resources/datasets/cd-8-plus-t-cells-of-healthy-donor-3-1-
standard-3-0-2.

28.   https://www.10xgenomics.com/resources/datasets/cd-8-plus-t-cells-of-healthy-donor-4-1-
standard-3-0-2.

29.   N. Tickotsky, T. Sagiv, J. Prilusky, E. Shifrut, N. Friedman, McPAS-TCR: a manually
curated catalogue of pathology-associated T cell receptor sequences. *Bioinformatics* **33**,
2924-2929 (2017).

30.   J. Schmidt *et al.*, Prediction of neo-epitope immunogenicity reveals TCR recognition
determinants and provides insight into immunoediting. *Cell Rep Med* **2**, 100194 (2021).

31.   M. Shugay *et al.*, VDJdb: a curated database of T-cell receptor sequences with known
antigen specificity. *Nucleic Acids Res* **46**, D419-D427 (2018).

32.   R. Vita *et al.*, The Immune Epitope Database (IEDB): 2018 update. *Nucleic Acids Res* **47**,
D339-D343 (2019).

33.   W. Zhang *et al.*, PIRD: Pan Immune Repertoire Database. *Bioinformatics* **36**, 897-903
(2020).

34.   M. B. Amin *et al.*, The Eighth Edition AJCC Cancer Staging Manual: Continuing to build
a bridge from a population-based to a more "personalized" approach to cancer staging.
CA Cancer J Clin **67**, 93-99 (2017).
35.   G. J. Tong *et al.*, Comparison of the eighth version of the American Joint Committee on
Cancer manual to the seventh version for colorectal cancer: A retrospective review of our
data. *World J Clin Oncol* **9**, 148-161 (2018).
36.   B. Q. T. Nguyen *et al.*, Improvement in neoantigen prediction via integration of RNA
sequencing data for variant calling. *Front Immunol* **14**, 1251603 (2023).
37.   V. J. Severine Catreux, Lisa Murray, Rami Mehio, Gavin Parnaby, Cooper Roddey,
Michael   Ruehle,   Wei-Ting   Chen,   Fan   Zhang.
https://www.illumina.com/science/genomics-research/articles/dragen-shines-again-
precisionfda-truth-challenge-v2.html.
38.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
39.   A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
40.   S. Andrews (2010) FastQC: a quality control tool for high throughput sequence data.
(Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom).
41.   D. C. Koboldt *et al.*, VarScan 2: somatic mutation and copy number alteration discovery
in cancer by exome sequencing. *Genome Res* **22**, 568-576 (2012).
42.   H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-2079 (2009).
43.   http://broadinstitute.github.io/picard.
44.   W. McLaren *et al.*, The Ensembl Variant Effect Predictor. *Genome Biol* **17**, 122 (2016).
45.   https://www.takarabio.com/learning-centers/next-generation-sequencing/bioinformatics-
resources/cogent-ngs-immune-profiler.
46.   A. Szolek *et al.*, OptiType: precision HLA typing from next-generation sequencing data.
Bioinformatics **30**, 3310-3316 (2014).
47.   G. Li *et al.*, Splicing neoantigen discovery with SNAF reveals shared targets for cancer
immunotherapy. *Sci Transl Med* **16**, eade2886 (2024).
48.   J. Hundal *et al.*, pVAC-Seq: A genome-guided in silico approach to identifying tumor
neoantigens. *Genome Med* **8**, 11 (2016).
49.   J. Hundal *et al.*, Accounting for proximal variants improves neoantigen prediction. *Nat
Genet* **51**, 175-179 (2019).
50.   T. Lu *et al.*, Deep learning-based prediction of the T cell receptor-antigen binding
specificity. *Nat Mach Intell* **3**, 864-875 (2021).
51.   C. Zhou *et al.*, pTuneos: prioritizing tumor neoantigens from next-generation sequencing
data. *Genome Med* **11**, 67 (2019).
52.   Z. Moodie *et al.*, Response definition criteria for ELISPOT assays revisited. *Cancer
Immunol Immunother* **59**, 1489-1501 (2010).
53.   F. Chen *et al.*, Neoantigen identification strategies enable personalized immunotherapy in
refractory solid tumors. *J Clin Invest* **129**, 2056-2070 (2019).
54.   S. Hey *et al.*, Analysis of CDR3 Sequences from T-Cell Receptor beta in Acute
Respiratory Distress Syndrome. *Biomolecules* **13** (2023).
55.   V. Roudko *et al.*, Shared Immunogenic Poly-Epitope Frameshift Mutations in
Microsatellite Unstable Tumors. *Cell* **183**, 1634-1649 e1617 (2020).

56.   S. Maleki Vareki, High and low mutational burden tumors versus immunologically hot
and cold tumors and response to immune checkpoint inhibitors. *J Immunother Cancer* **6**,
157 (2018).

57.   L. Laghi *et al.*, Prognostic and Predictive Cross-Roads of Microsatellite Instability and
Immune Response to Colon Cancer. *Int J Mol Sci* **21** (2020).

58.   M. B. Yassai, Y. N. Naumov, E. N. Naumova, J. Gorski, A clonotype nomenclature for T
cell receptors. *Immunogenetics* **61**, 493-502 (2009).

59.   A. Vitiello, M. Zanetti, Neoantigen prediction and the need for validation. *Nat Biotechnol*
35, 815-817 (2017).

60.   J. Xia *et al.*, NEPdb: A Database of T-Cell Experimentally-Validated Neoantigens and
Pan-Cancer Predicted Neoepitopes for Cancer Immunotherapy. *Front Immunol* **12**,
644637 (2021).

61.   W. J. Zhou *et al.*, NeoPeptide: an immunoinformatic database of T-cell-defined
neoantigens. *Database (Oxford)* **2019** (2019).

62.   X. Tan *et al.*, dbPepNeo: a manually curated database for human tumor neoantigen
peptides. *Database (Oxford)* **2020** (2020).

63.   G. Zhang, L. Chitkushev, L. R. Olsen, D. B. Keskin, V. Brusic, TANTIGEN 2.0: a
knowledge base of tumor T cell antigens and epitopes. *BMC Bioinformatics* **22**, 40
(2021).

64.   J. Wu *et al.*, TSNAdb: A Database for Tumor-specific Neoantigens from
Immunogenomics Data Analysis. *Genomics Proteomics Bioinformatics* **16**, 276-282
(2018).

65.   A. Montemurro *et al.*, NetTCR-2.0 enables accurate prediction of TCR-peptide binding
by using paired TCRalpha and beta sequence data. *Commun Biol* **4**, 1060 (2021).

66.   E. S. Borden, K. H. Buetow, M. A. Wilson, K. T. Hastings, Cancer Neoantigens:
Challenges and Future Directions for Prediction, Prioritization, and Validation. *Front*
Oncol **12**, 836821 (2022).

67.   K. Joshi *et al.*, Spatial heterogeneity of the T cell receptor repertoire reflects the
mutational landscape in lung cancer. *Nat Med* **25**, 1549-1559 (2019).

68.   R. Motta *et al.*, Immunotherapy in microsatellite instability metastatic colorectal cancer:
Current status and future perspectives. *J Clin Transl Res* **7**, 511-522 (2021).

69.   N. Xie *et al.*, Neoantigens: promising targets for cancer therapy. *Signal Transduct Target*
Ther **8**, 9 (2023).

70.   M. Muller *et al.*, Machine learning methods and harmonized datasets improve
immunogenic neoantigen prediction. *Immunity* 10.1016/j.immuni.2023.09.002 (2023).

71.   P. Malekzadeh *et al.*, Antigen Experienced T Cells from Peripheral Blood Recognize p53
Neoantigens. *Clin Cancer Res* **26**, 1267-1276 (2020).