1  **The similarities in microbial and chemical patterns of fermentation in two open**
2  **environments were promoted by using 150-year-old Nukadoko as starters**
3

4  **Authors**
5  Marin Yamaguchi[1,2,3], Kota Imai[1,4], Dominique Chen[5], Young ah Seong[6], Kazuhiro Jo[7],
6  Kohei Ito[1*]
7

**Affiliation**
1. BIOTA Inc., Tokyo, 101-0022, Japan
2. Faculty of Pharmaceutical Sciences, The University of Tokyo, Tokyo, 113-8654,
Japan
3. College of Arts and Sciences, The University of Tokyo, Tokyo, 153-8902, Japan
4. Graduate School of Life and Medical Sciences, Doshisha University, Kyoto, 610-
0394, Japan
5. Faculty of Letters, Arts and Sciences, Waseda University, Tokyo, 162-8644, Japan
6. Faculty of Engineering and Design, Hosei University, Tokyo, 102-8160, Japan
7. Faculty of Design, Kyushu University, Fukuoka, 815-8540, Japan
18

19   * Corresponding author
20  e-mail: kohei@biota.ne.jp
21

**Abstract**
Nukadoko, a fermented rice bran employed in traditional Japanese pickling, uses lactic
acid bacteria to ferment vegetables. Here, we report the microbial and chemical data of a
mixture of matured 150-year-old nukadoko and commercially available rice bran placed in two
open environments over 29 days. Across the two environments, *Loigolactobacillus* was
identified as the dominant microbial genera in the later stages of fermentation in nukadoko.
The period of increase in the relative abundance of *Loigolactobacillus* correlated with a
decrease in pH and Oxidation-Reduction Potential (ORP) values. While the two environments
showed a difference in the rate of change in microbial diversity, they shared the common
process through which *Loigolactobacillus* outcompeted adventitious bacteria in nukadoko, as
indicated by the alpha and beta diversity index. Thus, the similarities in microbial and chemical
data across two open environments during fermentation using starters indicate that starters
contribute to the stability of fermentation in open environments.
35

36  **Keywords**
37  fermentation, microbiota, Nukadoko, lactic acid bacteria, loigolactobacillus.
38

39  **Acknowledgements**
40       We would like to thank "Hyaku-goju-nen no Nukadoko Hozonkai" for providing their
41  matured Nukadoko and giving us permission to analyze its microbiota.

## Introduction

Nukadoko, a fermented rice bran employed in traditional Japanese pickling, uses lactic acid bacteria (LAB) to ferment vegetables (*Nukazuke*). Traditionally, the nukadoko preparation process involves the natural fermentation of rice bran, continuously renewed by adding fresh bran. A previous study reported that the initial fermentation time was thereby reduced using an aged nukadoko as a fermentation starter ready for use (Ono et al., 2014). Indeed, many modern fermentation practices often involve the use of starters, which also serve the purpose of preventing food spoilage in the fermentation process of meat, fish, and dairy products (2).

The microbiota of the fresh rice bran used in nukadoko is especially rich in microbial diversity, giving rise to the distinct flavor of nukadoko (Ono et al., 2014). The type of rice bran used, salinity levels, temperature, air exposure, and fermentation time can all be factors that distinguish the nukadoko microbiota (3). However, in a previous study involving four types of nukadoko, such microbial diversity eventually converges to represent similar microbial communities as the fermentation process continues (Ono et al., 2014). A similar phenomenon has also been observed in other fermented foods such as kimchi, where samples of various types of kimchi at later stages of fermentation were mainly composed of *Lactobacillales*, related to LAB (4). However, still little is known about the complete picture of the nukadoko microbiota, as well as the chemical changes during the fermentation process.

In this study, we examined the chemical and microbial changes during the fermentation of nukadoko in two different open environments over 29 days.

## Materials and Methods
### Preparation of *nukadoko* and sample collection

Nukadoko utilized in this study was prepared by mixing commercially available rice bran (Maruho Foods, Kagawa, Japan) with matured 150-year-old nukadoko. The long-aged (approximately 150 years) nukadoko used in this experiment were obtained from a nukadoko manufacturer (Hyaku-Gojunen-no-Nukadoko-Hozonkai) in Fukuoka City, Japan. The ingredients of the 150-year-old nukadoko were rice bran, salt, water, *kombu* (a type of edible seaweed), dried young sardines, bird's eye chili, Sichuan peppercorns, and rice malt. 250 g of the 150-year-old nukadoko was mixed with fresh rice bran (500 g), salts (125 g), and distilled water (500 mL). The mixed samples were placed in two locations for 29 days.

Three participants at both Location 1 and Location 2 were assigned to stir nukadoko on a daily basis at each location (Fig. 1). We chose Location 1, Hakko Department, a commercial establishment specializing in fermented foods and also offering dining options, and location 2, an exhibition at The National Museum of Emerging Science and Innovation, as the experimental locations. We deployed two sets of nukadoko vessel equipped with pH and ORP sensors (27) to each location. Among Location 2, a public space where temperature and humidity are strictly controlled, and Location 1, a more enclosed environment, we aimed to investigate whether the patterns of nukadoko fermentation vary among the two locations with such different characteristics and user demographics. The frequency of stirring was determined according to the hygiene standards of each facility and the operating patterns of the staff: once daily at Location 1 and twice daily at Location 2, once every morning and once every evening. For each stirring session, the participants were asked to mix nukadoko thoroughly from the bottom 2-3 times, with each session spanning approximately 3 minutes.

Sample collection took place on specific days, namely days 0, 3, 6, 9, 15, 17, 22, and 29 for Location 1, and on days 0, 3, 6, 9, 14, 18, 22, and 29 for Location 2, using individually wrapped disposable plastic spoons and 50mL polypropylene conical tubes. Thus, we obtained 16 samples in total from the two locations. Samples were stored in a freezer after collection and kept at -20°C until DNA extraction.

**Chemical analysis**

We employed high-precision pH and ORP sensors for robots developed by Atlas Scientific (USA) and inserted them into the nukadoko rice bran. These sensors were connected to electronic circuits (Raspberry Pi 4) that transmit the sensed data to a cloud database (Amazon Web Service RDS) each minute during the whole period.

**Total DNA extraction and high-throughput sequencing**

DNA isolation from 50 ml nukadoko samples was performed using the beat-beating method using the multi-beads shocker (Yasui Kikai, Osaka). The extracted DNA was quantified using Qubit™ (Thermo Fisher Scientific, Massachusetts). The DNA was further purified using a GenCheck® DNA Extraction Kit Type S/F. The first PCR cycle was conducted using ExTaq HS DNA Polymerase, where specific primers 341F (5'-CCTACGGGNGGCWGCAG-3') and 805R (5'-GACTACHVGGGTATCTAATCC-3') of the V3-V4 region of the 16S DNA were amplified (5). 2 μl of the first PCR product, purified using AMPure XP beads, served as a template for library preparation. The second PCR product was again purified using AMPure XP beads. Amplicon sequencing was performed using 460 bp paired-end sequencing on the Illumina MiSeq platform (Illumina Inc., San Diego, CA, USA) at FASMAC Co., Ltd (Kanagawa, Japan).

**Microbial analysis**

Microbial analysis was performed as previously regarding previous studies (6). Briefly, raw FASTQ data were imported into the QIIME2 platform (2022.8) as qza files (7). Denoising sequences and quality control were performed using the QIIME dada2, and finally, sequences were produced into amplicon sequence variants (ASVs) (8). ASVs were assigned to the SILVA database's SSU 138 using the QIIME feature-classifier classification scikit-learn package (9,10). ASVs classified as mitochondria, chloroplast, or unassigned were excluded from subsequent analysis. Subsampling was performed to reduce bias due to differences in read depth between samples. The beta diversity indices weighted and unweighted UniFrac distances were calculated, and the microbial community structure differences between the two groups were visualized following a principal coordinate analysis (PCoA). Data were visualized using R (version 4.0.4), ggplot2 (version 3.4.3), and ggprism (version 1.0.4) (11,12).

**Statistical analysis**

Mann–Whitney U tests were used to compare all combinations of elements within the two locations.

**Data Availability**

The datasets generated through 16S rRNA amplicon sequencing are available and deposited in the DDBJ Sequence Read Archive (SRA) database under accession numbers DRR514074-DRR514089 and BioProject PRJDB17051.

**Results**

**Changes in ORP and pH values during fermentation of nukadoko**

Fig. 2a shows the gradual decrease in pH values for both samples, indicating the fermentation rate. The most rapid decrease in pH occurred between days 14 and 16 for Location 1 and between days 3 and 7 for Location 2 (Fig. 2a). The pH values indicated the difference in the time period in which rapid fermentation occurs among the two sample locations. Finally,

the pH values at the two locations on the last day (day 29) were significantly different ($p < 0.05$).

The ORP values showed variability throughout the experiment. The decline in the ORP values from approximately -50 mV to -250 mV occurs in the Location 1 sample across 5 days (days 9-13), and the decline from approximately 200 mV to -100 mV occurs in the Location 2 sample across 8 days (days 0-7) (Fig. 2b). Similarly to the results obtained through the differences in pH values, the variation in ORP across the experiment is also indicative of the different nature of the fermentation process among Location 1 and Location 2. Finally, the ORP values at the two locations on the last day (day 29) were not significantly different ($p = 0.0507$).

**Increase of the relative abundance of LAB**

The number of raw reads obtained from 16 samples remaining after quality filtering was 323,217. After removing ASVs classified as mitochondrial, chloroplast, or unassigned, the total number of reads was 201,452. The ASVs from the samples were classified, with the most abundant genera being *Loigolactobacillus*, *Pantoea*, *Sphingomonas*, *Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium, Pseudomonas*, and *Staphylococcus* on day 0 (Fig. 3). Over time, *Loigolactobacillus* dominates the majority in both Location 1 (Day 0: 69.6 % and Day 29: 99.0 %) and Location 2 (Day0: 69.5 % and Day 29: 99.4 %) samples. Despite the similar pattern, the rate of decrease in microbial diversity follows different paths, as the Location 1 sample continued to exhibit a wider variety of microbes over a longer period, whereas, in the Location 2 sample, *Loigolactobacillus* dominated the majority at a much earlier phase, at around day 6 of the experiment.

The microbial community structures of both samples were compared using principal component analysis (PCoA) by weighted UniFrac distance metrics. At the beginning, both Location 1 and Location 2 show similar microbial compositions, consisting of a broad range of microbial genera, as shown in Fig. 3. This is indicated by the proximity with which the two sample points on day 0 are located on the PCoA plot. However, between days 3 and 9, Location 1 and Location 2 samples exhibited distinct microbial compositions and eventually showed similarities again towards the end of the experiment (Fig. 4a). Similarly to the results indicated by Fig. 2a, this indicates the difference in the process and rate of fermentation among the two sample locations. Indeed, at around day 3, the sample point from the Location 2 sample approaches the clustered sample groups, indicating the end of the fermentation process. However, in the Location 1 sample, the data points started to cluster together on day 14, displaying the gap in fermentation rate among the two sample locations. The sample points at the end of fermentation can be clustered together, where PC2 measures around 0%, indicating that the fermentation process exhibit similar microbial trends across the two locations over time.

The Shannon Entropy values also exhibit similar results, where the decrease in the alpha diversity occurs exponentially, and then eventually stabilizes. The alpha diversity measures high among the Location 1 sample until day 14 (Fig. 4b), and its decrease at the end coincides with the period *Loigolactobacillus* takes over as the most dominant genus. The days in which the alpha diversity decreases correlate with the days *Loigolactobacillus* dominates, occurring between days 14-15 for the Location 1 sample and between days 6-7 for the Location 2 sample. This period of change in microbial diversity aligns with the period of changes in pH values obtained through chemical analysis.

**Discussion**
**Changes in the ORP and the pH values during fermentation of nukadoko**

The ORP values are often used to evaluate the micro-aerobic status of the environments in which fermentation take place (13). In a previous study involving the

*Lactobacillus mucosae* species the ORP values of micro-aerobic fermentation was determined to be between -300 and -150 mV (14). The readings of ORP values obtained in this experiment mainly lie within such micro-aerobic fermentation level, indicating the normal fermentation process induced by the daily mixings and the concurrent air intake into nukadoko. This is also consistent with previous findings that ORP values reflect activities of both aerobic (positive ORP) and anaerobic (negative ORP) bacteria, and thus ORP values are indicative of the level of fermentation (15). While aerobic growth may promote oxidative stress in LAB species, oxygen availability is an important factor that determines the level of aromatic compounds produced by LAB (16). Aromatic compounds, mainly acetoin and diacetyl, are known to impact the flavor of dairy products during food fermentations with LAB (17), suggesting that the distinct flavors of nukadoko may also be supported by similar aromatic compounds. Thus, the micro-aerobic environments in the two locations in this experiment may have influenced the metabolic routes of LAB to produce acetoin and diacetyl, giving rise to the nukadoko flavor.

A previous study has also demonstrated the abundance of LAB and the accompanying pH decrease during the late fermentation period, similar to the results obtained in this experiment (18). Among the two sample locations, the time points of chemical and microbial changes are aligned, demonstrating the advancement of nukadoko fermentation. Changes in chemical values including pH values occurred between days 14-16 in Location 1 and days 3-7 in Location 2, which coincide with the days in which alpha diversity decreases for both locations. This suggests that the chemical and microbiota data of nukadoko fermentation are interrelated.

**Changes in the taxonomic composition during fermentation of nukadoko**

Our results showed that *Loigolactobacillus*, an anaerobic, homo-fermentative bacteria, became dominant in both environments. Similarly, in another study, *Loigolactobacillus* was identified as the dominant genus in nukadoko samples (19). The process through which adventitious bacteria are outcompeted by LAB at the later stages of fermentation is dependent on the types of LAB that dominate the microbiota (17). A previous study (18) investigated the chemical changes during the initial stage of fermentation across four samples made of different types of rice bran. Lactate concentration can be attributed to the differences between hetero-fermentative LAB during the early fermentation stage and homo-fermentative LAB at the later stages (19). The dominance of *Loigolactobacillus* in our experiment suggests the abundance of homo-fermentative bacteria in 150-year-old nukadoko, indicating its matured fermentation state. It must be noted that *Loigolactobacillus* is a genus deriving from the reclassification of *Lactobacillus* (20).

In another study, Sakamoto identified two predominant species of *Lactobacillus* that maintain a nukadoko environment that prevents contamination by other microorganisms, conserving the distinct nukadoko flavor and preventing further spontaneous fermentation by adventitious microbes (21). Indeed, within the architectural environment in which the samples are placed, various microbes, known as the Microbiome of the Built Environment (MoBE), exist, which can influence the production of fermented foods (22,23). MoBE has been found to contaminate the fermentation starter in a *sake* brewery (24). Similarly to the effects of *Lactobacillus* dominance of the microbiota in other experiments involving nukadoko, the presence of *Loigolactobacillus* in our experiment may also provide stability to the fermentation process and prevent adventitious microbes from contaminating the nukadoko environment. The high prevalence of *Loigolactobacillus* from day 0 may have resulted from an increased amount of readily matured nukadoko. We used an increased amount of starter compared to a previous study (18) in order to support fermentation in open environments.

Bacterial genus recently isolated from nukadoko in other findings, including Weisella (doi: 10.1128/MRA.01160-21) and Bacillus (doi: 10.1128/MRA.00705-21), were rarely

246 present in the nukadoko samples. This suggests the wide variety of microbiota in different
247 nukadoko environments.
248   We also examined the microbial composition of nukadoko over the fermentation period,
249 where most adventitious bacteria such as *Pantoea, Pseudomonas*, and *Staphylococcus*
250 originating from sufficiently aged nukadoko were not detected by the end of the fermentation
251 process in both Location 1 and Location 2 samples. Such changes in microbial composition
252 resemble data obtained through previous studies, where one (24) notes the changes from the
253 prevalence of *Enterobacter*, a type of gram-negative bacteria encompassing *Pantoea*, to that of
254 LAB in the later stages of fermentation. Thus, adventitious bacteria are continuously
255 outcompeted by *Lactobacillus* and/or *Loigolactobacillus* and their lactic acid products. The
256 LAB-dominant microbiota is not unique to nukadoko and can be seen in several fermented
257 foods (25).
258
259 **The microbial diversity of the two locations eventually became similar**
260   Fig. 4a indicates that the two groups exhibit differences in the sample microbiota at the
261 initial stages of fermentation but later show high similarities. Fig. 4b indicates the gradual
262 decrease in the Shannon Diversity index across two groups at the later stages of fermentation.
263 It is suggested that these changes in microbial diversity reflect the gradual dominance of
264 *Loigolactobacillus* throughout the fermentation process. This is consistent with the findings in
265 other fermented foods, including Suancai, a traditional sauerkraut in Northeast China. In an
266 experiment involving Suancai fermentation, 16S rRNA microbial analysis indicated that by
267 day 30 of fermentation, 100% of the sample bacteria comprised of LAB (26). Indeed, the
268 dominance of LAB at later stages of fermentation is often observed in various types of
269 fermented foods, which reflects the common modern fermentation practice of introducing
270 starters so that fermentation-contributing microbes can dominate the microbiota at a more rapid
271 pace (2).
272   It must be noted that *Loigolactobacillus* was detected at a relatively high abundance in
273 the nukadoko from the initial stage of fermentation, which may have been caused by an
274 excessive amount of starter. In addition, due to the analysis at the ASV levels, it is yet to be
275 known whether the detected *Loigolactobacillus* consists of a single or multiple species.
276
277 **Conclusion**
278   Through microbial and chemical analyses, we have demonstrated the similarities in
279 microbial and chemical patterns during nukadoko fermentation in two different open
280 environments. Such similarities in the fermentation process observed among the two sample
281 locations demonstrates how the 150-year-old nukadoko starter contributes to the creation of
282 the optimal environment for nukadoko fermentation, characterized by the dominance of
283 *Loigolactobacillus*.

284
285 **Figures**
286

287
288 **Fig. 1** Pictures demonstrating the maintenance of nukadoko in two open environments.
289

291
292 **Fig. 2** Regression plots of time-series data of OPR (a) and pH (b) values of the nukadoko
293 environment in Location 1 and Location 2 samples over 29 days. The red line represents data
294 from the Location 1 sample, and the blue line represents data from the Location 2 sample.
295
296

297
**Fig. 3** Stacked bar plots of the genus-level taxonomic composition during the fermentation of
nukadoko over 29 days. The top 15 genera are listed, and the rest are noted as Remainder.
300
301

**Fig. 4** Changes in the microbial diversity during the fermentation of nukadoko samples. (a) Principal-coordinate analysis (PCoA) by Weighted UniFrac distance metrics of Location 1 and Location 2 samples. The line connects the data points across 29 experimental days. The data points represented by the circles indicate Location 1, and data points represented by the triangles indicate Location 2. (b) Line plots of Shannon diversity index of Location 1 and Location 2 samples over 29 days.

309
310

## References

1. Ono H, Nishio S, Tsurii J, Kawamoto T, Sonomoto K, Nakayama J. Monitoring of the microbiota profile in nukadoko, a naturally fermented rice bran bed for pickling vegetables. J Biosci Bioeng [Internet]. 2014 Nov;118(5):520–5. Available from: http://dx.doi.org/10.1016/j.jbiosc.2014.04.017

2. Imai K, Niwa R, Fujioka M, Ito K. Understanding the quality and safety of food production through the lens of The Microbiome of The Built Environment. Biosci Biotechnol Biochem [Internet]. 2023 Nov 22; Available from: http://dx.doi.org/10.1093/bbb/zbad164

3. Osawa R, Kuroiso K, Goto S, Shimizu A. Isolation of tannin-degrading lactobacilli from humans and fermented foods. Appl Environ Microbiol [Internet]. 2000 Jul;66(7):3093–7. Available from: http://dx.doi.org/10.1128/AEM.66.7.3093-3097.2000

4. Park E-J, Chun J, Cha C-J, Park W-S, Jeon CO, Bae J-W. Bacterial community analysis during fermentation of ten representative kinds of kimchi with barcoded pyrosequencing. Food Microbiol [Internet]. 2012 May;30(1):197–204. Available from: http://dx.doi.org/10.1016/j.fm.2011.10.011

5. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. Nucleic Acids Res [Internet]. 2013 Jan 7;41(1):e1. Available from: http://dx.doi.org/10.1093/nar/gks808

6. Ito K, Niwa R, Yamagishi Y, Kobayashi K, Tsuchida Y, Hoshino G, et al. A unique case in which Kimoto-style fermentation was completed with Leuconostoc as the dominant genus without transitioning to Lactobacillus. J Biosci Bioeng [Internet]. 2023 Jun;135(6):451–7. Available from: http://dx.doi.org/10.1016/j.jbiosc.2023.03.007

7. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol [Internet]. 2019 Aug;37(8):852–7. Available from: http://dx.doi.org/10.1038/s41587-019-0209-9

8. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods [Internet]. 2016 Jul;13(7):581–3. Available from: http://dx.doi.org/10.1038/nmeth.3869

9. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome [Internet]. 2018 May 17;6(1):90. Available from: http://dx.doi.org/10.1186/s40168-018-0470-z

10. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res [Internet]. 2013 Jan;41(Database issue):D590-6. Available from: http://dx.doi.org/10.1093/nar/gks1219

11. Wickham H. ggplot2 [Internet]. Springer New York; [cited 2023 Nov 8]. 8 p. Available from: https://link.springer.com/book/10.1007/978-0-387-98141-3

12.   Creators Charlotte Dawson1 Show affiliations 1. University of Cambridge.
csdaw/ggprism: ggprism 1.0.2 [Internet]. Available from:
https://zenodo.org/record/4556067

13.   Barati rashvanlou R, Rezaee A, Farzadkia M, Gholami M, Kermani M. Effect of micro-
aerobic process on improvement of anaerobic digestion sewage sludge treatment: flow
cytometry and ATP assessment. RSC Adv [Internet]. 2020;10(59):35718–28. Available
from: http://dx.doi.org/10.1039/d0ra05540a

14.   Li D, Jiao C, He W, Yan Z, Yuan Y, Li Z, et al. Comparison of micro-aerobic and
anaerobic fermentative hydrogen production from corn straw. Int J Hydrogen Energy
[Internet]. 2016 Apr;41(12):5456–64. Available from:
http://dx.doi.org/10.1016/j.ijhydene.2016.01.141

15.   Chen D, Ogura H, Seong YA. NukaBot: Research and design of a human-microbe
interaction model. In: The 2019 Conference on Artificial Life [Internet]. Cambridge,
MA: MIT Press; 2019. Available from:
https://www.mitpressjournals.org/doi/abs/10.1162/isal_a_00138

16.   Zotta T, Parente E, Ricciardi A. Aerobic metabolism in the genus*Lactobacillus*: impact
on stress response and potential applications in the food industry. J Appl Microbiol
[Internet]. 2017 Apr;122(4):857–69. Available from:
http://dx.doi.org/10.1111/jam.13399

17.   Gänzle MG. Lactic metabolism revisited: metabolism of lactic acid bacteria in food
fermentations and food spoilage. Curr Opin Food Sci [Internet]. 2015 Apr;2:106–17.
Available from: http://dx.doi.org/10.1016/j.cofs.2015.03.001

18.   Nakayama J, Hoshiko H, Fukuda M, Tanaka H, Sakamoto N, Tanaka S, et al. Molecular
monitoring of bacterial community structure in long-aged nukadoko: pickling bed of
fermented rice bran dominated by slow-growing lactobacilli. J Biosci Bioeng [Internet].
2007 Dec;104(6):481–9. Available from: http://dx.doi.org/10.1263/jbb.104.481

19.   Niwa R, Chen D, Seong YA, Jo K, Ito K. Direct contact of fermented rice bran beds
promotes food-to-hand transmission of lactic acid bacteria. FEMS Microbiol Lett
[Internet]. 2023 Nov 13; Available from: http://dx.doi.org/10.1093/femsle/fnad120

20.   Zheng J, Wittouck S, Salvetti E, Franz CMAP, Harris HMB, Mattarelli P, et al. A
taxonomic note on the genus Lactobacillus: Description of 23 novel genera, emended
description of the genus Lactobacillus Beijerinck 1901, and union of Lactobacillaceae
and Leuconostocaceae. Int J Syst Evol Microbiol [Internet]. 2020 Apr;70(4):2782–858.
Available from: http://dx.doi.org/10.1099/ijsem.0.004107

21.   Sakamoto N, Tanaka S, Sonomoto K, Nakayama J. 16S rRNA pyrosequencing-based
investigation of the bacterial community in nukadoko, a pickling bed of fermented rice
bran. Int J Food Microbiol [Internet]. 2011 Jan 5;144(3):352–9. Available from:
http://dx.doi.org/10.1016/j.ijfoodmicro.2010.10.017

22.   Gilbert JA, Stephens B. Microbiology of the built environment. Nat Rev Microbiol
[Internet]. 2018 Nov;16(11):661–70. Available from: http://dx.doi.org/10.1038/s41579-
018-0065-5

23. Ito K. Understanding the characteristics of the microbiomes of the built environment and their effects on the human body. Indoor Environment [Internet]. 2023;26(1):29–42. Available from: http://dx.doi.org/10.7879/siej.26.29

24. Ito K, Niwa R, Kobayashi K, Nakagawa T, Hoshino G, Tsuchida Y. A dark matter in sake brewing: Origin of microbes producing a Kimoto-style fermentation starter. Front Microbiol [Internet]. 2023 Feb 2;14:1112638. Available from: http://dx.doi.org/10.3389/fmicb.2023.1112638

25. Rezac S, Kok CR, Heermann M, Hutkins R. Fermented Foods as a Dietary Source of Live Organisms. Front Microbiol [Internet]. 2018 Aug 24;9:1785. Available from: http://dx.doi.org/10.3389/fmicb.2018.01785

26. Yang H, Zou H, Qu C, Zhang L, Liu T, Wu H, et al. Dominant microorganisms during the spontaneous fermentation of Suan Cai, a Chinese fermented vegetable. Food Sci Technol Res [Internet]. 2014;20(5):915–26. Available from: http://dx.doi.org/10.3136/fstr.20.915

27. Dominique Chen, Young ah Seong, Hiraku Ogura, Kiichi Moriya, Yuto Mi- tani, and Naoto Sekiya. 2021. Nukabot: Design of Care for Human-Microbe Relationships. In CHI Conference on Human Factors in Computing Systems Extended Abstracts (CHI '21 Extended Abstracts). https://doi.org/10.1145/3411763. 3451605

## Statements & Declarations

### Funding

This work was supported by JSPS KAKENHI (Grant Number 21H03768 and Grant Number 21K18344), of which D.C. is the Principal Investigator.

### Competing Interests

K.I. is a board member at BIOTA Inc., Tokyo, Japan. M.Y. and K.I. are employed by BIOTA Inc. as a part-time developer. All others do not have any competing interests.

### Author Contributions

The study was conceived by all the authors. Chen Dominique and Kohei Ito designed the experiments. Chen Dominique collected samples. Marin Yamaguchi drafted the original manuscript. Kota Imai and Kohei Ito performed microbiome analysis. Young ah Seong, Kazuhiro Jo, and Chen Dominique edited the manuscript and supervised the study. All authors have contributed to the manuscript and approved the submitted version.

### Ethics Approval and Consent to Participate

The research protocol was approved by the Research Ethics Committee, Faculty of Design Engineering, Hosei University (Application Number:2022-4; approval date: March 28, 2023). All procedures were conducted according to the ethics committee's guidelines and regulations. Informed consent was obtained from all participants before starting this study.