Chromosome Structure I: Loop extrusion or boundary:boundary pairing?

Xinyang Bing[1,4]*, Wenfan Ke[2]*, Miki Fujioka[3]*, Amina Kurbidaeva[2,5], Sarah Levitt[2], Mike

Levine[1], Paul Schedl[2], and James B. Jaynes[3]

1: Lewis Sigler Institute, Princeton University, Princeton, NJ. 08544

2: Department of Molecular Biology, Princeton University, Princeton NJ 08544

3: Department of Biochemistry and Molecular Biology, Thomas Jefferson University,

Philadelphia, PA 19107

4: Present address: BlueRock Therapeutics, Cambridge, Ma 20142

5: Present address: Center for Genomics and Systems Biology, New York University, NY 10003

* These authors contributed equally.

Abstract

Two different models have been proposed to explain how the endpoints of chromatin looped

domains ("TADs") in eukaryotic chromosomes are determined. In the first, a cohesin complex

extrudes a loop until it encounters a boundary element roadblock, generating a stem-loop (and an

unanchored loop). In this model, boundaries are functionally autonomous: they have an intrinsic

ability to halt the movement of incoming cohesin complexes that is independent of the properties

of neighboring boundaries. In the second, loops are generated by boundary:boundary pairing. In

this model, boundaries are functionally non-autonomous, and their ability to form a loop depends

upon how well they match with their neighbors. Moreover, unlike the loop-extrusion model,

pairing interactions can generate both stem-loops and circle-loops. We have used a combination

of MicroC to analyze how TADs are organized and experimental manipulations of the *even*

skipped TAD boundary, *homie*, to test the predictions of the "loop-extrusion" and the "boundary-

pairing" models. Our findings are incompatible with the loop-extrusion model and instead

suggest that endpoints of TADs in flies are determined by a mechanism in which boundary

elements physically pair with their partners, either head-to-head, or head-to-tail, with varying

degrees of specificity. How the partners find each other is not clear but is unlikely to require a

loop extrusion mechanism.

Introduction

The chromosomes of multicellular animals have a regular and inheritable physical organization. This was first recognized in studies on the lampbrush chromosomes in amphibian oocytes and the polytene chromosomes of insects (Callan, 1963; Nora et al., 2020; Zhimulev and Belyaeva, 1991; Zhimulev et al., 1983). Subsequent research has shown that the architectural features inferred from analysis of lampbrush and polytene chromosomes are present in chromosomes throughout much of the animal kingdom. The key organizational principle is the subdivision of the chromatin fiber into a series of independent looped domains, commonly called "TADs" (Cavalheiro et al., 2021; Chetverina et al., 2017; Jerković et al., 2020; Matthews and White, 2019; Rowley and Corces, 2018). With important exceptions, the arrangement of TADs along a given chromosome tend to be invariant and are largely independent of the cell type or developmental stage. This regular and inheritable organization is a reflection of the underlying mechanism of TAD formation. TADs are separated from each other by special elements called boundaries or insulators. While these elements have been found in many different species, they have been most fully characterized in *Drosophila* (Cavalheiro *et al.*, 2021; Chetverina *et al.*, 2017). Fly boundaries span DNA sequences of 150 bp to 1.5 kb in length and contain one or more nucleosome-free nuclease-hypersensitive regions. These nuclease-hypersensitive regions are targets for a large collection of DNA binding proteins that have been implicated in boundary function. Some of the fly boundary factors are widely conserved (CTCF, BEN family proteins, GAF), while others appear to be restricted to insect linages (Su(Hw), Pita, Zw5, Pita, Mod(mdg4), BEAF) (Heger et al., 2013; Heger and Wiehe, 2014; Schoborg and Labrador, 2010).

Boundary elements in flies are not only responsible for organizing the chromatin fiber, they also have genetic activities. When interposed between enhancers or silencers and target

promoters, boundary elements block regulatory interactions (Bell et al., 2001; Chetverina et al.,

2014; Chetverina *et al.*, 2017). This insulating activity provides a mechanism for delimiting

units of independent gene activity: genes located between a pair of compatible boundaries are

subject to regulatory interactions with enhancers/silencers present in the same chromosomal

interval, while they are insulated from the effects of enhancers/silencers located beyond either

boundary in adjacent regulatory neighborhoods. Genetic studies suggest that the insulating

activity of boundary elements is a consequence of subdividing the chromosome into a series of

topologically independent domains (Cai and Shen, 2001; Gohl et al., 2011; Muravyova et al.,

2001). Organizing the chromatin fiber into looped domains enhances contacts between

sequences within the loop, while it suppresses contacts with sequences outside of the loop.

While it is not known whether a similar insulation mechanism is at play in mammals, the fact

that mammalian chromosomes are also subdivided into TADs by boundary-like elements

suggests that it might be.

A critical question in chromosome biology is the mechanism(s) responsible for

determining the endpoints of the loop domains, the TADs. In mammals, a novel loop-extrusion

model has been proposed to not only generate chromosomal loops but also determine the

endpoints of those loops (Alipour and Marko, 2012; Fudenberg et al., 2016; Guo et al., 2012;

Guo et al., 2015; Nuebler et al., 2018; Sanborn et al., 2015). In this model, a cohesin complex

initiates loop formation at a loading site within the "loop-to-be" and then extrudes a chromatin

loop until it bumps into CTCF-dependent roadblocks, one on each side of the extruding loop.

The location of these roadblocks combined with the processive action of the cohesin complex

determines the endpoints of each TAD (Davidson and Peters, 2021; Ghosh and Meyer, 2021;

Mirny and Dekker, 2022; Perea-Resa et al., 2021). A key assumption of the loop-extrusion

model is that mammalian boundaries are fully autonomous: they are roadblocks, and their

physical presence in and of itself is sufficient to define the loop endpoint, independent of the

functional properties of neighboring boundaries. In a more specific variant of this model, the

relative orientation of the paired CTCF roadblocks is also important—in order to halt cohesin-

mediated extrusion, the CTCF sites on each side of the loop must have a convergent orientation.

In this model the CTCF roadblocks are able to block an oncoming cohesin complex in only one

direction; however, their intrinsic blocking activity is independent of the orientation of other

CTCF sites in their neighborhood.

While loop extrusion is widely thought to be operative in mammals, the evidence in flies

is inconsistent with this mechanism. Genetics studies have shown that fly boundaries are

functionally non-autonomous and that their activities in both loop formation and gene regulation

depend upon their ability to engage in direct physical interactions with other boundaries

(Chetverina *et al.*, 2014; Chetverina *et al.*, 2017). That fly boundaries might function by

physical pairing was first suggested by studies which showed that the *gypsy* transposon *su(Hw)*

boundary and the bithorax complex (BX-C) *Mcp* boundary can mediate regulatory interactions

(enhancer/silencer: reporter) between transgenes inserted at sites separated by 15-32 MB or more

(Muller et al., 1999; Sigrist and Pirrotta, 1997; Vazquez et al., 1993). Consistent with direct

physical interactions, these distant transgenes co-localize *in vivo*, and in living tissue remain in

contact for extended periods of time (Chen et al., 2018; Li et al., 2011; Vazquez et al., 2006).

Some of the parameters that govern pairing interactions have been defined in two different

transgene assays, insulator bypass and boundary competition. In one version of the bypass

assay, a set of enhancers are placed upstream of two different reporters (Cai and Shen, 2001;

Kyrchanova et al., 2008a; Muravyova *et al.*, 2001). When a single boundary is introduced

between the enhancers and the closest reporter, the enhancers are unable to activate either

reporter. However, if the same boundary is placed downstream of the closest reporter, bypass is

observed. In this case the closest reporter, which is bracketed by the two boundaries, is still

insulated from the enhancers; however, the downstream reporter is activated (Muravyova *et al.*,

2001). Kyrchanova *et al.* (Kyrchanova *et al.*, 2008a) showed that bypass activity is, in most

cases, orientation-dependent. When the same boundary is placed in the upstream and

downstream position, they typically need to be introduced in the opposite orientation. The

reason for this is that self-pairing is head-to-head and this configuration generates a stem-loop

topology, bringing the upstream enhancers in close proximity to the downstream reporter. When

the two boundaries are introduced in the same orientation, the enhancers do not activate the

downstream reporter because a circle-loop rather than a stem-loop is formed (Chetverina *et al.*,

2017; Kyrchanova *et al.*, 2008a). Head-to-head pairing seems to be a common feature of self-

pairing interactions between fly boundaries and was observed for *scs*, *scs'*, *iA2* and *wari*

(Kyrchanova *et al.*, 2008a). A preference for head-to-head interactions is also observed for

heterologous pairing interactions between different boundaries in the *Abdominal-B* (*Abd-B*)

region of the BX-C (Kyrchanova et al., 2011; Kyrchanova et al., 2008b). There are of course

some exceptions. The BX-C *Fab-7* boundary pairs with itself and with its neighbor *Fab-8* in

both orientations (Kyrchanova *et al.*, 2011; Kyrchanova *et al.*, 2008b). Bypass activity also

depends upon finding the proper match. For example, when multimerized dCTCF or Zw5

binding sites are placed both upstream and downstream of the closest reporter, bypass is

observed. However, there is no bypass when multimerized dCTCF sites are tested with

multimerized Zw5 sites (Kyrchanova *et al.*, 2008a). Similar results have been reported for the

Zipic and Pita factors (Zolotarev et al., 2016). Boundary:boundary pairing preferences are also

observed in competition assays in which different boundaries are introduced into the same

transgene (Gohl *et al.*, 2011). In these assays, the insulating activity of a boundary in the

blocking position (between an enhancer and a reporter that is flanked by a 3' boundary) is

challenged by placing a heterologous boundary upstream of the enhancer. If the boundary

upstream of the enhancer is a better match for the boundary located downstream of the reporter,

then the insulating activity of the boundary in the blocking position can be compromised or lost

altogether. These and other experiments argue that fly boundaries are functionally non-

autonomous—that is, their activities depend upon their ability to physically interact with other

boundaries.

However, the differences between the two models are not limited to the mechanisms for

determining the endpoints of TADs. The models also differ in the possible topologies of the

chromatin loops that form TADs. Only two loop topologies can be generated by the cohesin-

extrusion mechanism. One is a stem-loop while the other is an unanchored loop. Depending

upon the location of the road-blocks that halt cohesin extrusion and the cohesin loading sites, the

chromatin fiber would be organized into a series of stem-loops. These stem-loops will be

separated from each other by unanchored loops (see Fig. 1B). This is the configuration that is

often illustrated in articles discussing the loop-extrusion model. If the relevant road-blocks are

very closely juxtaposed, then the unanchored connecting loops will disappear and be replaced by

a series of stem-loops which point in opposite orientations, as illustrated in Fig. 1C. Stem-loops

are also possible in the boundary-pairing model, and they will be formed when two heterologous

boundaries pair with each other head-to-tail (see Fig 1D) in *cis*. Like loop-extrusion, the stem-

loops could be separated from each other by unanchored loops (Fig. 1E). If the boundaries in the

neighborhood pair with each other head-to-tail, a series of connected stem-loops pointing in

opposite orientations will be generated (Fig. 1D). In this case, the main axis of the chromosome

would be a series of paired boundaries (Fig. 1F).

Stem-loops are not, however, the only loop topology that can be generated in the boundary-

pairing model. If the neighboring boundaries pair with each other head-to-head instead of head-

to-tail, a circle-loop structure will be generated (Fig 1G). Like stem-loops, the circle-loops could

be connected by unanchored loops (Fig. 1H). Alternatively, if boundaries pair with both

neighbors head-to-head this will generate a series of linked circle-loops oriented in (more or less)

the same direction (see Fig. 1I).

In the studies reported here we have critically evaluated these two models. We have

used a combination of MicroC to analyze how TADs are organized and experimental

manipulations to test the predictions of the "loop-extrusion" and the "boundary-pairing"

models. For these experimental manipulations, we have used the well-characterized *homie*

boundary from the *even skipped* (*eve*) locus. *homie* together with *nhomie* flank the 16 kb

eve locus, and these two elements share many of the properties of other fly boundaries

(Fujioka et al., 2013; Fujioka et al., 2009). Like the *gypsy* and *Mcp* boundaries, *nhomie* and

homie can mediate long-distance regulatory interactions (Fujioka et al., 2016). For

example, a *lacZ* reporter transgene containing *homie* (or *nhomie*) inserted on one homolog

can be activated by an *eve* enhancer in a *homie*- (or *nhomie*-) containing transgene inserted

on the other homolog over a distance of 2 Mb. Long-distance activation is also observed in

heterologous combinations of *nhomie* and *homie*, but not in combinations with phage

lambda DNA. Similarly, regulatory interactions are observed when a *homie*- or *nhomie*-

containing reporter is inserted at an attP landing site in the *hebe* gene, 142 kb upstream of

the *eve* locus (Chen *et al.*, 2018; Fujioka *et al.*, 2016; Fujioka *et al.*, 2009). In this case, the

eve enhancers drive reporter expression during development in a pattern that mimics the

stage- and tissue-specific expression of the endogenous *eve* gene. Activation also depends

upon the orientation of the boundary (5'→3') relative to the reporter in the transgene. For

homie, the reporter must be located "upstream" of the boundary (5'→3'), just like the *eve*

enhancers (and *eve* gene) are "upstream" of endogenous *homie* (see Fig. 2B). Little or no

activation is observed when the relative orientation is reversed so that the reporter is located

"downstream" of *homie*. For *nhomie*, the reporter must be located "downstream", in the

same relative position as that in which the *eve* enhancers and *eve* gene are located with

respect to endogenous *nhomie* (5'→3': Fig. 2B).

Results

The eve TAD

We used MicroC to probe the TAD organization of the *eve* locus and the genomic region

that extends from *eve* to just beyond the attP site at –142 kb. For this purpose, we chose nuclear

cycle 14 embryos. At this stage of development, most nuclei in our sample will have completed

replication and will be arrested in G2. As sister chromosomes in flies pair with each other in

register, there will be two juxtaposed copies of this region of the chromosome. The sister

chromosomes should be linked to each other by cohesin in preparation for the asynchronous

nuclear divisions at the end of NC14 (Collier and Nasmyth, 2022; Gligoris et al., 2014). While

homolog pairing is limited in pre-cellular blastoderm embryos, there are regions of the

chromosome that already show evidence of pairing (Fung et al., 1998). In fact, transvection

between *gypsy* insulator-containing transgenes has been observed in live imaging studies of

nuclear cycle 14 embryos (Lim et al., 2018). Since the NC14 nuclei are just emerging from an

earlier mitosis and a round of DNA synthesis, one would imagine that other key features of

chromosome organization might also be in the process of being established. Thus, it should be

possible to capture evidence of intermediates like those expected to be generated as the extruding

cohesin complexes move through the *eve* TAD and other nearby loci at this particular stage of

development.

Shown in Fig. 2A is a blow-up of the TAD organization of the *eve* locus. As expected, the

left and right boundaries of the *eve* TAD correspond to *nhomie* and *homie*, respectively, and

sequences containing these two boundaries converge at the apex of the *eve* interaction triangle.

All of the sequences within the TAD appear to come in contact with each other and define a high

density of internal contact domain (HDIC). These contacts are expected to be generated, at least

in part, by the sliding of the chromatin fiber within the TAD against itself, much like sequences

in circular/supercoiled DNA can interact with each other (Fig. 1E). There are also sequences

within the *eve* TAD that show enhanced interactions. For example, there is a darker interaction

triangle connecting *nhomie* and the promoter region, while another triangle connects *homie* (and

the PRE adjacent to *homie*) to the promoter region. These enhanced internal contacts would be

expected to help insulate the *eve* gene from external regulatory elements and at the same time

facilitate interactions between *eve* and its upstream and downstream enhancers.

As predicted by both the loop-extrusion and boundary-pairing models, the *eve* TAD is a

stem-loop, not a circle-loop (Fig. 2B). This is evident from the prominent "volcano plume" at

the apex of the *eve* TAD (volcano triangle). A volcano plume is observed because sequences

flanking *eve* are brought into close proximity when *homie* and *nhomie* are linked to each other by

either cohesin or by head-to-tail pairing (see Fig.1: see also Ke et al.). Fig. Supplemental 1

shows the hierarchy of interactions between TADs to either side of *eve* (L-M, K-M and J-M in

Sup. Fig. 1). This conclusion is supported by previous transgene studies. As first described in

the boundary bypass experiments of Cai and Shen (Cai and Shen, 2001) and Muravyova et al.

(Muravyova *et al.*, 2001), the formation of a stem-loop structure when two boundaries are linked

together brings sequences immediately beyond the linked boundaries into close proximity with

each other. In this assay, a stem-loop structure is required, as it enables enhancers flanking the

"upstream" boundary to interact with reporters flanking the downstream boundary (Kyrchanova

et al., 2008a). In contrast, a circle-loop configuration does not bring the neighbors to either side

of *eve* into close proximity.

While a stem-loop is the topology expected for the *eve* TAD in both models, one feature of

the TAD is inconsistent with the loop-extrusion model. According to this model, loops (TADs)

are generated by a cohesin complex that forms a small bubble at a loading site somewhere in the

TAD-to-be and then extrudes both strands of the growing loop at an equal rate until it bumps into

the boundary road-blocks (Davidson and Peters, 2021; Ghosh and Meyer, 2021; Mirny and

Dekker, 2022; Perea-Resa *et al.*, 2021). If the loading site were located in the middle of the

TAD-to-be, and the extrusion intermediates in a population of nuclei were captured by

crosslinking, one should observe a crosslink-generated stripe that begins at the loading site and

extends perpendicular to the chromosomal DNA until it reaches the apex that links the two TAD

boundaries with imaginary lines of 45° (Fig. Supplemental 2B). This is the pattern that would be

generated by a "broken zipper" when it transiently links the zipper teeth equidistant from the

initial "loading site". If the loading site is off center to the left, the crosslinking of the passing

cohesin complex in a population of nuclei will generate a perpendicular stripe until cohesin come

to a halt when it encounters the roadblock on the left (Fig. Supplemental 2C). Assuming that the

complex continues to extrude the right strand, it will then generate a stripe at 45° degrees that

extends to the apex of the TAD. If the loading site corresponds to one of the CTCF roadblocks

or is located close to the roadblock, the cohesin complex will generate a 45° stripe that comes to

a halt at the apex of the triangle when it encounters the neighboring CTCF roadblock (Fig.

Supplemental 2D and E). If there are multiple loading sites within the TAD, there should be a

series of perpendicular stripes that generate and sequentially reinforce the stripe(s) at 45° (Fig.

Supplemental 2F). However, in spite of the fact that NC14 fly embryos should provide by far

the best prospect of actually capturing loop extrusion intermediates, there are no vertical stripes

in the *eve* TAD, nor are there stripes at 45° that extend to the apex of the *eve* TAD. Nor do we

observe a population of perpendicular stripes that convert into a series of reinforced 45° stripes.

There is also no evidence of stripes that are initially tilted to the left or right of a "loading site",

as might be observed if the rate of extrusion were unequal on the two strands, and then

subsequently convert to a 45° stripe as cohesin comes to a halt at the first boundary encountered.

TADs in the eve environs

The attP insertion site at –142 kb is near the end of the first exon of the *hebe* transcription

unit (Fig. 2C). It is located just within a ~14 kb TAD, TB, that has a high density of internal

contacts, and extends from the *hebe* promoter to the 3' end of the neighboring *dila* gene. In

between *hebe* and the *eve* locus there at least are ten distinct TADs, TC-TL (Fig. 2C). The

endpoints of most of these TADs correspond to sequences that are associated with one or more

known chromosomal architectural factors (dots in Fig. 2C).

These TADs correspond to the fundamental building blocks for organizing the 3D

architecture of this chromosomal segment. Superimposed upon these TADs are regions that

exhibit lower density of contacts (LDC). For example, the *mef2* gene spans two TADs, TJ and

TK. TK contains the *mef2* distal promoter region and extends to the internal *mef2* promoter,

while TJ extends to near the promoter of the divergently transcribed *Etf-QO* gene (blue arrow).

Both of these TADs are linked together by a rectangular LDC domain J-K (see Fig.

Supplemental 2). Likewise, TJ is linked to its immediate neighbor TI by the LDC domain I-J. In

the next level, TADs separated by a single TAD are linked together. In the region immediately

above TJ, the LDC domain I-K links TI to TK (Fig. Supplemental 2). Similarly, TJ is linked to

TL by J-L while TJ is also linked to the large TAD TM (which contains *TER94* and the

neighboring gene, *pka-R2*) on the other side of *eve* by J-M. The same pattern of a hierarchical

series of LDC domains linking TAD neighbors, TAD next-door neighbors and next-next-door

neighbors is observed in the DNA segment that includes the *hebe* gene.

--cohesin-mediated loop extrusion: Like *eve*, the TADs in the region between *eve* and the attP

site in the *hebe* locus are defined by right angle triangles (volcanos) with a high density of

internal contacts. However, unlike *eve,* these volcano triangles do not have a plume, and instead

are surrounded by a series of LDC rectangular clouds. This is not the contact pattern that one

might expect for either a series of stem-loops connected by unanchored loops or a series of stem-

loops connected to each other. Moreover, there are no perpendicular or angled crosslinking

stripes (broken zippers) emanating from the base of these triangles, nor are their stripes along

their 45° legs. Thus, in spite of the fact that the NC14 nuclei used in this analysis are just

emerging from an earlier mitosis and a round of DNA synthesis and should be in the process of

assembling TADs, the crosslinking signatures that should be generated by the embrace of

passing cohesin complexes are completely absent.

In the loop extrusion model, the LDC domains would arise in a subset of nuclei because the

extruding cohesin complex breaks through one or more boundary roadblocks before its

progression is halted (Fig. Supplemental 3A versus B, C and D) (Hsieh et al., 2020; Krietenstein

et al., 2020). Breakthroughs could occur at only one roadblock on one side of the extruding loop

(Fig. Supplemental 3B and C), or at multiple roadblocks on one or both sides (Fig. Supplemental

3D), giving rise to a set of overlying LDC domains like that evident in the region between *hebe*

and *eve*. However, while there would have to be multiple breakthrough events to account for the

hierarchical array of LDC domains observed in these NC14 embryos, there is no evidence of the

45° crosslinking stripes that would be expected to mark the legs of the LDC domains following

the breakthrough. For example, the I-J LDC domain (Fig. Supplemental 2) could be generated

by cohesin complexes that initiated extrusion in either TI or TJ and then failed to halt at the

TI:TJ boundary (green arrowhead). In either case, there should be a 45° stripe of crosslinking

that marks (at least part of) the left (TI) and/or the right (TJ) leg of the I-J LDC domain and

extends to the apex of the TI-TJ LDC triangle; however, there is no evidence of such stripes.

Nor is there evidence of crosslinking stripes marking one or both legs of the J-K or I-K LDC

domains (Fig. Supplemental 2).

--boundary:boundary pairing: At least two different mechanisms are expected to account for the

TADs and LDC domains in the region between *eve* and the attP insertion site at -142 kb. First,

since none of the TADs in this DNA segment are topped by volcano plumes, they could

correspond to circle-loops. As circle-loops are generated by head-to-head pairing, the

boundaries in the region between *nhomie* and *hebe* would be expected to pair with each other

head-to-head. (The TAD to TAD contact maps generated by circle-loops and stem-loops are

considered further in the accompanying paper, Ke *et al.*)

As illustrated in Fig. Supplemental 3E-G, the coiled circle-loop TADs, though topologically

independent, are expected to be in relatively close proximity to each other. This means that in

addition to crosslinking events within the TADs, there will be crosslinking events between

neighboring TADs. These crosslinking events could generate the LDCs seen in Fig. 2C and Fig.

Supplemental 2. Since TADs next to each other (Fig. Supplemental 3E and F; Fig. Supplemental

2 I-J or J-K) would typically be expected to interact more frequently than TADs separated by

one, two or more TADs (Fig. Supplemental 3G; Fig. Supplemental 2 I-K or J-L), there should be

a progressive reduction in contact frequency at each step. This is generally what is seen (see Fig.

Supplemental 2). While connected stem-loops generated by cohesin extrusion should also bump

into one another, contacts between next-door neighbors might be expected to be less frequent

than contacts between next-next-door neighbors (Fig. 1B, C and Fig. Supplemental 3A). (TAD-

to-TAD crosslinking and the impact of loop topology is further examined in Ke et al., 2023.)

The second mechanism would be switching and/or combining pairing partners (Fig.

Supplemental 3H-J). Though imaging studies suggest that pairing interactions can be of

relatively long duration (>30 min: Chen *et al.*, 2018; Vazquez *et al.*, 2006)), they are not

permanent, and other nearby boundaries can compete for pairing interactions (Gohl *et al.*, 2011).

If switching/combining occurs in NC14 embryos (see Fig. Supplemental 3H-J), then the precise

pattern of TADs and LDC domains could also be impacted by the relative avidity of potential

pairing partners in the neighborhood and also the distances separating potential partners. For

example, the rectangle linking TH and TI (H-I in Fig. Supplemental 2) could represent partner

switching so that the boundary between TI and TJ (green arrowhead in Fig. Supplemental 2)

sometimes pairs with the boundary separating TH and TI (blue arrowhead) and sometimes with

the boundary separating TG and TH (red arrowhead in Fig. Supplemental 1). Distant boundary

elements could also preferentially interact with each other, generating a supra-TAD that contains

multiple TAD domains. For example, at the apex of the LDC domain B-I there is an interaction

dot (purple arrow, Fig. Supplemental 2) that links the TA-TB boundary to the TI-TJ boundary

(green arrowhead).

Activation of a distant reporter by the eve enhancers

In previous studies we found that a minimal 367 bp *homie* fragment can orchestrate

regulatory interactions between the *eve* enhancers and reporters inserted at an attP site in the first

intron of the *hebe* transcription unit, 142 kb upstream of the *eve* gene (Fujioka *et al.*, 2016). This

minimal fragment lacks the single *homie* CTCF site, but has sites for several other generic

boundary proteins including BEAF, Zipic, Su(Hw), Pita and Ibf2. In blastoderm stage embryos,

lacZ reporter expression was observed in a 7-stripe pattern that coincided with the stripes of the

endogenous gene. However, at this stage only a subset of *eve*-expressing nuclei expressed the

reporter. Later in development, during mid-embryogenesis, reporter expression was observed in

cells in the dorsal mesoderm, the ventral CNS, and in the anal plate region, in a pattern that

recapitulated *eve* gene expression (Fujioka *et al.*, 2016). At these later stages, most of the *eve*-

expressing cells also appear to express the reporter. The *eve*-dependent activation of the reporter

in a stripe pattern has also been visualized in live imaging of pre-cellular blastoderm-stage

embryos (Chen *et al.*, 2018). Within the 7 *eve* stripes, the majority of the nuclei (~80-85%) did

not express the *lacZ* reporter. In these nuclei, the transgene was found to be at a considerable

average physical distance from the *eve* locus (~700 nM). In the remaining nuclei, the transgene

was located in closer proximity to the *eve* locus (~350 nM, on average). Within this subset, the

transgene was expressed in most of the nuclei. Moreover, in those cases in which the transgene

was active, it remained in close proximity and continued to express *lacZ* for the duration of the

experiment (~30 min).

In the two models for TAD formation, quite different mechanisms must be invoked to

account for activation of the reporter at –142 kb by the *eve* enhancers. In the boundary-pairing

model, the transgene *homie* boundary at –142 kb loops over the intervening TADs and pairs with

the *nhomie:homie* complex flanking the *eve* TAD (c.f., Fig. 3B and C). In the loop-extrusion

model, a cohesin complex initiating loop extrusion in the *eve* TAD must break through the

nhomie roadblock at the upstream end of the *eve* TAD. It must then make its way past the

boundaries that separate *eve* from the *attP* site in the *hebe* gene, and come to a halt at the *homie*

boundary associated with the *lacZ* reporter. This would generate a novel TAD, *eveMammoth*

(*eveMa*), that extends from the *eve homie* all the way to the *homie* fragment at –142 kb, and

encompasses both the reporter and the *eve* gene, including its enhancers (c.f., Fig. 3D). Of

course, the *eveMa* TAD could also be generated by a cohesin complex that initiated in, for

example, TF. However, in this case, the runaway cohesin complex would have to break through

the intervening boundary roadblocks in both directions. In both the boundary-pairing and loop-

extrusion models, the configuration of the chromatin fiber would lead to the activation of *lacZ*

expression by the *eve* enhancers, while the reporter would be protected from the *hebe* enhancers

by the *homie* boundary.

To test these two models, we used a transgene that has two divergently transcribed

reporters, *lacZ* and *gfp*, that are each under the control of an *eve* promoter (Fig. 3A). Two

different versions of the transgene were generated. In the first, *LhomieG*, the *eve-lacZ* reporter is

located "upstream" of *homie* (with respect to the 5'→3'orientation of *homie* in the *eve* locus),

while *eve-gfp* is located downstream (Fig. 3A: top). In the second, the orientation of *homie*

within the transgene is reversed, so that *gfp* is upstream of *homie*, while *lacZ* is downstream

(*GhomieL*) (Fig. 3A: bottom). These two transgenes were then individually inserted into the *attP*

site at –142 kb, in either the Z5 or Z3 orientation. In the Z5 orientation, the *eve-lacZ* reporter is

on the *eve* side of the *homie* boundary, so that it is closer to the *eve* locus (Fig. 3A). The *eve-gfp*

reporter is on the opposite side of the *homie* boundary and is farther away from the *eve* locus.

This places the *eve-gfp* reporter next to a series of *hebe* enhancers located farther upstream, in

the intron of the *hebe* gene, and thus it should be subject to regulation by these enhancers (Fig.

2C). In the Z3 orientation, the entire transgene is inserted in the opposite orientation so that *eve-*

gfp is on the *eve* side of *homie*, while *eve-lacZ* is next to the *hebe* enhancers (Fig. 6A). As a

control, we also generated a Z5 transgene in which *lambda* DNA instead of *homie* was inserted

between the *eve-gfp* and *eve-lacZ* reporters.

LhomieG Z5 and GhomieL Z5

The results for transgenes *LhomieG Z5* and the control *LlambdaG Z5* are most

straightforward and will be considered first. In both the loop-extrusion and boundary-pairing

models, the *LlambdaG Z5* control is not expected to display any regulatory interactions with the

eve locus. Fig. 4C shows that this is the case: the *eve-lacZ* and *eve-gfp* reporters in *LlambdaG Z5*

embryos are silent at the blastoderm stage, and transcripts are only very rarely detected. Later in

development, both reporters are expressed in a repeating pattern along the dorsal midline (Fig.

6E). This expression is driven by the *hebe* gene enhancers located just beyond the *gfp*-reporter

(Fujioka *et al.*, 2009).

In the case of the *LhomieG Z5* insert, regulatory interactions between the transgene and the

eve locus will be observed if a stem-loop is established that links the transgene to *eve*. In the

loop-extrusion model, the breakthrough cohesin complexes would generate a stem-loop,

eveMammoth (*eveMa*), by coming to a halt at the *homie* boundary in the transgene and the *homie*

boundary in *eve*, as illustrated in Fig. 3D. This will place the *lacZ* reporter in close proximity to

the *eve* enhancers. It will also disrupt the *eve* TAD, as *nhomie* is no longer linked to the *homie*

boundary by cohesin. In the boundary-pairing model, the *lacZ* reporter in the transgene is

"upstream" of *homie* just like the *eve* gene and its enhancers are "upstream" of *homie* in the *eve*

locus. Since *homie* pairs with itself head-to-head, pairing between *homie* in the transgene and

homie in the *eve* locus will generate the same stem-loop as that generated by loop-extrusion.

However, because *homie* pairs with *nhomie* head-to-tail, a more complex structure like that in

Fig. 3B would be expected if the pairing interactions with the transgene do not disrupt the *eve*

TAD. Consistent with a tripartite structure of the sort shown here, previous studies have shown

that three *Mcp* transgenes on three different chromosomes can interact with each other

genetically (Muller *et al.*, 1999). Subsequent direct visualization of *Mcp*-mediated pairing

interactions in imaginal discs by Vazquez et al. (Vazquez *et al.*, 2006) showed that four *Mcp*-

containing transgene separated by Mbs (at Bridges cytogenetic intervals ~65 on the left arm of

chromosome 3, and ~83 and ~95 on the right arm of chromosome 3) and/or located on different

homologs, clustered in the same nuclear foci in 94% of the nuclei.

As predicted from the formation of a stem-loop linking the transgene to the *eve* locus, *lacZ*

is expressed in blastoderm-stage embryos in 7 stripes that coincide with the 7 stripes of the

endogenous *eve* gene. However, unlike *eve*, not all nuclei in the seven stripes express *lacZ*

(compare with the *eve* control in Fig. 4A). Later, during mid-embryogenesis, *lacZ* is expressed

in the mesoderm, the CNS, and the anal plate in a pattern that mimics the endogenous *eve* gene.

At this stage almost all *eve* positive cells are also positive for *lacZ*. Unlike the *LlambdaG Z5*

transgene, the *hebe* enhancers do not drive *eve-lacZ* expression, as they are beyond the *homie*

boundary (Fig. 4E). The *eve-gfp* reporter in the *LhomieG Z5* transgene responds differently. As

was observed for the *LlambdaG Z5* control, the *eve* enhancers drive little if any *gfp* expression,

either at the blastoderm stage or later in development (Fig. 5C, E). Instead, *gfp* is expressed in

the midline during mid-embryogenesis under the control of the *hebe* enhancers (Fig. 5E), which

should be in the same TAD as *eve-gfp*. These results would be consistent with both the loop-

extrusion (*eveMa*) and boundary-pairing models (Fig. 3B and D).

GhomieL Z5

The situation is more complicated for the *GhomieL Z5* transgene. This transgene is

inserted in the same chromosomal orientation as *LhomieG Z5*: *eve-lacZ* is on the *eve* side of the

homie boundary and *eve-gfp* is on the *hebe* enhancer side of the boundary. However, the

orientation of the boundary within the transgene is switched so that *eve-gfp* rather than *eve-lacZ*

is "upstream" of *homie*. In the boundary-pairing model, flipping the orientation of *homie* in the

transgene but keeping the same orientation of the transgene in the chromosome will have two

consequences. First, as illustrated in Fig. 3C, the topology of the chromatin loop connecting the

homie boundary in the transgene and the *eve* locus will change from a stem-loop to a circle-loop.

The reason for the change in loop topology is that *homie* pairs with *homie* head-to-head, and with

nhomie head-to-tail (Fujioka *et al.*, 2016). As before, the transgene *homie* will pair with *eve*

homie head-to-head to generate a simple circle-loop; however, if, as expected, it also pairs with

nhomie so that the *eve* TAD remains intact, a more complicated loop structure would be

generated (Fig. 3C). Second, the reporter that is preferentially activated by the *eve* enhancers

will switch from *eve-lacZ* to *eve-gfp*. In order to be activated by the *eve* enhancer, the reporter

must be "upstream" of the transgene *homie*. This physical relationship means that the *eve-gfp*

reporter will be activated by the *eve* enhancers independent of the orientation of the transgene in

the chromosome. On the other hand, the *eve-lacZ* reporter will not be activated by the *eve*

enhancers. In addition, it will still be insulated from the *hebe* enhancers by the *homie* boundary.

It is not possible to form a circle-loop in the loop extrusion model. Instead, the

breakthrough cohesin complex will come to a halt as before when it encounters the inverted

homie boundary at –142 kb (Fig. 3D). This means that the *eve-lacZ* reporter will be in the same

TAD (*eveMa*) as the *eve* enhancers and will be activated by the *eve* enhancers. In contrast, the

eve-gfp reporter will be in the neighboring TAD and will not be activated by the *eve* enhancers.

It will, however, be regulated by the nearby *hebe* enhancers.

Fig. 5 shows that the four predictions of the boundary-pairing model are correct: a) the *eve*

enhancers drive *eve-gfp* expression, b) the *hebe* enhancers drive *eve-gfp* expression, c) *eve-lacZ*

is not subject to regulation by the *eve* enhancers, and d) *eve-lacZ* is insulated from the *hebe*

enhancers. This would also mean that pairing of the transgene *homie* in the *GhomieL Z5* insert

with *homie* (and *nhomie*) in the *eve* locus generates a circle-loop, not a stem-loop, as was the

case for *LhomieG Z5*. In contrast, since loop-extrusion cannot generate circle-loops, the key

predictions of this model, namely that *eve-gfp* will be silent while *eve-lacZ* will be regulated by

the *eve* enhancers, are not satisfied. Instead, *eve-gfp* is activated by the *eve* enhancers, while *eve-*

lacZ is not.

While the results described above for the *GhomieL Z5* transgene are inconsistent with a

"simple" loop-extrusion model, there is one potential caveat: the 5'→3' orientation of the *homie*

boundary is inverted so that it is "pointing" towards the *eve* locus rather than away from it like

the endogenous *homie* boundary. In this orientation, it is possible that *homie* no longer functions

as an effective roadblock, and instead the cohesin complex emanating from the *eve* locus (or

from a loading site somewhere in between) slips past this *homie* fragment and continues until it

is blocked by a properly oriented boundary further upstream. In this case the newly formed

stem-loop, *eveGargantuan* (*eveGa*) (Fig. 3E), would include the *eve-gfp* reporter, and it would

then be activated by the *eve* enhancers. However, there are two problems with postulating this

novel *eveGa* TAD. First, *eve-lacZ* would be in the same *eveGa* TAD as *eve-gfp*, and it should

also be activated by the *eve* enhancers. Second, since the *homie* boundary was bypassed by the

cohesin complex to form the larger *eveGa* TAD, it should not be able to block the *hebe*

enhancers from activating *eve-lacZ*. Neither of these predictions is correct.

GhomieL Z3 and *LhomieG Z3*

To confirm these conclusions, we examined the expression patterns of the *GhomieL* and

LhomieG transgenes when they are inserted in the opposite orientation (Z3) in the –142 kb attP

site. In the boundary-pairing model, the orientation of *homie* in the transgene determines which

reporter will be preferentially activated, while the orientation of the transgene *homie* in the

chromosome determines whether a stem-loop or a circle loop will be formed. In *GhomieL Z3* a

stem-loop will be formed, and *eve-gfp* will be activated by the *eve* enhancers; however, since

eve-lacZ is adjacent to the *hebe* enhancers, they will activate it rather than the *eve-gfp* reporter

(Fig. 6B). In *LhomieG Z3* the topology of the loop will switch to a circle-loop, and *eve-lacZ* will

be activated by both the *eve* and *hebe* enhancers (Fig. 6C). In the first version of the loop-

extrusion model, the *eve-gfp* reporter in both *GhomieL Z3* (Fig. 6D) and *LhomieG Z3* (not

shown), should be activated by the *eve* enhancers, since it is included in the *eveMa* TAD when

the transgenes are inserted in the Z3 orientation, while *eve-lacZ* will be regulated by the *hebe*

enhancers (Fig. 6D). In the second version of the loop-extrusion model, the functioning of the

homie roadblock depends upon its orientation relative to endogenous *homie*. In this model,

GhomieL Z3 is expected to form the *eveMa* TAD (Fig. 6D), while *LhomieG Z3* will form the

larger *eveGa* TAD (Fig. 6E).

As would be predicted by all three models, the *eve* enhancers activate the *eve-gfp* reporter in

the *GhomieL Z3* insert (Fig. 6F and G). Likewise, *homie* blocks the *hebe* enhancers from

activating the *eve-gfp* reporter, while they turn on *eve-lacZ* expression instead. As predicted by

the boundary-pairing model, *eve*-dependent expression of the reporters switches from *gfp* to *lacZ*

in the *LhomieG Z3* insert (Fig. 6F and G). This would not be expected in the first version of the

loop-extrusion model, but it is predicted in the second version. However, as indicated in the

diagram in Fig. 6E, the two reporters, along with the *hebe* enhancers, are included in the larger

eveGa TAD, and for this reason both reporters should be activated equally by both the *eve* and

hebe enhancers. This is not the case. The *hebe* enhancers are blocked by the intervening *homie*

boundary, while there is little or no *eve*-dependent *gfp* expression.

TAD formation by transgenes inserted at –142 kb

We used MicroC to examine the TAD organization of the genomic region extending from

upstream of the attP site through the *eve* locus in 12-16 hr embryos carrying different transgene

insertions. At this stage, *eve* is expressed in only a small number of cells in the CNS, mesoderm

and anal plate, and this is also true for *hebe*. This means that most of the interactions detected by

MicroC are in cells in which the *eve* gene and the two reporters are inactive. In addition, while

the transgene co-localizes with *eve* in less than about a fifth of blastoderm stage nuclei, the

frequency of physical contact is expected to be much higher in 12-16 hr embryos (c.f., Vazquez

et al. Vazquez *et al.*, 2006)). This is because at this stage of development, cell cycles are much

longer, and homologs are fully paired. The key findings are summarized below.

LlambdaG Z5: While there is no evidence of contact between sequences in *eve* and

sequences around the attP site at –142 kb in wild-type embryos, this not true for embryos

carrying the dual reporter transgene. Perhaps the most unexpected finding is a weak, but clearly

discernable pattern of interaction linking the *LlambdaG Z5* transgene to the *eve* locus (Fig. 7A).

However, since the *LlambdaG Z5* transgene does not respond to the *eve* enhancers, this physical

interaction is not sufficient to drive detectable expression of the reporters. Though our

experiments do not allow us to unequivocally identify which sequences in the transgene are

responsible for this long-distance interaction, the most likely candidates are the two *eve*

promoters in the *LlambdaG Z5* transgene. As noted above, interaction triangles are observed

within the *eve* TAD spanning the regions between *nhomie* and *homie* (and/or the *eve* 3' PRE) and

the *eve* promoter. It seems possible that the factors supporting these intra-TAD interactions are

also responsible for connecting *LlambdaG Z5* to the *eve* locus. In this respect it is interesting to

note that *homie* and *nhomie* stimulate enhancer-promoter interactions in a transgene assay

(Yokoshi et al., 2020).

521   *LhomieG Z5*: Unlike *LlambdaG Z5*, the *eve-lacZ* reporter in the *LhomieG Z5* insert is

activated by the *eve* enhancers, and this regulatory interaction is perfectly paralleled by the

strong pattern of crosslinking between the transgene and the *eve* locus (Fig. 7B). On the *eve*

side, the interactions between the transgene and the *eve* locus generate a heavily populated band

of crosslinked sequences that span the 16 kb *eve* TAD. On the transgene side, there is an

unequal distribution of crosslinked sequences to either side of the transgene *homie* boundary. As

shown in the inset, the heaviest density of crosslinked sequences is on the *eve-lacZ* side of the

transgene, consistent with the activation of this reporter by *eve* enhancers. To confirm that the

interaction bias mimics the differences in activity of the two reporters in cells in which the

reporters are expressed, we analyzed the interaction profiles from viewpoints within either the

eve-lacZ or *eve-gfp* genes. Fig. 7D shows that *eve-lacZ* interacts with sequences extending

across the *eve* TAD, while *eve-gfp* interactions are largely restricted to the *eve homie* element.

While these results would be seemingly be consistent with both the boundary-pairing and loop-

extrusion models, there is no evidence of cohesin breaking through multiple intervening

boundaries into order to establish the novel *eve(Mam)* TAD. Instead, the organization of the

TADs and LDC domains in the interval in between the transgene and *eve* (Fig. 7B) resemble that

seen in wild-type NC14 embryos (see Fig. 2) (with the caveat that there are significantly fewer

reads in the transgene experiment). Equally important, the same is true for the *eve* TAD. In

order to generate the *eve(Mam)* TAD, the cohesin complex must break through the *nhomie*

boundary and, in so doing, disrupt the *eve* TAD. However, there are no obvious alterations in

either the *eve* volcano or plume with respect to their structure or relative contact density

compared to the *LlambdaG Z5* and wild type NC14 embryos. Likewise, there is no evidence of

novel vertical and 45° crosslinking stripes that would reflect the boundary breakthrough events

that are expected in the loop-extrusion model to accompany the formation of a physical

connection between the transgene and the *eve* locus.

We next used the transgene *homie* as the viewpoint to test the predictions of the

boundary-pairing and loop-extrusion models. Since *homie* can pair with itself (head-to-head)

and with *nhomie* (head-to-tail) in transvection assays (Fujioka *et al.*, 2016), a prediction of the

boundary-pairing model is that there will be physical interactions between the transgene *homie*

and both *nhomie* and *homie* in the *eve* locus (c.f., Vazquez et al., Vazquez *et al.*, 2006)). Also

based on what is known about how fly boundaries pair with each other (Chetverina *et al.*, 2014;

Chetverina *et al.*, 2017; Kyrchanova *et al.*, 2008a), self-interactions are expected to be stronger

than heterologous interactions. In contrast, in the loop-extrusion model, the cohesin complex

forms the *eveMa* TAD by linking *homie* in the transgene to *homie* in the *eve* locus, and this

requires the cohesin complex to break through *nhomie* and disrupt the connections between

nhomie and *homie* in the *eve* locus. In this case, *transgene:homie*←→*eve:homie* interactions

should be observed, while *transgene:homie*←→*-eve:nhomie* should not. Fig. 8A shows that, as

predicted by the boundary-pairing model, the transgene *homie* interacts with both *homie* and

nhomie in the *eve* locus, and the cross-linking frequency is greater for *homie:homie* interactions.

Since the *eve* TAD does not appear to be disturbed by the presence of the *LhomieG Z5*

transgene, these findings would suggest that the transgene *homie* likely pairs with both *nhomie*

and *homie* in a pattern somewhat like that illustrated in Fig. 3B. While the interactions with the

transgene are inconsistent with the loop-extrusion model, one could make the *ad hoc* assumption

that runaway cohesin complexes form not one but two different novel TADs. One would be

eveMa. In the other, *eveElephant* (*eveEl*), a cohesin complex that initiates in, for example, the

TE TAD would come to a halt on one side at the *homie* boundary in the transgene, and on the

other side at the *nhomie* boundary in the *eve* locus (Fig. Supplemental 4). However, this

configuration would not explain how the transgene reporter is activated by *eve* enhancers. Since

the *eve* enhancers in *eveEl* are located in a different TAD than the TADs containing *eve-lacZ* and

eve-gfp, they would not be able to activate either reporter.

GhomieL Z5: When *homie* is reversed in the *GhomieL Z5* transgene, the *eve-gfp* reporter

responds to the *eve* enhancers instead of *eve-lacZ*. In the boundary-pairing model, *eve-gfp* is

activated because the self- and heterologous-pairing interactions between *nhomie* and *homie* are

orientation-dependent. As a result, the topology of the loop formed between the transgene and

eve switches from a stem-loop to a circle-loop, and this brings the *eve-gfp* reporter in close

proximity to the *eve* enhancers instead of the *eve-lacZ* reporter. Since circle-loops cannot be

formed in the loop-extrusion model, activation of the distal *eve-gfp* reporter would require the

formation of a larger *eveGa* TAD in order to encompass the reporters in the transgene and the

eve enhancers in the same looped domain (Fig. 3E). However, as shown in Fig. 7C, we do not

observe the cross-linking pattern expected for a new *eveGa* TAD that includes the entire

transgene at –142 kb and then extends to an even more distal boundary. Nor are there any novel

vertical/45° crosslinking stripes or obvious perturbations in the basic TAD organization in the

region beyond –142 kb, or between –142 kb and the *eve* locus, as might be expected for cohesin

complexes breaking through multiple intervening roadblocks. Instead, as was the case for

LhomieG Z5, the *GhomieL Z5* transgene is physically linked to the *eve* locus, and there is a

prominent band of crosslinked transgene sequences that spans the *eve* TAD (Fig. 7C).

Consistent with the expression patterns of the two reporters, the heaviest density of crosslinked

sequences is on the *eve-gfp* side of the transgene (see inset). This is confirmed by the interaction

profiles generated using either the *eve-gfp* or *eve-lacZ* gene body as the viewpoint: the most

frequent contacts are between sequences within *eve* and the *eve-gfp* reporter (Fig. 7D). In

addition, as was the case for *LhomieG Z5*, when the transgene *homie* is used as the viewpoint,

interactions are observed between both *homie* and *nhomie* in the *eve* locus, with the most

frequent corresponding to self-pairing interactions (Fig. 8B). Other than postulating the *ad hoc*

eveEl TAD (Fig. Supplemental 4), this would not be expected for either version of the loop-

extrusion model.

LhomieG Z3 and *GhomieL Z3*: To confirm the findings for transgenes inserted in the Z5

orientation, we used MicroC to analyze the TAD organization of *LhomieG Z3* and *GhomieL Z3*.

(Sup. Figs. 5 and 6). Like their Z5 counterparts, both show a strong band of interaction between

the transgene and sequences spanning the *eve* locus. In the case of *GhomieL Z3*, which is

predicted to form a stem-loop, bringing *eve-gfp* in contact with the *eve* enhancers, the *eve-gfp*

sequences interact more frequently with the *eve locus* (see inset in Fig. Supplemental 5A).

homie is in the opposite orientation in the transgene and in the chromosome in *LhomieG Z3*, and

in the boundary pairing model is expected to generate a circle-loop. While the sample

preparation for MicroC was of poorer quality than the others, the inset in Sup. Fig.6A shows that,

as expected, *eve-lacZ* interacts more frequently with sequences in the *eve* locus than *eve-gfp*. On

the other hand, there is no evidence of a larger *evaGa* TAD even though *eve-lacZ* is activated by

the *eve* enhancers. These interactions are confirmed when either *eve-lacZ* or *eve-GFP* are used

as viewpoints to plot the physical contacts between the *eve* TAD and the two reporters in

LhomieG Z3 and *GhomieL Z3* (Sup. Figs. 5B and 6B). In addition, as was the case for the two

Z5 transgenes, when the transgene *homie* is used as the viewpoint, interactions are observed

between *homie* in the transgene and *both homie* and *nhomie* in the *eve* locus (Fig. 8A and B).

Again, the most frequent interactions are self-pairing interactions between *homie* in the transgene

and the *eve homie*.

Loop topology and interactions with neighbors: The results in the previous sections demonstrate

that the topology of the loop formed between the transgene and the *eve* locus depends upon the

orientation of *homie* in the transgene *and* the orientation of the transgene in the chromosome.

When *homie* in the transgene is pointing away from the *eve* locus, the topology is a stem-loop.

When *homie* in the transgene is pointing towards the *eve* locus, the topology is a circle-loop.

Changing the topology also impacts how sequences flanking the transgene and flanking *eve*

interact with each other. With the important caveat that the depth of our MicroC sequencing is

limited, the interaction profiles reflect the topology of the transgene-induced loop. For *homie*

pointing away from *eve* (*LhomieG Z5* and *GhomieL Z3*), sequences to the *TER94* side of *eve* and

to the *hebe* enhancer side of the transgene are more frequently crosslinked with each other (Fig.

Supplemental 7). These interactions would be expected for a stem-loop structure (Fig. 3B). For

homie pointing towards *eve*, sequences on the *TER94* side of *eve* interact with sequences on the

eve side of the transgene. This would be expected for a circle-loop structure (Fig. 3C).

Discussion

Two different mechanisms have been proposed to explain how TADs are formed and their

endpoints determined. In the first, a cohesin complex first induces a small loop at a loading site

within the TAD-to-be and then progressively extrudes one or both strands until the complex

encounters boundary roadblocks. If the loading site is located asymmetrically within the TAD-

to-be, extrusion will continue on the unblocked strand until a roadblock is encountered. The

endpoints of the loops were initially thought to be determined by the location of the first two

roadblocks encountered by the cohesin complex (Alipour and Marko, 2012). In a refinement of

this model, the orientation of a given roadblock relative to the direction of movement of the

cohesin complex determines whether it will come to a halt (Davidson and Peters, 2021;

Fudenberg *et al.*, 2016; Guo *et al.*, 2012; Guo *et al.*, 2015; Sanborn *et al.*, 2015). It has also been

suggested that cohesin complexes can break through a roadblock and come to a halt at more

distant roadblocks, so that in a population of cells analyzed by Hi-C or MicroC, the basic units of

organization, the TADs, are overlaid by a hierarchy of LDC domains (Hsieh *et al.*, 2020;

Krietenstein *et al.*, 2020). However, the only loop topologies that are possible in the loop-

extrusion model are a stem-loop and an unanchored loop.

In the other mechanism, TAD formation is governed by boundary:boundary pairing

interactions. In this case, the critical factors are the pairing and orientation preferences of

potential partners, together with proximity. Since boundary pairing interactions often exhibit an

orientation preference, the topology of the loop formed by paired boundaries can be either a

stem-loop or a circle-loop, depending on their relative orientation in the chromosome. In this

respect it differs from the loop-extrusion model in that switching the orientation of the boundary

does not preclude the formation of a TAD, whereas it can in some versions of the loop-extrusion

model.

In the studies reported here, we have used MicroC to analyze the TAD organization of a

~150 kb chromosomal segment flanking the *eve* locus, and have experimentally tested the

predictions of these two models for the mechanisms of TAD formation. Our studies do not

support a loop-extrusion mechanism, and are inconsistent with this model on multiple levels.

Manipulating "TAD" formation: In our experimental paradigm, a transgene containing two

divergently transcribed reporters and the *eve* boundary *homie* is inserted 142 kb upstream of the

eve TAD. Depending upon the orientation of the *homie* boundary within the transgene, either

eve-gfp or *eve-lacZ* is activated by developmentally regulated enhancers in the *eve* TAD. Since

there are a series of well-defined TADs and their associated boundary elements separating the

transgene from *eve*, a version of the loop-extrusion model in which the cohesin complex

invariably arrests upon encountering *nhomie* and *homie* at the ends of the *eve* TAD cannot

account for this long-distance regulation. Instead, the extruding cohesin complex must be able to

break through the boundaries located between *homie* in the transgene and *homie* in the *eve* TAD.

This would generate a novel TAD, *eveMa*, with endpoints corresponding to the two *homie*

boundaries (Fig. 3D). In this case, the reporter on the *eve* side of the transgene *homie* would be

in the same domain as the *eve* enhancers, and thus potentially subject to regulation. While

eveMa could account for reporter expression in inserts in which the transgene *homie* is pointing

away from *eve* (and also away from the reporter on the *eve* side of the transgene insert), it does

not explain reporter expression when the transgene *homie* is pointing towards *eve* (and in this

case towards the reporter on the *eve* side of the transgene insert). In the latter case, the reporter

activated by the *eve* enhancers lies outside of the *eveMa* (cf., Fig. 5D), and thus is not included in

the same TAD as the *eve* enhancers. In order to account for activation of the distal reporter, one

must imagine that the cohesin complex breaks through the *homie* boundary in the transgene and

continues on until extrusion is halted by an even more distal boundary. In this revision of the

loop-extrusion model, both reporters would be included in the same TAD, *eveGa* (c.f., Fig. 5E),

as the *eve* enhancers, and both reporters should be activated. However, this is not observed.

TAD organization: loop-extrusion versus boundary pairing: Analysis of the TAD organization

in the region spanning the transgene insertion site and *eve* also does not fit with the expectations

of the loop-extrusion model. To account for reporter activation, the cohesin complex must be

able to break through the *nhomie* boundary. This breakthrough could be engineered by a cohesin

complex that initiated loop extrusion either from within the *eve* TAD or from one of the TADs

located between the *nhomie* boundary and the transgene at –142 kb. However, in the absence of

the transgene, there is no indication that *nhomie* is particularly prone to break-through events. In

fact, the *eve* TAD differs from most of the TADs in between *eve* and the transgene in that it is

incorporated into larger LDC interaction triangles much less frequently. There is also no reason

to imagine that the presence of a *homie* (or *nhomie*: see Fujioka *et al.*, 2016)) transgene at –142

kb would somehow induce *nhomie* break-through events.

As expected from the regulatory interactions, *homie*-containing transgenes at –142 kb

generate a strong band of crosslinking events that physically link the transgene to the *eve* locus.

In the loop-extrusion model, this strong band of interaction should generate a novel interaction

triangle (LDC or HDIC) with endpoints corresponding to the *homie* elements in the transgene

and *eve*. Contrary to this expectation, a triangle of increased interaction frequencies spanning the

entire region between –142 kb and *homie* is not observed for any of the transgene inserts. In

viewpoints from the *homie* element in the *LhomieG Z5* and *GhomieL Z3* transgenes, there are

two peaks in the *eve* TAD. The more prominent peak maps to *homie*, while the less prominent

peak maps to *nhomie*. To account for these interaction peaks, the loop-extrusion model would

have to assume that the presence of the *LhomieG Z5* and *GhomieL Z3* transgenes induces the

formation of two novel TADs, the *eveMa* (*homie:homie*) TAD in Fig. 5 and another TAD,

eveElephant, that links the transgene *homie* to the *nhomie* element in the *eve* locus (Fig.

Supplemental 4). However, while the *eveMa* TAD could potentially explain the activation of

reporters on the proximal side of the transgene *homie* by the *eve* enhancers, these TADs do not

explain how the *eve* enhancers are able to activate reporters on the distal (*hebe* enhancer) side of

the *GhomieL Z5* and *LhomieG Z3* transgene. This would require the formation of yet another

TAD, *eveGa*, that encompasses *eve*, the transgene, and one or more TADs beyond the *hebe* gene

(Fig. 3E). However, there is no indication of physical interactions between the *homie* boundary

in *eve* and a boundary located beyond the *hebe* gene. Moreover, like *eveMa*, *eveGa* would also

have to be accompanied by the formation of *eveEl*, since the *homie* element in *GhomieL Z5* and

LhomieG Z3 interacts with both *nhomie* and *homie*.

While these results are inconsistent with the expectations of the loop-extrusion model,

they dovetail nicely with many of the predictions of the boundary-pairing model. First, in all

four transgenes, *homie* interacts with both *homie* and *nhomie* in the *eve* locus, but does not

interact with the multiple boundary elements located in between. This is consistent with other

studies which show that boundaries often have strong partner preferences (Gohl *et al.*, 2011;

Kyrchanova *et al.*, 2011; Kyrchanova *et al.*, 2008b). Second, interaction frequencies are greater

for self-pairing than for heterologous pairing. This fits with what is known about how partner

preferences are determined (Blanton et al., 2003; Chetverina *et al.*, 2017; Erokhin et al., 2021;

Kyrchanova *et al.*, 2008a; Zolotarev *et al.*, 2016). Third, the pattern of reporter activation in the

four transgenes is precisely as expected from previous studies on the orientation dependence of

homie pairing with itself (head-to-head) and with *nhomie* (head-to-tail) (Chen *et al.*, 2018;

Fujioka *et al.*, 2016). This orientation dependence means that the reporter located "upstream" of

homie is the one that is activated, independent of the orientation of the transgene in the

chromosome. Fourth, the pattern of physical interactions between the *eve* locus and the

transgenes recapitulates the pattern of reporter activation. That is, when *eve-lacZ* sequences are

preferentially crosslinked to sequences in the *eve* locus, then the *eve-lacZ* reporter is activated by

the *eve* enhancers. Likewise, *eve-gfp* is preferentially crosslinked to sequences in *eve* when it is

activated by the *eve* enhancers. Again this is dependent on the orientation of *homie* in the

transgene, but not on the orientation of the transgene in the chromosome. Fifth, unlike the loop-

extrusion model where only stem-loops or disconnected loops can be generated, both stem-loops

and circle-loops can be, and, as we have shown here, actually are generated by

boundary:boundary pairing. The topology of the loop formed between the transgene and *eve*

depends on the orientation of the transgene *homie* in the chromosome. When *homie* in the

transgene insert is pointing away from the *eve* locus, a stem-loop is generated. When *homie* in

the transgene insert is pointing towards the *eve* locus, a circle-loop is generated. That both loop

topologies can be generated by boundary:boundary pairing is confirmed by not only by the

pattern of expression of the two reporters in the four transgene inserts but also by their pattern of

physical interactions evident in our MicroC experiments.

Are the pairing interactions of homie and nhomie unusual? There are two reasons to think that

the properties of most fly boundaries are similar to those of *homie* and *nhomie*. First, all of the

fly boundaries that have been tested in assays that are expected to require pairing do have this

activity (Blanton *et al.*, 2003; Cai and Shen, 2001; Erokhin et al., 2011; Gohl *et al.*, 2011;

Kyrchanova *et al.*, 2008a; Kyrchanova *et al.*, 2011; Kyrchanova et al., 2007; Kyrchanova *et al.*,

2008b; Muravyova *et al.*, 2001). Second, multimerized binding sites for generic polydactyl zinc

finger proteins like Su(Hw), Pita, dCTCF, Zw5, and Zipic that are found associated with many

fly boundaries are not only able to block enhancer/silencer action but can also mediate pairing

interactions in insulator bypass and transvection assays (Erokhin *et al.*, 2021; Kyrchanova *et al.*,

2008a; Zolotarev *et al.*, 2016). Moreover, as is the case for *homie* and *nhomie*, pairing

interactions require the appropriate partners: multimerized sites for Zw5 do not pair with

multimerized sites for dCTCF (Erokhin *et al.*, 2021; Kyrchanova *et al.*, 2008a; Zolotarev *et al.*,

2016). This makes good sense, since these proteins form dimers/multimers, as do other fly

boundary factors like BEAF, Mod(mdg4), the BEN-domain protein Insensitive and the Elba

complex (Avva and Hart, 2016; Bonchuk et al., 2021; Bonchuk et al., 2011; Bonchuk et al.,

2020; Fedotova et al., 2018; Fedotova et al., 2019; Fedotova et al., 2017; Hart et al., 1997). The

ability to multimerize means that boundaries that share binding sites for the same protein, i.e., for

Pita, Zipic or CTCF, can be physically linked to each other by a single protein complex.

Boundary pairing orientation and loop topology: Our analysis of the TAD organization in the

eve neighborhood suggests that the predominant loop topology may be a circle-loop, not a stem-

loop. Only the *eve* TAD (and perhaps the very small TAD, TA) is clearly a stem-loop. It is

topped by a distinctive volcano plume and is not overlaid by a hierarchy of LDC domains,

indicating that it must be "insulated" from crosslinkable interactions with other nearby TADs.

By contrast, none of the TADs in the region spanning *eve* and the *hebe* gene have volcano

plumes at their apex. Instead, all of these TADs are overlaid by a hierarchal organization of

LDC domains (clouds) that diminish in crosslinking intensity as the number of in-between TADs

increases. As discussed above, the contact pattern of the TADs to the left of *eve* are likely to

arise from partner switching and/or head-to-head boundary pairing interactions that generate

circle-loops. Pairing orientation versus loop topology is considered further in the accompanying

paper (Ke *et al.*).

Orientation and long-distance regulatory interactions: Our experiments highlight an important

constraint on long-distance regulatory interactions that are mediated not only by boundary

elements like *homie* and *nhomie* but also by the GAF-dependent tethering and bypass elements

(Batut et al., 2022; Kyrchanova et al., 2023; Kyrchanova et al., 2018; Kyrchanova et al., 2019a;

Kyrchanova et al., 2019b; Levo et al., 2022; Li et al., 2023; Mohana et al., 2023), namely the

orientation of the physical interactions that bring distant enhancers (or silencers) into close

proximity with their gene targets. As we have shown here, the orientation of the physical

interactions that bring distant sequences close together determines whether the enhancer will be

able to activate its target gene. In principle the effects of orientation could be mitigated by

increasing the distance between the boundary/tethering elements and either the enhancers

(silencers) or the promoter. However, distance can be a critical if not a rate limiting step in

transcriptional activation. For example, live imaging experiments have shown that the frequency

and amplitude of transcriptional bursts depend on the distance between the enhancer and target

promoter. Nearly continuous bursting and high amplitude transcription was observed when the

enhancer was located close the promoter (Fukaya et al., 2016). However, at distances as little as

6 kb, transcription became discontinuous with discrete bursts, while at 9 kb, the bursts were

much less frequent, and the amplitude/output was substantially reduced (Yokoshi *et al.*, 2020).

If similar distance-dependent reductions in transcriptional efficiency are common, then it would

not be possible to overcome the topological constraints on regulatory interactions by simply

increasing the distance between the boundary/tethering elements and the enhancers/promoters.

While our studies do not directly address how chromosome architecture impacts regulatory

interactions in vertebrates, it seems clear that long-distance regulatory interactions in vertebrates

(c.f., Tan et al., 2023) will be subject to the same types of topological constraints as those that

apply in flies.

Acknowledgements: We would like to thank the Gordon Grey for running the fly food

facility at Princeton, members of the Lewis Sigler Genomics Core facility for their invaluable

assistance with DNA sequencing and Sergey Ryabichko for his help with smFISH probes. We

would also like to thank members of MOL431 for creative input. Special thanks to Olga

Kyrchanova, Daria Chetverina, Maksim Erokhin, Pavel Georigev, Tsutomu Aoki, Girish

Deshpande, Airat Ibragimov, Sergey Ryabichko, Yuri Pritykin and Alex Ostrin for stimulating

discussions and sharing unpublished data.

  Materials and methods

804 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
n-Heptane	Fisher Chemical	O3008-4
Paraformaldehyde 20% solution, EM Grade	Electron Microscopy Sciences	15713-S
Formaldehyde, 16%, methanol free, Ultra Pure	Polysciences Inc	18814-10
PBS - Phosphate-Buffered Saline (10X) pH 7.4, RNase-free	Thermal Fisher	AM9624
Tween 20	Sigma	P1379
Triton X-100	Bio-Rad	161-0407
Tris base	Sigma	11814273001
Methanol	Fisher Chemical	203403
SSC, 20X	Thermal Fisher	15557044
Formamide	Thermal Fisher	17899
Dextran sulfate	Sigma	D8906
Salmon Sperm DNA	Thermal Fisher	AM9680
Ribonucleoside Vanadyl Complex	NEB	S1402S
Nuclease-free BSA	Sigma	126609
Triethylammonium Acetate	Sigma	625718
dGTP (100 MM)	VWR	76510-208
dTTP (100 MM)	VWR	76510-224
Lonza NuSieve 3:1 Agarose	Thermal Fisher	BMA50090
T4 DNA ligase	NEB	M0202L
Biotin-11-dCTP	Jen Bioscience	NU-809-BIOX

Biotin-14-dATP	Jen Bioscience	NU-835-BIO14
Qubit dsDNA HS Assay Kit	Life Technologies Corporation	Q32851
Atto 633 NHS ester	Sigma	01464
Phase Lock Gel™, QuantaBio - 2302830, Phase Lock Gel Heavy	VMR	10847-802
NEBNext Ultra II DNA Library Prep Kit for Illumina	NEB	E7645S
Ampure Xp 5ml Kit	Thermal Fisher	NC9959336
Hifi Hotstart Ready Mix	Thermal Fisher	501965217
Dynabeads MyOne Streptavidin C1	Life Technologies Corporation	65001
cOmplete™, EDTA-free Protease Inhibitor Cocktail	Sigma	11873580001
N,N-Dimethylformamide	Sigma	227056
Potassium acetate solution	Sigma	95843
DSG (disuccinimidyl glutarate)	Thermal Fisher	PI20593
T4 Polynucleotide Kinase - 500 units	NEB	M0201S
DNA Polymerase I, Large (Klenow) Fragment - 1000 units	NEB	M0210L
End-it DNA End Repair Kit	Thermal Fisher	NC0105678
Proteinase K recomb. 100 mg	Sigma	3115879001
Nuclease Micrococcal (s7)	Thermal Fisher	NC9391488
EGS (ethylene glycol bis(succinimidyl succinate))	Thermal Fisher	PI21565
Atto 565 NHS ester	Sigma	72464
Oligonucleotides		
WK_eve_probeset Sequence see "Supplemental Data S1"	Biosearch Technologies	

WK_LacZ_probeset Sequence see "Supplemental Data S1"	Biosearch Technologies	
WK_GFP_probeset Sequence see "Supplemental Data S1"	Biosearch Technologies	
NEBNext Multiplex Oligos for Illumina	NEB	E7335S
Software and algorithms		
Fiji (ImageJ)	Schindelin et al., 2012	fiji.sc
NIS element	Nikon	microscope.healthca re.nikon.com/product s/software/nis-elements
Graphpad Prism 8	GraphPad Software	graphpad.com
HiGlass	Kerpedjiev et al., 2018	higlass.io
bwa	Li H. and Durbin R., 2009	bio-bwa.sourceforge.net/
samtools	Github/open source	samtools.github.io
pairsamtools	Github/open source	github.com/mirnylab/ pairsamtools
pairix	Github/open source	github.com/4dn-dcic/pairix
cooler	Abdennur, N., and Mirny, L., 2019	github.com/mirnylab/ cooler
Miniconda	Anaconda	docs.conda.io/en/lat est/miniconda.html
Snakemake	Github/open source	snakemake.github.io

Plasmid construction and transgenic lines

The dual reporters construct was described previously (Fujioka *et al.*, 2016). In short, both reporters contain the *eve* basal promoter (-275 to +106 bp relative to *eve* start site), the *lacZ* (*eve-lacZ*) or *EGFP* (*eve-GFP*) coding region, and the *eve* 3' UTR (+1300 to +1525 bp). These two reporters are divergently transcribed. The 367 bp *homie* fragment, or a 500 bp fragment from *lambda* phage DNA, was placed between the two reporters. The *homie* fragment was placed in both orientations: in *LhomieG*, the *eve-lacZ* reporter is located upstream of the *homie* fragment, while in *GhomieL*, *eve-GFP* is located upstream of the *homie* fragment (see Fig 2A).

The -142 kb attP landing site was described previously (Fujioka *et al.*, 2009). The -142 kb site contains two attP target sites for phiC31 recombinase-mediated cassette exchange (RMCE) (Bateman et al., 2006) and *mini-white* as a marker. RMCE can result in either orientation of an inserting transgene (Z5 and Z3; see text). RMCE events were identified by loss of *mini-white*, and the orientation of each insert was determined by PCR.

smFISH probe preparation

The sequences of target genes were obtained from Flybase (flybase.org)(Gramates et al., 2022). To design probes, the target gene sequences were submitted to the Biosearch Technologies Stellaris RNA FISH probe designer tool (biosearchtech.com), with parameters set to 20nt probes, 48 probes per gene, and mask level 5 for *Drosophila melanogaster*. The sequence of the probes used are listed in Supplemental Data S1. The probes with 3' MdC(TEG-Amino) modifications were then ordered from Biosearch Technologies in 96-well plate format, cartridge purification and salt free conditions and in 5 nmol scale. To prepare the fluorophore for probe labeling, the Sigma Atto NHS ester fluorophore (Atto 565 NHS or Atto 633 NHS) were dissolved in dimethylformamide at 5mg/mL as the stock solution. For probe labeling, 1 nmol of each probe was combined and mixed with Atto NHS ester fluorophore at 1mg/mL in 500μL of

0.1M sodium bicarbonate (pH 8.0-8.5) for 2 hours, and the probes were then precipitated with

0.5M potassium acetate (pH 5.2) in 75% EtOH. After that, the probes were dried and resuspend

in 50μL of DePC-treated H_2O. From there, the probes were injected into HPLC in which the

percentage 0.1M triethylammonium acetate varied from 7% to 30% with a flow rate of 1mL/min.

The detector to monitor coupled probes during the HPLC purification were set to 260nm for

DNA and the absorption wavelength of the coupled fluorophore. The coupled probes were

collected when both 260nm and the absorption wavelength channel were detected. The coupled

probes were dried and diluted into appropriate concentration in DePC-treated H_2O for smFISH

experiments.

smFISH

smFISH methods were adopted and modified from previous studies (Little and Gregor,

2018; Trcek et al., 2017). 100-200 flies were placed in a cage with an apple juice plate at the

bottom of the cage. Early stages of embryos were for 7hours, while for later stages embryos,

collection was overnight. Embryos from each plates were washed into collection mesh and

dechorionated in bleach for 2min, then fixed in 5mL of 4% paraformaldehyde in 1X PBS and

5mL of heptane for 15min with horizontal shaking. The paraformaldehyde were then removed

and replaced with 5mL methanol. The embryos were then devitellinized by vortexing for 30s,

and washed in 1mL of methanol twice. Methanol were then removed and replaced by PTw(1X

PBS with 0.1% Tween-20) through serial dilution as 7:3, 1:1, and 3:7 methanol:PTw. The

embryos were washed twice in 1mL of PTw and then twice in 1mL smFISH wash buffer (4X

SSC, 35% formamide and 0.1% Tween-20). Embryos were incubated with ~5nM coupled

smFISH probes in hybridization buffer (0.1g/mL dextran sulfate, 0.1mg/mL salmon sperm

ssDNA, 2mM ribonucleoside vanadyl complex, 20μg/mL RNAse-free BSA, 4X SSC, 1%

Tween-20 and 35% formamide) for 12-16h. Embryos were then washed twice for 2 hours in

1mL smFISH wash buffer, followed by 4X washing in 1mL PTw. For DAPI/Hoechst staining,

the embryos were stained with 1ug/mL DAPI or Hoechst in PTw, and washed 3X with 1mL

PTw. Finally, the embryos were mounted on microscope slides with Aqua PolyMount and #1.5

coverslip for imaging.

Imaging, image analysis and statistics

Embryos from smFISH were imaged by using a Nikon A1 confocal microscope system,

Plan Apo 20X/0.75 DIC objective. Z-stack images were taken with interval of 2μm, 4X average,

1024X1024 resolution, and the appropriate laser power and gain were set for 405, 561, and 640

channels to avoid overexposure. Images were processed by ImageJ and the maximum projection

was applied to each of the stack images. To measure the stripe intensity of early embryos, multi-

channel images were first split into single channels and the stripe signal was highlighted and

selected by the MaxEntropy thresholding method. For APR intensity measurement, the ROI tool

was used to crop out the APR region of late-stage embryos. The cells with APR signal were also

highlighted and selected by MaxEntropy thresholding. The particle measurement tool was used

to measure the average intensity of all cells that have the signal. At the same time, the

background signal (average intensity) was measured on cells without a signal in the same

embryo. The relative intensity (signal to background) for each embryo was calculated by using

the stripe signal and the background from the same embryo. To make comparisons between

independent biological replicates, the average background signal of all embryos from each

replicate was calculated. The relative intensity of each embryo from each replicate were

normalized based on the average background signal of all embryos from same replicates.

GraphPad Prism was used for data visualization and statistical analysis. To compare intensity

from embryos in different groups, different signals from the same embryo (e.g., *lacZ* and *GFP*)

were paired and paired two-tailed t-tests were used to calculate p-values. All raw measurements

and normalized data are included in Supplemental Data S2.

MicroC library construction

Embryos were collected on yeasted apple juice plates in population cages for 4 hours,

incubated for 12 hours at 25°C, then subjected to fixation as follows. Embryos were

dechorionated for 2 mins in 3% sodium hypochlorite, rinsed with deionized water, and

transferred to glass vials containing 5 mL PBST (0.1% Triton-X100 in PBS), 7.5 mL n-heptane,

and 1.5mL fresh 16% formaldehyde. Crosslinking was carried out at room temperature for

exactly 15 mins on an orbital shaker at 250rpm, followed by addition of 3.7 mL 2M Tris-HCl

pH7.5 and shaking for 5 mins to quench the reaction. Embryos were washed twice with 15 mL

PBST and subjected to secondary crosslinking. Secondary crosslinking was done in 10mL of

freshly prepared 3mM final DSG and ESG in PBST for 45 mins at room temperature with

passive mixing. The reaction was quenched by addition of 3.7mL 2M Tris-HCl pH7.5 for 5

mins, washed twice with PBST, snap-frozen, and stored at -80°C until library construction.

Micro-C libraries were prepared as previously described (Batut *et al.*, 2022) with the following

modifications: 50uL of 12-16hr embryos were used for each biological replicate. 60U of MNase

was used for each reaction to digest chromatin to a mononucleosome:dinucleosome ratio of 4.

Libraries were barcoded, pooled and subjected to paired-end sequencing on an Illumina Novaseq

S1 100 nt Flowcell (read length 50 bases per mate, 6-base index read).

Micro-C data processing

MicroC data for *D. melanogaster* were aligned to custom genomes edited from the

Berkeley Drosophila Genome Project (BDGP) Release 6 reference assembly (dos Santos et al.,

2015) with BWA-MEM (Li and Durbin, 2009) using parameters **-S -P -5 -M**. Briefly, the

custom genomes are simply insertions of the transgenic sequence into the –142kb integration

site, as predicted from perfect integration. These events were confirmed using PCR post-

integration. The resultant BAM files were parsed, sorted, de-duplicated, filtered, and split with

Pairtools (https://github.com/mirnylab/pairtools). We removed pairs where only half of the pair

could be mapped, or where the MAPQ score was less than three. The resultant files were

indexed with Pairix (https://github.com/4dn-dcic/pairix). The files from replicates were merged

with Pairtools before generating 100bp contact matrices using Cooler (Abdennur and Mirny,

2020). Finally, balancing and Mcool file generation was performed with Cooler's Zoomify tool.

Virtual 4C profiles were extracted from individual replicates using FAN-C (Kruse et al.,

2020) at 400bp resolution. The values were summed across replicates and smoothed across three

bins (1.2kb). Viewpoints were determined based on the most informative region for

interpretation. Ultimately we decided to use 800bp regions in the gene body of either *GFP* or

lacZ, moving downstream from the *eve* promoter. However, similar results were obtained using

viewpoints between transgenic *homie* and the promoters of either gene. For *homie* viewpoints,

due to transgenic *homie* being masked, some loss of signal is expected at the viewpoint (Figure

6).

Figure Legends

Figure 1. Diagram of the possible loop topologies generated by loop-extrusion and boundary:boundary pairing. A) Cohesin embraces a loop at a loading site somewhere within the TAD-to-be and then extrudes a loop at an equal rate on both strands. Extrusion continues until cohesin encounters boundary roadblocks on both strands. B) In one model, the orientation of the roadblock is important. As a consequence, the chromatin fiber will be organized into a series of stem-loops separated from each other by unanchored loops. C) If the presence of a boundary roadblock, but not its orientation, is sufficient to halt extrusion, the chromatin fiber will be organized into a series of linked stem-loops. The main axis of the chromosome will be defined by the cohesin:CTCF roadblocks. D) The pairing of two boundaries head-to-tail generates a stem-loop. E) If boundary pairing interactions are strictly pairwise, head-to-tail pairing will generate a series of stem-loops separated from each other by unanchored loops. F) If boundaries can engage in multiple head-to-tail pairing interaction, the chromosome will be organized into a series of linked stem-loops. The main axis of the chromosome will be defined by a series paired boundaries. G) The pairing of two boundaries head-to-head generates a circle-loop. H) If boundary pairing interactions are strictly pairwise there will be a series of circle-loops separated from each other by unanchored loops. I) If boundaries can engage in multiple head-to-head pairing interactions, the chromatin fiber will be organized into a series of circle-loos connected to each other at their base. Pairing interactions between boundaries #1 and #2 need not be in register with pairing of boundaries #2 and #3. In this case, the main axis of the chromosome may bend and twist and this could impact the relative orientation of the circle-loops.

Figure 2. TAD organization of the *even-skipped* locus and neighboring sequences. A) The *eve* skipped TAD is a volcano with a plume that is anchored by *nhomie* and *homie*. ChIP-seq data below the MicroC map indicate many of the known fly chromosomal architectural proteins proteins are associated with the two *eve* boundary elements *in vivo* (Cuartero et al., 2014; Duan et al., 2021; Gaskill et al., 2021; Li et al., 2015; Sun et al., 2015; Ueberschär et al., 2019; Zolotarev *et al.*, 2016). B) The *eve* locus forms a stem-loop structure. In this illustration *nhomie* pairs with *homie* head-to-tail, and this configuration forms a stem-loop which brings sequences upstream *nhomie* and downstream of *homie* in contact as is observed in insulator bypass assays (Kyrchanova *et al.*, 2008a). The *eve* locus is shown assembled into a coiled "30 nM" chromatin fiber. C) MicroC contact pattern for the chromosomal region spanning the attP site at –142 kb and the *eve* locus. Like the *eve* volcano, this contact pattern was generated using aggregated previously published NC14 data (Batut *et al.*, 2022; Levo *et al.*, 2022). A black arrow indicates the –142kb locus where transgenes are integrated into the genome. A blue arrow indicates position of the *Etf-QO* gene. Note the numerous TADs separating the –142kb *hebe* locus from the *eve* "volcano TAD". Individual TADs are labeled TA—TM. Black boxes indicate positions of sequences that, based on ChIP experiments, are bound by one or more known insulator proteins *in vivo*.

Figure 3. Schematics of boundary pairing and loop extrusion. A) *LhomieG Z5* and *GhomieL Z5* transgenes and the *eve* locus on a linear map. The same color codes are used throughout. In the *eve* locus, *nhomie* (blue arrow) and *homie* (red arrow) are oriented, by convention, so that they are pointing towards the right. This convention is maintained in the two transgenes. In *LhomieG Z5 homie* (top) is in the opposite orientation from *homie* in the *eve* locus is pointing away from the *eve* locus. In *GhomieL Z5 homie* (bottom) *homie* is in the same orientation as the

homie in the *eve* locus and is pointing towards the *eve* locus. B) Predicted boundary pairing
interactions between *LhomieG Z5* and the *eve* locus. *homie* in the transgene pairs with *homie* in
the *eve* locus head-to-head. Since *homie* in the transgene is pointing in the opposite orientation
from *homie* in the *eve* locus, a stem-loop will be generated. If *homie* in the transgene also pairs
with *nhomie* in the *eve* locus head-to-tail, a loop structure like that shown in B will be generated.
In this topology, *eve-lacZ* is in close proximity to *eve* enhancers and *eve-gfp* is in contact with
the *hebe* enhancer. C) Predicted boundary pairing between *GhomieL Z5* and the *eve* locus. The
GhomieL Z5 transgene is inserted in the same chromosomal orientation as *LhomieG Z5*;
however, the *homie* boundary is inverted so that it is pointing towards the *eve* in the same
orientation as the *homie* in the *eve* locus. *Homie* in the transgene will pair with *homie* in the *eve*
locus head-to-head, and this generates a circle-loop. If *homie* in the transgene also pairs with
nhomie in the *eve* loucs a loop structure like that shown in C will be generated. In this topology
eve-gfp will be activated by both the *eve* and *hebe* enhancers. D) Loop extrusion model for
LhomieG Z5. Transgene *homie* and endogenous *homie* determine the endpoints of the extruded
eveMammoth (*eveMa*) loop. In this topology, *eve-lacZ* is in close proximity to the *eve*
enhancers, while *eve-gfp* is close to the *hebe* enhancers. E) Loop extrusion model for *GhomieL*
Z5. The cohesin complex bypasses transgenic *homie* and is stopped at an upstream boundary X
to generate a novel *eveGarantuan* (*eveGa*) loop. Both *eve-gfp* and *eve-lacZ* are located within
the same *eveGa* TAD, and thus should interact with *eve*. Since the *hebe* enhancers are within
this TAD as well, they would also be able to activate both reporters.
984
Figure 4. The *lacZ* reporter is activated by *eve* enhancers in the *LhomieG Z5* transgene. A)
Expression patterns of endogenous *eve* in early (stage 4-5) and late (stage 14-16) embryos by
smFISH. *eve* (Atto 633) probes were used to hybridize with *eve* mRNA. *eve* is shown in yellow
and DAPI in blue. N>3. B) Schematics of transgenes. C) Expression patterns of transgenic
reporters in early stages embryos. Top: *LlambdaG Z5*. Bottom: *LhomieG Z5*. *GFP* is in green,
lacZ is in orange, and DAPI is in blue, here and in E. D) Quantification of normalized stripe
signals for transgenes shown in C. The reporter smFISH intensity (signal to background), here
and in F, was measured, normalized, and plotted as described in Methods. N>3. n=27 for
LlambdaG Z5 and n=42 for *LhomieG Z5*. E) Expression patterns of transgenic reporters in late
stages embryos. Top: *LlambdaG Z5*. Bottom: *LhomieG Z5*. F) Quantification of normalized
stripe signals for transgenes shown in E. N>3. n=27 for *LlambdaG Z5* and n=56 for *LhomieG*
Z5. Scale bars = 200μm. N = # independent biological replicates. n = # animals. The paired
two-tailed t-test were used for statistical analysis. *p < 0.05, **p < 0.01, ***p < 0.001, ****p <
0.0001, ns: not significant. Raw measurements are available in the Source Data files.
999
Figure 5. The *GFP* reporter is activated by both *eve* and *hebe* enhancers in the *LhomieL*
Z5 transgene. A) Schematics of transgenes. B) Expression patterns of transgenic reporters in
early stage embryos. Top: *LlambdaG Z5*. Bottom: *GhomieL Z5*. *GFP* is in green, *lacZ* is in
orange, and DAPI is in blue, here and in D. C) Quantification of normalized stripes signal as
represented in B. The reporter smFISH intensity (signal to background) was measured,
normalized, and plotted as described in Methods, here and in E. N>3. n=27 for *LlambdaG Z5*,
and n=46 for *LhomieG Z5*. D) Expression patterns of transgenic reporters in late stage embryos.
Top: *LlambdaG Z5*. Bottom: *GhomieL Z5*. E) Quantification of normalized stripe signals for
transgenes shown in D. N>3. n=27 for *LlambdaG Z5*, and n=59 for *LhomieG Z5*. Scale bars =
200μm. N = # independent biological replicates. n = # animals. The paired two-tailed t-test
were used for statistical analysis. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns: not
significant. Raw measurements are available in the Source Data files.

1012
Figure 6. Expression of reporters in Z3 transgenes. A) Schematics of Z3 transgenes. B)
Boundary pairing for *GhomieL Z3* transgene long-range interactions. In this topology, *GFP* is in
close proximity to *eve* enhancers, and *lacZ* is in contact with the *hebe* enhancer region. C)
Boundary pairing for the *LhomieG Z3* transgene. In this topology, *lacZ* is in close proximity to
both the *eve* enhancers and the *hebe* enhancers. D) Loop extrusion model for *GhomieL Z3*.
Transgenic *homie* and endogenous *homie* flank the *eveMammoth* loop. In this topology, *GFP* is
in close proximity to *eve* enhancers, and *lacZ* is close to the *hebe* enhancers. E) Loop extrusion
model for the *LhomieG Z3* transgene. The cohesin complex bypasses transgenic *homie* and is
stopped at an upstream boundary X to create the *eveGarantuan* loop. Both *GFP* and *lacZ* are at
a similar physical distance from *eve* enhancers and *hebe* enhancers in this topology. F)
Expression patterns of transgenic reporters in early stage embryos. Top: *GhomieL Z3*. Bottom:
LhomieG Z3. *GFP* is in green, *lacZ* is in orange, and DAPI is in blue. N=3, n=24 for both
GhomieL Z3 and *LhomieG Z3*. **G)** Expression patterns of transgenic reporters in late stage
embryos. Top: *GhomieL Z3*. Bottom: *LhomieG Z3*. N=3, n=24 for both *GhomieL Z3* and
LhomieG Z3. Scale bars = 200μm, N = # independent biological replicates, n = # animals.
1028
Figure 7. *homie* in Z5 transgene *LhomieG* and *GhomieL* Z3 mediates long-range
interactions with the *eve* locus *Small dark blue arrows* – insulators upstream and downstream
of *hebe* that appear important to demarcate interaction domains between –142kb and *eve*. *Large*
blue arrow – endogenous *nhomie*. *Large red arrow* – *homie*, either endogenous or in the
transgene. The direction of arrows follow established convention on *nhomie/homie*, and do not
reflect orientation of insulator protein binding motifs *per se*. *Gray boxes* – enhancers. A)
MicroC map of the control line *LlambdaG Z5*. Scaled cartoons of the two loci of interest are
shown directly below the MicroC map, and an unscaled blowup of the elements of interest at
each locus is provided. Within the MicroC map of the entire locus, a zoom-in of the off-diagonal
interaction between the –142kb insertion cassette and the endogenous *eve* locus is shown. Note a
slight increase in interaction frequency (compare to Figure 1A). B, C**)** MicroC map of *LhomieG*
Z5 and *GhomieL Z5*, respectively. The only difference between the two lines is the orientation of
transgenic *homie*, as indicate below each MicroC map. Blow up of the off-diagonal interaction
between –142kb and *eve*, including scaled cartoons denoting features of interest in the top right
corner. Note the changed pattern of interaction with the endogenous *eve* locus due to the
orientation switch. D) "Virtual 4C" maps obtained from MicroC maps of "B" (top panel), and
"C" (bottom panel). Viewpoints are shown in both panels from either the *lacZ* gene (orange) or
the *GFP* gene (green). Note in each cassette the altered interaction of either gene with
endogenous *eve*, depending on the change of orientation of *homie*.
1048
Figure 8. *homie* in the transgene interacts with *eve homie* and *nhomie*. "Virtual 4C" maps
of the *LhomieG* (A) or *GhomieL* (B) transgenes, for the Z5 (blue) and Z3 (orange) orientations.
Viewpoints are taken from the transgenic *homie* sequence. Importantly, note that transgenic
homie preferentially interacts with endogenous *homie* no matter the orientation. Intriguingly, the
orientation switch of transgenic *homie* leads to an apparent change in its degree of preference for
endogenous *homie* over *nhomie*.

1060
1061 **Supplemental Figures:**
1062
1063
Figure Sup. 1. TAD organization of the *even-skipped* locus and neighboring sequences
MicroC contact pattern for the chromosomal region spanning the attP site at –142 kb and the *eve*
locus. Like the *eve* volcano, this contact pattern was generated using aggregated previously
published NC14 data (Batut *et al.*, 2022; Levo *et al.*, 2022). Arrow indicates the –142kb locus
where transgenes are integrated into the genome. LDC domains are labeled according to the
interacting TADs: for example, H-I indicates crosslinking events generated by physical
interactions between sequences in two neighboring TADs, TH and TI, while I-J indicates
interactions between sequences in TI and TJ. Likewise, G-I indicates crosslinking events
generated by physical interactions between sequences TADs TG and TI, which are separated
from each other by TAD TH. Note that unlike the TADs to the left of *eve*, the *eve* TAD does not
interact frequently with its neighbors. Purple arrow indicates potential interaction dot between
the boundary marking the border of TA and TB and the boundary marking the border of TI and
TJ. Red, blue and green arrowheads mark boundaries that could engage in partner switching: see
text.
1078
Figure Sup. 2. Trouble ahead, trouble behind. A) Illustration of a broken zipper—zipping in
front but not closing behind. B) Crosslinking pattern expected for cohesin loading at a site very
close to the middle of the TAD-to-be: The vertical line of linked DNA is generated when the two
chromatin fibers in the TAD-to-be are transiently linked together as cohesin passes. When
cohesin bumps into the CTCF roadblocks (BE) at the boundaries of the TAD, the vertical line
stops. C) Cohesin loading site is offset from center of TAD-to-be to the left. The crosslinking
initiates at the loading site and extends vertically until the cohesin complex bumps into the
CTCF roadblock on the left. At that point extrusion of the left chromatin fiber halts, while it
continues to extrude the chromatin fiber on the right. The line of crosslinked sequences then
extends at a 45º angle until cohesin comes to a halt at the CTCF roadblock on the right. D and
E) The loading site is at or close to the boundary on the left (D) or the right (E). In this case
extrusion will be unidirectional and the DNAs crosslinked as cohesin passes will form a line that
extends at a 45º angle until cohesin comes to a halt at the CTCF roadblock. F). Cohesin is able
to load at multiple sites in the TAD-to-be. In this case there will be a series of vertical lines
whose intensity will be correlated with the relative frequency that a give loading site is used. The
vertical lines will extend in each case until cohesin comes to a halt at the closest CTCF
roadblock. At that point, the line of crosslinked DNA will extend at a 45º angle until that
particular cohesin complex comes to halt at the CTCF roadblock that is farther away. The 45º
line of crosslinked DNA will become progressive more intense as it approaches the apex of the
triangle.
1099
1100
1101
Figure Sup. 3. Models for the formation of LDC domains. Several different mechanisms
could explain the LDC domains. A-D). In the loop extrusion model, the LDC domains arise
because cohesin sometimes breaks through a CTCF roadblock. In A) there are three primary
TADs, TAD1, TAD2, and TAD3. B and C) Occasional breakthrough events can generate a
secondary TAD, LDC1-2 (TAD1+TAD2) or LDC2-3 (TAD2+TAD3). D) Even less frequent
breakthrough events can generate a tripartite TAD (TAD1+TAD2+TAD3) that will correspond

to LDC1-3. E-G). LDC domains arise because TADs sometimes bump into their neighbors. In
E, TAD1 bumps into TAD2 to give LDC1-2, while in F TAD2 bumps into TAD3 to give
LDC2:3. G) Less frequently TAD1 bumps into TAD3 to give LDC1:3. H-J) LDC domains arise
because boundaries can switch partners. H and I) In H boundary A (dark green) skips boundary
B (red) and pairs with boundary C (tan). This generates a new TAD, TAD1+TAD2 which
corresponds to LDC domain 1-2. In I, boundary B (red) skips boundary C (tan) and pairs with D
(blue). This generates a new TAD, TAD2+TAD3 which corresponds to LDC domain 2-3. J) In
J, boundary A (dark green) skips boundary B (red) and boundary C (tan) and pairs with boundary
D (blue). This generates a new TAD, TAD1+TAD2+TAD3 which corresponds to LDC domain
1-3.
1118
1119
Figure Sup. 4. Loop extrusion: *eveElephant*. A) Loop extrusion orientation dependent model
for LhomieG Z5 transgene as the sketched in Figure 2E. This configuration generates *eveMa*. B)
Instead running through *nhomie*, the cohesin complex comes to a halt at the *nhomie* boundary,
generating *eveElephant*. A second cohesin complex generates the *eve* TAD by loading at a site
within the *eve* locus. Both *eveElephant* and the *eve* TAD are stem-loops. However, while this
double stem-loop configuration would explain why the transgene *homie* makes contact with both
nhomie and *homie*, the *eve* enhancers are isolated in their own loop and would not make contact
with the two reporters in the transgene.
1128
Figure Sup. 5. MicroC of *GhomieL Z3*. A) MicroC map of *GhomieL Z3*. Gray lines are
masked regions due to repeated sequences. In this case, focusing on the zoom-in map from top
to bottom, most likely P-element remnants 3' of the *lacZ* gene, transgenic *homie*, and P-element
remnants 3' of the *GFP* gene. Compare directly to *GhomieL Z5* for changes in interaction
preference due to change of *homie* orientation (Figure 4C). B) "Virtual 4C" map from either
lacZ or *GFP* gene body viewpoints.
1135
1136   **Figure Sup. 6. Micro C of *LhomieG Z3*.** A) MicroC map of *LhomieG Z3*. Gray lines are
1137   masked regions due to repeated sequences. In this case, most possibly *P-element* remnants 3' of
1138   the *lacZ* gene. Compare directly to *GhomieL Z3* for changes in interaction preference due to
1139   change of homie orientation (Figure 4B). B) "Virtual 4C" map from either *lacZ* or *GFP* gene
1140   body viewpoints.
1141
1142
1143
Figure Sup. 7. Blowup of interactions generated by the transgene with sequences
neighboring –142 kb and the *eve* TAD. The interaction pattern indicates that the loop topology
changes as a result of the orientation of transgenic *homie*. Blow-up of off-diagonal interactions,
focused on the interaction between the endogenous *eve* locus and the –142kb transgenic insert
with A) *LhomieG Z3,* B) *LhomieG Z5,* C) *GhomieL Z3,* or D) *GhomieL Z5*. Note the switch in
preferential interaction between regions adjacent to transgenic *homie* and endogenous *homie*
(indicated by arrow brackets). In cassettes where transgenic *homie* is in the same orientation in
the genome as endogenous *homie* (A and D), regions upstream of transgenic *homie* preferentially
interact with regions downstream of endogenous *homie*. In contrast, when transgenic *homie* is in
the opposite orientation in the genome as endogenous *homie* (B and C), regions downstream of
transgenic *homie* preferentially interact with regions downstream of endogenous *homie*. This

suggests a different loop topology when the orientation of the transgene *homie* element in the
chromosome is flipped.
1157
1158
1159

<u>REFERENCES</u>

Abdennur, N., and Mirny, L.A. (2020). Cooler: scalable storage for Hi-C data and other
genomically labeled arrays. Bioinformatics *36*, 311-316. 10.1093/bioinformatics/btz540.
Alipour, E., and Marko, J.F. (2012). Self-organization of domain structures by DNA-loop-
extruding enzymes. Nucleic Acids Res. *40*, 11202-11212. 10.1093/nar/gks925.
Avva, S.V., and Hart, C.M. (2016). Characterization of the Drosophila BEAF-32A and BEAF-
32B Insulator Proteins. PLoS One *11*, e0162906. 10.1371/journal.pone.0162906.
Bateman, J.R., Lee, A.M., and Wu, C.T. (2006). Site-specific transformation of Drosophila via
phiC31 integrase-mediated cassette exchange. Genetics *173*, 769-777.
Batut, P.J., Bing, X.Y., Sisco, Z., Raimundo, J., Levo, M., and Levine, M.S. (2022). Genome
organization controls transcriptional dynamics during development. Science *375*, 566-570.
10.1126/science.abi7178.
Bell, A.C., West, A.G., and Felsenfeld, G. (2001). Insulators and boundaries: versatile regulatory
elements in the eukaryotic genome. Science *291*, 447-450. 10.1126/science.291.5503.447.
Blanton, J., Gaszner, M., and Schedl, P. (2003). Protein:protein interactions and the pairing of
boundary elements in vivo. Genes Dev. *17*, 664-675.
Bonchuk, A., Boyko, K., Fedotova, A., Nikolaeva, A., Lushchekina, S., Khrustaleva, A., Popov,
V., and Georgiev, P. (2021). Structural basis of diversity and homodimerization specificity of
zinc-finger-associated domains in Drosophila. Nucleic Acids Res. *49*, 2375-2389.
10.1093/nar/gkab061.
Bonchuk, A., Denisov, S., Georgiev, P., and Maksimenko, O. (2011). Drosophila BTB/POZ
domains of "ttk group" can form multimers and selectively interact with each other. J. Mol. Biol.
412, 423-436. 10.1016/j.jmb.2011.07.052.
Bonchuk, A., Kamalyan, S., Mariasina, S., Boyko, K., Popov, V., Maksimenko, O., and
Georgiev, P. (2020). N-terminal domain of the architectural protein CTCF has similar structural
organization and ability to self-association in bilaterian organisms. Sci Rep *10*, 2677.
10.1038/s41598-020-59459-5.
Cai, H.N., and Shen, P. (2001). Effects of cis arrangement of chromatin insulators on enhancer-
blocking activity. Science *291*, 493-495.
Callan, H.G. (1963). THE NATURE OF LAMPBRUSH CHROMOSOMES. Int. Rev. Cytol. *15*,
1-34. 10.1016/s0074-7696(08)61114-6.
Cavalheiro, G.R., Pollex, T., and Furlong, E.E. (2021). To loop or not to loop: what is the role of
TADs in enhancer function and gene regulation? Curr. Opin. Genet. Dev. *67*, 119-129.
10.1016/j.gde.2020.12.015.
Chen, H., Levo, M., Barinov, L., Fujioka, M., Jaynes, J.B., and Gregor, T. (2018). Dynamic
interplay between enhancer–promoter topology and gene activity. Nat. Genet. *50*, 1296.
Chetverina, D., Aoki, T., Erokhin, M., Georgiev, P., and Schedl, P. (2014). Making connections:
insulators organize eukaryotic chromosomes into independent cis-regulatory networks.
Bioessays *36*, 163-172.

Chetverina, D., Fujioka, M., Erokhin, M., Georgiev, P., Jaynes, J.B., and Schedl, P. (2017).
Boundaries of loop domains (insulators): Determinants of chromosome form and function in
multicellular eukaryotes. Bioessays *39*, 1600233.
Collier, J.E., and Nasmyth, K.A. (2022). DNA passes through cohesin's hinge as well as its
Smc3-kleisin interface. Elife *11*. 10.7554/eLife.80310.
Cuartero, S., Fresan, U., Reina, O., Planet, E., and Espinas, M.L. (2014). Ibf1 and Ibf2 are novel
CP190-interacting proteins required for insulator function. EMBO J. *33*, 637-647.
Davidson, I.F., and Peters, J.M. (2021). Genome folding through loop extrusion by SMC
complexes. Nat Rev Mol Cell Biol *22*, 445-464. 10.1038/s41580-021-00349-7.
dos Santos, G., Schroeder, A.J., Goodman, J.L., Strelets, V.B., Crosby, M.A., Thurmond, J.,
Emmert, D.B., Gelbart, W.M., and FlyBase, C. (2015). FlyBase: introduction of the Drosophila
melanogaster Release 6 reference genome assembly and large-scale migration of genome
annotations. Nucleic Acids Res. *43*, D690-697.
Duan, J., Rieder, L., Colonnetta, M.M., Huang, A., McKenney, M., Watters, S., Deshpande, G.,
Jordan, W., Fawzi, N., and Larschan, E. (2021). CLAMP and Zelda function together to promote
Drosophila zygotic genome activation. Elife *10*. 10.7554/eLife.69937.
Erokhin, M., Davydova, A., Kyrchanova, O., Parshikov, A., Georgiev, P., and Chetverina, D.
(2011). Insulators form gene loops by interacting with promoters in Drosophila. Development
138, 4097-4106.
Erokhin, M., Gorbenko, F., Lomaev, D., Mazina, M.Y., Mikhailova, A., Garaev, A.K.,
Parshikov, A., Vorobyeva, N.E., Georgiev, P., Schedl, P., and Chetverina, D. (2021). Boundaries
potentiate polycomb response element-mediated silencing. BMC Biol. *19*, 113. 10.1186/s12915-
021-01047-8.
Fedotova, A., Aoki, T., Rossier, M., Mishra, R.K., Clendinen, C., Kyrchanova, O., Wolle, D.,
Bonchuk, A., Maeda, R.K., and Mutero, A. (2018). The BEN domain protein insensitive binds to
the Fab-7 chromatin boundary to establish proper segmental identity in Drosophila. Genetics
210, 573-585.
Fedotova, A., Clendinen, C., Bonchuk, A., Mogila, V., Aoki, T., Georgiev, P., and Schedl, P.
(2019). Functional dissection of the developmentally restricted BEN domain chromatin boundary
factor Insensitive. Epigenetics & chromatin *12*, 2.
Fedotova, A.A., Bonchuk, A.N., Mogila, V.A., and Georgiev, P.G. (2017). C2H2 Zinc Finger
Proteins: The Largest but Poorly Explored Family of Higher Eukaryotic Transcription Factors.
Acta Naturae *9*, 47-58.
Fudenberg, G., Imakaev, M., Lu, C., Goloborodko, A., Abdennur, N., and Mirny, L.A. (2016).
Formation of Chromosomal Domains by Loop Extrusion. Cell Rep *15*, 2038-2049.
10.1016/j.celrep.2016.04.085.
Fujioka, M., Mistry, H., Schedl, P., and Jaynes, J.B. (2016). Determinants of Chromosome
Architecture: Insulator Pairing in *cis* and in *trans*. PLoS Genet. *12*, e1005889.
Fujioka, M., Sun, G., and Jaynes, J.B. (2013). The Drosophila *eve* insulator Homie promotes *eve*
expression and protects the adjacent gene from repression by Polycomb spreading. PLoS Genet.
9, e1003883.
Fujioka, M., Wu, X., and Jaynes, J.B. (2009). A chromatin insulator mediates transgene homing
and very long-range enhancer-promoter communication. Development *136*, 3077-3087.
Fukaya, T., Lim, B., and Levine, M. (2016). Enhancer control of transcriptional bursting. Cell
166, 358-368.
Fung, J.C., Marshall, W.F., Dernburg, A., Agard, D.A., and Sedat, J.W. (1998). Homologous
chromosome pairing in Drosophila melanogaster proceeds through multiple independent
initiations. J. Cell Biol. *141*, 5-20.

Gaskill, M.M., Gibson, T.J., Larson, E.D., and Harrison, M.M. (2021). GAF is essential for
zygotic genome activation and chromatin accessibility in the early Drosophila embryo. Elife *10*.
10.7554/eLife.66668.
Ghosh, R.P., and Meyer, B.J. (2021). Spatial Organization of Chromatin: Emergence of
Chromatin Structure During Development. Annu. Rev. Cell. Dev. Biol. *37*, 199-232.
10.1146/annurev-cellbio-032321-035734.
Gligoris, T.G., Scheinost, J.C., Bürmann, F., Petela, N., Chan, K.L., Uluocak, P., Beckouët, F.,
Gruber, S., Nasmyth, K., and Löwe, J. (2014). Closing the cohesin ring: structure and function of
its Smc3-kleisin interface. Science *346*, 963-967. 10.1126/science.1256917.
Gohl, D., Aoki, T., Blanton, J., Shanower, G., Kappes, G., and Schedl, P. (2011). Mechanism of
chromosomal boundary action: roadblock, sink, or loop? Genetics *187*, 731-748.
Gramates, L.S., Agapite, J., Attrill, H., Calvi, B.R., Crosby, M.A., Dos Santos, G., Goodman,
J.L., Goutte-Gattat, D., Jenkins, V.K., Kaufman, T., et al. (2022). FlyBase: a guided tour of
highlighted features. Genetics *220*. 10.1093/genetics/iyac035.
Guo, Y., Monahan, K., Wu, H., Gertz, J., Varley, K.E., Li, W., Myers, R.M., Maniatis, T., and
Wu, Q. (2012). CTCF/cohesin-mediated DNA looping is required for protocadherin α promoter
choice. Proc Natl Acad Sci USA *109*, 21081-21086. 10.1073/pnas.1219280110.
Guo, Y., Xu, Q., Canzio, D., Shou, J., Li, J., Gorkin, D.U., Jung, I., Wu, H., Zhai, Y., and Tang,
Y. (2015). CRISPR inversion of CTCF sites alters genome topology and enhancer/promoter
function. Cell *162*, 900-910.
Hart, C.M., Zhao, K., and Laemmli, U.K. (1997). The scs' boundary element: characterization of
boundary element-associated factors. Mol Cell Biol. *17*, 999-1009.
Heger, P., George, R., and Wiehe, T. (2013). Successive gain of insulator proteins in arthropod
evolution. Evolution *67*, 2945-2956.
Heger, P., and Wiehe, T. (2014). New tools in the box: an evolutionary synopsis of chromatin
insulators. Trends Genet. *30*, 161-171.
Hsieh, T.S., Cattoglio, C., Slobodyanyuk, E., Hansen, A.S., Rando, O.J., Tjian, R., and Darzacq,
X. (2020). Resolving the 3D Landscape of Transcription-Linked Mammalian Chromatin
Folding. Mol. Cell *78*, 539-553.e538. 10.1016/j.molcel.2020.03.002.
Jerković, I., Szabo, Q., Bantignies, F., and Cavalli, G. (2020). Higher-Order Chromosomal
Structures Mediate Genome Function. J. Mol. Biol. *432*, 676-681. 10.1016/j.jmb.2019.10.014.
Ke, W., Fujioka, M., Schdle, P., and Jaynes, J.B. Chromosome Structure II: Stem-loops and
circle-loops.
Krietenstein, N., Abraham, S., Venev, S.V., Abdennur, N., Gibcus, J., Hsieh, T.S., Parsi, K.M.,
Yang, L., Maehr, R., Mirny, L.A., et al. (2020). Ultrastructural Details of Mammalian
Chromosome Architecture. Mol. Cell *78*, 554-565.e557. 10.1016/j.molcel.2020.03.003.
Kruse, K., Hug, C.B., and Vaquerizas, J.M. (2020). FAN-C: a feature-rich framework for the
analysis and visualisation of chromosome conformation capture data. Genome Biol *21*, 303.
10.1186/s13059-020-02215-9.
Kyrchanova, O., Chetverina, D., Maksimenko, O., Kullyev, A., and Georgiev, P. (2008a).
Orientation-dependent interaction between Drosophila insulators is a property of this class of
regulatory elements. Nucleic Acids Res. *36*, 7019-7028.
Kyrchanova, O., Ibragimov, A., Postika, N., Georgiev, P., and Schedl, P. (2023). Boundary
bypass activity in the abdominal-B region of the Drosophila bithorax complex is position
dependent and regulated. Open Biol *13*, 230035. 10.1098/rsob.230035.
Kyrchanova, O., Ivlieva, T., Toshchakov, S., Parshikov, A., Maksimenko, O., and Georgiev, P.
(2011). Selective interactions of boundaries with upstream region of Abd-B promoter in

Drosophila bithorax complex and role of dCTCF in this process. Nucleic Acids Res. *39*, 3042-
3052.
Kyrchanova, O., Kurbidaeva, A., Sabirov, M., Postika, N., Wolle, D., Aoki, T., Maksimenko, O.,
Mogila, V., Schedl, P., and Georgiev, P. (2018). The bithorax complex iab-7 Polycomb response
element has a novel role in the functioning of the Fab-7 chromatin boundary. PLoS Genet. *14*,
e1007442.
Kyrchanova, O., Sabirov, M., Mogila, V., Kurbidaeva, A., Postika, N., Maksimenko, O., Schedl,
P., and Georgiev, P. (2019a). Complete reconstitution of bypass and blocking functions in a
minimal artificial Fab-7 insulator from Drosophila bithorax complex. Proceedings of the
National Academy of Sciences, 201907190.
Kyrchanova, O., Toshchakov, S., Parshikov, A., and Georgiev, P. (2007). Study of the functional
interaction between Mcp insulators from the Drosophila bithorax complex: effects of insulator
pairing on enhancer-promoter communication. Mol. Cell. Biol. *27*, 3035-3043.
Kyrchanova, O., Toshchakov, S., Podstreshnaya, Y., Parshikov, A., and Georgiev, P. (2008b).
Functional interaction between the Fab-7 and Fab-8 boundaries and the upstream promoter
region in the Drosophila Abd-B gene. Mol. Cell. Biol. *28*, 4188-4195.
Kyrchanova, O., Wolle, D., Sabirov, M., Kurbidaeva, A., Aoki, T., Maksimenko, O.,
Kyrchanova, M., Georgiev, P., and Schedl, P. (2019b). Distinct Elements Confer the Blocking
and Bypass Functions of the Bithorax Fab-8 Boundary. Genetics, genetics. 302694.302019.
Levo, M., Raimundo, J., Bing, X.Y., Sisco, Z., Batut, P.J., Ryabichko, S., Gregor, T., and
Levine, M.S. (2022). Transcriptional coupling of distant regulatory genes in living embryos.
Nature *605*, 754-760. 10.1038/s41586-022-04680-7.
Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics *25*, 1754-1760. 10.1093/bioinformatics/btp324.
Li, H.-B., Muller, M., Bahechar, I.A., Kyrchanova, O., Ohno, K., Georgiev, P., and Pirrotta, V.
(2011). Insulators, not Polycomb response elements, are required for long-range interactions
between Polycomb targets in Drosophila melanogaster. Mol. Cell. Biol. *31*, 616-625.
Li, L., Lyu, X., Hou, C., Takenaka, N., Nguyen, H.Q., Ong, C.T., Cubeñas-Potts, C., Hu, M.,
Lei, E.P., Bosco, G., et al. (2015). Widespread rearrangement of 3D chromatin organization
underlies polycomb-mediated stress-induced silencing. Mol. Cell *58*, 216-231.
10.1016/j.molcel.2015.02.023.
Li, X., Tang, X., Bing, X., Catalano, C., Li, T., Dolsten, G., Wu, C., and Levine, M. (2023).
GAGA-associated factor fosters loop formation in the Drosophila genome. Mol. Cell *83*, 1519-
1526.e1514. 10.1016/j.molcel.2023.03.011.
Lim, B., Heist, T., Levine, M., and Fukaya, T. (2018). Visualization of transvection in living
Drosophila embryos. Mol. Cell *70*, 287-296. e286.
Little, S.C., and Gregor, T. (2018). Single mRNA Molecule Detection in Drosophila. Methods
Mol Biol *1649*, 127-142. 10.1007/978-1-4939-7213-5_8.
Matthews, N.E., and White, R. (2019). Chromatin Architecture in the Fly: Living without
CTCF/Cohesin Loop Extrusion?
Mirny, L., and Dekker, J. (2022). Mechanisms of chromosome folding and nuclear organization:
their interplay and open questions. Cold Spring Harbor Perspectives in Biology *14*, a040147.
Mohana, G., Dorier, J., Li, X., Mouginot, M., Smith, R.C., Malek, H., Leleu, M., Rodriguez, D.,
Khadka, J., Rosa, P., et al. (2023). Chromosome-level organization of the regulatory genome in
the Drosophila nervous system. Cell *186*, 3826-3844.e3826. 10.1016/j.cell.2023.07.008.
Muller, M., Hagstrom, K., Gyurkovics, H., Pirrotta, V., and Schedl, P. (1999). The mcp element
from the Drosophila melanogaster bithorax complex mediates long-distance regulatory
interactions. Genetics *153*, 1333-1356.

Muravyova, E., Golovnin, A., Gracheva, E., Parshikov, A., Belenkaya, T., Pirrotta, V., and
Georgiev, P. (2001). Loss of insulator activity by paired Su(Hw) chromatin insulators. Science
291, 495-498.
Nora, E.P., Caccianini, L., Fudenberg, G., So, K., Kameswaran, V., Nagle, A., Uebersohn, A.,
Hajj, B., Saux, A.L., Coulon, A., et al. (2020). Molecular basis of CTCF binding polarity in
genome folding. Nat Commun *11*, 5612. 10.1038/s41467-020-19283-x.
Nuebler, J., Fudenberg, G., Imakaev, M., Abdennur, N., and Mirny, L.A. (2018). Chromatin
organization by an interplay of loop extrusion and compartmental segregation. Proc Natl Acad
Sci USA *115*, E6697-e6706. 10.1073/pnas.1717730115.
Perea-Resa, C., Wattendorf, L., Marzouk, S., and Blower, M.D. (2021). Cohesin: behind
dynamic genome topology and gene expression reprogramming. Trends Cell Biol. *31*, 760-773.
10.1016/j.tcb.2021.03.005.
Rowley, M.J., and Corces, V.G. (2018). Organizational principles of 3D genome architecture.
Nature Reviews Genetics *19*, 789-800.
Sanborn, A.L., Rao, S.S., Huang, S.-C., Durand, N.C., Huntley, M.H., Jewett, A.I., Bochkov,
I.D., Chinnappan, D., Cutkosky, A., and Li, J. (2015). Chromatin extrusion explains key features
of loop and domain formation in wild-type and engineered genomes. Proceedings of the National
Academy of Sciences *112*, E6456-E6465.
Schoborg, T.A., and Labrador, M. (2010). The phylogenetic distribution of non-CTCF insulator
proteins is limited to insects and reveals that BEAF-32 is Drosophila lineage specific. J. Mol.
Evol. *70*, 74-84. 10.1007/s00239-009-9310-x.
Sigrist, C.J., and Pirrotta, V. (1997). Chromatin insulator elements block the silencing of a target
gene by the Drosophila polycomb response element (PRE) but allow trans interactions between
PREs on different chromosomes. Genetics *147*, 209-221.
Sun, Y., Nien, C.Y., Chen, K., Liu, H.Y., Johnston, J., Zeitlinger, J., and Rushlow, C. (2015).
Zelda overcomes the high intrinsic nucleosome barrier at enhancers during Drosophila zygotic
genome activation. Genome Res. *25*, 1703-1714. 10.1101/gr.192542.115.
Tan, L., Shi, J., Moghadami, S., Parasar, B., Wright, C.P., Seo, Y., Vallejo, K., Cobos, I.,
Duncan, L., Chen, R., and Deisseroth, K. (2023). Lifelong restructuring of 3D genome
architecture in cerebellar granule cells. Science *381*, 1112-1119. 10.1126/science.adh3253.
Trcek, T., Lionnet, T., Shroff, H., and Lehmann, R. (2017). mRNA quantification using single-
molecule FISH in Drosophila embryos. Nat Protoc *12*, 1326-1348. 10.1038/nprot.2017.030.
Ueberschär, M., Wang, H., Zhang, C., Kondo, S., Aoki, T., Schedl, P., Lai, E.C., Wen, J., and
Dai, Q. (2019). BEN-solo factors partition active chromatin to ensure proper gene activation in
Drosophila. Nat Commun *10*, 5700. 10.1038/s41467-019-13558-8.
Vazquez, J., Farkas, G., Gaszner, M., Udvardy, A., Muller, M., Hagstrom, K., Gyurkovics, H.,
Sipos, L., Gausz, J., Galloni, M., and et al. (1993). Genetic and molecular analysis of chromatin
domains. Cold Spring Harb Symp Quant Biol *58*, 45-54. 10.1101/sqb.1993.058.01.008.
Vazquez, J., Muller, M., Pirrotta, V., and Sedat, J.W. (2006). The Mcp element mediates stable
long-range chromosome-chromosome interactions in Drosophila. Molecular Biology of the Cell
17, 2158-2165.
Yokoshi, M., Segawa, K., and Fukaya, T. (2020). Visualizing the Role of Boundary Elements in
Enhancer-Promoter Communication. Mol. Cell *78*, 224-235.e225. 10.1016/j.molcel.2020.02.007.
Zhimulev, I.F., and Belyaeva, E.S. (1991). Chromomeric organization of polytene chromosomes.
Genetica *85*, 65-72. 10.1007/bf00056107.
Zhimulev, I.F., Pokholkova, G.V., Bgatov, A.V., Semeshin, V.F., Umbetova, G.H., and
Belyaeva, E.S. (1983). Genetic interpretation of polytene chromosomes banding pattern. Mol.
Biol. Rep. *9*, 19-23. 10.1007/bf00777469.

Zolotarev, N., Fedotova, A., Kyrchanova, O., Bonchuk, A., Penin, A.A., Lando, A.S., Eliseeva,
I.A., Kulakovskiy, I.V., Maksimenko, O., and Georgiev, P. (2016). Architectural proteins Pita,
Zw5,and ZIPIC contain homodimerization domain and support specific long-range interactions
in Drosophila. Nucleic Acids Res. *44*, 7228-7241. 10.1093/nar/gkw371.
1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

FIGURES

1417

1418
1419 **Figure 1. Diagram of the possible loop topologies generated by loop-extrusion and**
1420 **boundary:boundary pairing.**

1421

1422

1423

1424
1425 **Figure 2. TAD organization of the *even-skipped* locus and neighboring sequences.**
1426
1427
1428
1429

1430
1431 **Figure 3. Schematics of boundary pairing and loop extrusion.**

1441
1442
1443 **Figure 4. The *lacZ* reporter is activated by *eve* enhancers in the *LhomieG Z5* transgene.**

1458

1459
1460 **Figure 5. The *GFP* reporter is activated by both *eve* and *hebe* enhancers in the *LhomieG***
1461 ***Z5* transgene.**

1470
1471 **Figure 6. Expression of reporters in Z3 transgenes.**

Figure 7. *homie* in Z5 transgene *LhomieG* and *GhomieL* Z3 mediates long-range interactions with the *eve* locus.

1498
1499 **Figure 8.** *homie* **in the transgene interacts with** *eve homie* **and** *nhomie.*
1500