Determinants of spring migration departure dates in a New World sparrow: weather variables

reign supreme

Allison J. Byrd[1,2*], Katherine M. Talbott[2], Tara M. Smiley[3], Taylor B. Verrett[5], Michael S. Gross[4],

Michelle L. Hladik[4], Ellen D. Ketterson[1,2], Daniel J. Becker[5*]

[1]Environmental Resilience Institute, Indiana University, Bloomington, IN

[2]Department of Biology, Indiana University, Bloomington, IN

[3]Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY

[4]U.S. Geological Survey, California Water Science Center, Sacramento, CA, 95819

[5]School of Biological Sciences, University of Oklahoma, Norman, OK

*These authors contributed equally.

Corresponding author: Allison Byrd, e-mail: abyrd.in.uni@gmail.com, present address: Biology

Building, 1001 E 3rd St, Bloomington, IN 47405

Funding

Funds were provided by Indiana University's Grand Challenge Initiative, Prepared for

Environmental Change to EDK and TMS and from the University of Oklahoma to DJB.

Conflict of interest

All authors declare they have no conflicts of interest.

Keywords: movement ecology, songbirds, haemosporidians, Motus, stable isotopes

Abstract

Numerous factors influence the timing of spring migration in birds, yet the relative importance of intrinsic and extrinsic variables on migration initiation remains unclear. To test for interactions among weather, migration distance, parasitism, and physiology in determining spring departure date, we used Dark-eyed Juncos (*Junco hyemalis hyemalis*) as a model migratory species known to harbor diverse and common haemosporidian parasites. Prior to spring migration departure from their wintering grounds in Indiana, USA, we quantified the intrinsic variables of fat, body condition (i.e., mass~tarsus residuals), physiological stress (i.e., ratio of heterophils to lymphocytes), cellular immunity (i.e., leukocyte composition and total count), migration distance (i.e., distance to the breeding grounds) using stable isotopes of hydrogen from feathers, and haemosporidian parasite intensity. We then attached nanotags to determine the timing of spring migration departure date using the Motus Wildlife Tracking System. We used additive Cox proportional hazard mixed models to test how risk of spring migratory departure was predicted by the combined intrinsic measures, along with meteorological predictors on the evening of departure (i.e., average wind speed and direction, relative humidity, and temperature). Model comparisons found that the best predictor of spring departure date was average nightly wind direction and a principal component combining relative humidity and temperature. Juncos were more likely to depart for spring migration on nights with largely southwestern winds and on warmer and drier evenings (relative to cooler and more humid evenings). Our results indicate that weather conditions at take-off are more critical to departure decisions than the measured physiological and parasitism variables.

Introduction

Over-wintering duration and timing of departure for avian spring migration is a process influenced by a

host of extrinsic and intrinsic factors (Covino et al., 2015; Hurlbert and Liang, 2012; Satterfield et al.,

2018). The timing of spring migration must balance the advantages of early arrival on the breeding

grounds, including higher quality mates and territories (Gunnarsson et al., 2006; Møller, 1994; Newton,

2010; Rotics et al., 2018; Smith and Moore, 2005), with the risk of arriving before adequate food

resources are available (Gunnarsson et al., 2006; Møller, 1994; Newton, 2010; Rotics et al., 2018; Smith

and Moore, 2005). Stressors experienced on the wintering grounds can also affect the timing and duration

of spring migration. Extrinsic factors such as changing weather conditions (Marra et al., 2005) and

intrinsic factors such as high parasite intensity (Dietsch, 2005; Reed et al., 2003) can alter stopover

duration as birds recover from or accommodate these additional physiological burdens (Schmaljohann et

al. 2022; Skrip et al. 2015; Linscott and Senner 2021). The relative contribution of extrinsic and intrinsic

factors experienced at the wintering grounds in shaping spring departure remains less well understood,

and studies addressing these interactions are especially needed as anthropogenic pressures continue to

influence the climate and natural habitat (Kubelka et al., 2022; Visser et al., 2009; Wilcove and Wikelski,

2008).

For decades, ornithologists have known that weather variables influence avian migration timing in both

spring and fall (see:(Richardson, 1978). Factors that influence the probability of migratory departure

include wind speed (Chapman et al., 2016; Drake et al., 2014; Kemp et al., 2010; Liechti, 2006;

Nussbaumer et al., 2022), wind direction (Covino et al., 2015; Hebrard, 1971; Horton et al., 2016; Kemp

et al., 2010; Lack, 1963; Lack and Eastwood, 1962; Sinelschikova et al., 2007), temperature (Hüppop and

Winkel, 2006; Marra et al., 2005; Saino et al., 2007; Tøttrup et al., 2010; Usui et al., 2017), and relative

humidity (Klaassen et al., 2012; Liechti, 2006; Schmaljohann et al., 2009; Serra-Cobo et al., 1998; Zhang

and Wu, 2018). The high correlation among these weather variables adds to the difficulty in disentangling

the influence of these factors from potential additional stressors. For example, warm air can hold more

moisture than cold air, thus at the same absolute humidity, cooler air (perhaps counterintuitively) has

higher relative humidity than warmer air (Lawrence, 2005).

The relationship between wind and bird migration has been studies observed for decades, and prevailing

winds were a likely a selection factor in that shaping the evolution of migratory patterns (Alerstam 1979;

Alerstam 1979; Evans 2009; Able 1972; Richardson 1978). Birds that are highly selective of favorable

winds can maximize flight speed and minimize energy expenditure across both long and short-distance

migratory flights (Alerstam 1979). Migrants have faster ground and air-speeds in spring migration than

fall (Horton et al. 2016), and spring migration is typically completed over a shorter duration than fall

(Newton 2010; La Sorte et al. 2013; La Sorte et al. 2016; Nilsson et al. 2013). As such, models that

forecast bird migration rely heavily on wind data (Erni et al., 2002; Van Doren and Horton, 2018).

Intrinsic factors have also been recognized to affect readiness for migratory departure. Given the

energetic costs of migration (Wikelski et al., 2003), adequate fat stores are integral in providing fuel for

long-distance flight and larger stores can be a reliable predictor of migratory readiness(Price, 2010;

Ramenofsky, 1990; Weber et al., 1994; Witter and Cuthill, 1993); correspondingly, measures of body

condition (i.e., mass~tarsus residuals) allow for standardized body size comparisons among individuals

(Labocha and Hayes, 2012). The degree of physiological stress and immune state of individuals at the

wintering grounds can also affect spring migratory timing. Preparation for long-distance migration

increases plasma concentrations of corticosterone (the primary avian glucocorticoid responsible for

maintaining homeostasis) in many songbirds (Holberton, 1999; Landys et al., 2004), which can facilitate

earlier departure from the wintering grounds or spring stopover (Eikenaar et al., 2013, 2017). Elevated

corticosterone can shift the composition of leukocytes in blood, subsequently elevating the ratio of

heterophils to lymphocytes (HL ratios; Davis et al., 2008). While plasma corticosterone levels are highly

sensitive to the acute stress of capture, stress-induced changes in leukocyte profiles occur more slowly,

such that HL ratios can serve as a more tractable approximation of energetic costs prior to migration

(Davis and Maney, 2018). In a similar fashion, the energetic cost of migratory preparation can induce

trade-offs with the immune system (Lochmiller and Deerenberg, 2000), such that migrants may

downregulate immune activity or rely primarily on less costly immune defenses. For example, several

thrush species show lower total leukocyte counts upon arrival at stopover sites in spring, reflecting such

trade-offs (Owen and Moore, 2008). Variation in immune investment may in turn affect departure

decisions; recent work demonstrated that songbirds with higher titers of natural antibodies and

immunoglobulin Y have longer spring stopovers (Brust et al., 2022).

Related to seasonal variation in physiological stress and immunity, songbirds may also experience

additional overwintering stressors that affect spring migration. For example, exposure to toxins such as

neonicotinoids in wintering food sources (Eng et al., 2017; Goulson, 2014) can delay migration departure

or increase stopover duration as birds recover from or accommodate these burdens. An additional such

stressor could be through parasite infection, such as that from dipteran-borne haemosporidian blood

parasites (i.e., the genera *Plasmodium*, *Haemoproteus*, and *Leucocytozoon*). Experimental infections of

captive birds have shown migration-relevant physiological costs of infection, such as reduced general

activity (Mukhin et al., 2016; Yorinks and Atkinson, 2000) and body condition (Atkinson et al., 1995) as

well as shifts in the development of migratory restlessness (Kelly et al., 2016, 2020). Observational

studies have also identified impacts of haemosporidian infection on body condition in migrating birds

(Garvin et al., 2006; Merrill et al., 2018), on the timing of arrival to the breeding grounds (Asghar et al.,

2011; Santiago-Alarcon et al., 2013), and on autumn departure timing (Ágh et al., 2019). However, it

remains unclear how haemosporidian infections interact with extrinsic and other intrinsic factors to affect

migration timing.

In this study, we used overwintering migratory Dark-eyed Juncos (*Junco hyemalis hyemalis;* hereafter

"junco") to test the hypothesis that physiological state and haemosporidian infection interact with weather

conditions and migration distance to shape spring migratory departure timing. Haemosporidia have been

well characterized in Dark-eyed juncos, with chronic (i.e., long- term) infections detected in wintering

and breeding populations (Becker et al. 2020; Slowinski et al. 2018; Talbott et al. 2022; Martínez-Renau

et al. 2022; Becker et al. 2019; Deviche, Greiner, and Manteca 2001; Ferrer 2022). We used stable

isotopes of hydrogen from feathers to estimate the distance to breeding grounds for each individual

(Bowen et al., 2014; Hobson, 1999; Hobson et al., 2012; Rubenstein and Hobson, 2004; Wunder, 2010)

and the Motus Wildlife Tracking System to determine departure date (Taylor et al., 2017). We predicted

that birds with higher fat reserves, low intensity or no haemosporidian parasitism, lower HL ratios, and

lower total leukocyte counts would depart earlier in the year and during evenings of following winds (i.e.,

tailwind). We also predicted that wind direction (following winds) would be more important than wind

speed, and that headwinds would be the strongest deterrent to departure, even more so than intrinsic

predictors.

Methods

Junco sampling and captive housing

We caught wild juncos from 31 January to 26 February 2020 at four locations within 15 km of

Bloomington, Indiana. None of the birds were captured at the release site, and we systematically rotated

capture efforts through all four sites to avoid confounding results by capture date or location. We captured

juncos using baited mist nets and walk-in traps, took standard morphometric measurements including

mass and body fat (Fudickar et al., 2016; Jawor et al., 2006; Singh et al., 2019), and fitted each bird with

an aluminum US Fish & Wildlife Service leg band (federal permit # 20261, Indiana state permit 20-528).

While capture efforts were ongoing, juncos were held at an indoor aviary at Indiana University to prevent

additional exposure to arthropod vectors of haemosporidia. Juncos were provided *ad libitum* water and

food (mealworms, organic millet, organic sunflower seeds, and a blended mixture of organic millet,

organic carrots, and organic blueberries) and could fly freely in 6.4 x 3.2 m rooms. Light cycles for rooms

reflected the natural photoperiod of Bloomington, Indiana at the time of the study.

As a potential additional extrinsic factor, juncos in this study were randomly dosed with either a

neonicotinoid (imidacloprid suspended in sunflower seed oil) or control (sunflower seed oil) to

experimentally test impacts on migration timing. However, imidacloprid metabolites of 5-OH-imidacloprid

and imidacloprid-olefin were detected in only one bird's post-dose plasma at 46.67 ng/mL and 72.23 ng/mL.

This individual was removed from all subsequent analyses. Imidacloprid or metabolites were not detected in

pre-dose or post-dose (6 hours after) plasma of any other individuals ($n = 37$; Supp. Appx A, Table S1).

Therefore, we did not include imidacloprid exposure as a possible predictor of migratory timing. Possible

explanations for the absence of imidacloprid detection in plasma samples are that the dose was

regurgitated or rapidly metabolized by all birds.

On the days of release (3 and 4 March 2020), we attached a nanotag to each bird (Lotek model NTQB2-2,

11 x 5 x 4 mm, 0.32g, $\leqq$ 0.5 g total mass including leg harness). Birds were released at Kent Farm

Research Station (located nine miles east of Bloomington, IN), an area frequented by overwintering

juncos; we provided organic seed in feeders and on the ground in an attempt to minimize departure from

the area due to resource limitation (Bridge et al., 2010) .

Hematological analyses

At capture, we collected $\leq$ 150 μL blood from each bird by pricking the brachial vein with a sterile

needle, followed by collection with heparinized capillary tubes. We also collected the first secondary

feather from each bird for stable isotope analysis (Fudickar et al., 2016). We separated plasma and red

blood cells (centrifuged for 10 min at 10,000 rpm) and stored samples at -20°C until DNA extraction

using a Maxwell RSC Whole Blood DNA Kit (Promega). Any birds unable to be sexed by wing length

and plumage were verified using sexing PCR of extracted DNA (Griffiths et al., 1998). Only males

($n=37$) were included to control for sex differences in migration timing, as female juncos migrate earlier

than males (Nolan and Ketterson, 1990).

Because birds were held in captivity for variable lengths between capture and release with nanotags

($\bar{x}$=25, range of 13 to 33 days), we collected additional blood samples 24 hours prior to release, using the

blood collection protocol described above. We prepared thin blood smears on glass slides stained with

Wright–Giemsa (Quick III, Astral Diagnostics). We then evaluated leucocyte profiles and

haemosporidian intensity using light microscopy (AmScope, B120C-E1). A single observer (TBV)

recorded the number of leukocytes under 400X magnification across 10 random fields to estimate

inflammatory state and investment in cellular immunity. A differential count was then performed by

recording the identity of the first 100 leukocytes (heterophils, lymphocytes, monocytes, eosinophils, and

basophils) at 1000X magnification (oil immersion;(Campbell, 1995). We then screened 100 fields of view

at 1000X magnification for *Plasmodium*, *Haemoproteus*, and *Leucocytozoon (Becker et al., 2019, 2020;*

Cosgrove, 2005; Valkiunas et al., 2008). We derived the mean total leukocyte count, HL ratios, and the

total number of haemosporidian-infected erythrocytes (parasite intensity) as three predictor variables.

Usable blood smears were available for 34 of our 37 tagged individuals.

Hydrogen isotope analysis

We used stable isotopic analysis of hydrogen (δ^2H) in feathers to infer probable breeding latitude and

median distance from capture point(Bowen et al., 2014; Hobson, 1999; Hobson et al., 2012; Rubenstein

and Hobson, 2004; Wunder, 2010). Feathers were cleaned using a 2:1 chloroform:methanol solution to

remove external oils and contaminants. We used a forced isotopic equilibration procedure to ensure

exchangeable hydrogen with a water vapor of known isotopic composition in a flow-through chamber

system at 115°C (Sauer et al., 2009; Schimmelmann, 1991). Samples were analyzed using a thermal

conversion element analyzer coupled with a ThermoFinnigan Delta Plus XP isotope ratio mass

spectrometer at the Indiana University Stable Isotope Research Facility. Isotopic data are reported in

standard per mil notation (‰) relative to VSMOW (Vienna Standard Mean Oceanic Water) using two

reference materials: USGS77 (polyethylene powder) and hexatriacontane 2 (C_{36} n-alkane 2). Analytical

precision was ± 1.0‰ for δ^2H values. We calculated the isotopic composition of the non-exchangeable

hydrogen per sample, assuming a 17% exchangeability rate for feathers (Schimmelmann, 1991;

Schimmelmann et al., 1999).

We performed Bayesian geographic assignments for the breeding location of individual birds based on

feather δ^2H values using the *assignR* package (Ma et al., 2020). To calibrate our precipitation-feather

isoscape, we used growing season precipitation isoscape rasters from waterisotopes.org (Bowen et al.,

2005; Bowen and Revenaugh, 2003) and the isotopic composition of non-migratory Dark-eyed juncos

from previous studies(Becker et al., 2019; Hobson et al., 2012). Median latitudinal and distance estimates

were calculated from geographic assignment probability maps by extracting the geographic cell

coordinates with the highest posterior probability (top 10%) using the *raster* package (Hijmans, 2019).

Migration distance was defined as the distance from the release site to the centroid of the estimated

breeding polygon (Becker et al., 2022; Wanamaker et al., 2020).

Motus data processing

Motus data (project 240; available at https://motus.org/data) were downloaded on 25 July 2020, four

months after the known period of spring migration in Indiana of wintering juncos (Ketterson and Nolan,

1982). Data were filtered and cleaned following a standard protocol using the *motus* package in R(Taylor

et al., 2017). Specifically, we removed runs with a low probability of being a true detection (i.e.,

motusFilter=0) and adopted a moderately strict filter to be more conservative about detection inclusions.

We then derived the departure date as the last day (ordinal date) a bird was detected at the Kent Farm

Research Station Motus station. We additionally report cases in which birds were detected at other Motus

stations between banding and our data download date.

Weather data

We obtained relative humidity data from ncei.noaa.gov (Diamond et al., 2013) and all remaining weather

variables (humidity, wind speed, wind direction [converted to wind rose] and weather type) from

timeanddate.com (Thorsen, 1995-2023). We averaged hourly data from 1800-2400 on departure date

evenings (thus, variables are labeled average humidity and so on).

Because average relative humidity and average temperature are dependent (Lawrence, 2005), we

conducted a principal components analysis (PCA) of these two variables, with variables centered and

scaled to have unit variance. The first PC (hereafter weather PC1) explained 56% of the variation and was

loaded negatively by average relative humidity (–0.71) and positively by average temperature (0.71);

resulting values indicate increasingly warm and dry evening weather. This variable was used in

downstream analyses alongside average wind speed and average wind direction.

Statistical analyses

Prior to analyses of migration timing, we used a linear mixed effects model with a random effect of site to

derive an index of body condition through the residuals of a regression of mass (measured prior to

release) on tarsus length (Schulte-Hostedde et al., 2005; Wanamaker et al., 2020).

We modeled days until spring migratory departure under an event-time analysis framework, using

additive Cox proportional hazard mixed models (CPHMMs, (Therneau and Grambsch, 2000) and

Grambsch, 2000). These semi-parametric models allow determining how the risk of departure changes

with covariates that can also vary with time, and such models have been applied to study migration timing

in both avian and non-avian systems (Castro-Santos and Haro, 2003; Dossman et al., 2015). We used the

mgcv package to fit CPHMMs with a random effect of site. To test the direct and interactive effects of

extrinsic (i.e., weather PC1, wind direction, wind speed) and intrinsic predictors (i.e., fat, body condition,

total leukocytes, HL ratios, haemosporidian intensity) on risk of spring departure (Wood, 2017).

Including all predictor variables and biologically relevant interactions in a single full model was not

possible given our sample size ($n = 34$, excluding birds without hematology data). We instead built 10

candidate CPHMMs representing *a priori* hypotheses of additive and interactive effects while restricting

models to at most three fixed effects to limit overfitting (Burnham and Anderson, 2002). Most predictors

showed low collinearity (ρ ranged from –0.63 to 0.62, x̄ = 0.01), and moderately correlated predictors

(i.e., weather PC1 and wind speed, ρ = –0.63; body fat and condition, ρ = 0.62) were excluded from the

same models. All predictors were modeled using thin plate splines with smoothing penalty with the

exception of wind direction, which used a cyclic cubic spline to account for circular data. Interaction

terms were modeled as tensor products. Our models represented additive effects of intrinsic variables only

(e.g., fat, haemosporidian intensity, and total leukocytes), additive effects of extrinsic variables only (e.g.,

wind direction and weather PC1), additive effects of both intrinsic and extrinsic variables (e.g.,

haemosporidian intensity, HL ratios, and weather PC1), interactive effects of intrinsic variables (e.g.,

effects of haemosporidian intensity dependent on HL ratios), interactive effects of extrinsic variables

(e.g., effects of wind speed depend on wind direction), and interactive effects of intrinsic and extrinsic

variables (e.g., effects of weather PC1 depend on haemosporidian intensity). We compared CPHMMs fit

with maximum likelihood using Akaike information criterion adjusted for small sample size (AICc) with

the *MuMIn* package(Bartoń, 2013). We also derived Akaike weights (w_i) to facilitate comparison and

considered models within two ΔAICc of the top model to be competitive. Competitive models were

refitted to the full dataset, with CPHMM predictions visualized as relative hazards with 95% confidence

intervals (Nakagawa and Schielzeth, 2013).

Results

Hydrogen isotopes and likely breeding origins

Hydrogen isotopic composition of junco feathers ranged from -158.9 to -95.8%, reflecting median

breeding locations from 66 to 51°N, respectively (Fig. 1). Maximum and minimum latitudinal estimates

spanned from 70 to 36 °N. Estimated distances to breeding grounds accordingly ranged from

approximately 2086 to 3900 kilometers (x̄=3297, SE=78.4). Incidentally, we detected six of our 37 tagged

juncos at other Motus towers following their departure from Indiana (Fig. 1). Five birds were detected at

the Lake Petite station in Wisconsin (42.5117°, -88.5488°) and one bird was detected at the Werden

station (42.7551°, -80.2724°) in Ontario, Canada.

Haemosporidian infection

We identified haemosporidian infections in 11 of 34 juncos prior to release (32.4%, 95% CI: 19.1–

49.2%). Only two individuals had detectable *Haemoproteus* spp. infection (intensity=16–23 infected

erythrocytes from 100 fields of view), and nine individuals had detectable *Leucocytozoon* spp. infection

(intensity=1–7 infected erythrocytes from 100 fields of view); no individuals harbored co-infecting

haemosporidian parasites. We also opportunistically detected *Trypanosoma* spp. in a single bird (1/34,

two parasites were detected from 100 fields of view), and this individual was not infected by either

Haemoproteus spp. nor *Leucocytozoon* spp. parasites.

Extrinsic and intrinsic predictors of migration timing

All birds departed 8–31 days after release ($\bar{x}$=19.11 ± 0.81 SE), between March 11 and April 3 2020 (Fig.

2). Comparison among 10 CPHMMs predicting risk of departure date as a function of extrinsic and

intrinsic factors identified only one competitive model, which included the nonlinear additive effects of

wind speed and weather PC1 (w_i=0.61; Table 1). Wind direction had the greatest relative importance

among predictors (99%), followed by weather PC1 (62%); wind speed, the interaction between wind

speed and weather PC1, and estimated migration distance all had lesser importance (38%, 22%, and 16%,

respectively). All intrinsic predictors (i.e., fat, body condition, total leukocytes, HL ratios) and their

interactions were unimportant (≤ 1%).

Our top model explained 30.5% of the deviance in spring migration risk, with both wind direction

($\chi^2$1.8,2 = 17.42, p < 0.001) and weather PC1 ($\chi^2$1.4,3 = 29.61, p < 0.001) being significant non-linear

predictors of departure. Specifically, the risk of spring departure was greatest with average nightly

southwestern winds and for humid and cool nights (Fig. 3).

Discussion

Numerous factors influence the timing of spring departure in birds, yet the relative influence of

intrinsic and extrinsic variables on migration initiation remains a topic of debate. In this study, we

tested the relative influence of these diverse factors on the risk of spring departure.in a modestly-

sized group of wild-caught juncos. Weather variables outperformed all tested intrinsic variables.

We predicted that both wind direction and speed would have the greatest influence on departure;

however, only wind direction was supported by our top model. Temperature and relative humidity

(i.e., weather PC1) also influenced departures, with humid and cool nights having greater risk of

departure than warmer, drier nights. While we predicted that birds with intense haemosporidian

infection would depart later than uninfected individuals or those with low-intensity infections,

infection intensity did not influence departure risk. Additionally, higher fat reserves, lower

leukocyte counts, and lower HL ratios did not affect departure risk. Our results suggest that the

advantage of following winds outweigh the cost of haemosporidian infections and the other

measured physiological variables during spring migration.

Many migratory birds harbor haemosporidian infections throughout the year, including during migration,

(Ricklefs et al. 2017; Cornelius et al. 2014; Pulgarín-R et al. 2019) and ongoing research continues to

clarify the effects of avian malaria on migration timing, duration, and distance. Prior work on juncos

found greater prevalence of haemosporidian infections in a non-migratory subspecies compared to

sympatric overwintering migrants following fall migration, suggesting that infected birds may be more

likely to experience disease-induced mortality during migration (Slowinski et al., 2018). Similarly, juncos

with longer migrations were more likely to show elevated HL ratios upon arrival to their wintering

grounds; this effect was stronger in haemosporidian-infected birds, suggesting interactive effects of

migration and infection on host energetics (Becker et al., 2019). Therefore, we predicted that juncos with

more intense infections might delay migration initiation to mitigate these potential physiological costs.

However, our results indicate that weather has a strong impact on migration irrespective of

haemosporidian intensity. Although birds must navigate both intrinsic and extrinsic factors when

choosing when to depart for migration, the cost of initiating spring migration under sub-optimal wind

conditions may be greater than the cost of departing with a more intense haemosporidian infection.

Importantly, our study focused on spring migration in adult juncos. While chronic haemosporidian

infections may not impact migration timing during this life stage, it is unclear if this pattern would hold

true for juncos during autumn migration, especially juveniles. In fact, juvenile European robins (Erithacus

rubecula) with haemosporidian infections arrive at wintering ground later than uninfected subadults (Ágh

et al. 2019). Thus, season, age, and stage of infection may influence whether haemosporidian infections

delay migration initiation. Experimental infections in conjunction with departure tracking would provide

more conclusive data on how weather interacts with parasite intensity to influence migration initiation.

Our analysis also indicates that migration distance and mean wind speed were relatively less informative

predictors of spring departure date. Regarding migration distance, we expected that birds with longer

spring migrations would depart earlier, as these individuals would potentially need to undertake

prolonged stopovers to rest and refuel on the way to their breeding grounds. However, research suggests

that long-distance migrants have a greater ability to modify migratory behavior while en route and thus

departure date may not be directly related to migration distance as migration speed and routes are flexible

(La Sorte and Fink, 2017; Marra et al., 2005). While juncos have been well-studied in regard to

differential migration and migration timing (Cristol et al., 1999; Ketterson and Nolan, 1976), more

research is needed to understand how distance to breeding ground influences departure date. Isotopic

feather analysis is advancing work in this field, as is the improvement and proliferation of nanotag

technology (e.g., Motus, Cellular Tracking Technologies), which will help further distinguish the various

influences on migration distance as it relates to departure.

Past studies have shown that both wind speed and direction are important predictors of departure

and that tail winds can significantly increase flight times and greatly decrease energy requirements

(T. Alerstam, 1979; Bloch and Bruderer, 1982; Kemp et al., 2010; Liechti, 2006). There are several

possibilities as to why these variables were less important in our analysis. Optimal wind selectivity

is a product of many factors, including the size of the bird; rate of fat use; distance, duration, and

altitude of migration; and even the wind pattern itself (Thomas Alerstam, 1979). Lastly, birds here

departed over a relatively short time frame (a minimum of eight days after release, up to 31 days);

therefore, there may not have been sufficient variation in wind speed and direction to outrank other

predictors in our models.

In contrast to our expectations about physiological state and departure date, body condition, fat score, HL

ratios, and leukocyte counts were also all uninformative predictors of spring migration date. Higher fat

reserves are instrumental to migratory performance in songbirds (Price 2010) and may promote the onset

of migratory restlessness, advancing their departure (Lupi et al., 2017; Studds and Marra, 2005). The lack

of a relationship between physiological measures and departure date observed here could reflect

individual variation in migration strategies; for example, strong selective pressure to arrive early at the

breeding grounds may motivate some birds to leave in relatively poor body condition and with high HL

ratios and then compensate during stopover (Prop et al., 2003). Lastly, the time between release from

captivity and the onset of spring migration (8-31 days) may have precluded a true representation of some

physiological measures at departure; for example, maximum fat deposition rates in migratory songbirds

have been reported as high as 12% differences in lean body mass in one day (Lindström, 1991).

The findings presented here add to the large body of information on the importance of weather on

migration departure in songbirds. Importantly, our study focused on migratory departure in male juncos,

while females may be differentially impacted by the variables measured. Indeed, this species shows sex-

based variation in overwintering latitudes, with females migrating earlier and further south than males

(Nolan and Ketterson 1990). Our sample size is a significant limitation in understanding and asking these

questions, as well. A more robust sample size would have increased the confidence in our findings.

Additional data are also needed to understand whether the results of our study are context dependent; for

example, intrinsic factors might become important during years with low food availability or inclement

weather. This will be increasingly important for wildlife conservation going forward, as climate change is

associated with increasingly severe temperatures and unpredictable weather events (X. Zhang et al. 2013;

Stott 2016).

Data availability

The data that support the findings of this study are openly available in Dryad (https://datadryad.org) at

https://datadryad.org/stash/share/Ne24zAcw7YGshc_QeUDPeWFrXNzJaPPGd-0kctpgK7Y.

Ethics

Trapping, banding, and transmitter attachment was overseen by AJB and performed by AJB, DJB, and

KMT under United States Federal Permit # 20261, Indiana state permit # 20-528, and approved under

Bloomington Institutional Animal Care and Use Committee (BIACUC) # 18-030-15, Section 1. A Doctor

of Veterinary Medicine (DVM) from Indiana University's Laboratory Animal Resources (LAR) taught

the oral gavage technique to AJB, DJB and KMT and this procedure was approved under the same

protocol (BIACUC #18-030-15, Section 30).

Competing interests

The authors declare they have no competing interests.

Acknowledgements

We thank Benjamin Higgins, Indiana University for his dedication in the lab. We also thank

Peter Sauer and Kathyrn Evans in the Stable Isotope Research Facility at Indiana University for

372 assistance in isotopic preparation and analysis. Lastly, we thank members of the Ketterson

373 laboratory at Indiana University and two anonymous reviewers for helpful feedback on previous

374 versions of this manuscript.

References

Ágh N, Piross IS, Majoros G, et al. (2019) Malaria infection status of European Robins seems to associate
with timing of autumn migration but not with actual condition. *Parasitology* 146(6): 814–820.

Alerstam T. (1979) Optimal use of wind by migrating birds: combined drift and overcompensation.
Journal of theoretical biology 79(3). Elsevier: 341–353.

Alerstam Thomas (1979) Wind as Selective Agent in Bird Migration. *Ornis Scandinavica* 10(1). [Nordic
Society Oikos, Wiley]: 76–93.

Asghar M, Hasselquist D and Bensch S (2011) Are chronic avian haemosporidian infections costly in
wild birds? *Journal of avian biology* 42(6). Wiley: 530–537.

Atkinson CT, Woods KL, Dusek RJ, et al. (1995) Wildlife disease and conservation in Hawaii:
pathogenicity of avian malaria (Plasmodium relictum) in experimentally infected iiwi (Vestiaria
coccinea). *Parasitology* 111 Suppl. cambridge.org: S59–69.

Baillie J, Hilton-Taylor C, Stuart SN, et al. (2004) *2004 IUCN Red List of Threatened Species: A Global
Species Assessment*. IUCN.

Bartoń K (2013) MuMIn: Multi-Model Inference, version 1.9. 0. *R package*. Epub ahead of print 2013.

Beaudoin RL, Applegate JE, Davis DE, et al. (1971) A model for the ecology of avian malaria. *Journal of
wildlife diseases* 7(1). meridian.allenpress.com: 5–13.

Becker DJ, Talbott KM, Smiley TM, et al. (2019) Leukocyte profiles vary with breeding latitude and
infection status in a seasonally sympatric songbird. *Animal Migration*. Available at:
http://dx.doi.org/10.1515/ami-2019-0004.

Becker DJ, Singh D, Pan Q, et al. (2020) Artificial light at night amplifies seasonal relapse of
haemosporidian parasites in a widespread songbird. *Proceedings. Biological sciences / The Royal
Society* 287(1935): 20201831.

Becker DJ, Byrd A, Smiley TM, et al. (2022) Novel Rickettsia spp. in two common overwintering North
American songbirds. *Emerging microbes & infections* 11(1). Taylor & Francis: 2746–2748.

Bloch R and Bruderer B (1982) The air speed of migrating birds and its relationship to the wind.
Behavioral ecology and sociobiology 11(1). Springer: 19–24.

Bowen GJ and Revenaugh J (2003) Interpolating the isotopic composition of modern meteoric
precipitation. *Water resources research* 39(10). American Geophysical Union (AGU).

Bowen GJ, Wassenaar LI and Hobson KA (2005) Global application of stable hydrogen and oxygen
isotopes to wildlife forensics. *Oecologia* 143(3). Springer: 337–348.

Bowen GJ, Liu Z and Vander Zanden HB (2014) Geographic assignment with stable isotopes in IsoMAP.
Methods in ecology and evolution / British Ecological Society. Wiley Online Library. Epub ahead of
print 2014. DOI: 10.1111/2041-210X.12147.

Bridge ES, Kelly JF, Bjornen PE, et al. (2010) Effects of nutritional condition on spring migration: do
migrants use resource availability to keep pace with a changing world? *The Journal of experimental*

biology 213(Pt 14): 2424–2429.

Brust V, Eikenaar C, Packmor F, et al. (2022) Do departure and flight route decisions correlate with
immune parameters in migratory songbirds? *Functional ecology* 36(12). John Wiley & Sons, Ltd:
3007–3021.

Burnham KP and Anderson DR (2002) A practical information-theoretic approach. *Model selection and*
multimodel inference, 2nd ed. Springer, New York 2.

Campbell TW (1995) Avian hematology and cytology. *Journal of zoo animal medicine* 19(4). JSTOR:
244.

Castro-Santos T and Haro A (2003) Quantifying migratory delay: a new application of survival analysis
methods. *Canadian journal of fisheries and aquatic sciences. Journal canadien des sciences*
halieutiques et aquatiques 60(8). Canadian Science Publishing: 986–996.

Chapman JW, Nilsson C, Lim KS, et al. (2016) Adaptive strategies in nocturnally migrating insects and
songbirds: contrasting responses to wind. *The Journal of animal ecology* 85(1). Wiley Online
Library: 115–124.

Cornelius EA, Davis AK and Altizer SA (2014) How important are hemoparasites to migratory
songbirds? Evaluating physiological measures and infection status in three neotropical migrants
during stopover. *Physiological and biochemical zoology: PBZ* 87(5). journals.uchicago.edu: 719–
728.

Cosgrove C (2005) Avian malaria parasites and other haemosporidia.—Gediminas valkiunas. 2004. CRC
press, Boca Raton, Florida, USA. 932 pp. ISBN 0-415-30097-5. US$170 (hardcover). *Systematic*
biology 54(5). Oxford University Press (OUP): 860–863.

Covino KM, Holberton RL and Morris SR (2015) Factors influencing migratory decisions made by
songbirds on spring stopover. *Journal of avian biology* 46(1): 73–80.

Cristol DA, Baker MB and Carbone C (1999) Differential Migration Revisited. In: Nolan V, Ketterson
ED, and Thompson CF (eds) *Current Ornithology*. Boston, MA: Springer US, pp. 33–88.

Davis AK and Maney DL (2018) The use of glucocorticoid hormones or leucocyte profiles to measure
stress in vertebrates: What's the difference? *Methods in ecology and evolution / British Ecological*
Society 9(6). Wiley: 1556–1568.

Davis AK, Maney DL and Maerz JC (2008) The use of leukocyte profiles to measure stress in vertebrates:
a review for ecologists. *Functional ecology* 22(5). Wiley: 760–772.

de Angeli Dutra D, Fecchio A, Braga ÉM, et al. (2022) Migratory behaviour does not alter
cophylogenetic congruence between avian hosts and their haemosporidian parasites. *Parasitology*.
cambridge.org: 1–8.

Deviche P, Greiner EC and Manteca X (2001) Seasonal and age-related changes in blood parasite
prevalence in Dark-eyed Juncos (Junco hyemalis, Aves, Passeriformes). *The Journal of experimental*
zoology 289(7). Wiley: 456–466.

Diamond HJ, Karl TR, Palecki MA, et al. (2013) U.S. Climate Reference Network after One Decade of
Operations: Status and Assessment. *Bulletin of the American Meteorological Society* 94(4).
American Meteorological Society: 485–498.

Dietsch TV (2005) Seasonal variation of infestation by ectoparasitic chigger mite larvae (Acarina: Trombiculidae) on resident and migratory birds in coffee agroecosystems of Chiapas, Mexico. *The Journal of parasitology* 91(6): 1294–1303.

Dossman BC, Mitchell GW, Norris DR, et al. (2015) The effects of wind and fuel stores on stopover departure behavior across a migratory barrier. *Behavioral ecology: official journal of the International Society for Behavioral Ecology* 27(2). Oxford Academic: 567–574.

Drake A, Rock CA, Quinlan SP, et al. (2014) Wind speed during migration influences the survival, timing of breeding, and productivity of a neotropical migrant, Setophaga petechia. *PloS one* 9(5). journals.plos.org: e97152.

Eikenaar C, Fritzsch A and Bairlein F (2013) Corticosterone and migratory fueling in Northern wheatears facing different barrier crossings. *General and comparative endocrinology* 186. Elsevier: 181–186.

Eikenaar C, Müller F and Leutgeb C (2017) Corticosterone and timing of migratory departure in a songbird. *Royal Society B ….* royalsocietypublishing.org. Epub ahead of print 2017.

Eng ML, Stutchbury BJM and Morrissey CA (2017) Imidacloprid and chlorpyrifos insecticides impair migratory ability in a seed-eating songbird. *Scientific reports* 7(1): 15176.

Ephrath JE, Goudriaan J and Marani A (1996) Modelling diurnal patterns of air temperature, radiation wind speed and relative humidity by equations from daily characteristics. *Agricultural systems* 51(4). Elsevier: 377–393.

Erni B, Liechti F, Underhill LG, et al. (2002) Wind and rain govern the intensity of nocturnal bird migration in Central Europe- a log-linear regression analysis. *Ardea* 90(1). researchgate.net: 155–166.

Ferrer MM (2022) *AVIAN HAEMOSPORIDIAN BLOOD PARASITE DIVERSITY, PREVALENCE, AND DISTRIBUTION IN MICHIGAN'S WESTERN UPPER PENINSULA*. Michigan Technological University. Available at: https://digitalcommons.mtu.edu/etdr/1470/ (accessed 7 June 2023).

Fudickar AM, Greives TJ, Atwell JW, et al. (2016) Reproductive Allochrony in Seasonally Sympatric Populations Maintained by Differential Response to Photoperiod: Implications for Population Divergence and Response to Climate Change. *The American naturalist* 187(4): 436–446.

Garvin MC, Szell CC and Moore FR (2006) BLOOD PARASITES OF NEARCTIC–NEOTROPICAL MIGRANT PASSERINE BIRDS DURING SPRING TRANS-GULF MIGRATION: IMPACT ON HOST BODY CONDITION. *The Journal of parasitology* 92(5). Allen Press: 990–996.

Griffiths R, Double MC, Orr K, et al. (1998) A DNA test to sex most birds. *Molecular ecology* 7(8). Wiley Online Library: 1071–1075.

Gunnarsson TG, Gill JA, Atkinson PW, et al. (2006) Population-scale drivers of individual arrival times in migratory birds. *The Journal of animal ecology* 75(5): 1119–1127.

Hahn S, Bauer S, Dimitrov D, et al. (2018) Low intensity blood parasite infections do not reduce the aerobic performance of migratory birds. *Proceedings. Biological sciences / The Royal Society* 285(1871). royalsocietypublishing.org.

Hallmann CA, Foppen RPB, van Turnhout CAM, et al. (2014) Declines in insectivorous birds are associated with high neonicotinoid concentrations. *Nature* 511(7509): 341–343.

Hebrard JJ (1971) The nightly initiation of passerine migration in spring: a direct visual study. *The Ibis*.
Wiley Online Library. Epub ahead of print 1971. DOI: 10.1111/j.1474-919X.1971.tb05119.x.

Hijmans RJ (2019) Introduction to the'raster'package (version 2.8-19). mran.microsoft.com. Epub ahead
of print 2019.

Hobson KA (1999) Tracing origins and migration of wildlife using stable isotopes: a review. *Oecologia*
120(3): 314–326.

Hobson KA, Van Wilgenburg SL, Wassenaar LI, et al. (2012) Linking hydrogen ($\delta2H$) isotopes in
feathers and precipitation: sources of variance and consequences for assignment to isoscapes. *PloS*
one 7(4): e35137.

Holberton RL (1999) Changes in patterns of corticosterone secretion concurrent with migratory fattening
in a neotropical migratory bird. *General and comparative endocrinology* 116(1). Elsevier: 49–58.

Horton KG, Van Doren BM, Stepanian PM, et al. (2016) Seasonal differences in landbird migration
strategies. *The Auk* 133(4). Oxford Academic: 761–769.

Hüppop O and Winkel W (2006) Climate change and timing of spring migration in the long-distance
migrant Ficedula hypoleuca in central Europe: the role of spatially different temperature changes
along migration routes. *Journal of ornithology / DO-G* 147(2). Springer: 344–353.

Hurlbert AH and Liang Z (2012) Spatiotemporal variation in avian migration phenology: citizen science
reveals effects of climate change. *PloS one* 7(2): e31662.

Jawor JM, McGlothlin JW, Casto JM, et al. (2006) Seasonal and individual variation in response to
GnRH challenge in male dark-eyed juncos (Junco hyemalis). *General and comparative*
endocrinology 149(2). Elsevier: 182–189.

Kelly TR, MacGillivray HL, Sarquis-Adamson Y, et al. (2016) Seasonal migration distance varies with
natal dispersal and predicts parasitic infection in song sparrows. *Behavioral Ecology and*
Sociobiology. Available at: http://dx.doi.org/10.1007/s00265-016-2191-2.

Kelly TR, Rubin BD, MacDougall-Shackleton SA, et al. (2020) Experimental Malaria Infection Affects
Songbirds' Nocturnal Migratory Activity. *Physiological and biochemical zoology: PBZ* 93(2). The
University of Chicago Press: 97–110.

Kemp MU, Shamoun-Baranes J, Van Gasteren H, et al. (2010) Can wind help explain seasonal
differences in avian migration speed? *Journal of avian biology* 41(6). Wiley: 672–677.

Ketterson ED and Nolan V (1982) The Role of Migration and Winter Mortality in the Life History of a
Temperate-Zone Migrant, the Dark-Eyed Junco, as Determined from Demographic Analyses of
Winter Populations. *The Auk* 99(2). Oxford Academic: 243–259.

Ketterson ED and Nolan V Jr (1976) Geographic variation and its climatic correlates in the sex ratio of
eastern-wintering dark-eyed juncos (junco hyemalis hyemalis). *Ecology* 57(4). Wiley: 679–693.

Klaassen M, Hoye BJ, Nolet BA, et al. (2012) Ecophysiology of avian migration in the face of current
global hazards. *Philosophical transactions of the Royal Society of London. Series B, Biological*
sciences 367(1596): 1719–1732.

Kubelka V, Sandercock BK, Székely T, et al. (2022) Animal migration to northern latitudes:

environmental changes and increasing threats. *Trends in ecology & evolution* 37(1). Elsevier: 30–41.

Labocha MK and Hayes JP (2012) Morphometric indices of body condition in birds: a review. *Journal of*
ornithology / DO-G 153(1). Springer: 1–22.

Lack D (1963) Weather factors initiating migration. *Proc. XIII Int. Ornithol. Congr.* Epub ahead of print
1963.

Lack D and Eastwood E (1962) Radar films of migration over eastern England. *British birds; an*
illustrated magazine devoted to the birds on the British list. Epub ahead of print 1962.

Landys MM, Wingfield JC and Ramenofsky M (2004) Plasma corticosterone increases during migratory
restlessness in the captive white-crowned sparrow Zonotrichia leucophrys gambelli. *Hormones and*
behavior 46(5). Elsevier: 574–581.

La Sorte FA and Fink D (2017) Migration distance, ecological barriers and en-route variation in the
migratory behaviour of terrestrial bird populations. *Global ecology and biogeography: a journal of*
macroecology 26(2). Wiley: 216–227.

Lawrence MG (2005) The Relationship between Relative Humidity and the Dewpoint Temperature in
Moist Air: A Simple Conversion and Applications. *Bulletin of the American Meteorological Society*
86(2). American Meteorological Society: 225–234.

Liechti F (2006) Birds: blowin' by the wind? *Journal of ornithology / DO-G* 147(2). Springer Science and
Business Media LLC: 202–211.

Lindström Å (1991) Maximum Fat Deposition Rates in Migrating Birds. *Ornis Scandinavica* 22(1).
[Nordic Society Oikos, Wiley]: 12–19.

Lochmiller RL and Deerenberg C (2000) Trade-offs in evolutionary immunology: just what is the cost of
immunity? *Oikos* 88(1). Wiley: 87–98.

Lopez-Antia A, Ortiz-Santaliestra ME, Mougeot F, et al. (2013) Experimental exposure of red-legged
partridges (Alectoris rufa) to seeds coated with imidacloprid, thiram and difenoconazole.
Ecotoxicology 22(1). Springer: 125–138.

Lopez-Antia A, Ortiz-Santaliestra ME, Mougeot F, et al. (2015) Imidacloprid-treated seed ingestion has
lethal effect on adult partridges and reduces both breeding investment and offspring immunity.
Environmental research 136. Elsevier: 97–107.

Lupi S, Maggini I, Goymann W, et al. (2017) Effects of body condition and food intake on stop-over
decisions in Garden Warblers and European Robins during spring migration. *Journal of ornithology*
/ DO-G 158(4). Springer: 989–999.

Ma C, Vander Zanden HB, Wunder MB, et al. (2020) assignR : An r package for isotope-based
geographic assignment. *Methods in ecology and evolution / British Ecological Society* 11(8). Wiley:
996–1001.

Marra PP, Francis CM, Mulvihill RS, et al. (2005) The influence of climate on the timing and rate of
spring bird migration. *Oecologia* 142(2): 307–315.

Martínez-Renau E, Rojas-Estévez N, Friis G, et al. (2022) Haemosporidian parasite diversity and
prevalence in the songbird genus Junco across Central and North America. *Ornithology* 139(3).

566   Oxford Academic: ukac022.

567 Mason R, Tennekes H, Sánchez-Bayo F, et al. (2013) Immune suppression by neonicotinoid insecticides
568   at the root of global wildlife declines. *J Environ Immunol Toxicol* 1(1): 3–12.

569 Merrill L, Levengood JM, England JC, et al. (2018) Blood parasite infection linked to condition of
570   spring-migrating Lesser Scaup (Aythya affinis). *Canadian journal of zoology* 96(10). Canadian
571   Science Publishing: 1145–1152.

572 Møller AP (1994) Phenotype-dependent arrival time and its consequences in a migratory bird. *Behavioral*
573   *ecology and sociobiology* 35(2). Springer Nature: 115–122.

574 Mukhin A, Palinauskas V, Platonova E, et al. (2016) The Strategy to Survive Primary Malaria Infection:
575   An Experimental Study on Behavioural Changes in Parasitized Birds. *PloS one* 11(7).
576   journals.plos.org: e0159216.

577 Nakagawa S and Schielzeth H (2013) A general and simple method for obtaining R2 from generalized
578   linear mixed-effects models. *Methods in ecology and evolution / British Ecological Society* 4(2).
579   Wiley Online Library: 133–142.

580 Newton I (2010) *The Migration Ecology of Birds*. Elsevier.

581 Nolan V Jr and Ketterson ED (1990) Timing of autumn migration and its relation to winter distribution in
582   dark-eyed juncos. *Ecology* 71(4). Wiley: 1267–1278.

583 Nussbaumer R, Schmid B, Bauer S, et al. (2022) Favorable winds speed up bird migration in spring but
584   not in autumn. *Ecology and evolution* 12(8). Wiley Online Library: e9146.

585 Owen JC and Moore FR (2008) Relationship between energetic condition and indicators of immune
586   function in thrushes during spring migration. *Canadian journal of zoology* 86(7). Canadian Science
587   Publishing: 638–647.

588 Price ER (2010) Dietary lipid composition and avian migratory flight performance: Development of a
589   theoretical framework for avian fat storage. *Comparative biochemistry and physiology. Part A,*
590   *Molecular & integrative physiology* 157(4). Elsevier: 297–309.

591 Prop J, Black JM and Shimmings P (2003) Travel schedules to the high arctic: barnacle geese trade-off
592   the timing of migration with accumulation of fat deposits. *Oikos* 103(2). Wiley: 403–414.

593 Pulgarín-R PC, Gómez C, Bayly NJ, et al. (2019) Migratory birds as vehicles for parasite dispersal?
594   Infection by avian haemosporidians over the year and throughout the range of a long-distance
595   migrant. *Journal of biogeography* 46(1). Wiley: 83–96.

596 Ramenofsky M (1990) Fat storage and fat metabolism in relation to migration. *Bird migration:*
597   *physiology and ecophysiology*. Springer-Verlag New York: 214–231.

598 Reed KD, Meece JK, Henkel JS, et al. (2003) Birds, migration and emerging zoonoses: west nile virus,
599   lyme disease, influenza A and enteropathogens. *Clinical medicine & research* 1(1): 5–12.

600 Richardson WJ (1978) Timing and Amount of Bird Migration in Relation to Weather: A Review. *Oikos*
601   30(2). [Nordic Society Oikos, Wiley]: 224–272.

602 Ricklefs RE, Medeiros M, Ellis VA, et al. (2017) Avian migration and the distribution of malaria

parasites in New World passerine birds. *Journal of biogeography* 44(5). Wiley: 1113–1123.

Rotics S, Kaatz M, Turjeman S, et al. (2018) Early arrival at breeding grounds: Causes, costs and a trade-
off with overwintering latitude. *The Journal of animal ecology* 87(6): 1627–1638.

Rubenstein DR and Hobson KA (2004) From birds to butterflies: animal movement patterns and stable
isotopes. *Trends in ecology & evolution* 19(5): 256–263.

Saino N, Rubolini D, Jonzén N, et al. (2007) Temperature and rainfall anomalies in Africa predict timing
of spring migration in trans-Saharan migratory birds. *Climate Research* 35. Inter-Research Science
Center: 123–134.

Santiago-Alarcon D, Mettler R, Segelbacher G, et al. (2013) Haemosporidian parasitism in the
blackcapSylvia atricapillain relation to spring arrival and body condition. *Journal of avian biology*
44(6). Wiley: 521–530.

Satterfield DA, Marra PP, Sillett TS, et al. (2018) Responses of migratory species and their pathogens to
supplemental feeding. *Philosophical transactions of the Royal Society of London. Series B,*
Biological sciences 373(1745).

Sauer PE, Schimmelmann A, Sessions AL, et al. (2009) Simplified batch equilibration for D/H
determination of non-exchangeable hydrogen in solid organic material. *Rapid communications in*
mass spectrometry: RCM 23(7). Wiley: 949–956.

Schimmelmann A (1991) Determination of the concentration and stable isotopic composition of
nonexchangeable hydrogen in organic matter. *Analytical chemistry* 63(21). American Chemical
Society: 2456–2459.

Schimmelmann A, Lewan MD and Wintsch RP (1999) D/H isotope ratios of kerogen, bitumen, oil, and
water in hydrous pyrolysis of source rocks containing kerogen types I, II, IIS, and III. *Geochimica et*
cosmochimica acta 63(22). Elsevier: 3751–3766.

Schmaljohann H, Liechti F and Bruderer B (2009) Trans-Sahara migrants select flight altitudes to
minimize energy costs rather than water loss. *Behavioral ecology and sociobiology* 63(11). Springer:
1609–1619.

Schulte-Hostedde AI, Zinner B, Millar JS, et al. (2005) Restitution of mass–size residuals: Validating
body condition indices. *Ecology* 86(1). Wiley: 155–163.

Serra-Cobo J, Sanz-Trullén V and Martínez-Rica JP (1998) Migratory movements of Miniopterus
schreibersii in the north-east of Spain. *Acta theriologica* 43(3). rcin.org.pl: 271–283.

Sinelschikova A, Kosarev V, Panov I, et al. (2007) The influence of wind conditions in Europe on the
advance in timing of the spring migration of the song thrush (Turdus philomelos) in the south-east
Baltic region. *International journal of biometeorology* 51(5). Springer: 431–440.

Singh D, Reed SR, Kimmitt AA, et al. (2019) Breeding at higher latitude as measured by stable isotope is
associated with higher photoperiod threshold and delayed reproductive development in a songbird.
bioRxiv. Available at: https://www.biorxiv.org/content/10.1101/789008 (accessed 22 October 2023).

Slowinski SP, Fudickar AM, Hughes AM, et al. (2018) Sedentary songbirds maintain higher prevalence
of haemosporidian parasite infections than migratory conspecifics during seasonal sympatry. *PloS*
one 13(8): e0201563.

Smith RJ and Moore FR (2005) Arrival timing and seasonal reproductive performance in a long-distance migratory landbird. *Behavioral ecology and sociobiology* 57(3). Springer Science and Business Media LLC: 231–239.

Stott P (2016) How climate change affects extreme weather events. *Science* 352(6293). science.org: 1517–1518.

Studds CE and Marra PP (2005) Nonbreeding habitat occupancy and population processes: An upgrade experiment with a migratory bird. *Ecology* 86(9). Wiley: 2380–2385.

Talbott KM, Becker DJ, Soini HA, et al. (2022) Songbird preen oil odour reflects haemosporidian parasite load. *Animal behaviour* 188. Elsevier: 147–155.

Taylor P, Crewe T, Mackenzie S, et al. (2017) The Motus Wildlife Tracking System: a collaborative research network to enhance the understanding of wildlife movement. *Avian Conservation and Ecology/Ecologie et Conservation des Oiseaux* 12(1). The Resilience Alliance.

Therneau TM and Grambsch PM (2000) *Modeling Survival Data: Extending the Cox Model*. Springer US.

Thorsen S (1995-2023) Past Weather in Bloomington, Indiana, USA — March 2020. Available at: https://www.timeanddate.com/ (accessed 1 November 2022).

Tonra CM, Marra PP and Holberton RL (2011) Migration phenology and winter habitat quality are related to circulating androgen in a long-distance migratory bird. *Journal of avian biology* 42(5). Wiley: 397–404.

Tøttrup AP, Rainio K, Coppack T, et al. (2010) Local temperature fine-tunes the timing of spring migration in birds. *Integrative and comparative biology* 50(3). academic.oup.com: 293–304.

Usui T, Butchart SHM and Phillimore AB (2017) Temporal shifts and temperature sensitivity of avian spring migratory phenology: a phylogenetic meta-analysis. *The Journal of animal ecology* 86(2). Wiley: 250–261.

Valkiunas G, Iezhova TA, Krizanauskiene A, et al. (2008) A comparative analysis of microscopy and PCR-based detection methods for blood parasites. *The Journal of parasitology* 94(6): 1395–1401.

Van Doren BM and Horton KG (2018) A continental system for forecasting bird migration. *Science* 361(6407). science.org: 1115–1118.

Visser ME, Perdeck AC, van BALEN JH, et al. (2009) Climate change leads to decreasing bird migration distances. *Global change biology* 15(8). Wiley: 1859–1865.

Wanamaker SM, Singh D, Byrd AJ, et al. (2020) Local adaptation from afar: migratory bird populations diverge in the initiation of reproductive timing while wintering in sympatry. *Biology letters* 16(10): 20200493.

Wang X, Anadón A, Wu Q, et al. (2018) Mechanism of Neonicotinoid Toxicity: Impact on Oxidative Stress and Metabolism. *Annual review of pharmacology and toxicology* 58: 471–507.

Weber TP, Houston AI and Ens BJ (1994) Optimal departure fat loads and stopover site use in avian migration: an analytical model. *Proceedings. Biological sciences / The Royal Society* 258(1351). The Royal Society: 29–34.

Wikelski M, Tarlow EM, Raim A, et al. (2003) Costs of migration in free-flying songbirds. *Nature*
423(6941). Nature Publishing Group: 704–704.

Wilcove DS and Wikelski M (2008) Going, going, gone: is animal migration disappearing. *PLoS biology*
6(7). journals.plos.org: e188.

Witter MS and Cuthill IC (1993) The ecological costs of avian fat storage. *Philosophical transactions of*
the Royal Society of London. Series B, Biological sciences 340(1291). royalsocietypublishing.org:
73–92.

Wood SN (2017) *Generalized Additive Models: An Introduction with R, Second Edition*. CRC Press.

Wunder MB (2010) Using isoscapes to model probability surfaces for determining geographic origins.
Isoscapes. Springer. Epub ahead of print 2010.

Yorinks N and Atkinson CT (2000) Effects of Malaria on Activity Budgets of Experimentally Infected
Juvenile Apapane (Himatione Sanguinea). *The Auk* 117(3). Oxford Academic: 731–738.

Zhang J and Wu L (2018) The influence of population movements on the urban relative humidity of
Beijing during the Chinese Spring Festival holiday. *Journal of cleaner production* 170. Elsevier:
1508–1513.

Zhang X, Wan H, Zwiers FW, et al. (2013) Attributing intensification of precipitation extremes to human
influence. *Geophysical research letters* 40(19). American Geophysical Union (AGU): 5252–5257.

<u>Figures</u>

Figure 1. Estimated breeding locations and migratory distances based on junco feather δ^2H. Orange

shading represents overlapping geographic breeding assignments of individuals. Estimates reflect cells

within the top 10% highest posterior probability (Bowen et al., 2014; Ma et al., 2020; Wunder, 2010).

Points on the map represent the centroid of each estimated breeding polygon, colored by distance to

location of capture (star; Bloomington, IN). Lines radiating from the star represent subsequent Motus

detections after departure. The seasonal breeding range for *Junco hyemalis(Baillie et al., 2004)* is shown

in light grey (the species' full range map was used to construct geographic assignment maps).

Figure 2. (A) Release and departure timing inferred from Motus at the Kent Farm Research Station for

all 37 tagged juncos. (B) Mean probabilities of spring departure as inferred from Kaplan–Meier survival

curves (i.e., 1 – survival probabilities) using the *survival* package alongside 95% confidence intervals.

712 **Figure 3.** Relative hazards of spring departure estimated from the most competitive CPHMM as selected

713 through AICc ($w_i = 0.61$). The predicted relative hazard and 95% confidence intervals from this model are

714 displayed for mean wind direction and weather PC1.

716 **Tables**

717 **Table 1**. Comparison of CPHMMs predicting junco risk of spring departure; all models include a random

718 effect of site. Candidate models are ranked by ΔAICc alongside Akaike weights (w_i) and deviance

719 explained (DE).

Fixed effects	ΔAICc	w_i	DE
~ s(wind direction) + s(weather PC1)	0.00	0.61	31%
~ s(wind speed) + s(wind direction) + ti(wind speed, wind direction)	2.09	0.22	3%
~ s(wind speed) + s(wind direction) + s(migration distance)	2.69	0.16	4%
~ s(intensity) + s(weather PC1) + ti(intensity, weather PC1)	8.04	0.01	20%
~ s(intensity) + s(HL ratios) + s(weather PC1)	12.71	<0.01	10%
~ s(intensity) + s(total leukocytes) + ti(intensity, total leukocytes)	26.54	<0.01	25%
~ s(body condition) + s(migration distance)	27.87	<0.01	5%
~ s(fat score) + s(migration distance)	27.87	<0.01	5%
~ s(intensity) + s(total leukocytes) + s(HL ratios)	28.15	<0.01	4%
~ s(intensity) + s(HL ratios) + ti(intensity, HL ratios)	28.15	<0.01	4%

720