N-Glycans on the extracellular domain of the Notch1 receptor control Jagged-1 induced Notch signalling and myogenic differentiation of S100β resident vascular stem cells

Eoin Corcoran[1], Abidemi Olayinka[1], Mariana di Luca[1], Yusof Gusti[1],

Roya Hakimjavadi[1], Brendan O'Connor[2], Eileen M. Redmond[3] and Paul A. Cahill[1]*

[1] Vascular Biology and Therapeutics Laboratory, School of Biotechnology Faculty of Science and Health, Dublin City University, Dublin, Ireland.

[2] School of Biotechnology Faculty of Science and Health, Dublin City University, Dublin, Ireland

[3] Department of Surgery, University of Rochester, 601 Elmwood Ave, Rochester, NY 14642, United States

Abstract

Notch signalling, critical for development and postnatal homeostasis of the vascular system, is highly regulated by several mechanisms including glycosylation. While the importance of O-linked glycosylation is widely accepted, the structure and function of N-glycans has yet to be defined. Here, we take advantage of lectin binding assays in combination with pharmacological, molecular, and site-directed mutagenetic approaches to study N-glycosylation of the Notch1 receptor. We find that several key oligosaccharides containing bisecting or core fucosylated structures decorate the receptor, control expression and receptor trafficking, and dictate Jagged-1 activation of Notch target genes and myogenic differentiation of multipotent S100β vascular stem cells. N-glycans at asparagine (N) 1241 and 1587 protect the receptor from accelerated degradation, while the oligosaccharide at N888 directly affects signal transduction. Conversely, N-linked glycans at N959, N1179, N1489 do not impact canonical signalling but inhibit differentiation. Our work highlights a novel functional role for N-glycans in controlling Notch1 signalling and differentiation of vascular stem cells.

Teaser

A sweet development in Notch regulation of vascular smooth muscle cell differentiation

Introduction

Notch signalling is crucial for the development and postnatal homeostasis of the vascular system [1]. It has been implicated in cardiovascular disease, cancer, and neurological and

kidney disease. The Notch pathway is activated by interaction of receptors (Notch1–4) and ligands (Delta-like 1, 3, 4, and Jagged 1, 2) present on the surface of adjacent cells [3]. This interaction triggers sequential cleavage events resulting in release of the active form of Notch i.e., Notch intracellular domain (NICD). NICD translocates into the nucleus and, in combination with the transcription factor recombinant binding protein for the immunoglobulin region kJ (RBPJ, also known as CBF1/Suppressor of Hairless/LAG1, CSL) and the Mastermind-like (MAML 1–3) adaptor proteins, activates target genes including HES (Hairy and Enhancer of Split) and HEY (Hairy/enhancer-of-split related) that are essential modulators of neuronal, endocrine, and cardiovascular cell fate [4]. The Notch pathway may also be activated in the absence of 'canonical' ligands and independent of γ-secretase proteolytic cleavage; this non-canonical Notch signalling modulates Wnt/β-catenin, mammalian target of the rapamycin 2 complex (mTORC2)/Akt, Nuclear Factor kappa B (NF-κB), IkB kinase (IKK)-α/β and phosphatase and tensin homolog (PTEN)-induced kinase (PINK1) [5].

Notch impacts arterial pathology including atherosclerotic plaque development [6–10]. There is compelling evidence for the involvement of (i) differentiated medial vascular smooth muscle cells (SMC) that are stem cell antigen-1 positive (Sca1+) [11-12] and (ii) various adventitial [13-14] and medial progenitors [15–18] [19-20] in progressing vascular lesion formation through the accumulation of 'SMC-like' cells. Notch signalling decreases SMC differentiation marker expression [21] and promotes SMC-like cell accumulation within lesions [10]. Multiple studies describe the ability of Notch signalling to induce myogenic differentiation of stem cell populations such as C3H 10T1/2 embryonic fibroblasts, mesenchymal stromal stem cells and neural crest stem cells [22–24]. While the exact mechanism and cell(s) involved by which Notch and its target genes affect lesion formation remains incompletely understood, there is substantial evidence suggesting that the Notch1 receptor in particular [8, 10], plays a pivotal role in this process.

The Notch signalling pathway is tightly regulated by several mechanisms including cis-inhibition, negative feedback loops and PEST ubiquitination [25, 26] and by glycosylation of the Notch receptor [27]. There are several highly conserved glycosylation sites along the extracellular domain of the Notch receptor, most of which harbour O-linked glycans. Conserved sites of O-glycosylation on Notch were first discovered at the turn of the century in Drosophila melanogaster [28] and this gave rise to a wave of research centred on O-glycosylation of Notch, resulting in many more consensus sites and several important glycosyltransferases, such as Fringe, being identified [29, 30]. In contrast, very little is known about the specifics of N-linked glycosylation of Notch. It has been established that there are six N-glycosylation consensus sites along the extracellular domain of Notch [31] and that N-glycans are certainly present on the Notch receptor [34] but the number of these sites that are glycosylated, the structures of these N-glycans and their potential role in Notch signalling, remains unknown.

The impact of N-glycan decoration of key proteins on a given process can be revealed using chemical inhibitors of N-glycosylation. Tunicamycin (TNC), an antagonist for the GlcNAc phosphotransferase (GPT) enzyme, catalyses a critical step in the protein N-glycosylation pathway, linking the growing polypeptide to a dolichol precursor [32]. 1-deoxymannojirimycin (DMJ) blocks the trimming of the 14-sugar dolichol precursor oligosaccharide by inhibiting the mannosidase I enzyme and preventing the synthesis of complex or hybrid-type N-glycans [33]. Because the specificities of these inhibitors are broad, a more targeted approach for regulating N-glycans can be achieved using siRNA

92	knockdown of key glycosyltransferases, Fut8 and Mgat3. A family of glycosyltransferases
93	known as N-acetylglucosaminyl transferases (GnT's) are responsible for the addition of
94	GlcNAc residues to N-linked glycans. There are six members of this family (GnT-I to
95	GnT-VI), all of which operate in the Golgi apparatus [34]. GnT-III (coded by the Mgat3
96	gene) catalyses the addition of a GlcNAc residue through a β-1,4 linkage to the β-mannose
97	in the tri-mannosyl core of a growing N-linked oligosaccharide, creating a structure
98	known as a bisecting N-glycan [35]. This structure has been shown to inhibit the addition
99	of sugar residues by other glycosyltransferases such as GnT-II, GnT-IV and Fut8 [36, 37].
100	Bisecting N-glycans are heavily involved in the regulation of several cellular processes,
101	including cell adhesion, protein degradation and growth factor signalling [38]. Members
102	of the fucosyltransferase (FUT) family of enzymes facilitate the addition of fucose
103	residues to the core N-glycan structure [39] in a process known as core fucosylation. Fut8
104	catalyses the transfer of fucose from a donor substrate to the asparagine-149 bound
105	GlcNAc at the base of an N-glycan via an α1,6-linkage [40]. Core-fucosylated N-glycans
106	regulate a wide array of cellular processes such as the expression of essential cadherins
107	and the function of specific types of immunoglobulins [41-42]. As of yet, it has not been
108	established whether N-glycans on the Notch1 receptor are core fucosylated, though it has
109	been noted that aberrant activity of both Fut8 and Notch is regularly observed in cancer
110	[43].

111	Of note, lectins have become powerful tools to determine the N-glycan profile of a
112	glycoprotein, as they are highly specific carbohydrate-binding proteins with no catalytic
113	activity [44]. They can bind mono- or oligosaccharides and can recognise and distinguish
114	between distinct branching and linkage patterns [45]. Several assays have been developed
115	and optimised which exploit the high level of specificity seen in lectin-oligosaccharide
116	binding to determine the glycosylation profile of a given glycoprotein sample. The
117	enzyme-linked lectin assay (ELLA) is a simple and rapid method of glycan analysis that,
118	unlike techniques such as mass spectrometry and HPLC, does not require the chemical or
119	enzymatic removal of carbohydrates from proteins [46]. In this way, lectins can be used to
120	detect changes in N-glycosylation patterns following inhibitory interventions.

121	The aim of this study was to characterise the profile of N-glycosylation of the Notch1
122	receptor and explore the impact of these N-glycans on Notch signalling, receptor
123	trafficking and Notch-mediated myogenic differentiation of stem cells *in vitro*.

Results

Jagged-1 (Jag-1) activation of the Notch1 receptor promotes myogenic differentiation

127	The phenotype of mouse aortic vascular stem cells (mVSCs) and C3H10T1/2 embryonic
128	fibroblasts were assessed by immunocytochemistry. Cultured mouse smooth muscle cells
129	(SMCs) (MOVAS-SMCs) served as a positive control for myogenic SMC differentiation
130	marker expression. The mVSCs and C3H 10T1/2 cells were both positive for S100β and
131	Sca1 expression, while negative for SMC markers, Cnn1 and Myh11 [Suppl Figure 1(a),
132	Figure 1(a)]. The stemness of isolated S100β mVSCs was confirmed by demonstrating
133	increased telomere length and telomerase activity when compared to aortic tissue and
134	cultured SMCs [Suppl Figure 1(b), (c)]. Adipogenic (*Lpl* and *Fabp4*) and osteogenic gene
135	expression (*Alpl* and *Sost*) was significantly increased following treatment of S100β
136	mVSCs with lineage specific differentiation media, when compared to the maintenance
137	media control [Suppl Figure 1(d), (e)]. These data confirm the phenotype and stemness of

the S100β mVSCs as distinct from freshly isolated aortic SMCs and aortic SMCs in
culture.

Immobilised recombinant Notch ligand, Jagged-1 increased the number of SMC-
like/myogenic Cnn1 positive cells (40.5 ± 5%, p<0.05 n=6) in S100β mVSCs cultures,
when compared to the Fc-IgG control (0.1 ± 0.4%) [Figure 1(b), (c)] concomitant with a
significant increase in the expression of Notch target gene, *Hey1* and the SMC
differentiation marker gene, *Myh11*, an effect attenuated following pre-treatment of the
cells with the γ-secretase inhibitor, DAPT [Figure 1(d),(e)]. As the Notch1 receptor has
been implicated in arteriosclerotic lesion formation [8, 10, 49], the effect of Notch1
receptor depletion following siRNA knockdown on Jagged-1 induced *Hey1* and *Myh11*
expression was also assessed in S100β mVSCs [Figure 1(g), (h)]. The transfection
efficiency of the siRNA's was confirmed at > 80% under optimised conditions using a
fluorescently- tagged siRNA (TYE 563 DsiRNA) [Suppl Figure 2(a)]. *Notch1* mRNA
levels [Suppl Figure 2(b)] and Notch1 receptor protein levels [Suppl Figure 2(c)] were
significantly reduced when compared to the scrambled control concomitant with a
significant inhibition of Jagged-1 induced *Hey1* [Figure 1(d)] and *Myh11* [Figure 1(e)]
expression.

These data confirm that multipotent S100β mVSCs and C3H10T1/2 cells respond to
Notch ligand-induced activation of Notch signalling to promote myogenic differentiation
of these cells via activation of the Notch1 receptor.

Chemical inhibition of N-glycosylation impairs Jagged-1 induced Notch signalling

N-glycan analysis was first assessed in whole cell lysates before immunoprecipitated
Notch1 was prepared, and lectin binding evaluated using a series of commercial lectins
(Table I). Ectopic overexpression of a plasmid-encoded Notch1 was achieved using cells
transfected with either *pCS2 Notch1 Full Length-6MT* or *pCS2 Notch1 ICv-6MT* plasmids
[Suppl Figure 3(a)]. Samples were immuno-probed using an anti-myc-tag antibody and
stained for either the myc-tagged Notch1 or myc-tagged NICD, or an anti-β-actin
antibody, which served as the loading control [Suppl Figure 3(b)].

TNC and DMJ are both chemical inhibitors of N-glycosylation [32-33]. Cells were treated
with different concentrations of TNC (0.3 - 0.7μg/mL) or DMJ (2.0 - 4.0μg/mL) before
lectin binding was assessed in whole cell lysates using the ELLA assay. The binding
characteristics of each lectin have previously been described (Table I, [46]). There was a
dose-dependent decrease in absorbance, indicating a reduction in lectin binding, following
treatment of S100β mVSCs with TNC when compared to the non-treated control [Figure
2(a)]. Maximum inhibition was observed at 0.5μg/mL (in the cases of WGA and E-PHA).
Similarly, DMJ treatment caused a significant reduction in binding across all lectins
compared to the non-treated control with maximal inhibition observed at 4.0μg/mL
[Figure 2(c)]. A similar dose-dependent decrease in lectin binding was observed in whole
lysates following treatment of C3H10T1/2 cells with TNC [Suppl Figure 4(a)] and DMJ
[Suppl Figure 4(b)], but only in WGA and E-PHA. No significant change in binding was
detected at any DMJ concentration with ConA or LCA [Suppl Figure 4(b)]. These data
confirm the optimal concentrations of TNC and DMJ required to reduce lectin binding in
whole cell lysates. After establishing the effect of TNC and DMJ in whole cell lysates,
their effect on the N-glycan profile of the immunoprecipitated Notch1 receptor was
assessed.

Immunoprecipitated Notch1 was prepared following optimisation of the immunoprecipitation protocol from whole cell lysates [Suppl Figure 2(d)]. Lectin binding was significantly reduced in the immunoprecipitated Notch1 preparation from S100β mVSCs treated with both TNC [Figure 2(b)] and DMJ [Figure 2(d)] across all lectins other than WGA. Inhibition by DMJ was as pronounced as that of TNC. The repercussions of removing these N-glycans from the Notch1 receptor on Jagged 1-induced Notch signalling and myogenic differentiation of S100β mVSCs and C3H10T1/2 cells was assessed. Increased *Hey1* and *Myh11* expression following treatment with recombinant Jagged-1, when compared to the control IgG-Fc, was significantly reduced in TNC [Figure 2(e),(f)] and DMJ [Figure 2(g),(h)] treated S100β mVSCs. In a similar manner, recombinant Jagged-1 significantly increased *Hey1* and *Myh11* expression in C3H10T1/2 cells when compared to IgG-Fc, an effect that was significantly reduced in TNC treated [Suppl Figure 4(d), (e)] and DMJ treated cells [Suppl Figure 4(f), (g)]. Interestingly, TNC treatment also led to a significant decrease in *Hey1* expression in the Fc-treated control sample [Suppl Figure 4(d)]. The decrease in *Hey1* expression following treatment of S100β mVSCs with TNC and DMJ was not due to changes in the expression of the Notch1 receptor as there was no significant change in either *Notch1* transcript levels or Notch1 protein expression following TNC or DMJ treatment, compared to the non-treated control [Suppl Figure 3(c), (d)].

Notch intracellular domain (NICD) can activate Notch signalling independent of ligand binding; this function was exploited to determine whether TNC and DMJ affected key Notch signalling downstream of receptor activation. Ectopic expression of the Notch1 NICD [Suppl Figure 3(a), (b)] significantly increased the expression of the Notch target gene *Hey1* in S100β mVSCs [Figure 3(a), (c)]. There was no significant change in this response following TNC or DMJ treatment of these cells [Figure 3(a), (c)]. Similarly, expression of the myogenic marker *Myh11* was increased following ectopic expression of NICD [Figure 3(b), (d)]. However, this response was significantly attenuated by treatment with either TNC or DMJ [Figure 3(b), (d)). These data suggest that Notch pathway elements between the cleavage of the NICD and the activation of *Hey1* gene are unaffected by TNC or DMJ, but components of the myogenic differentiation pathway are altered by these treatments.

Collectively, these data show that inhibition of global glycosyltransferase activity with TNC and DMJ reduced the level of N-glycan structures on cellular glycoproteins within whole lysates from S100β mVSCs and C3H10T1/2 cells. This includes Notch1 as both inhibitors had a significant effect on the N- glycosylation profile of the purified immunoprecipitated receptor, supporting the presence of N-glycans on Notch1. Removal of these N-glycans caused a significant reduction in ligand-induced Notch signalling and myogenic differentiation in these cells. As downstream components of Notch signalling and myogenic differentiation pathways following ectopic expression of Notch1 NICD were not altered by *TBC* or *DMJ*, their effect appears to be primarily at the level of the Notch1 receptor.

GnT-III and Fut8 depletion impairs Jagged-1 induced Notch signalling

Mgat3 and *Fut8* genes code for the glycosyltransferases GnT-III and Fut8, respectively. To deplete their levels, targeted siRNA knockdown was performed in S100β mVSCs [Suppl Figure 3(e), (f)] and C3H10T1/2 cells [Suppl Figure 5(a), (b)], respectively, before the N-glycan profile of whole lysates [Figure 4(a)] and immunoprecipitated Notch1

[Figure 4(b)] was assessed by lectin binding using the ELLA assay. There was a
significant reduction in lectin binding of whole lysates following *Mgat3* knockdown for
ConA and E-PHA lectins in both S100β mVSCs [Figure 4(a)] and C3H10T1/2 cells
[Suppl Figure 4(c)] compared to the scrambled control, while no significant reduction in
binding was detected with the LCA or WGA lectins. In the case of the *Fut8* knockdown,
this led to a significant decrease in LCA binding in both cell lines, while any changes in
binding with the other three lectins was statistically insignificant [Figure 4(a), Suppl
Figure 4(c)). In parallel cultures, *Mgat3* knockdown in S100β mVSCs resulted in a
significant reduction in lectin binding of immunoprecipitated Notch1 with both the ConA
and E-PHA lectins compared to the scrambled control, while no significant reduction in
binding was detected with LCA or WGA lectins [Figure 4(b)]. Meanwhile, *Fut8*
knockdown led to significant decreases in ConA and LCA binding, but not for WGA or E-
PHA lectins [Figure 4(b)].

The repercussions of *Mgat3* and *Fut8* knockdown on Jagged-1 induced Notch signalling
and myogenic differentiation revealed a significant reduction in Jagged-1 stimulated *Hey1*
and *Myh11* expression in S100β mVSCs and C3H10T1/2 cells following both *Mgat3*
knockdown [Figure 4(c), (d), Suppl Figure 5(c), (d)] and *Fut8* knockdown, respectively
[Figure 4(e), (f), Suppl Figure 5(e), (f)], when compared to the scrambled controls. In
contrast, Notch1 NICD induced Notch signalling and myogenic differentiation of S100β
mVSCs was unaffected by *Mgat3* [Figure 5(a), (b)] and *Fut8* knockdown [Figure 5(c),
(d)], when compared to the empty vector (EV) controls. Moreover, the decrease in *Hey1*
expression following *Fut8* knockdown in S100β mVSCs was not due to changes in the
expression of the Notch1 receptor as there was no significant change in either Notch1
protein expression [Figure 5(e, f)] or *Notch1* mRNA levels [Figure 5(g)] following
knockdown, compared to scrambled control. In contrast, while there was a slight albeit
significant change in the expression of Notch 1 protein levels following *Mgat3*
knockdown [Figure 5(e), (f)] this was not reflected in changes to *Notch1* mRNA levels
[Figure 5(g)].

Inhibition of N-glycosylation impairs Notch receptor trafficking and degradation.

N-glycosylation has been shown to be an essential modification for correct protein
trafficking from the endoplasmic reticulum (ER) and Golgi apparatus (GA) to their site of
action within the cell [50-51]. Therefore, the possibility that inhibition of N-glycosylation
by either chemical inhibitor (TNC or DMJ) or knockdown of *Mgat3* and *Fut8* interferes
with Notch1 protein trafficking and localisation to the membrane was explored.
Membrane fractionation of crude lysate from S100β mVSCs revealed the presence of
distinct protein band profiles between each fraction, confirmed using pan-cadherin, a
ubiquitously expressed membrane-bound protein [Figure 6(a)]. Fractionation revealed that
the transmembrane Notch1 protein is exclusively present in the membrane fraction and
absent in the cytosolic fraction [Figure 6(a)]. Following treatment of the cells with TNC
or DMJ for 48 h, there was a significant reduction in the expression of Notch1 in the
membrane fraction [Figure 6(b), (d)]. Brefeldin A (BFA), as a potent inhibitor of protein
transport in the Golgi apparatus, served as a positive control. Since cellular Notch1
protein levels remained unchanged following treatment with TNC and DMJ [Suppl Figure
3(c), (d)], it is unlikely that TNC and DMJ treatment was causing endocytosis and
degradation of the receptor [Figure 6(b)]. This observation was further supported by
immunocytochemical analysis of Notch1 localisation *in situ*. Golgin97 served as a control
to visualise the subcellular location of the GA [Figure 7(a),(c)]. The ratio of both the

fluorescence intensity and density significantly decreased following treatment with TNC, DMJ or BFA [Figure 7(c)] wherein a significant decrease in the ratio of membrane:perinuclear localisation of Notch1 was observed following treatment [Figure 7(a), (c)]. Lower ratios indicate a higher level of fluorescence in the perinuclear region compared to that at the membrane. This analysis suggests an accumulation of Notch1 in the perinuclear region of the cell in response to TNC and DMJ treatment, similar to BFA [Figure 7(a), (c)] and supports a significant reduction of Notch1 protein trafficking following inhibition of N-glycosylation.

In a similar manner, membrane fractionation was used to determine whether knockdown of *Mgat3* and *Fut8* also reduced Notch1 localisation to the membrane. There was a significant reduction in the expression of Notch1 in the membrane fraction following *Mgat3* knockdown, compared to the scrambled control. In contrast, there was no significant change following *Fut8* knockdown [Figure 6(c), (e)]. This analysis suggests that Notch1 trafficking from the GA to the plasma membrane is inhibited by *Mgat3* knockdown. However, in parallel experiments, immunocytochemical analysis of Notch1 localisation *in situ* following knockdown of *Mgat3* and *Fut8* demonstrated that there was no significant change in the ratio of either the fluorescence intensity or density following knockdown whereas both ratios decreased significantly following treatment BFA [Figure 7(b), (d)]. Therefore, it is more likely that this reduction is due to the decrease in total Notch1 protein in the cell following *Mgat3* knockdown [Figure 5(e)].

In summary, knockdown of the *GnT-III* and *Fut8* glycosyltransferases provided a more specific insight into how two groups of N-glycans (bisected and core fucosylated) contributed to Notch signalling and myogenic differentiation. Knockdown of the *Mgat3* transcript (which codes for the GnT-III protein) led to a reduction in E-PHA binding, while *Fut8* knockdown resulted in a decreased level of LCA binding. These findings provide evidence for the presence of both complex bisecting and core fucosylated N-glycans on the Notch1 receptor. Moreover, knockdown of *GnT-III* and *Fut8* glycosyltransferases demonstrated that both groups of N-glycans play a major role in activation of both Notch target gene expression and myogenic marker expression following Jagged-1 activation of mVSCs.

Site-directed mutagenesis of six N-glycosylation impairs Jagged-1 induced Notch signalling

Specific targeting of individual glycans to further define the function of both bisecting and core fucosylated N-glycans was performed through site-directed mutagenesis of the six N-glycosylation consensus sequences along the Notch1 ECD. Site-directed mutagenesis of plasmid-encoded Notch1 was chosen, and the proportion of transfected cells was maximised in each population through puromycin selection [Suppl Figure 6(a), (b)]. The optimum concentration of plasmid DNA was assessed using a green fluorescent protein (GFP)-coding plasmid [Suppl Figure 6(b)]. Using the *pCS2 Notch1 Full Length-6MT* plasmid, six single base substitutions were carried out separately on the Notch1 gene to alter asparagine residues at amino acid positions 888, 959, 1179, 1241, 1489 and 1587 to isoleucine residues. Mutant plasmids were sequenced and aligned with the wild- type plasmid sequence using Multalin to confirm that the correct single base substitution had been incorporated in each case. In each case an essential adenine base was correctly converted to a thymine base [Suppl Figure 7(a)]. Chromatograms were also generated to

determine the quality of both the sequencing read and the sample plasmid and confirmed high quality with clearly defined peaks [Suppl Figure 7(b)].

Mutations at N888I, N1489I and N1587I all resulted in a significant decrease in ConA, LCA and E-PHA binding compared to that observed in the wild-type cells but did not significantly alter WGA binding. N959I led to a significant reduction only in ConA binding, while N1179I caused a significant attenuation in only the binding of LCA [Figure 8(a)]. In parallel cultures, Jagged-1 significantly increased the expression of *Hey1* in control wild-type Notch1 transfected cells, an effect that was significantly attenuated by mutations N888I, N1241I and N1587I of the Notch1 ECD [Figure 8(b)] concomitant with a significant reduction in Jagged-1 induced *Myh11* expression [Figure 8(b)].

The changes in *Hey1* or *Myh1*1 expression were not attributed to a discrepancy in efficiency of plasmid transfection across mutants as there was no significant difference between levels of Notch1-coding plasmid DNA across all mutant plasmid samples compared to the wild-type control [Suppl Figure 8(a)]. However, a significant decrease in Notch1 protein levels was observed in cells transfected with mutants N1241I and N1587I, respectively, while no significant change was detected across all other mutants, compared to the wild-type control [Figure 8(d)]

These data indicate that all the mutations incorporated into the Notch1 gene, except N1241, had a measurable effect on receptor glycosylation. Lectin binding indicated the presence of a range of different N-glycan structures on the Notch1 ECD, including core fucosylated and bisecting N-glycans, and some additional structures that could not be identified with the limited range of lectins used. Mutation of asparagine residues at Notch1 N-glycosylation sites reduced *Hey1* expression (N888I, N1241I and N1587I) and *Myh11* expression (all mutations) following Jagged-1 stimulation. This was not due to altered plasmid transfection efficiency, but for some mutants (N1241I and N1587I), altered expression and/or degradation of Notch1 may be involved. It is likely that the N-glycans at the four other mutation sites may influence Notch1 processing and signalling, such as receptor cleavage or trafficking.

Discussion

Glycosylation is an essential regulator of many cellular processes and Notch signalling is no different [52]. In contrast to the well documented role of O-glycosylation, very little is known about the structure and function of Notch1 N-linked glycans. Here we report on bisecting and core fucosylated N-glycans decorating the extracellular domain (ECD) of the Notch1 receptor and their putative role in ligand-induced Notch signalling and myogenic differentiation of resident vascular stem cells *in vitro*.

O-glycosylation is well established as influencing the folding, trafficking, activation, and cleavage of Notch1 [29, 53-55]. Although it has been shown that there are six highly conserved N-glycosylation sites on the Notch receptor [31], there is currently no data available to determine how many of these sites are glycosylated and what type of N-glycans reside on the receptor, other than data that suggests the Notch1 receptor contains complex bisecting N-glycans [56]. The Notch signalling pathway, and the Notch1 receptor in particular, plays an important role in lineage specification *in vitro* and

vascular lesion formation *in vivo* [10, 22]. Therefore, this study sought to characterise the profile of N-glycosylation of the Notch1 receptor and explore the impact of removing these N-glycans on Notch signalling and on Notch-mediated myogenic differentiation of S100β stem cells.

The Notch signalling pathway drives myogenic specification of a range of stem cells, including S100β resident mVSCs [22-24, 42, 57]. Here, immobilised recombinant Jagged-1 was used to activate the Notch receptor and stimulate the expression of both Notch target genes and myogenic markers in S100β mVSCs. Canonical Notch signalling operates via the NICD-RBP-Jκ complex which, along with other transcription factors, activates Notch target gene expression [3]. The NICD-RBP-Jκ complex is essential for maintaining SMC differentiation [48, 58] through interaction with the serum response factor (SRF)-myocardin transcription activator complex [59]. However, several studies have described the ability of Notch target genes, such as *Hey1* and *Hey2*, to inhibit myocardin-induced SMC differentiation, although many of these studies differ in their description of the mechanism of inhibition, with some citing a direct interaction between Notch target genes and SRF, while others suggest that this occurs via an independent parallel pathway [21, 60]. While it has been well documented that Notch target genes such as *Hey1* and *Hes1* contain RBP-Jκ binding domains at specific GTGGGAA consensus sequences within their promoter regions [61-62], bioinformatic analysis of the *Myh11* and *Cnn1* promoter regions suggests this consensus sequence is also present within their promoter [63]. Moreover, CBF1 (the human form of RBP-Jκ) interacts with SMAD2/3 in the TGF-β signalling pathway to promote transcription of SMC genes independent of myocardin, such as *Cnn1* and *SM22α,* albeit this interaction was not observed with Notch1 [63]. While NICD induced Notch signalling in S100β mVSCs promotes Notch target gene and myogenic gene expression in S100β mVSCs, the Jagged-1 response was dependent on canonical Notch signalling as the γ-secretase inhibitor, DAPT effectively abrogate this response.

Mammalian cells express four different types of Notch receptor (Notch1 - 4) and each one can initiate downstream Notch signal transduction in target cells [64]. Notch1 plays a role in the process of arterial remodelling and lesion formation [8, 10, 49]. Here, knockdown of *Notch1* mRNA in mVSCs using siRNA significantly attenuated Jagged-1-induced Notch target gene and myogenic gene expression confirming the pivotal role of Notch1 in lineage specification in these cells. Future studies that selectively delete the Notch1 receptor in resident S100β mVSCs using *S100β-CreERT2* transgenic mice crossed with Notch1 floxed mice before vascular injury will reveal the importance of this data *in vivo,* in addition with studies that explore the role of the other three Notch receptors in this process. In this context, loss of Notch 2 in SMCs leads to protein profile changes during lesion formation but does not alter overall lesion morphology or cell proliferation [65]. Nevertheless, the current findings highlight the potential of Notch1 as a therapeutic target for controlling the accumulation of SMC-like cells from mVSCs in arteriosclerotic lesion progression.

To date, evidence regarding the profile of N-linked glycans on the Notch1 receptor suggests that six consensus sites are present and highly conserved along the ECD, and that at least some of the N-glycans at these sites contain a bisecting structure [31, 66]. In the current study, ELLA analysis of the purified Notch1 receptor confirmed the presence of N-glycans, and of bisecting N-glycans, whilst also revealing the existence of oligosaccharides with a fucose residue bound to the core GlcNAc by an α-1,6 linkage

(referred to as core fucosylation). Treatment with global N-glycosylation inhibitors tunicamycin (TNC) and 1-deoxymannojirimycin (DMJ) further supported these findings as each inhibitor promoted a reduction in binding of the E-PHA (specific for bisecting structures) and LCA (specific for core fucosylated structures) lectins to the purified Notch1 protein. However, many lectins have weak affinity for oligosaccharide structures other than their main binding target, including both E-PHA and LCA [67, 68]. To further confirm the type of structures bound by these lectins, a strategy of siRNA knockdown was implemented to inhibit the expression of the glycosyltransferases responsible for the formation of bisecting and core fucosylated N-glycans, namely GnT-III and Fut8, respectively.

Knockdown of the *Mgat3* and *Fut8* glycosyltransferase-coding genes reduced the level of N-glycan structures on cellular glycoproteins within whole lysates. This includes Notch1 as both knockdowns had a significant effect on bisecting or core fucosylated N-glycans of the purified IP receptor, supporting the TNC and DMJ inhibition studies on the presence of both bisecting and core fucosylated N-glycans decorating the Notch1 receptor. *Mgat3* and *Fut8* knockdown reduced the level of ligand mediated Notch signalling and myogenic differentiation in both mVSCs and C3H 10T1/2 cells. As downstream components of Notch signalling and myogenic differentiation pathways following ectopic expression of Notch1 NICD were not altered by *Mgat3* or *Fut8* knockdown, their effect appears to be primarily on the Notch1 receptor as Notch pathway components downstream of receptor cleavage were not affected by a reduction in synthesis of bisecting or core fucosylated N-glycans. Furthermore, as Notch1 transcript levels were not affected by either knockdown, but Notch1 protein expression was significantly reduced by *Mgat3* expression, these data suggests that bisecting N-glycans may have a role in preventing the degradation of the Notch1 receptor. The function of core fucosylated N-glycans is more difficult to decipher as reduction of their synthesis through *Fut8* knockdown had no effect on Notch1 protein expression or on receptor trafficking but caused a significant attenuation of Notch signalling and myogenic differentiation. There may be an alternative function for core fucosylated N-glycans on the Notch1 receptor, such as regulating cleavage events.

However, as with the chemical inhibition strategies used in this study, this siRNA knockdown strategy could also alter the N-glycosylation of proteins other than the Notch1 receptor and it is possible that this could influence Notch downstream signalling capacity. Indeed, nicastrin, for example, is part of the γ-secretase complex which cleaves the NICD from the Notch1 transmembrane domain (TMD) following receptor activation, and it contains bisecting N-glycans which may be crucial for its role in the cleavage process [66, 69]. Similarly, *Fut9* knockdown of core fucosylated N-glycans impairs the proliferation of S100β neural stem cells via inhibition of Musashi-1 (Msi-1), an RNA-binding protein and activator of the Notch signalling pathway, that binds to *Numb* mRNA and inhibits its translation. *Fut9* knockdown is coupled with increased Numb levels eventually resulting in the suppression of Notch signalling [70]. Therefore, the reduction in Notch target gene and myogenic gene expression following *Mgat3* and F*ut8* knockdown may have been due, at least in part, to removal of N-glycans (bisected and core fucosylated) from other key proteins impacting on Notch signalling in these cells. For this reason, it was important to further explore the mechanisms by which core fucosylated and bisecting N-glycans were affecting the Notch signalling process. Collectively, the data confirm a role for N-linked bisecting glycosylation in the ligand activation of Notch1 in S100β mVSCs. Our finding that inhibition of GnT-III blocks Notch activation suggests that targeting GnT-III or blocking bisecting glycosylation sites on the Notch1 receptor may lead to new

therapeutics targeting neointimal formation. Overall, the evaluation of GnT-III expression levels could also have clinical utility for stratifying patient cohorts that could benefit from therapies that target Notch1 signalling.

The impact of N-linked glycosylation on O-fucose glycosylation of Notch is not known. The N-linked sites within Notch1 ECD are in close proximity to O-fucose modified sites. O-Fucose glycosylation of Notch EGF repeats has been shown to create a code to either inhibit or activate Notch ligand-dependent activation [71]. It will be important in the future to determine how the bisecting glycan structures on the Notch1 receptor in mVSCs alter this O-fucose code. While the importance of O-linked glycosylation in Notch receptor-ligand binding and receptor processing and trafficking is well documented [58, 60], little information is available on the role of N-glycans in this context. While activation of Notch target genes by the ectopically expressed NICD was not affected by TNC and DMJ, components of the myogenic differentiation pathway downstream of Notch activation appeared altered leading to inhibition of that process. These data did not necessarily implicate Notch1 N-glycosylation directly in the process, as many other glycoproteins would be modified by the chemical inhibitors. However, further investigation into how TNC and DMJ attenuated the Jagged-1 response revealed that it was mediated by preventing Notch1 receptor trafficking out of the ER-Golgi complex to the plasma membrane. A similar effect on Notch 3, where TNC treatment resulted in accumulation of the receptor in the ER-Golgi complex has been reported [72]. While this may be due to changes in glycosylation of the numerous proteins involved in receptor trafficking, multiple previous reports have demonstrated that inhibiting the N-glycosylation of developing proteins can block their transport to the membrane [73–75]. Therefore, it is likely that some of the N-glycans on the extracellular domain of the Notch1 receptor are required for correct trafficking of the protein to the plasma membrane. Unlike TNC and DMJ treatment, knockdown of GnT-III and Fut8 did not affect downstream elements of either Notch signalling or myogenic differentiation pathways, indicating that bisecting and core fucosylated N-glycans exert their influence on the Jagged-1 response either at, or before, the point of Notch receptor activation. Analysis of receptor trafficking revealed that inhibition of the glycosyltransferase did not result in accumulation of Notch1 in the ER-Golgi complex, as was seen with chemical inhibition of N-glycosylation. This suggests that the inhibition of trafficking observed following TNC and DMJ treatment was either due to altered glycosylation of transport proteins or removal of N-glycans on the Notch1 receptor which do not contain bisecting or core fucosylated structures. Combination of receptor trafficking data and measurement of Notch1 protein levels revealed that bisecting N-glycans prevent the accelerated degradation of the Notch1 receptor. This supports previous studies showing Notch1 receptor colocalization with lysosomal degradation proteins following *Mgat3* knockdown [56]. The role of core fucosylated N-glycans may be more directly related to the activation of the Notch1 receptor as knockdown of the *Fut8* glycosyltransferase inhibited Notch signalling and myogenic differentiation but did not affect Notch1 expression, trafficking, degradation, or downstream signalling.

Site-directed mutagenesis of the six N-glycosylation consensus sequences along the Notch1 ECD further defined the function of both bisecting and core fucosylated N-glycans. While initial analysis indicated that the amino acid site N1241 may not be glycosylated, additional information indicates that it is more likely that the N-glycan at this site was not detected due to the limited range of lectins used and the low binding affinity of some lectins *in vitro* [76]. This is further emphasised by the fact that the N1179

mutation did not lead to a significant change in ConA binding, but did significantly reduce the level of LCA binding, indicating that a core fucosylated oligosaccharide may be present at this site which was not detected by ConA. Similarly, analysis of the N959 mutant indicated that an N-linked oligosaccharide is present at this site but does not contain structures required for binding to LCA, WGA or E-PHA. Meanwhile, the glycan at site N1179 was found to contain a core fucosylated structure. Finally, it is noteworthy that N-glycans at N888, N1489 and N1587 shared a similar lectin binding profile with significant reduction in the level of binding with ConA, LCA and E-PHA with these mutants, suggesting that glycans at those sites are both core fucosylated and have a bisecting structure. Therefore, it is possible that they share similar structural characteristics and that the N-glycans at those sites may be both core fucosylated and have a bisecting structure. However, this is unlikely as it has previously been demonstrated that GnT-III modification of an oligosaccharide actively inhibits the addition of a fucose by Fut8 [77]. This discrepancy may be explained by the alternative binding specificities of certain lectins. E-PHA predominantly binds to N-glycans containing a bisecting structure but has also been shown to have affinity for oligosaccharides with terminal galactose residues [70,78]. Similarly, LCA has very high binding affinity for α1,6-fucosylated glycans, but can also bind the mannose core of an N-glycan in the absence of 213 core fucosylation [67]. Further investigation will be required to decipher which structures they possess as data from the ELLA suggests the presence of either a core fucosylated N-glycan with a possible terminal galactose residue, or a bisecting N-glycan. Data from previous studies [56] along with glycosyltransferase knockdown and chemical inhibition experiments in this study, suggests that at least one of these three sites contains an N-glycan with a bisecting structure, with the most likely candidate being N1587 as mutation of this residue resulted in the greatest reduction in E-PHA binding. Future studies using additional mutations on both the essential asparagine residue and on other proximal non-essential residues could reveal the impact of the amino acid alteration on protein expression, function, and stability. Many studies which involve the mutation of N-glycosylation sites on a glycoprotein also use simultaneous multi-site mutation to investigate whether the combined impact of losing two or more N-glycans differs from the effect observed following loss of a single N-glycan [75, 79, 80]. This is a strategy that could reveal crucial information about the synergistic roles of N-glycans on the Notch1 ECD.

Regarding the mutations at sites N888, N1241 and N1587, their effect specifically prevents canonical Notch signalling. ELLA data suggests that the N-glycan at N888 possessed either a bisecting or core fucosylated structure and combining this with evidence from glycosyltransferase knockdown experiments, this means that its absence does not lead to accumulation of Notch1 in the ER-Golgi complex. Furthermore, since the mutation at this site did not result in any change to Notch1 protein levels in the cell, i.e., did not accelerate Notch1 degradation, this suggests that it is most likely a core fucosylated N-glycan (possibly with a terminal galactose residue), rather than a bisecting N-glycan. Taken together, these data suggest that the N-glycan at N888 has a direct influence on Notch1 signal activation at the membrane. This may occur through regulation of processes such as receptor folding or cleavage, mechanisms which are regularly affected by loss of an individual N-glycan on other membrane-bound proteins [81-83]. The N1241I mutation resulted in a dramatic decrease in Notch1 protein levels in the cell, implying that the Notch1 receptor was rapidly degraded, and as a result, Notch target gene expression was inhibited. ELLA data suggested that there may have been no N-glycan present at N1241 but evidence from this mutation experiment is a strong indication that an

N-glycan does indeed reside at this site but does not contain bisecting or core fucosylated structures. Removal of the N-glycan at N1587 also reduced Notch1 protein levels, most likely through receptor degradation. Combined with ELLA data and evidence from siRNA knockdown experiments, the cumulative data suggests that the oligosaccharide which resides at this site contains a complex bisecting structure.

Notably, all six mutations resulted in a reduction of the myogenic differentiation of stem cells in response to Jagged-1 stimulation. This further supports lectin binding data by confirming that N-glycans reside at each of the six consensus sites and reinforces data from chemical inhibition and glycosyltransferase knockdown experiments by demonstrating that all these glycans are involved in the process of Notch-mediated myogenic differentiation. However, only the mutations at sites N888, N1241 and N1587 significantly attenuated the expression of the Notch target gene *Hey1*, suggesting that the other three mutations at sites N959, N1179 and N1489 have an indirect effect on myogenic differentiation which is independent of canonical Notch signalling. In addition, none of these three mutations led to significant changes in Notch1 receptor protein levels in the cell. Their effect on myogenic differentiation may be related to non-canonical Notch signalling pathways, such as the Wnt/β-catenin pathway [84-86]. The Notch1 receptor has been shown to interact with activated β-catenin at the membrane [87]. While this interaction occurs at the intracellular C- terminal of the Notch1 receptor and it is unlikely that any direct interaction takes place between these glycans and β-catenin, it is possible that their absence leads to altered processing/folding of the receptor which results in dysregulation of Wnt signalling and myogenic differentiation (Table II).

Data limitations include the narrow range of available lectins which are specific for N-linked oligosaccharides. Most commercially available lectins exhibit their highest affinity for a particular monosaccharide in a specific branching/linkage format. However, many of these structures can be found in both O-linked and N-linked glycans, such as Erythrina critagalli lectin (ECL) that bind strongly to the terminal galactose of an N-acetyllactosamine-containing N-glycan, but will also bind to a terminal β1,4-galactose in O-glycans [88]. In addition, while mutation of specific asparagine residues is a highly specific method, a single change in nucleotide or amino acid sequence can have a dramatic effect on the expression, processing and/or function of the target protein [80, 89]. In this study, isoleucine was chosen as the substitute due the ease of mutation (only a single base substitution required for each Asn site) and the non-reactive nature of the side chain. However, the change in side-chain composition may alter the conformation of the receptor or result in issues with steric hindrance [90]. While glutamine as a substitute is widely used since its structure is the most similar to asparagine, the polar amide group of the glutamine residue may lead to unwanted molecular reactions [91]. This can be circumvented by mutating the Asn residue to a number of different amino acids, mutating the serine/threonine residue in the consensus sequence, or mutating a proximal amino acid which is not part of the consensus sequence [92]. Finally, alternative chemical inhibitors such as swainsonine, which blocks α-mannosidase-II activity, could provide greater insights into the role of N-glycans while knockdown of additional glycosyltransferases such as GnT-IV and ST3Gal1 could be useful in deciphering the exact structures of N-linked oligosaccharides on the Notch1 receptor [52, 93].

In conclusion, activation of the Notch1 receptor by its ligand, Jagged-1, stimulates myogenic differentiation of resident S100β mVSCs. Furthermore, N-glycosylation intervention strategies, including chemical inhibition, siRNA knockdown of the GnT-III and Fut8 glycosyltransferases, and site-directed mutagenesis of N-glycosylation sites, reveal that this signalling is heavily regulated by N-glycans which reside at six consensus

sequences along the extracellular domain of Notch1. Several key oligosaccharides at these sites were found to contain bisecting or core fucosylated structures. Notch1 expression and trafficking analysis indicated that N-glycans at asparagine (N) 1241 and 1587 protect the receptor from accelerated degradation, while the oligosaccharide at N888 directly affects signal transduction. Conversely, N-linked glycans at the other three sites do not impact canonical Notch1 signalling but still play a role in Notch-mediated myogenic differentiation, possibly via a non-canonical Notch signalling pathway.

The putative role of bisecting and core fucosylated N-glycans in ligand induced activation of the Notch1 receptor may present as a novel therapeutic target for controlling Notch1 signalling and myogenic differentiation of S100β resident vascular stem cells during subclinical atherosclerosis where these cells have been reported to play a major role in early lesion development [16-18].

Materials and Methods

Materials

A full list of the materials is available in Supplementary Materials

Cell Culture

Cells were grown in tissue culture flasks (25-175 cm^2) or in 6-well plates in a humidified incubator and were visualised by a Nikon Eclipse TS100 phase contrast microscope. S100β mouse vascular stem cells (mVSCs) were isolated from the aortic arch of female C57BL/6J mice (used in accordance with EU legislation (Directive 2010/63/EU)) as previously described and grown in maintenance medium (MM1) containing DMEM supplemented with 2% chick embryo extract, 1% FBS, 0.02 µg/ml, bFGF basic protein, B-27 Supplement, N-2 supplement, 1% Penicillin-Streptomycin, 50 nM 2-Mercaptoethanol and 100 nM retinoic acid [16]. These stem cells exhibited a uniform neural-like morphology in low density culture adopting a dendritic-like tree shape and retaining their morphological characteristics at low density throughout repeated passage. Cells were fed with MM1 every 2-3 days and passaged every 3-4 days or when ~70% confluent. Cells were routinely grown in MM and dissociated with 0.5X TrypLE and expanded. Cells were used at passage numbers below p25. C3H/10T1/2 embryonic fibroblasts have been extensively characterised as multipotent mesenchymal cells [47]. Clone 8 - CCL-226 cells were grown in EMEM basal media supplemented with 10% heat inactivated FBS and 1% Penicillin Streptomycin. Cells were used at passage numbers below P20 and were always sub-cultured before confluency surpassed 70% to prevent post-confluence inhibition. Freezing media consisted of EMEM with 10% FBS, 1% Pen-Strep and 5% DMSO.

Adipogenesis Differentiation

Adipogenic differentiation of mVSCs was performed using StemPro® adipogenesis differentiation medium in accordance with the manufacturer's recommendations. Briefly, cells were seeded at a density of 1 x 10^4 cells/cm^2 in their maintenance medium and left for two days before being washed with PBS twice followed by feeding with StemPro® adipogenesis differentiation media for 14 days. Cells were harvested and adipogenic differentiation was evaluated by measuring the expression level of lipoprotein lipase (*Lpl*) and fatty acid binding protein 4 (*Fabp4*).

Osteogenesis Differentiation

Osteogenic differentiation of mVSCs was performed using the StemPro® osteogenic differentiation medium in accordance with the manufacturer's recommendations. Briefly, cells were seeded at a density of 1 x 10^4 cells/cm^2 in their maintenance medium and left for two days before being washed with PBS twice followed by feeding with StemPro® osteogenic differentiation media for 21 days. Cells were harvested and osteogenic differentiation was evaluated by measuring the expression level of alkaline phosphatase (*Alpl*) and Sclerostin (*Sost*).

Telomerase Activity Assay

Cell lysates were prepared using the RIPA buffer method and diluted to a protein concentration of 10-750 ng/µL before telomerase activity was measured using a TRAPeze® XL Telomerase Detection Kit. Briefly, PCR was carried out using fluorescein-labelled primers to amplify telomeric repeats in the sample DNA. Higher telomerase activity meant that there were more telomeric repeats to be amplified by the Taq polymerase, resulting in a more fluorescent product. The PCR products were analysed in a Tecan Safire II fluorescent plate reader. Telomerase activity was calculated by comparing the level of fluorescence detected in each sample to a standard curve generated from a set of telomeric repeat standards. The telomerase activity of a sample was given in TPG units.

Plasmid Amplification and Purification

Transformation: LB Agar (20g/L LB, 12g/L Agar) was made up in dH$_2$O and autoclaved to sterilise. After cooling to 55°C, appropriate antibiotics were added (ampicillin at 100µg/mL or kanamycin at 50µg/mL). The agar was poured into sterile petri dishes, allowed to harden, and stored at 4°C. A sample (100ng) of plasmid DNA was mixed with 50µL of competent JM109 bacterial cells in a sterile microfuge tube. Tube was placed on ice for 30 mins. Cells were then heat-shocked at 42°C for 45 secs to allow plasmid DNA to cross the cell membrane. Tube was placed back on ice for 2 mins. 500µL of SOC medium (20% tryptone, 5% yeast extract, 0.5% NaCl, 1% 0.25M KCl, 2% 1M glucose) was added to the tube which was then incubated at 37°C for 45 mins. The contents of the tube were transferred to a LB Agar plate with appropriate antibiotics. Plates were incubated overnight at 37°C. A single colony was picked from the transformation plate and inoculated in 5mL of LB broth (20g/L LB) with appropriate antibiotics. This culture was incubated at 37°C in a shaking incubator for 6-8 hours. 1.2mL of this culture was then transferred to a sterile conical culture flask containing 250mL of LB broth with appropriate antibiotic. Flask was incubated at 37°C overnight in a shaking incubator. 50mL was removed from the flask for plasmid purification. The rest was used to make stocks by supplementing with 50% glycerol and storing at -80°C.

Plasmid DNA Purification

Plasmid DNA was purified from transformed bacterial cells using the Invitrogen PureLink™ HiPure Plasmid Filter Midiprep Kit according to the manufacturer's guidelines. A Nanodrop® ND-1000 Spectrophotometer was used to quantitate the DNA concentration before DNA samples were interrogated by agarose gel electrophoresis and visualised with the SynGene G-Box.

RNA Extraction

RNA was extracted from cells using the Promega Reliaprep™ RNA Miniprep System according to the provided instructions. A Nanodrop® ND-1000 Spectrophotometer was used to quantitate the RNA concentration before RNA samples were analysed by *Quantitative Reverse-Transcription Polymerase Chain Reaction (qRT-PCR)*

Quantitative Reverse-Transcription Polymerase Chain Reaction (qRT-PCR)

Quantitative RT-PCR was performed using the Rotor Gene Q from Corbett Research, as previously described [16]. A ubiquitously expressed reference gene, hypoxanthine guanine phosphoribosyltransferase (*Hprt)* was used as a control. Two different kits were used for this method depending on the primer set: Rotor-Gene SYBR® Green RT-PCR Kit (Qiagen, 204174) and SensiFAST™ SYBR® No-ROX One-Step Kit (BioLine, BIO-72001). Samples were run in triplicate for each primer set.

Immunocytochemistry

Immunocytochemical analysis of protein expression in cells was performed, as described previously [16]. Fixed cells on coverslips were stained for expression of the intermediate myogenic differentiation marker Calponin1 (Cnn1) and the calcium binding protein, S100β by incubating overnight at 4°C with antibody diluted to an appropriate concentration with 1% BSA in PBS. Cells were washed x3 and incubated at room temperature for 1 hr with the secondary antibody (diluted with 1% BSA in PBS), protected from light. Nuclear staining was performed with DAPI which was diluted to 1μg/mL. Coverslips were mounted before cell nuclei and antibody staining was visualised and imaged using the Olympus DP-50 fluorescent microscope. The number of positive cells was counted for each image and divided by the total number of cells in that image to give a ratio, and subsequently a percentage, of antigen-positive cells.

Transient Transfections

Transfection of both plasmid DNA and single strand siRNA duplexes into cells was performed using the Mirus Bio TransIT-X2® dynamic delivery system, according to the manufacturer's guidelines after optimization. Cells were grown to ~ 80% confluency in the maintenance medium before transfection. The transfection mix was made up in a sterile microfuge tube according to kit protocol and incubated at room temperature for 30 minutes to allow transfection complexes to form. This mixture was then added dropwise to the cell culture. Cells were incubated at 37°C for 48 hours before harvesting. Transfection efficiency was assessed using a control GFP-plasmid or TYE 563 DsiRNA (Mirius Bio), a fluorescently tagged siRNA. A non-transfected sample was used as a negative control. Cells were incubated at 37°C for 48 h before being fixed with 3.7% formaldehyde. Nuclei were stained with DAPI, and coverslips were mounted onto slides before being imaged using a Olympus DP-50 fluorescent microscope. The number of cells containing GFP, or dye were counted and graphed as a percentage of the total number of cells. A transfection efficiency of 99.6% was achieved in the dye-transfected sample, compared to 0% in the non-transfected sample.

Genomic DNA Extraction

Genomic DNA was extracted from cells using the Bioline ISOLATE II Genomic DNA Kit according to the manufacturer's instructions. Briefly, cells were washed and scraped into PBS and pelleted by centrifugation at 1500 rpm for 5 minutes. The pellet was resuspended in lysis buffer GL. Proteinase K and Lysis Buffer G3 were then added to the sample before incubation at 70ºC for 15 minutes. 99% ethanol was added to each sample to adjust DNA binding conditions. Samples were loaded into columns containing a silica membrane and centrifuged at 11,000 x g for 1 minute. The flow-through was discarded and columns were washed twice in wash buffer. Elution buffer was then added to each column. Samples were incubated at room temperature for 1 minute before centrifugation at 11,000 x g for 1 minute. The flow-through contained the purified DNA.

Protein Isolation and Quantitation

Cells at >80% confluency were scraped into culture media and pelleted by centrifugation. The supernatant was discarded, cells were resuspended in an appropriate volume of RIPA buffer and incubated on ice for 30 minutes to induce cell lysis. The lysate was then clarified by centrifugation at 8,000 x g for 20 minutes at 4ºC. The supernatant containing cellular proteins was retained while the cell debris pellet was discarded. The Pierce™ BCA Protein Assay Kit was used to quantitate the concentration of protein in the clarified cell lysate.

Membrane Protein Isolation

Isolation of proteins from the cell membrane was carried out using the Mem-PER Plus Membrane Protein Extraction Kit according to the manufacturer's instructions.

Membrane Fractionation

Cells were harvested and the Mem-PER™ Plus Membrane Protein Extraction Kit was used to perform membrane fractionation on the crude lysate according to the manufacturer's instructions. Protein concentration in both the cytoplasmic fraction and membrane fraction was quantified by BCA Assay. Both samples were run on an SDS-PAGE precast protein gel and subsequently transferred to a nitrocellulose membrane. Staining with Ponceau S confirmed the difference in the presence of protein band profiles between the two samples. Immunoblotting was then carried out with an anti-Notch1 antibody and chemiluminescent staining was used to visualise bands.

Effect of TNC and DMJ on Membrane Localisation of Notch1

Cells were treated in the absence or presence of TNC (0.5μg/mL), DMJ (4μg/mL), 1 x Brefeldin A (BFA) at 37ºC for 48 h before being harvested. Membrane fractions were isolated using the Mem-PER™ Plus Membrane Protein Extraction Kit, according to the manufacturer's instructions. Protein concentration in the membrane fractions was measured by BCA Assay. Western blot was then carried out and samples were immuno-probed using an anti-Notch1 antibody. Chemiluminescent staining was used to visualise immunoblotting. Densitometry was carried out using ImageJ to measure the intensity of Notch1 bands while β-actin, pan-cadherin and Ponceau S staining was used as loading controls.

Trafficking of Notch1 Receptor

Cells were seeded at 5,000 cells/well onto sterile coverslips and treated with or without TNC (0.5μg/mL), DMJ (4μg/mL), 1 x BFA at 37°C for 48 h before being fixed with 3.7% formaldehyde. Immunocytochemistry was then carried out and cells were stained using an anti-Notch1 antibody with an IgG control antibody used as a secondary control. A control sample was also stained with an anti-Golgin-97 antibody to visualise the subcellular location of the golgi apparatus (GA). Nuclei were stained with DAPI. Samples were imaged using a fluorescent microscope. ImageJ was used to calculate the mean fluorescence intensity and the integrated density of fluorescence at both the cell membrane and in the perinuclear region (adjacent to the nucleus, i.e., where the GA is located). Samples were graphed as a ratio of the fluorescence at the membrane to the fluorescence in the perinuclear region.

Purification of Notch1 by Immunoprecipitation

Protein lysates were prepared, and the protein concentration measured by BCA assay. Immunoprecipitation was carried out using Protein A-conjugated magnetic beads and either an anti-Notch1 rabbit monoclonal antibody, a rabbit IgG control antibody, or a mouse IgG control antibody. Following incubation with the antibody, supernatants (containing unbound proteins) were transferred to a fresh tube and appropriately stored. Bound protein was eluted off the beads by either 1X Laemmli Buffer or a 0.1M Glycine buffer (pH 2.0). Samples eluted with the glycine buffer were neutralised using a 1M Tris buffer (pH 9.0). Both eluted bound proteins and previously stored unbound protein samples were then run on an SDS-PAGE precast protein gel, transferred to a nitrocellulose membrane, immuno-probed using an anti-Notch1 antibody and visualised by TMB staining of the HRP-conjugated secondary antibody.

Ectopic Expression of Myc-tagged Notch1 and NICD

The pCS2 Notch1 full length-6MT and pCS2 Notch1 ICv-6MT plasmids were purchased from Addgene (LGC Standards, Teddington, UK) as bacterial stabs. Plasmid-containing bacteria were transferred to a 5mL culture with selective antibiotic at 37°C for 8 hrs. 1.2mL of this culture was then transferred to a 250mL secondary culture with selective antibiotics which was incubated overnight at 37°C. Plasmid DNA was isolated from this culture using the Invitrogen PureLink™ HiPure Plasmid Filter Midiprep Kit. To verify plasmid length and quality, plasmid DNA was digested with an appropriate restriction enzyme. This served to linearise the DNA before running plasmid samples through gel electrophoresis on an agarose gel. Visualisation of this gel under UV light confirmed that plasmids were the correct length and that the stocks contained no contaminating DNA. To validate the ectopic expression of both proteins, cells were grown to ~80% confluency in 3 wells of a 6-well plate and transfected with either the Notch1 full-length plasmid, the NICD plasmid or an empty vector plasmid (pCDNA3.1) using the TransIT-X2 Dynamic Delivery System. Cells were incubated at 37°C for 48 h in MM1 before being harvested. Protein was isolated and quantified by BCA assay. Western blot was then carried out, with all samples being immuno-probed using an anti-Myc-tag antibody to stain for either the myc-tagged Notch1 or myc-tagged NICD, and an anti-β-actin antibody was used as the loading control.

Western Blot

Following protein isolation and quantitation, samples were prepared for SDS-PAGE using a precast SDS 10% polyacrylamide gel to perform electrophoresis and the Pierce™ G2 Fast Blotter for protein transfer, as previously described [48]. Ponceau S staining confirmed that the transfer had been successful. Blocking buffer (5% dried milk in TBS with 0.1% Tween) was added to the membrane and incubated at room temperature for 1-2 h on a rocker before the membrane was washed in TBST (50mM Tris, 150mM NaCl, 1mM $MnCl_2$, 1mM $CaCl_2$, 1mM $MgCl_2$, 0.1% Tween-20) and incubated with the primary antibody (diluted with 5% BSA in TBST) overnight on a rocker at 4°C. Primary antibody was removed, membrane washed 3 times (5 minutes each) with TBST, before incubation with an HRP-conjugated secondary antibody (diluted with 5% BSA in TBST) on a rocker for 1 hr at room temperature. Fresh TMB was added to the membrane which was then incubated on a rocker at room temperature for approx. 10 mins before the blot was imaged using BioRad GelDoc EZ Gel System. Alternatively, Pierce™ ECL Western Blotting Substrate was added to the membrane which was then incubated on a rocker at room temperature for 5 minutes. The blot was then imaged using Syngene G-Box.

Enzyme-Linked Lectin Assay (ELLA)

All samples were diluted to the same concentration with 50μL of treated sample or control sample added in triplicate to each well of a 96-well plate. The plate was incubated overnight at 4°C with gentle shaking. Wells were emptied and blocking buffer (0.5% PVA in PBS) added to each well. The plate was incubated for 2 hrs at room temperature with gentle shaking before wells were washed with TBST (50mM Tris, 150mM NaCl, 1mM $MnCl_2$, 1mM $CaCl_2$, 1mM $MgCl_2$, 0.1% Tween-20). Biotinylated lectins (50μL, 5μg/mL) were added to each well and incubated at room temperature for 1 hr with gentle shaking before wells were emptied and washed again with TBST. HRP-conjugated anti-biotin antibody (50μL, 1:10,000) was then added to each well and incubated at room temperature for 1 hour with gentle shaking. Wells were washed with TBST before a TMB substrate solution (90μL) was added and incubated for 10 minutes at room temperature with gentle shaking. Stop solution (10% H_2SO_4) was added to each well before the absorbance was measured at 450 nm in a plate reader.

N-glycosylation Profile of Purified Notch1

Cells were grown to ~80% confluency in 3 wells per 6-well plate. One well was not transfected while the TransIT-X2 Dynamic Delivery System was used to transfect the other two wells with either the Notch1 full-length plasmid or an empty vector control plasmid. Cells were incubated at 37°C for 48 h in maintenance media (MM1) before being harvested. Protein was isolated and quantified by BCA assay. IP was carried out to purify the ectopically expressed Notch1 using Protein G-conjugated magnetic beads and an anti-Myc-tag antibody. The non-transfected sample was immunoprecipitated with a mouse IgG control antibody. Bound proteins were eluted off the magnetic beads using a 0.1M Glycine buffer (pH 2.0) and a 1M Tris buffer (pH 9.0) was used to neutralise the purified protein fraction. BCA assay was carried out again to determine the concentration of protein in each sample post-IP. ELLA was then applied to detect the presence of N-glycans in all three samples. Control wells containing PBS rather than a protein sample were used to determine the baseline for non-specific absorbance, while the mouse IgG control sample

acted as the baseline for non-specific lectin- or antibody-binding to the IP eluate. Both baseline values were subtracted from sample absorbance values to decipher the true level of specific lectin binding. TBST acted as the vehicle control for all lectins.

Site-directed mutagenesis

Mutation of N-glycosylation sites on a plasmid-encoded Notch1 receptor was achieved using the Agilent QuikChange Lightning Site Directed Mutagenesis Kit, according to the manufacturer's instructions. Briefly, primers were specific for a region of the Notch1 gene containing an N-glycosylation site and were designed to incorporate a single base substitution in copies of the plasmid DNA template. PCR tubes were placed in an Applied Biosystems® Veriti® Thermal Cycler. A DpnI restriction enzyme was added to each tube of PCR products to degrade any parental template DNA. Mutant plasmids were then transformed into ultracompetent XL10-Gold bacterial cells via heat-shock at 42°C. After allowing the cells to recover for 1 hour in NZY$^+$ broth at 37°C, each transformation reaction was spread on an LB agar plate with Ampicillin (50μg/mL) and incubated at 37°C overnight. Colonies were picked and inoculated in liquid LB. Transformed bacteria were grown and mutant plasmids were purified before sending them for sequencing with Source Biosciences to confirm that the correct single base substitution had been incorporated in each case.

Data Analysis and Statistics

The mean and standard deviation (or standard error margin) was calculated for all the technical and biological replicates in each experiment and these values were used to plot the data. The Kolmogorov-Smirnov normality test was used to analyse the spread of the data as it uses the most stringent conditions. The outcome of the normality test (along with the number of samples and parameters) dictated what type of significance test was carried out. In the case of data containing only two samples, an unpaired t-test with Welch's correction was performed on normally distributed data. For non-parametric (skewed) data, the Wilcoxon test was carried out to calculate statistical significance. For data containing more than two samples, ANOVA testing was used to perform multiple comparisons between all samples. This could be carried out using an ordinary one-way (one parameter to compare) or two-way (two parameters to compare) ANOVA for normalised data, or the Kruskal-Wallis test for non-parametric data. This would reveal the level of statistical significance for both the data as a whole and between each sample.
The Materials and Methods section should provide sufficient information to allow replication of the results. Begin with a section titled Experimental Design describing the objectives and design of the study as well as prespecified components.

References

1. MacGrogan D, Münch J, de la Pompa JL. Notch and interacting signalling pathways in cardiac development, disease, and regeneration. Nat Rev Cardiol. 2018;15:685–704.

2. Talora C, Campese AF, Bellavia D, Felli MP, Vacca A, Gulino A, et al. Notch signaling and diseases: an evolutionary journey from a simple beginning to complex outcomes. Biochim Biophys Acta. 2008;1782:489–97.

3. Bray SJ. Notch signalling in context. Nat Rev Mol Cell Biol. 2016;17:722–35.

4. Siebel C, Lendahl U. Notch Signaling in Development, Tissue Homeostasis, and Disease. Physiol Rev. 2017;97:1235–94.

5. Katoh M. Canonical and non-canonical WNT signaling in cancer stem cells and their niches: Cellular heterogeneity, omics reprogramming, targeted therapy and tumor plasticity (Review). Int J Oncol. 2017;51:1357–69.

6. Caolo V, Schulten HM, Zhuang ZW, Murakami M, Wagenaar A, Verbruggen S, et al. Soluble Jagged-1 inhibits neointima formation by attenuating Notch-Herp2 signaling. Arterioscler Thromb Vasc Biol. 2011;31:1059–65.

7. Lindner V, Booth C, Prudovsky I, Small D, Maciag T, Liaw L. Members of the Jagged/Notch gene families are expressed in injured arteries and regulate cell phenotype via alterations in cell matrix and cell-cell interaction. Am J Pathol. 2001;159:875–83.

8. Li Y, Takeshita K, Liu P-Y, Satoh M, Oyama N, Mukai Y, et al. Smooth muscle Notch1 mediates neointimal formation after vascular injury. Circulation. 2009;119:2686–92.

9. Sakata Y, Xiang F, Chen Z, Kiriyama Y, Kamei CN, Simon DI, et al. Transcription factor CHF1/Hey2 regulates neointimal formation in vivo and vascular smooth muscle proliferation and migration in vitro. Arterioscler Thromb Vasc Biol. 2004;24:2069–74.

10. Redmond EM, Liu W, Hamm K, Hatch E, Cahill PA, Morrow D. Perivascular delivery of Notch 1 siRNA inhibits injury-induced arterial remodeling. PLoS One. 2014;9:e84122.

11. Chappell J, Harman JL, Narasimhan VM, Yu H, Foote K, Simons BD, et al. Extensive Proliferation of a Subset of Differentiated, yet Plastic, Medial Vascular Smooth Muscle Cells Contributes to Neointimal Formation in Mouse Injury and Atherosclerosis Models. Circ Res. 2016;119:1313–23.

12. Dobnikar L, Taylor AL, Chappell J, Oldach P, Harman JL, Oerton E, et al. Disease-relevant transcriptional signatures identified in individual smooth muscle cells from healthy mouse vessels. Nat Commun. 2018;9:4567.

13. Majesky MW, Horita H, Ostriker A, Lu S, Regan JN, Bagchi A, et al. Differentiated Smooth Muscle Cells Generate a Subpopulation of Resident Vascular Progenitor Cells in the Adventitia Regulated by Klf4. Circ Res. 2017;120:296–311.

14. Kramann R, Goettsch C, Wongboonsin J, Iwata H, Schneider RK, Kuppe C, et al. Adventitial MSC-like Cells Are Progenitors of Vascular Smooth Muscle Cells and Drive Vascular Calcification in Chronic Kidney Disease. Cell Stem Cell. 2016;19:628–42.

15. Wan M, Li C, Zhen G, Jiao K, He W, Jia X, et al. Injury-activated transforming growth factor β controls mobilization of mesenchymal stem cells for tissue remodeling. Stem Cells. 2012;30:2498–511.

16. Di Luca M, Fitzpatrick E, Burtenshaw D, Liu W, Helt J-C, Hakimjavadi R, et al. The calcium binding protein S100β marks hedgehog-responsive resident vascular stem cells within vascular lesions. NPJ Regen Med. 2021;6:10.

17. Molony C, King D, Di Luca M, Kitching M, Olayinka A, Hakimjavadi R, et al. Disease-Relevant Single Cell Photonic Signatures Identify S100β Stem Cells and their Myogenic Progeny in Vascular Lesions. Stem Cell Rev Rep. 2021;17:1713–40.

18. Yuan F, Wang D, Xu K, Wang J, Zhang Z, Yang L, et al. Contribution of Vascular Cells to Neointimal Formation. PLoS One. 2017;12:e0168914.

19. Tang J, Wang H, Huang X, Li F, Zhu H, Li Y, et al. Arterial Sca1+ Vascular Stem Cells Generate De Novo Smooth Muscle for Artery Repair and Regeneration. Cell Stem Cell. 2020;26:81-96.e4.

20. Lyu L, Li Z, Wen Z, He Y, Wang X, Jiang L, et al. Fate mapping RNA-sequencing reveal Malat1 regulates Sca1+ progenitor cells to vascular smooth muscle cells transition in vascular remodeling. Cell Mol Life Sci. 2023;80:118.

21. Proweller A, Pear WS, Parmacek MS. Notch signaling represses myocardin-induced smooth muscle cell differentiation. J Biol Chem. 2005;280:8994–9004..

22. Kurpinski K, Lam H, Chu J, Wang A, Kim A, Tsay E, et al. Transforming growth factor-beta and notch signaling mediate stem cell differentiation into smooth muscle cells. Stem Cells. 2010;28:734–42.

23. Doi H, Iso T, Sato H, Yamazaki M, Matsui H, Tanaka T, et al. Jagged1-selective notch signaling induces smooth muscle differentiation via a RBP-Jkappa-dependent pathway. J Biol Chem. 2006;281:28555–64.

24. High FA, Zhang M, Proweller A, Tu L, Parmacek MS, Pear WS, et al. An essential role for Notch in neural crest during cardiovascular development and smooth muscle differentiation. J Clin Invest. 2007;117:353–63.

25. Kopan R, Ilagan MXG. The canonical Notch signaling pathway: unfolding the activation mechanism. Cell. 2009;137:216–33.

26. Fortini ME. Notch signaling: the core pathway and its posttranslational regulation. Dev Cell. 2009;16:633–47.

27. Matsumoto K, Luther KB, Haltiwanger RS. Diseases related to Notch glycosylation. Mol Aspects Med. 2021;79:100938.

28. Moloney DJ, Shair LH, Lu FM, Xia J, Locke R, Matta KL, et al. Mammalian Notch1 is modified with two unusual forms of O-linked glycosylation found on epidermal growth factor-like modules. J Biol Chem. 2000;275:9604–11.

29. Acar M, Jafar-Nejad H, Takeuchi H, Rajan A, Ibrani D, Rana NA, et al. Rumi is a CAP10 domain glycosyltransferase that modifies Notch and is required for Notch signaling. Cell. 2008;132:247–58.

30. Shao L, Moloney DJ, Haltiwanger R. Fringe modifies O-fucose on mouse Notch1 at epidermal growth factor-like repeats within the ligand-binding site and the Abruptex region. J Biol Chem. 2003;278:7775–82.

31. Takeuchi H, Haltiwanger RS. Significance of glycosylation in Notch signaling. Biochem Biophys Res Commun. 2014;453:235–42.

32. Heifetz A, Keenan RW, Elbein AD. Mechanism of action of tunicamycin on the UDP-GlcNAc:dolichyl-phosphate Glc-NAc-1-phosphate transferase. Biochemistry. 1979;18:2186–92.

33. Elbein AD. Inhibitors of the biosynthesis and processing of N-linked oligosaccharide chains. Annu Rev Biochem. 1987;56:497–534.

34. Rademacher TW, Parekh RB, Dwek RA. Glycobiology. Annual Review of Biochemistry. 1988;57:785–838.

35. Narasimhan S. Control of glycoprotein synthesis. UDP-GlcNAc:glycopeptide beta 4-N-acetylglucosaminyltransferase III, an enzyme in hen oviduct which adds GlcNAc in beta 1-

4 linkage to the beta-linked mannose of the trimannosyl core of N-glycosyl oligosaccharides. J Biol Chem. 1982;257:10235–42.

36. Gu J, Nishikawa A, Tsuruoka N, Ohno M, Yamaguchi N, Kangawa K, et al. Purification and characterization of UDP-N-acetylglucosamine: alpha-6-D-mannoside beta 1-6N-acetylglucosaminyltransferase (N-acetylglucosaminyltransferase V) from a human lung cancer cell line. J Biochem. 1993;113:614–9.

37. Longmore GD, Schachter H. Product-identification and substrate-specificity studies of the GDP-L-fucose:2-acetamido-2-deoxy-beta-D-glucoside (FUC goes to Asn-linked GlcNAc) 6-alpha-L-fucosyltransferase in a Golgi-rich fraction from porcine liver. Carbohydr Res. 1982;100:365–92.

38. Song Y, Aglipay JA, Bernstein JD, Goswami S, Stanley P. The bisecting GlcNAc on N-glycans inhibits growth factor signaling and retards mammary tumor progression. Cancer Res. 2010;70:3361–71.

39. Staudacher E, Altmann F, Wilson IB, März L. Fucose in N-glycans: from plant to man. Biochim Biophys Acta. 1999;1473:216–36.

40. Wilson JR, Williams D, Schachter H. The control of glycoprotein synthesis: N-acetylglucosamine linkage to a mannose residue as a signal for the attachment of L-fucose to the asparagine-linked N-acetylglucosamine residue of glycopeptide from alpha1-acid glycoprotein. Biochem Biophys Res Commun. 1976;72:909–16.

41. Okazaki A, Shoji-Hosaka E, Nakamura K, Wakitani M, Uchida K, Kakita S, et al. Fucose depletion from human IgG1 oligosaccharide enhances binding enthalpy and association rate between IgG1 and FcgammaRIIIa. J Mol Biol. 2004;336:1239–49.

42. Osumi D, Takahashi M, Miyoshi E, Yokoe S, Lee SH, Noda K, et al. Core fucosylation of E-cadherin enhances cell-cell adhesion in human colon carcinoma WiDr cells. Cancer Sci. 2009;100:888–95.

43. Pakkiriswami S, Couto A, Nagarajan U, Georgiou M. Glycosylated Notch and Cancer. Front Oncol. 2016;6:37.

44. Lis H, Sharon N. Lectins: Carbohydrate-Specific Proteins That Mediate Cellular Recognition. Chem Rev. 1998;98:637–74.

45. Ghazarian H, Idoni B, Oppenheimer SB. A glycobiology review: carbohydrates, lectins and implications in cancer therapeutics. Acta Histochem. 2011;113:236–47.

46. Thompson R, Creavin A, O'Connell M, O'Connor B, Clarke P. Optimization of the enzyme-linked lectin assay for enhanced glycoprotein and glycoconjugate analysis. Anal Biochem. 2011;413:114–22.

47. Zhao L, Li G, Chan K-M, Wang Y, Tang P-F. Comparison of multipotent differentiation potentials of murine primary bone marrow stromal cells and mesenchymal stem cell line C3H10T1/2. Calcif Tissue Int. 2009;84:56–64.

48. Morrow D, Scheller A, Birney YA, Sweeney C, Guha S, Cummins PM, et al. Notch-mediated CBF-1/RBP-J{kappa}-dependent regulation of human vascular smooth muscle cell phenotype in vitro. Am J Physiol Cell Physiol. 2005;289:C1188-1196.

49. Chen Q, Yang F, Guo M, Wen G, Zhang C, Luong LA, et al. miRNA-34a reduces neointima formation through inhibiting smooth muscle cell proliferation and migration. J Mol Cell Cardiol. 2015;89 Pt A:75–86.

50. Fan H, Meng W, Kilian C, Grams S, Reutter W. Domain-specific N-glycosylation of the
membrane glycoprotein dipeptidylpeptidase IV (CD26) influences its subcellular
trafficking, biological stability, enzyme activity and protein folding. Eur J Biochem.
1997;246:243–51.

51. Li L-B, Chen N, Ramamoorthy S, Chi L, Cui X-N, Wang LC, et al. The role of N-
glycosylation in function and surface trafficking of the human dopamine transporter. J Biol
Chem. 2004;279:21012–20.

52. Stanley P. What Have We Learned from Glycosyltransferase Knockouts in Mice? J Mol
Biol. 2016;428:3166–82.

53. Boskovski MT, Yuan S, Pedersen NB, Goth CK, Makova S, Clausen H, et al. The
heterotaxy gene GALNT11 glycosylates Notch to orchestrate cilia type and laterality.
Nature. 2013;504:456–9.

54. Matsumoto K, Ayukawa T, Ishio A, Sasamura T, Yamakawa T, Matsuno K. Dual Roles of
O-Glucose Glycans Redundant with Monosaccharide O-Fucose on Notch in Notch
Trafficking. J Biol Chem. 2016;291:13743–52.

55. Ogawa M, Tashima Y, Sakaguchi Y, Takeuchi H, Okajima T. Contribution of extracellular
O-GlcNAc to the stability of folded epidermal growth factor-like domains and Notch1
trafficking. Biochem Biophys Res Commun. 2020;526:184–90.

56. Allam H, Johnson BP, Zhang M, Lu Z, Cannon MJ, Abbott KL. The glycosyltransferase
GnT-III activates Notch signaling and drives stem cell expansion to promote the growth and
invasion of ovarian cancer. J Biol Chem. 2017;292:16351–9.

57. Tang Z, Wang A, Yuan F, Yan Z, Liu B, Chu JS, et al. Differentiation of multipotent
vascular stem cells contributes to vascular diseases. Nat Commun. 2012;3:875.

58. Morrow D, Guha S, Sweeney C, Birney Y, Walshe T, O'Brien C, et al. Notch and vascular
smooth muscle cell phenotype. Circulation Research. 2008;103:1370–82.

59. Noseda M, Fu Y, Niessen K, Wong F, Chang L, McLean G, et al. Smooth Muscle alpha-
actin is a direct target of Notch/CSL. Circ Res. 2006;98:1468–70.

60. Doi H, Iso T, Yamazaki M, Akiyama H, Kanai H, Sato H, et al. HERP1 inhibits myocardin-
induced vascular smooth muscle cell differentiation by interfering with SRF binding to
CArG box. Arterioscler Thromb Vasc Biol. 2005;25:2328–34.

61. Maier MM, Gessler M. Comparative analysis of the human and mouse Hey1 promoter: Hey
genes are new Notch target genes. Biochem Biophys Res Commun. 2000;275:652–60.

62. Chang P-J, Boonsiri J, Wang S-S, Chen L-Y, Miller G. Binding of RBP-Jkappa (CSL)
protein to the promoter of the Kaposi's sarcoma-associated herpesvirus ORF47 (gL) gene is
a critical but not sufficient determinant of transactivation by ORF50 protein. Virology.
2010;398:38–48.

63. Tang Y, Urs S, Boucher J, Bernaiche T, Venkatesh D, Spicer DB, et al. Notch and
transforming growth factor-beta (TGFbeta) signaling pathways cooperatively regulate
vascular smooth muscle cell differentiation. J Biol Chem. 2010;285:17556–63.

64. Hasan SS, Fischer A. Notch Signaling in the Vasculature: Angiogenesis and Angiocrine
Functions. Cold Spring Harb Perspect Med. 2023;13:a041166.

65. Peterson SM, Turner JE, Harrington A, Davis-Knowlton J, Lindner V, Gridley T, et al. Notch2 and Proteomic Signatures in Mouse Neointimal Lesion Formation. Arterioscler Thromb Vasc Biol. 2018;38:1576–93.

66. Allam H, Aoki K, Benigno BB, McDonald JF, Mackintosh SG, Tiemeyer M, et al. Glycomic analysis of membrane glycoproteins with bisecting glycosylation from ovarian cancer tissues reveals novel structures and functions. J Proteome Res. 2015;14:434–46.

77. Maupin KA, Liden D, Haab BB. The fine specificity of mannose-binding and galactose-binding lectins revealed using outlier motif analysis of glycan array data. Glycobiology. 2012;22:160–9.

68. Nagae M, Soga K, Morita-Matsumoto K, Hanashima S, Ikeda A, Yamamoto K, et al. Phytohemagglutinin from Phaseolus vulgaris (PHA-E) displays a novel glycan recognition mode using a common legume lectin fold. Glycobiology. 2014;24:368–78.

69. Yu G, Nishimura M, Arawaka S, Levitan D, Zhang L, Tandon A, et al. Nicastrin modulates presenilin-mediated notch/glp-1 signal transduction and betaAPP processing. Nature. 2000;407:48–54.

70. Yagi H, Saito T, Yanagisawa M, Yu RK, Kato K. Lewis X-carrying N-glycans regulate the proliferation of mouse embryonic neural stem cells via the Notch signaling pathway. J Biol Chem. 2012;287:24356–64.

71. Kakuda S, Haltiwanger RS. Deciphering the Fringe-Mediated Notch Code: Identification of Activating and Inhibiting Sites Allowing Discrimination between Ligands. Dev Cell. 2017;40:193–201.

72. Stahl M, Uemura K, Ge C, Shi S, Tashima Y, Stanley P. Roles of Pofut1 and O-fucose in mammalian Notch signaling. J Biol Chem. 2008;283:13638–51.

73. Sun X, Tie HC, Chen B, Lu L. Glycans function as a Golgi export signal to promote the constitutive exocytic trafficking. J Biol Chem. 2020;295:14750–62.

74. Skrenkova K, Lee S, Lichnerova K, Kaniakova M, Hansikova H, Zapotocky M, et al. N-Glycosylation Regulates the Trafficking and Surface Mobility of GluN3A-Containing NMDA Receptors. Front Mol Neurosci. 2018;11:188.

75. Janezic EM, Lauer SM-L, Williams RG, Chungyoun M, Lee K-S, Navaluna E, et al. N-glycosylation of α1D-adrenergic receptor N-terminal domain is required for correct trafficking, function, and biogenesis. Sci Rep. 2020;10:7209.

76. Weis WI, Drickamer K. Structural basis of lectin-carbohydrate recognition. Annu Rev Biochem. 1996;65:441–73.

77. Zhao Y, Sato Y, Isaji T, Fukuda T, Matsumoto A, Miyoshi E, et al. Branched N-glycans regulate the biological functions of integrins and cadherins. FEBS J. 2008;275:1939–48.

78. Yamamoto K, Ito S, Yasukawa F, Konami Y, Matsumoto N. Measurement of the carbohydrate-binding specificity of lectins by a multiplexed bead-based flow cytometric assay. Anal Biochem. 2005;336:28–38.

79. Lee TK, Koh AS, Cui Z, Pierce RH, Ballatori N. N-glycosylation controls functional activity of Oatp1, an organic anion transporter. Am J Physiol Gastrointest Liver Physiol. 2003;285:G371-381.

80. Ding Y, Uchida K, Aizawa S-I, Murphy K, Berezuk A, Khursigara CM, et al. Effects of N-glycosylation site removal in archaellins on the assembly and function of archaella in Methanococcus maripaludis. PLoS One. 2015;10:e0116402.

81. West MB, Wickham S, Quinalty LM, Pavlovicz RE, Li C, Hanigan MH. Autocatalytic cleavage of human gamma-glutamyl transpeptidase is highly dependent on N-glycosylation at asparagine 95. J Biol Chem. 2011;286:28876–88.

82. Hsiao C-C, Cheng K-F, Chen H-Y, Chou Y-H, Stacey M, Chang G-W, et al. Site-specific N-glycosylation regulates the GPS auto-proteolysis of CD97. FEBS Lett. 2009;583:3285–90.

83. Branza-Nichita N, Negroiu G, Petrescu AJ, Garman EF, Platt FM, Wormald MR, et al. Mutations at critical N-glycosylation sites reduce tyrosinase activity by altering folding and quality control. J Biol Chem. 2000;275:8169–75.

84. Brack AS, Conboy IM, Conboy MJ, Shen J, Rando TA. A temporal switch from notch to Wnt signaling in muscle stem cells is necessary for normal adult myogenesis. Cell Stem Cell. 2008;2:50–9.

85. Acosta H, López SL, Revinski DR, Carrasco AE. Notch destabilises maternal beta-catenin and restricts dorsal-anterior development in Xenopus. Development. 2011;138:2567–79.

86. Tanaka S, Terada K, Nohno T. Canonical Wnt signaling is involved in switching from cell proliferation to myogenic differentiation of mouse myoblast cells. J Mol Signal. 2011;6:12.

87. Hayward P, Brennan K, Sanders P, Balayo T, DasGupta R, Perrimon N, et al. Notch modulates Wnt signalling by associating with Armadillo/beta-catenin and regulating its transcriptional activity. Development. 2005;132:1819–30.

88. Itakura Y, Nakamura-Tsuruta S, Kominami J, Sharon N, Kasai K-I, Hirabayashi J. Systematic comparison of oligosaccharide specificity of Ricinus communis agglutinin I and Erythrina lectins: a search by frontal affinity chromatography. J Biochem. 2007;142:459–69.

89. Delorme E, Lorenzini T, Giffin J, Martin F, Jacobsen F, Boone T, et al. Role of glycosylation on the secretion and biological activity of erythropoietin. Biochemistry. 1992;31:9871–6.

90. Ishikawa H, Uchida T, Takahashi S, Ishimori K, Morishima I. Ligand migration in human myoglobin: steric effects of isoleucine 107(G8) on O(2) and CO binding. Biophys J. 2001;80:1507–17.

91. van Berkel PH, van Veen HA, Geerts ME, de Boer HA, Nuijens JH. Heterogeneity in utilization of N-glycosylation sites Asn624 and Asn138 in human lactoferrin: a study with glycosylation-site mutants. Biochem J. 1996;319 (Pt 1) Pt 1:117–22.

92. Flack MR, Froehlich J, Bennet AP, Anasti J, Nisula BC. Site-directed mutagenesis defines the individual roles of the glycosylation sites on follicle-stimulating hormone. J Biol Chem. 1994;269:14015–20.

93. Jacob GS. Glycosylation inhibitors in biology and medicine. Curr Opin Struct Biol. 1995;5:605–11.

Acknowledgments

Funding:

Irish Research Council (IRC) GOIPG/2016/1529 (EC)
Irish Research Council (IRC) GOIPG/2014/43 (MDL)
Government of Saudi Arabia ID: IR15208 (JG)
National Institutes of Health R01-AA024082 (EMR)
Science Foundation Ireland (SFI) award 11-PI-1128 (PAC)
Health Research Board (HRB), Ireland HRA-POR-2015-1315 (PAC)
1217
1218   **Author contributions**
1219   Conceptualization: PAC, EC
1220   Methodology: EC, AO, BOC, YG, RH, MDL
1221   Investigation: EC, AO,
1222   Visualization: SFB
1223   Supervision: PAC
1224   Writing—original draft: EC, AO
1225   Writing—review & editing: AO, EMR, PAC
1226
1227   **Competing interests:** Authors declare that they have no competing interests.
1228
1229
1230   **Data and materials availability:** The datasets used and/or analysed during the current study are
1231   available from the corresponding author on reasonable request. All data are available in
1232   the main text or the supplementary materials.

Figures and Tables

Figure 1: Jagged-1 activation of Notch1 signalling and SMC differentiation in S100β mVSCs.
Immunocytochemical analysis of (a) Myh11 and S100β protein expression in untreated mVSCs and (b) Cnn1
expression in aortic S100β mVSCs treated with recombinant immobilised Jagged-1 (Jag-1 Fc) or IgG-Fc at
1μg/mL in MM1 media after 7 days. Scale bar = 25μm. (c) Mean percentage of Myh11, S100β and Cnn1
positive cells. DAPI was used to stain nuclei. Positive cells were counted and compared against total DAPI
cell count. Data are the mean ± SEM and representative of n=3 *p ≤ 0.05. (d-g) Real-time qRT-PCR analysis
of *Hey1* and *Myh11* mRNA levels in S100β mVSCs treated with recombinant immobilised Jagged-1 (Jag-1
Fc) or IgG-Fc (Fc) at 1μg/mL for 48 hrs with or without prior treatment of cells with a γ-secretase inhibitor,
DAPT versus DMSO control (d, e) and following Notch1 receptor knockdown (f-g) versus scrambled
control. The gene, hypoxanthine phosphoribosyltransferase *(hprt)* was used as a housekeeping control. Data
are the mean ± SEM and representative of n=3 *p ≤ 0.05

Figure 2: Chemical inhibition of N-glycosylation (TNC and DMJ) on lectin binding and Jagged-1 activation of Notch1 signalling and SMC differentiation in S100β mVSCs. ELLA analysis of N-glycan specific lectin binding of whole cell lysates (a, c) and immunoprecipitated ectopic expression of the Notch1 receptor (b, d) following treatment of S100β mVSC with a range of TNC (a, b) and DMJ (c, d) concentrations for 48 h at 37ºC. Data are presented as the mean absorbance value at 450 nm normalised to the negative control TBST sample. Data are the mean ± SEM and representative of n=3 *p ≤ 0.05 compared to the non-treated sample probed with the same lectins. Real-time qRT-PCR analysis of *Hey1* and *Myh11* mRNA levels in S100β mVSCs treated with recombinant immobilised Jagged-1 (Jag-1 Fc) or IgG-Fc (Fc) at 1µg/mL in MM1 media after 48 hrs following ectopic expression of the Notch1 receptor with or without prior treatment of cells with TNC (0.5µg/mL) (e, f) and DMJ (4µg/mL) (g-h). The gene, hypoxanthine phosphoribosyltransferase *(hprt)* was used as a housekeeping control. Data are the mean ± SEM and representative of n=3 *p ≤ 0.05

Figure 3: **Chemical inhibition of N-glycosylation (TNC and DMJ) on Notch1 intracellular domain (NICD) activation of Notch signalling and SMC differentiation in S100β mVSCs**. Real-time qRT-PCR analysis of *Hey1* (a, c) and *Myh11* (b, d) mRNA levels in S100β mVSCs following ectopic expression of the Notch1 NICD after 48 hrs with or without prior treatment of cells with TNC (0.5μg/mL) (a, b) or DMJ (4μg/mL) (c-d). The gene, hypoxanthine phosphoribosyltransferase *(hpr*t) was used as a housekeeping control. Data are the mean ± SEM and representative of n=3 *p ≤ 0.05.

Figure 4: *Mgat3* and *Fut8* knockdown on lectin binding, Notch signalling, and SMC differentiation in **S100β mVSCs**. ELLA analysis of N-glycan specific lectin binding of whole cell lysate (a) and immunoprecipitated ectopic expression of the Notch1 receptor (b) following *Mgat3* and *Fut8* knockdown with specific siRNA duplexes (25nM) versus scrambled control. Data are presented as the mean absorbance value at 450 nm normalised to the negative control TBST sample. Data are the mean ± SEM and representative of n=3 *p ≤ 0.05. compared to the non-treated sample probed with the same lectins. Real-time qRT-PCR analysis of *Hey1* (c, e) and *Myh11* (d, f) mRNA levels in S100β mVSCs treated with recombinant immobilised Jagged-1 (Jag-1 Fc) or IgG-Fc (Fc) at 1μg/mL after 48 hrs following *Mgat3* (c, d) or *Fut8* (e, f) knockdown with specific siRNA duplexes (25nM) versus scrambled control. The gene, hypoxanthine phosphoribosyltransferase *(hpr*t) was used as a housekeeping control. Data are the mean ± SEM and representative of n=3 *p ≤ 0.05

Figure 5: *Mgat3* and *Fut8* **knockdown on Notch1 intracellular domain (NICD) activation of Notch signalling and SMC differentiation in S100β mVSCs.** Real-time qRT-PCR analysis of *Hey1* (a, c) and *Myh11* (b, d) mRNA levels in S100β mVSCs stimulated by ectopic expression of Notch1 NICD after 48 hrs following *Mgat3* (a, b) and *Fut8* (c, d) knockdown with specific siRNA duplexes (25nM) versus scrambled control. The gene, hypoxanthine phosphoribosyltransferase *(hprt)* was used as a housekeeping control. Immunoblot analysis of Notch1 protein levels (e, f) and real-time qRT-PCR analysis of *Notch1* mRNA levels (g) following *Mgat3* or *Fut8* knockdown with specific siRNA duplexes (25nM) versus scrambled control. (e) a representative immunoblot image of Notch1 bands at 120kDa and β-actin bands at 42kDa. (f) Densitometry analysis of the intensity of Notch1 bands normalised to the β-actin loading control. Notch1 protein levels given as ratio of expression relative to the untreated negative control sample. (g) *Notch1* mRNA levels. Data are the mean ± SEM and representative of n=3, ns or *p ≤ 0.05.

Figure 6: Chemical inhibition of N-glycosylation (TNC and DMJ) and *Mgat3* and *Fut8* knockdown on membrane localisation of the Notch1 receptor. (a) Membrane fractionation was carried out on S100β mVSCs lysates before Notch1 and Pan-Cadherin protein expression was assessed in cytoplasmic and membrane fractions, respectively. A representative immunoblot image of Notch1 at 120kDa (upper panel) and Ponceau S staining (lower panel) in each fraction. (b) Notch 1 protein expression in the membrane fractions of whole lysates from S100β mVSCs treated with TNC (0.5μg/mL), DMJ (4μg/mL) or 1 X BFA for 48 h at 37°C. (c) Notch 1 protein expression in the membrane fraction of whole lysates from S100β mVSCs following *Mgat3* or *Fut8* knockdown compared to scrambled control or cells treated with 1 X BFA. (d, e) Densitometry analysis of the intensity of Notch1 bands normalised to the Ponceau S loading control. Notch1 protein levels in membrane fraction presented as the ratio of expression relative to the untreated negative control or scrambled control. Data are the mean ± SEM and representative of n=3, ns or *p ≤ 0.05

Figure 7: Chemical inhibition of N-glycosylation (TNC and DMJ) and *Mgat3* and *Fut8* knockdown on Notch1 receptor trafficking. Localization of Notch 1 receptor expression in S100β mVSCs treated with (a) TNC (0.5μg/mL), DMJ (4μg/mL) or 1 X BFA or (b) following *Mgat3* or *Fut8* knockdown, compared to untreated negative control sample, scrambled control, or treated with 1 X BFA for 48 h at 37°C. Representative images at 60X magnification. Location of the GA was confirmed using Golgin-97. Scale bar = 10μm. (c-d) ImageJ used to calculate the ratio of Membrane: Perinuclear fluorescence using (c) integrated density and (d) mean fluorescence intensity values. Data are the mean ± SEM and representative of n=3, ns or *p ≤ 0.05.

Figure 8: Site-directed mutagenesis of asparagine residues on the Notch1 ECD receptor on lectin binding, Notch signalling, and SMC differentiation in C3H10T1/2 cells. (a) ELLA analysis of N-glycan specific lectin binding of immunoprecipitated wild-type and mutant Notch1 receptors (N888I, N959I, N1179I, N1241I, N1489I and N1587I) from puromycin selected C3H10T1/2 cells. Data are presented as the mean absorbance value at 450nm normalised to the negative control TBST sample and the IgG immunoprecipitation control. (b, c) Real-time qRT-PCR analysis of Jag-1 stimulated *Hey1* (b) and *Myh11* (c) mRNA levels in C3H10T1/2 cells following ectopic expression of wild-type and mutant Notch1 receptors. Cells were treated with recombinant immobilised Jagged-1 (Jag-1 Fc) or IgG-Fc (Fc) at 1μg/mL in MM1 media after 48 hrs. The gene, hypoxanthine phosphoribosyltransferase *(hpr*t) was used as a housekeeping control. (d) Immunoprecipitated wildtype (WT) and mutant Notch1 receptor expression. A representative immunoblot image of full-length Notch1 bands at 300kDa and β-actin bands at 42kDa. (e) Densitometry analysis of the intensity of Notch1 bands normalised to the β-actin loading control. Full-length Notch1 protein levels presented as the ratio of expression relative to the wild-type Notch1 control. Data are the mean ± SEM and representative of n=3, ns or *p ≤ 0.05.

Lectin	Binding specificity
ConA	Internal α-linked mannose
LCA	Internal α-linked mannose. Strongest affinity for mannose core with α-1,6 linked fucose bound to primary GlcNAc residue
WGA	High affinity for terminal GlcNAc and dimers/trimers of GlcNAc. Also reacts with chitobiose core of N-linked glycans, specifically the Man β(1,4)GlcNAc β(1,4)GlcNAc trisaccharide. Can also bind GlcNAc in O-glycans
E-PHA	Binds complex N-glycans. Usually at branching point from mannose core, specifically Galβ4GlcNAcβ2Manα6; (GlcNAcβ4); (GlcNAcβ4Manα3). Strong affinity for complex bisecting N-glycan structure.

Table I: The binding specificities of lectins. The binding specificity of Concanavalin A (ConA), Lens Culinaris Agglutinin (LCA), Wheat germ agglutinin (WGA) and Phaseolus vulgaris lectin E (E-PHA) on the ELLA assay.

Asn Site	N-Glycan Structures	Possible Role of Glycan
N888		Regulation of receptor folding/cleavage
N959		Impacts non-canonical Notch signalling
N1179		Impacts non-canonical Notch signalling
N1241	No N-glycan present	Regulation of receptor degradation
N1489		Impacts non-canonical Notch signalling
N1587		Regulation of receptor degradation

GlcNAc   Mannose   Galactose   Fucose

* Indicates most likely structure based on Notch1 expression and signalling data

Table II: N-Glycan Structures and Possible Roles on the Notch1 Receptor. The N-glycan structures and possible role of each glycan in regulating Notch signalling at specific asparagine residues on the Notch 1 ECD.