The role of desmoplakin for neuronal function in the dentate gyrus and anxiety-

related behavior

Keisuke Otsubo[a,#], Naoko Sakashita[a,#], Yuki Nishimoto[a,#], Yo Sato[a], Katsunori

Kobayashi[b], Kanzo Suzuki[a], Eri Segi-Nishida[a,*]

[a]Department of Biological Science and Technology, Faculty of Advanced Engineering,

Tokyo University of Science, Tokyo, Japan

[b] Department of Pharmacology, Graduate School of Medicine, Nippon Medical School,

Tokyo 113-8602, Japan

[#] these authors equally contributed to this work

*Correspondence to:

Eri Segi-Nishida,

6-3-1 Niijuku, Katsushika-ku, Tokyo, Japan, 125-8585

Email: eri.segi.nishida@rs.tus.ac.jp

Running title: Role of desmoplakin in DG

Abstract

Desmoplakin (Dsp) is a component of desmosomal cell-cell junctions that interacts

with the cadherin complex and cytoskeletal intermediate filaments. In addition to its

function as an adhesion component, Dsp is involved in various biological processes,

such as gene expression, differentiation, and migration. Dsp is specifically expressed in

the hippocampal dentate gyrus (DG) in the central nervous system. However, it is

unclear how Dsp impacts hippocampal function and its related behaviors. Using an

adeno-associated virus knockdown system in mice, we provide evidence that Dsp in

the DG maintains hippocampal functions, including neuronal activity and adult

neurogenesis, and contributes to anxiolytic effects. Dsp protein is mostly localized in

mature granule cells in the adult DG. Dsp knockdown in the DG resulted in a lowered

expression of an activity-dependent transcription factor FosB, and an increased

expression of mature neuronal markers, such as calbindin. In addition, the suppression

of Dsp decreases serotonin responsiveness at the DG output mossy fiber synapses and

impairs adult neurogenic processes in the subgranular zone of the DG. Moreover, DG-

specific Dsp knockdown mice showed an increase in anxiety-like behaviors. Taken

together, this research uncovers an unexplored function for Dsp in the central nervous

system and suggests that Dsp in the DG may function as a regulator to maintain proper

neuronal activation and contribute to the adaptation of emotion-related behavior.

Keywords: anxiety; dentate gyrus; desmoplakin; hippocampus; knockdown;

neurogenesis

Introduction

Desmosomes are adhesive intercellular junctions that provide strong adhesion between cells and are abundant in tissues exposed to mechanical stress, such as the heart and skin. Mutations in these factors cause skin and heart vulnerability (Kowalczyk and Green, 2013). Desmosomal components comprise transmembrane cadherins, such as desmoglein and desmocollin, and intracellular proteins, such as plakoglobin, plakophilin, and desmoplakin (Dsp) (Garrod and Chidgrey, 2008; Broussard et al., 2015). Dsp is a family of plakin proteins that interacts with the cadherin/plakoglobin/plakophilin complex through the N-terminal domain and with an intermediate filament cytoskeleton through the C-terminal domain (Delva et al., 2009). In addition to mediating cell-to-cell adhesion, desmosomes function as cell signaling regulators involved in cell proliferation and differentiation (Johnson et al. 2014). In particular, Dsp is involved in processes such as gene expression, differentiation, and microtubule dynamics, indicating its diverse biological functions in addition to its function as an adhesion component (Bendrick et al., 2018; Dubash et al., 2016; Lombardi et al., 2016; Patel et al., 2014).

In the central nervous system, Dsp is specifically expressed in the hippocampal dentate gyrus (DG) in the brain (Yamasaki et al., 2008). The DG gates the flow of neocortical information into the hippocampal CA3 region. The DG is one of the few areas of the mammalian brain where adult neurogenesis occurs (Drew et al., 2013; Zhao et al., 2008). Neurogenesis is a multi-step process consisting of proliferation, differentiation, survival, and maturation and is regulated by various factors, including signals from adjacent granule cells (Kempermann et al., 2015). The neurogenic region in the DG is the subgranular zone (SGZ), which is adjacent to the mature granule cell layer. It has been reported that Dsp is characterized as a molecular maker for neuronal

maturation in the DG of hippocampus (Kobayashi et al., 2010). The expression level of

Dsp in the DG increases with postnatal development, similarly to the patterns of other

DG maturation markers, calbindin and tryptophan-2,3-dioxygenase 2 (Tdo2). Dsp

transcription is directly regulated by Bcl11b, a transcription factor that plays an

important role in postnatal development of the DG (Simon et al., 2012). Forebrain

neuron-specific knockout of Dsp in Emx1-Cre mice impairs progenitor proliferation

and neuronal differentiation at 14 days postnatal age, indicating that Dsp contributes to

the development of DG neurons (Simon et al., 2012). However, whether Dsp regulates

the function of mature DG neurons and the mechanisms through which Dsp affects the

process of adult neurogenesis in the neighboring SGZ region remain unclear.

The hippocampus is a limbic structure implicated in the regulation of stress and

pathophysiology/treatment of anxiety and depression (Segi-Nishida and Suzuki, 2022).

We have previously demonstrated that long-term antidepressant treatment suppresses

the expression of *Dsp* in the DG, indicating that Dsp may be an important molecule in

antidepressant effect (Kobayashi et al., 2010; Imoto et al., 2017). Neuron-specific

knockout mice of norbin, a known G-protein regulator, show depression-like behavior

and decreased expression of *Dsp* in the DG. These previous studies indicate that Dsp

plays a key role in emotion-related behaviors in the DG. Nevertheless, whether Dsp in

the DG during adulthood is involved in emotion-related behaviors such as depression

and anxiety is unclear.

In this study, we investigated the effect of Dsp knockdown in the DG on

neuronal activation, adult hippocampal neurogenesis, and emotion-related behaviors

using an adeno-associated virus (AAV) knockdown system. We found that Dsp in the

adult hippocampal DG maintains the function of DG neurons, regulates neuronal

differentiation during adult neurogenesis, and behaviorally contributes to anxiolytic

effects.

Materials and Methods

Experimental animals

Seven-to-eight-week-old male C57BL/6N mice (23–25 g) were purchased from Japan

SLC (RRID5295404; Hamamatsu, Japan). The mice were housed in groups of six per

cage (32 cm length × 11 cm width × 13.5 cm height). All the mice were housed under

standard conditions (temperature: 24 ± 2 °C, humidity: 55% ± 5%) with a 12 h

light/dark cycle and *ad libitum* access to water and food (MR standard, Nosan

Corporation, Yokohama, Japan). The weight gain of each mouse was recorded during

the experiments to monitor their physical condition. All mice were habituated for longer

than one week before the experimental procedures were performed. Animal use and

procedures were conducted in according the Tokyo University of Science regulations

on animal care and use in research and were approved by the Animal Care and Use

Committee of Tokyo University of Science (approval numbers K20009, K21007, and

K22007).

AAV preparation and hippocampal administration

AAVs (serotype rh10) expressing artificial microRNAs (miRNAs) targeting Dsp or

control miRNAs with Emerald Green Fluorescent Protein (EmGFP) under the EF1α

promoter were generated as follows: We designed the sequences for artificial

microRNAs for Dsp according to the BLOCK-iT RNAi Designer website (Thermo

Fisher Scientific). First, the DNA fragment encoding miRNA for *Dsp* (5'-TGC TGT

TGA GAT GGA CAT CAA TGC ACG TTT TGG CCA CTG ACT GAC GTG CAT

TGG TCC ATC TCA A-3' and 5'-CCT GTT GAG ATG GAC CAA TGC ACG TCA

GTC AGT GGC CAA AAC GTG CAT TGA TGT CCA TCT CAA C-3') was subcloned

into pcDNA6.2-GW-EmGFP-miR using BLOCK-iT Pol II miR RNAi Expression

Vector Kit (Invitrogen). The EmGFP and Dsp miRNA fragments were amplified by

PCR from this vector and inserted between the KpnI and EcoRV sites of the pAAV-

EF1α-DIO EYFP vector (#27056; Addgene). The resulting plasmid was named pAAV-

EF1α-EmGFP-miR-Dsp. The DNA fragment encoding control microRNA (5′-TGC

TGA AAT GTA CTG CGC GTG GAG ACG TTT TGG CCA CTG ACT GAC GTC

TCC ACG CAG TAC ATT T-3′ and 5′-CCT GAA ATG TAC TGC GTG GAG ACG

TCA GTC AGT GGC CAA AAC GTC TCC ACG CGC AGT ACA TTT C-3′), which

can form a hairpin structure processed into mature microRNA but is predicted not to

target any known vertebrate gene, was used to generate an AAV vector expressing

control microRNA. TheBLOCK-iT Pol II miR RNAi Expression Vector Kit was used

instead of those encoding Dsp miRNA. AAV particles were produced using the

previously described procedure with minor modifications (Shinohara et al., 2018).

Briefly, the AAVrh10 expression plasmid (Addgene #112866), pHelper vector

(#240071, Agilent Technologies, Santa Clara, CA), and the constructed AAV-vector

plasmid were transfected into AAV-293 cells (#240073, Agilent) using Lipofectamine

2000 (Thermo Fisher Scientific). Transfected AAV-293 cells were collected 72 h after

transfection, resuspended in artificial cerebrospinal fluid, frozen, and thawed three

times. Cell debris was eliminated by centrifugation at $10,000 \times g$ for 10 min and the

supernatant containing the virus was collected. The viral suspension was incubated with

benzonase (Merck Millipore, Burlington, MA, USA) to degrade residual DNA. The

prepared AAV solution was aliquoted and stored at -80 °C. The AAV titer was

quantified using PCR (5′-TGA GTC ACC CAC ACA AAG GA-3′ and 5′-CCA AGC

TGG CCT AAC TTC AG-3′) after proteinase K treatment (Merck Millipore). Under

anesthesia with a mixture of medetomidine (0.3 mg/kg, ZENOAQ, Japan), midazolam

(4 mg/kg, Sandoz, Japan), and butorphanol (5 mg/kg, Meiji, Japan), the mouse brain

was fixed with stereotaxic instruments (Narishige, Tokyo, Japan). An AAV solution of

500 nL per injection site (5.0×10^8 copies) was stereotaxically injected at two sites in

each hemisphere into the DG using a PV-820 Pneumatic PicoPump (World Precision

Instruments, Hessen, Germany) through a glass micropipette made with a PN-30

micropipette puller (Narishige). The stereotaxic coordinates were targeted to the ventral

DG: 3.2 mm posterior to the bregma, 2.7 mm lateral to the midline, and 3.0 mm ventral

to the skull surface at the bregma, according to the mouse brain atlas (Paxinos and

Franklin, 2012). Data were excluded from the analysis if fluorescence was not observed

in the DG.

Electroconvulsive stimulation and drug treatment

Bilateral ECS (current, 25–35 mA; shock duration, 1 sec; frequency, 100 pulse/sec;

pulse width, 0.5 msec) was administered with a pulse generator (ECT Unit; Ugo Basile,

VA, Italy) to mice as described previously (Imoto et al., 2017). ECS was administered

4 times a week. Fluoxetine hydrochloride (LKT Labs, St. Paul, MN, USA) was

dissolved in drinking water and orally applied at a dose of 22 mg/kg/day for 4 weeks

as described previously (Kobayashi et al., 2010).

RNA extraction and real-time PCR

Seven to eight weeks after AAV infection, the mice were decapitated under deep

anesthesia, coronal brain slices (1 mm) were cut using a tissue slicer, and the DG of the

hippocampus was dissected. Total RNA was extracted using the Reliaprep RNA Cell

Miniprep System (Promega, Madison, WI, USA) and reverse transcribed using

ReverTra Ace (Toyobo, Osaka, Japan). Subsequent, real-time PCR was performed

using the StepOne system (Applied Biosystems, Foster City, CA) and Thunderbird

SYBR qPCR mix (Toyobo). The expression levels of each gene were quantified using

standardized external dilutions. The relative expression levels of the target genes were

normalized to those of 18S rRNA or *Gapdh*. The specificity of each primer set was

confirmed by melt-curve analysis, and the product size was examined using gel

electrophoresis. Table 1 summarizes the primer sequences for each gene.

Electrophysiological analysis

Mice were decapitated under deep anesthesia 7–8 weeks after AAV infection, and both

hippocampi were isolated. Transverse hippocampal slices (380 μm) were cut in ice-cold

saline using a tissue slicer (7000smz, Campden Instruments Ltd., Leics, UK). The slices

were then incubated for 30 min at 30 °C and maintained in a humidified interface-

holding chamber at room temperature before use. Intense GFP fluorescence in the DG

and the mossy fiber (MF) tract was confirmed, and electrophysiological recordings

were performed as described (Kobayashi et al., 2010, 2020). Electrophysiological

recordings were made in a submersion-type chamber maintained at 27.0–27.5 °C and

superfused at 2 mL/min with recording saline composed of (in mM): NaCl, 125; KCl,

2.5; NaH_2PO_4, 1.0; $NaHCO_3$, 26.2; glucose, 11; $CaCl_2$, 2.5; $MgCl_2$, 1.3 (equilibrated

with 95% O_2 / 5% CO_2). Field excitatory postsynaptic potentials (EPSPs) arising from

the MF synapses were evoked by stimulating the dentate granule cell layer with bipolar

tungsten electrodes and were recorded from the stratum lucidum of CA3 using a glass

pipette filled with 2 M NaCl. The amplitude of field EPSPs was measured with a 0.5-

ms window positioned at 70–80% of the peak of the baseline field EPSPs. More than

85% block of EPSP by an agonist of group II metabotropic glutamate receptors (2S,2'R,3'R)-2-(2',3'-dicarboxycyclopropyl) glycine (DCG-IV, 1 μM) was the criterion used to identify MF input. A single electrical stimulation was delivered at a frequency of 0.05 Hz. The initial slope of the EPSPs was measured through analysis. All recordings were performed using a Multiclamp 700 B amplifier (Molecular Devices, Sunnyvale, CA, USA), filtered at 2 kHz, and stored on a personal computer via an interface (digitized at 10 kHz). Data were obtained from distinct samples. Serotonin hydrochloride was purchased from Sigma-Aldrich.

Sample isolation for immunohistochemistry

Five'-bromo-2'-deoxyuridine (BrdU, 150 mg/kg, Nacalai Tesque, Japan) was intraperitoneally administered to mice once 2 h before sacrifice to evaluate cell proliferation to label the dividing cells, or four times on 14 and 15 days (twice a day) before sacrifice to evaluate cell survival.

For Dsp immunostaining, the mice were deeply anesthetized and transcardially perfused with cold saline. The brains were then stored at -80°C until further use. Serial fresh frozen sections (18 μm thick) were cut through the entire hippocampus using a cryostat (Leica 1510, Leica Microsystems, Wetzlar, Germany) and stored at -80°C. For other immunostaining procedures, mice were deeply anesthetized and perfused with cold saline followed by 4% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4) and were post-fixed at 4 °C for 48 h. The brains were then cryoprotected in 20% sucrose for 48 h and in 30% sucrose for 24 h and stored at -80°C until further use. Serial sections (30 μm thick) were cut through the entire hippocampus using a cryostat and stored in 30% glycerol and 30% ethylene glycol in 0.02 M phosphate buffer (pH 7.4) at -20 °C until staining.

Immunohistochemistry

For Dsp or Dsp/NeuN immunostaining, fresh frozen sections were fixed with 100%

methanol for 20 min at 4 °C. After fixation, the sections were washed with phosphate-

buffered saline (PBS), blocked using 10% equine serum in PBS, and incubated with

monoclonal mouse anti-Dsp (1:50; Progen, 61003, RRID:AB_2920666), rabbit anti-

NeuN (1:1000, Abcam, ab177487, RRID:AB_2532109) at 4 °C overnight. After

washing with PBS, the sections were incubated with donkey anti-mouse IgG antibodies

conjugated to Alexa Fluor 647 (1:300; Thermo Fisher Scientific, RRID:AB_162542)

and donkey anti-rabbit IgG antibodies conjugated to Alexa Fluor 555 (1:1000; Thermo

Fisher Scientific, RRID:AB_2535853) for 60 min. The sections were then incubated

with 4',6-diamidino-2-phenylindole (DAPI; 1:10,000; Merck Millipore) and mounted

on slides with Mowiol (Merck Millipore) after washing with PBS.

For FosB, NeuroD1, and doublecortin (DCX) immunostaining, the 30 μm thick

floating sections were washed with PBS and blocked using 10% equine serum

(SH30074; Cytiva) in PBS containing 0.3% Triton X-100 (PBST) at room temperature

for 60 min, followed by overnight incubation with rabbit anti-FosB (1:1000; Abcam,

ab184938, RRID: AB_2721123), rabbit anti-NeuroD1 (1:1000; Abcam, ab213725,

RRID:AB_2801303), rabbit anti-doublecortin (DCX) (1:1000; Cell Signaling

Technology, Danvers, MA, USA; RRID:AB_561007) at 4 °C. For BrdU

immunostaining, the floating sections were incubated in 50% formamide in 2×saline

sodium citrate buffer (SSC) for 2 h at 60 °C, followed by incubation in 2 M hydrogen

chloride at 37 °C for 30 min, and neutralized with 0.1 M boric acid (pH 8.5) at room

temperature for 10 min before blocking. After blocking, the sections were incubated

overnight with monoclonal rat anti-BrdU (1:8000; Abcam, ab6326, RRID:AB_305426)

at 4°C. After washing with PBST, the sections were incubated with donkey anti-rabbit

IgG antibody conjugated to Alexa Fluor 555 (1:1000 for NeuroD1 and FosB

immunostaining; Thermo Fisher Scientific, RRID:AB_2535853) and donkey anti-rat

IgG antibody conjugated to Alexa Fluor 555 (1:1000 for BrdU immunostaining;

Abcam, RRID: AB_2813834) for 60 min. After washing with PBST, the sections were

incubated with DAP (1:10000) and mounted on slides with Mowiol after washing with

PBS. For DCX immunostaining, the sections were incubated with biotinylated goat

anti-rabbit IgG (1:200; Vector Laboratories Inc., Burlingame, CA, USA; BA1000,

RRID: AB_2313606) for 60 min. The sections were washed with PBST and incubated

with the ABC Vectastain Kit (Vector), and antigen detection was performed with 0.06%

3,3'-diaminobenzidine (Wako) staining. After washing, the sections were mounted on

slides using Entellan New (Merck Millipore).

Quantification of BrdU-, NeuroD1-, DCX-, and FosB-positive cells

Computer-assisted image analysis (ImageJ, NIH, Bethesda, MD, USA; QuPath,

Bankhead et al., 2017) was used for cell analysis. For quantification of FosB-positive

cells, three sections of DG-expressing EmGFP were photographed under a microscope

(BZX-700, Keyence, Osaka, Japan). The number of FosB-positive cells was counted

and presented as the number of cells per 10,000 μm^2 area in the granule cell layer of

the DG. For BrdU-labeled cell quantification, 6–8 sections of DG-expressing EmGFP

were photographed under a microscope. To evaluate cell proliferation, the number of

BrdU-positive cells was counted in the SGZ and presented as the number of cells per

1,000 μm along the SGZ. To evaluate cell survival, the number of BrdU-positive cells

in the granule cell layer of the DG was counted and presented as the number of cells

per 100,000 μm^2 of the granule cell layer. For NeuroD-positive cell quantification, four

sections of the DG expressing EmGFP were photographed using a microscope. The

number of NeuroD1-positive cells in the SGZ was counted and presented as the number

of cells per 1,000 μm along the SGZ. For DCX-positive cell quantification, 2–3 sections

of DG-expressing EmGFP were photographed using a microscope. DCX-positive cells

in which dendrites reached the molecular layer and underwent complex processes were

subcategorized as DCX-positive cells with long and branched dendrites. The total

number of DCX-positive and DCX-positive cells with long dendrites was counted in

the SGZ and presented as the number of cells per 100 μm along the SGZ.

Behavioral experiments

The mice were habituated to the testing room for at least 30 min before the behavioral

test. During the test, the movement of each mouse was recorded using a digital video

camera. All records were stored and analyzed using the SMART 3.0 video tracking

software (Panlab, Barcelona, Spain, RRID: SCR_002852). The apparatus was

thoroughly cleaned with 70% ethanol after the removal of each mouse. The time course

of each experiment is shown in Fig. 6A.

The open field test (OFT) was performed for 10 min using an apparatus

composed of opaque white walls and a floor (50 cm × 50 cm × 20 cm) illuminated at

an intensity of 80 lux in the center of the floor. Each mouse was placed in the center of

an open-field arena, and locomotor activity was recorded for 10 min. The total distance

moved, relative time spent in the central zone, and entry time into the central zone were

measured. The floor was divided into 25 squares, and the four central squares were

defined as the central zone.

The forced swim test (FST) was performed in a cylindrical container (13 cm

diameter, 25.5 cm height) filled with water to a height of 18 cm. Water (23–25 °C) was

replaced between the trials. The FST was performed and recorded for 10 min using a

digital video camera. The duration of immobility was then quantified.

The spontaneous locomotor activity (LMA) test was performed in a transparent

home cage without bedding (20 cm length × 12.5 cm width) over 60 min. The total

distance moved during the last 30 min was tracked.

The compartment social interaction test (SIT) was performed in an SIT

apparatus (LE894T, Panlab) consisting of three chambers (42 cm × 60 cm × 22 cm) and

two partitions (5.5 cm × 0.15 cm × 17 cm). Each mouse was placed in the middle

chamber and allowed to explore the chamber freely for 5 min. Immediately after

habituation, an unfamiliar mouse (a 7–8-week-old male C57BL/6N mouse) was

introduced into one of the two side chambers enclosed in a wire cage. An identical

empty wire cage was placed on the opposite side of the chamber. After placement, the

side doors were opened simultaneously, and the test mouse was allowed to explore the

entire three-chamber arena for 10 min. The time spent in each compartment was

measured.

An elevated plus maze (EPM) test was performed in an EPM apparatus (EPM-

X, Muromachi Kikai, Japan) consisting of two open arms (30 cm × 6 cm, 40 lux) and

two closed arms (30 cm × 6 cm × 15 cm, 20 lux) extending from a central platform (6

cm × 6 cm). The maze was elevated to 40 cm above the floor. Each mouse was placed

in the center of the maze, facing the open arm, and allowed to explore freely for 10 min.

The total distance and time spent in the open arms were measured using Smart 3.0.

Statistical analyses

All data are presented as the mean ± standard error of the mean. Statistical analyses

were performed using an unpaired Student's t-test, Welch's t-test, or two-way analysis

317 of variance (ANOVA). Statistical significance was set at $P < 0.05$. Before performing

318 t-test, normality of the data was examined using the Shapiro–Wilk test. If the data did

319 not pass the normality test, a nonparametric test, the Mann–Whitney U test, was

320 performed. Detailed statistical data are presented in Supplementary Table 1 (Table S1).

321 All analyses were performed using the PRISM 9 software (GraphPad, San Diego, CA,

322 USA).

Results

Desmoplakin is localized around the cell bodies of granule cells in the hippocampal DG and is responsible for chronic antidepressant stimulation

Dsp, a granule cell-specific maturation marker (Yamasaki et al., 2008; Kobayashi et al., 2010), is highly and specifically expressed in the DG of the hippocampus of the central nervous system (Supplemental Fig. 1, Allen Institute for Brain Science, 2004). Although we previously observed that the gene expression level of Dsp significantly decreased in the DG of mice treated with antidepressant stimulation (Kobayashi et al., 2010; Imoto et al., 2017), the localization of Dsp in the DG was not confirmed. Therefore, we initially examined the cellular localization and cell type of Dsp protein expression. Immunohistological analysis revealed that Dsp was localized around the cell bodies (Fig. 1A and 1C). Double immunohistochemical staining with Dsp and NeuN showed that Dsp immunoreactivity signals were detected in NeuN-positive cells in the outer DG, whereas Dsp signals were sparse in NeuN-negative and NeuN-positive cells in the inner DG, which predominantly contains neuronal progenitors and immature neurons (Fig. 1A [a], [b]). These findings suggest that Dsp is mostly localized in mature granule cells in the DG. Dsp immunostaining was only observed after methanol fixation; therefore, the localization of other neuronal markers was not investigated.

Furthermore, we investigated whether Dsp expression was regulated by antidepressant stimulation at the protein level. The expression of *Dsp* was reduced by chronic administration of electroconvulsive seizure (ECS) or a selective serotonin reuptake inhibitor (SSRI) (Fig. 1B), as previously reported (Imoto et al., 2017). Immunohistological analysis confirmed that the cellular localization of Dsp was reduced after repeated administration of ECS (Fig. 1C).

Dsp expression knockdown in the DG by an adeno-associated virus (AAV) carrying Dsp miRNA

To explore the role of Dsp in hippocampal function, we performed an AAV-mediated knockdown of Dsp in the DG. For this purpose, we generated AAVrh10 expressing EGFP and an artificial miRNA targeting Dsp mRNA (AAV-Dsp miRNA; Fig. 2A). The AAV rh10 serotype shows high transduction efficiency into granule cells in the DG (Sehara et al., 2022). Seven to eight weeks after the injection of AAV-Dsp or control miRNA into the DG, EGFP signals, which were co-expressed with Dsp miRNA, were observed in the granule cells but not in the immature neuronal marker DCX-positive cells in the DG (Fig. 2A). *Dsp* mRNA expression significantly decreased in the AAV-Dsp miRNA group (Fig. 2B). Moreover, immunohistological analysis showed a decrease in Dsp immunoreactivity in the DG following AAV-Dsp miRNA injection (Fig. 2C), demonstrating that the sequence of artificial miRNAs targeting Dsp used in this study has the potential to knockdown Dsp expression.

Changes in activity- and mature-related marker expression in the DG and alteration the sensitivity to serotonin at the MF synapse by Dsp knockdown

First, we examined the direct effect of Dsp knockdown on the activity of mature DG neurons. FosB, including ΔFosB, is induced in response to many types of chronic stimuli and accumulates because of its high stability (Nestler et al., 1999; Lamothe-Molina et al., 2022). Consequently, we performed immunohistochemical analysis using an anti-FosB antibody to monitor neuronal activity in the DG under normal home-cage conditions. FosB immunoreactivity in the granule cell layer of the DG significantly decreased in the Dsp knockdown group (Fig. 3A).

Because Dsp is a granule cell-specific DG maturation marker (Yamasaki et al.,

2008; Kobayashi et al., 2010), we examined the expression of other markers, including

Calb, Tdo2, Ntf3, and Il1r1, in the DG. The expression of *Calb, Tdo2,* and *Ntf3*

significantly increased in the DG of the Dsp knockdown group (Fig. 3B). These results

indicate that Dsp in the DG is involved in the regulation of neuronal activity and

expression of maturation markers.

Dsp interacts with the desmosomal component γ-catenin (plakoglobin). We

hypothesized that the loss of Dsp may regulate catenin-related transcription via

enhancement of γ-catenin translocation into the nucleus (Garcia-Gras et al., 2006).

Therefore, we examined the gene expression of *Hdac2* and *Tiam1*, which are subject to

transcriptional regulation by catenin, and found that their expression increased in the

Dsp knockdown group (Fig. 3C).

Antidepressant treatments substantially change the physiological properties of

the synapses formed by DG output MFs onto CA3 pyramidal cells, concomitantly with

the reduction in Dsp expression (Kobayashi et al., 2010; Imoto et al., 2017).

Electrophysiological analyses of MF-CA3 synaptic transmission were then performed

using acute hippocampal slices (Fig. 4A). No significant changes in the ratio of field

EPSP to fiber volleys were observed (Fig. 4B), suggesting that the basal transmission

efficacy at the MF synapse was not altered by Dsp knockdown. Antidepressant

treatment attenuates the synaptic facilitation induced by repetitive stimulation and

augments serotonergic modulation at the MF synapse (Kobayashi et al., 2010; Imoto et

al., 2017; Kobayashi et al., 2020). Dsp knockdown did not change synaptic facilitation

after 1 Hz stimulation (Fig. 4C and D). Bath-applied serotonin induced the potentiation

of field EPSPs in the AAV control group (Fig. 4E), consistent with the findings of

previous studies. The magnitude of serotonin-induced potentiation was significantly

reduced in the Dsp knockdown group (Fig. 4E and F). Collectively, these findings

indicate that suppression of Dsp alters neuronal activity in the DG and sensitivity to

serotonin at the MF synapse.

Changes in neurogenic activity in the DG by Dsp knockdown

The activation state of mature neurons in the DG affects the adjacent neurogenic

processes (Segi-Nishida and Suzuki, 2022). Since the above results suggest that Dsp in

mature DG neurons regulates neuronal activation, we explored the influence of Dsp

knockdown on hippocampal neurogenesis. Neurogenesis involves cell proliferation,

neuronal differentiation, and survival. First, we examined the effect of Dsp knockdown

on cell proliferation. BrdU was administered 2 h before the sacrifice to label

proliferating cells (Fig. 5A, left). BrdU-positive cells were detected in the SGZ of the

DG (Fig. 5A, middle), and no difference was observed in the number of proliferating

cells between the control and Dsp knockdown groups (Fig. 5A, right). We then

examined the expression of Neuro D1, which is expressed in progenitors/neuroblasts

and immature neurons in the adult DG. The number of NeuroD1-positive cells in the

SGZ was significantly decreased in the Dsp knockdown group (Fig. 5B). We further

examined the expression of the immature neuronal marker, DCX, in the DG. The total

number of DCX-positive cells and those with long dendrites decreased in the Dsp

knockdown group (Fig. 5C). Because most newborn cells in the DG die within 2–3

weeks following proliferation in mice (Ueno et al., 2019), we evaluated the effect of

Dsp knockdown on the survival of newborn cells 2 weeks after BrdU administration

(Fig. 5D, left). The number of BrdU-positive cells in the DG did not differ between the

control and Dsp knockdown groups (Fig. 5D, right).

Influence of Dsp knockdown on anxiety-related behaviors

The ventral DG is a key brain region involved in emotional behaviors, including anxiety- and depression-like behaviors (Sahay and Hen, 2007; Levone et al. 2015; Anacker et al. 2018). To examine the influence of Dsp knockdown in the DG on emotion-related behaviors, we conducted three sets of behavioral experiments (Fig. 6A). First, we performed an open-field test to assess behavior in an anxiety-provoking environment. Dsp knockdown in the DG suppressed locomotor activity in the open field box, and a suppressive trend was observed; however, it was not significant for the time spent in the central area or the number of entries into the central area (Fig. 6B). Next, we evaluated immobility time in the FST to assess depression-like behavior. No difference in immobility time between the control and Dsp knockdown groups was observed (Fig. 6C). Another set of experiments was conducted on locomotor activity in a home cage, social interaction test, and elevated plus maze test. Contrary to the results of the open field box, no change in locomotor activity between the control and knockdown groups was observed (Fig. 6D), indicating normal locomotor activity under ordinal circumstances and decreased locomotor activity in anxiety-provoking environments following Dsp knockdown. In the three-chamber social interaction test, the control mice spent more time around the area with an unfamiliar mouse than around the area with empty cages. This social interaction behavior was comparable to that observed in the Dsp knockdown mice (Fig. 6E). In the elevated plus maze test, a reduction in both the time spent in the open arm and locomotor activity was observed in Dsp knockdown mice (Fig. 6F). These results demonstrate that Dsp knockdown mice in the DG shows anxiety-like behavior. Also, these findings can implicate that expression of Dsp in the DG plays an anti-anxiety role.

Discussion

This study aimed to investigate the role of Dsp expressed in the DG in hippocampal function and emotion-related behaviors. We showed that Dsp is localized mostly in mature granule cells in the DG and that knockdown of Dsp decreases the expression of neuronal activity marker and serotonin responsiveness in DG neurons while enhancing the expression of mature neuronal markers. In addition, we demonstrated that Dsp knockdown in the DG impairs adult neurogenic processes in the adjacent SGZ and that these hippocampal changes contribute to anxiety-like behaviors. These findings suggest that Dsp in the DG maintains the activity of mature neurons, regulates neuronal differentiation in the adult neurogenic process, and contributes to anxiolytic effects. This study is the first to demonstrate the role of Dsp in neuronal activity and emotion-related behaviors in the DG.

Expression of desmoplakin in the DG

Since Dsp could play an important role in cell-to-cell adhesion and subsequent intracellular regulatory signaling, we initially confirmed its localization in the brain. In the central nervous system, *Dsp* was specifically expressed in the hippocampal DG (Supplemental Fig. 1). Immunohistological studies showed that Dsp was localized in the cell bodies of most dentate granule cells (Fig. 1A). Moreover, Dsp expression was lower in the inner DG than in the inner DG. *Dsp* expression in the adult DG was similar to that of the DG maturation markers *Calb* and *Tdo2* (Kobayashi et al., 2010), suggesting that Dsp is expressed in mature DG neurons. These results are consistent with those of a previous study that demonstrated that Dsp colocalizes with the post-mitotic neuron marker Bcl11b but not with the neural stem cell/progenitor marker SOX2 in the DG at 14 days of age (Simon et al., 2012).

Desmosome components include transmembrane cadherins and intracellular

proteins such as plakoglobin, plakophilin, and Dsp (Garrod and Chidgrey, 2008;

Broussard et al., 2015). Electron microscopy images of the rat DG showed no

desmosome structures between the cell bodies (Laatsch and Cowan, 1966). Therefore,

Dsp in the DG may form a complex different from that in epithelial desmosomes. The

expression and interactions of each desmosome component in the DG will be examined

to elucidate the localization and function of Dsp in neurons.

Role of desmoplakin in mature DG neurons

We found that Dsp knockdown in the DG suppressed FosB expression (Fig. 3A); FosB

is useful for monitoring steady-state activation since FosB, including ΔFosB, is induced

in response to many types of stimuli and shows high stability (Nestler et al., 1999;

Lamothe-Molina et al., 2022). In contrast, Dsp knockdown increased the expression of

the mature neuronal markers *Calb1*, *Tdo2*, and *Ntf3* (Fig. 3B), which are negatively

regulated by strong neuronal activation, such as in the ECS and epileptic seizures

(Imoto et al., 2017; You et al., 2017). Electrophysiological analysis showed that Dsp

suppression did not alter the basal transmission efficiency at the MF-CA3 synapses;

however, it reduced serotonin-induced potentiation (Fig. 4). Synaptic potentiation by

serotonin is mediated by presynaptic 5-HT4 receptor-cAMP signaling in the MF

(Kobayashi et al., 2008; 2020). Collectively, reduced neuronal activity due to Dsp

knockdown may enhance mature marker expression and alter serotonin-activated

intracellular signaling in the DG neurons.

Dsp is involved in processes such as gene expression, differentiation, and cell

migration (Bendrick et al., 2018; Dubash et al., 2016; Lombardi et al., 2016; Patel et

al., 2014). Dsp interacts with plakoglobin, also known as γ-catenin (Delva et al., 2009).

Suppression of Dsp expression in atrial myocyte cell lines leads to nuclear localization

of plakoglobin (Garcia-Gras et al., 2006). We showed that the expression of *Hdac2* and

Tiam1, which are transcriptionally regulated by catenin signaling, was increased by the

suppression of Dsp expression (Fig. 3C). HDAC2 negatively regulates gene expression

involved in synapse formation and synaptic plasticity in neurons (Guan et al., 2009;

Wang et al., 2018); Tiam1 is a Rho-guanine nucleotide exchange factor (Rho-GEF),

important for synaptic structure and function in the hippocampal DG (Rao et al., 2019).

Further detail analysis will be required to characterize the Dsp-mediated signaling

pathway. In addition, confirmation of the nuclear translocation of plakoglobin (γ-

catenin) will clarify whether changes in the expression of these factors are mediated by

catenin signaling.

Influence of desmoplakin knockdown on the neurogenic process in the DG

Adult neurogenesis occurs in the SGZ of the DG. Neurogenesis is regulated by various

signals from surrounding neurons (Kempermann et al., 2015). We investigated the role

of Dsp in the DG neurons in adult neurogenesis. Suppression of Dsp reduced the

number of NeuroD1-positive neural progenitors and DCX-positive immature neurons,

whereas the number of proliferating cells remained unchanged (Fig. 5). EGFP signals,

which were coexpressed with Dsp miRNA, were not observed in DCX-positive cells of

the DG (Fig. 2A), suggesting that Dsp in DG neurons indirectly regulates neurogenic

processes in the adjacent SGZ. Forebrain neuron-specific ablation of Dsp impairs

hippocampal neurogenesis on postnatal day 14 (Simon et al., 2012). During postnatal

development in this mutant line, cell proliferation was suppressed, whereas the number

of NeuroD1-positive progenitors increased. All these findings suggest that Dsp is

important for the regulation of cell proliferation during developmental neurogenesis

and for neural differentiation during both developmental and adult neurogenesis. The

reduction in the number of neural progenitors and immature neurons observed in this

study may be due to the inhibition of the initial process of neural differentiation or

acceleration of the differentiation rate, which allowed them to progress to maturity

without sufficient differentiation. Subsequent examination of neuronal differentiation

using surviving BrdU-labeled cells will clarify the role of Dsp in adult neurogenesis.

How does Dsp indirectly regulate neuronal differentiation in post-mitotic

neurons? Neuronal activation, such as that in the ECS, promotes the expression of adult

neurogenic factors such as *Vegf*, *Fgf2*, and *Wnt2* in the DG (Newton et al., 2003;

Warner-Schmidt et al., 2007). Therefore, the suppression of neuronal activity in the DG

by inhibiting Dsp expression may reduce the expression of these growth factors.

Furthermore, intracellular skeletal regulation by Dsp may also affect neurogenic niche

formation. The neurogenic niche provides a unique milieu containing an extracellular

matrix, humoral factors, and cell-cell contacts that allow controlled neuronal

development (Kempermann et al., 2015). Altering the cytoskeleton of dentate granule

cells may affect intercellular contacts and extracellular matrix adhesion and inhibit

neurogenic niche formation.

Desmoplakin in the DG as a key regulator for emotion-related behaviors

The DG of the hippocampus controls emotion-related behaviors (Segi-Nishida and

Suzuki, 2023). Although Dsp knockdown in the DG did not affect home cage activity,

depression-like behavior, or social behavior, it reduced the time spent in the open arm

in the EPM test and reduced activity in the novel open field and EPM environment (Fig.

6), indicating an anxious phenotype. The hippocampus plays functionally distinct roles

in the dorsal and ventral regions (Fanselow and Dong, 2010). The dorsal region is

involved in stress-induced learning and memory changes, and the ventral region

contributes to stress responses and emotions such as anxiety. Because AAV was injected

into the ventral DG area in this study (as described in the Methods section), Dsp in the

ventral DG may play an anxiolytic role.

Chronic antidepressant treatment induces anxiolytic-like effects (Santarelli et

al., 2003) and promotes the differentiation of newborn neurons in the DG (Wang et al.,

2008; Ueno et al., 2019), while suppressing the expression of mature markers, including

Dsp, in the DG (Fig. 1; Kobayashi et al., 2010; Imoto et al., 2017). The loss of Dsp

function in the DG may lead to anxiolytic effects; however, surprisingly, anxiety-like

behavior was observed (Fig. 6). Adult neurogenesis in the DG affects anxiety-like

behaviors. Transgenic mice in which adult neurogenesis was suppressed by the

expression of the pro-apoptotic protein BAX in neural progenitor cells show increased

anxiety-related behaviors (Revest et al., 2009). Therefore, suppression of adult

neurogenesis in the DG by Dsp knockdown may promote anxiety-like behaviors.

Furthermore, acute activation of the ventral DG by channelrhodopsin in

proopiomelanocortin (POMC)-Cre mice suppresses anxiety-like behavior in the EPM

(Kheirbek et al., 2013). Because our findings showed that Dsp knockdown in the DG

reduced the expression of the neuronal activity marker FosB (Fig. 3), the suppression

of neuronal activity in the DG by Dsp knockdown may also contribute to anxiety-like

behavior.

Taken together, our findings demonstrate that Dsp in the DG regulates neuronal

function and contributes to anxiolytic effects. This research uncovers an unexplored

function for Dsp in the central nervous system and implies that Dsp in the DG may

function as a regulator maintaining proper neuronal activation. Excessive

antidepressant treatment can exacerbate neuronal activation, resulting in increased

motor activity (Kobayashi et al., 2010; 2011). Considering the results of these studies

and our findings, Dsp knockdown in the DG may function as a negative feedback

mechanism for excessive neuronal activation rather than merely mimicking the effects

of antidepressants or anxiolytics. This is because chronic antidepressant treatment

reduces Dsp expression. Further studies are needed to investigate how Dsp knockdown

alters stress-induced responsibility and emotion-related behaviors. Furthermore, the

elucidation of the regulation of intracellular signaling by Dsp will enhance

understanding of the development and neuronal function of the DG.

Data availability statement

All datasets analyzed in the current study are available from the corresponding author

upon reasonable request.

Ethics statement

Animal use and procedures were conducted in accordance with the Tokyo University

of Science regulations on animal care and use in research and were approved by the

Animal Care and Use Committee of Tokyo University of Science (approval number

K20009, K21007, and K22007).

Author contributions: KO, NS, YN, YS and KK designed the study, conducted the

experiments, analyzed the data, and drafted the manuscript. KS designed the study and

drafted the manuscript. ESN designed the study, analyzed the data, and drafted the

manuscript. All the authors have read and approved the final manuscript.

Funding statements: This work was supported in part by MEXT KAKENHI Grants

20K07090 (ESN), 22K20697 (KS), 23H02801(KK and ESN), and 23K18425(KK).

Kato Memorial Bioscience Foundation (KS), Brain Science Foundation (KS), Takeda

Science Foundation (KS).

Conflict of interest: The authors have no potential conflicts of interest to declare

relevant to this article.

Acknowledgements: We would like to thank *Editage* (www.editage.jp) for the English

language editing.

References

Allen Institute for Brain Science (2004). Allen Mouse Brain Atlas [https://mouse.brain-map.org/experiment/show/69201504]. Available from mouse.brain-map.org.

Anacker, C., Luna, V.M., Stevens, G.S., Millette, A., Shores, R., Jimenez, J.C., Chen, B., and Hen, R. (2018). Hippocampal neurogenesis confers stress resilience by inhibiting the ventral dentate gyrus. *Nature* 559, 98-102. doi:10.1038/s41586-018-0262-4

Bendrick, J.L., Eldredge, L.A., Williams, E.I., Haight, N.B., and Dubash, A.D. (2019). Desmoplakin Harnesses Rho GTPase and p38 Mitogen-Activated Protein Kinase Signaling to Coordinate Cellular Migration. *J Invest Dermatol* 139, 1227-1236. doi:10.1016/j.jid.2018.11.032

Broussard, J.A., Getsios, S., and Green, K.J. (2015). Desmosome regulation and signaling in disease. *Cell Tissue Res* 360, 501-512. doi:10.1007/s00441-015-2136-5

Delva, E., Tucker, D.K., and Kowalczyk, A.P. (2009). The desmosome. *Cold Spring Harb Perspect Biol* 1, a002543. doi:10.1101/cshperspect.a002543

Drew, L.J., Fusi, S., and Hen, R. (2013). Adult neurogenesis in the mammalian hippocampus: why the dentate gyrus? *Learn Mem* 20, 710-729. doi:10.1101/lm.026542.112

Dubash, A.D., Kam, C.Y., Aguado, B.A., Patel, D.M., Delmar, M., Shea, L.D., and Green, K.J. (2016). Plakophilin-2 loss promotes TGF-beta1/p38 MAPK-dependent fibrotic gene expression in cardiomyocytes. *J Cell Biol* 212, 425-438. doi:10.1083/jcb.201507018

Fanselow, M.S., and Dong, H.W. (2010). Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* 65, 7-19. doi:10.1016/j.neuron.2009.11.031

Garcia-Gras, E., Lombardi, R., Giocondo, M.J., Willerson, J.T., Schneider, M.D., Khoury, D.S., and Marian, A.J. (2006). Suppression of canonical Wnt/beta-catenin signaling by nuclear plakoglobin recapitulates phenotype of arrhythmogenic right ventricular cardiomyopathy. *J Clin Invest* 116, 2012-2021. doi:10.1172/JCI27751

Garrod, D., and Chidgey, M. (2008). Desmosome structure, composition and function. *Biochim Biophys Acta* 1778, 572-587. doi:10.1016/j.bbamem.2007.07.014

Guan, J.S., Haggarty, S.J., Giacometti, E., Dannenberg, J.H., Joseph, N., Gao, J., Nieland, T.J., Zhou, Y., Wang, X., Mazitschek, R., Bradner, J.E., DePinho, R.A.,

Jaenisch, R., and Tsai, L.H. (2009). HDAC2 negatively regulates memory
formation and synaptic plasticity. *Nature* 459, 55-60. doi:10.1038/nature07925

Imoto, Y., Segi-Nishida, E., Suzuki, H., and Kobayashi, K. (2017). Rapid and stable
changes in maturation-related phenotypes of the adult hippocampal neurons by
electroconvulsive treatment. *Mol Brain* 10, 8. doi:10.1186/s13041-017-0288-9

Johnson, J.L., Najor, N.A., and Green, K.J. (2014). Desmosomes: regulators of cellular
signaling and adhesion in epidermal health and disease. *Cold Spring Harb Perspect*
Med 4, a015297. doi:10.1101/cshperspect.a015297

Kempermann, G., Song, H., and Gage, F.H. (2015). Neurogenesis in the Adult
Hippocampus. *Cold Spring Harb Perspect Biol* 7, a018812.
doi:10.1101/cshperspect.a018812

Kheirbek, M.A., Drew, L.J., Burghardt, N.S., Costantini, D.O., Tannenholz, L.,
Ahmari, S.E., Zeng, H., Fenton, A.A., and Hen, R. (2013). Differential control of
learning and anxiety along the dorsoventral axis of the dentate gyrus. *Neuron* 77,
955-968. doi:10.1016/j.neuron.2012.12.038

Kobayashi, K., Ikeda, Y., Haneda, E., and Suzuki, H. (2008). Chronic fluoxetine
bidirectionally modulates potentiating effects of serotonin on the hippocampal
mossy fiber synaptic transmission. *J Neurosci* 28, 6272-6280.
doi:10.1523/JNEUROSCI.1656-08.2008

Kobayashi, K., Ikeda, Y., Sakai, A., Yamasaki, N., Haneda, E., Miyakawa, T., and
Suzuki, H. (2010). Reversal of hippocampal neuronal maturation by serotonergic
antidepressants. *Proc Natl Acad Sci U S A* 107, 8434-8439.
doi:10.1073/pnas.0912690107

Kobayashi, K., Ikeda, Y., and Suzuki, H. (2011). Behavioral destabilization induced
by the selective serotonin reuptake inhibitor fluoxetine. *Mol Brain* 4, 12.
doi:10.1186/1756-6606-4-12

Kobayashi, K., Mikahara, Y., Murata, Y., Morita, D., Matsuura, S., Segi-Nishida, E.,
and Suzuki, H. (2020). Predominant Role of Serotonin at the Hippocampal Mossy
Fiber Synapse with Redundant Monoaminergic Modulation. *iScience* 23, 101025.
doi:10.1016/j.isci.2020.101025

Kowalczyk, A.P., and Green, K.J. (2013). Structure, function, and regulation of
desmosomes. *Prog Mol Biol Transl Sci* 116, 95-118. doi:10.1016/B978-0-12-
394311-8.00005-4

Laatsch, R.H., and Cowan, W.M. (1966). Electron microscopic studies of the dentate

gyrus of the rat. I. Normal structure with special reference to synaptic organization.

J Comp Neurol 128, 359-395. doi:10.1002/cne.901280305

Lamothe-Molina, P.J., Franzelin, A., Beck, L., Li, D., Auksutat, L., Fieblinger, T.,

Laprell, L., Alhbeck, J., Gee, C.E., Kneussel, M., Engel, A.K., Hilgetag, C.C.,

Morellini, F., and Oertner, T.G. (2022). DeltaFosB accumulation in hippocampal

granule cells drives cFos pattern separation during spatial learning. *Nat Commun*

13, 6376. doi:10.1038/s41467-022-33947-w

Levone, B.R., Cryan, J.F., and O'Leary, O.F. (2015). Role of adult hippocampal

neurogenesis in stress resilience. *Neurobiol Stress* 1, 147-155.

doi:10.1016/j.ynstr.2014.11.003

Lombardi, R., Chen, S.N., Ruggiero, A., Gurha, P., Czernuszewicz, G.Z., Willerson,

J.T., and Marian, A.J. (2016). Cardiac Fibro-Adipocyte Progenitors Express

Desmosome Proteins and Preferentially Differentiate to Adipocytes Upon Deletion

of the Desmoplakin Gene. *Circ Res* 119, 41-54.

doi:10.1161/CIRCRESAHA.115.308136

Nestler, E.J., Kelz, M.B., and Chen, J. (1999). DeltaFosB: a molecular mediator of

long-term neural and behavioral plasticity. *Brain Res* 835, 10-17.

doi:10.1016/s0006-8993(98)01191-3

Newton, S.S., Collier, E.F., Hunsberger, J., Adams, D., Terwilliger, R., Selvanayagam,

E., and Duman, R.S. (2003). Gene profile of electroconvulsive seizures: induction

of neurotrophic and angiogenic factors. *J Neurosci* 23, 10841-10851.

doi:10.1523/JNEUROSCI.23-34-10841.2003

Patel, D.M., Dubash, A.D., Kreitzer, G., and Green, K.J. (2014). Disease mutations in

desmoplakin inhibit Cx43 membrane targeting mediated by desmoplakin-EB1

interactions. *J Cell Biol* 206, 779-797. doi:10.1083/jcb.201312110

Paxinos, G.F., K. B. J., 2012. 4th ed. Academic Press., Cambridge, MA.

Rao, S., Kay, Y., and Herring, B.E. (2019). Tiam1 is Critical for Glutamatergic

Synapse Structure and Function in the Hippocampus. *J Neurosci* 39, 9306-9315.

doi:10.1523/JNEUROSCI.1566-19.2019

Revest, J.M., Dupret, D., Koehl, M., Funk-Reiter, C., Grosjean, N., Piazza, P.V., and

Abrous, D.N. (2009). Adult hippocampal neurogenesis is involved in anxiety-

related behaviors. *Mol Psychiatry* 14, 959-967. doi:10.1038/mp.2009.15

Sahay, A., and Hen, R. (2007). Adult hippocampal neurogenesis in depression. *Nat*
Neurosci 10, 1110-1115. doi:10.1038/nn1969

Santarelli, L., Saxe, M., Gross, C., Surget, A., Battaglia, F., Dulawa, S., Weisstaub, N.,
Lee, J., Duman, R., Arancio, O., Belzung, C., and Hen, R. (2003). Requirement of
hippocampal neurogenesis for the behavioral effects of antidepressants. *Science*
301, 805-809. doi:10.1126/science.1083328

Segi-Nishida, E., and Suzuki, K. (2022). Regulation of adult-born and mature neurons
in stress response and antidepressant action in the dentate gyrus of the hippocampus.
Neurosci Res. doi:10.1016/j.neures.2022.08.010

Sehara, Y., Hayashi, Y., Ohba, K., Uchibori, R., Urabe, M., Inutsuka, A., Shimazaki,
K., Kawai, K., and Mizukami, H. (2022). Higher Transduction Efficiency of AAV5
to Neural Stem Cells and Immature Neurons in Gerbil Dentate Gyrus Compared to
AAV2 and rh10. *Hum Gene Ther* 33, 76-85. doi:10.1089/hum.2021.106

Shinohara, R., Taniguchi, M., Ehrlich, A.T., Yokogawa, K., Deguchi, Y., Cherasse,
Y., Lazarus, M., Urade, Y., Ogawa, A., Kitaoka, S., Sawa, A., Narumiya, S., and
Furuyashiki, T. (2018). Dopamine D1 receptor subtype mediates acute stress-
induced dendritic growth in excitatory neurons of the medial prefrontal cortex and
contributes to suppression of stress susceptibility in mice. *Mol Psychiatry* 23, 1717-
1730. doi:10.1038/mp.2017.177

Simon, R., Brylka, H., Schwegler, H., Venkataramanappa, S., Andratschke, J.,
Wiegreffe, C., Liu, P., Fuchs, E., Jenkins, N.A., Copeland, N.G., Birchmeier, C.,
and Britsch, S. (2012). A dual function of Bcl11b/Ctip2 in hippocampal
neurogenesis. *EMBO J* 31, 2922-2936. doi:10.1038/emboj.2012.142

Ueno, M., Sugimoto, M., Ohtsubo, K., Sakai, N., Endo, A., Shikano, K., Imoto, Y.,
and Segi-Nishida, E. (2019). The effect of electroconvulsive seizure on survival,
neuronal differentiation, and expression of the maturation marker in the adult mouse
hippocampus. *J Neurochem* 149, 488-498. doi:10.1111/jnc.14691

Wang, H., Warner-Schmidt, J., Varela, S., Enikolopov, G., Greengard, P., and Flajolet,
M. (2015). Norbin ablation results in defective adult hippocampal neurogenesis and
depressive-like behavior in mice. *Proc Natl Acad Sci U S A* 112, 9745-9750.
doi:10.1073/pnas.1510291112

Wang, J.W., David, D.J., Monckton, J.E., Battaglia, F., and Hen, R. (2008). Chronic
fluoxetine stimulates maturation and synaptic plasticity of adult-born hippocampal
granule cells. *J Neurosci* 28, 1374-1384. doi:10.1523/JNEUROSCI.3632-07.2008

Wang, S.E., Ko, S.Y., Kim, Y.S., Jo, S., Lee, S.H., Jung, S.J., and Son, H. (2018).
Capsaicin upregulates HDAC2 via TRPV1 and impairs neuronal maturation in
mice. *Exp Mol Med* 50, e455. doi:10.1038/emm.2017.289

Warner-Schmidt, J.L., and Duman, R.S. (2007). VEGF is an essential mediator of the
neurogenic and behavioral actions of antidepressants. *Proc Natl Acad Sci U S A* 104,
4647-4652. doi:10.1073/pnas.0610282104

Yamasaki, N., Maekawa, M., Kobayashi, K., Kajii, Y., Maeda, J., Soma, M., Takao,
K., Tanda, K., Ohira, K., Toyama, K., Kanzaki, K., Fukunaga, K., Sudo, Y.,
Ichinose, H., Ikeda, M., Iwata, N., Ozaki, N., Suzuki, H., Higuchi, M., Suhara, T.,
Yuasa, S., and Miyakawa, T. (2008). Alpha-CaMKII deficiency causes immature
dentate gyrus, a novel candidate endophenotype of psychiatric disorders. *Mol Brain*
1, 6. doi:10.1186/1756-6606-1-6

You, J.C., Muralidharan, K., Park, J.W., Petrof, I., Pyfer, M.S., Corbett, B.F.,
LaFrancois, J.J., Zheng, Y., Zhang, X., Mohila, C.A., Yoshor, D., Rissman, R.A.,
Nestler, E.J., Scharfman, H.E., and Chin, J. (2017). Epigenetic suppression of
hippocampal calbindin-D28k by DeltaFosB drives seizure-related cognitive
deficits. *Nat Med* 23, 1377-1383. doi:10.1038/nm.4413

Zhao, C., Deng, W., and Gage, F.H. (2008). Mechanisms and functional implications
of adult neurogenesis. *Cell* 132, 645-660. doi:10.1016/j.cell.2008.01.033

Table 1. List of primers used for qPCR analysis.

Gene	Forward (5' to 3')	Reverse (5' to 3')
Dsp	GCTGAAGAACACTCTAGCCCA	ACTGCTGTTTCCTCTGAGACA
Calb1	TCTGGCTTCATTTCGACGCTG	ACAAAGGATTTCATTTCCGGTGA
Tdo2	ATGAGTGGGTGCCCGTTTG	GGCTCTGTTTACACCAGTTTGAG
Ntf3	AGTTTGCCGGAAGACTCTCTC	GGGTGCTCTGGTAATTTTCCTTA
Il1r1	GTGCTACTGGGGCTCATTTGT	GGAGTAAGAGGACACTTGCGAAT
Hdac2	GGAGGAGGCTACACAATCCG	TCTGGAGTGTTCTGGTTTGTCA
Tiam1	AGACTGAACGCACCTATGTGA	ACTCGACTTGAAACTCCACCAT
18S	GAGGCCCTGTAATTGGAATGAG	GCAGCAACTTTAATATACGCTATTGG
Gapdh	AGGTCGGTGTGAACGGATTTG	TGTAGACCATGTAGTTGAGGTCA

Figure 1. Expression of Dsp in the dentate gyrus (DG) of the hippocampus

(A) Representative images of immunoreactivity for Dsp (green) and NeuN (red) in the DG. Magnified images showing Dsp-positive (a; arrows) or -negative (b; arrowheads) cells. DAPI (blue) is utilized for nuclear staining. (B) Relative gene expression of *Dsp* at 24 h after 11 times of electroconvulsive seizure (ECS) stimulation (left) or 4-weeks treatment of selective serotonin reuptake inhibitor (SSRI) fluoxetine. Data are expressed using dot plots and means $\pm$ standard error of mean. *, $P<0.05$; **, $P<0.01$. The n number is given in graph. (C) Representative images of immunoreactivity for Dsp (DSP; green), and DAPI (blue) in the DG after 20 times of ECS stimulation. Scale bars: 20 μm.

Figure 2. Knockdown of Dsp in the DG by adeno-associated virus (AAV) carrying Dsp artificial miRNA

(A) Structure of AAVrh10 carrying EGFP and artificial microRNA sequence for Dsp (DSP miRNA). Time course of experimental procedure (Left). EGFP and immunoreactivity for an immature neuronal marker doublecortin (red) in the DG indicating mature neuronal infection of the AAV after 7–8 weeks of injection. (B) Gene expression of *Dsp* 7–8 weeks after AAV infection. Data are expressed using dot plots and means ± standard error of mean. *, *P*<0.05. The n number is given in graph. (C) Representative images of immunoreactivity for Dsp (DSP; red) and DAPI (blue) in the DG 7–8 weeks after AAV infection. Scale bars: 100 μm.

Figure 3. Expression changes in the DG by Dsp knockdown

(A) Effect of Dsp knockdown in the DG on the expression of FosB protein. Representative images of immunoreactivity for FosB in the DG (left). Scale bars: 100 μm. Quantification of FosB-positive cell numbers (right) in the granule cell layer. (B) Effect of Dsp knockdown in the DG on the expression of maturation markers in the DG. Expression of *Calb1, Tdo2, Ntf3,* and *Il1r1*. (C) Effect of Dsp knockdown in the DG on the expression of *Hdac2* and *Tiam1* which are regulated by β-catenin signaling in the DG. Data are expressed using dot plots and means ± standard error of mean. The n number is given in graph. *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; n.s., not significant.

Figure 4. Electrophysiological changes in the DG by Dsp knockdown

(A) Diagram showing the electrode arrangement for recording field excitatory post synaptic potentials (EPSPs) at the mossy fiber (MF)–CA3 synapse. (B) Ratios for MF field EPSPs to fiber volley (FV) at the baseline stimulus intensity. (C and D) Effect of Dsp knockdown on synaptic facilitation induced by repetitive stimulation at the MF synapse. (E) Effect of Dsp knockdown on potentiation of MF synaptic transmission induced by serotonin (5 μM, 5 min) applied in the bath. Sample recordings show averages of nine consecutive field EPSPs before and at the end of serotonin application in control DG and knockdown DG. (F) Effect of Dsp knockdown on serotonin-induced synaptic potentiation. Data are expressed using dot plots and means ± standard error of mean. The n number of data given in graph represents the number of slices. **, $P<0.01$.

Figure 5. Adult neurogenic changes in the DG by Dsp knockdown
(A) Effect of Dsp knockdown on proliferation in the DG. Left: Time course of the experimental procedure. Middle: Representative images of immunoreactivity for BrdU. The arrows represent BrdU-positive cells in the SGZ. Scale bars: 20 μm. Right: Quantification of BrdU-positive cell numbers in the SGZ. (B) Effect of Dsp knockdown on neuronal differentiation in the DG. Left: Representative images of immunoreactivity for NeuroD1 in the DG. Scale bars: 100 μm. Right: Quantification of NeuroD1-positive cell numbers in the SGZ. (C) Effect of Dsp knockdown on cell number of doublecortin (DCX)-positive immature neurons in the DG. Left: Representative images of immunoreactivity for DCX in the DG. Scale bars: 40 μm. Middle: Quantification of DCX-positive cell numbers in the SGZ. Right: Quantification of DCX-positive cells with long dendrites in the SGZ. (D) Effect of Dsp knockdown on survival of 2-weeks of cell age in the DG. Left: Time course of the experimental procedure. Right: Quantification of BrdU-positive cells of 2-weeks (2 wks) of cell age in the DG. Data are expressed using dot plots and means ± standard error of mean. The n number is given in graph. *, $P<0.05$; **, $P<0.01$. n.s. not significant.

Figure 6. Behavioral changes in the DG by Dsp knockdown

(A) Time course of the experimental procedure. (B) Exploratory behaviors of open field test (OFT). Total distance (left) in the open field. Time spent (middle) and entry times (right) in the center of the open field. (C) Immobility time in the forced swim test (FST). (D) Total distance of locomotor activity (LMA) in the home cage. (E) Social interaction behavior in the 3-chamber social interaction test (SIT). A stranger mouse within a wire cup placed in one chamber (social) and an empty wire cup placed in another chamber (Non-Social). (F) Time spent on the open arms (left) and total distance (right) in the elevate plus maze (EPM) test. Data are expressed using dot plots and means ± standard error of mean. The n number is given in graph. *, $P<0.05$; **, $P<0.01$; n.s. not significant.