Noncanonical Activity of Med4 as a Gatekeeper of Metastasis through Epigenetic Control of Integrin Signaling

Authors:

Seong-Yeon Bae[1-4*], Yi Chen[2,5-6], Hong Chen[3,7], Dhiraj Kumar[1-3], Spyros Karaiskos[4], Jane Xu[2,5-6], Chao Lu[2,4], Aaron D. Viny[2, 4-6*], Filippo G. Giancotti[1-4†]

Affiliations:

[1]Cancer Metastasis Initiative, Herbert Irving Comprehensive Cancer Center, Columbia University Irving Medical Center, New York, New York 10032, USA

[2]Department of Genetics and Development, Vagelos College of Physicians and Surgeons, Columbia University Irving Medical Center, New York, New York 10032, USA

[3]Department of Cancer Biology, The University of Texas MD Anderson Cancer Center, Houston, Texas 77054, USA

[4]Herbert Irving Comprehensive Cancer Center, Columbia University Irving Medical Center, New York, New York 10032, USA

[5]Columbia Stem Cell Initiative, Vagelos College of Physicians and Surgeons, Columbia University Irving Medical Center, New York, New York 10032, USA

[6]Division of Hematology and Oncology, Department of Medicine, Columbia University Irving Medical Center, New York, New York 10032, USA

[7]Department of Thoracic/Head and Neck Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, Texas 77054, USA

[†]Deceased

*Correspondence: sb4595@cumc.columbia.edu (S.B.), adv2105@cumc.columbia.edu (A.D.V.)

Authors' Disclosures

A.D. Viny is in the scientific advisory board for Arima Genomics. All other authors declare no competing interests.

Abstract

Breast cancer metastatic relapse after a latency period, known as metastatic dormancy. Through
genetic screening in mice, we identified the mediator complex subunit 4 (*Med4*) as a novel
tumor-cell intrinsic gatekeeper in metastatic reactivation. *Med4* downregulation effectively
awakened dormant breast cancer cells, prompting macroscopic metastatic outgrowth in the lungs.
Med4 depletion results in profound changes in nuclear size and three-dimensional chromatin
architecture from compacted to relaxed states in contrast to the canonical function of the
Mediator complex. These changes rewire the expression of extracellular matrix proteins,
integrins, and signaling components resulting in integrin-mediated mechano-transduction and
activation of YAP and MRTF. The assembly of stress fibers pulls on the nuclear membrane and
contributes to reinforcing the overall chromatin modifications by *Med4* depletion. MED4 gene
deletions were observed in patients with metastatic breast cancer, and reduced *MED4* expression
correlates with worse prognosis, highlighting its significance as a potential biomarker for
recurrence.

Significance: This work establishes the first link between Med4 and its role as an epigenetic and
transcriptional repressor of ECM programs that reactivates dormant tumor cells in breast cancer
metastasis. We highlight the distinct noncanonical role of Med4, in contrast to Med1, as well as
the functional implications *in vitro* and *in vivo*. We have functionally validated the aberrant
signaling pathways activated through both genetic add back and chemical inhibition and
correlate our findings with Med4 expression, metastatic potential, and overall outcome in
retrospective patient cohorts.

INTRODUCTION

Survivorship for patients with breast cancer is predicated not by the disease burden in the breast
itself, but rather the capability of cells to metastasize outside the breast. Over 90% of patients
with breast cancer succumb as a result of distant metastasis. Clinical experience and
experimental data in both patients and mouse models show that metastatic disease is the
overwhelming cause of breast cancer deaths. Cellular and molecular studies of metastatic breast
cancer suggest that disseminating cells from the primary tumor undergo a variable period of
dormancy at pre-metastatic sites before undergoing reactivation and outgrowth in macroscopic
and life-threatening lesions (1). These data point to a distinct stage of tumor progression whereby
organ tropism and quiescence may hallmark both a biomarker of high-risk disease as well as a
key therapeutic target. Albeit a small window of opportunity, particularly when the period of
clinical latency is short (e.g., TNBC) (2,3) this cellular transformation may represent the
keystone to prevention of distant metastasis.

Genomic studies have suggested that macroevolutionary shifts, such as chromosomal instability,
contribute to the emergence of cells with metastatic potential from within primary tumors (4).
However, their subsequent dissemination and final outgrowth in target organs may be driven by
epigenetic mechanisms regulating key features of cellular identity as in epithelial-to-
mesenchymal transition (EMT) and stemness/self-renewal (5). Despite their clinical importance,
metastatic dormancy and reactivation are poorly understood due to the complexity of the
underlying biology. To deconvolve this complexity, we developed a forward genetic screening
strategy in the mouse to unveil key regulators of metastatic reactivation (6). Screening of open
reading frame (ORF) libraries enabled us to identify several positive regulators of metastatic

reactivation of breast cancer, including Coco, which binds to and inhibits BMP in the lung
microenvironment, and the tetraspanin TM4SF1, which mediates non-canonical activation of the
JAK2-STAT3 signaling pathway (7,8). In addition, we have found that the master epigenetic
regulator PRC1 coordinates stemness and immune suppression at bone metastatic sites in
prostate cancer (9).

Collectively, our prior studies suggest that metastatic outgrowth is initiated by a subpopulation of
cancer stem cells with metastatic potential (metastasis-initiating cells), which undergo dormancy
and reactivation in response to then activation of signaling pathways and transcriptional
networks similar to those that govern these processes in adult stem cells (2). However, numerous
modulators involved in regulating metastatic dormancy remain to be identified due to the
extensive heterogeneity observed in the molecular, genetic, and clinical characteristics of breast
cancer metastases. To unveil additional modulators associated with metastatic dormancy, we
conducted an *in vivo* CRISPR-Cas9 screen aimed at identifying genetic factors that contribute to
breast cancer reactivation in dormant tumor cells. Here we present *mediator complex subunit 4*
(*Med4*) as a novel tumor-cell intrinsic gatekeeper in metastatic reactivation.

The Mediator complex regulates Pol II-mediated transcription by adjoining enhancer-bound
transcription factors and the general transcription machinery in *cis* (10,11). Mediator, comprising
a 26-subunit core organized into head, middle, and tail modules, and a four-subunit dissociable
kinase module, manifests functional adaptability due to the variable arrangement of its subunits
(12,13). The subunit-specific composition and function of Mediator in its interaction with
transcription factors (TF) imply that individual subunits of the complex regulate distinct
transcriptional responses (14,15). The molecular basis for the disparate functional roles of
individual subunits in various biological contexts remains incompletely understood (16-18),
particularly as it pertains to *Med4*. We demonstrated that genetic perturbation of *Med4* awakens
dormant tumor cells in pre-metastatic niches. The biological relevance of our findings is
supported by the observation that low *MED4* expression is associated with increased risk of
metastatic relapse in patients with breast cancer. Taken together, we propose an innovative
epigenetic regulatory mechanism whereby *Med4* loss rewires integrin signaling through
decompaction of 3-dimensional chromatin architecture.

RESULTS

Med4 **enforces dormancy.**

To determine novel entities enforcing dormancy, we performed a genome-wide loss-of-function
screen using the GeCKO v2 sgRNA library (6,19) (Fig. 1A). Following tail vein injection and
lung infiltration, 4T07 cells that had acquired the ability to exit dormancy underwent reactivation
and generated macrometastases (Supplementary Fig. S1A). After isolating tumor cells from each
metastasis, we cloned and sequenced the integrated sgRNA inserts. Intriguingly, we identified
sgRNAs targeting 3 genes (*Med4*, *Nipbl; Nipped-B-like protein*, and *Esco1; Establishment of*
sister chromatid cohesion N-acetyltransferase 1) involved in the formation of super-enhancers
(Fig. 1B). Whereas MED4 is a core subunit of the Mediator complex, which connects super-
enhancers and promoters in *cis*, NIPBL and ESCO1 aid in loading cohesin onto DNA and hence
physically stabilize the assembly of the super-enhancer-promoter complex. We decided to focus
on *Med4* because of its inferred biological function, its biological potency *in vivo*, and its
potential relevance to human breast cancer. To confirm that *Med4* gene was genuinely edited to

promote metastasis, we performed indel PCR analysis and western blotting in the clone
originally isolated from the screen. The results indicated that the clone which contains a sgRNA
targeting *Med4*, carried heterozygous instead of homozygous deletion (one allele of the clone
carries a nonsense mutation) (Supplementary Fig. S1B-S1E). We hypothesized that *Med4* is an
essential gene and examination of the DepMap database supports this hypothesis (Supplementary
Fig. S1F) (20). In agreement with the essentiality of *Med4*, we were not able to generate *Med4*
KO clones presumably because of strong counterselection. To validate the biological potency of
Med4 in vivo, we used MirE shRNA targeting *Med4* in 4T07-TGL cells (Supplementary Fig.
S1G-S1I) and another syngeneic dormant system, D2A1-d-TGL cells (Fig. 1C-1F). Upon *Med4*
silencing, the tumor cells acquired the capacity to efficiently colonize the lung when
intravenously injected into immunocompetent mice. To determine if *Med4* silencing specifically
induces reactivation from dormancy at the pre-metastatic niche, we used a doxycycline-inducible
promoter to express shRNAs targeting *Med4*. Mice were treated with doxycycline either
immediately or 2 weeks after tail vein injection (Fig. 1G). Whereas immediate onset of *Med4*
knockdown induced metastatic outgrowth at around 2 weeks, a 2-week delay in shRNA
expression caused a similar delay in metastatic outgrowth (Fig. 1H-1J), confirming that *Med4*
downregulation re-awakens mammary tumor cells from dormancy. Taken together, these
findings suggest that *Med4* serves as a sentinel of dormant tumor cells from reactivation.

Lower levels of *MED4* are associated with metastatic relapse in patients.

To examine the potential relevance of our findings to human breast cancer, we first examined the
expression of MED4 in multiple metastatic breast cancer datasets from cBioportal (21,22).
Analysis of patient datasets indicated that *MED4* has shallow deletions in almost 40% of human
breast cancer metastasis samples (Fig. 1K, Supplementary Fig. S2A and S2B) (23). The
proportion of breast tumors with deep deletions of *MED4* is extremely low, presumably because
MED4 is an essential gene as aforementioned. Consistently with genomic analysis, paired RNA-
seq revealed that mRNA levels of *MED4* are lower but not undetectable in lung, liver, and brain
metastases than in matched primary tumors (Fig. 1L) (24). Intriguingly, Kaplan-Meier analysis
revealed that *MED4* expression levels could be used to stratify patients into high- and low-risk
groups in terms of distant metastasis-free survival; lower MED4 levels were strongly correlated
with poor prognosis in patients with advanced breast cancer across subtypes (Fig. 1M, and
Supplementary Fig. S2C-S2E) (25). Furthermore, lower levels of *MED4* mRNA could serve as a
prognostic indicator for poor overall survival, particularly in subgroups of patients with estrogen
receptor-negative (ER-) or human epidermal growth factor 2-positive (HER2+) breast cancer
(Supplementary Fig. S2F and S2G). Collectively, these results suggest that MED4's expression
level could be a prognostic marker to predict metastatic relapse in breast cancer.

***Med4* loss promotes cancer stem cell traits.**

Phenotypic and functional studies indicated that the *Med4*-silenced cells proliferate modestly
slower as compared to control cells under standard culture conditions (Supplementary Fig. S3A
and S4A). However, we considered the potential role of *Med4* in regulating the expression of
cancer stem cell traits given that the metastasis-initiating cells exhibit functional similarities to
cancer stem cells (26). Intriguingly, *Med4*-silenced cells exhibited a substantial increase in their
self-renewal capacities, demonstrated by the increased number and size of tumor spheres
(Supplementary Fig. S3B and S4B), enhanced clonogenicity (Supplementary Fig. S3C and S4C),

and improved anchorage-independent growth (Supplementary Fig. S3D and S4D). These
findings suggest that the silencing of *Med4* induces metastatic outgrowth by augmenting cancer
stem cell traits, as observed previously for overexpression of Coco or TM4SF1 (7,8). In addition,
Med4-deficient cells exhibited increased migratory (Supplementary Fig. S3E and S4E) and
invasive (Supplementary Fig. S3F) capacities which are common characteristics of cells
undergoing epithelial to mesenchymal transition (EMT) (27). Overall, these results are consistent
with the role of *Med4* as a dormancy enforcer and suggest that *Med4* silencing endows dormant
tumor cells with self-renewal capacities to produce metastatic outgrowths.

Med4 downregulation reprograms the transcriptional and epigenetic landscape leading to
the reinforcement of desmoplastic niche formation, in contrast to the effects of _Med1_.

To obtain a comprehensive view of the transcriptional reprograming in dormancy and metastatic
reactivation, we performed bulk RNA sequencing (RNA-seq) of D2A1 (metastatic) cells and its
dormant variant D2A1-d cells (28). We identified D2A1_metastatic gene signature (225
differentially upregulated genes in D2A1) and D2A1_dormant gene signature (207 differentially
upregulated genes in D2A1-d), respectively (Supplementary Fig. S5A). We employed both
signatures in analyzing bulk RNA-seq data from D2A1-d cells expressing two different shRNAs
targeting *Med4*, in comparison to a control shRNA. Gene set enrichment analysis (GSEA)
indicates that the D2A1_metastatic signature was enriched in *Med4*-silenced cells while the
D2A1_dormant signature was enriched in control cells (Supplementary Fig. S5B). This result
indicates that *Med4* depletion reprograms transcriptional profile from a dormant to a metastatic
state. In addition, RNA-seq revealed that *Med4* silencing skews the transcriptional profile to an
active state, antithetical to the canonical function of the Mediator in transcription (14) (Fig. 2A).
To determine if the changes in the transcriptional landscape resulting from *Med4* silencing are a
general feature of the Mediator complex or specific to *Med4*, we employed MirE shRNA to
target the mediator complex subunit 1 (*Med1*) (Supplementary Fig. S5C). MED1 has been the
most extensively investigated among the Mediator subunits and is recognized as a prominent
transcriptional coactivator and a constituent of super-enhancers (SEs) (10,29). Intriguingly, the
silencing of *Med1* resulted in a skewed transcriptional profile towards repression which contrasts
with the effect observed with *Med4* silencing (Supplementary Fig. S5D). GSEA using metastatic
and dormant signatures also indicated that *Med1* reprograms the transcriptional profile in the
opposite direction compared to *Med4* (Supplementary Fig. S5E). Globally the knockdown of
either *Med4* or *Med1* did not affect the expression of other mediator complex subunits,
suggesting that the observed effects are related to subunit-specific function rather than overall
Mediator stoichiometry (Supplementary Fig. S5F-S5H). Thus, we sought to investigate the
impact of *Med4* depletion on the three-dimensional (3D) chromatin structure. To map chromatin
accessibility comprehensively, we first performed micrococcal nuclease (MNase) digestion assay
(30). A titration of MNase revealed that a susceptibility for MNase was elevated in *Med4*-
silenced cells especially at the poly-nucleosome part (Fig. 2B). This result suggested that *Med4*
silencing decompacts heterochromatic structure in which DNA is tightly packed with
nucleosomes. On the contrary, *Med1* silencing resulted in heightened resistance to MNase,
suggesting that the downregulation of *Med1* caused chromatin structure to become more compact
(Supplementary Fig. S5I). We also confirmed changes in the genome-wide chromatin
accessibility landscape by *Med4* silencing using the assay for transposase-accessible chromatin
with sequencing (ATAC-seq). The results clearly demonstrated that *Med4* depletion primarily
leads to a substantial increase in chromatin accessibility (Fig. 2C). Conversely, *Med1* silencing

had minimal impact on chromatin accessibility (Supplementary Fig. S5J). To delve deeper into the consequences of *Med4* silencing on chromatin structure and transcriptional output, we integrated RNA-seq and ATAC-seq results. Upon depleting *Med4*, the majority of genes exhibited significant changes in chromatin accessibility and gene expression, with increased levels in both accessibility and transcription (Fig. 2D). GSEA revealed that the downregulation of *Med4* induces the expression of signatures reflective of core extracellular matrix proteins (ECMs), assembly of the ECMs, integrin signaling, muscle contraction processes and epithelial to mesenchymal transformation (EMT) (Supplementary Fig. S5K and S5L). Consistent with the results from GSEA, gene ontology (GO) analyses of genes showing concomitant changes in accessibility and gene expression (located in the upper quadrant of the intersection between RNA-seq and ATAC-seq) also indicated significant enrichment of ECM-related and cell-adhesion-related gene signatures (Fig. 2E). In particular, the majority of differentially expressed genes (DEGs) after *Med4* silencing was collagen- (e.g., *Col1a1, Col1a2, Col6a1*), actomyosin-related (e.g., *Acta2, Myl9*), and ECM-remodeling genes (e.g., *Loxl3, Mmp2, Adamts5*) (Fig. 2F). This suggests that *Med4* depletion is associated with stiffening matrices surrounding cancer cells, so called desmoplasia, which is a critical manifestation of cancer progression (31-33). Consistently, immunohistochemistry (IHC) of α-SMA (a protein coded by *Acta2*) and Masson's trichrome stain, which labels collagen, validated the substantial desmoplastic stromal reaction in metastases induced by *Med4* silencing compared to the dormant niche (Fig. 2G). Collectively, these results indicate that *Med4* deficiency bolsters matrix stiffening, which creates a favorable environment conducive to the metastatic outgrowth.

Med4 silencing alters nuclear architecture and decompacts 3D chromatin organization by increasing cytoskeletal tensile forces.

Although a growing body of evidence indicates that Mediator exerts on enhancer-promoter looping functionally rather than structurally, Mediator has been implicated in super-enhancer formation and promoter-enhancer looping orchestrating with the cohesin ring (11,34). It is thereby conceivable that Mediator plays a role in both maintaining and modifying chromatin architecture. Thus, we sought to visualize the chromatin structure by staining HP1α, a component of constitutive heterochromatin (35). Intriguingly, *Med4*-deficient cells displayed a sparse distribution of heterochromatin foci accompanied by an increase in nuclear size, whereas the control cells exhibited a condensed heterochromatin structure (Fig. 3A, and Supplementary Movies 1-3). In contrast, *Med1*-deficient cells did not exhibit significant changes in the distribution of heterochromatin foci (Supplementary Fig. S6A, and Supplementary Movies 4-6). These findings indicate that *Med4* plays a role in controlling chromatin and nuclear architectures. Given that the nuclear lamina is known to anchor the heterochromatin structure to the nuclear periphery and be associated with gene silencing, we hypothesized that the nuclear lamina plays a crucial role in the alterations of heterochromatin structure caused by *Med4* silencing (36,37). To investigate this, we co-stained the cells with LAMIN A/C and H3K9me3, another marker for heterochromatin. Interestingly, we observed that H3K9me3 was dislodged from the nuclear lamina in *Med4*-depleted cells, whereas it remained colocalized in control cells (Fig. 3B). Overall, these findings indicate that the depletion of *Med4* leads to the decompaction of the 3-dimensional chromatin structure. Furthermore, in *Med4*-depleted cells, we observed a reduction in the densities of LAMIN A/C at the nuclear periphery, accompanied by an enlargement in nuclear size (Fig. 3C and 3D). Notably, the overexpression of *MED4* had the ability to reverse nuclear enlargement and restore the distorted nuclear lamina. However, such changes were not

observed in *Med1*-deficient cells (Supplementary Fig. S6B). These results led us to hypothesize
that *Med4* could potentially alter the solubility of LAMIN A/C. To investigate this, we detected
p-LAMIN A/C (Ser^{22}) (38,39) and total LAMIN A/C, followed by sub-fractionating the
nucleoplasm and nuclear envelope (Supplementary Fig. S6C). However, our results did not
reveal any changes in LAMIN solubility caused by *Med4* silencing. Since the stress fibers are
connected to the matrix via integrins and to the nuclear membrane through the LINC complex
(40), heightened tension in the actin cytoskeleton could conceivably increase the size of the
nucleus by exerting radial forces on the nuclear membrane. To investigate the impact of
cytoskeletal tensile forces on the alterations in nuclear size and architecture resulting from *Med4*
depletion, we used actin polymerization inhibitors, cytochalasin D, or latrunculin B (41).
Remarkably, upon treating the cells with these actin cytoskeleton disruptors, we observed a
reversal of the changes in nuclear size and architecture caused by *Med4* silencing (Fig. 3E).
Furthermore, disrupting the actin cytoskeleton resulted in the reversal of the dispersed
heterochromatin structure observed in *Med4*-deficient cells (Fig. 3F and Supplementary Fig.
S6D). These results suggest the reduction of MED4 leads to elevated expression of ECM
proteins and the assembly of stress fibers. These stress fibers apply tension to the nuclear
membrane, potentially facilitating or reinforcing alterations in chromatin structure.

***Med4* deficiency promotes cell growth within a desmoplastic niche by activating integrin**
signaling.

Considering the increasing body of evidence supporting the impact of biochemical and
biomechanical cues derived from the extracellular matrix (ECM) on cancer cell metastatic
abilities, we sought to examine if *Med4* mediates outgrowth in response to the stiffness and
composition of matrices. To model the normal ductal microenvironment characterized by a
basement membrane-rich ECM, and the stiff tumor microenvironment enriched in collagen I
(32,42), we placed control and *Med4*-silenced breast cancer cells in two different conditions: (1)
3D Matrigel, representing the basement membrane-rich ECM, and (2) 3D Matrigel plus collagen
I, which recapitulates the tumor microenvironment. Strikingly, *Med4* silencing promoted tumor
organoid growth and invasion in stiff 3D matrices (Matrigel + collagen I), while cells maintained
a spherical growth pattern in the soft 3D matrices (Matrigel-only) regardless of their *Med4*
expression (Fig. 4A). Ki67 staining indicated that *Med4* inactivation increases cell cycle
progression in stiff 3D matrices but slightly decreases it in 2D culture (Fig. 4B). These data
indicate that cells in a *Med4*-deficient background can preferentially grow in the desmoplastic
niche. Collectively, these results indicate that *Med4* deficiency not only induces matrix stiffening,
fostering a favorable environment for metastatic outgrowth, but also stimulates the tumor-cell
intrinsic survival and proliferative abilities within the established metastatic niche.

To exit dormancy, metastatic tumor cells are required to extend lamellipodia and establish
productive interactions with the matrix. Adhesion molecules, particularly integrins, play a crucial
role in transmitting external stimuli from the ECM (43). These integrins are integrated into focal
adhesion (FA) complexes, which connect the cytoskeleton to the ECM, enabling the reciprocal
transmission of forces between cells and the surrounding microenvironment (44). Given the
pivotal role of integrins in cell adhesion and signaling, along with the transcriptional alterations
in the ECM composition (Fig. 2E) and integrin repertoire (Supplementary Fig. S7A) by *Med4*
silencing, we hypothesized that *Med4* deficiency promotes metastatic reactivation by activating
integrin-mediated signaling. Immunofluorescence analysis revealed that *Med4*-silenced cells
exhibit a more organized actin cytoskeleton and more numerous focal adhesions containing

activated FAK (p-FAK Y^{397}), paxillin, and vinculin (Fig. 4C). Also, p130^{Cas}, JNK, p38, AKT,
and MLC2 phosphorylation were significantly increased upon *Med4* depletion (Fig. 4D).
Furthermore, we validated that *Med4*-deficient cells exhibited increased formation of aligned
stress fibers and focal adhesions in a Matrigel + collagen I environment, indicating that the
downregulation of *Med4* enhanced integrin-mediated adhesion in the 3D culture setting (Fig. 4E).
In order to corroborate the implication of integrin signaling activation in the growth of organoids
due to *Med4* silencing, we employed defactinib, a targeted inhibitor of FAK (45). Remarkably,
the application of defactinib demonstrated a marked reduction in the invasive proliferation of
Med4-deficient cells within the 3D Matrigel + collagen I matrix (Fig. 4F). Collectively, these
findings provide evidence indicating that *Med4* silencing triggers the activation of integrin
signaling.

Med4 loss promotes metastatic reactivation by activating integrin-mediated nuclear
mechanosensing.

To identify the top transcription factors controlling the transcriptome changes by *Med4* silencing,
we applied ChIP-X Enrichment Analysis 3 (ChEA3) transcription factor enrichment analysis to
the RNA-seq data (46). Notably, we found that SRF, and TEAD 1/2 cistromes govern the
transcriptional reprogram by *Med4* depletion along with EMT cistromes (Fig. 5A). Also, *de novo*
motif enrichment analysis performed on genes exhibiting concurrent increases in accessibility
and gene expression following *Med4* silencing confirmed the enrichment of TEAD, TEAD4, and
SRF motifs (Fig. 5B). SRF, a co-factor for myocardin-related transcription factor (MRTF), and
TEADs, transcriptional co-activators for YAP and TAZ, are major mechanosensitive
transcriptional factors that transduce mechanical cues by the cytoskeleton to gene expression in
response to integrin-mediated adhesion. GSEA results revealed that the gene signatures for
MRTF/SRF (47) and for YAP (48) are enriched in *Med4*-deficient cells (Fig. 5C). In response to
both biochemical and mechanical signals, YAP undergoes translocation from the cytoplasm to
the nucleus, where it interacts with and triggers the activation of the TEAD transcription factor
(49). MRTF binds stoichiometrically to soluble G-actin and is released as a consequence of actin
polymerization and formation of stress fibers. Once released from G-actin, MRTF enters into the
nucleus, where it functions as a co-factor for SRF, which drives expression of cytoskeletal genes
in a feed forward loop (50). Hence, we sought to examine the cellular localization of these
transcription factors. Immunofluorescence results revealed that *Med4* silencing led to the
predominant nuclear localization of both MRTF-A and YAP (Fig. 5D). Consistent with these
results, expression of MRTF/SRF target genes(51,52), YAP/TEAD target genes (53), and shared
MRTF and YAP target genes were activated in *Med4*-deficient cells, confirming these
mechanosensitive transcription factors were activated by Med4 depletion (Supplementary Fig.
S7B). Restoring MED4 expression by overexpressing MED4 cDNA led to the relocation of
MRTF-A to the cytoplasm and resulted in the reversal of the activation of MRTF-A/SRF target
gene expression (Fig. 5E, 5F and Supplementary Fig. S7C). Immunohistochemistry of lung
sections unveiled that a notable portion of macrometastases originating from *Med4*-deficient
cells exhibited positive staining for nuclear MRTF-A and YAP, in contrast to micrometastases
originating from control cells which exhibited negative nuclear staining of these factors
(Supplementary Fig. S7D). To determine if these integrin-mediated mechanosensitive
transcription factors play a pivotal role for *in vivo* metastatic reactivation by *Med4* loss, we
ablated these mechanosensitive transcription factors though shRNA silencing and injected into
immunocompetent BALB/c recipients (Supplementary Fig. S7E). Silencing *Mrtfa*, *Srf,* or *Yap* +

Taz abrogated the metastatic capacity of *Med4*-silenced cells, pointing to a requirement for these
transcriptional regulators in the process of metastatic reactivation (Fig. 5G). These findings
indicate that the depletion of MED4 triggers metastatic reactivation through the initiation of
integrin-driven mechano-transduction, consequently leading to the activation of MRTF-SRF and
YAP/TAZ signaling pathways.

DISCUSSION

The Mediator complex is implicated in multiple molecular functions encompassing epigenetic
regulation, transcriptional elongation, termination, mRNA processing, noncoding RNA
activation and super-enhancer formation(54); yet the exact mechanisms through which this
plastic and multimeric complex coordinates multiple transcription factors remain unclear. In
addition, whereas MED1 is likely involved in the formation of super-enhancers and
transcriptional activation, which drive the expression of key cell-identity genes (10,29) as well as
oncogenic transcription factors (55-57), both the physiological and pathophysiological roles of
MED4 is poorly understood. Interestingly, in public datasets from large scale transcriptional
profiling efforts, down-regulation of *MED4* has been correlated with cervical cancer and
retinoblastoma (58-60) . This study represents the first description of MED4 as a non-canonical
Mediator component and transcriptional repressor and the first description of MED4 being
implicated in metastasis, highlighting its potential role as a gatekeeper for metastatic relapse.
Intriguingly, the epigenetic and transcriptional reprogramming effects observed by *Med4*
silencing were in stark contrast to the canonical function of the Mediator complex and
specifically of *Med1*. This component-specific disparity identifies a unique and directionally
oppositional function that MED4 serves within the context of Mediator-mediated transcription.
These findings highlight tissue-specific and gene-specific epigenetic mechanisms of control,
hereby in this setting, resulting in aberrant ECM remodeling, integrin-mediated mechano-
transduction, and facilitating the emergence of tumor cells from dormancy within metastatic
niches. Our results demonstrate that MED4 inactivation contributes to the stiffening of the
extracellular matrix (ECM) and the activation of integrin-mediated mechano-transduction. This
is in concert with the euchromatinization and activation of MRTF/SRF and YAP/TAZ
transcription factor programs. Taken together, we posit that *Med4* loss results in decompaction of
the three-dimensional chromatin structure, hereby unlocking ECM gene expression programs.
We show that this epigenetic and transcriptional activation facilitates a functional and permissive
cellular tropism both *in vitro* and *in vivo*.

We demonstrate that *Med4* deficiency has the potential to initiate metastatic reactivation within
the context of breast cancer. Interestingly, our analysis reveals that *Med4* silencing
predominantly stimulates a transcriptional profile consisting of ECM-related genes, which stands
in contrast to the Mediator's conventional role in transcription. Conversely, the silencing of *Med1*
displayed a distinct pattern of transcriptome regulation, highlighting the considerable flexibility
of the complex contingent on its subunit composition. Sun and colleagues conducted a shRNA
screening targeting individual subunits of transcription factor IID (TFIID) and the Mediator
complex (34). Disrupting various subunits yielded distinct effects on the transcriptome of mouse
embryonic stem cells (mESCs), even within subunits of the same complex. Their shRNA-based
screen, similar to our findings, revealed that *Med4* shRNA altered transcriptome towards
activation, and gene ontology analysis of the up-regulated gene list highlighted the enrichment of
ECM signatures. However, when they employed the auxin degron system for the acute

degradation of MED4, it minimally affected the integrity of the Mediator complex and resulted
in transcriptome downregulation by decreasing RNA polymerase II recruitment to the promoters.
Hence, it is plausible that a dosage effect whereby *Med4* knockdown as opposed to complete
knockout drives the pathological phenotype.

The spatial arrangement of chromosomes within the nucleus represents an important layer of
transcriptional regulation. Recent research has shown that heterochromatin is connected to the
nuclear lamina, and mechanical forces are conveyed to interactions between the lamina and
chromatin(61,62). The nucleus has been proposed as a mechanosensor that detects biochemical
and mechanical signals from cellular adhesions through the actin cytoskeleton (63). This
interaction directly leads to the stretching of chromatin, resulting in enhanced transcription (64).
Our data revealed that *Med4*-depleted cells exhibited larger nucleus and decondensed chromatin
structure (Fig. 3A). Given that stress fibers link the matrix through integrins and the nuclear
membrane via the linker of nucleoskeleton and cytoskeleton (LINC) complex (40), increased
tension within the actin cytoskeleton could potentially enlarge the nucleus by applying radial
forces to the nuclear membrane. These enlarged nuclei could potentially lead to the relaxation of
the three-dimensional chromatin structure and increase expression of genes encoding ECM and
cytoskeletal components. The mechanism by which the force-induced activation is selectively
applied to particular genes remains elusive. One possibility is transcriptional feedback of
MRTF/SRF on mechanoadaptation (52,65,66). Upon cytosolic F-actin polymerization, the
MRTF/SRF system is activated and stimulates the expression of genes associated with
extracellular matrix (ECM), adhesion, and the cytoskeleton. Thus, it is conceivable that this
mechanoadaptation through mechanosensitive transcription factors contributes to the
reinforcement of the phenotype resulting from Med4 silencing. In order to articulate this feed-
forward loop between cytoskeletal tensile forces and ECM gene expression by *Med4* silencing, a
comprehensive analysis of the transcriptome and epigenetics after disrupting the LINC complex
by introducing a dominant-negative KASH (DN-KASH) could provide further insights.

We noted that cells with reduced *Med4* levels exhibit decreased fitness when cultured in a 2D
environment (Supplementary Fig. S3A and S4A). However, under conditions involving 3D
Matrigel + collagen I or tumorsphere growth, the *Med4*-silenced cells displayed notable
enhancement in growth and the formation of spheres (Fig. 4A and 4B). In addition, *Med4*-
deficient cells showed higher proliferative capacity and ECM and collagen deposition levels in
the lung parenchyma (Fig. 2G). These findings suggest that a *Med4*-deficient tumor cell not only
leads to create a desmoplastic niche that supports the growth of metastatic cells but also
stimulates the intrinsic survival and proliferative capabilities of tumor cells within the established
metastatic microenvironment. However, the precise mechanism behind this heightened
adaptability in a 3D culture context has yet to be elucidated. A recent study about *MED12 Gly-
44* mutations in the context of leiomyoma revealed findings in line with our results (67). While
MED12 Gly-44 mutant cells displayed a proliferation deficiency under 2D culture conditions,
these mutants formed notably larger spheroids in 3D spheroid environments compared to wild-
type cells. The *MED12* mutations led to changes in the transcriptome, resulting in the expression
of collagen and ECM genes, and these transcriptomic alterations were further magnified in 3D
conditions. Thus, it is plausible that the Mediator might play a pivotal role in modulating cellular
adaptability within the tumor microenvironment through ECM remodeling though the precise
molecular pathways triggered by *Med4* deficiency, which contribute to this increased
adaptability, are yet to be precisely identified. Among potential mechanisms, one notable avenue

is that *Med4* potentially alters the integrin repertoire, leading to an augmentation in the
expression of collagen receptors such as integrin α2β1 (Supplementary Fig. S7A). Further
insights could be gained through additional transcriptome and epigenetic analyses conducted
under varying culture conditions.

Current adjuvant breast cancer therapeutic regimens are aimed to prevention of metastatic
relapse, and despite the prognostic value of pathological, genomic, and transcriptional testing,
late metastasis occurs in nearly 25% of patients (68). Elucidating the molecular mechanisms that
enable dormant tumor cells to evade the immune system and survive under stress at pre-
metastatic sites, as well as identifying subsets of patients who could benefit from therapy
targeting these mechanisms, would greatly reduce the mortality related to metastatic relapse. We
found that a large fraction of metastatic breast cancer samples from patients have shallow *MED4*
deletions (Fig. 1K, Supplementary Fig. S2A and S2B), and lower *MED4* levels correlated with
poor prognosis in patients with advanced breast cancer across subtypes (Fig. 1M and
Supplementary Fig. S2C), indicating that MED4 likely has a substantial role in governing
metastasis in breast cancer patients.
442
Our findings highlight the potential and mechanistic basis for *MED4* expression and related gene
signatures as a biomarker of metastatic dormancy. Future studies will be aimed to determine if
MED4 expression may be able to stratify patients prone to recurrence, enabling the identification
of subsets at high risk for subclinical metastasis and may uniquely benefit from therapeutic
intensification. Moreover, these data lay the foundational groundwork for novel strategies
targeting effectors activated by MED4 loss including early phase inhibitors of the FAK or
YAP/TEAD pathways.
450

METHODS

Animal studies

Mouse studies were conducted in accordance with protocols approved by the Institutional Animal Care and Use Committee of Columbia University and MD Anderson Cancer Center. Six-week-old female mice BALB/c were purchased from The Jackson Laboratory.

To examine lung colonization, 4T07 (3.0×10^5), D2A1-d (2.0×10^5), and D2A1 (2.0×10^5) mammary carcinoma cells were resuspended in 100 µl of PBS and injected into the lateral tail vein.

Doxycycline (Dox) was administered to mice through water *ad libitum* (2 mg/ml). For inducible shRNA experiments, Dox was administered either immediately after injection (Dox day 0) of the cancer cells or 14 days later (Dox day 14). Bioluminescence (BLI) was used to examine metastatic outgrowth. Metastatic lesions were confirmed by postmortem dissection.

For bioluminescent imaging, mice were anesthetized and injected retro-orbitally with 1.5 mg (75 mg/kg) of D-luciferin (Gold Biotechnology; LUCK-1G) at the indicated times after injecting cells. Animals were imaged in an IVIS Spectrum chamber within 5 min after D-luciferin injection, and data were recorded using Living Image software (PerkinElmer). To measure lung colonization, photon flux was calculated for each mouse by using a circular region of interest encompassing the thorax of the mouse. After subtracting a background value obtained from a control mouse injected only with D-luciferin, photon flux was normalized to the value obtained 30 min after injection of the tumor cells. This latter value was arbitrarily set at 100.

Cell lines

The 4T07 cell line was generously provided by Dr. Fred Miller (Wayne State University, Detroit, MI) and cultured as described (69). The D2A1-d and D2A1 cell lines were generously provided by Dr. Robert Weinberg (MIT, Boston, MA) and cultured as described(28). HEK293FT cells (from Thermo Fisher Scientific) was cultured per the manufacturer's protocol. For bioluminescent tracking, cell lines were infected with a lentiviral vector encoding herpes simplex 3 virus thymidine kinase 1, green fluorescent protein (GFP) and firefly luciferase. GFP-positive cells were isolated by FACS.

Genome-wide CRISPR-Cas9 screening.

A forward genetic screening was performed as described in with a minor modification (6). The GeCKO v2 library A was obtained from Addgene (1000000052) and was amplified and packaged in lentiviruses as described previously(19). The viral titer was determined by limiting dilution-colony counting. 4T07 cells were transduced with the library at the MOI of 0.3 to ensure one sgRNA/cell. Library-transduced cells were selected with 2 µg/ml of puromycin for 7 days, and then injected through tail vein of 6-week-old female BALB/c mice. Mice were monitored weekly by bioluminescent imaging and lung metastases were confirmed by postmortem dissection at 3-4 weeks after injecting library-transduced cells. Macroscopic lung nodules were dissected separately and digested in Collagenase Type IV (Stem Cell technologies; 07909) at 37°C for 1.5 h with rocking. After centrifuging at 1,000 rpm for 3 min, pellets were resuspended in the complete medium and filtered through 35 µm strainers. Isolated clonogenic tumor cells were expanded in 4 µg/ml of puromycin medium. Genomic DNAs were isolated using Maxwell®

RSC Cultured Cells DNA Kit (Promega; AS1620) according to the manufacturer's protocol.
Inserted sgRNA sequences were amplified by PCR with the following primers and
thermocycling   parameters   (v2Adaptor-F: 5'-
AATGGACTATCATATGCTTACCGTAACTTGAAAGTATTTCG-3',   v2Adaptor-R: 5'-
TCTACTATTCTTTCCCCTGCACTGTTGTGGGCGATGTGCGCTCTG-3'; 95°C for 2 min,
30 cycles of (98°C for 1 s, 60°C for 5 s, 72°C for 35 s), and 72°C for 1 min)
All PCR was performed using KAPA HiFi HotStart ReadyMix (Kapa Biosystems; KK2602).
The PCR products were isolated by QIAquick PCR Purification Kit (Qiagen; 28104) and
subjected to Sanger sequencing. To validate hits, the sgRNAs were cloned in lentiCRISPR v2
vector, and transduced into 4T07-TGL cells. Transduced clones were injected through tail vein
of 6-week-old female BALB/c mice and lung metastases were monitored weekly by
bioluminescent imaging.
505
Indel analysis for validating a knockout screening.

A pair of primers was designed to amplify the 250-300 bp region centered around the sgRNA cut
site (GRCm39 chr.14 73752726-73752745). 300 ng of genomic DNA isolated in *Genome-wide*
CRISPR-Cas9 screening was amplified by PCR with the following primers and thermocycling
parameters (*Med4* indel primer_F: 5'-CACGGCCATGGCTTTCTAAC-3', *Med4* indel primer_R:
5'- GAGGGTTGTGAGCTACGTGT-3'; 95°C for 2 min, 30 cycles of (98°C for 1 s, 60°C for 5 s,
72°C for 35 s), and 72°C for 1 min). PCR was performed using NEBNext® High-Fidelity 2X
PCR Master Mix (NEB; M0541S). The purified PCR products were separated by a 4-20% TBE
gel (Invitrogen; EC62252BOX) and stained with GelRed for 1 hr. The bands were visualized by
ChemiDoc (Bio-Rad).
516
DNA constructs for overexpression and silencing.

The cDNA encoding human *MED4* was obtained from Addgene (15425). The cDNA was sub-
cloned in FLAG-HA-pcDNA3.1- (Addgene; 52535) to generate N-terminally Flag-HA tagged
MED4 cDNA. The N-term Flag-HA-MED4 cDNA was sub-cloned in pCDH-EF1-MCS-IRES-
Neo (System Biosciences; CD502A-1) and verified by sequencing. The shRNAs were designed
using the SplashRNA algorithm(70). shRNA sequences are shown in Supplementary Table S1.

Cell growth assay

Cells were seeded at a density of 1.0 x 10^5 cells onto 6 cm dishes (D2A1-d cells) or 10 cm dishes
(4T07 cells) at Day 0. Cells were trypsinized and counted at the indicated.
526
Tumorsphere assay

Single cells suspension of 4T07, D2A1-d, and D2A1 cells (10,000 cells/ 2 ml) were plated on 6-
well ultra-low attachment plates for tumorsphere cultures. Cells were cultured in serum free
MEBM supplemented with 1:50 B27, 10 ng/ml bFGF (BD Biosciences;) and 20 ng/ml EGF (BD
Biosciences;) and 4 ug/ml insulin, 4 ug/ml heparin, and 0.4% BSA for 10 days. Tumorsphere
were visualized under phase contrast microscope, photographed and counted by GelCounter and
represented graphically.
534

Soft agar colony formation assay

6-well culture plates were coated with 1 ml of 0.6 % Low Melt Agarose (Bio-Rad; 1613111). Cells were diluted at a density of 1.0×10^4 cells/ml in media containing 0.3% Low Melt Agarose and seeded 1 ml/ well. After 10 days, cells were stained with 0.005% crystal violet in 10% neutral buffered formalin (NBF) for 30 min at R.T. Soft agar was destained with distilled water until the background got clear. Colonies were scanned for visualization and counted manually.

3D culture

8-well chamber slides were coated with either 100 µl of 100% Matrigel (Corning; 354263) or 50% Matrigel + 50% Collagen I (Corning; 354236) at 37°C for 30 min prior to cell seeding (71). Cells were seeded onto the matrix at a density of 400 cells/100 µl/well.

Clonogenicity assay

For measuring clonogenicity capacities, D2A1-d and 4T07 cells were seeded onto 10 cm dishes at a density of 1,000 cells. After 6 days, the morphology and size of colonies were captured by a phase-contrast microscopy. Cells were fixed with 100% methanol for 20 min and stained with 0.5% crystal violet in 20% methanol for 5 min. Then plates were destained with PBS until the background got clear. The images of whole plates were taken by a scanner and colonies were counted manually.

Transwell migration/invasion assay

Migration assays were performed as previously described in (72) with minor modification. using uncoated cell culture inserts with 8 µm pores (Corning; 3422) according to manufacturer's instructions. Briefly, D2A1-d and 4T07 cells were harvested with Trypsin/EDTA and resuspended in serum-free medium at a density of 1.0×10^5 cells/ml for D2A1-d and 2.0×10^5 cells/ml for 4T07. 100 µl of cells were seeded into the upper chamber of uncoated cell culture inserts with 8 µm pores (Corning; 3422). Lower chambers were filled with 650 µl medium containing 10% FBS as a chemoattractant. After 18 hours, cells were fixed and permeabilized with 70% ethanol for 15 min and stained with 0.1% crystal violet/10% ethanol for 20 min. The non-migrating cells on the upper surface of the filter were removed by cotton swab. The number of migratory cells was measured by counting at $\times$ 200 magnifications using a microscope. For invasion assays, Matrigel was diluted in ice-cold serum-free medium to a concentration of 300 µg/ml. 100 µl of diluted Matrigel solution was added into the upper chamber and incubated at 37°C for 30 min to coat the membrane. D2A1-d (1.0×10^4 cells) and 4T07 (3.0×10^4 cells) cells were seeded into the upper chamber and incubated for 24 hours. The following procedure was performed as described above.

Micrococcal nuclease digestion assay (MNase digestion assay)

Global chromatin accessibility was assessed by MNase digestion assay as described previously (73) with minor modification. 1 million cells were cross-linked with 1% formaldehyde for 10 min at R. T. and quenched with 125 mM glycine for 5 min. Cells were washed once with cold PBS and collected by centrifuging at 1,000 rpm for 5 min at 4°C. Fixed cells were re-suspended

in lysis buffer (10 mM Tris-HCl, pH7.5, 10 mM NaCl, 0.5% NP-40) and incubated on ice for 10
min. After centrifugation, the cell pellets ware washed once with MNase digestion buffer (20
mM Tris-HCl, pH7.5, 15 mM NaCl, 60 mM KCl, 1 mM CaCl$_2$). After re-suspended in 250 µl
MNase digestion buffer with proteinase inhibitors, 62.5 units per 1 million cells MNase (NEB;
M0247S) was added to resuspended pellets. After incubation at 37°C for the indicated time, 250
µl of STOP buffer (100 mM Tris-HCl, pH 8.1, 20 mM EDTA, 200 mM NaCl, 2% Triton X-100,
0.2% sodium deoxycholate) was added. Lysates were sonicated for 4 cycles (30 sec on and 30
sec off) in Bioruptor and supernatant was collected after centrifugation. The supernatant was
reverse-crosslinked at 65°C overnight and treated with RNase A at 37°C for 1 h and Proteinase
K at 55°C for 2 h. The DNA fragments were purified by ethanol precipitation and analyzed by
agarose gel electrophoresis.
588

Immunofluorescence

For staining cells on 2D culture, cells were rinsed with PBS once and fixed with 4%
paraformaldehyde (PFA) in PBS for 15 min. Cells were washed three times with PBS for 5 min
followed by permeabilizing with 0.5% Triton X-100/PBS for 5 min. After rinsing once with PBS,
cells were blocked with 1% BSA/0.3% Triton X-100/PBS for 1 h. Next, cells were incubated in
primary antibodies diluted in the blocking buffer at 4°C overnight followed by incubating in the
secondary antibodies for 1 h R.T. After washing once with PBS, nuclei were counter-stained
with DAPI (1:1000 in PBS) for 5 min followed by mounting.
For staining cells on 3D culture, cells were fixed with 2% PFA in PBS supplemented with 50
mM glycine for 30 min at R.T. After washing three times for 20 min with PBS supplemented
with 50 mM glycine, cells were blocked with 10% normal goat serum in the IF buffer (0.1%
BSA, 0.2% Triton X-100, 0.05% Tween-20 in PBS) at R.T. for 1 h. Primary antibodies were
diluted in the blocking buffer and subjected to cells. After overnight incubation at 4°C, cells
were washed with the IF buffer for 20 min three times and followed by incubating with
secondary antibodies (diluted in the blocking buffer) for 1 h at R.T. After washing once with the
IF buffer, nuclei were counter stained with DAPI for 5 min, followed by mounting.
605

Immunohistochemistry

Immunostaining of mouse lungs was performed as previously described (8). In brief, tissues
were fixed in 10% neutral buffered formalin for 24 hours, embedded in paraffin, sectioned, and
stained with hematoxylin and eosin (H&E) for histopathological evaluation. For
immunohistochemistry, unstained slides of sectioned tissues were deparaffinized and
epitope/antigen retrieval was performed with 10 mM sodium citrate (pH 6.0). The slides were
treated with 3% H2O2 to block endogenous peroxidase. The slides were then incubated in 5%
normal goat serum/5% BSA/0.3% Triton X-100/PBS and stained with the appropriate primary
antibodies. After an overnight incubation in primary antibodies, slides were stained with
secondary antibodies and DAPI before mounting with the Vectashield mounting medium (Vector
Laboratories).

Western blot

Protein lysates were harvested with RIPA lysis buffer (Sigma Aldrich; R0278) containing
protease inhibitor cocktail (Thermo Fisher Scientific; 78429), phosphatase inhibitor 2 and 3

(Sigma Aldrich; P5726and P0044). Samples were standardized for protein concentration using
the Pierce BCA protein assay (VWR 23227), and incubated at 100 °C for 10 min under
reducing conditions. After denaturation, samples were separated by Bolt 4–12% Bis-Tris Plus
Gels (Thermo Fisher Scientific NW04125BOX) and transferred onto a PVDF membrane using
iBlot Transfer Stacks (Thermo Fisher Scientific IB401001). Blots were blocked with 5% bovine
serum albumin (BSA) in TBST for 1 h at room temperature. Blots were then probed with
different primary antibodies (listed in Supplementary Table S2) in the blocking buffer overnight
at 4 °C. Blots were washed with TBST for 10 min three times before incubation with secondary
antibodies [Anti-rabbit IgG-HRP (Cell Signaling Technology 7074S, 1:5000), Anti-mouse IgG-
HRP (Cell Signaling Technology 7076S, 1:5000), anti-goat IgG-HRP (Santa Cruz Biotechnology
sc-2354, 1:5000)] in 2.5% BLOT-QuickBlocker (G Biosciences 786-011) in TBST for 1 h at
room temperature. Blots were washed with TBST and imaged using chemiluminescent substrate
[Pierce ECL (Thermo Fisher Scientific 32209), SuperSignal West Pico PLUS (Thermo Fisher
Scientific 34577), or SuperSignal West Femto (Thermo Fisher Scientific 34096)] on the
ChemiDox XRS + (Bio-Rad).

Real time PCR analysis

Total RNA was extracted using the Maxwell® RSC simplyRNA Cells and RNase-free DNase
Kit (Promega; AS1390) using an automated RNA/DNA isolation instrument (Maxwell,
Promega). cDNAs were synthesized using High-Capacity cDNA Reverse Transcription Kit
(Applied Biosystems; 4368814). cDNA corresponding to approximately 10 ng of starting RNA
was used for one reaction. qRT-PCR was performed with PowerUp SYBR Green Master Mix
(Applied Biosystems) on QuantStudio™ 7 Flex Real-Time PCR System. All quantifications
were normalized to endogenous GAPDH. Primer sequences used to amplify genes were
purchased from Sigma-Aldrich and are listed in Supplementary Table S3.

Bulk RNA-sequencing and analysis

Total RNA was isolated as described above. RNA-sequencing was performed by the BGI
sequencing facility in Hong Kong. RNA quality, including RNA concentration, RIN value,
28S/18S ratio, and fragment length distribution was confirmed with an Agilent 2100 Bio
analyzer (Agilent RNA 6000 Nano Kit). Libraries were prepared by using the standard
methodology and run on a BGISEQ-500 platform. Raw reads were quality-checked and
subsequently mapped to the mouse genome (mm10_UCSC) through the Spliced Transcripts
Alignment to a Reference (STAR)(74), which outputs a BAM file containing both the sequence
and its genomic coordinate. Together with a General Transfer Format (GTF) the file containing
all the genomic annotations were used to determine the total number of reads per transcript using
the HTSeq Count program.(75) Differential gene expression was analyzed by using the DESeq2
algorithm(76) with |fold change| $\geq$ 2 and FDR < 0.05. Gene set enrichment analysis (GSEA) was
performed on a pre-ranked gene list that was generated based on the gene expression changes
between shMed4 and control cells(77). Gene expression data were transformed in logarithmic
scale and normalized accordingly. Heatmaps showing the gene expression differences between
control and Med4 KD cells were generated by using the Morpheus online tool
(https://software.broadinstitute.org/morpheus/).

ATAC-sequencing and analysis

Chromatin accessibility assays utilizing the bacterial Tn5 transposase were performed as
described with minor modifications (78). Cells (5.0 x 10^4) were lysed and incubated with
transposition reaction mix for 30 minutes at 37°C. Samples were PCR-amplified and sequenced
on an Illumina NextSeq 500.

Bulk ATAC-seq raw data were pre-processed using the nfcore/atacseq pipeline version 2.0
(10.5281/zenodo.2634132), implemented in Nextflow (version 22.10.2). Reads were trimmed for
both quality and Illumina adapter sequences using 'trim_galore' then aligned to mouse assembly
mm10 with BWA using the default parameters. Aligned reads with the same start site and
orientation were removed using the Picard tool MarkDuplicates
(http://broadinstitute.github.io/picard). Density profiles were created by extending each read to
the average library fragment size and then computing density using the BEDTools suite
(http://bedtools.readthedocs.io). Reads were not extended when generating ATAC-seq read
density. Enriched regions were discovered using MACS2 and scored against matched input
libraries (|fold change| > 2 and p-value < 0.005). Peaks were then filtered against genomic
'blacklisted' regions and filtered peaks within 500 bp were merged to create a full peak atlas.
Raw read counts were tabulated over this peak atlas using featureCounts
(http://subread.sourceforge.net). All genome browser tracks and read density tables were
normalized to a sequencing depth of ten million mapped reads. Peaks were annotated using
linear genomic distance, with a gene assigned to a peak if it was within 50 kb upstream or
downstream of the gene start or end, respectively. Motif signatures were obtained using the 'de
novo' approach in Homer v4.7.2 (http://homer.ucsd.edu). The 'DESeq2' package in R was
employed to identify differential gene expressions and accessibility with the threshold of
absolute log2 FoldChange > log2 (1.5) and FDR ≤ 0.1. When a single gene was linked to
multiple ATAC-seq peaks, all associated peaks were retained for analysis. A contingency plot
was used to visualize the integrated results from ATAC-Seq and RNA-Seq data. Each quadrant
in the plot indicates the statistical significance of either high or low expression for gene
expression or accessibility.

Statistics and reproducibility

GraphPad 8 software and R were used for statistical analysis. Results were represented as means
±□standard deviations (SDs). The normality of the population distribution will be assessed by
Shapiro-Wilk test (a significance level of α = 0.05) followed by visualizing with Q-Q plot or
histogram. For *in vitro* studies, statistical significance between groups was performed by using
an unpaired Student's t-test with a minimum of three biologically independent samples. For *in*
vivo studies, between-group differences were assessed using the Mann-Whitney U test as a
nonparametric method with a minimum of five biologically independent samples. Statistical
significance was determined when p value was less than 0.05.
698

References

1. Early Breast Cancer Trialists' Collaborative G. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. Lancet **2005**;365(9472):1687-717 doi 10.1016/S0140-6736(05)66544-0.
2. Giancotti FG. Mechanisms governing metastatic dormancy and reactivation. Cell **2013**;155(4):750-64 doi 10.1016/j.cell.2013.10.029.
3. Zhang XH, Giuliano M, Trivedi MV, Schiff R, Osborne CK. Metastasis dormancy in estrogen receptor-positive breast cancer. Clin Cancer Res **2013**;19(23):6389-97 doi 10.1158/1078-0432.CCR-13-0838.
4. Turajlic S, Swanton C. Metastasis as an evolutionary process. Science **2016**;352(6282):169-75 doi 10.1126/science.aaf2784.
5. Lambert AW, Pattabiraman DR, Weinberg RA. Emerging Biological Principles of Metastasis. Cell **2017**;168(4):670-91 doi 10.1016/j.cell.2016.11.037.
6. Gao H, Chakraborty G, Lee-Lim AP, Mavrakis KJ, Wendel HG, Giancotti FG. Forward genetic screens in mice uncover mediators and suppressors of metastatic reactivation. Proc Natl Acad Sci U S A **2014**;111(46):16532-7 doi 10.1073/pnas.1403234111.
7. Gao H, Chakraborty G, Lee-Lim AP, Mo Q, Decker M, Vonica A, *et al.* The BMP inhibitor Coco reactivates breast cancer cells at lung metastatic sites. Cell **2012**;150(4):764-79 doi 10.1016/j.cell.2012.06.035.
8. Gao H, Chakraborty G, Zhang Z, Akalay I, Gadiya M, Gao Y, *et al.* Multi-organ Site Metastatic Reactivation Mediated by Non-canonical Discoidin Domain Receptor 1 Signaling. Cell **2016**;166(1):47-62 doi 10.1016/j.cell.2016.06.009.
9. Su W, Han HH, Wang Y, Zhang B, Zhou B, Cheng Y, *et al.* The Polycomb Repressor Complex 1 Drives Double-Negative Prostate Cancer Metastasis by Coordinating Stemness and Immune Suppression. Cancer Cell **2019**;36(2):139-55 e10 doi 10.1016/j.ccell.2019.06.009.
10. Whyte WA, Orlando DA, Hnisz D, Abraham BJ, Lin CY, Kagey MH, *et al.* Master transcription factors and mediator establish super-enhancers at key cell identity genes. Cell **2013**;153(2):307-19 doi 10.1016/j.cell.2013.03.035.
11. El Khattabi L, Zhao H, Kalchschmidt J, Young N, Jung S, Van Blerkom P, *et al.* A Pliable Mediator Acts as a Functional Rather Than an Architectural Bridge between Promoters and Enhancers. Cell **2019**;178(5):1145-58 e20 doi 10.1016/j.cell.2019.07.011.
12. Verger A, Monte D, Villeret V. Twenty years of Mediator complex structural studies. Biochem Soc Trans **2019**;47(1):399-410 doi 10.1042/BST20180608.
13. Abdella R, Talyzina A, Chen S, Inouye CJ, Tjian R, He Y. Structure of the human Mediator-bound transcription preinitiation complex. Science **2021**;372(6537):52-6 doi 10.1126/science.abg3074.
14. Soutourina J. Transcription regulation by the Mediator complex. Nat Rev Mol Cell Biol **2018**;19(4):262-74 doi 10.1038/nrm.2017.115.
15. Allen BL, Taatjes DJ. The Mediator complex: a central integrator of transcription. Nat Rev Mol Cell Biol **2015**;16(3):155-66 doi 10.1038/nrm3951.
16. Ito M, Yuan CX, Okano HJ, Darnell RB, Roeder RG. Involvement of the TRAP220 component of the TRAP/SMCC coactivator complex in embryonic development and thyroid hormone action. Mol Cell **2000**;5(4):683-93 doi 10.1016/s1097-2765(00)80247-6.

745   17.   Stevens JL, Cantin GT, Wang G, Shevchenko A, Shevchenko A, Berk AJ. Transcription
746   control by E1A and MAP kinase pathway via Sur2 mediator subunit. Science
747   **2002**;296(5568):755-8 doi 10.1126/science.1068943.
748   18.   Barbieri CE, Baca SC, Lawrence MS, Demichelis F, Blattner M, Theurillat JP, *et al.*
749   Exome sequencing identifies recurrent SPOP, FOXA1 and MED12 mutations in prostate
750   cancer. Nat Genet **2012**;44(6):685-9 doi 10.1038/ng.2279.
751   19.   Joung J, Konermann S, Gootenberg JS, Abudayyeh OO, Platt RJ, Brigham MD, *et al.*
752   Genome-scale CRISPR-Cas9 knockout and transcriptional activation screening. Nat
753   Protoc **2017**;12(4):828-63 doi 10.1038/nprot.2017.016.
754   20.   Tsherniak A, Vazquez F, Montgomery PG, Weir BA, Kryukov G, Cowley GS, *et al.*
755   Defining a Cancer Dependency Map. Cell **2017**;170(3):564-76 e16 doi
756   10.1016/j.cell.2017.06.010.
757   21.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, *et al.* The cBio cancer
758   genomics portal: an open platform for exploring multidimensional cancer genomics data.
759   Cancer Discov **2012**;2(5):401-4 doi 10.1158/2159-8290.CD-12-0095.
760   22.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, *et al.* Integrative
761   analysis of complex cancer genomics and clinical profiles using the cBioPortal. Sci
762   Signal **2013**;6(269):pl1 doi 10.1126/scisignal.2004088.
763   23.   Lefebvre C, Bachelot T, Filleron T, Pedrero M, Campone M, Soria JC, *et al.* Mutational
764   Profile of Metastatic Breast Cancers: A Retrospective Analysis. PLoS Med
765   **2016**;13(12):e1002201 doi 10.1371/journal.pmed.1002201.
766   24.   Siegel MB, He X, Hoadley KA, Hoyle A, Pearce JB, Garrett AL, *et al.* Integrated RNA
767   and DNA sequencing reveals early drivers of metastatic breast cancer. J Clin Invest
768   **2018**;128(4):1371-83 doi 10.1172/JCI96153.
769   25.   Gyorffy B, Lanczky A, Eklund AC, Denkert C, Budczies J, Li Q, *et al.* An online
770   survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer
771   prognosis using microarray data of 1,809 patients. Breast Cancer Res Treat
772   **2010**;123(3):725-31 doi 10.1007/s10549-009-0674-9.
773   26.   Valastyan S, Weinberg RA. Tumor metastasis: molecular insights and evolving
774   paradigms. Cell **2011**;147(2):275-92 doi 10.1016/j.cell.2011.09.024.
775   27.   Pastushenko I, Blanpain C. EMT Transition States during Tumor Progression and
776   Metastasis. Trends Cell Biol **2019**;29(3):212-26 doi 10.1016/j.tcb.2018.12.001.
777   28.   De Cock JM, Shibue T, Dongre A, Keckesova Z, Reinhardt F, Weinberg RA.
778   Inflammation Triggers Zeb1-Dependent Escape from Tumor Latency. Cancer Res
779   **2016**;76(23):6778-84 doi 10.1158/0008-5472.CAN-16-0608.
780   29.   Hnisz D, Abraham BJ, Lee TI, Lau A, Saint-Andre V, Sigova AA, *et al.* Super-enhancers
781   in the control of cell identity and disease. Cell **2013**;155(4):934-47 doi
782   10.1016/j.cell.2013.09.053.
783   30.   Mieczkowski J, Cook A, Bowman SK, Mueller B, Alver BH, Kundu S, *et al.* MNase
784   titration reveals differences between nucleosome occupancy and chromatin accessibility.
785   Nat Commun **2016**;7:11485 doi 10.1038/ncomms11485.
786   31.   Ishihara S, Haga H. Matrix Stiffness Contributes to Cancer Progression by Regulating
787   Transcription Factors. Cancers (Basel) **2022**;14(4) doi 10.3390/cancers14041049.
788   32.   Paszek MJ, Zahir N, Johnson KR, Lakins JN, Rozenberg GI, Gefen A, *et al.* Tensional
789   homeostasis and the malignant phenotype. Cancer Cell **2005**;8(3):241-54 doi
790   10.1016/j.ccr.2005.08.010.

33. Ueno H, Shinto E, Hashiguchi Y, Shimazaki H, Kajiwara Y, Sueyama T, *et al.* In rectal
cancer, the type of desmoplastic response after preoperative chemoradiotherapy is
associated with prognosis. Virchows Arch **2015**;466(6):655-63 doi 10.1007/s00428-015-
1756-1.
34. Sun F, Sun T, Kronenberg M, Tan X, Huang C, Carey MF. The Pol II preinitiation
complex (PIC) influences Mediator binding but not promoter-enhancer looping. Genes
Dev **2021**;35(15-16):1175-89 doi 10.1101/gad.348471.121.
35. Wreggett KA, Hill F, James PS, Hutchings A, Butcher GW, Singh PB. A mammalian
homologue of Drosophila heterochromatin protein 1 (HP1) is a component of constitutive
heterochromatin. Cytogenet Cell Genet **1994**;66(2):99-103 doi 10.1159/000133676.
36. van Steensel B, Belmont AS. Lamina-Associated Domains: Links with Chromosome
Architecture, Heterochromatin, and Gene Repression. Cell **2017**;169(5):780-91 doi
10.1016/j.cell.2017.04.022.
37. Reddy KL, Zullo JM, Bertolino E, Singh H. Transcriptional repression mediated by
repositioning of genes to the nuclear lamina. Nature **2008**;452(7184):243-7 doi
10.1038/nature06727.
38. Peter M, Nakagawa J, Doree M, Labbe JC, Nigg EA. In vitro disassembly of the nuclear
lamina and M phase-specific phosphorylation of lamins by cdc2 kinase. Cell
1990;61(4):591-602 doi 10.1016/0092-8674(90)90471-p.
39. Heald R, McKeon F. Mutations of phosphorylation sites in lamin A that prevent nuclear
lamina disassembly in mitosis. Cell **1990**;61(4):579-89 doi 10.1016/0092-
8674(90)90470-y.
40. Janota CS, Calero-Cuenca FJ, Gomes ER. The role of the cell nucleus in
mechanotransduction. Curr Opin Cell Biol **2020**;63:204-11 doi
10.1016/j.ceb.2020.03.001.
41. Medjkane S, Perez-Sanchez C, Gaggioli C, Sahai E, Treisman R. Myocardin-related
transcription factors and SRF are required for cytoskeletal dynamics and experimental
metastasis. Nat Cell Biol **2009**;11(3):257-68 doi 10.1038/ncb1833.
42. Egeblad M, Rasch MG, Weaver VM. Dynamic interplay between the collagen scaffold
and tumor evolution. Curr Opin Cell Biol **2010**;22(5):697-706 doi
10.1016/j.ceb.2010.08.015.
43. Kechagia JZ, Ivaska J, Roca-Cusachs P. Integrins as biomechanical sensors of the
microenvironment. Nat Rev Mol Cell Biol **2019**;20(8):457-73 doi 10.1038/s41580-019-
0134-2.
44. Case LB, Waterman CM. Integration of actin dynamics and cell adhesion by a three-
dimensional, mechanosensitive molecular clutch. Nat Cell Biol **2015**;17(8):955-63 doi
10.1038/ncb3191.
45. Jones SF, Siu LL, Bendell JC, Cleary JM, Razak AR, Infante JR, *et al.* A phase I study of
VS-6063, a second-generation focal adhesion kinase inhibitor, in patients with advanced
solid tumors. Invest New Drugs **2015**;33(5):1100-7 doi 10.1007/s10637-015-0282-y.
46. Keenan AB, Torre D, Lachmann A, Leong AK, Wojciechowicz ML, Utti V, *et al.*
ChEA3: transcription factor enrichment analysis by orthogonal omics integration.
Nucleic Acids Res **2019**;47(W1):W212-W24 doi 10.1093/nar/gkz446.
47. Esnault C, Stewart A, Gualdrini F, East P, Horswell S, Matthews N, *et al.* Rho-actin
signaling to the MRTF coactivators dominates the immediate transcriptional response to
serum in fibroblasts. Genes Dev **2014**;28(9):943-58 doi 10.1101/gad.239327.114.

48.   Cordenonsi M, Zanconato F, Azzolin L, Forcato M, Rosato A, Frasson C*, et al.* The
Hippo transducer TAZ confers cancer stem cell-related traits on breast cancer cells. Cell
2011;147(4):759-72 doi 10.1016/j.cell.2011.09.048.
49.   Zhao B, Ye X, Yu J, Li L, Li W, Li S*, et al.* TEAD mediates YAP-dependent gene
induction and growth control. Genes Dev **2008**;22(14):1962-71 doi 10.1101/gad.1664408.
50.   Posern G, Treisman R. Actin' together: serum response factor, its cofactors and the link to
signal transduction. Trends Cell Biol **2006**;16(11):588-96 doi 10.1016/j.tcb.2006.09.008.
51.   Calvo F, Ege N, Grande-Garcia A, Hooper S, Jenkins RP, Chaudhry SI*, et al.*
Mechanotransduction and YAP-dependent matrix remodelling is required for the
generation and maintenance of cancer-associated fibroblasts. Nat Cell Biol
2013;15(6):637-46 doi 10.1038/ncb2756.
52.   Foster CT, Gualdrini F, Treisman R. Mutual dependence of the MRTF-SRF and YAP-
TEAD pathways in cancer-associated fibroblasts is indirect and mediated by cytoskeletal
dynamics. Genes Dev **2017**;31(23-24):2361-75 doi 10.1101/gad.304501.117.
53.   Dupont S, Morsut L, Aragona M, Enzo E, Giulitti S, Cordenonsi M*, et al.* Role of
YAP/TAZ in mechanotransduction. Nature **2011**;474(7350):179-83 doi
10.1038/nature10137.
54.   Yin JW, Wang G. The Mediator complex: a master coordinator of transcription and cell
lineage development. Development **2014**;141(5):977-87 doi 10.1242/dev.098392.
55.   Rasool RU, Natesan R, Deng Q, Aras S, Lal P, Sander Effron S*, et al.* CDK7 Inhibition
Suppresses Castration-Resistant Prostate Cancer through MED1 Inactivation. Cancer
Discov **2019**;9(11):1538-55 doi 10.1158/2159-8290.CD-19-0189.
56.   Yang Y, Leonard M, Zhang Y, Zhao D, Mahmoud C, Khan S*, et al.* HER2-Driven Breast
Tumorigenesis Relies upon Interactions of the Estrogen Receptor with Coactivator
MED1. Cancer Res **2018**;78(2):422-35 doi 10.1158/0008-5472.CAN-17-1533.
57.   Tsang FH, Law CT, Tang TC, Cheng CL, Chin DW, Tam WV*, et al.* Aberrant Super-
Enhancer Landscape in Human Hepatocellular Carcinoma. Hepatology **2019**;69(6):2502-
17 doi 10.1002/hep.30544.
58.   Lyng H, Lando M, Brovig RS, Svendsrud DH, Johansen M, Galteland E*, et al.*
GeneCount: genome-wide calculation of absolute tumor DNA copy numbers from array
comparative genomic hybridization data. Genome Biol **2008**;9(5):R86 doi 10.1186/gb-
2008-9-5-r86.
59.   Albrecht P, Ansperger-Rescher B, Schuler A, Zeschnigk M, Gallie B, Lohmann DR.
Spectrum of gross deletions and insertions in the RB1 gene in patients with
retinoblastoma and association with phenotypic expression. Hum Mutat **2005**;26(5):437-
45 doi 10.1002/humu.20234.
60.   Mitter D, Ullmann R, Muradyan A, Klein-Hitpass L, Kanber D, Ounap K*, et al.*
Genotype-phenotype correlations in patients with retinoblastoma and interstitial 13q
deletions. Eur J Hum Genet **2011**;19(9):947-58 doi 10.1038/ejhg.2011.58.
61.   Dechat T, Pfleghaar K, Sengupta K, Shimi T, Shumaker DK, Solimando L*, et al.* Nuclear
lamins: major factors in the structural organization and function of the nucleus and
chromatin. Genes Dev **2008**;22(7):832-53 doi 10.1101/gad.1652708.
62.   Tajik A, Zhang Y, Wei F, Sun J, Jia Q, Zhou W*, et al.* Transcription upregulation via
force-induced direct stretching of chromatin. Nat Mater **2016**;15(12):1287-96 doi
10.1038/nmat4729.

63.   Iyer KV, Pulford S, Mogilner A, Shivashankar GV. Mechanical activation of cells
induces chromatin remodeling preceding MKL nuclear transport. Biophys J
2012;103(7):1416-28 doi 10.1016/j.bpj.2012.08.041.
64.   Sun J, Chen J, Mohagheghian E, Wang N. Force-induced gene up-regulation does not
follow the weak power law but depends on H3K9 demethylation. Sci Adv
2020;6(14):eaay9095 doi 10.1126/sciadv.aay9095.
65.   Olson EN, Nordheim A. Linking actin dynamics and gene transcription to drive cellular
motile functions. Nat Rev Mol Cell Biol **2010**;11(5):353-65 doi 10.1038/nrm2890.
66.   O'Connor JW, Riley PN, Nalluri SM, Ashar PK, Gomez EW. Matrix Rigidity Mediates
TGFbeta1-Induced Epithelial-Myofibroblast Transition by Controlling Cytoskeletal
Organization and MRTF-A Localization. J Cell Physiol **2015**;230(8):1829-39 doi
10.1002/jcp.24895.
67.   Buyukcelebi K, Chen X, Abdula F, Elkafas H, Duval AJ, Ozturk H, *et al.* Engineered
MED12 mutations drive leiomyoma-like transcriptional and metabolic programs by
altering the 3D genome compartmentalization. Nat Commun **2023**;14(1):4057 doi
10.1038/s41467-023-39684-y.
68.   Berry DA, Cronin KA, Plevritis SK, Fryback DG, Clarke L, Zelen M, *et al.* Effect of
screening and adjuvant therapy on mortality from breast cancer. N Engl J Med
2005;353(17):1784-92 doi 10.1056/NEJMoa050518.
69.   Aslakson CJ, Miller FR. Selective events in the metastatic process defined by analysis of
the sequential dissemination of subpopulations of a mouse mammary tumor. Cancer Res
1992;52(6):1399-405.
70.   Pelossof R, Fairchild L, Huang CH, Widmer C, Sreedharan VT, Sinha N, *et al.* Prediction
of potent shRNAs with a sequential classification algorithm. Nat Biotechnol
2017;35(4):350-3 doi 10.1038/nbt.3807.
71.   Nguyen-Ngoc KV, Shamir ER, Huebner RJ, Beck JN, Cheung KJ, Ewald AJ. 3D culture
assays of murine mammary branching morphogenesis and epithelial invasion. Methods
Mol Biol **2015**;1189:135-62 doi 10.1007/978-1-4939-1164-6_10.
72.   Bae SY, Kim HJ, Lee KJ, Lee K. Translationally controlled tumor protein induces
epithelial to mesenchymal transition and promotes cell migration, invasion and metastasis.
Sci Rep **2015**;5:8061 doi 10.1038/srep08061.
73.   Cheng L, Zhang X, Wang Y, Gan H, Xu X, Lv X, *et al.* Chromatin Assembly Factor 1
(CAF-1) facilitates the establishment of facultative heterochromatin during pluripotency
exit. Nucleic Acids Res **2019**;47(21):11114-31 doi 10.1093/nar/gkz858.
74.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, *et al.* STAR: ultrafast
universal RNA-seq aligner. Bioinformatics **2013**;29(1):15-21 doi
10.1093/bioinformatics/bts635.
75.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput
sequencing data. Bioinformatics **2015**;31(2):166-9 doi 10.1093/bioinformatics/btu638.
76.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol **2014**;15(12):550 doi 10.1186/s13059-014-
0550-8.
77.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, *et al.*
Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide
expression profiles. Proc Natl Acad Sci U S A **2005**;102(43):15545-50 doi
10.1073/pnas.0506580102.

78. Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native
chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding
proteins and nucleosome position. Nat Methods **2013**;10(12):1213-8 doi
10.1038/nmeth.2688.
932

Acknowledgments

This study is dedicated to the memory of our mentor, teacher, and friend, Filippo Giancotti, a pioneer in the metastasis field known for his brilliance, rigor, and kindness. This work was supported by NIH grant R35 CA197566 and partially by CPRIT Recruitment of Established Investigators Award RR160031 (F.G.G.). Use of the Core Facilities was supported by NIH grants P30CA008748 (MSKCC), P30 CA016672 (MDACC), and P30 CA013696 (Columbia University).

Author's Contributions

S. Bae: Conceptualization, validation, investigation, methodology, writing-original draft, writing-review and editing. **F.G. Giancotti:** Conceptualization, funding acquisition **Y. Chen:** Formal analysis and validation. **H. Chen:** Formal analysis and validation. **S. Karaiskos:** Formal analysis and validation. **J.J. Xu:** Formal analysis and validation. **C. Lu:** Formal analysis, investigation and validation. **A.D. Viny:** Validation, investigation, writing-original draft, writing-review and editing.

Figure Legends

Figure 1. *Med4* **was identified as a gatekeeper for metastatic reactivation by forward**
genetic screening using CRISPR-Cas9 system. A, Schematic representation of the *in vivo*
CRISPR-Cas9 screening strategy. **B,** The scheme of chromatin looping for enhancer-promoter
interaction. **C,** RT-qPCR analysis and immunoblots of *Med4* expression in D2A1-d-TGL cells
expressing MirE shRNA targeting either non-targeting control (NTC) or *Med4*; Mean values (±
S.D.) p values; unpaired Student's t-test. **D-F,** D2A1-d-TGL cells transduced with either NTC or
Med4 shRNAs were inoculated *i.v.* in syngeneic mice. Lung metastasis was measured by
bioluminescent (BLI) imaging at the indicated time. The panels show representative images (D),
the graphs of the normalized BLI photon flux (E), and the graphs of the counts of
macrometastases (F). Mean values (± S.D.) p values; Mann-Whitney U test (E and F). **G-J,**
Schematic representation of inducible expression experiment (G), Immunoblot analysis of
MED4 in D2A1-d-TGL cells transduced with doxycycline (Dox)-inducible MirE shRNA
targeting NTC or *Med4* (H), D2A1-d-TGL cells expressing Dox-inducible MirE shRNA
targeting NTC or *Med4* were inoculated *i.v.* in syngeneic mice. Dox was administrated either
immediately after injection (Dox day 0) or 14 days after inoculation of the cells (Dox day 14).
Lung metastasis was measured by BLI normalized photon flux at the indicated time. The panels
show representative images (I), and the graphs of the normalized BLI photon flux (J); Mean
values (± S.D.) p values; Mann-Whitney U test. **K,** Copy number alterations in metastatic breast
cancer patients (from INSERM) were analyzed by cBioportal database. **L,** *MED4* mRNA levels
were measured by paired RNA-seq of primary tumors and multiple metastases. **M,** Kaplan-Meier
analysis of relapse-free survival (RFS) of all patients in publicly available breast cancer datasets
(source KM Plotter for breast cancer). Patients were stratified according to *MED4* expression
using a probe set, 222438_at as indicated. HR, hazard ratio.

Figure 2. The downregulation of *Med4* **leads to the reprogramming of the transcriptional**
and epigenetic landscape, bolstering matrix stiffening. A, Heatmap of differentially expressed
genes (DEGs; |Fold change| ≥ 2, FDR < 0.05) in NTC versus *Med4*-silenced D2A1-d-TGL cells.
B, MNase digestion assay was performed in nuclei isolated from the indicated cells for
increasing incubation time. The panels show the representative image (upper), and the
quantification graph (lower). **C,** Differential accessibility ($\log_2$ fold change in reads per
accessible region) plotted against the mean reads per region. **D,** Intersection of RNA-seq and
ATAC-seq plotted as $-\log_{10}$(FDR)*sign of the $\log_2$ fold change of *Med4*-sileced D2A1-d-TGL
cells compared to NTC and each point representing one ATAC peak. **E,** Gene ontology (GO)
analysis of the genes located in the upper quadrant of (D) from the intersection of RNA-seq and
ATAC-seq. **F,** Heatmap shows the expression level of matrisome genes amongst the top
differentially expressed genes in NTC versus *Med4*-depleted cells. **G,** Immunohistochemistry for
α-SMA and Masson's trichrome staining on lung metastases in mice that were intravenously
injected with D2A1-d-TGL cells transduced with either non-targeting control (NTC) or *Med4*
shRNAs.

Figure 3. *Med4* **silencing decompacts 3D chromatin structure by pulling on the nuclear**
lamina through stress fibers. A, D2A1-d cells expressing either NTC or *Med4* shRNAs were
stained with HP1α (red). The panels display representative images (left) and fluorescence
intensity profiles across one cell (right). **B,** Cells were co-stained with H3K9me3 (red) and

LAMIN A/C (green). The panels show representative images (left) and fluorescence intensity profile of H3K9me3 across one cell (right). **C** and **D,** D2A1-d cells expressing either NTC or *Med4* shRNAs were co-transduced with either empty vector (EV) or Flag-HA tagged MED4 cDNA to rescue MED4 expression. Immunoblots of MED4 and HA (C). Cells were stained with LAMIN A/C (red) and nuclei were counter-stained with DAPI (blue). The panel shows representative images (left) and quantification of nucleus sizes in diameter (right) (D); Mean values (± S.D.) p values; unpaired Student's t-test. **E,** D2A1-d cells expressing either NTC or *Med4* shRNAs were treated with 0.1% DMSO, 200 nM cytochalasin D, or 100 nM latrunculin B for 20 min as indicated. Cells were stained with LAMIN A/C (green) and F-actin (red). Nuclei were counter-stained with DAPI (blue). The panel shows representative images (left) and quantification of nucleus sizes in volume (right); Mean values (± S.D.) p values; unpaired Student's t-test. **F,** D2A1-d cells expressing either NTC or *Med4* shRNAs were treated with 0.1% DMSO or 200 nM cytochalasin D for 20 min. Cells were stained with HP1α (red) and F-actin (green). The panels show representative images (left) and fluorescence intensity profile of HP1α across one cell (right).

Figure 4. *Med4*-deficient cells exhibit a propensity to grow within a desmoplastic niche, through the activation of integrin signaling. A, D2A1-d cells expressing MirE shRNA targeting *Med4* or NTC were cultured on 3D Matrigel with or without collagen I. **B,** Cells were cultured either on 2D or on 3D Matrigel + collagen I matrix, and stained with Ki67 (purple). Nuclei were counterstained with DAPI (blue). The panel shows representative images (top) and quantification graphs for numbers of Ki67+ cells (bottom). **C,** D2A1-d cells expressing MirE shRNA targeting Med4 cultured on 2D were stained for focal adhesions, and the nuclei were stained with DAPI (blue). **D,** Total cell lysates were subjected to immunoblotting for integrin-signaling molecules. **E,** D2A1-d cells expressing either NTC or Med4 shRNAs were cultured either on 3D Matrigel + collagen I matrix, and stained with either F-actin or p-FAK (Y^{397}). Nuclei were counter-stained with DAPI. **F,** D2A1-d cells treated with or without 2.5 μM defactinib were cultured either on 3D Matrigel + collagen I matrix.

Figure 5. *Med4* loss activates integrin-mediated mechano-transduction MRTF/SRF and YAP/TAZ pathways. A, ChEA3 TF enrichment analysis of the promoters of the genes upregulated in *Med4*-deficient cells compared to NTC. **B,** HOMER motif analysis of genes in the upper right quadrant of (Figure 2D) which represents the intersection of RNA-seq and ATAC-seq data in *Med4*-silenced D2A1-d-TGL cells compared to non-targeting control (NTC). p-values were generated using the binomial test. **C,** Gene signature enrichment analysis (GSEA) of MRTF, SRF, and YAP transcriptional signatures. NES, normalized enrichment score; FDR, false discovery rate (n = 3 biological replicates per condition). **D,** D2A1-d cells expressing MirE shRNA targeting *Med4* cultured on 2D were stained for MRTF-A (red), or for YAP (purple). The nuclei were counterstained with DAPI (blue). **E** and **F,** D2A1-d cells expressing either NTC or *Med4* shRNAs were co-transduced with either empty vector (EV) or Flag-HA tagged MED4 cDNA to rescue MED4 expression. Cells were stained with MRTF-A (red) and counter-stained with DAPI (blue). The panel shows representative images (E) and quantification of subcellular localization (F). **G,** D2A1-d-TGL cells co-transduced with indicated MirE shRNAs were inoculated *i.v.* in syngeneic mice. Lung metastasis was measured by bioluminescent (BLI) imaging at the indicated time. The panels show representative images (left), and the graphs of the normalized BLI photon flux (right). Mean values (± S.D.) p values; Mann-Whitney U test.

Supplemental figure Legends

Supplementary Figure S1. *Med4* **was hemizygously deleted to induce metastatic reactivation.**
(A) 4T07-TGL cells either mock control or expressing GeCKO v2 library were inoculated *i.v.* into BALB/c syngeneic mice. Lung metastasis was measured by bioluminescent (BLI) imaging at the indicated time.
(B) Schematics of the mouse chromosomal region encompassing *Med4*, location of the sequence complementary to the gRNA isolated from the screening, and the primer pairs used for genomic DNA PCR.
(C) The isolated clone from the screening which harbors a gRNA targeting *Med4* was subjected to an indel analysis using genomic DNA PCR (left) and immunoblotting (right).
(D) Sequences of the fragments of DNA deleted by CRISPR-Cas9. Red boxes indicate the junction. (E) Sequence alignment results of translated amino acids from each allele to wild type MED4.
(F) DepMap analysis of the dependency of tumor cell line panels in CRISPR (blue) and RNAi (violet) databases on MED4. X-axis: dependency scores.
(G) RT-qPCR analysis of *Med4* expression in 4T07-TGL cells expressing MirE shRNA targeting either non-targeting control (NTC) or *Med4*; Mean values (± S.D.) p-values; unpaired Student's t-test.
(H and I) 4T07-TGL cells transduced with either NTC or *Med4* shRNAs were inoculated *i.v.* in syngeneic mice. Lung metastasis was measured by bioluminescent (BLI) imaging at the indicated time. The panels show representative images (H), and the graphs of the normalized BLI photon flux (I). Mean values (± S.D.) p values; Mann-Whitney U test.

Supplementary Figure S2. *MED4* **loss serves as a predictor for breast cancer relapse.**
(A and B) cBioportal analysis of copy number alterations in metastatic breast cancer patients from the Metastatic Breast Cancer Project; Provisional (A) and Archived (B).
(C) Kaplan-Meier analysis of RFS in patients stratified according to *MED4* expression using a probe set, 222438_at as indicated. Patients were subdivided by breast cancer subtypes according to PAM50.
(D and E) Kaplan-Meier analysis of RFS in patients stratified according to *MED4* expression using a probe set, 222438_at as indicated. Patients were subdivided by estrogen receptor (ER) expression level (D) or by human epidermal growth factor receptor 2 (*HER2*) expression level determined by Affymetrix microarray (E).
(F and G) KM plotter analysis of overall survival (OS) based on *MED4* expression level from RNA-seq data of breast cancer patients. Patients were subdivided by estrogen receptor (ER) expression level (F) or by human epidermal growth factor receptor 2 (*HER2*) expression level determined by Affymetrix microarray (G). HR, hazard ratio.

Supplementary Figure S3. *Med4* **depletion promotes cancer stem cell traits in D2A1-d cells.**
(A) Cell growth of D2A1-d-TGL cells expressing MirE shRNA targeting either NTC or *Med4* was measured by counting cell numbers at the indicated time; Mean values (± S.D.) p values; unpaired Student's t-test (n = 3 per group, *P = 0.0171, **P = 0.0017, ***P = 0.0004).
(B) Control or Med4-silenced D2A1-d-TGL cells were subjected to tumorsphere assay. The panel shows representative images of wells and microscopic images of tumor spheres (left) and

mean numbers of organoid ($\pm$ S.D.; right) per 10,000 cells. p values; unpaired Student's t-test (n
= 3 per group, *P = 0.0122, **P = 0.0089).
(C) Indicated D2A1-d-TGL cells were subjected to clonogenicity assay. The panel shows
representative images of dishes and microscopic images of colonies (left) and mean numbers of
colonies ($\pm$ S.D.; right) per 1,000 cells. p values; unpaired Student's t-test (n = 3 per group, **P
= 0.0056, ***P = 0.0006).
(D) Indicated D2A1-d-TGL cells were subjected to soft agar assay. The panel shows
representative images of wells and microscopic images of colonies (left) and mean numbers of
colonies ($\pm$ S.D.; right) per 10,000 cells. p values; unpaired Student's t-test (n = 3 per group, *P
= 0.0118, ****P < 0.0001).
(E and F) Control or *Med4*-silenced D2A1-d-TGL cells were subjected to transwell migration
assay (E) and transwell invasion assay (F). The panel shows representative images of fields (left)
and mean numbers of cells ($\pm$ S.D.; right) per field. p values; unpaired Student's t-test ((E) n = 3
per group, ***P = 0.0007, ***P = 0.0009, (F) n = 3 per group, **P = 0.0018, ***P = 0.0007).
1100
Supplementary Figure S4. *Med4* depletion promotes cancer stem cell traits in 4T07 cells.
(A) Cell growth of 4T07-TGL cells expressing MirE shRNA targeting either NTC or *Med4* was
measured by counting cell numbers at the indicated time; Mean values ($\pm$ S.D.) p values;
unpaired Student's t-test (n = 3 per group, **P = 0.0040, **P = 0.0067, ***P = 0.0003).
(B) Tumorsphere assay results in control and *Med4*-silenced 4T07-TGL cells. The panel shows
representative images of wells and microscopic images of tumor spheres (left) and mean
numbers of organoid ($\pm$ S.D.; right) per 10,000 cells. p values; unpaired Student's t-test (n = 3
per group, *P = 0.0158, **P = 0.0055).
(C) Clonogenicity assay results in control and *Med4*-depleted 4T07-TGL. The panel shows
representative images of dishes and microscopic images of colonies (left) and mean numbers of
colonies ($\pm$ S.D.; right) per 1,000 cells. p values; unpaired Student's t-test (n = 3 per group, ***P
= 0.0002, ****P < 0.0001).
(D) Soft agar assay results in the indicated 4T07-TGL cells. The panel shows representative
images of wells and microscopic images of colonies (left) and mean numbers of colonies ($\pm$ S.D.;
right) per 1,000 cells. p values; unpaired Student's t-test (n = 3 per group, **P = 0.0018, **P =
0.0027). (E) Transwell migration results of control and *Med4*-silenced 4T07-TGL cells. The
panel shows representative images of fields (left) and mean numbers of cells ($\pm$ S.D.; right) per
field. p values; unpaired Student's t-test (n = 3 per group, ***P = 0.0001, ****P < 0.0001).
1119
Supplementary Figure S5. *Med1* alters the transcriptional and epigenetic landscape
discordantly with *Med4* while *Med4*-loss driven transcriptional program is necessary for
metastasis.
(A) MA plot from RNA-seq analysis in D2A1 versus D2A1-d cells (Fold change $\geq$ 2, FDR <
0.05)
(B) GSEA plots of D2A1_metastatic and D2A1_dormant signatures in Med4-depleted cells as
compared with control cells.
(C) RT-qPCR analysis and immunoblots of *Med1* expression in D2A1-d-TGL cells expressing
MirE shRNA targeting either non-targeting control (NTC) or *Med1*; Mean values ($\pm$ S.D.) p
values; unpaired Student's t-test.
(D) Heatmap of differentially expressed genes (DEGs; Fold change $\geq$ 2, FDR < 0.05) in NTC
versus *Med1*-silenced D2A1-d-TGL cells.

(E) GSEA plots of D2A1_metastatic and D2A1_dormant signatures in *Med1*-silenced cells as
compared with control cells.
(F) The heatmap showing the $\log_2$ fold change (Log2FC) in the gene expression of selected
Mediator subunits when comparing sh*Med1* and *shMed4* to NTC, respectively. Each cell in the
heatmap contains a numerical value representing the Log_2FC, rounded to one decimal place.
(G) RT-qPCR analysis of *Med4* and *Med1* expression in D2A1-d-TGL cells expressing MirE
shRNA targeting either non-targeting control (NTC) or *Med4*; Mean values ($\pm$ S.D.) p values;
unpaired Student's t-test.
(H) RT-qPCR analysis of *Med4* and *Med1* expression in D2A1-d-TGL cells expressing MirE
shRNA targeting either non-targeting control (NTC) or *Med1*; Mean values ($\pm$ S.D.) p values;
unpaired Student's t-test.
(I) MNase digestion assay was performed in nuclei isolated from the indicated cells for
increasing incubation time. The panels show the representative image (upper), and the
quantification graph (lower).
(J) MA plot of ATAC-seq results comparing NTC versus *Med1*-silenced D2A1-d-TGL cells.
Differential accessibility ($\log_2$ fold change in reads per accessible region) plotted against the
mean reads per region.
(K) The dot plot represents the gene set enrichment analysis (GSEA) across four datasets
(Hallmarks, KEGG, Reactome, and GO). The indicated pathways were enriched in *Med4*-
silenced D2A1-d cells.
(L) GSEA plots show enrichment of HALLMARK_EMT, REACTOME_ECM organization, and
KEGG_Focal adhesion in NTC versus sh*Med4*.
1154
Supplementary Figure S6. The downregulation of *Med1* did not produce discordant
alterations in nuclear and chromatin structure, in contrast to the effects observed with
***Med4* silencing.**
(A) D2A1-d cells expressing either NTC or *Med1* shRNAs were stained with HP1α (red). The
panels show representative images (left) and fluorescence intensity profile across one cell (right).
(B) Cells were stained with LAMIN A/C (red) and counter-stained with DAPI (blue).
(C) D2A1-d cells expressing either NTC or Med4 shRNAs were subcellular fractionized into the
cytoplasm, nucleoplasm, and nuclear envelope (NE). Total lysates were subsequently analyzed
by immunoblotting using the indicated antibodies.
(D) D2A1-d cells expressing either NTC or *Med4* shRNAs were treated with 100 nM
Latrunculin B for 20 min. Cells were stained with HP1α (red) and F-actin (green). The panels
show representative images (left) and fluorescence intensity profile of HP1α across one cell
(right).
1168
Supplementary Figure S7. Med4 alters the repertoire of integrin subunits and mediates the
signaling pathway of integrin-mediated mechano-transduction.
(A) A GSEA plot of integrin complex (GO_CC; top). Heatmap shows the level of expression of
integrin subunit genes amongst the top differentially expressed genes in control versus *Med4*-
silenced D2A1-d-TGL cells (bottom).
(B) RT-qPCR results were indicated as relative mean expression of MRTF/SRF only target
genes, YAP only target genes and shared target genes in either control or Med4-silenced D2A1-d
cells. (C) RT-qPCR results were indicated as relative mean expression of MRTF/SRF only target
genes in D2A1-d cells expressing either NTC or Med4 shRNAs were co-transduced with either

empty vector (EV) or Flag-HA tagged MED4 cDNA. p values; unpaired Student's t-test (n = 3
per group, *P = 0.0449, **P = 0.0016, ***P = 0.001, ****P < 0.0001).
(D) D2A1-d-TGL cells transduced with either NTC or Med4 shRNAs were inoculated *i.v.* in
syngeneic mice. Lung sections were stained as indicated.
(E) D2A1-d-TGL cells were co-transduced with MirE shRNA targets for Med4 and MirE
shRNA targets the mechanosensitive transcription factors as indicated. Total lysates were
subjected to immunoblotting with the indicated antibodies.
1185

Figure 1 Bae et al.

Figure 2 Bae et al.

Figure 4 Bae et al.

Figure S2 Bae et al.

A Cell growth (D2A1-d)

B Tumorspheres (D2A1-d)

C Clonogenecity (D2A1-d)

D Soft agar (D2A1-d)

E Migration (D2A1-d)

F Invasion (D2A1-d)

A Cell growth (4T07)

B Tumorspheres (4T07)

C Clonogenecity (4T07)

D Soft agar (4T07)

E Migration (4T07)