CAnDI: a new tool to investigate conflict in homologous gene trees and explain convergent trait evolution

Holly M. Robertson[1,2], Joseph F. Walker[1,3*], and Edwige Moyroud[1,2*]

[1] The Sainsbury Laboratory, Bateman Street, CB2 1LR, University of Cambridge, UK

[2] Department of Genetics, Downing site, CB2 3EJ, University of Cambridge, UK

[3] Department of Biological Sciences, University of Illinois at Chicago, Chicago, IL, USA

* Corresponding authors

Email: em500@cam.ac.uk (EM) and jfw52@uic.edu (JFW)

Running Title: Gene conflict explains phenotypic convergence

Keywords: plant carnivory, trait evolution, conflict analyses, homolog trees, convergence

Abstract

Phenotypic convergence is found across the tree of life, and morphological similarities in distantly related species are often presumed to have evolved independently. However, clarifying the origins of traits has recently highlighted the complex nature of evolution, as apparent convergent features often share similar genetic foundations. Hence, the tree topology of genes that underlie such traits frequently conflicts with the overall history of species relationships. This conflict creates both a challenge for systematists and an exciting opportunity to investigate the rich, complex network of information that connects molecular trajectories with trait evolution. Here we present a novel conflict identification program named CAnDI (Conflict And Duplication Identifier), which enables the analysis of conflict in homologous gene trees rather than inferred orthologs. We demonstrate that the analysis of conflicts in homologous trees using CAnDI yields more comparisons than in ortholog trees in six datasets from across the eukaryotic tree of life. Using the carnivorous trap of Caryophyllales, a charismatic group of flowering plants, as a case study we demonstrate that analysing conflict on entire homolog trees can aid in inferring the genetic basis of trait evolution: by dissecting all gene relationships within homolog trees, we find genomic evidence that the molecular basis of the pleisiomorphic mucilaginous sticky trap was likely present in the ancestor of all carnivorous Caryophyllales. We also show that many genes whose evolutionary trajectories group species with similar trap devices code for proteins contributing to plant carnivory and identify a *LATERAL ORGAN BOUNDARY DOMAIN* transcription factor as a possible candidate for regulating sticky trap development.

Introduction

High-throughput sequencing technologies have allowed large multi-gene phylogenetic data (phylogenomic datasets) to emerge as the standard for inferring species relationships (Delsuc et al. 2005; Jeffroy et al. 2006). These datasets have become increasingly common and provide a wealth of information for investigating contentious regions in the tree of life (Dunn et al. 2008; Chiari et al. 2012; Wickett et al. 2014). Efforts to analyse the complete set of gene phylogenies in a genome, the "phylome" (Sicheritz-Pontén and Andersson 2001; Huerta-Cepas et al. 2007), are emerging as a new field.

In addition to species tree inference, these datasets also provide unprecedented opportunities to explore evolutionary history beyond species relationships. By comparing gene evolutionary histories to that of the species, it is possible to uncover patterns that reflect historical events resulting from ancient population structures (Parins-Fukuchi et al. 2021; Stull et al. 2021). In particular, these phylogenomic datasets may shed some light on the question of how similar adaptations emerge repeatedly during evolution.

Convergent evolution can be described as the evolution of phenotypic similarity (Doolittle 1994; Stern 2013; Arbuckle et al. 2014) but it can also be defined at a molecular level by considering the genetic basis of alike traits. Phenotypic convergence can reflect molecular convergence when parallel changes occur in genes underlying a given feature, but it can also arise from the recruitment of existing ancestral allelic

46   polymorphisms or the introduction of an allele by hybridisation (either ancient or recent)

(Stern 2013). Both recruitment and hybridisation are reflected in the phylome as gene

48   tree conflict, and investigating the conflicting evolutionary histories in genomic data can

provide valuable biological insights (Doyle 1992; Maddison 1997; Rokas et al. 2003).

50   Examples of this include ancestral populations of butterflies that underwent introgressive

hybridization to produce shared mimicry genes (Dasmahapatra et al. 2012), incomplete

52   or random lineage sorting resulting in unusually high levels of homoplasy in cacti

(Copetti et al. 2017), the distribution of karyotypic features in mammals (Robinson et al.

54   2008), and the horizontal transfer of large DNA regions that explains the phylogenetic

distribution of C4 photosynthesis in grasses (Dunning et al. 2019). An adaptive role for

56   genes in conflict was also established in tomato plants (Pease et al. 2016), and more

recently in Marsupials (Feng et al. 2022) and Diatoms (Roberts et al. 2023). Hence,

58   investigating conflicting patterns can illuminate the evolution of complex traits and

uncover molecular players that could contribute to novelty.

To capture the true extent of genome-wide patterns of discordance and all relevant

62   biological signals, it is essential to analyse as much of the phylome as possible; and

currently, the most cost-effective approach to accessing nearly complete phylomes is

64   through transcriptomes. However, current studies and methods for investigating conflict

often focus on analysing orthologs inferred from homolog trees. Focusing solely on the

66   inferred orthologs results in the loss of information and potentially interesting biological

signals. Paralogs (gene copies that arise within a population due to gene duplication)

68 must still, however, be accounted for, as they can mislead both inferred species trees and

gene tree conflict analyses.

A case study in phenotypic convergence: the carnivorous Caryophyllales

72 The flowering plant order Caryophyllales contains several emblematic species of

carnivorous plants, including the Southeast Asian pitcher plants, sundews, and the Venus

74 flytrap. While carnivory is the predicted ancestral condition in the clade of all

carnivorous Caryophyllales (Cameron et al. 2002; Heubl et al. 2006; Renner and Specht

76 2011), not all species in this clade are carnivorous. The Ancistrocladaceae family, for

example, is entirely non-carnivorous.

The clade of carnivorous Caryophyllales exhibits several instances of apparent

80 phenotypic convergence. Trap type and stem morphology are two particularly interesting

examples of major, complex adaptations involving many genes. Underlying both is the

82 placement of the clade of Ancistrocladaceae and Dioncophyllaceae as sister to the

monotypic family Drosophyllaceae, a relationship supported by phylogenomic datasets

84 composed of hundreds or even thousands of genes (Walker et al. 2017, 2018; Baker et al.

2022). This inferred relationship is of note as the families Ancistrocladaceae and

86 Drosophyllaceae exhibit striking morphological similarities not with each other but with

other, more distantly related, species in the clade.

Specifically, Drosophyllaceae, a family containing only one species, the Portuguese

90 sundew (*Drosophyllum lusitanicum*), is not sister to the genus *Drosera* (commonly

known as the sundew family) even though both have a sticky trap mechanism of

92   capturing prey and a herbaceous stem morphology. Moreover, there is also

Triphyophyllum peltatum, a liana in the Dioncophyllaceae, which is native to West

94   Africa and develops sundew like carnivorous sticky trap leaves under phosphorus

deficient conditions (Winkelmann et al. 2023).

This polyphyletic distribution of sticky traps within the carnivorous Caryophyllales

98   indicates that either true convergence has occurred or a precursor to the sticky trap

existed ~75 mya, with the extant phylogenetic distribution of these traps resulting from

100   incomplete lineage sorting, horizontal gene transfer, or introgression.

Another implication of Drosophyllaceae being sister to the clade of Ancistrocladaceae

102   and Dioncophyllaceae is that this relationship places the lianas of Ancistrocladaceae as

more closely related to the herbaceous Drosophyllaceae rather than the carnivorous

104   pitcher plant family Nepenthaceae, which is also a liana. This relationship would indicate

that either an ancestral population that existed approximately 67 mya exhibited a woody

106   stem morphology later lost for Drosophyllaceae or that such a lifestyle has been gained

twice independently (Cameron et al. 2002; Heubl et al. 2006; Walker et al. 2017).

Here we introduce CAnDI (Conflict And Duplication Identifier), a new program we

110   developed to analyse all gene relationships in an unbiased way and investigate conflict

across the phylome. This program dissects all relationships within homolog trees and

112   analyses them for conflict, concordance, or origination by duplication. This method

expands upon previous work in conflict detection, as it analyses and reports every

114   instance of conflict in a dataset and allows users to search for conflicts of interest. In

addition, it allows users to search for patterns of interest within homolog trees. We

116   demonstrate that CAnDI can identify evolutionary patterns that cannot be detected using

current approaches by applying it to a series of datasets from various closely and

118   distantly related groups. We further show that the use of orthologs accurately

demonstrates the amount of conflict expected from the whole phylome. Following this,

120   we conduct a more detailed case study in the carnivorous Caryophyllales, where our new

method enables us to provide genomic evidence that molecular variants shared by extant

122   species with a mucilaginous sticky trap were already present in the ancestor of all

carnivorous Caryophyllales. Using CAnDI, we also identify a set of genes that could

124   have participated in the emergence of the sticky trap morphology during the evolution of

the Caryophyllales. In particular, our analyses, combined with quantification of gene

126   expression levels in sundew tissues, single out the transcription factor *LBD4-like* as a

candidate regulator of sticky trap development.

Materials and Methods

130   **Comparison of whole homolog trees with ortholog trees for identifying**

gene tree conflict in six datasets

132   We compiled six datasets of species trees, ortholog trees, and gene families (homolog

trees) from published studies that broadly span the tree of life. We used the species

134   relationships and homolog trees inferred in the initial studies as a basis for each conflict

analysis. If available, we used rooted homolog trees from the original study. If the rooted

136   homolog trees were not available, we used the extract_clades.py script from (Yang et al.

2015) to extract them.

All of the datasets, both homologs and orthologs, were analysed using CAnDI to count

140   the total number of nodes assessed corresponding to each node in the species, where

each node was assessed as either conflicting with the species tree or concordant with the

142   species tree. Gene duplication nodes were not counted.

144   We extracted 12,172 rooted homologs from the unrooted homologs previously inferred

for the plant family Amaranthaceae (AMAR) by (Morales-Briones et al. 2021), using a

146   threshold of 40 ingroup taxa. The AMAR dataset did not have support values for

homolog trees. The 12,699 orthologs of the AMAR dataset with appropriate outgroup

148   representation for rooting were from the original study where they were extracted using

the monophyletic outgroup (MO) method (Yang and Smith, 2014). The Diatom (DIAT)

150   dataset from (Parks et al. 2018) had duplications extracted with a minimum of 10

ingroup taxa, resulting in 18,914 homolog trees. The support values for this dataset were

152   from a rapid bootstrap analysis; thus, we applied a cutoff of $\geq 70\%$. The 197 orthologs

from the original study, that were extracted using the Rooted Tree (RT) method (Yang

154   and Smith, 2014), were used for the ortholog comparison for the DIAT dataset. The

Legumes dataset (LEGU), consisting of 8,038 homolog trees from (Koenen et al. 2020),

156   had been extracted in the original study, and support values on the nodes were from a

rapid bootstrap analysis; thus, a cutoff of (≥ 70) was used. For the LEGU orthologs, we

158   used the ortholog trees extracted with the RT method from the original study. To extract

the homologs we used the same outgroup taxa, resulting in 1014 trees with outgroup

160   representation. The Hymenoptera dataset (HYMN) was originally published by (Johnson

et al. 2013) but was subsequently re-analysed by (Smith et al. 2015), who identified

162   5,863 homolog trees and calculated rapid bootstrap support; therefore, we used a support

cutoff of $\geq 70\%$ for the HYMN dataset. To generate orthologs for the HYMN dataset, we

164   used the RT method with a minimum cutoff of four taxa using the

prune_from_rooted_tree.py script of (Morales-Briones et al. 2021), resulting in 6519

166   orthologs. The 6902 homolog trees from the Ericales (ERIC) dataset from (Larson et al.

2020) were analysed using a support value of (SH-aLRT ≥ 80). Conflict analysis on

168   ortholog trees from the ERIC dataset was conducted using the 382 rooted ortholog trees

from the original study that had been extracted using the RT method. The details of the

170   carnivorous Caryophyllales (CARN) dataset, also used in this comparison, are described

below.

Investigating the relationship of conflict with trait distribution in the

174   **carnivorous Caryophyllales (CARN) dataset**

We used a minimum threshold of four ingroup taxa to extract 6,006 rooted homolog

176   trees from the unrooted homolog trees inferred by Walker et al. (Walker et al. 2017). We

term this the carnivorous Caryophyllales (CARN) dataset. Information regarding this

178   and other datasets as well as all supplementary material is available on DRYAD

(doi:10.5061/dryad.g4f4qrfzq). The 1237 orthologs we used for the CARN dataset were

180   those from the original study, which were extracted using one-to-one orthology with

100% gene occupancy required. All trees in the CARN dataset had SH-aLRT support

182 values; thus, we used a support cutoff of ≥80%. We then used CAnDI to identify all

homolog trees in the CARN dataset that contained relationships (bipartition) of interest

184 and the number of times the relationship of interest occurred in a single homolog tree.

We identified trees with the bipartition *Drosophyllum + Drosera* and all trees with the

186 bipartition *Drosophyllum + Ancistrocladus*. We also used CAnDI to identify all

bipartitions that contained only *Nepenthes ampullaria*, *Nepenthes alata,* and

188 *Drosophyllum lusitanicum*. The same procedure was performed to identify all instances

in homolog trees where Ancistrocladaceae (Ancistrocladus) was sister to Nepenthaceae

190 (*N. alata* and *N. ampullaria*). For each group of taxa investigated, we compared the

number of bipartitions that contained only those taxa against the number of instances

192 where an additional species was also included in the bipartition to eliminate any bias

caused by differences in gene recovery in different samples.

Investigating gene function in the CARN dataset

196 Annotations for the CARN dataset were inferred using the coding sequences of the

genome of *Arabidopsis thaliana* (TAIR10; Downloaded from Ensembl Plants on Jan

198 27[th], 2021). The best blast hit (e-value 1e-3) to the transcriptome of *Nepenthes alata*

from each homolog cluster was used to predict the function of each cluster. Using this

200 approach, 137 out of the 146 clusters supporting the monotypic family Drosophyllaceae

sister to the genus *Drosera* could be annotated (9 clusters failed to return a blast hit

202 against *A. thaliana*. These 137 annotations represented 132 distinct Arabidopsis genes.

Gene ontology analyses with PANTHER16.0 were performed on this list of 132 *A.*

204 *thaliana* genes using the PANTHER Overrepresentation Test (Released 20210224) and

the GO Ontology database (DOI: 10.5281/zenodo.5228828 Released 2021-08-18) and

206 either the "GO biological process complete" or the "GO molecular function complete"

dataset. The Fisher's exact test was performed with False Discovery Rate correction for

208 each analysis. A *Chitinase-like protein 1* [homolog to *Arabidopsis CTL1*; labelled

cluster2518 by Walker et al. (Walker et al. 2017)] was further investigated since this

210 gene is associated with carnivory. We spot-checked the annotation for this gene by

testing the blast annotation of other homologs within cluster2518 against the NCBI

212 RefSeq non-redundant protein database to ensure that this was not the result of a

spurious hit. All sequences from cluster2518 matched a Chitinase gene. We then

214 performed the reverse mapping feature with CAnDI ("-r") and mapped the coalescent-

based species tree topology for the CARN dataset to the cluster2518 homolog tree.

216 **Plant material**

Drosera admirabilis, *Drosera aliciae* and *Drosera coccicaulis* were purchased from

218 Wack's Wicked plants (UK), grown in ambient conditions (natural light, with

temperature kept between 18-25°C) and watered from the base with distilled water.

220 Tissues were harvested from emerging young leaves (unfurled primordia <1cm long,

trichomes absent or not yet able to secrete mucilage), mature leaves representing fully

222 developed sticky traps densely covered with trichomes (stalked glandular hairs) secreting

mucilage droplets (Fig. S5) and inflorescences and immediately frozen in liquid nitrogen.

224 For each species tissues were harvested from three distinct individuals to constitute three

biological replicates for the gene expression analysis. All tissue samples were stored at -

226 80°C until RNA extraction.

Gene expression analysis

Frozen tissues (from three distinct individuals for each species, constituting three

230 biological replicates) were ground to a fine powder using a mortar and pestle. RNA was

extracted using the Spectrum Plant Total RNA kit (Sigma-Aldrich) and retro-transcribed

232 using SuperscriptIII reverse transcriptase (Invitrogen) following the manufacturer's

instructions. Quantitative real time PCR was performed using the Luna Universal qPCR

234 Mastermix (NEBiolabs) and the Light Cycler 480 system (Roche). For each species and

tissue type, three independent biological replicates were used (i.e., RNA extracted from

236 three distinct individuals) and three technical replicates were performed for each

condition (i.e., three repeats of the qRT-PCR reaction assessing gene expression for a

238 given tissue type, of a given biological replicate for each species). *Actin* and *eIF4A*

homologs were used as reference genes as established by Arai and collaborators (Arai et

240 al. 2021). The primer efficiencies were 100% for *Actin*, 93% for *eIF4* and 109% for

LBD4-like homolog genes. Primers were designed to match gene regions conserved

242 between the three different species of *Drosera*. Primer sequences are given in the

Supplementary Spreadsheet 3. The gene expression was calculated relative to the

244 housekeeping gene actin and the common base method, which accounts for the measured

efficiency of each primer pair, was used to calculate relative expression levels (Ganger et

246 al. 2017). Welch's t-tests or ANOVA followed by Tukey's HSD post-hoc tests were

conducted in R [https://www.r-project.org, (R Core Team 2017)] and used to test for the

248 likelihood that LBD4-like average expression levels were equivalent across tissues.

Results and Discussion

CAnDI provides a procedure to dissect relationships and analyse conflict in whole homolog trees

Homolog trees contain all gene sequences with similarities resulting from their shared evolutionary history, such as those originating through gene duplication (paralogs) or speciation (orthologs). Gene duplication can occur due to local tandem duplications, chromosome duplications, or genome-wide duplication events (Lynch and Force 2000). Ancient whole genome duplications are particularly widespread in plants due to their tolerance for polyploidy (Clark and Donoghue 2018), meaning patterns of gene duplication and loss particularly characterize many plant gene families. However, gene duplication, by any mechanism, is also widespread in other eukaryotes, and gene family analyses are becoming more popular throughout evolutionary biology (Iñiguez and Hernández 2017; Copley 2020). Hence, homolog trees widen the available data for assessment and analysis of phylogenetic conflict compared to approaches that focus on only ortholog trees. With this and the widespread nature of gene duplications in the eukaryotes in mind, methods that detect and account for gene duplications on a gene family tree, allowing for the analysis of the whole homologous gene tree and not just a part of it, have widespread applicability.

Existing approaches for exploring conflict in large, multi-gene phylogenetic datasets

272 include counting gene concordance frequencies (Ane et al. 2006; Baum 2007; Smith et

al. 2015; Minh et al. 2020), dividing the tree into quartets (Pease et al. 2018; Zhou et al.

274 2020), and comparing gene tree likelihoods either through Bayes factors or gene-wise

likelihoods (Gatesy and Baker 2005; Brown and Thomson 2016; Shen et al. 2017;

276 Walker et al. 2018). However, these approaches tend to rely on inferred orthology. This

is due to the fact that any homolog tree containing a duplication could not be analysed in

278 entirety for conflict or concordance with a species tree, with the existence of multiple

tips from the same taxon confounding this process. Previous analyses have used various

280 orthology inference methods to mitigate such a loss of information (Zmasek and Eddy

2002; Hejnol et al. 2009; Yang and Smith 2014). However, all previous methods discard

282 at least some information about speciation events that occur post-duplication, meaning

many speciation nodes are not assessed for conflict even though they may be

284 informative.

286 To perform conflict analyses on entire homolog trees, we developed the novel program

CAnDI (Conflict And Duplication Identifier,

288 https://github.com/HollyMaeRobertson/CAnDI). This program allows for the analysis of

conflict in multiple ways. Firstly, as in PhyParts, (Smith et al., 2015) rooted homolog

290 trees and ortholog trees can be mapped to a rooted species tree, generating count data for

conflicts and concordances at each node in the species tree. In addition to this, a rooted

292 species tree can also be mapped to a rooted homolog tree, labelling each node in that

homolog tree as one of three mutually exclusive categories: a concordant node, a

conflicting node, or a duplication node. In either case, analysis of whole homolog trees requires the specification of each node in each tree as one of either a duplication or a speciation node. Duplication nodes are those where sister clades contain at least one overlapping taxon, resulting from gene duplication. All non-duplication nodes, inferred to have arisen by speciation, can be analysed for conflict or concordance with the species tree.

In order to conduct such an analysis, CAnDI decomposes the homolog tree into subtrees by the following process (Fig. 1): first, every child node of a duplication node is copied into a list of subtrees along with its downstream structure (Fig. 1A-B). Then, in any subtree itself containing a duplication, that duplication node is collapsed (Fig. 1C): every downstream node is removed, and the node is marked with a non-repeating list of all the taxa originally present. This ensures that each speciation node is only analysed once: otherwise, a speciation node downstream of (for example) two duplications would be analysed twice, once in the larger subtree originating at the first duplication and a second time in the subtree originating at the second duplication. Following the collapsing of duplications, any subtrees where the root node leads directly to a tip (i.e., the tree contains one taxon) and any subtrees where the root node is a duplication (all speciation nodes in these trees will be analysed in the two child subtrees of that duplication) are excluded (Fig. 1D), leaving a set of subtrees that form a nonoverlapping, complete description of every speciation node in the original homolog tree in a form that can be analysed using a bipartition method (Fig. 1E) (Salichos and Rokas 2013).

318   **Figure 1. Overview of the decomposition process of a whole homolog tree into**

subtrees. **A-D**: Steps required for subtree decomposition and subsequent processing. The

320   homolog tree, containing homologs from 5 distinct species (a, b, c, d and e), is first

decomposed into preliminary subtrees by splitting at each duplication node (red stars)

322   (**A**), with every child node of every duplication in the tree being added to the subtree list

(**B**). Any duplications located within each of these subtrees (except at the root node) are

324   collapsed and marked as not to be analysed (dashed branch as in the leftmost center

subtree, **C**). The outgroups of each subtree are also included but labelled as not to be

326 analysed to enable searching for missing taxa (dotted branches, **C**). Any subtrees whose

root node is a duplication or leads directly to a tip are finally excluded (**D**), leaving only

328 the informative subtrees with no overlap. **E**: The initial homolog tree and the final

subtrees to be analysed are shown side-by-side.

Bipartitions form the basis for conflict and concordance in the same manner as PhyParts

332 (Smith et al. 2015). As relationships arise from both speciation and duplication in the

homolog tree, the left side of the bipartition (ingroup subtree) is evaluated against all

334 bipartitions in the species tree (Fig. 1). The right side of the bipartition from the

homolog tree is all taxa found before the duplication and any taxa within the ingroup

336 subtree that are not in the left side of the bipartition. Any taxa in the species tree not

found in either the left or right side of the homolog tree bipartition are removed from the

338 species tree bipartitions to account for missing data. Concordance is identified as a

species tree bipartition that perfectly matches a homolog tree bipartition. Conflict occurs

340 when a homolog bipartition overlaps with a species tree bipartition but contains a non-

matching set of taxa. If a support value is specified, bipartitions in the homolog tree with

342 support below that value are not evaluated and are treated as uninformative.

344 CAnDI also identifies alternative conflict: under some circumstances, two non-nested

nodes in a single subtree can conflict with the same node at the species tree. A single

346 node from the homolog tree can also conflict with multiple species tree nodes. In the

former case, both conflicting nodes are reported, but only one instance of conflict is

348 counted. In the latter case, CAnDI reports the node at which conflict first occurs.

350   Finally, CAnDI allows users to search for a particular relationship of interest. This

searching procedure first creates subtrees from each homolog tree as above, then

352   searches for the bipartition corresponding to the conflict of interest in the subtrees. How

often a given homolog tree has the conflict and how many homolog trees have the same

354   conflict can both be reported.

356   Whole-homolog approaches and methods for analysing the phylome have been used for

other kinds of analysis than conflict [reviewed by (Smith and Hahn 2021)]. Entire

358   homolog trees have been shown to provide a holistic approach to species tree

reconciliation without the need to pre-infer orthology and are capable of recapitulating

360   the same species relationships as using only orthologs (Smith and Hahn 2021).

ASTRAL-Pro, a coalescent-based approach to infer species relationships, uses a quartet

362   tree decomposition strategy that separates speciation and duplication nodes (Zhang et al.

2020).

Furthermore, two approaches have been developed to decompose the homolog trees and

366   analyse them with respect to species relationships. Both methods use bipartitions as the

basis for the analysis. The TreeKO method (Marcet-Houben and Gabaldón 2011),

368   implemented in ETE3 (Huerta-Cepas et al. 2016), calculates the Robinson-Foulds (RF)

distance between a species tree and a homolog tree(s). The RF distance involves

370   decomposing each tree into bipartitions and counting the total number of bipartitions that

are unique to each tree. As this involves separating duplication from speciation nodes,

372   the tree dissection approach is similar to the one utilized here [e.g., the entire phylome

(Sicheritz-Pontén and Andersson 2001)]. The biggest difference is the TreeKO method

374   only provides a support score for the species tree. In contrast, the gene search approach

in CAnDI allows the user to identify specific relationships in gene trees. CAnDI is also

376   designed for detailed analysis of homolog trees, such as mapping species relationships

onto the homolog tree and providing the number of conflicting relationships with a given

378   species tree node.

380   The summary of gene trees from CAnDI is most similar to the second method, Phyparts

(Smith et al. 2015). Phyparts takes a homolog tree-centric approach that involves

382   decomposing the entire homolog tree into its respective bipartitions. The decomposition

allows the program to determine whether the homolog tree conflicts or not. However, it

384   bypasses the orthology detection procedure entirely by stipulating that each homolog

tree can only be concordant, conflicting, or both at a single time with the species tree.

386   Furthermore, as Phyparts is not designed for searching for specific conflicts, the

bipartitions focus on the deepest conflicting bipartition in the tree as a summary of

388   conflict. In short, this means that an individual homolog tree will not be summarized as

having multiple conflicts or concordances.

TreeKO and PhyParts allow users to input species trees, but only to map homolog trees

392   onto a species tree, rather than also allowing users to map the species tree onto a

homolog tree. The method we developed for this study and implemented in CAnDI

394   enables users to investigate all conflicts and concordances from a single homolog tree,

unlike previous methods that can only infer whether a tree is conflicting and not the

396   number of nodes at which an individual homolog tree is concordant or conflicting.

However, we stress that Phyparts and TreeKO function for their respective purposes, and

398   the differences outlined are not criticisms but explanations of how CAnDI has built upon

these authors' work.

Using homolog trees as a general basis for exploring conflict allows for

402   **more comparisons**

We applied CAnDI to datasets from six distinct groups of eukaryotes (hymenopterous

404   insects, unicellular diatoms, the carnivorous Caryophyllales and three other groups of

flowering plants) to compare conflict analysis based on homolog trees to that of inferred

406   ortholog trees. We found that using entire gene families allowed significantly more

relationships to inform the analysis (Fig.2). As the orthologs from the original study

408   were used, if the study required 100% taxon occupancy for all genes then each node was

analysed the same number of times.

Figure 2. Using entire gene families provides more relationships to inform analyses and uncover conflicts. The graphs correspond to datasets from six distinct groups: the Hymenoptera order of insects (HYMN); the diatoms, a lineage of microbial eukaryotes (DIAT) and four different plants groups: the carnivorous Caryophyllales (CARN), Fabaceae (LEGU), Ericales (ERIC) and Amaranthaceae (AMAR). The graphs show the ortholog bipartitions in the species tree, ordered along the x-axis according to the number of intersecting bipartitions (fewest to most). The filled in circles are the number of times the bipartition is analysed using homologs, and the open circles are the number of times the bipartition is analysed using orthologs.

422   In these analyses, we extracted rooted homolog trees using the same approach as Yang and colleagues (Yang et al. 2015). This approach required a cutoff for the minimum

424   number of ingroup taxa. However, recent work on non-time reversible models should allow users to extend even further rooting homologs without the need to extract

426   duplicates from the homolog tree. Homolog trees rooted in this way extend the method,

allowing users to analyse the entire phylome for any taxa with a sequenced genome.

Using inferred orthology is not an issue for species tree inference (Smith and Hahn

430   2021) or identifying processes associated with speciation patterns among genes (e.g.,

incomplete lineage sorting), where conflicts of interest are usually the most common

432   ones. However, this information loss can be costly when investigating topologies that

rarely arise, such as in this study, where we compare the relationship between trait

434   distribution (distinct trap or stem morphologies) and conflict. Moreover, relying on

inferred orthology is often insufficient for studies focusing on one gene family,

436   especially in cases where that gene family has undergone significant expansion in

multiple lineages. The approach we implemented in CAnDI provides a solution to

438   overcome those limitations.

Evolutionary conflict helps explain trait distribution in the carnivorous

442   **Caryophyllales**

To further investigate the potential value of CAnDI as an analytical tool, and with the

444   specific goal of identifying adaptive conflict, i.e. those specific conflicts which appear to

correspond with a phenotype of interest's distribution on the species tree, we turned to

446   the carnivorous Caryophylalles.

Existing consensus (Cameron et al. 2002; Heubl et al. 2006; Renner and Specht 2011)

448   states that the last common ancestor of the carnivorous Caryophyllales clade was

carnivorous. As a precaution, we revisited this hypothesis with increased taxon sampling

450   (Supplementary Methods) and our analysis predicted an ancestor of all the carnivorous

Caryophyllales with a sticky trap (Fig.S1). This is consistent with the findings of Renner

452   and Specht (Renner and Specht 2011), who used evidence from the different types of

glands present across the carnivorous Caryophyllales to argue that the snap traps of

454   *Dionaea muscipula* and *Aldrovanda vesiculosa* evolved from a simpler modified leaf

similar to the modern sticky trap. A similar evolutionary path could also account for the

456   emergence of *Nepenthes* pitcher traps. All carnivorous traps within the Caryophyllales

are modified leaves, but snap traps and pitfall traps both exhibit elaborated

458   morphologies, less reminiscent of a typical leaf than the sticky traps of genus *Drosera* or

the species *Drosophyllum lusitanicum* (monotypic family Drosophyllaceae) and

460   *Triphyophyllum peltatum* (family Dioncophyllaceae). However, this remains a claim

made on phenotypic evidence, which can be misleading (Corbett-Detig et al. 2020), with

462   no investigation of the evolutionary histories of the genes contributing to sticky trap

development. Regardless of the mechanisms behind the subsequent elaboration of the

464   pitfall and snap trap forms from an ancestral sticky trap, our results indicate that the

presence of a sticky trap in unrelated sundews is unlikely to be a true molecular

466   convergence (i.e., independent molecular basis) but could instead be the product of

introgression or incomplete lineage sorting. Thus, the presence of the sticky trap in

468   phylogenetically distant lineages is likely an example of hemiplasy, the topological

discordance between a gene tree and a species tree that can give the illusion of

470   homoplasy (Dasmahapatra et al. 2012).

472 To further investigate this hemiplasy, we generated rooted homolog trees from the

unrooted homolog trees of the carnivorous Caryophyllales dataset we already used in

474 Fig. 2 and originally published by Walker et al. in 2017. We then identified specific

conflicts with the established consensus species tree in these homologs using CAnDI

476 (Fig.3).

Figure 3. Patterns of conflict and ancestral traits are not found in all descendants

482 **("pleisiomorphic traits") in the carnivorous Caryophyllales.** Pie charts depict the

amount of gene tree conflict observed, with the blue, red and grey slices representing,

484 respectively, the proportion of homolog gene trees concordant, conflicting and

uninformative (SH-alrt ≤ 80 or missing taxon) at each node in the species tree. In case of

486 conflict, the proportions of gene trees supporting two main alternative topologies are

represented with a dotted or a striped red slice, respectively. The proportion of gene

488   trees supporting various alternative topologies is represented with a solid red slice. The

type of trap found in each carnivorous species, as well as the herbaceous or woody stem

490   morphology, is depicted with a symbol on the right of the phylogeny. Picture credit:

Drosophyllum lusitanicum (incidencematrix), *Ancistrocladus heyneanus* (Vinayaraj) and

492   *Drosera capensis* (NoahElhardt) via Wikimedia Commons.

494   We found that an order of magnitude more conflicting gene trees support *D. lusitanicum*

as sister to *Drosera binata* than the evolutionarily equivalent conflicting relationship

496   *Ancistrocladus robertsoniorum* sister to *D. binata* (Fig.4A). When we tested if we could

recover this pattern using the orthologs inferred by the original study, we only found one

498   ortholog tree placed *D. lusitanicum* and *D. binata* sister to one another. This pattern is

only evident in homolog trees, thereby emphasising the importance of analysing as much

500   of the phylome as possible when making biological inferences.

Figure 4. Distribution of conflicting gene trees according to the alternative species relationship they support, focusing on (A) carnivory and (B) life-history. The relationship being identified is drawn as a tree (top), while the number of conflicting gene trees supporting that relationship are shown below in homolog trees, clades within homolog trees and ortholog trees. The number of trees/subtrees for each alternative sister pair is shown in blue, followed by the number of well-supported ($80 \geq$ SH-alrt) trees/subtrees in grey. Homolog trees refers to the number of trees containing the relationship, while clades in homolog trees refer to every time the relationship occurred, including multiple occurrences in the same homolog tree. Conflicts that place species with shared stem and trap morphology as sister species are more common than evolutionarily equivalent conflicts that do not.

516   Using this dataset, we also investigated the apparent convergence of the woody vine life

        history phenotype in *Nepenthes* and *Ancistrocladus* (Fig.3 and Supplementary Fig. S2).

518   We found that, regardless of whether whole homologs or only ortholog trees were

        considered, more than twice as many conflicting genes placed the woody vines clades as

520   sister to one another, compared to the less similar herbaceous Drosophyllaceae as sister

        to Ancistrocladaceae (Fig.4B). Using homolog trees makes the pattern more easily

522   detectable and robust (trend based on 98 gene trees rather than only 16), again

        highlighting the importance of analysing as much data as possible (Fig.4).

        It should be noted that gene trees are prone to estimation errors. Our use of homolog

526   trees allows for the best-informed relationships, as taxon sampling is well known to

        increase the accuracy of phylogenetic estimation (Zwickl and Hillis 2002). Using

528   homolog trees, which are informed by the maximum number of samples possible, we

        should have increased accuracy compared to extracted orthologs. To account for errors

530   that may have arisen in the tree estimation procedure, we applied a support filter for the

        bipartitions, in this case requiring SH-alrt of at least 80. When the analysis was run with

532   and without support, our results identified a biased number of genes supporting the

        placement of sticky traps sister to each other and woody plants sister to each other

534   (Fig.4). This bias, even with the filters, helps indicate this is an actual biological signal

        and not an artifact of a lack of information. Although some gene tree errors are

536   inevitable, the functional association of the genes in this study provides further support

        for the signal and is an important consideration for any gene identification approach.

538   It is also worth noting that the family Dioncophyllaceae, which contains only three

known species, two non-carnivorous lianas and the likely-facultatively carnivorous liana

540   *Triphyophyllum peltatum*, is not represented in this dataset, but the literature strongly

supports Dioncophyllaceae sister to Ancistrocladaceae (Baker et al. 2022). An analysis

542   including *T. peltatum* and seeking conflicts that cluster *T. peltatum*, *Drosera* and

Drosophyllum could provide additional information regarding life history and trap

544   morphology evolution.

546   There are several reasons these homolog trees might display conflict. The probability of

true molecular convergence is extremely low, given the large number of genes and sites

548   displaying this pattern. Moreover, despite the short branch lengths separating *Drosera*

and *Drosophyllum* in many gene trees (Supplementary Fig. S3), recent introgression is

550   similarly unlikely, given the distant relationship between these species (*Drosera* and

Drosophyllum diverged ~75 mya). However, distinguishing ancient reticulation from

552   ancient incomplete lineage sorting would be challenging, given the age of the

relationships in question. Regardless of the mechanism of acquisition, it remains that

554   adaptive selection on these genes is likely responsible for the retention of specific allelic

variants post-acquisition.

Genes that support Drosophyllaceae as sister to *Drosera* tend to be

558   **associated with the carnivorous syndrome**

Genes whose evolutionary history conflicts with the species tree but places species with

560   similar morphologies (i.e., trap type) sister to each other could help identify some of the

molecular bases associated with the emergence of a given trait and likely to have

562   contributed to adaptation (Lee et al. 2011). We therefore performed gene ontology (GO)

term analyses on the gene trees that CAnDI identified as conflicting with the species

564   tree, which instead placed the sundews *Drosera* and *Drosophyllum* as sister to one

another. These analyses revealed that the functional categories associated with genes

566   exhibiting this conflicting phylogenetic relationship are often related to carnivory

(Tables 1 and 2). Cellular response to ozone (GO:0071457) and to reactive oxygen

568   species (GO:0034614), and superoxide dismutase activity (GO:0004784), emerged as

the most overrepresented biological process and molecular function, respectively.

570   Carnivorous species have been shown to use free radicals during the digestion process

(Chia et al. 2004), but these can also cause cellular damage. Hence, genes with

572   enzymatic activities capable of scavenging superoxide radicals (Bowler et al. 1991)

could play a key protective role in carnivorous plants. GO terms associated with nitrogen

574   utilization (i.e., GO:0019676, GO:0019740, GO:0004356) are also among the top

categories enriched in the set of genes exhibiting conflict and could relate to the

576   assimilation of nutrients following prey digestion. Our analyses also reveal enrichment

patterns for GO terms connected with photosynthesis and respiration (i.e., GO:0016168,

578   GO:0046933, GO:0009768). Producing the morphology and molecules that allow

carnivory (i.e., pigment to attract insects or traps to catch them) is energetically

580   demanding, and carnivory impacts both respiration and photosynthesis (Adamec et al.,

2021; Pavlovič, 2022). Indeed, trap development and activity are generally accompanied

582   by spatio-temporal activation of respiration and inactivation of photosynthesis.

Following this, photosynthesis can increase as a result of feeding (Pavlovič and

Saganová 2015). Our analysis also points toward other interesting categories. For

instance, genes connected to salicylic acid binding (GO:1901149) are overrepresented

among the genes in conflict with the species tree that group the two genera of sundews

as sisters to each other (Table 1). Salicylic acid (SA) and jasmonic acid (JA) are well-

known to work together to regulate plant immunity (Pieterse et al., 2012). Interestingly,

recent results revealed that JA signaling for plant defense has likely been co-opted by

some Caryophyllales to support carnivory (Pavlovič and Saganová, 2015; Bemm et al.,

2016; Pavlovič and Mithöfer, 2019; Pavlovič et al., 2024). Whether this indicates that the

crosstalk between SA and JA can possibly mediate carnivory-related function remains to

be established.

Table 1. Result of Gene Ontology analyses (Molecular Function) performed on genes in conflict with the species tree but supporting the *Drosera + Drosophyllum* relationship. The top ten most enriched categories from the 'Molecular Function' database are indicated. An extended version of this table, including subcategories, is provided in Supplementary Spreadsheet 1. FE, Fold Enrichment; FDR, False Discovery Rate.

GO Term	Description	Expected (Observed)	FE	Raw P-value	FDR
GO:0004784	Superoxide dismutase activity	0.04(3)	78.02	1.72e-05	1.74e-03
GO:0004356	glutamate-ammonia ligase activity	0.03(2)	69.35	6.24e-04	3.90e-02
GO:0030060	L-malate dehydrogenase activity	0.05(3)	62.41	2.95e-05	2.74e-03
GO:0003746	translation elongation factor activity	0.08(5)	61.19	5.73e-08	9.78e-06
GO:0016168	Chlorophyll binding	0.13(7)	52.01	2.94e-10	7.94e-08
GO:0008266	Poly(U) RNA binding	0.13(6)	46.23	1.06e-08	2.16e-06
GO:1901149	salicylic acid binding	0.13(5)	37.15	4.95e-07	8.03e-05

GO:0046933	proton-transporting ATP synthase activity, rotational mechanism	0.12(3)	26.01	2.88e-04	1.99e-02
GO:0016762	xyloglucan:xyloglucosyl transferase activity	0.15(3)	19.50	6.26e-04	3.84e-02
GO:0004601	Peroxidase activity	0.37(6)	16.42	2.78e-06	3.61e-04

602 **Table 2. Result of Gene Ontology analyses (Biological Process) performed on genes in conflict with the species tree but supporting the *Drosera* + *Drosophyllum* sister**
604 **relationship.** The main categories enriched from the 'Biological Process' database are indicated, along with selected related categories (italicized). An extended version of this
606 table, including all subcategories, is provided in Supplementary Spreadsheet 2. FE, Fold Enrichment; FDR, False Discovery Rate.

Go Term	Description	Expected (Observed)	FE	Raw P-value	FDR
GO:0071457	Cellular response to ozone	0.01(2)	>100	2.25e-04	1.09e-04
GO:0034614	*cellular response to reactive oxygen species*	*0.16(4)*	*25.22*	*2.92e-05*	*2.00e-03*
GO:0019676	Ammonia assimilation cycle	0.03(2)	69.35	6.24e-04	2.73e-02
GO:0006541	*glutamine metabolic process*	*0.08(3)*	*39.01*	*9.80e-05*	*5.25e-03*
GO:0019740	*nitrogen utilisation*	*0.03(2)*	*59.44*	*8.00e-04*	*3.40e-02*
GO:0009768	Photosynthesis, light harvesting in photosystem I	0.12(7)	60.68	1.017-10	2.24e-08
GO:0006542	Glutamine biosynthetic process	0.03(2)	59.44	8.00e-04	3.45e-02
GO:0071493	Cellular response to UV-B	0.03(2)	59.44	8.00e-04	3.42e-02
GO:0019430	Removal of superoxide radicals	0.05(3)	56.74	3.75e-05	2.40e-03
GO:0009645	Response to low light intensity stimulus	0.09(5)	54.75	9.17e-08	1.27e-05
GO:0030388	Fructose 1,6-bisphosphate metabolic process	0.06(3)	52.01	4.67e-05	2.86e-03
GO:0071329	Cellular response to sucrose stimulus	0.04(2)	46.23	1.21e-03	4.78e-02
GO:0046500	S-adenosylmethionine metabolic process	0.04(2)	46.23	1.21e-03	4.75e-02

Next, we conducted a detailed examination of the families to which these conflicting
610 gene trees belong. Plant carnivory is typified by a set of characters that enable them to attract, capture, digest and assimilate animal prey. This cluster of attributes is known as

612   the carnivorous syndrome. Within the genes exhibiting conflict, we found families

participating in each of the four facets of the carnivorous syndrome (Table 3 and

614   Supplementary Fig. S3); for example, three gene clusters code for enzymes participating

in anthocyanin synthesis. Red anthocyanin pigments colour the sticky trap of both

616   *Drosera* and *Drosophyllum*. Furthermore, while the purpose of anthocyanin

pigmentation in carnivorous species is debated, red-pigmented traps are widely found in

618   nature, and pigment accumulation is unmistakably associated with carnivory (Schefer

and Ruxton 2008; Jürgens et al. 2015; Gilbert et al. 2018). Sticky trap leaves and their

620   glandular emergences (tentacles) can bend toward stuck prey to secure their capture.

Such movements are generally the product of acid-growth processes (Poppinga et al.

622   2013). Acid-growth allows cell extension and growth following acidification of the cell

wall. Three gene clusters in the set encode Xyloglucan

624   endotransglucosylases/hydrolases, proteins capable of regulating cell wall extensibility

in response to wall acidification (Majda and Robert 2018). Enzymes found in trap

626   secretions and mediating prey digestion (i.e., chitinase) or protecting plant cells from

damages caused by free radicals used during the digestion process (i.e., peroxidases) are

628   also overrepresented among conflicting genes. Finally, genes encoding proteins

contributing to the utilisation of nitrogen (for growth and development) or to the use of

630   amino acids (alternative respiratory substrate to fuel the energy demand of carnivory)

from digested prey are also present (Table 3). As *Drosera* and *Drosophyllum* resemble

632   each other in terms of morphology and the action mechanism of their traps, these genes

with histories that conflict with the species tree represent candidate genes that could

634   have participated in the emergence of the sticky trap phenotype ~75 Mya, or that may

have mediated the parallel evolution of sticky traps in both groups of sundews.

Table 3. **Genes in conflict with species relationships that support the relationship**

638   *Drosera + Drosophyllum* **and encoding protein belonging to families with carnivory**

associated functions. The first column indicates the aspect of the carnivorous syndrome

640   associated with selected clusters.

	Cluster	Gene Family	Molecular Function	Role in Carnivory	Reference
Prey attraction	2986	Cinnamate-4-hydroxylase (C4H)	Catalyses the second step of the main phenylpropanoid pathway	Production of red anthocyanin pigment colouring the sticky trap hairs (tentacles) may lure insect prey to traps and/or may reduce the risk of a pollinator–prey conflict	(Schaefer and Ruxton 2008; Jürgens et al. 2015)
	82	Chalcone synthase (CHS)	Flavonoid biosynthesis		
	1332	Anthocyanidin synthase (ANS/LDOX)	Anthocyanin biosynthesis		
Retention by traps	1112	Xyloglucan endotransglucosylases/ hydrolase	Regulation of cell wall extensibility	Participate in the movement of sticky trap leaves and their glandular emergence (tentacles) that bend toward stuck prey due to the acid-growth process	(Poppinga et al. 2013; Majda and Robert 2018)
	810				
	3821				
Degradation of prey by digestive enzymes	2518	Chitinase	Degradation of the polysaccharide chitin	The digestive enzyme found in trap fluid or secretions Degradation of chitin from insect prey	(Matušíková et al. 2005; Renner and Specht 2013)
	2136	Formate dehydrogenase	Catalyse the oxidation of formate to carbon dioxide	Enzymes detected in the secretion of digestive fluids	(Schulze et al. 2012)
	160	Fructose-biphosphate aldolase	Plays a crucial role in glycolysis and gluconeogenesis		
	4896				
	1196	Peroxidase	Removal of H_2O_2, oxidation of toxic reductants	Oxidation of prey molecules Protect cells and enzymes from oxidative damage	(Fukushima et al. 2017; Ravee et al. 2018)
	677				
	435				
	3495	Superoxide dismutase	Scavenging of superoxide radicals	Reduce cellular damage mediated by oxygen radicals used during the prey digestion process	(Bowler et al. 1991; Chia et al. 2004; Schulze et
	999				
	3714				

					al. 2012)
Uptake of soluble compounds	4294 926 315	Dicarboxylate transporter1 Glutamine synthetase	Nitrogen Utilisation	Proteins of prey can be used as a source of nitrogen for growth and development	(Small et al. 1977; Scherzer et al. 2013)
	478 632	Malate dehydrogenase	Respiration	Prey-derived amino acids can be used as an alternative respiratory substrate to fuel the energy demand for prey digestion	(Fasbender et al. 2017)

An in-depth analysis of *Chitinase* (*CTL1*) reveals a history of duplication and conflict

We found a class I chitinase among the genes placing *Drosophyllum* and *Drosera* as

646 sister to one another (Table 3). The class I chitinase gene family plays an essential role in

plant evolutionary defense against fungal pathogens. Producing chitinase allows plants

648 to break down the chitinous fungal cell wall (Samac et al. 1990; Brogue et al. 1991).

Two subclasses of class I chitinases have been identified in the carnivorous

650 Caryophyllales, with subclass Ia participating in the pathogenesis response. In contrast,

members of subclass Ib help break down the chitinous exoskeleton of insects, supporting

652 carnivory (Renner and Specht 2012).

Although a commonly performed mapping approach is gene tree to species tree, the

656 ability to map a species tree onto a gene tree provides a powerful tool for single gene

family analyses. Species tree reconciliation algorithms such as NOTUNG (Chen et al.

658 2000) map duplications and other events to gene trees, but CAnDI does so in homolog

trees with a focus on conflict. By mapping the species tree onto the class I chitinase

homolog tree (Fig.5A), we found evidence for the previously characterised duplication

of class I chitinases at the base of the carnivorous Caryophyllales (Renner and Specht

2012). Although the carnivorous Caryophyllales are noted for their high propensity of

genome duplication (Walker et al. 2017; Palfalvi et al. 2020), *Drosophyllum* does not

have identifiable evidence for a whole genome duplication (Walker et al. 2017). As this

class I chitinase duplication is also conserved in *Drosophyllum*, it is likely not a

byproduct of whole genome duplication, though it remains possible that a combination

of whole genome duplication and local duplication assisted the spread of the chitinases

through this clade.

Figure 5. Gene duplication and conflict in the evolution of (A) the chitinase I gene (left) and (B) the *LBD4-like* gene (right). black = concordance, red = conflict, blue = duplication. The type of trap found in each carnivorous species is depicted with a symbol on the right of the homolog tree.

An LBD transcription factor may play a key role in the development of the carnivorous trap

We first discussed the chitinase gene family due to its critical role in the evolution of plant carnivory and the focus other researchers have placed on this family in distantly related clades of carnivorous plants (Fukushima et al. 2017; Lan et al. 2017; Saul et al. 2023). However, our interest was drawn to the bias in patterns of conflict, which indicates some form of selection driving the mucilaginous trap in the carnivorous Caryophyllales. This is also suggested by the GO term analysis (Table 3). Stochastic mapping on species trees has helped researchers infer the possible ancestral states that have led to carnivory (Renner and Specht 2011). We show that complementing this work with examinations of the evolutionary history of individual gene trees can provide further insight into the evolution of plant carnivory. Such analyses can be performed on genes already known to participate in the trait of interest, such as class I chitinases, and further test previous hypotheses. Importantly, CAnDI can also single out genes with conflicting histories that have not previously been linked to carnivory, providing useful starting points to investigate their potential role.

For instance, we found one gene coding for a transcription factor among the set of 146 conflicting genes that support a sister relationship between the sundews *Drosophyllum* and *Drosera* (Fig.5B). This transcription factor belongs to the *LATERAL ORGAN BOUNDARY DOMAIN* (*LBD*) gene family, and its closest relative in the model plant Arabidopsis is *LBD4*. It will hereafter be referred to as *LBD4-like*, since we have not

determined whether this caryophyllid gene and Arabidopsis *LBD4* form a true clade.

702   Members of the *LBD* family are regulators of plant development, controlling, for

instance, leaf morphogenesis (Iwakawa et al. 2002; Shuai et al. 2002). They also fulfil

704   important roles in regulating anthocyanin and nitrogen metabolism (Rubin et al. 2009).

Interestingly, LBD homologues have also been implicated in nyctinasty (Chen et al.

706   2012), a behaviour characterised by pronounced leaf movements in response to the

day/night cycle. In 2013, Poppinga and colleagues highlighted the need to investigate the

708   function of *LBD* genes in other plant families also producing motor organs, including the

snap trap of the carnivorous *Dionea*, in order to understand better the origins and

710   evolution of plant lateral motor organs (Poppinga et al. 2013). Regulators of sticky trap

formation should start being expressed early during leaf development in young

712   primordia that are yet to form the morphological features characteristic of the sticky trap.

To test whether the spatio-temporal expression patterns of sundews *LBD4-like* homologs

714   were compatible with a possible role in trap elaboration, we characterised its expression

in young and mature leaves of three species producing a sticky trap: *Drosera*

716   *admirabilis*, *Drosera aliciae* and *Drosera coccicaulis* (Fig.6A-C). As inflorescences

were available in the *D. coccicaulis* individuals, we also measured gene expression in

718   those tissues for comparison (Fig.6D).

Figure 6. The expression dynamics of *Drosera LBD4-like* genes are compatible with a role during trap formation. (A-D) Young leaf primordia (boxed region) and mature traps from *Drosera admirabilis* (A), *Drosera aliciae* (B) and *Drosera coccicaulis* (C). The inflorescence of *D. coccicaulis* is partially visible in (D), indicated by a white arrowhead. **(E-F)** Relative expression levels of *LBD4-like* homologs in young leaf primordia and mature sticky traps of *D. admirabilis* (E) and *D. aliciae* (F). * $p < 0.05$, ** $p < 0.01$, Welch's t-test. Individual dots represent distinct biological replicates (n = 3), and red lines indicate mean relative expression. (**G**) Relative expression levels of *LBD4-like* homolog in young leaf primordia, mature sticky traps and inflorescence of *D.*

coccicaulis. Red lines indicate mean relative expression. Letters a, b, c, indicate the

734 results of a post hoc Tukey's HSD test, with group means associated with different

letters being significantly different from one another ($p < 0.01$).

In all three species, we found that the *LBD4-like* gene was expressed in developing traps

738 and that its expression decreases as development progresses. On average, transcript

levels were two to ten times higher in young leaf primordia compared to mature ones

740 (Fig.6E-G). In *D. coccicaulis, LBD4-like* homolog expression was significantly weaker

in reproductive tissues (inflorescence), as transcripts levels were more than eight times

742 higher in emerging leaf primordia and almost 4 times as high in mature traps (Fig.6G).

Taken together, our results indicate that the expression dynamic of *LBD4-like* genes is

744 compatible with a possible role in trap development in *Drosera*. Hence, *LBD4-like* is a

promising candidate for functional investigation as it could be involved in shaping the

746 unique morphology of sticky trap leaves and/or contribute to their ability to produce

pigments or assimilate nitrogen following prey capture.

Conclusion

750 We developed the new program CAnDI to systematically investigate patterns of gene

tree conflict across the whole phylome, specifically enabling the unbiased analysis of

752 conflict in whole homologous trees. We demonstrated that CAnDI allows for far more

comparisons between gene trees and the species tree to be made using phylogenomic

754 datasets, regardless of the identity of the organisms considered, in six previously

published phylogenomic datasets. We then focused on the carnivorous Caryophyllales,

756 showing that both carnivory and woodiness were traits associated with gene tree conflict

by identifying 146 genes supporting a sister relationship between the two genera of

758 sundews, directly conflicting with the established phylogeny of the carnivorous

Caryophyllales. This supports existing evidence that phenotypic convergence and/or new

760 adaptations frequently rely on recruiting ancestral genetic variation (Pease et al. 2016;

Schweizer et al. 2019). We uncovered a clear link between those 146 genes with

762 conflicting histories and all aspects of the carnivory syndrome. We also discovered that

an *LBD4* homolog is a possible candidate regulator of sticky trap development. Hence,

764 CAnDI enables focused examination of genes previously known to participate in a trait

of interest but also provides promising means to identify novel candidates for functional

766 investigation. This illustrates how examining patterns of gene tree conflict can further

our understanding of how evolution shapes developmental processes and generates

768 phenotypic convergences.

Supplementary Material

The Supplementary Material file contains:

772   **Fig. S1** Ancestral state estimation of the trap morphology in the carnivorous
            Caryophyllales supports a sticky trap ancestor.
774   **Fig. S2** Ancestral state estimation of the growth habit in the carnivorous
            Caryophyllales.
776   **Fig. S3** Graphical representation of all the trees listed in Table 3 of the main
            text.
778   **Fig. S4** Example of examining the relationship between a homolog tree and a
            species tree.
780   **Fig. S5** Picture of young leaf and mature trap tissues used for qRT-PCR assay.

Supplementary Method – Ancestral state reconstruction

782   **Supporting References**

784   Other supporting materials for this manuscript include the following:

786   **Supplementary Spreadsheet 1.** Outcome of Gene Ontology Analysis
         (Molecular Function)

788   **Supplementary Spreadsheet 2.** Outcome of Gene Ontology Analysis
         (Biological Process)

790   **Supplementary Spreadsheet 3.** List of Primers used in this study

Acknowledgments and Funding

We would like to thank Yuxi Liang for drawing the three silhouettes representing the

794   different trap systems used in Fig.3-5 and Fig. S1-S2. Stephen Smith for helpful

discussions on the best procedure for identifying duplication events. Andrew Alverson,

796   Erik Koenen, Diego Morales-Briones, and Drew Larson for assistance with dataset

acquisition and formatting.

798   We are also grateful to Drew Larson, Chris Whitewoods and the members of the

Moyroud lab and Walker lab for useful discussions and comments on earlier versions of

800   this manuscript. Our research is supported by the Gatsby Charitable Foundation, the

Isaac Newton Trust/Wellcome Trust ISSF, Murray Edwards College and the SBS-DTP

802   PhD program at the University of Cambridge and start-up funding from the University of

Illinois at Chicago.

References

806   Adamec, L., Matušíková, I. and Pavlovič, A., 2021. Recent ecophysiological, biochemical and evolutional insights into plant carnivory. Ann. Bot., 128:241-259.

808   Ane C., Larget B., Baum D.A., Smith S.D., Rokas A. 2006. Bayesian Estimation of Concordance among Gene Trees. Mol. Biol. Evol. 24:412–426.

810   Arai N., Ohno Y., Jumyo S., Hamaji Y., Ohyama T. 2021. Organ-specific expression and epigenetic traits of genes encoding digestive enzymes in the lance-leaf sundew (Drosera adelae).
812   J. Exp. Bot. 72:1946–1961.

    Arbuckle K., Bennett C.M., Speed M.P. 2014. A simple measure of the strength of convergent
814   evolution. Methods Ecol. Evol. 5:685–693.

    Baker W.J., Bailey P., Barber V., Barker A., Bellot S., Bishop D., Botigué L.R., Brewer G.,
816   Carruthers T., Clarkson J.J., Cook J., Cowan R.S., Dodsworth S., Epitawalage N.,

    Françoso E., Gallego B., Johnson M.G., Kim J.T., Leempoel K., Maurin O., Mcginnie C.,
818   Pokorny L., Roy S., Stone M., Toledo E., Wickett N.J., Zuntini A.R., Eiserhardt W.L., Kersey P.J., Leitch I.J., Forest F. 2022. A Comprehensive Phylogenomic Platform for Exploring the
820   Angiosperm Tree of Life. Syst. Biol. 71:301–319.

    Baum D.A. 2007. Concordance trees, concordance factors, and the exploration of reticulate
822   genealogy. TAXON. 56:417–426.

    Bemm F, Becker D, Larisch C, et al. . 2016. Venus flytrap carnivorous lifestyle builds on
824   herbivore defence strategies. Genome Research 26, 812–825.

    Bettisworth B., Stamatakis A. 2021. Root Digger: a root placement program for phylogenetic
826   trees. BMC Bioinformatics. 22:225.

    Bowler C., Slooten L., Vendenbranden S., De Rycke R., Botterman J., Sybesma C., Van Montagu
828   marc, Inzé D. 1991. Manganese superoxide dismutase can reduce cellular damage mediated by oxygen radicals in transgenic plants. EMBO J. 10:1723–1732.

830   Brogue K., Chet I., Holliday M., Cressman R., Biddle P., Knowlton S., Mauvais C.J., Broglie R. 1991. Transgenic Plants with Enhanced Resistance to the Fungal Pathogen *Rhizoctonia solani*.
832   Science. 254:1194–1197.

    Brown J.M., Thomson R.C. 2016. Bayes factors unmask highly variable information content,
834   bias, and extreme influence in phylogenomic analyses. Syst. Biol.:syw101.

    Brown J.W., Walker J.F., Smith S.A. 2017. Phyx: phylogenetic tools for unix. Bioinformatics.
33:1886–1888.

Cameron K.M., Wurdack K.J., Jobson R.W. 2002. Molecular evidence for the common origin of snap-traps among carnivorous plants. Am. J. Bot. 89:1503–1509.

Chen J., Moreau C., Liu Y., Kawaguchi M., Hofer J., Ellis N., Chen R. 2012. Conserved genetic determinant of motor organ identity in Medicago truncatula and related legumes. Proc. Natl. Acad. Sci. 109:11723–11728.

Chen K., Durand D., Farach-Colton M. 2000. Notung: dating gene duplications using gene family trees. Proc. Fourth Annu. Int. Conf. Comput. Mol. Biol.:96–106.

Chia T.F., Aung H.H., Osipov A.N., Goh N.K., Chia L.S. 2004. Carnivorous pitcher plant uses free radicals in the digestion of prey. Redox Rep. 9:255–261.

Chiari Y., Cahais V., Galtier N., Delsuc F. 2012. Phylogenomic analyses support the position of turtles as the sister group of birds and crocodiles (Archosauria). BMC Biol. 10:65.

Clark J.W., Donoghue P.C.J. 2018. Whole-Genome Duplication and Plant Macroevolution. Trends Plant Sci. 23:933–945.

Copetti D., Búrquez A., Bustamante E., Charboneau J.L.M., Childs K.L., Eguiarte L.E., Lee S., Liu T.L., McMahon M.M., Whiteman N.K., Wing R.A., Wojciechowski M.F., Sanderson M.J.
2017. Extensive gene tree discordance and hemiplasy shaped the genomes of North American columnar cacti. Proc. Natl. Acad. Sci. 114:12003–12008.

Copley S.D. 2020. Evolution of new enzymes by gene duplication and divergence. FEBS J. 287:1262–1283.

Corbett-Detig R.B., Russell S.L., Nielsen R., Losos J. 2020. Phenotypic Convergence Is Not Mirrored at the Protein Level in a Lizard Adaptive Radiation. Mol. Biol. Evol. 37:1604–1614.

Dasmahapatra K.K., Walters J.R., Briscoe A.D., Davey J.W., Whibley A., Nadeau N.J., Zimin A.V., Hughes D.S.T., Ferguson L.C., Martin S.H., Salazar C., Lewis J.J., Adler S., Ahn S.-J.,
Baker D.A., Baxter S.W., Chamberlain N.L., Chauhan R., Counterman B.A., Dalmay T., Gilbert L.E., Gordon K., Heckel D.G., Hines H.M., Hoff K.J., Holland P.W.H., Jacquin-Joly E.,
Jiggins F.M., Jones R.T., Kapan D.D., Kersey P., Lamas G., Lawson D., Mapleson D., Maroja L.S., Martin A., Moxon S., Palmer W.J., Papa R., Papanicolaou A., Pauchet Y., Ray D.A.,
Rosser N., Salzberg S.L., Supple M.A., Surridge A., Tenger-Trolander A., Vogel H., Wilkinson P.A., Wilson D., Yorke J.A., Yuan F., Balmuth A.L., Eland C., Gharbi K., Thomson M., Gibbs
R.A., Han Y., Jayaseelan J.C., Kovar C., Mathew T., Muzny D.M., Ongeri F., Pu L.-L., Qu J., Thornton R.L., Worley K.C., Wu Y.-Q., Linares M., Blaxter M.L., ffrench-Constant R.H., Joron
M., Kronforst M.R., Mullen S.P., Reed R.D., Scherer S.E., Richards S., Mallet J., Owen McMillan W., Jiggins C.D., The Heliconius Genome Consortium. 2012. Butterfly genome
reveals promiscuous exchange of mimicry adaptations among species. Nature. 487:94–98.

Delsuc F., Brinkmann H., Philippe H. 2005. Phylogenomics and the reconstruction of the tree of
872 life. Nat. Rev. Genet. 6:361–375.

Doolittle R.F. 1994. Convergent evolution: the need to be explicit. Trends Biochem. Sci. 19:15–18.

Doyle J.J. 1992. Gene Trees and Species Trees: Molecular Systematics as One-Character Taxonomy. Syst. Bot. 17:144–163.

Dunn C.W., Hejnol A., Matus D.Q., Pang K., Browne W.E., Smith S.A., Seaver E., Rouse G.W., Obst M., Edgecombe G.D., Sørensen M.V., Haddock S.H.D., Schmidt-Rhaesa A., Okusu A., Kristensen R.M., Wheeler W.C., Martindale M.Q., Giribet G. 2008. Broad phylogenomic sampling improves resolution of the animal tree of life. Nature. 452:745– 749.

Dunning L.T., Olofsson J.K., Parisod C., Choudhury R.R., Moreno-Villena J.J., Yang Y., Dionora J., Quick W.P., Park M., Bennetzen J.L., Besnard G., Nosil P., Osborne C.P., Christin P.-A. 2019. Lateral transfers of large DNA fragments spread functional genes among grasses. Proc. Natl. Acad. Sci. 116:4416–4425.

Fasbender L., Maurer D., Kreuzwieser J., Kreuzer I., Schulze W.X., Kruse J., Becker D., Alfarraj S., Hedrich R., Werner C., Rennenberg H. 2017. The carnivorous Venus flytrap uses prey-derived amino acid carbon to fuel respiration. New Phytol. 214:597–606.

Feng S., Bai M., Rivas-González I., Li C., Liu S., Tong Y., Yang H., Chen G., Xie D., Sears K.E., Franco L.M., Gaitan-Espitia J.D., Nespolo R.F., Johnson W.E., Yang H., Brandies P.A., Hogg C.J., Belov K., Renfree M.B., Helgen K.M., Boomsma J.J., Schierup M.H., Zhang G. 2022. Incomplete lineage sorting and phenotypic evolution in marsupials. Cell. 185:1646-1660.e18.

Foster C.S.P., Sauquet H., van der Merwe M., McPherson H., Rossetto M., Ho S.Y.W. 2017. Evaluating the Impact of Genomic Data and Priors on Bayesian Estimates of the Angiosperm Evolutionary Timescale. Syst. Biol. 66:338–351.

Fukushima K., Fang X., Alvarez-Ponce D., Cai H., Carretero-Paulet L., Chen C., Chang T.-H., Farr K.M., Fujita T., Hiwatashi Y., Hoshi Y., Imai T., Kasahara M., Librado P., Mao L., Mori H., Nishiyama T., Nozawa M., Pálfalvi G., Pollard S.T., Rozas J., Sánchez-Gracia A., Sankoff D., Shibata T.F., Shigenobu S., Sumikawa N., Uzawa T., Xie M., Zheng C., Pollock D.D., Albert V.A., Li S., Hasebe M. 2017. Genome of the pitcher plant Cephalotus reveals genetic changes associated with carnivory. Nat. Ecol. Evol. 1:1–9.

Ganger M.T., Dietz G.D., Ewing S.J. 2017. A common base method for analysis of qPCR data and the application of simple blocking in qPCR experiments. BMC Bioinformatics. 18:534.

Gatesy J., Baker R.H. 2005. Hidden Likelihood Support in Genomic Data: Can Forty-Five Wrongs Make a Right? Syst. Biol. 54:483–492.

Gilbert K.J., Nitta J.H., Talavera G., Pierce N.E., 2018. Keeping an eye on coloration: ecological correlates of the evolution of pitcher traits in the genus *Nepenthes* (Caryophyllales), Biol. J. Linn. Soc. 123:321–337

Hejnol A., Obst M., Stamatakis A., Ott M., Rouse G.W., Edgecombe G.D., Martinez P., Baguñà J., Bailly X., Jondelius U., Wiens M., Müller W.E.G., Seaver E., Wheeler W.C., Martindale
M.Q., Giribet G., Dunn C.W. 2009. Assessing the root of bilaterian animals with scalable phylogenomic methods. Proc. R. Soc. B Biol. Sci. 276:4261–4270.

Heubl G., Bringmann G., Meimberg H. 2006. Molecular Phylogeny and Character Evolution of Carnivorous Plant Families in Caryophyllales - Revisited. Plant Biol.:821–830.

Huerta-Cepas J., Dopazo H., Dopazo J., Gabaldón T. 2007. The human phylome. Genome Biol. 8:R109.

Huerta-Cepas J., Serra F., Bork P. 2016. ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic Data. Mol. Biol. Evol. 33:1635–1638.

Iñiguez L.P., Hernández G. 2017. The Evolutionary Relationship between Alternative Splicing and Gene Duplication. Front. Genet. 8.

Iwakawa H., Ueno Y., Semiarti E., Onouchi H., Kojima S., Tsukaya H., Hasebe M., Soma T., Ikezaki M., Machida C., Machida Y. 2002. The ASYMMETRIC LEAVES2 Gene of Arabidopsis
thaliana, Required for Formation of a Symmetric Flat Leaf Lamina, Encodes a Member of a Novel Family of Proteins Characterized by Cysteine Repeats and a Leucine Zipper. Plant Cell
Physiol. 43:467–478.

Jeffroy O., Brinkmann H., Delsuc F., Philippe H. 2006. Phylogenomics: the beginning of
926 incongruence? Trends Genet. 22:225–231.

Johnson B.R., Borowiec M.L., Chiu J.C., Lee E.K., Atallah J., Ward P.S. 2013. Phylogenomics
928 Resolves Evolutionary Relationships among Ants, Bees, and Wasps. Curr. Biol. 23:2058– 2062.

Jürgens A., Witt T., Sciligo A., El-Sayed A.M. 2015. The effect of trap colour and trap-flower
930 distance on prey and pollinator capture in carnivorous Drosera species. Funct. Ecol. 29:1026– 1037.

932 Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods. 14:587–
934 589.

Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
936 Improvements in Performance and Usability. Mol. Biol. Evol. 30:772–780.

Koenen E.J.M., Ojeda D.I., Steeves R., Migliore J., Bakker F.T., Wieringa J.J., Kidner C., Hardy
O.J., Pennington R.T., Bruneau A., Hughes C.E. 2020. Large-scale genomic sequence data resolve the deepest divergences in the legume phylogeny and support a nearsimultaneous
evolutionary origin of all six subfamilies. New Phytol. 225:1355–1369.

Kozlov A.M., Darriba D., Flouri T., Morel B., Stamatakis A. 2019. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics. 35:4453–4455.

Lan T., Renner T., Ibarra-Laclette E., Farr K.M., Chang T.-H., Cervantes-Pérez S.A., Zheng C., Sankoff D., Tang H., Purbojati R.W., Putra A., Drautz-Moses D.I., Schuster S.C., Herrera-
Estrella L., Albert V.A. 2017. Long-read sequencing uncovers the adaptive topography of a carnivorous plant genome. Proc. Natl. Acad. Sci. 114:E4435–E4441.

Larson D.A., Walker J.F., Vargas O.M., Smith S.A. 2020. A consensus phylogenomic approach highlights paleopolyploid and rapid radiation in the history of Ericales. Am. J. Bot. 107:773–789.

Lee E.K., Cibrian-Jaramillo A., Kolokotronis S.-O., Katari M.S., Stamatakis A., Ott M., Chiu J.C., Little D.P., Stevenson D.W., McCombie W.R., Martienssen R.A., Coruzzi G., DeSalle R.
2011. A Functional Phylogenomic View of the Seed Plants. PLOS Genet. 7:e1002411.

       Lynch M., Force A. 2000. The Probability of Duplicate Gene Preservation by
954   Subfunctionalization. Genetics. 154:459–473.

       Maddison W.P. 1997. Gene Trees in Species Trees. Syst. Biol. 46:523–536.

956   Maddison W.P., Maddison D.R. 2021. Mesquite: a modular system for evolutionary analysis.

       Majda M., Robert S. 2018. The Role of Auxin in Cell Wall Expansion. Int. J. Mol. Sci. 19:951.

958   Marcet-Houben M., Gabaldón T. 2011. TreeKO: a duplication-aware algorithm for the comparison of phylogenetic trees. Nucleic Acids Res. 39:e66.

960   Matušíková I., Salaj J., Moravčíková J., Mlynárová L., Nap J.-P., Libantová J. 2005. Tentacles of in vitro-grown round-leaf sundew (Drosera rotundifoliaL.) show induction of chitinase activity
962   upon mimicking the presence of prey. Planta. 222:1020–1027.

       Minh B.Q., Hahn M.W., Lanfear R. 2020. New Methods to Calculate Concordance Factors for
964   Phylogenomic Datasets. Mol. Biol. Evol. 37:2727–2733.

       Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington S.F.,
966   Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2021. Disentangling Sources of Gene Tree Discordance in Phylogenomic Data Sets: Testing Ancient Hybridizations in Amaranthaceae s.l.
968   Syst. Biol. 70:219–235.

       Naser-Khdour S., Minh B.Q., Lanfear R. 2021. Assessing Confidence in Root Placement on
970   Phylogenies: An Empirical Study Using Non-Reversible Models for Mammals. 2020.07.31.230144.

972 Nguyen L.-T., Schmidt H.A., von Haeseler A., Minh B.Q. 2015. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol. Biol. Evol.
974 32:268–274.

Palfalvi G., Hackl T., Terhoeven N., Shibata T.F., Nishiyama T., Ankenbrand M., Becker D.,
Förster F., Freund M., Iosip A., Kreuzer I., Saul F., Kamida C., Fukushima K., Shigenobu S., Tamada Y., Adamec L., Hoshi Y., Ueda K., Winkelmann T., Fuchs J., Schubert I., Schwacke R.,
Al-Rasheid K., Schultz J., Hasebe M., Hedrich R. 2020. Genomes of the Venus Flytrap and Close Relatives Unveil the Roots of Plant Carnivory. Curr. Biol. 30:2312-2320.e5.

Parins-Fukuchi C., Stull G.W., Smith S.A. 2021. Phylogenomic conflict coincides with rapid morphological innovation. Proc. Natl. Acad. Sci. 118:e2023058118.

Parks M.B., Nakov T., Ruck E.C., Wickett N.J., Alverson A.J. 2018. Phylogenomics reveals an extensive history of genome duplication in diatoms (Bacillariophyta). Am. J. Bot. 105:330-347.

Pavlovič, A., 2022. Photosynthesis in Carnivorous Plants: From Genes to Gas Exchange of Green Hunters. Critical Reviews in Plant Sciences, 41:305-320.

Pavlovič, A., Koller, J., Vrobel, O., Chamrád, I., Lenobel, R. and Tarkowski, P., 2024. Is the co-option of jasmonate signalling for botanical carnivory a universal trait for all carnivorous plants?
Journal of Experimental Botany, 75:334-349.

Pavlovič A, Mithöfer A. 2019. Jasmonate signalling in carnivorous plants: copycat of plant
990 defence mechanisms. Journal of Experimental Botany 70:3379–3389.

Pavlovič A., Saganová M. 2015. A novel insight into the cost–benefit model for the evolution of
992 botanical carnivory. Ann. Bot. 115:1075–1092.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling
distinguishes lack of support from conflicting support in the green plant tree of life. Am. J. Bot. 105:385–403.

Pease J.B., Haak D.C., Hahn M.W., Moyle L.C. 2016. Phylogenomics Reveals Three Sources of Adaptive Variation during a Rapid Radiation. PLOS Biol. 14:e1002379.

Pieterse, C. M. J., Van der Does, D., Zamioudis, C., Leon-Reyes, A., and Van Wees, S. C. M. (2012). Hormonal modulation of plant immunity. Annu. Rev. Cell Dev. Biol. 28, 489–521.

Poppinga S., Masselter T., Speck T. 2013. Faster than their prey: New insights into the rapid movements of active carnivorous plants traps. BioEssays. 35:649–657.

R Core Team. 2017. R: A language and environment for statistical computing. R Found. Stat. Comput. Vienna Austria.

Ravee R., Salleh F. 'Imadi M., Goh H.-H. 2018. Discovery of digestive enzymes in carnivorous plants with focus on proteases. PeerJ. 6:e4914.

Renner T., Specht C.D. 2011. A Sticky Situation: Assessing Adaptations for Plant Carnivory in the Caryophyllales by Means of Stochastic Character Mapping. Int. J. Plant Sci. 172:889–901.

Renner T., Specht C.D. 2012. Molecular and Functional Evolution of Class I Chitinases for Plant Carnivory in the Caryophyllales. Mol. Biol. Evol. 29:2971–2985.

Renner T., Specht C.D. 2013. Inside the trap: gland morphologies, digestive enzymes, and the evolution of plant carnivory in the Caryophyllales. Curr. Opin. Plant Biol. 16:436–442.

Roberts W.R., Ruck E.C., Downey K.M., Pinseel E., Alverson A.J. 2023. Resolving marine–freshwater transitions by diatoms through a fog of discordant gene trees. :2022.08.12.503770.

Robinson T.J., Ruiz-Herrera A., Avise J.C. 2008. Hemiplasy and homoplasy in the karyotypic phylogenies of mammals. Proc. Natl. Acad. Sci. 105:14477–14481.

Rokas A., Williams B.L., King N., Carroll S.B. 2003. Genome-scale approaches to resolving incongruence in molecular phylogenies. Nature. 425:798–804.

Rubin G., Tohge T., Matsuda F., Saito K., Scheible W.-R. 2009. Members of the LBD Family of Transcription Factors Repress Anthocyanin Synthesis and Affect Additional Nitrogen Responses
in Arabidopsis. Plant Cell. 21:3567–3584.

    Samac D.A., Hironaka C.M., Yallaly P.E., Shah D.M. 1990. Isolation and Characterization of
1022   themGenes Encoding Basic and Acidic Chitinase in Arabidopsis thaliana. Plant Physiol. 93:907–914.

1024   Saul, F., Scharmann, M., Wakatake, T., Rajaraman S., Marques A., Freund M., Bringmann G., Channon L., Becker D., Carroll E., Low Y.W., Lindqvist C., Gilbert K.J., Renner T., Masuda S.,
1026   Richter M., Vogg G., Shirasu K., Michael T.P., Hedrich R., Albert V.A., Fukushima K., 2023. Subgenome dominance shapes novel gene evolution in the decaploid pitcher plant Nepenthes
1028   gracilis. Nat. Plants 9:2000–2015.

    Schaefer H.M., Ruxton G.D. 2008. Fatal attraction: carnivorous plants roll out the red carpet to
1030   lure insects. Biol. Lett. 4:153–155.

    Scherzer S., Krol E., Kreuzer I., Kruse J., Karl F., von Rüden M., Escalante-Perez M., Müller T.,
1032   Rennenberg H., Al-Rasheid K.A.S., Neher E., Hedrich R. 2013. The Dionaea muscipula Ammonium Channel DmAMT1 Provides NH4+ Uptake Associated with Venus Flytrap's Prey
1034   Digestion. Curr. Biol. 23:1649–1657.

    Schulze W.X., Sanggaard K.W., Kreuzer I., Knudsen A.D., Bemm F., Thøgersen I.B., Bräutigam
1036   A., Thomsen L.R., Schliesky S., Dyrlund T.F., Escalante-Perez M., Becker D., Schultz J.,

1038 Karring H., Weber A., Højrup P., Hedrich R., Enghild J.J. 2012. The Protein Composition of the Digestive Fluid from the Venus Flytrap Sheds Light on Prey Digestion Mechanisms. Mol. Cell. Proteomics. 11:1306–1319.

1040 Schweizer M., Warmuth V., Alaei Kakhki N., Aliabadian M., Förschler M., Shirihai H., Suh A., Burri R. 2019. Parallel plumage colour evolution and introgressive hybridization in wheatears. J.
1042 Evol. Biol. 32:100–110.

Shen X.-X., Hittinger C.T., Rokas A. 2017. Contentious relationships in phylogenomic studies
1044 can be driven by a handful of genes. Nat. Ecol. Evol. 1:1–10.

Shuai B., Reynaga-Peña C.G., Springer P.S. 2002. The Lateral Organ Boundaries Gene Defines a
1046 Novel, Plant-Specific Gene Family. Plant Physiol. 129:747–761.

Sicheritz-Pontén T., Andersson S.G.E. 2001. A phylogenomic approach to microbial evolution.
1048 Nucleic Acids Res. 29:545–552.

Small G.C., Onraët A., Grierson D.S., Reynolds G. 1977. Studies on Insect-Free Growth,
1050 Development and Nitrate-Assimilating Enzymes of Drosera Aliciae Hamet. New Phytol. 79:127–133.

1052 Smith M.L., Hahn M.W. 2021. New Approaches for Inferring Phylogenies in the Presence of Paralogs. Trends Genet. 37:174–187.

1054 Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals conflict, concordance, and gene duplications with examples from animals and plants. BMC Evol.
1056 Biol. 15:150.

Smith S.A., Walker J.F. 2019. PyPHLAWD: A python tool for phylogenetic dataset construction.
1058 Methods Ecol. Evol. 10:104–108.

Stern D.L. 2013. The genetic causes of convergent evolution. Nat. Rev. Genet. 14:751–764.

Stull G.W., Qu X.-J., Parins-Fukuchi C., Yang Y.-Y., Yang J.-B., Yang Z.-Y., Hu Y., Ma H., Soltis P.S., Soltis D.E., Li D.-Z., Smith S.A., Yi T.-S. 2021. Gene duplications and phylogenomic
conflict underlie major pulses of phenotypic evolution in gymnosperms. Nat. Plants. 7:1015–1025.

Walker J.F., Yang Y., Feng T., Timoneda A., Mikenas J., Hutchison V., Edwards C., Wang N., Ahluwalia S., Olivieri J., Walker-Hale N., Majure L.C., Puente R., Kadereit G., Lauterbach M.,
Eggli U., Flores-Olvera H., Ochoterena H., Brockington S.F., Moore M.J., Smith S.A. 2018. From cacti to carnivores: Improved phylotranscriptomic sampling and hierarchical homology
inference provide further insight into the evolution of Caryophyllales. Am. J. Bot. 105:446–462.

Walker J.F., Yang Y., Moore M.J., Mikenas J., Timoneda A., Brockington S.F., Smith S.A. 2017. Widespread paleopolyploidy, gene tree conflict, and recalcitrant relationships among the carnivorous Caryophyllales. Am. J. Bot. 104:858–867.

Wickett N.J., Mirarab S., Nguyen N., Warnow T., Carpenter E., Matasci N., Ayyampalayam S., Barker M.S., Burleigh J.G., Gitzendanner M.A., Ruhfel B.R., Wafula E., Der J.P., Graham S.W., 1074 Mathews S., Melkonian M., Soltis D.E., Soltis P.S., Miles N.W., Rothfels C.J., Pokorny L., Shaw A.J., DeGironimo L., Stevenson D.W., Surek B., Villarreal J.C., Roure B., Philippe H., 1076 dePamphilis C.W., Chen T., Deyholos M.K., Baucom R.S., Kutchan T.M., Augustin M.M., Wang J., Zhang Y., Tian Z., Yan Z., Wu X., Sun X., Wong G.K.-S., Leebens-Mack J. 2014. 1078 Phylotranscriptomic analysis of the origin and early diversification of land plants. Proc. Natl. Acad. Sci. 111:E4859–E4868.

1080 Winkelmann T., Bringmann G., Herwig A., Hedrich R. 2023. Carnivory on demand: phosphorus deficiency induces glandular leaves in the African liana Triphyophyllum peltatum. New Phytol. 1082 239:1140–1152.

Yang Y., Moore M.J., Brockington S.F., Soltis D.E., Wong G.K.-S., Carpenter E.J., Zhang Y., 1084 Chen L., Yan Z., Xie Y., Sage R.F., Covshoff S., Hibberd J.M., Nelson M.N., Smith S.A. 2015. Dissecting Molecular Evolution in the Highly Diverse Plant Clade Caryophyllales Using 1086 Transcriptome Sequencing. Mol. Biol. Evol. 32:2001–2014.

Yang Y., Smith S.A. 2014. Orthology Inference in Nonmodel Organisms Using Transcriptomes 1088 and Low-Coverage Genomes: Improving Accuracy and Matrix Occupancy for Phylogenomics. Mol. Biol. Evol. 31:3081–3092.

1090 Yao G., Jin J.-J., Li H.-T., Yang J.-B., Mandala V.S., Croley M., Mostow R., Douglas N.A., Chase M.W., Christenhusz M.J.M., Soltis D.E., Soltis P.S., Smith S.A., Brockington S.F., Moore 1092 M.J., Yi T.-S., Li D.-Z. 2019. Plastid phylogenomic insights into the evolution of Caryophyllales. Mol. Phylogenet. Evol. 134:74–86.

1094 Yap V.B., Speed T. 2005. Rooting a phylogenetic tree with nonreversible substitution models. BMC Evol. Biol. 5:2.

1096 Zhang R., Fu X., Zhao C., Cheng J., Liao H., Wang P., Yao X., Duan X., Yuan Y., Xu G., Kramer E.M., Shan H., Kong H. 2020. Identification of the Key Regulatory Genes Involved in Elaborate 1098 Petal Development and Specialized Character Formation in Nigella damascena (Ranunculaceae). Plant Cell. 32:3095–3112.

1100 Zhou X., Lutteropp S., Czech L., Stamatakis A., Looz M.V., Rokas A. 2020. Quartet-Based Computations of Internode Certainty Provide Robust Measures of Phylogenetic 1102 Incongruence. Syst. Biol. 69:308–324.

Zmasek C.M., Eddy S.R. 2002. RIO: Analyzing proteomes by automated phylogenomics 1104 using resampled inference of orthologs. BMC Bioinformatics. 3:14.

Zwickl D.J., Hillis D.M. 2002. Increased Taxon Sampling Greatly Reduces Phylogenetic Error.

1106   Syst. Biol. 51:588–598.

Figure Captions

Figure 1. Overview of the decomposition process of a whole homolog tree into subtrees. A-D: Steps required for subtree decomposition and subsequent processing. The homolog tree, containing homologs from 5 distinct species (a, b, c, d and e), is first decomposed into preliminary subtrees by splitting at each duplication node (red stars) (**A**), with every child node of every duplication in the tree being added to the subtree list (**B**). Any duplications located within each of these subtrees (except at the root node) are collapsed and marked as not to be analysed (dashed branch as in the leftmost center subtree, **C**). The outgroups of each subtree are also included but labelled as not to be analysed to enable searching for missing taxa (dotted branches, **C**). Any subtrees whose root node is a duplication or leads directly to a tip are finally excluded (**D**), leaving only the informative subtrees with no overlap. **E**: The initial homolog tree and the final subtrees to be analysed are shown side-by-side.

Figure 2. Using entire gene families provides more relationships to inform analyses and uncover conflicts. The graphs correspond to datasets from six distinct groups: the Hymenoptera order of insects (HYMN); the diatoms, a lineage of microbial eukaryotes (DIAT) and four different plants groups: the carnivorous Caryophyllales (CARN), Fabaceae (LEGU), Ericales (ERIC) and Amaranthaceae (AMAR). The graphs show the ortholog bipartitions in the species tree, ordered along the x-axis according to the number of intersecting bipartitions (fewest to most). The filled in circles are the number

of times the bipartition is analysed using homologs, and the open circles are the number

of times the bipartition is analysed using orthologs.

Figure 3. Patterns of conflict and ancestral traits are not found in all descendants ("pleisiomorphic traits") in the carnivorous Caryophyllales. Pie charts depict the

amount of gene tree conflict observed, with the blue, red and grey slices representing, respectively, the proportion of homolog gene trees concordant, conflicting and

uninformative (SH-alrt ≤ 80 or missing taxon) at each node in the species tree. In case of conflict, the proportions of gene trees supporting two main alternative topologies are

represented with a dotted or a striped red slice, respectively. The proportion of gene trees supporting various alternative topologies is represented with a solid red slice. The

type of trap found in each carnivorous species, as well as the herbaceous or woody stem morphology, is depicted with a symbol on the right of the phylogeny. Picture credit:

Drosophyllum lusitanicum (incidencematrix), *Ancistrocladus heyneanus* (Vinayaraj) and *Drosera capensis* (NoahElhardt) via Wikimedia Commons.

Figure 4. Distribution of conflicting gene trees according to the alternative species relationship they support, focusing on (A) carnivory and (B) life-history. The

relationship being identified is drawn as a tree (top), while the number of conflicting gene trees supporting that relationship are shown below in homolog trees, clades within

homolog trees and ortholog trees. The number of trees/subtrees for each alternative sister pair is shown in blue, followed by the number of well-supported ($80 \geq$ SH-alrt)

trees/subtrees in grey. Homolog trees refers to the number of trees containing the relationship, while clades in homolog trees refer to every time the relationship occurred,

1152 including multiple occurrences in the same homolog tree. Conflicts that place species

 with shared stem and trap morphology as sister species are more common than

1154 evolutionarily equivalent conflicts that do not.

1156 **Figure 5. Gene duplication and conflict in the evolution of (A) the chitinase I gene**

 (left) and (B) the *LBD4-like* gene (right). black = concordance, red = conflict, blue =

1158 duplication. The type of trap found in each carnivorous species is depicted with a symbol

 on the right of the homolog tree.

1160

 Figure 6. The expression dynamics of *Drosera LBD4-like* genes are compatible with

1162 **a role during trap formation. (A-D)** Young leaf primordia (boxed region) and mature

 traps from *Drosera admirabilis* (A), *Drosera aliciae* (B) and *Drosera coccicaulis* (C).

1164 The inflorescence of *D. coccicaulis* is partially visible in (D), indicated by a white

 arrowhead. **(E-F)** Relative expression levels of *LBD4-like* homologs in young leaf

1166 primordia and mature sticky traps of *D. admirabilis* (E) and *D. aliciae* (F). * $p < 0.05$, **

 $p < 0.01$, Welch's t-test. Individual dots represent distinct biological replicates (n = 3),

1168 and red lines indicate mean relative expression. (**G**) Relative expression levels of *LBD4-*

 like homolog in young leaf primordia, mature sticky traps and inflorescence of *D.*

1170 *coccicaulis*. Red lines indicate mean relative expression. Letters a, b, c, indicate the

 results of a post hoc Tukey's HSD test, with group means associated with different

1172 letters being significantly different from one another ($p < 0.01$).