Maintenance of cell wall remodeling and vesicle production are connected in *Mycobacterium tuberculosis*

Vivian Salgueiro[1]*, Jorge Bertol[2]*, Claude Gutierrez[3], Jose L. Serrano-Mestre[1], Noelia Ferrer-Luzon[2], Lucía Vázquez-Iniesta[1], Ainhoa Palacios[4], Laia Pasquina-Lemonche[5], Akbar Espaillat[6], Laura Lerma[1], Brian Weinrick[7], Jose L. Lavin[8], Felix Elortza[4], Mikel Azkalgorta[4], Alicia Prieto[9], Pilar Buendía-Nacarino[10], José L. Luque-García[10], Olivier Neyrolles[3], Felipe Cava[6], Jamie K. Hobbs[5], Joaquín Sanz[2]# and Rafael Prados-Rosales[1]#

[1]Department of Preventive Medicine and Public Health and Microbiology, Universidad Autónoma de Madrid, Madrid, Spain;[2]Institute for Bio-computation and Physics of Complex Systems BIFI, Department of Theoretical Physics, University of Zaragoza, Zaragoza, Spain; [3]Institut de Pharmacologie et de Biologie Structurale, IPBS, Université de Toulouse, CNRS, UPS, Toulouse, France; [4]CIC bioGUNE, , Basque Research and Technology Alliance (BRTA), Bizkaia Science and Technology Park, Derio, Spain; [5]Department of Physics and Astronomy, University of Sheffield, Sheffield, UK; [6]Department of Molecular Biology and Laboratory for Molecular Infection Medicine Sweden, Umeå Centre for Microbial Research, SciLifeLab, Umeå University, Umeå, Sweden; [7]Trudeau Institute, Saranac Lake, NY, USA; [8]Bioinformatics Unit, Neiker-Tecnalia, Derio, Spain; [9]Department of Microbial & Plan Biotechnology, Centro de Investigaciones Biológicas Margarita Salas, Spanish National Research Council (CSIC),Madrid, Spain; [10]Department of Analytical Chemistry, Universidad Complutense de Madrid, Madrid, Spain.

Key words: extracellular vesicles, *Mycobacterium tuberculosis*, vesiculogenesis, peptidoglycan, membrane, permeability.

#To whom correspondence should be addressed:

Dr. Rafael Prados-Rosales– rafael.prados@uam.es

Dr. Joaquín Sanz- jsanz@bifi.es

*Equal contribution

Running title: *Mycobacterium* vesicle biogenesis mechanism.

Abstract

Pathogenic and nonpathogenic mycobacteria secrete extracellular vesicles (EVs) under various conditions. EVs produced by *Mycobacterium tuberculosis* (*Mtb*) have raised significant interest for their potential in cell communication, nutrient acquisition, and immune evasion. However, the relevance of vesicle secretion during tuberculosis infection remains unknown due to the limited understanding of mycobacterial vesicle biogenesis. We have previously shown that a transposon mutant in the LCP-related gene *virR* (*virR^mut^*) manifested a strong attenuated phenotype during experimental macrophage and murine infections, concomitant to enhanced vesicle release. In this study, we aimed to understand the role of VirR in the vesicle production process in *Mtb*. We employ genetic, transcriptional, proteomics, ultrastructural and biochemical methods to investigate the underlying processes explaining the enhanced vesiculogenesis phenomenon observed in the *virR^mut^*. Our results establish that VirR is critical to sustain proper cell permeability via regulation of cell envelope remodeling possibly through the interaction with similar cell envelope proteins, which control the link between peptidoglycan and arabinogalactan. These findings advance our understanding of mycobacterial extracellular vesicle biogenesis and suggest that these set of proteins could be attractive targets for therapeutic intervention.

Introduction

Mycobacterium tuberculosis (*Mtb*) is one of the most successful human pathogens with the capacity to persist within the host establishing long-lasting infections. *Mtb* possesses multiple highly evolved mechanisms to manipulate host cellular machinery and evade the host immune system. Similar to other intracellular pathogens, *Mtb* releases proteins and lipids, that interfere with host cellular processes, via canonical or specialized secretion systems (Majlessi et al., 2015). How mycobacteria confined within phagosomes delivers virulence factors to other cellular compartments and/or into the extracellular milieu to affect the physiology of neighboring cells is currently a matter of investigation. Understanding the physiology of this phenomenon during the infection process might be critical to develop alternative therapies against tuberculosis (TB).

Like most forms of life, *Mtb* uses an alternative antigen release process based on extracellular vesicles (EVs) (Prados-Rosales et al., 2011). Mycobacterial EVs (MEVs) are membrane walled spheres of 60-300 nm in diameter, released by live bacteria *in vitro* and *in vivo*. MEVs contain iron scavenging molecules, immunologically active complex lipids and lipoproteins, and classical virulence factors such as the *Mycobacterium ulcerans* toxin mycolactone (Lee et al., 2015; Marsollier et al., 2007; Prados-Rosales et al., 2014b, 2011). Several studies have clearly established a role of MEVs in immunomodulation and shown that MEVs deliver factors that impair macrophage effector functions, inhibit T cell activation, and modify the response of host cells to infection (Athman et al., 2015, 2017; Prados-Rosales et al., 2011, 2014b, 2014a; Rath et al., 2013). MEVs have also shown potential as biomarkers of *Mtb* infection (Schirmer et al., 2022; Ziegenbalg et al., 2013). Although the importance of MEVs has been recognized, hardly anything is known regarding both how such vesicles are exported across the mycobacterial cell wall and the molecular mechanisms underlying vesicle formation in mycobacteria. To date, several modulators of MEVs have been described. Our group identified two conditions leading to stimulation of vesicle production, namely: iron starvation (Prados-Rosales et al., 2014b) and the deletion of *virR* (Rv0431, vesiculogenesis and immune response Regulator) (Rath et al., 2013). In addition, it was shown that the Pst/SenX3-RegX3 signal transduction system regulates MEVs production independent from VirR (White et al., 2018), suggesting an alternative mechanism of vesicle production. We have recently demonstrated that both iron starvation and deletion of VirR in *Mtb*, two hypervesiculating conditions, trigger the induction of the isoniazid-induced gene operon *iniBAC* (Gupta et al., 2023). We determined that both IniA and IniC are dynamin-like proteins (DLP) presumably assisting MEV release through the fusion of the cell membrane (Gupta et al., 2023; Wang et al., 2019). Understanding how defects

in VirR lead to hypervesiculation through DLPs is important to provide mechanistic
explanation for the vesiculation process in mycobacteria.

Several lines of evidence support the notion that VirR might be involved in
maintaining cell wall integrity: (i) First, sequence analysis of the VirR protein indicates
that it contains a conserved LytR_C domain (**Fig S1**), which is usually found in
combination with another domain named LytR-Cps2A-Psr (LCP), and proteins carrying
both domains have been associated with the remodeling of the cell wall in Gram-positive
bacteria (Hubscher et al., 2008). *Mtb* has six genes encoding for LCP proteins, three with
both LCP and LytR_C domains (Rv3267, Rv3484 and Rv0822), two with a solo LytR_C
domain (VirR and Rv2700) and one protein with a solo LCP domain (Rv3840) (**Fig S1**).
The LCP family includes enzymes that transfer glycopolymers from membrane-linked
precursors to peptidoglycan (PG) or cell envelope proteins and are central to cell
envelope integrity (Kawai et al., 2011). In mycobacteria its was demonstrated that some
members of the (LCP) family of proteins are responsible for the linkage between
arabinogalactan (AG) and PG (Grzegorzewicz et al., 2016; Harrison et al., 2016).
Rv3267, Rv3484 and Rv0822 appear to have overlapping functions in cell wall assembly,
Rv3267 being the primary ligase (Grzegorzewicz et al., 2016). Interestingly, while
knocking down the gene encoding for the main AG-PG ligase (cg0847, *lcpA*) in
Corynebacterium glutamicum leads to the release of outer membrane material to the
extracellular medium (Baumgart et al., 2016), this phenotype is not observed in any
knockout strain in the *Mtb* orthologs (Grzegorzewicz et al., 2016), probably reflecting the
ability of the mutants to negatively regulate the biosynthesis of cell wall constituents in
response to a decrease in ligase activity; (ii) Second, we and others have determined
the increased susceptibility of the transposon *virR* mutant (*virR^{mut}*) with respect to the
WT (Ballister et al., 2019; Rath et al., 2013), and the mutant in the *virR* ortholog, Rv2700
(cell envelope integrity, *cei*) to cell wall targeting drugs such as meropenem (Ballister et
al., 2019). Based on these results, we hypothesize that LytR_C solo domain proteins
including VirR and Cei Rv2700 have critical roles in the maintenance of cell wall
homeostasis and their absence provokes cell wall alterations leading to enhanced
vesiculation.

In this work we show that the absence of VirR leads to ultrastructural changes in
the cell envelope as observed by cryo-electron microscopy (cryo-EM). These changes
are compatible with enhanced permeability and an enlargement of the PG layer, as
measured by high resolution atomic force microscopy (AFM). We demonstrate that VirR
interacts with canonical LCP proteins and propose a model for such interaction. These
results indicate that VirR is a central scaffold at the cell envelope remodeling process

assisting the PG-AG linking process and that this function is important for vesicle
production in *Mtb*.

164
165
166   **Results**
167

High EV production linked to the lack of VirR is associated with an altered cell
envelope in *Mtb* in the absence of cell lysis. While Gram-negative bacteria can
release outer membrane vesicles directly into the extracellular environment from their
outermost compartment, Gram-positive bacteria and mycobacteria have thick cell
envelopes surrounding the cytoplasmic membrane that acts as a permeability barrier.
The mycobacterial cell envelope is composed of PG covalently linked to AG, which in
turn is decorated with exceptionally long-chain mycolic acids (MA). These MA, together
with intercalating glycolipids, form the unique mycobacterial outer membrane (OM)
(Kalscheuer et al., 2019). Previous analysis of mycobacterial vesicle associated lipids
showed predominantly polar lipids, consistent with the cytoplasmic membrane being the
likely origin of the vesicles (Prados-Rosales et al., 2011). This is also true in EV produced
by Fe-limited *Mtb* whose lipid content consists mainly of polar lipids and the lipidic
siderophore, mycobactin (Prados-Rosales et al., 2014b). Given that these vesicles must
traffic from their point of origin in the plasma membrane through the cell envelope, we
hypothesized that local remodeling of the cell envelope is necessary for EV release, and
that this process may be exacerbated in a *virR* mutant (*virR^{mut}*). To test this, we examined
the cell envelope structure of *virR^{mut}* grown in minimal medium (MM), as well as wild type
(WT) and *virR^{mut}* complemented (*virR^{mut}*-C) strains (Rath et al., 2013) (**Fig 1**). Using cryo-
electron microscopy (cryo-EM) on whole cells, which allows the study of mycobacterial
cell surface-associated compartments in a close-to-native state (Sani et al., 2010), we
could easily discern three layers comprising the mycobacterial cell envelope: i) the outer
membrane (OM); ii) an intermediate layer composed of two sublayers (L1 and L2,
previously assigned to the mycolate-PG-AG (mAGP) network (Sani et al., 2010)); and iii)
the cytoplasmic membrane (CM) (**Fig 1**). We observed a significant increase in the
thickness of the overall cell envelope (OM-CM) in the *virR^{mut}* strain compared to the WT.
This difference was largely explained by the expansion of the CM-L1 layer, which
appeared also more granulated, suggesting cell wall alterations closer to the cell
membrane and indicating aberrant production of either AG, PG or both.
Complementation of *virR* restored normal cell envelope structure (**Fig 1A-C**).

197   In Gram-negative bacteria increased vesiculation has been observed in
198   conditions associated with either loss or maintenance of periplasm integrity (McMillan

and Kuehn, 2021). Moreover, a novel mechanism of vesicle production common to both Gram-negative and Gram-positive bacteria has recently been described, in which a cell explosion phenomenon driven by bacteriophages leads to the formation of EVs in culture when bacteria are under genotoxic stress (Nagakubo et al., 2021). Consequently, we tested whether the enhanced vesiculation described in $virR^{mut}$ (Rath et al., 2013) could be linked to cell lysis events. To do this, we analyzed the presence of cytoplasmic IdeR, a transcription factor which is not present in EVs and is not detected in secreted proteins preparations, in culture supernatants of Mtb strains, assuming that its detection could be linked to a loss of envelope integrity (Gupta et al., 2023). We first confirmed that the $virR^{mut}$ strain manifested an enhanced vesiculation by measuring EVs using nanoparticle tracking analysis (NTA) (**Fig S2A**). We did not detect differences in size distribution of isolated EVs between strains. The same supernatants and cell lysates were used to detect the presence of IdeR by immunoblot using a specific polyclonal antibody (Pandey and Rodriguez, 2014). As a control for normal release of an extracellular protein, we used a monoclonal antibody against Ag85b (Prados-Rosales et al., 2017). We could not detect IdeR in the supernatant of WT and $virR^{mut}$ strain, but a faint band corresponding to IdeR could be observed in the SN of the $virR$ complemented strain ($virR^{mut}$-Comp). These results suggest that the absence of $virR$ does not lead to cell lysis but its overexpression may compromise cell envelope integrity (**Fig S2B**). We validated this approach by serially diluting WT, $virR^{mut}$ and $virR^{mut}$-Comp strains on 7H10 solid media. As shown in **Fig S2D**, we did not observe differences in the number of cells across the diluted spots between strains. Taken together, the observed changes in the cell envelope of high EV producing bacteria suggest that cell envelope alterations in response to lack of VirR and increased membrane vesicle biogenesis may be connected in Mtb.

The absence of LytR_C solo domain proteins in *Mtb* leads to enhanced cell envelope permeability concomitant to hypervesiculation. The collective cryoEM observations that indicated an enlargement of the compartment next to the cell membrane may be linked to a more permeable cell envelope. Supporting this notion is the recent finding that $virR^{mut}$ or the mutant in the $virR$ ortholog cei (Rv2700) manifest an enhanced uptake of ethidium bromide (EtBr) relative to WT strain (Ballister et al., 2019). We could reproduce these results and add that $virR^{mut}$-Comp strain restores the permeability levels of the WT strain (**Fig 2A**). This enhanced permeability was concomitant to an increased sensitivity to different antitubercular drugs, including meropenem and vancomycin (Ballister et al., 2019). In agreement with these results, we could measure a significantly enhanced sensitivity of $virR^{mut}$ to the PG-targeting drug vancomycin relative to WT and $virR^{mut}$-Comp strains (**Fig 2B**). Interestingly, the

magnitude of the sensitivity to cell wall targeting drugs was either moderate (ethambutol)
or absent (isoniazid) in *virR^{mut}* (Ballister et al., 2019).
Furthermore, we observed enhanced incorporation of fluorescent D-amino acids
(FDAAs) in *virR^{mut}* relative to the WT and complemented strains (**Fig 2C**). These probes
can freely enter cells and may be incorporated into PG by at least three different
mechanisms, depending on the species: through the cytoplasmic steps of PG
biosynthesis and via two distinct transpeptidation reactions taking place in the periplasm
(Kuru et al., 2019). Consequently, the differential labeling observed in *virR^{mut}* relative to
WT strain may be a consequence of altered PG turnover in the mutant.
We then tested whether the enhanced permeability and vesiculation of *virR^{mut}* are also
connected in an *Mtb* mutant lacking *cei* (Ballister et al., 2019). To do this, we generated
cei conditional mutants using the CRISPR interference (CRISPRi) technology (Rock et
al., 2017). Further, we also generated *virR* CRISPRi conditional mutants to properly
compare both strains (**Fig 2D**). When these strains were cultured in the presence of
anhydrotetracycline (ATc) to induce the transcriptional silencing of target genes, we
could measure enhanced vesiculation by analyzing vesicle concentration in the
supernatant by NTA (**Fig 2E**), relative to strains cultured in the absence of ATc. Overall,
these results validate the enhanced permeability phenotype of *Mtb* mutants in the
LytR_C solo domain proteins VirR and Cei and establish that this subfamily of proteins
participates in the regulation of vesicle production in *Mtb*. Moreover, the higher
susceptibility to PG-cross-linking inhibitors in both mutants and the enhanced
incorporation of FDAAs in the *virR* mutant strain suggests defects in PG remodeling.
258
Transcriptional profiling of *virR^{mut}* showcases systemic alterations in cell wall
architecture and metabolism. To identify *virR*-dependent mediators of vesicle
production, we carried out a transcriptional profiling of *virR^{mut}* and compared it to those
from a WT and *virR^{mut}*-Comp strains. Previous efforts to define the transcriptional profile
of *virR^{mut}* showed no significant differences compared to WT (Rath et al., 2013).
However, those experiments were carried out with bacteria grown in rich 7H9 medium,
while here we use a defined high-Fe minimal medium (MM), which is typically used for
vesiculation experiments (Prados-Rosales et al., 2011). We extracted RNA from four
independent cultures (average optical densities (OD) of 0.40, 0.38 and 0.42 for WT,
virR^{mut}, and *virR^{mut}* -Comp, respectively) one day before vesicle isolation, and conducted
RNA-seq (**Fig 3A**). Principal component analysis (PCA) was then performed on the
transcriptomic data, after filtering out lowly expressed genes, as well as genes containing
outlier observations (N=3985 genes tested, see Methods), and revealed significant
differences between strains. The first principal component, explaining 49% of the

variance, separated *virR^mut* strain samples from strains bearing functional copies of *virR*
(**Fig 3B**, WT and *virR^mut*-Comp vs *virR^mut*, one-tailed Mann-Whitney test p=2.02E-2).
Furthermore, the second principal component (28.3% variance) captured *virR*-
independent effects of mutagenesis on transcriptional profiles, as it separated WT
samples from *virR^mut*-Comp and *virR^mut* (one-tailed Mann-Whitney test p=2.02E-2).
After PCA, we conducted differential expression analysis between strains to find 1808,
and 2230 genes differentially expressed in *virR^mut* compared to either WT, or *virR^mut*-
Comp strains, respectively (**Supplementary file S1**, **Fig 3C**, Benjamini-Hochberg
FDR<0.05), representing 45.4% and 56.0% of all genes tested. Although transcriptional
profiles of the WT and *virR^mut*-Comp strains are also significantly diverse, with up to 2184
differentially expressed genes between them, the differences in expression between
virR^mut when compared to either WT or *virR^mut*-Comp are significantly correlated (r=0.63,
p<2.2e-16), and the sets of both upregulated and downregulated genes in *virR^mut* vs any
of the other strains are mutually, directionally enriched (**Fig 3D**). Finally, we defined a
set of N=399 genes that are simultaneously upregulated in *virR^mut* with respect to any of
the two strains containing functional copies of *virR* (logFC>0 and FDR<0.05 for both
virR^mut vs WT and *virR^mut* vs *virR^mut*-Comp), as well as a set of N=502 genes
downregulated in *virR^mut* vs both *virR*-containing strains. This way, the differential
expression signature found for these genes in the *virR^mut* strain is most likely VirR-
dependent and not-induced by non-specific consequences of the mutagenesis
procedures.
Next, we looked for gene ontologies (GO) enriched among these *virR*-dependent genes,
stratified by the direction of the effects (399 up-regulated, and 502 down-regulated in
virR^mut relative to WT and *virR^mut*-Comp alike). A total of 51 GO terms related with
biological processes and cell compartment appeared significantly enriched among genes
upregulated in the *virR^mut* strain (FDR=10%), while 85 terms were enriched among
downregulated genes (FDR=10%, see Methods, and **supplementary file S2**). When
focusing on the most prominent of these enrichments (OR>2), the analyses highlighted
several groups of biological processes and cell compartment ontologies that appeared
altered in *virR^mut* with respect to both WT and *virR^mut*-Comp strains (**Fig 3E**).
First, among genes that were expressed at lower levels in *virR^mut*, we found terms related
to several stresses associated with the intra-phagosomal environment (Schnappinger et
al., 2003), such as response to hypoxia (FDR=1.7E-5), heat (FDR = 2.2E-2) or response
to metal ions (FDR=3.4E-2), suggesting that responses to host-induced stress are
compromised in the mutant, as it is also suggested by the weaker, albeit significant
enrichments of responses to host immune response (OR=1.88, FDR=7.47e-02, not
highlighted in **Fig 3E**). Furthermore, the mutant strain showed a reduction in transcripts

related with secretion (FDR=7.6e-03), and, specifically, protein secretion by the type VII secretion system (FDR=5.2 e-03). Contributing to this enrichment, we found that the key antigen ESAT-6 was significantly down-regulated in $virR^{mut}$ with respect to both WT (log2FC=-0.17, FDR=4.1E-3) and $virR^{mut}$-Comp strains (log2FC=-0.19, FDR=2.6E-3), while $cfp10$ (Rv3874), was similarly less expressed in $virR^{mut}$ than in WT (log2FC=-0.24, FDR=2.6E-3).

Second, we found enriched terms related with lipid metabolism and transport both among the sets of genes up and downregulated in the $virR^{mut}$ strain. On the one hand, genes downregulated in $virR^{mut}$ appeared enriched in fatty acid biosynthetic processes (FDR=6.7E-2), suggesting that the regulation of both anabolic and catabolic pathways of lipids, key for bacterial survival and pathogenesis upon infection (Ghazaei, 2018), are, at least in part, VirR-dependent. Furthermore, the mutant strain also shows a significant induction of a number of GO terms related with cell wall composition (e.g. cell wall organization: FDR=3.9e-2), including terms associated with the biosynthesis key components of the mycobacterial cell wall such as PDIMs (DIM/DIP cell wall layer assembly enrichment FDR=9.5e-03) and importantly peptidoglycan (peptidoglycan-based cell wall biogenesis FDR=2.2e-2) (**Fig 3F**).

Third, we found a series of enrichments related with global regulation of gene expression (positive: FDR=4.4e-2, negative: FDR=5.4e-2), and RNA biosynthesis (positive: FDR=5.4e-2, negative: FDR=5.8e-2), among genes downregulated in $virR^{mut}$, suggesting a global alteration of the transcriptional landscape in the mutant.

Fourth, we found that metabolic reprogramming in the $virR^{mut}$ strain is not restricted to lipids but also involves amino acids (cellular amino acid metabolic processes FDR=2.1e-3), and carbohydrates (cellular carbohydrate biosynthetic process FDR=6.8e-3), enriched among genes upregulated in the mutant. Similarly, energy production also appears altered in $virR^{mut}$, with related GO terms enriched among downregulated genes, such as electron transfer activity, (FDR=4.2e-2), as well as terms enriched among upregulated genes in the mutant such as NADH dehydrogenase activity, (FDR=3.3e-3). All together, these results combined with the differences concerning lipid metabolism discussed above, suggest a global remodeling of metabolic networks in $virR^{mut}$.

***virR*-dependent genes intersect the regulons of key transcriptional regulators of responses to stress, dormancy, and cell wall remodeling.** To shed light on the specific transcriptional circuits whose disruption in the $virR^{mut}$ strain may lead to the differential gene expression patterns described, we resorted to the regulon annotations previously reported (Minch et al., 2015). In this work, authors collected ChIP-seq data on binding events between a panel of transcription factors (TFs) and DNA to annotate

interactions between 143 regulators and the promoters of 7248 target genes (peaks between -150 and +70 nucleotides from transcription staring sites). Capitalizing on these data, we interrogated whether the sets of genes up- (N=399) and downregulated in the $virR^{mut}$ strain (N=502) were significantly enriched among each of the regulons reported by Minch et al. A total number of 10 regulons appeared enriched among either down (9) or upregulated genes (1) in $virR^{mut}$ (**Supplementary file S3** and **Fig 4A**, highlighted TFs at OR>2 FDR<0.1). Furthermore, the expression of 7 out of 10 of the TFs controlling these regulons was itself found to be $virR$-dependent in our transcriptional data (**Fig 4B**). Genes more highly expressed in the $virR^{mut}$ strain appeared significantly enriched among Rv0339 targets (OR=9.1, FDR=7.5E-2, **Fig 4A**), named IniR because of its role as key regulator of the isoniazid-induced operon $iniBAC$, a key element of the response to cell wall biosynthesis inhibition induced by isoniazid (Boot et al., 2017) and linked to vesicle biogenesis in Mtb (Gupta et al., 2023). Several additional transcription factors known to regulate cell wall biosynthesis and homeostasis were also enriched among genes downregulated in the $virR^{mut}$ strain. These include SigF (OR=5.8, FDR=8.1E-2), a sigma factor controlling cell envelope organization (Williams et al., 2007), $mtrA$ (OR=2.2, FDR=6.0E-2), a TF controlling peptidoglycan cleavage and alterations in the context of regulation of cell growth and division, whose disruption has recently been shown to lead to increased permeability (Peterson et al., 2023); as well as Rv0472c (OR=11.7, FDR=3.5E-2), a transcription factor involved in the regulation of mycolate remodeling pathways (Peterson et al., 2019), a process used by the pathogen upon phagocytosis to evade macrophage-induced stress and enter dormancy. Interestingly, two genes under the control of Rv0472c, that are key to enable this process, are the mycolate desaturases $desA1$ and $desA2$ (Peterson et al., 2019), which appear downregulated in the $virR^{mut}$ strain with respect to WT (logFC=-0.4, -0.1; FDR=2.3E-11, 3.2E-2, for $desA1$, and $desA2$, respectively), as well as with respect to $virR^{mut}$-Comp (logFC=-0.4, -0.4; FDR=7.2E-12, 5.3E-21). Furthermore, Rv0472c has also been linked with the regulation of key proteins in the response to several sources of stress in Mtb, such as the putative nitroreductase Rv3131 (response to oxidative/nitrosative stress), as well as the triacylglycerol synthase $tgs1$, responsible for the synthesis and accumulation of triacylglycerol, a major energy source for the bacterium during dormancy (Chauhan and Tyagi, 2009), both of which are strongly downregulated in the $virR^{mut}$ strain vs both WT: (logFC=-1.2, -1.7; FDR=3.1E-19, 3.2E-14) and $virR^{mut}$-Comp (logFC=-0.4, -1.9; FDR=3.9E-5, 1.4E-17).

The adjacent genes Rv3131 and $tgs1$ (Rv3030c) appear under the control of other key regulators of the response to intracellular stress in Mtb and the transition to dormancy, whose regulons appear as well enriched among genes downregulated in $virR^{mut}$ (**Fig**

4C). These include *devR* (OR=2.7, FDR=1.9E-5), the canonical regulator of the response to hypoxia and transition to dormancy in *Mtb* (Park et al., 2003), as well as Rv1985c (OR=4.1, FDR=4.5E-11), a key regulator of the homeostasis of the threonine biosynthesis pathway, a metabolic route needed by the bacterium for persistence *in vivo* (Hasenoehrl et al., 2019). Three of the four remaining TFs whose regulons we found enriched among *virR^{mut}* downregulated genes have been similarly linked to the response to assorted sources of stress that are cardinal features of the phagosome environment, such as oxidative stress, response to toxic metals, and low pH (Schnappinger et al., 2003). These include the regulator *lsr2* (OR=2.3, FDR=5.1E-4), a global histone-like protein responsible from protecting the DNA physically from oxidative stress (Bartek et al., 2014), the zinc-uptake regulator Zur (*furB*) (Maciąg et al., 2007) (OR=8.8, FDR=4.5E-2), and *tcrX* (OR=2.6, FDR=5.0E-2), part of the recently characterized acid-sensing two-component system tcrXY (Stupar et al., 2024). Notably, we observed that removing catalase from culture medium prior to submitting *virR^{mut}* to a source of oxidative stress like H202 for 8 h, significantly affected its growth relative to WT and *virR^{mut}*-C strains (**Fig S3**). This result validates the transcriptomics data reflecting a defect in regulating oxidative stress in *virR^{mut}* and encourage to investigate its connection with vesicle production.

Cell envelope permeability is a major driver of MEV release. If enhanced cell permeability is a global consequence of *virR* disruption that leads to enhanced vesiculation, we hypothesized that other conditions either inducing or decreasing cell envelope permeability could lead to alterations in vesicle production in *Mtb*. To test this, we treated *Mtb* with sublethal concentrations of thioridazine (TRZ), a known enhancer of cell permeability (De Keijzer et al., 2016); and, alternatively, we cultured the bacillus with cholesterol as a sole carbon source, a condition which reduces mycobacterial cell permeability (Brzostek et al., 2009) (**Fig 5A**). The enrichment of the DevR regulon among the genes that are downregulated in *virR^{mut}* is also suggestive of the existence of a convergent regulatory route controlling both cell permeability and vesiculation under control of this transcription factor, whose activation has been described as a key part of the pathogen's transcriptomic response to cholesterol (Soto-Ramirez et al., 2017), concomitant to a reduction in cellular permeability (Brzostek et al., 2009). Importantly, neither cholesterol nor TRZ altered bacterial viability as measured by the absence of the release of the transcription factor IdeR, as an indication of cell lysis or by spotting serially diluted cultures on solid medium from the corresponding conditions (**Fig S2C,D**). We then measured vesicle levels in culture supernatants and found that *Mtb* significantly releases more EVs upon treatment with TRZ, and significantly reduces vesicle release

in the presence of cholesterol as a sole carbon source (**Fig 5B**), as measured by dot blot

after normalization to colony forming units (CFUs). These results strongly suggest that

cell envelope permeability determines the magnitude of the vesiculation phenomenon in

Mtb. We next investigated whether common transcriptional features underpin the

regulation of cell envelope permeability in the tested conditions. To do that, we compared

the transcriptional response of *virR^{mut}* with that of available datasets including *Mtb*

cultured in the presence of 0.01% of cholesterol as a sole carbon source (Pawełczyk et

al., 2021), and *Mtb* treated with 6 µg ml-1 of TRZ (Dutta et al., 2010). Experimentally, we

retrieved the set of genes up and downregulated in *Mtb* grown in cholesterol vs glucose-

rich culture media (Brzostek et al., 2009), and interrogated whether those genes were

enriched among the sets of differentially expressed genes in *virR^{mut}* vs WT, and *virR^{mut}*

vs *virR^{mut}*-Comp reported in this study (**Fig 5C**). Interestingly, we found that those genes

whose expression changed in the same direction in conditions associated with low, or

high permeability, in both studies were mutually enriched, while the cross-comparisons

between sets with opposite responses to low-permeability-associated treatments in both

studies led to significant depletion statistics. These results validate the initial hypothesis

according to which changes in permeability induced by these two treatments are linked

to the activation of a common transcriptional program. The inspection of genes that are

simultaneously upregulated in treatments associated with low permeability in both

studies identified DevR as a putative regulator associated with the orchestration of this

transcriptional footprint of low cell permeability. The two-component system DevS-DevR

(also known as DosS-DosR) has a well-known role as a global regulator of the transition

to dormancy, and its regulon is consistently upregulated in non-permeable dormant

bacteria (Honaker et al., 2009).

Interestingly, authors found that low-permeability bacteria cultured in a cholesterol

medium, albeit featuring a regular growing phenotype, activated the DosR regulon,

suggesting that cell wall remodeling leading to a decrease in permeability linked with the

induction of DosR can be triggered independently to the transcriptional regulatory

programs leading to growth arrest and dormancy in *Mtb* (Pawełczyk et al., 2021).

Importantly, the integration of the genes that are downregulated in *virR^{mut}* validates this

hypothesis (**Fig 5D and FigS4**), since not only the intersection between the DevR

regulon and the set of genes downregulated in *virR^{mut}* are significantly larger than

expected by chance (see **Fig 4A**, OR=2.67, p=2.7E-7), but also the intersection between

DevR regulon and the genes that are activated by cholesterol in Brzostek et al. data

(OR=1.89, p=3.5e-5). 30 out of the 173 genes annotated in the DosR regulon described

in (Minch et al., 2015) are simultaneously upregulated in conditions linked to low

permeability in both studies.

A similar analytic approach for the comparison of the response of $virR^{mut}$ and the
response to the drug TRZ, a treatment well-known for inducing an increase in
permeability in *Mtb*-treated cells (Dutta et al., 2010) (GSE16626), led to conceptually
similar results, with genes simultaneously downregulated in both TRZ and $virR^{mut}$
(OR=1.96, p=9.1E-4) being enriched. Interestingly, among the 22 genes showing the
largest, and most significant repression effects by both treatments (log2FC>0.25 &
FDR<0.05 simultaneously, for TRZ effects, $virR^{mut}$ vs WT), we found 4 genes whose
functions are related with metal nutrients uptake and homeostasis, such as genes
encoding the bacterioferritin BrfB, the metal exporter CtpJ, the cupper regulator CsoR
and Moew, involved in Molibdenum-cofactors biosynthesis (**Fig 5E**). Taken together,
these analyses highlight the transcriptional parallels between $virR^{mut}$ and both TRZ and
cholesterol conditions, which strongly suggests the existence of a common
transcriptional signature associated with the control of cell permeability.

Modulation of EV levels and permeability in $virR^{mut}$ by cholesterol and TRZ. We
next wondered about the effect of culturing $virR^{mut}$ on both cholesterol or TRZ could have
on cell growth, permeability and EV production. In the case of cholesterol, it has also
been shown to affect other aspects of physiology (redox, respiration, ATP), which can
directly affect permeability (Lu et al., 2017). We monitored $virR^{mut}$ growth in MM
supplemented with either glycerol, cholesterol as a sole carbon source, or TRZ at 3 ug
ml^{-1} for 20 days. While cholesterol significantly enhanced the growth $virR^{mut}$ after 5 days
relative to glycerol medium, supplementation of glycerol medium with TRZ restricted
growth during the whole time-course (**Fig S5A**). The study of cell permeability in the
same conditions indicated that the enhanced cell permeability observed in glycerol MM
was reduced when $virR^{mut}$ when cultured with cholesterol as sole carbon source.
Conversely, the presence of TRZ increased cell permeability relative to the medium
containing solely glycerol (**Fig S5C**). As we have previously observed for the WT strain,
either condition (Chol or TRZ) also modified vesiculation levels in the mutant accordingly
(**Fig S5B**). Interestingly, TLC analysis of cell membrane polar lipids showed that $virR^{mut}$
grown in cholesterol restores the previously observed higher Ac_2PIM_2 levels relative to
WT when growing in glycerol (**Fig S8A**). These results strongly indicates that other
aspects of mycobacterial physiology besides permeability are also affected in the
absence of VirR and may contribute to the observed enhanced vesiculation.

virR* regulates the protein content of EVs in *Mtb. To capture the impact of
transcriptional adaptation to the proteome, and the degree of divergence between
transcriptional shifts and proteins changes observed in the absence of *virR*, we next

performed label-free peptide mass spectrometry (LF-MS) of whole cell lysates (WCLs)
of $virR^{mut}$, WT, and $virR^{mut}$-Comp strains. Additionally, we isolated EVs from axenic
cultures and obtained their proteomic composition to examine for potential VirR-
dependent differences in protein cargo that may alter the functionality of EVs (**Fig 6A**).
As a result, we obtained proteomic profiles across strains and cell compartments for a
total of N=935 proteins. Upon PCA analysis of the proteomic data (**Fig 6B**), we found
that the first principal component (PC1), explaining 36.9% of total variance, separated
samples from WCLs and EVs in all strains (PC1 differs across cell compartments: one-
tailed Mann Whitney test p=2.1e-05), while the second PC (PC2) (13.6% of total variance
explained) captured strain effects on the proteomic footprint only among EVs (one tailed
Mann Whitney t test for PC2 differences between $virR^{mut}$ EVs and those from either WT
or $virR^{mut}$ -Comp strains (one-tailed Mann Whitney p=1.2 e-02). These results suggest
that differences in protein abundances between EVs and WCLs are larger than
differences found between strains, and that, concerning the differences across strains,
these are more important among EVs. To verify this, we conducted differential protein
expression analyses to find barely 27 proteins differentially expressed between $virR^{mut}$
and any of the two $virR$-expressing strains, which represents 2.9% of the total of proteins
under analyses (**supplementary file S4**). Despite the scarcity of significant differences
in expression across strains in WCLs, some of the features most differentially expressed
in $virR^{mut}$ are, besides VirR itself (**Fig 6C**), proteins involved in key processes for bacterial
survival and virulence, including key metabolic enzymes such as the Acetyl-coenzyme
A synthetase Acs (logFC = -3.92 (FDR=4.5e-3) and -2.96 (FDR=5.2E-2) for the
differences between $virR^{mut}$ and WT, and between $virR^{mut}$ and $virR^{mut}$-Comp,
respectively), as well as FprA, (logFC= -3.12, FDR=4.5e-3 e-02 for $virR^{mut}$-WT and
logFC= -2.16, FDR=5.9 e-02 for $virR^{mut}$- $virR^{mut}$-Comp), involved in the reduction of
NADP+ into NADPH, both downregulated in $virR^{mut}$.
Conversely to the WCL dataset, we did find more frequent differences in protein
expression across strains in EVs, with as many as 361, and 382 proteins differentially
expressed in $virR^{mut}$ EVs when compared to those from WT and $virR^{mut}$-Comp strains,
respectively (**supplementary file S4**, **Fig 6D**). This represents 38.6 and 40.9% of all
proteins tested, respectively. Similarly to what was observed at the level of RNA
expression, differences in expression for the contrasts ($virR^{mut}$-WT) and ($virR^{mut}$- $virR^{mut}$-
Comp) were directionally enriched (**Fig 6E**), thus motivating the definition of two sets of
proteins that 1) were observed at higher levels in the EVs isolated from $virR^{mut}$ than in
those from both WT and $virR^{mut}$ -Comp strains (N=117 $virR^{mut}$ -upregulated proteins); or
2) were observed at lower levels in the EVs from $virR^{mut}$ than in those from any of the
other two strains (N=108 $virR^{mut}$-downregulated proteins). Interestingly, as many as 21

out of the 117 proteins that appeared enriched in $VirR^{mut}$ EVs were also upregulated in $virR^{mut}$ at the transcriptional level (One-sided Fisher exact test of enrichment OR=2.02, p=7.0e-3) suggesting that at least a part of these differences in EVs' protein cargo are a consequence of differences in gene expression already observed at a transcriptional level. Notwithstanding that, proteins appearing at lower concentrations in $virR^{mut}$ EVs were not enriched among genes bearing significant transcriptional effects in the mutant. Regarding downregulated EV proteins in $virR^{mut}$ we observed robust enrichments in GO terms related with protein biosynthesis, including translation (FDR=9.8 e-22), ribosome (FDR=5.3 e-29), and ribosome biogenesis (FDR=4.1 e-27, **Fig 6F** and **supplementary file S5**). Consistent with these results, 44 ribosomal proteins (out of 49 detected) were found to be significantly less expressed in the $virR^{mut}$ strain relative to either WT or $virR^{mut}$-Comp strains (FDR<0.05 for at least one comparison), while no ribosomal protein appeared significantly upregulated in $virR^{mut}$. Furthermore, 30 of these proteins were simultaneously downregulated in $virR^{mut}$ when compared to both VirR-containing strains (**Fig S6**) implying a stark ribosomal depletion in the EVs in the absence of VirR. Furthermore, we found that the two bacterioferritin proteins BfrA and BfrB were simultaneously less abundant in $virR^{mut}$ than in both WT and $virR^{mut}$-Comp strains (logFC<-1.5 and FDR<7.3E-4, for differences $virR^{mut}$ vs WT and $virR^{mut}$ vs $virR^{mut}$-Comp in both genes, **Fig 6G**, and **supplementary file S4**), together contributing to the enrichment of the GO term ferroxidase activity (FDR=6.4 e-02, **Fig 6F**). This result is suggestive of a possible defect of $virR^{mut}$ EVs in iron sequestration (Mohammadzadeh et al., 2021; Prados-Rosales et al., 2014b). We also found a series of metabolic pathways enriched among proteins upregulated in the EVs of the $virR^{mut}$ strain, including enzymes involved in metabolism of nucleotides (e.g. purine nucleotide biosynthetic process FDR=2.0e-02, amino acids (alpha amino acids FDR=8.3e-04), as well as carbohydrates (hexose metabolic process FDR=6.3e-2), whose catabolism appears also enriched among proteins downregulated in the mutant (e.g. glycolytic process FDR=2.2e-2 and carbohydrate catabolic process FDR=4.7E-2). The simultaneous enrichments of these metabolic pathways observed both among upregulated and downregulated vesicular proteins (**Fig S7**), indicates a complex, VirR-dependent, rewiring of metabolic routes in the vesicles from the mutant strain.

Further, proteins upregulated in $virR^{mut}$ EVs (N=117) were enriched in terms related to mycolic acids metabolic processes (FDR=2.1E-2), and peptidoglycan-based cell-wall biogenesis (FDR=9.0 e-02). Proteins such as InhA (logFC=3.71, 2.26 FDR=2.20 e-05, and 1.4E-3 for comparisons $virR^{mut}$ vs WT and $virR^{mut}$ vs $virR^{mut}$-Comp, respectively), and KasB (logFC=1.83; 2.49 FDR=2.23 e-02; 4.4e-3), involved in mycolic acid metabolism were also upregulated in $virR^{mut}$ (**Fig 6H**).

The upregulation of these proteins, along with the enrichments at the transcriptional level
of DIM/DIP biosynthetic pathways (upregulated in *virR^{mut}*), suggests a shift in the control
of the pool of propionyl-CoA, both at the level of its production, -as a byproduct of the
catabolism of methyl-branched lipids, odd-chain-length fatty acids as well as cholesterol-
and its detoxification by the bacterium through different routes, such as the
methylmalonyl pathway, the glyoxylate cycle and the methylcitrate cycles (Lee et al.,
2013). To shed light on these changes at the metabolic level, we conducted a metabolic
profiling to quantify the concentration of metabolites that are central in these different
detoxification routes including propionyl-CoA: acetyl-CoA, pyruvate, methylcitrate, and
methylmalonyl-CoA (**Fig 6I**). Only methylmalonyl-CoA was found to be significantly
repressed in *virR^{mut}* (p=0.028). We also observed differences between *virR^{mut}* and *virR^{mut}*
-Comp strains in methylcitrate. This result, along with the trend towards increased
protein, and especially gene expression that is observed for genes involved in the
biosynthesis of PDIMs from methylmalonyl (including phenolpthiocerol synthesis type-I
polyketide synthases A-E, (ppsA-E, **Fig 6I**), among others), suggests that the
methylmalonyl pool is depleted in the mutant strain because of an increased activity of
the PDIM biosynthetic pathway. Furthermore, most of the subunits of the
acetyl/propionyl-CoA carboxylase Acc (AccA1, AccA2, AccA3, AccD1, AccD5 appear
repressed in *virR^{mut}* EVs (**Fig 6I**), suggesting that the production of methylmalonyl Co-A
from the degradation of propionyl CoA is inhibited within EVs of the mutant strain.
Taken together, the omics analyses align with the structural and functional differences
observed in *virR^{mut}*. We reasoned that the increased expression of PDIMs biosynthetic
genes and the reduced expression of genes related to protein synthesis may be triggered
to compensate the enhanced permeability measured in *virR^{mut}*. In addition, this enhanced
permeability leads to an increase in the release of EV, which seems to be depleted in
ribosomes and iron storage proteins.
595
The absence of VirR leads to an aberrant enlargement of PG in *Mtb*. To elucidate
the specific molecular defects that explain the periplasm enlargement, the enhanced
permeability, and the antibiotic sensitivity phenotypes, we performed a biochemical
analysis of WT and *virR^{mut}* cell envelopes. First, we analyzed the total lipid profile by
liquid chromatography-mass spectrometry (LC-MS) and observed significant differences
in prevalent polar lipids linked to the cell membrane including, lysophosphatidic and
phosphatidic acid, species of monoacylated and diacylated glycerol,
phosphatidylethanolamines and phosphatidyl-myo-inositol mannoside (PIM) species,
being more abundant in *virR^{mut}* relative to WT and complementing strain. (**Fig 7A**). To
validate these results, we ran thin layer chromatography (TLC) experiments on the same

lipid extracts and found that the mutant carries more phospholipids in the membrane
including more AcPIM$_2$ and PE (**Fig 7B**). Moreover, we could not measure major
differences in the level of mycolic acid species (**Fig 7C**), while significantly more PDIMs
were detected in the mutant relative to the other strains (**Fig 7D**). This later result partially
validates proteomics and metabolomic data and suggest that the methylmalonyl pool is
depleted in the mutant strain because of an increased activity of the PDIM biosynthetic
pathway. However, other lipids including SL1 or PAT/DAT that share a common
precursor (propionyl- CoA) with PDIM may also explain the depletion of methylmalonyl.
Notably, we observed a reduction in the level of these lipids (SL1 or PAT/DAT) in $virR^{mut}$
cultured in glycerol relative to WT and complemented strains, suggesting that the excess
of PDIM synthesis can take away methyl malonyl CoA from the biosynthesis of SL-1 and
PAT/DAT in the absence of VirR (**Fig S8B**).
When analyzing apolar lipids, we also observed increased levels of free mycolic acids
(FMA) and free fatty acids (FFA) upon *virR* deletion (**Fig 7E**). The enhanced release of
these apolar lipids has also been documented in a *C. glutamicum* mutant in Cg0847
(Baumgart et al., 2016), a canonical LCP protein. Importantly, we could not detect any
of these lipids in EVs from $virR^{mut}$ (**Fig 7F**), while major polar phospholipids could be
resolved in EVs from all strains (**Fig 7G**), as previously reported (Prados-Rosales et al.,
2011). This result indicates that *Mtb* releases free apolar lipids independently from
membrane associated EVs.
Next, we performed a glycosyl composition analysis of isolated cell walls (including
mycolyl-arabinogalactan-peptidoglycan complex) and determined that there were no
significant differences between strains in AG content as well as in the ratio between
arabinose and galactose (A/G) (**Fig 8A**). Similarly, no changes were observed in the
galactan length as measured by the ratio between galactose (G) and rhamnose (R) G/R)
(Justen et al., 2020). We noticed that the ratio between N-acetylglucosamine (NAGc)
and rhamnose (R) (NAGc/R) was significantly higher in $virR^{mut}$ relative to WT, suggesting
an increase in PG content in this strain (**Fig 8A**). The validation of these results was
performed independently by measuring the amount of diaminopimelic acid (DAP) in
isolated cell walls, as a surrogate of PG content, and determined that $virR^{mut}$ carries two
times more PG than WT or complemented strains (**Fig 8B**).
We next obtained pure PG preparations of each strain and performed an analysis by
atomic force microscopy (AFM) (**Fig 8C and Fig S9**). This technique has been recently
used to investigate the nanometric features of isolated Gram-positive bacterial cell walls
(Pasquina-Lemonche et al., 2020). Semi-automated analysis of liquid AFM images from
mycobacterial isolated PG samples revealed that: (i) the absence of VirR leads to a
significant reduction in PG pore diameter (down to 9.2 ± 0.6 nm) relative to WT ($16 \pm$

3.3 nm) and complemented strains (10 ± 0.8 nm) (**Fig 8C, D and Fig S9**); (ii) the number

of pores is significantly higher in *virR^mut* compared to WT and complemented strain (**Fig**

8E and Fig S9); (iii) manual analysis of AFM images in ambient conditions revealed that

virR^mut PG is significantly thicker than that of the other strains of the study (**Fig 8F**). These

results show for the first time the nanometric architecture of isolated PG in mycobacteria

by AFM and suggest a role for VirR in regulating PG thickness and porosity. Strikingly,

we found that the PG enlargement observed in the *virR^mut* correlated with increased

susceptibility to the muramidase lysozyme (**Fig S10**), indicating modifications in PG

remodeling in *virR^mut*. In fact, a deeper MS analysis of PG showed a significant increase

in the abundance of deacetylated muropeptides in *virR^mut* relative WT and *virR^mut*-Comp

strain (**Fig 8G, H**). This PG feature has been linked to variability in lysozyme susceptibility

in other Gram positive bacterial species (Bera et al., 2005). Collectively, although the

structural changes at nanoscopic level of the peripheral areas of the cell observed by

AFM leads to a more dense and thick structure harder to penetrate, the chemical

changes in the peptidoglycan structure observed and biochemical assays suggest that

changes associated with alterations in VirR-regulated PG remodeling makes the wall

more permeable and support increased release of EV in *Mtb*.

VirR interacts with canonical LCP proteins. As mentioned above VirR is a LytR_C

solo domain protein with no catalytic LCP domain. Other members of the LCP protein

family like Rv3267 and Rv3287 are the main PG-AG ligases (Grzegorzewicz et al., 2016;

Harrison et al., 2016) and also carry a LytR_C domain. We have previously shown that

VirR interacts with itself (Rath et al., 2013). Consequently, we reasoned that if VirR

contributed to PG-AG ligation it could do it via interaction with canonical LCP proteins

through the LytR_C domain. Although VirR is one of the most abundant proteins found

in tuberculin preparations (Prasad et al., 2013), we determined that it localizes at sites

of nascent cell wall by fluorescence microscopy (**Fig S11**). Therefore, its location could

initially allow for interactions with other cell surface proteins like LCP. Both recombinant

VirR and the main AG-PG ligase Lcp1 (Rv3267) oligomerize in a blue-native PAGE gel

(**Fig 9A**). This observation suggests that they interact with proteins sharing similar

domain features. We next performed a flotation assay where we examined the behavior

of VirR on an iodixanol gradient in the presence or absence of Lcp1. Using a specific

antibody raised against VirR, we observed that the presence of Lcp1 displaces the

relative location of VirR toward upper fractions of the gradient, indicating a potential

interaction of both proteins and the formation of higher molecular weight complexes (**Fig**

9B). To investigate the interaction of virR and Lcp1, along with other Lcp proteins, in a

more physiological environment we used the bipartite splitGFP approach (Cabantous et

al., 2013) (**Fig 9C and Fig S12**), which has been recently utilized to demonstrate the co-
localization and interaction of the membrane-bound metal ATPase CtpC with the
chaperon PacL1 (Boudehen et al., 2022). We first recapitulated the VirR self-interaction
and added that such interaction implicates the LytR_C domain since no differences in
fluorescence were measured between full and truncated versions of VirR (VirRsol)
lacking the first 41 amino acids. As negative controls, we included interactions between
VirR and the cell surface associated metal ATPase CtpC (**Fig 9C**). When testing
interactions between VirR and other Lcp proteins (Rv0822c, Lcp1 (Rv3267) and Rv3284)
we could not measure any GFP signal when full Lcp proteins were used in the assay.
Conversely, a significant increase in fluorescence was measured for all combinations
when truncated Lcp proteins including either the C-terminal domain or specifically the
LytR_C domain were tested (**Fig 9C**). These results strongly suggest that VirR interacts
with Lcp proteins and that these interactions occur via LytR_C terminal domains.
693

DISCUSSION
We reported *virR* as the first gene implicated in the biogenesis of EVs in *Mtb* ten years
back (Rath et al., 2013). A transposon mutant in *virR* (*virR^{mut}*) overproduces MEVs and
this provokes augmentation of cytokine responses in mouse and human macrophages.
This mutant manifested an attenuated phenotype in experimental infections on
macrophages and mice (Beaulieu et al., 2010; Rath et al., 2013). Such studies
established that VirR has a role in virulence by negatively modulating the release of
MEVs. These studies provided evidence that vesiculogenesis in *Mtb* is genetically
regulated. Since then, conditions such as iron starvation (Prados-Rosales et al., 2014b)
and other genes, including the bacterial dynamin-like *iniAC* (Gupta et al., 2023), and the
Pst/SenX3-RegX3 signal transduction system (White et al., 2018) have been shown to
contribute to the biogenesis of MEVs. Interestingly, VirR, a LytR_C solo domain protein
(**Fig S1**), shares homology with Lcp proteins, which in *Mtb* are known to participate in
the enzymatic connection between AG and PG (Grzegorzewicz et al., 2016; Harrison et
al., 2016). This suggests VirR function might be linked to cell wall remodeling.
709   In this study, we investigated how VirR can control the magnitude of MEV release
in *Mtb*. We did this by studying the underlying enhanced vesiculogenesis phenomenon
observed in *virR^{mut}* using genetic, transcriptiomics, proteomics and ultrastructural and
biochemical methods. It was clear from our cryo-EM studies that *virR^{mut}* has an aberrant
cell wall as the layer above the cell membrane was significantly larger than that of WT
and complemented strains. The enlarged compartment showed higher granular texture,
a feature that has been previously associated to PG precursors in unrelated Gram-
positive bacteria (Zuber et al., 2006). This observation and previous studies showing the

increased sensitivity of *VirR^{mut}* to vancomycin (Ballister et al., 2019) strongly suggested
that the cell wall defect is connected to PG remodeling. The use of AFM on isolated PG
confirmed this and showed for the first time the nanostructure of PG in *Mtb*, providing
information on PG thickness, pore size and number of pores. AFM indicated that *virR^{mut}*
PG is thicker, has significantly more pores of small size. Considering the pore sizes
measured across strains (ranging from 9-16 nm), we initially ruled out the notion that
EVs (which are spheres of 50-300 nm in diameter) are released through the peripheral
cell wall. Therefore, *Mtb* must have an alternative mechanism to release EVs, possible
during the division process where the PG is most fragile, presumably due to defects in
crosslinking, something that awaits further experimental verification. These events were
not possible to visualize in AFM given that PG fragments from all the strains were lacking
the poles or clearly defined division sites but could be the subject of further studies in the
future. However, we cannot exclude that MEV plasticity could allow for crossing the cell
wall via PG pores.

A recent study showed that *Corynebacterium glutamicum* can produce and
release different types of MVs through different routes, including mycomembrane
blebbing or bubbling cell death, depending on the stress at which *C. glutamicum* is
exposed (Nagakubo et al., 2021). In the present study, we have shown that *virR^{mut}*
overproduces EVs under normal growing conditions in the absence of cell lysis,
according to our assay which measured cytoplasmic leaking of the transcription factor
IdeR or serially diluted spots on solid medium. Overall, we show that EV production in
Mtb under our growing conditions, is not linked to cell death. This fact largely excludes
the bubbling cell death mechanism. Moreover, our lipidomic analysis now and before
(Prados-Rosales et al., 2011) indicates the absence of mycomembrane lipids in isolated
MEVs from all strains, ruling out that MEVs originate at the mycomembrane. These
apparent disagreements with the *C. glutamicum* study (Nagakubo et al., 2021) might
indicate that *Mtb* EV biogenesis may differ from that of *C glutamicum*, and other mycolic
acid-containing bacteria with shorter fatty acids. Additional variables that may explain
differences between studies is the growth media and the time at which vesicles are
isolated (Gupta et al., 2023). Of note, *M. smegmatis* EVs were isolated at day 6
(Nagakubo et al., 2021), a time at which this fast-growing mycobacteria is already in
stationary phase.

The transcriptional profile of *virR^{mut}* can provide a partial explanation for the
attenuated phenotype of the mutant due to a downregulation of genes activated upon
stresses associated to the intraphagosomal environment. It is particularly interesting to
observe the specific reduction in transcript levels of the two most important effectors of
the esx-1 secretion system. On the other hand, genes involved in PDIM synthesis were

upregulated. It is possible that the specific upregulation of PDIM synthesis in absence of
virR is part of compensatory response to a reduced permeability. We were surprised to
find no differences between protein profiles in whole cells between WT and *virR^mut*
strains, despite difference in transcriptomics. A biological interpretation of these results
would point to the existence of global compensatory mechanisms of post-transcriptional
regulation used by the bacteria to partially recover comparable proteomic profiles in
virR^mut to those observed in the WT.

Contrary to whole cell lysates, EV protein profiles between *virR^mut* and WT were
largely different. In this context, it is accepted that certain proteins and lipids are enriched
in bacterial EVs and that preferential inclusion of such antigens in MVs supports a
specific vesicle biogenesis mechanism. In Gram positive bacteria and mycobacteria both
membrane and cytosolic proteins are included in EVs since the origin of these structures
is linked to the cell membrane (Prados-Rosales et al., 2011) and, presumably, the
incorporation of intracellular proteins may occur during the process of vesicle formation.
We found a higher abundance of proteins related to PG and MA synthesis in *virR^mut* EVs
relative to WT. Therefore, these two processes seem to contribute to EV biogenesis in
Mtb. Additional data presented in this study, which includes ultrastructural and chemical
analyses confirm this. Both PG and MA synthesis are complex processes, including a
diverse set of genes. What parts of the biogenesis of these molecules are important for
the formation of EVs is a matter of future investigations.

Another finding of the study is the connection between permeability and the
magnitude of the vesiculation process in *Mtb*. We linked these two phenomena first, by
measuring enhanced vesiculation and permeability in the same strain (*virR^mut*) and
second, by modifying the permeability using chemical (via sublethal exposure to TRZ)
and nutritional supplements (via providing cholesterol as sole carbon source) in WT *Mtb*.
TRZ, a repurposed drug that belongs to the class of phenothiazines, is known to affect
respiration in *Mtb* (Amaral et al., 1996). It is also assumed that TRZ can inhibit efflux
pumps leading to an increased susceptibility to first line antitubercular drugs when tested
in combination (De Knegt et al., 2014). This provides an explanation for the synergistic
effects of such combinations. An independent explanation of such synergy may also
come from the increased permeability, as it has been recently shown by a quantitative
proteomics analysis of *Mtb* cells exposed to sublethal concentration of TRZ (De Keijzer
et al., 2016). When we grew *Mtb* with cholesterol as a sole carbon source, a condition
that slows mycobacterial growth but also reduces cell permeability via cholesterol
accumulation in the cell wall (Brzostek et al., 2009), we observed the opposite trend on
MEV biogenesis. We consider this finding of high relevance since *Mtb* permeability is
one the major concerns when developing potential antimycobacterial drugs.

Nevertheless, bacterial cell permeability is a complex property that it is controlled by
physiochemical, biological, and chemical processes such as stereochemistry,
lipophilicity, saturation and unsaturation, flexibility, viscosity, fluidity, pressure,
temperature and physiological conditions (Nagamani and Sastry, 2021). How these
variables change to allow the release of MEVs is not known.
This study also proposes a model VirR function whereby VirR may work as a
scaffold for canonical Lcp proteins, which in *Mtb* links PG to AG. Based on the
demonstration that VirR interacts with canonical Lcp proteins, we believe that the
absence of VirR deregulates the localization of canonical Lcp proteins. Although we
cannot rule out the contribution of other phenomena including alteration of the PG
molecule, VirR is a central scaffold in the cell wall remodeling process. Taking into
consideration the fact that *VirR^{mut}* is attenuated in virulence on preclinical models of
infections and manifests an enhanced vesiculogenesis and permeability, VirR may
represent a novel and interesting drug target. We envision that targeting VirR may not
only make *Mtb* weaker and more permeable to other antitubercular drugs, but to
stimulate the immune system, via MEVs, to fight infection.
807
808 **MATERIALS AND METHODS**
809
Bacterial Strains, media and growth conditions. *Mtb* H37Rv (ATCC) and derivative
strains *virR^{mut}* and *virR^{mut-}C* (complemented strain) (Rath et al., 2013) were used in this
study. Bacteria were initially grown in Middlebrook 7H9 medium (7H9) supplemented
with 10% (v/v) oleic acid-albumin-dextrose-catalase (OADC) enrichment (Becton
Dickinson Microbiology Systems, Spark, MD, United States), 0.5% (v/v) glycerol and with
Tyloxapol 0.05% (v/v; Sigma-Aldrich, Burlington, MA, United States) prior to inoculation
in a minimal medium (MM) consisting of KH_2PO_4 1 g/l, Na_2HPO_4 2.5 g/l, asparagine
0.5 g/l, ferric ammonium citrate 50 mg/l, $CaCl_2$ 0.5 mg/l, $ZnSO_4$ 0.1 mg/l, without
Tyloxapol 0.05% (v/v), containing 0.7% (v/v) glycerol, pH 7.0.
Mtb was also cultured in MM in the presence of sublethal concentrations of thirodazine
(TRZ) of 6 µgml^{-1} as previously described. We determined that this concentration does
not affect cell growth. *Mtb* was also cultured in MM with cholesterol as a sole carbon
source as previously described (Brzostek et al., 2009).
Mtb expressing fluorescent VirR was generated by cloning VirR into pMV261-Venus
using fast cloning approach (Li et al., 2011).
M. smegmatis mc^{2}155 (ATCC 700084) and recombinants derived from this strain were
grown at 37°C in 7H9 medium (Difco) supplemented with 10% albumin-dextrose-catalase
(ADC, Difco) and 0.05% Tween-80 (Sigma-Aldrich), or on complete 7H11 solid medium

(Difco) supplemented with 10% OADC(Difco). When needed, streptomycin (25 μg ml^{-1})

was added to the culture media. *Escherichia coli* strains StellarTM (Takara bio, San Jose,

CA, United States) were grown at 37°C in LB broth (Difco) or on L-agar plates (Difco)

supplemented with streptomycin (25 μg ml^{-1}) when required.

***M. smegmatis* expression vectors for split GFP experiment.** Plasmids used for split

GFP experiments were constructed by modifying the plasmid pGMCS-PpacL1-pacL1-

Lk2-GFP11-ctpC(1-85)-Lk15-GFP(1-10) (Boudehen et al., 2022) (**Table S1**). Plasmids

pGMCS, conferring resistance to streptomycin, are shuttle vectors, episomal in *E. coli*

and integrative in mycobacteria through insertion at the attL5 mycobacteriophage

insertion site in the glyV tRNA gene. Lk2 (LEGSGGGGSGGGS) and Lk15

(GPGLSGLGGGGGSLG) are flexible linkers separating the protein domains of interest

from the last 11th b-sheet or the first ten b-sheets of GFP, respectively. First, the Zinc-

inducible PpacL1 promoter was replaced by P1, a strong constitutive promoter from *M.*

smegmatis (Ariyachaokun et al., 2020), resulting in pGMCS-P1-pacL1-Lk2-GFP11-

ctpC(1-85)-Lk15-GFP(1-10). Second, the pacL1 (Rv3269) sequence was replaced by

that of virR (encoding amino-acids 1 to 164) or virRsol (encoding amino-acids 42 to 164)

by In-Fusion HD cloning (Takara). PCR fragments were amplified using appropriate

primer pairs (**Table S1**) and Phusion High-Fidelity PCR Master Mix with GC Buffer

(Thermo Scientific, Walthman, MA, United States). Linear fragments were purified on

agarose gels and inserted into pGMCS backbone linearized by PCR amplification with

appropriate primers, using the In-Fusion HD Cloning Kit (Takara), following the

manufacturer's instructions. Finally, the ctpC sequence was replaced on the resulting

plasmids by the virR or virRsol sequences, or that encoding either the C-terminal part or

only the LytR_C domains of the Rv0822c, Rv3267 or Rv3484 proteins from *M.*

tuberculosis H37Rv. Plasmids were constructed by transformation into *E. coli* Stellar

recipient cells. All plasmids were verified by sequencing before introduction by

electroporation into *M. smegmatis* mc²155 strain.

H$_2$0$_2$ sensitivity assays. Bacterial strains were initially cultured in 7H9+ADN until they

reached an OD600nm of 0.5. Cells were washed in PBS and transferred to roller bottles

including the same medium and 13 mM H$_2$0$_2$. After 8 h, bacterial cultures were serially

diluted and spotted onto 7H10 solid medium.

Generation of the CRISPR mutants. To generate the conditional CRISPR mutants we

took advantage of the CRISPR interference (CRISPRi) technology (Rock et al., 2017).

Briefly, we designed sgRNA oligonucleotides containing the target Rv0431 (*virR*) and

Rv2700 (*cei*) genes sequences (**Table S1**) located immediately upstream of the PAM

sequence. The strengths of the PAM sequence were chosen following authors

recommendation to get high and intermediate gene repression, 158.1 or 42.2-fold

repression for *virR* and 145.2 and 47.3 for *cei*, respectively. The primers were annealed

and ligated into the gel purified BsmBI-digested CRISPRi vector (pIJR965 plasmid)

followed by transformation of competent *Mtb* cells via electroporation. Transcriptional

knockdown of genes were quantified by qRT-PCR; and protein levels were examined by

immunoblot using specific antibodies raised against VirR (Rath et al., 2013) and Cei

(Ballister et al., 2019). Complementation of *virR* mutant strain was performed using the

plasmid previously generated by Gateway® Cloning Technology (Invitrogen) (Rath et al.,

2013). Importantly, neither *virR^{mut}* nor the *virR^{mut}*-C showed altered expression of

downstream genes including Rv0432 or Rv0433 relative to WT strain (**Fig S13**).

EV isolation and purification. *Mtb*-EV were prepared and purified as previously

described. Briefly, the culture filtrate of 1 L cultures of *Mtb* grown in low iron MM without

detergent for 14 days was processed for vesicle isolation by differential centrifugation. In

parallel a 2 ml culture in MM supplemented with 0.05% Tyloxapol to disperse bacterial

clumps was used to determine viability by plating culture dilutions onto 7H10 agar plates

and enumerating colony forming unit (CFU) at the time of culture filtrate collection.

The membranous pellet containing vesicles obtained after ultracentrifugation of the

culture filtrate at 100,000 × g for 2 h at 4°C, was resuspended in 1 ml sterile phosphate-

buffered saline (PBS) and overlaid with a series of Optiprep (Sigma-Aldrich) gradient

layers with concentrations ranging from 30% to 5% (w/v). The gradients were centrifuged

at 100,000 × g for 16 h. At the end 1 ml fractions were removed from the top, diluted to

20 ml with PBS and purified vesicles recovered by sedimentation at 100,000 × g for 1 h.

Vesicle pellets were suspended in PBS before analysis. Protein concentration was

measured by Bradford assay (Bio Rad, Hercules, CA, United States) according to

manufacturer instructions. For all conditions test from which EVs were isolated,

normalization was performed by referring EVs concentration to colony forming units

(CFUs).

Nano particle tracking analysis (NTA). NTA was conducted using ZetaView (Particle

Metrix, Winning am Ammersee, Germany). Instrument calibration was performed prior

to EV analysis using 102 nm polystyrene beads (Thermo Fisher Scientific), according to

manufacturer instructions. Measurements were performed using a 405 nm 68 mW laser

and CMOS camera by scanning 11 cell positions and capturing 60 frames per position

at 25 C with camera sensitivity 85, shutter speed 100, autofocus and automatic

scattering intensity. Samples were diluted in pre-filtered PBS to approximately 106-107 particles·ml^{-1} in Millipore DI water. Analysis was performed using ZetaView Software version 8.05.12 SP1 with a minimum brightness of 30, maximum brightness of 255, minimum area of 5, maximum area of 1000, and s minimum trace length 15. Triplicate videos of each sample were taken in light scatter mode. Particle size and concentration were analyzed using the built-in EMV protocol and plotted using Prism software, version 8.0.1 (GraphPad Inc., San Diego, CA, United States).

Electron-Microscopy. Cells were fixed with 2% glutaraldehyde in 0.1 M cacodylate at room temperature for 24 h, and then incubated overnight in 4% formaldehyde, 1% glutaraldehyde, and 0.1% PBS. For scanning microscopy, samples were then dehydrated through a graded series of ethanol solutions before critical-point drying using liquid carbon dioxide in a Toumisis Samdri 795 device and sputter-coating with gold-palladium in a Denton Vacuum Desk-2 device. Samples were examined in a Zeiss Supra Field Emission Scanning Electron Microscope (Carl Zeiss Microscopy, LLC, North America), using an accelerating voltage of 5 KV.

For Cryo-EM, grids were prepared following standard procedures and observed at liquid nitrogen temperatures in a JEM-2200FS/CR transmission electron microscope (JEOL Europe, Croissy-sur-Seine, France) operated at 200 kV. An in-column omega energy filter helped to record images with improved signal/noise ratio by zero-loss filtering. The energy selecting slit width was set at 9 eV. Digital images were recorded on an UltraScan4000 CCD camera under low-dose conditions at a magnification of 55,058 obtaining a final pixel size of 2.7 Å/pixel.

Density profiles were calculated along rectangular selections with the ImageJ software (NIH, Bethesda, MD, United States). The number of cells analyzed varied from n = 20 (WT), n = 15 ($virR^{mut}$) and n = 23 ($virR^{mut}$-C). The comparisons extracted from this data is statistically relevant with t test with Welch's correction performed. For each cell three representative measurements were obtained. Mean and standard error was then calculated for measurements pooled for each strain.

Western Dot blot. Two µl of EV isolates and two-fold serial dilutions were spotted onto a nitrocellulose membrane (Abcam, Cambridge, United Kingdom) and process for dot blot using anti-EV polyclonal murine serum (Prados-Rosales et al., 2014a) as primary antibody and goat anti-mouse-HRP conjugated as secondary antibody and ECL prime Western Blotting Detection chemiluminescent substrate (GE Healthcare, Chicago, IL, United States). The signal was visualized in a Chemidoc MP imaging system (Bio Rad)s).

RNA sequencing. *Mtb* strains were grown in MM supplemented with 0.05% Tyloxapol at 37ºC until they reached an OD595nm of 0.3 and harvested by centrifugation. The cell pellets were resuspended in 1 ml Qiagen RNA protect reagent (Qiagen, Venlo, Netherlands) and incubated for 24 h at room temperature. Cells were disrupted by mechanical lysis in a FastPrep-24 instrument (MP Biomedicals, Santa Ana, CA, United States) in Lysing Matrix B tubes and RNA was purified with the Direct-zol RNA miniprep kit (Zymo Research, Irvine, CA, United States). The quantity and quality of the RNAs were evaluated using Qubit RNA HS Assay Kit (Thermo Fisher Scientific) and Agilent RNA 6000 Nano Chips (Agilent Technologies, Santa Clara, CA, United States), respectively. Sequencing libraries were prepared using the Ribo-Zero rRNA Removal Kit (Gram-positive Bacteria) (Illumina Inc., San Diego, CA, United Stated) and the TruSeqStranded mRNA library prep kit (Illumina Inc), following the Ribo-Zero rRNA Removal kit Reference guide and the "TruSeqStranded mRNA Sample Preparation Guide. Briefly, starting from 1 µg of total RNA, bacterial rRNA was removed and the remaining RNA was cleaned up using AgencourtRNAClean XP beads (Beckman Coulter). Purified RNA was fragmented and primed for cDNA synthesis. cDNA first strand was synthesized with SuperScript-II Reverse Transcriptase (Thermo Fisher Scientific) for 10 min at 25°C, 15 min at 42°C, 15 min at 70°C and pause at 4°C. cDNA second strand was synthesized with Illumina reagents at 16°C for 1 hr. Then, A-tailing and adaptor ligation were performed. Libraries enrichment was achieved by PCR (30 sec at 98°C; 15 cycles of 10 sec at 98°C, 30 sec at 60°C, 30 sec at 72°C; 5 min at 72°C and pause at 4°C). Afterwards, libraries were visualized on an Agilent 2100 Bioanalyzer using Agilent High Sensitivity DNA kit (Agilent Technologies) and quantified using Qubit dsDNA HS DNA Kit (Thermo Fisher Scientific). Library sequencing was carried out on an Illumina HiSeq2500 sequencer with 50 nucleotides single end reads and at least 20 million reads per individual library were obtained.

RNA-sequencing data analysis. Quality Control of sequenced samples was performed by FASTQC software (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Reads were mapped against the *M. tuberculosis* H37Rv strain (GCF_000195955.2_ASM19595v2) reference genome using Tophat (Trapnell et al., 2009) with --bowtie1 option, to align 50 bp reads. The resulting BAM alignment files for the samples were the input to Rsubread's (Liao et al., 2019) featureCounts function to generate a table of raw counts required for the Differential Expression (DE) analysis. Data preprocessing and differential expression analysis of the transcriptomic data was performed using DeSeq2 (Love et al., 2014). Genes with a median of expression lower than 1 CPM, and genes containing outlier observations were filtered before statistical

modeling, and the remaining N=4004 genes were used for differential expression
analysis. P-values were corrected for multiple testing using the R package, implementing
the Storey-Tibshirani FDR correction method (Storey et al., 2023). Genes with an
FDR<0.05 were deemed as differentially expressed. Data were deposited at Gene
expression Omnibus (GEO); GSE143996.

Quantitative Real-Time PCR
For reverse transcription cDNA was obtained from 10 µl of isolated RNA. Each reaction
included an RT control, where RNA was replaced by 10 µl of DEPC-treated water.
Reaction mixture 1 (volumes refer to a single sample) was prepared, containing 2 µl of
random primers 500 ng/µl; 2 µl of dNTPs mix 10 mM; and 12.5 µl of DEPC-treated water.
A total of 16.5 µl of this mixture was added to 10 µl of each sample, and the samples
were incubated for 5 minutes at 65 °C to remove secondary structures. After 2 minutes
on ice, samples were centrifuged at 12,000 rpm for 5 minutes. Reaction mixture 2
(volumes refer to a single sample) was prepared by mixing 8 µl of 5X reverse
transcription buffer; 2 µl of 0.1 M DTT; 2 µl of RNAseOUT inhibitor; and 1.5 µl of
SuperScript IV reverse transcriptase 200 U/µl. This reverse transcriptase was chosen for
its high specificity in synthesizing cDNA, particularly for nucleic acids with high G+C
content, as is the case with *M. tuberculosis*. A total of 13.5 µl of mixture 2 was added to
each sample, obtaining a final volume of 40 µl. The samples were incubated for 10
minutes at 25 °C and then at 55 °C for 30 minutes to synthesize cDNA. Finally, the
reaction was stopped by inactivating the enzyme for 10 minutes at 80 °C and centrifuging
briefly to recover the full volume. The samples were stored at -20 °C until use.
Amplification reactions were conducted by preparing a PCR reagent mixture containing,
per sample: 1 µl of MgCl2; 1 µl of SYBR Green (previously mixed with L.C. Fast Start
containing the polymerase); 0.25 µl of oligonucleotides specific to the sequences of
interest (Table 1) at a concentration of 20 µM; and 4.5 µl of H2O. In this case, the
sequences of interest are the Rv0432 (Forward primer: GCGTCTACAGTTCCGGGTAC;
Reverse primer: AGGGTACTGGTCAGGCTCTG) and Rv0433 (Forward primer:
GCTGATCTGGGGTGTACACG; Reverse primer: GCCAACAGATGCGGGTAGTA)
genes. The reaction was performed in a Light Cycler using capillaries containing 7 µl of
the PCR mixture and 3 µl of each sample. For the negative control, the 3 µl were replaced
with water, and for the positive control, they were replaced with 0.1 ng/µl DNA from the
reference strain of *M. tuberculosis*. After a brief centrifuge pulse, the volume was
introduced into the capillaries before analysis in the Light Cycler. Gene fold induction
normalized to 16S rRNA was calculated using the $2^{-\Delta\Delta Ct}$ method.

1012

Label-free mass spectrometry analysis. Total cell proteins and MEV-associated proteins were submitted to LC-MS label-free analysis using a Synapt G2Si ESI Q-Mobility-TOF spectrometer (Waters) coupled online to a NanoAcquity nano-HPLC (Waters), equipped with a Waters BEH C18 nano-column (200mm x 75 um ID, 1.8um). Samples were incubated and digested following the filter-aided sample preparation (FASP) protocol (Wiśniewski et al., 2009) Trypsin was added to a trypsin:protein ratio of 1:50, and the mixture was incubated overnight at 37°C ,dried out in a RVC2 25 speedvac concentrator (Christ), and resuspended in 0.1% FA. Peptides were desalted and resuspended in 0.1% FA using C18 stage tips (Millipore). Digested samples (500 ng) were loaded onto the LC system and resolved using a 60 min gradient (5 to 60% ACN). Data was acquired in HDDA mode that enhances signal intensities using the ion mobility separation step. Protein identification and quantification were carried out using Progenesis LC-MS (version 2.0.5556.29015, Nonlinear Dynamics). One of the runs was used as the reference to which the precursor masses in all other samples were aligned to. Only features comprising charges of 2+ and 3+ were selected. The raw abundances of each feature were automatically normalized and logarithmized against the reference run. Samples were grouped in accordance with the comparison being performed, and an ANOVA analysis was performed. A peak list containing the information of all the features was generated and exported to the Mascot search engine (Matrix Science Ltd.). This file was searched against a Uniprot/Swissprot database, and the list of identified peptides was imported back to Progenesis LC-MS. Protein quantitation was performed based on the three most intense non-conflicting peptides (peptides occurring in only one protein), except for proteins with only two non-conflicting peptides.

Proteomics data analysis. Normalization and imputation of the proteomics data was done using the DEP package for R (Zhang et al., 2018). Normalization was done using the vsn method (Huber et al., 2002), which transforms data tackling heteroskedasticity. Imputation of missing values was performed using the bpca method (Oba et al., 2003), which leans on principal component analysis to impute missing values.

Data was analyzed according to a design where differences between strains (*virR^mut* vs -WT, as well as *virR^mut* vs *virR^mut*-Comp) were estimated independently within each cell compartment, (whole cell lysates or EVs). according to a nested design: Expression ~ Compartment + Strain:Compartment.

Differential expression analysis of the proteomics data was performed using limma (Ritchie et al., 2015) and p-values were corrected for multiple testing according to the Storey-Tibshirani method using the R package qvalue (Storey et al., 2003). Proteins with an FDR<0.05 were selected as differentially expressed.

1050

Gene ontology (GO) enrichments. Enrichment analyses of the sets of differentially
expressed genes and proteins were performed in R using the terms of the Gene
Ontology database for testing, version 11.02.2020. Gene sets corresponding to the
biological process and cellular component ontologies between levels 4 to 6 (N=2245)
were selected and filtered according to their size (minimum size =10 in the RNAseq
analyses; and =3 in the proteomics analyses). The selected sets were tested for
enrichment using Fisher's exact test among genes, or proteins consistently up, or down
regulated in *virR^{mut}* with respect to both *virR*-expressing strains in each case. Multiple
testing correction was performed using the qvalue R package (Storey et al., 2023).
Terms with an FDR < 0.1 and an odds ratio > 2 were selected for visualization in figures
3E, and 6F, where node sizes was proportional to the significance of the enrichments. In
these visualizations, two terms were connected depending on the amount of sharing
between the groups of genes contributing to the enrichments in each of them, drawing a
link whenever the intersection between the groups of genes was larger than 50% of the
smallest of the two gene sets involved. Ontology term clusters, represented by colors,
were then assigned using the Louvain's algorithm for modularity optimization (Blondel et
al., 2008), as implemented in the R package igraph (Csardi and Nepusz, 2006).
Visualization of the resulting networks was done using the open-source software Gephi
(Bastian et al., 2009).

Targeted metabolomics. LC-MS analysis was carried out on a Schimazdu LC/MS-8030
coupled with a triple quadrupole mass analyser (QqQ) provided with an electrospray
source in a negative ionization mode. Extracted metabolites were quantified in Multiple
Reaction Monitoring (MRM) mode. A stock standard solution prepared in 17% (v/v)
methanol in water and containing all metabolites: acetyl-CoA, methyl citrate,
methylmalonyl-CoA, pyruvate, was used as external standard. Metabolites were
identified by ion pairing liquid chromatography. Acetyl-CoA, methyl citrate and
methylmalonyl-CoA were separated by reverse phase with a poroshell 120 Phenyl-Hexyl
analytical column (2.1 x 50mm, 2.7 µm, Agilent) and a binary gradient (**Table S2**)
consisting of a mobile phase A with water-methanol 97:3 (v/v), 10 mM tributylamine and
3 mM acetic acid, and mobile phase B composed of pure methanol. Pyruvate was
separated by normal phase using a Kinetex HILIC analytical column (2.1 x 150 mm, 2.6
µm, Phenomenex) and a binary gradient (**Table S3**) consisting of pure water as mobile
phase A and pure methanol as mobile phase B.

The flow rate was maintained at 0.4 ml/min and the injection volume was 20 µl.
Nitrogen was used as nebulizing and drying gas at a flow rate of 1.5 and 15 ml/min,

respectively. The desolvation line (DL) temperature was set at 250°C and the ionization voltage was fixed at 4.5 kV. Argon was used as collision gas to perform collision-induced dissociation at a pressure of 230 kPa. The dwell time was set at 100 ms and the detection was performed in MRM mode (detector voltage of 1.8 Kv), monitoring three transitions for each compound. The transition with higher intensity was selected as quantitative purposes (**Table S4**), while the other two were used for confirmation of the identity.

Lipidomic analysis by UPLC-MS. The effect of virR on relative abundance of different cell wall lipids was characterized as had been done previously for MEVs (Prados-Rosales et al., 2014b). Briefly, cell wall lipids were extracted from triplicate 10 ml cultures of wild type, *virR^{mut}*, and *virR^{mut}* complemented (*virR^{mut}*-C) strains overnight into 3 ml 2:1 chloroform:methanol at room temperature, extracts were dried under nitrogen before being dissolved in 0.75 ml LC-MS grade 2:1:1 isopropanol:acetonitrile:water. Extracts were separated on a Waters ACQUITY BEH C18 column UPLC heated to 55°C and eluted at a flow rate of 0.4 ml/minute with 60:40 acetonitrile:water as mobile phase A and 90:10 isopropanol:acetonitrile as mobile phase B, both containing 10 mM ammonium formate and 0.1% formic acid. Initial condition was 40%, B which increased to 43% at 2 minutes, 50% at 2.1 minutes, 54% at 12 minutes, 70% at 12.1 minutes, 90% at 18 minutes before returning to initial condition at 18.1 minutes to complete the run at 20 minutes. The coupled Waters Synapt G2 q-TOF MS was operated in positive ion resolution mode with the following conditions: capillary voltage 2kV, cone voltage 30V, desolvation gas temperature 550°C, gas flow 900L/hour, source temperature 120°C. Mass spectra were acquired in centroid mode from 100-3000 m/z with scan times of 0.5 second. Leucine enkephaline was used as reference. Relative abundances of Mass-retention time pairs were normalized to total ion current and lipid species were identified using Mycomass (Layre et al., 2011) and Mtb LipidDB databases (Sartain et al., 2011).

Lipidomic analysis by thin layer chromatography (TLC). Lipidomic analysis of both whole cells and EVs by TLC was performed as previously described (Boldrin et al., 2021). *Mtb* strains were grown at 37°C in standing Middlebrook 7H9 and when they reached an OD595 of 0.2, cells were washed and transferred to MM. Cells were grown until they reached an OD595 of 0.5. Further, cells were harvested by centrifugation at 3,500 rpm for 5 min, and pellets were heat-inactivated at 100°C for 1 h and used for lipid extraction. Five milliliters 2:1 (vol/vol) chloroform-methanol was added to the cell pellet and incubated at room temperature (RT) for 12 h with constant stirring. The organic extract was separated by centrifugation (1,000 × g, 5 min, RT) and decantation. The obtained pellet was extracted with 5 ml 2:1 (vol/vol) chloroform-methanol for 1 h at RT

with constant stirring and separated under the same conditions. The combined two
organic extracts (total lipids) were dried under a stream of nitrogen gas at RT and saved
for lipid analysis.

Polar lipids were separated by 2D-TLC by spotting 200 μg of total lipids on 60
F254 silica gel plates (Merck) using chloroform, methanol, water (50:30:6) as a mobile
phase 1 (first dimension), and chloroform, acetic acid, methanol, and water (40:25:3:6)
as mobile phase 2 (second dimension). Analysis of PDIMs and TAGs was performed in
a 1D-TLC format. Two hundred μg of total lipids were spotted on the silica plates and
separated using 9:1 petroleum ether-diethyl ether as mobile phase. Free mycolates were
separated by 1D-TLC using chloroform/methanol/ammonium hydroxide (80:20:2) as
mobile phase. Mycolic acids were extracted and separated from cells as previously
described (Vilchèze and Jacobs, 2007). Sulfolipids (SLs) and diacyltrehaloses (DATs)
were separated chloroform/methanol/water (100:14:0.8) (first dimension) and chloro-
form/acetone/methanol/water (50:60:2.5:3) (second dimension). EV-associated lipids
were separated and visualized following the same procedures as for WCLs.
Development of TLC plates was performed by repeating the process of pulverizing the
staining solution and heating the plate at 100°C 4 times. The staining solution contained
10.5 ml 15% ethanolic solution of 1-naphthol, 6.5 ml 97% sulfuric acid, 40.5 ml ethanol,
and 4 ml water.

1143

PG isolation. To obtain mycobacterial cell walls, whole cell pellets were resuspended in
PBS and boiled for 15 min. Boiled cells were transferred to 2 ml Lysing matrix tubes with
0.1 mm silica beads (M.P. Biomedical, Santa Ana, CA, United States). And mechanically
disrupted in a FastPrep-24TM (M.P. Biomedical) giving 12 cycles at predetermined
speed of 6 during 30 sec. Samples were allowed to chill on ice for 3 min between cycles.
Cell breakage was monitored by optical microscopy (using Methylene blue as stain), if
95% of the cells had not broken perform additional FastPrep-24TM cycles were
performed. Lysed cells were centrifuged at 170 × g for 30 sec to separate them from
beads and the supernatant were transferred to a clean 1.5 ml tube. The suspensions
were further digested with 10 µg of DNase and RNase/ml for 1 h at 4°C to obtain a cell
wall-enriched fraction after centrifugation at 27,000 × g for 10 min. Cell walls were
resuspended in PBS containing 2% sodium dodecyl sulfate (SDS) and the suspension
was incubated for 1 h at 90°C with constant stirring and recentrifuged at 27,000 × g for
30 min, and the supernatant was discarded. This process was repeated twice. The
resulting pellet was washed trice in distilled water, 80% acetone and acetone and
lyophilized. Pellets were resuspended in 0.9 ml Tris-HCL (50 mM, pH 7) buffer, plus 0.1
ml of protease K stock solution. The mixture was incubated at 60°C for 90 min, to digest

any remaining surface proteins attached to the CW and centrifuged again for 3 min at

20,000 × g and resuspend pellet in distilled water. The mixture was then heated at 90°C

for 1 h before centrifugation at 27,000 × g for 30 min. The supernatant was discarded

and washed twice with PBS and four times with deionized water to remove SDS. Pure

PG was obtained from isolated cell walls. Briefly, pure cell walls were resuspended in

0.5% KOH in methanol and stirred at 37°C for 4 days to hydrolyze mycolic acids (MA).

The mixture was centrifuged (27,000 × g 20 min) and the pellet was washed twice with

methanol. MA were separated from cell walls using diethyl ether and recentrifuged at

27,000 × gfor 20 min. The supernatant contained MAMEs. The extraction process with

diethyl ether was repeated twice and let dry. The resulting arabinogalactan (AG)-PG was

digested with 0.2 N H$_2$SO$_4$ at 85°C for 30 min and neutralized with BaCO$_3$ to remove the

arabinogalactan. Treated AG-PG was centrifuged at 27,000 × g for 20 min. Supernatant

contained soluble AG. The resulting insoluble PG was washed four times by

centrifugation in deionized water and stored at 4°C or room temperature.

1175

Analysis of diaminopimelic acid. For DAP quantification (Tsuruoka et al., 1985),

samples were hydrolyzed overnight in 6N HCl at 100°C in 1 ml capacity-crystal ampoules

(Wheaton). Samples were dried and resuspended in water and mixed in a 1:1:1 ratio

with pure acetic acid and ninhydrin reagent (250 mg ninhydrin, 4 ml phosphoric acid 0.6

M, and 6 ml pure acetic acid). Samples were measured at OD 434nm, and the amounts

of DAP were calculated based on a standard curve with pure DAP. The data displays

the average ± standard deviations n=3.

1183

Glycosyl composition of isolated cell walls. 500 µg of purified cell walls were mixed

with 10 µg of inositol and hydrolyzed with 0.1 M trifluoroacetic acid (TFA) for 2 h at 121°C.

The monosaccharides released were then converted into alditol acetates by reduction

with NaBH$_4$ and acetylation with acetic anhydride in pyridine. The products were

analyzed by gas chromatography-mass spectrometry in a 7890A/5975C system from

Agilent, using a DB-5ht (30 m x 0.25 mm x 0.1 µm) fused silica capillary column. The

compounds were identified by comparing the retention times of the peaks with those

recorded for standards analyzed under identical conditions. Quantification was carried

out considering the area of the peaks and the response factors of each compound with

respect to inositol.

1194

1195   **Atomic force microscopy and data analysis**. The dry pellets of purified peptidoglycan

1196   from the three strains were dissolved in pure water and further broken by tip sonication

1197   at 2 mA for 30 sec, three cycles. Then 10 µl of the PG suspension was added to standard

AFM substrate made of mica coated with 0.01 % Poly-L-lysine by incubating the sample during 90 min to achieve good coverage. The sample was further rinsed with water and dried with Nitrogen flow. The AFM in ambient conditions (i.e. air) was performed using Tapping mode in a Bruker FastScan Bio machine (Santa Barbara, CA, United States) with Nunano SCOUT 350 - Silicon AFM probes (spring constant: 42 N/m, Resonance frequency: 350 kHz). The images were taken using a free amplitude of 10 nm with set point of 70-80 % of free amplitude (e.g. 7 nm). For AFM high resolution imaging in liquid the images were acquired in Peak force Tapping mode in imaging buffer made of 10 mM Tris and 150 mM KCl, pH: 8, with the FastScan Bio machine using Bruker Fastscan-D AFM probes (spring constant: 0.1 N/m, Resonance frequency: 70 kHz) at the force range of 1-3 nN peak force set point. The imaging parameters used are as follows; Scan rate: 1 Hz; Scan angle: 0 °; Peak force amplitude: 80-100 nm and with high pixel resolution (512-880).

The thickness measurements were performed with standard Gwyddion 1D height distribution tool (selecting an area containing background and a single leaflet of peptidoglycan). The number of cells analysed varied from n = 23 (for Mtb WT), n = 36 (for $virR^{mut}$) and n = 25 (for $virR^{mut}$-C). The comparisons extracted from this data is statistically relevant with t test with Welch's correction performed. The analysis of the high-resolution images obtained in liquid was performed using the semi-automated custom-made routine from (Pasquina-Lemonche, et al. "Pore AFM", 2024) where the pore area, later converted in diameter, and the pore number were measured from the 2D slice of each image containing the maximum number of pores (see **Fig. S8**). The number of images similar to images shown in **Fig. 5** were n = 8 for all strains, analyzing both pore size and pore number, the statistical comparisons were also performed using a t test with Welch's correction.

Blue native PAGE. Oligomerization of VirR and Lcp1 was analyzed by Blue native PAGE (Invitrogen, Waltham, MA, United States). Recombinant VirR and Lcp1 proteins were prepared in 1xLB Native PAGE including DDM at 0.5% final concentration. After incubation at 4⁰C for 30 min, samples were centrifuged at 20,000 × g for 30 min at 4°C. The supernatant was recovered and mixed with G-250 additive following manufacturer instructions (Invitrogen) and loaded onto a Novex® Bist-Tris 4-16% gradient gel and run following manufacturer instructions. Gel was first fixed and then stained with Coomassie.

Liposome flotation assay. Individual or combined VirR and Lcp1 were prepared at 2 μM in buffer A in a final volume of 50 μl and incubated for 30 min at 37°C. This volume was mixed with 100 μl of 60% optiprep (Sigma-Aldrich) to a final concentration of 40%

and overlaid with 100 µl of subsequent optiprep solutions at 35%, 30%, 20% and 10%

(optiprep solutions were prepared in Buffer A). The samples were centrifugated at 70000

rpm for 2 h at 4°C in an Optima TLX Ultracentrifuge (Beckman Coulter, Indianapolis, IN

United States). Eleven fractions of 50 µl were recovered and examined by dot blot

(performed as previously described). Polyclonal antibodies raised against VirR and Lcp1

were used as primary antibodies. Quantification of dotblots was performed in ImageJ as

previously described (Schirmer et al., 2022). Briefly, integrated densities were calculated

after background subtraction, and referred to wild type values to obtain the relative

integrated intensities.

1244

1245   **Lysozyme and PG and AG susceptibility assays.** *Mtb* WT and *virR^{mut}* strains were

1246   grown in 7H9 media supplemented with ADC and Tween 80 to an OD 540 of 0.2 and

1247   then diluted to an OD540 of 0.1 with 7H9 containing 0 or 50 µgml^{-1} lysozyme (Sigma).

1248   The cultures were incubated with agitation at 37°C and OD540 was monitored daily.

1249

Fluorescence microscopy. The *Mtb* strain bearing the plasmid VirR-pMV261-Venus

was grown in MM to an OD595nm of 0.4. At this stage, cells were harvested, washed

three times in 1 × PBS (pH 7.4), resuspended in 1/3 – 1/6 volume of 4%

paraformaldehyde (PFA) and incubated for 2 h at 37 $^\circ$C. Finally, the cells were washed

twice in 1 × PBS and resuspended in 100–200 µl of PBS. Microscope slides were

covered with 5 µl poly-L-lysine (Sigma) and excess poly-L-lysine was removed away with

distillated water. A 20 – 40 µl aliquot was applied to pre-treated slides, allowed to air dry

and covered with antifade solution (Prolong Gold, Invitrogen) and visualized under the

fluorescence microscope using a 100x oil immersion objective on an Orca 12-ERG digital

CCD camera (Hamamatsu Photonics) mounted on a Nikon E600 microscope.

1260

FACS analysis for split-GFP experiments. *M. smegmatis* mc²155 transformed with the

pGMCS derivatives were inoculated in complete 7H9 medium. After overnight incubation

at 37°C, bacteria were collected and analyzed by fluorescence activated cell sorting

using a BD FACS LSRFortessa X20 flow cytometer. Flow cytometry data analysis was

performed using FlowJo software (Version 10; Becton, Dickinson and Company,

Ashland, OR, United States). The gating strategy is displayed in supplementary figure

10 (**Fig. S10**).

1268

1269   **Statistical analysis.** The statistical significance of the difference between experimental

1270   groups was determined by the two-tailed Student's test using PRISM 5.0. P values ≤

than 0.05 were considered significant. Statistical analysis of RNAseq is detailed in the corresponding section of Materials and Methods.

Data Availability. The Datasets produced in this study are available in the following database: RNA-seq data: Gene expression Omnibus, GSE143996 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE143996).

Funding Information. R.P.-R. acknowledges support by MINECO/FEDER EU contracts SAF2016–77433-R, PID2019-110240RB-I00, PID2022-136611OB-I00; and NIH-R01AI162821. CICbioGUNE thanks MINECO for the Severo Ochoa Excellence Accreditation (SEV-2016–0644). L.P.-L and J.K.H acknowledges support by the Wellcome Trust (212197/Z/19/Z) which funded part of this research. For the purpose of open access, the authors have applied and will apply a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. J.S. acknowledges support from the Spanish Ministry of Science and Innovation (MICINN) through grant PID2019-106859GA-I00 and Ramón y Cajal research grant RYC-2017-23560, as well as to the Government of Aragón, Spain, through grant B49-23R (NeuroBioSys). FC lab research is supported by the Swedish Research Council (VR2018-02823 and VR2018-05882), the Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, the Knut and Alice Wallenberg Foundation (KAW) grant KAW2012.0184 and the Kempe Foundation.

Acknowledgments. We are very grateful to Carl Nathan for sharing mutant *Mtb* strains. We thank G. Marcela Rodriguez for sharing the a-IdeR antibody. We thank Heran Darwin for sharing a-cei polyclonal antibody. We thank Emmanuelle Näser (Génotoul TRI-IPBS platform, Toulouse) for helping with flow cytometry. We thank Dr. Luke Alderwick (University of Birmingham) for the donation of the plasmid to recombinantly express Lcp1 protein. We thank M. Gutiérrez for assistance with figures. The authors have no conflict of interest to declare.

References

Amaral L, Kristiansen JE, Abebe LS, Millett W. 1996. Inhibition of the respiration of multi-drug resistant clinical isolates of Mycobacterium tuberculosis by thioridazine: Potential use for initial therapy of freshly diagnosed tuberculosis. *J Antimicrob Chemother* **38**. doi:10.1093/jac/38.6.1049

Ariyachaokun K, Grabowska AD, Gutierrez C, Neyrolles O. 2020. Multi-stress induction of the Mycobacterium tuberculosis MbcTA bactericidal toxin-antitoxin system.

Toxins (Basel) **12**. doi:10.3390/toxins12050329

Athman JJ, Sande OJ, Groft SG, Reba SM, Nagy N, Wearsch PA, Richardson ET,
Rojas R, Boom WH, Shukla S, Harding C V. 2017. Mycobacterium tuberculosis
Membrane Vesicles Inhibit T Cell Activation. *J Immunol* **198**:2028–2037.
doi:10.4049/jimmunol.1601199

Athman JJ, Wang Y, McDonald DJ, Boom WH, Harding C V, Wearsch PA. 2015.
Bacterial Membrane Vesicles Mediate the Release of Mycobacterium tuberculosis
Lipoglycans and Lipoproteins from Infected Macrophages. *J Immunol* **195**:1044–
1053. doi:10.4049/jimmunol.1402894

Ballister ER, Samanovic MI, Darwin KH. 2019. Mycobacterium tuberculosis rv2700
contributes to cell envelope integrity and virulence. *J Bacteriol* **201**.
doi:10.1128/JB.00228-19

Bartek IL, Woolhiser LK, Baughn AD, Basaraba RJ, Jacobs WR, Lenaerts AJ, Voskuil
MI. 2014. Mycobacterium tuberculosis Lsr2 is a global transcriptional regulator
required for adaptation to changing oxygen levels and virulence. *MBio* **5**.
doi:10.1128/mBio.01106-14

Bastian M, Heymann S, Jacomy M. 2009. Gephi: An Open Source Software for
Exploring and Manipulating Networks. *Proc Int AAAI Conf Web Soc Media* **3**.
doi:10.1609/icwsm.v3i1.13937

Baumgart M, Schubert K, Bramkamp M, Frunzke J. 2016. Impact of LytRCpsA-Psr
proteins on cell wall biosynthesis in Corynebacterium glutamicum. *J Bacteriol* **198**.
doi:10.1128/JB.00406-16

Beaulieu AM, Rath P, Imhof M, Siddall ME, Roberts J, Schnappinger D, Nathan CF.
2010. Genome-wide screen for Mycobacterium tuberculosis genes that regulate
host immunity. *PLoS One* **5**:e15120. doi:10.1371/journal.pone.0015120

Bera A, Herbert S, Jakob A, Vollmer W, Götz F. 2005. Why are pathogenic
staphylococci so lysozyme resistant? The peptidoglycan O-acetyltransferase OatA
is the major determinant for lysozyme resistance of Staphylococcus aureus. *Mol*
Microbiol **55**. doi:10.1111/j.1365-2958.2004.04446.x

Blondel VD, Guillaume J-L, Lambiotte R, Lefebre E. 2008. Fast unfolding of community
hierarchies in large networks. *J Stat Mech Theory Exp* **10**.

Boldrin F, Anso I, Alebouyeh S, Sevilla IA, Geijo M, Garrido JM, Marina A, Mazzabò
LC, Segafreddo G, Guerin ME, Manganelli R, Prados-Rosales R. 2021. The
phosphatidyl-myo-inositol dimannoside acyltransferase pata is essential for
mycobacterium tuberculosis growth in vitro and in vivo. *J Bacteriol* **203**.
doi:10.1128/JB.00439-20

Boot M, van Winden VJC, Sparrius M, van de Weerd R, Speer A, Ummels R, Rustad T,

Sherman DR, Bitter W. 2017. Cell envelope stress in mycobacteria is regulated by
the novel signal transduction ATPase IniR in response to trehalose. *PLoS Genet*
13:e1007131. doi:10.1371/journal.pgen.1007131

Boudehen YM, Faucher M, Maréchal X, Miras R, Rech J, Rombouts Y, Sénèque O,
Wallat M, Demange P, Bouet JY, Saurel O, Catty P, Gutierrez C, Neyrolles O.
2022. Mycobacterial resistance to zinc poisoning requires assembly of P-ATPase-
containing membrane metal efflux platforms. *Nat Commun* **13**.
doi:10.1038/s41467-022-32085-7

Brzostek A, Pawelczyk J, Rumijowska-Galewicz A, Dziadek B, Dziadek J. 2009.
Mycobacterium tuberculosis is able to accumulate and utilize cholesterol. *J*
Bacteriol **191**. doi:10.1128/JB.00488-09

Cabantous S, Nguyen HB, Pedelacq JD, Koraïchi F, Chaudhary A, Ganguly K, Lockard
MA, Favre G, Terwilliger TC, Waldo GS. 2013. A new protein-protein interaction
sensor based on tripartite split-GFP association. *Sci Rep* **3**.
doi:10.1038/srep02854

Chauhan S, Tyagi JS. 2009. Powerful induction of divergent tgs1-Rv3131 genes in
Mycobacterium tuberculosis is mediated by DevR interaction with a high-affinity
site and an adjacent cryptic low-affinity site. *J Bacteriol* **191**.
doi:10.1128/JB.00310-09

Csardi G, Nepusz T. 2006. The igraph software package for complex network
research. *InterJournal Complex Syst* **Complex Sy**.

De Keijzer J, Mulder A, De Haas PEW, De Ru AH, Heerkens EM, Amaral L, Van
Soolingen D, Van Veelen PA. 2016. Thioridazine alters the cell-envelope
permeability of mycobacterium tuberculosis. *J Proteome Res* **15**.
doi:10.1021/acs.jproteome.5b01037

De Knegt GJ, Ten Kate MT, Van Soolingen D, Aarnoutse R, Boeree MJ, Bakker-
Woudenberg IAJM, De Steenwinkel JEM. 2014. Enhancement of in vitro activity of
tuberculosis drugs by addition of thioridazine is not reflected by improved in vivo
therapeutic efficacy. *Tuberculosis* **94**. doi:10.1016/j.tube.2014.09.002

Dutta NK, Mehra S, Kaushal D. 2010. A Mycobacterium tuberculosis sigma factor
network responds to cell-envelope damage by the promising anti-mycobacterial
thioridazine. *PLoS One* **5**. doi:10.1371/journal.pone.0010069

Ghazaei C. 2018. Mycobacterium tuberculosis and lipids: Insights into molecular
mechanisms from persistence to virulence. *J Res Med Sci*.
doi:10.4103/jrms.JRMS_904_17

Grzegorzewicz AE, de Sousa-d'Auria C, McNeil MR, Huc-Claustre E, Jones V, Petit C,
Angala SK, Zemanova J, Wang Q, Belardinelli JM, Gao Q, Ishizaki Y, Mikusova K,

Brennan PJ, Ronning DR, Chami M, Houssin C, Jackson M. 2016. Assembling of

1383   the Mycobacterium tuberculosis Cell Wall Core. *J Biol Chem* **291**:18867–18879.

1384   doi:10.1074/jbc.M116.739227

1385   Gupta S, Bhagavathula, M Sharma V, Sharma N, Sharma N, Biswas A, Palacios A,

1386   Salgueiro V, Lavín J, Dogra N, Salgame P, Prados-Rosales, R Rodríguez G.

1387   2023. Dynamin-like proteins mediate extracellular vesicle secretion in

1388   Mycobacterium tuberculosis. *EMBO Rep* **24**:e55593.

1389   Harrison J, Lloyd G, Joe M, Lowary TL, Reynolds E, Walters-Morgan H, Bhatt A,

1390   Lovering A, Besra GS, Alderwick LJ. 2016. Lcp1 is a phosphotransferase

1391   responsible for ligating arabinogalactan to peptidoglycan in mycobacterium

1392   tuberculosis. *MBio* **7**. doi:10.1128/mBio.00972-16

1393   Hasenoehrl EJ, Rae Sajorda D, Berney-Meyer L, Johnson S, Tufariello JAM, Fuhrer T,

1394   Cook GM, Jacobs WR, Berney M. 2019. Derailing the aspartate pathway of

1395   Mycobacterium tuberculosis to eradicate persistent infection. *Nat Commun* **10**.

1396   doi:10.1038/s41467-019-12224-3

1397   Honaker RW, Leistikow RL, Bartek IL, Voskui MI. 2009. Unique roles of DosT and

1398   DosS in DosR regulon induction and Mycobacterium tuberculosis dormancy. *Infect*

1399   *Immun* **77**. doi:10.1128/IAI.01449-08

1400   Huber W, Heydebreck A Von, Ultmann HS¨, Poustka A, Vingron M. 2002. Variance

1401   stabilization applied to microarray data calibration and to the quantification of

1402   differential expression. *BIOINFORMATICS*.

1403   Hubscher J, Luthy L, Berger-Bachi B, Stutzmann Meier P. 2008. Phylogenetic

1404   distribution and membrane topology of the LytR-CpsA-Psr protein family. *BMC*

1405   *Genomics* **9**:617. doi:10.1186/1471-2164-9-617

1406   Justen AM, Hodges HL, Kim LM, Sadecki PW, Porfirio S, Ultee E, Black I, Chung GS,

1407   Briegel A, Azadi P, Kiessling LL. 2020. Polysaccharide length affects

1408   mycobacterial cell shape and antibiotic susceptibility. *Sci Adv* **6**.

1409   doi:10.1126/sciadv.aba4015

1410   Kalscheuer R, Palacios A, Anso I, Cifuente J, Anguita J, Jacobs Jr. WR, Guerin ME,

1411   Prados-Rosales R. 2019. The Mycobacterium tuberculosis capsule: a cell

1412   structure with key implications in pathogenesis. *Biochem J* **476**:1995–2016.

1413   doi:10.1042/BCJ20190324

1414   Kawai Y, Marles-Wright J, Cleverley RM, Emmins R, Ishikawa S, Kuwano M, Heinz N,

1415   Bui NK, Hoyland CN, Ogasawara N, Lewis RJ, Vollmer W, Daniel RA, Errington J.

1416   2011. A widespread family of bacterial cell wall assembly proteins. *EMBO J*

1417   **30**:4931–4941. doi:10.1038/emboj.2011.358

Kuru E, Radkov A, Meng X, Egan A, Alvarez L, Dowson A, Booher G, Breukink E,

Roper DI, Cava F, Vollmer W, Brun Y, Vannieuwenhze MS. 2019. Mechanisms of
Incorporation for D -Amino Acid Probes That Target Peptidoglycan Biosynthesis.
ACS Chem Biol **14**. doi:10.1021/acschembio.9b00664
Layre E, Sweet L, Hong S, Madigan CA, Desjardins D, Young DC, Cheng TY, Annand
JW, Kim K, Shamputa IC, McConnell MJ, Debono CA, Behar SM, Minnaard AJ,
Murray M, Barry CE, Matsunaga I, Moody DB. 2011. A comparative lipidomics
platform for chemotaxonomic analysis of mycobacterium tuberculosis. *Chem Biol*
18. doi:10.1016/j.chembiol.2011.10.013
Lee JE, Moon PG, Lee IK, Baek MC. 2015. Proteomic Analysis of Extracellular
Vesicles Released by Adipocytes of Otsuka Long-Evans Tokushima Fatty
(OLETF) Rats. *Protein J* **34**:220–235. doi:10.1007/s10930-015-9616-z
Lee W, VanderVen BC, Fahey RJ, Russell DG. 2013. Intracellular Mycobacterium
tuberculosis exploits host-derived fatty acids to limit metabolic stress. *J Biol Chem*
288:6788–6800. doi:10.1074/jbc.M112.445056
Li C, Wen A, Shen B, Lu J, Huang Y, Chang Y. 2011. FastCloning: A highly simplified,
purification-free, sequence- and ligation-independent PCR cloning method. *BMC*
Biotechnol **11**. doi:10.1186/1472-6750-11-92
Liao Y, Smyth GK, Shi W. 2019. The R package Rsubread is easier, faster, cheaper
and better for alignment and quantification of RNA sequencing reads. *Nucleic*
Acids Res **47**. doi:10.1093/nar/gkz114
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**. doi:10.1186/s13059-
014-0550-8
Lu R, Schaefer CM, Nesbitt NM, Kuper J, Kisker C, Sampson NS. 2017. Catabolism of
the Cholesterol Side Chain in Mycobacterium tuberculosis Is Controlled by a
Redox-Sensitive Thiol Switch. *ACS Infect Dis* **3**. doi:10.1021/acsinfecdis.7b00072
Maciąg A, Dainese E, Rodriguez GM, Milano A, Provvedi R, Pasca MR, Smith I, Palù
G, Riccardi G, Manganelli R. 2007. Global analysis of the Mycobacterium
tuberculosis Zur (FurB) regulon. *J Bacteriol* **189**. doi:10.1128/JB.01190-06
Majlessi L, Prados-Rosales R, Casadevall A, Brosch R. 2015. Release of
mycobacterial antigens. *Immunol Rev* **264**:25–45. doi:10.1111/imr.12251
Marsollier L, Brodin P, Jackson M, Kordulakova J, Tafelmeyer P, Carbonnelle E, Aubry
J, Milon G, Legras P, Andre JP, Leroy C, Cottin J, Guillou MLJ, Reysset G, Cole
ST, Korduláková J, Tafelmeyer P, Carbonnelle E, Aubry J, Milon G, Legras P,
Saint André JP, Leroy C, Cottin J, Guillou MLJ, Reysset G, Cole ST. 2007. Impact
of Mycobacterium ulcerans biofilm on transmissibility to ecological niches and
Buruli ulcer pathogenesis. *PLoS Pathog* **3**:e62. doi:10.1371/journal.ppat.0030062

McMillan HM, Kuehn MJ. 2021. The extracellular vesicle generation paradox: a
bacterial point of view. *EMBO J* **40**. doi:10.15252/embj.2021108174

Minch KJ, Rustad TR, Peterson EJR, Winkler J, Reiss DJ, Ma S, Hickey M, Brabant W,
Morrison B, Turkarslan S, Mawhinney C, Galagan JE, Price ND, Baliga NS,
Sherman DR. 2015. The DNA-binding network of Mycobacterium tuberculosis. *Nat*
Commun **6**. doi:10.1038/ncomms6829

Mohammadzadeh R, Ghazvini K, Farsiani H, Soleimanpour S. 2021. Mycobacterium
tuberculosis extracellular vesicles: exploitation for vaccine technology and
diagnostic methods. *Crit Rev Microbiol*. doi:10.1080/1040841X.2020.1830749

Nagakubo T, Tahara YO, Miyata M, Nomura N, Toyofuku M. 2021. Mycolic acid-
containing bacteria trigger distinct types of membrane vesicles through different
routes. *iScience* **24**. doi:10.1016/j.isci.2020.102015

Nagamani S, Sastry GN. 2021. Mycobacterium tuberculosis cell wall permeability
model generation using chemoinformatics and machine learning approaches. *ACS*
Omega **6**. doi:10.1021/acsomega.1c01865

Nečas D, Klapetek P. 2012. Gwyddion: An open-source software for SPM data
analysis. *Cent Eur J Phys*. doi:10.2478/s11534-011-0096-2

Oba S, Sato MA, Takemasa I, Monden M, Matsubara KI, Ishii S. 2003. A Bayesian
missing value estimation method for gene expression profile data. *Bioinformatics*
19:2088–2096. doi:10.1093/bioinformatics/btg287

Pandey R, Rodriguez GM. 2014. IdeR is required for iron homeostasis and virulence in
Mycobacterium tuberculosis. *Mol Microbiol* **91**. doi:10.1111/mmi.12441

Park HD, Guinn KM, Harrell MI, Liao R, Voskuil MI, Tompa M, Schoolnik GK, Sherman
DR. 2003. Rv3133c/dosR is a transcription factor that mediates the hypoxic
response of Mycobacterium tuberculosis. *Mol Microbiol* **48**. doi:10.1046/j.1365-
2958.2003.03474.x

Pasquina-Lemonche L, Burns J, Turner RD, Kumar S, Tank R, Mullin N, Wilson JS,
Chakrabarti B, Bullough PA, Foster SJ, Hobbs JK. 2020. The architecture of the
Gram-positive bacterial cell wall. *Nature* **582**. doi:10.1038/s41586-020-2236-6

Pawełczyk J, Brzostek A, Minias A, Płociński P, Rumijowska-Galewicz A, Strapagiel D,
Zakrzewska-Czerwińska J, Dziadek J. 2021. Cholesterol-dependent transcriptome
remodeling reveals new insight into the contribution of cholesterol to
Mycobacterium tuberculosis pathogenesis. *Sci Rep* **11**. doi:10.1038/s41598-021-
91812-0

Peterson EJ, Bailo R, Rothchild AC, Arrieta-Ortiz ML, Kaur A, Pan M, Mai D, Abidi AA,
Cooper C, Aderem A, Bhatt A, Baliga NS. 2019. Path-seq identifies an essential
mycolate remodeling program for mycobacterial host adaptation. *Mol Syst Biol* **15**.

doi:10.15252/msb.20188584

Peterson EJR, Brooks AN, Reiss DJ, Kaur A, Do J, Pan M, Wu WJ, Morrison R,

Srinivas V, Carter W, Arrieta-Ortiz ML, Ruiz RA, Bhatt A, Baliga NS. 2023. MtrA

modulates Mycobacterium tuberculosis cell division in host microenvironments to

mediate intrinsic resistance and drug tolerance. *Cell Rep* **42**.

doi:10.1016/j.celrep.2023.112875

Prados-Rosales R, Baena A, Martinez LR, Luque-Garcia J, Kalscheuer R,

Veeraraghavan U, Camara C, Nosanchuk JD, Besra GS, Chen B, Jimenez J,

Glatman-Freedman A, Jacobs Jr. WR, Porcelli SA, Casadevall A, Jacobs WR,

Porcelli SA, Casadevall A. 2011. Mycobacteria release active membrane vesicles

that modulate immune responses in a TLR2-dependent manner in mice. *J Clin*

Invest **121**:1471–1483. doi:10.1172/JCI44261

Prados-Rosales R, Carreno L, Cheng T, Blanc C, Weinrick B, Malek A, Lowary TL,

Baena A, Joe M, Bai Y, Kalscheuer R, Batista-Gonzalez A, Saavedra NA,

Sampedro L, Tomas J, Anguita J, Hung SC, Tripathi A, Xu J, Glatman-Freedman

A, Jacobs Jr. WR, Chan J, Porcelli SA, Achkar JM, Casadevall A. 2017.

Enhanced control of Mycobacterium tuberculosis extrapulmonary dissemination in

mice by an arabinomannan-protein conjugate vaccine. *PLoS Pathog*

13:e1006250. doi:10.1371/journal.ppat.1006250

Prados-Rosales R, Carreño LJ, Batista-Gonzalez A, Baena A, Venkataswamy MM, Xu

J, Yu X, Wallstrom G, Mitchell Magee D, LaBaer J, Achkar JM, Jacobs WR, Chan

J, Porcelli SA, Casadevall A. 2014a. Mycobacterial membrane vesicles

administered systemically in mice induce a protective immune response to surface

compartments of mycobacterium tuberculosis. *MBio* **5**:e01921-14.

doi:10.1128/mBio.01921-14

Prados-Rosales R, Weinrick BC, Pique DG, Jacobs Jr. WR, Casadevall A, Rodriguez

GM, Piqué DG, Jacobs WR, Casadevall A, Rodriguez GM. 2014b. Role for

Mycobacterium tuberculosis membrane vesicles in iron acquisition. *J Bacteriol*

196:1250–1256. doi:10.1128/JB.01090-13

Prasad TSK, Verma R, Kumar S, Nirujogi RS, Sathe GJ, Madugundu AK, Sharma J,

Puttamallesh VN, Ganjiwale A, Myneedu VP, Chatterjee A, Pandey A, Harsha HC,

Narayana J. 2013. Proteomic analysis of purified protein derivative of

mycobacterium tuberculosis. *Clin Proteomics* **10**. doi:10.1186/1559-0275-10-8

Rath P, Huang C, Wang T, Wang T, Li H, Prados-Rosales R, Elemento O, Casadevall

A, Nathan CF. 2013. Genetic regulation of vesiculogenesis and

immunomodulation in Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A*

110:E4790-7. doi:10.1073/pnas.1320118110

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. Limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**:e47. doi:10.1093/nar/gkv007

Rock JM, Hopkins FF, Chavez A, Diallo M, Chase MR, Gerrick ER, Pritchard JR, Church GM, Rubin EJ, Sassetti CM, Schnappinger D, Fortune SM. 2017. Programmable transcriptional repression in mycobacteria using an orthogonal CRISPR interference platform. *Nat Microbiol* **2**:16274. doi:10.1038/nmicrobiol.2016.274

Sani M, Houben ENG, Geurtsen J, Pierson J, De Punder K, Van Zon M, Wever B, Piersma SR, Jiménez CR, Daffé M, Appelmelk BJ, Bitter W, Van Wel N Der, Peters PJ. 2010. Direct visualization by Cryo-EM of the mycobacterial capsular layer: A labile structure containing ESX-1-secreted proteins. *PLoS Pathog* **6**. doi:10.1371/journal.ppat.1000794

Sartain MJ, Dick DL, Rithner CD, Crick DC, Belisle JT. 2011. Lipidomic analyses of Mycobacterium tuberculosis based on accurate mass measurements and the novel "Mtb LipidDB." *J Lipid Res* **52**. doi:10.1194/jlr.M010363

Schirmer S, Rauh L, Alebouyeh S, Delgado-Velandia M, Salgueiro VC, Lerma L, Serrano-Mestre JL, Azkargorta M, Elortza F, Lavín JL, García MJ, Tórtola-Fernández MT, Gola S, Prados-Rosales R. 2022. Immunogenicity of Mycobacterial Extracellular Vesicles Isolated From Host-Related Conditions Informs About Tuberculosis Disease Status. *Front Microbiol* **13**:907296. doi:10.3389/fmicb.2022.907296

Schnappinger D, Ehrt S, Voskuil MI, Liu Y, Mangan JA, Monahan IM, Dolganov G, Efron B, Butcher PD, Nathan C, Schoolnik GK. 2003. Transcriptional adaptation of Mycobacterium tuberculosis within macrophages: Insights into the phagosomal environment. *J Exp Med* **198**:693–704. doi:10.1084/jem.20030846

Soto-Ramirez MD, Aguilar-Ayala DA, Garcia-Morales L, Rodriguez-Peredo SM, Badillo-Lopez C, Rios-Muñiz DE, Meza-Segura MA, Rivera-Morales GY, Leon-Solis L, Cerna-Cortes JF, Rivera-Gutierrez S, Helguera-Repetto AC, Gonzalez-y-Merchand JA. 2017. Cholesterol plays a larger role during Mycobacterium tuberculosis in vitro dormancy and reactivation than previously suspected. *Tuberculosis* **103**. doi:10.1016/j.tube.2016.12.004

Storey JD, Bass AJ, Dabney A, Robinson D. 2023. qvalue: Q-value estimation for false discovery rate control.

Stupar M, Tan L, Kerr ED, De Voss CJ, Forde BM, Schulz BL, West NP. 2024. TcrXY is an acid-sensing two-component transcriptional regulator of Mycobacterium tuberculosis required for persistent infection. *Nat Commun* **15**.

doi:10.1038/s41467-024-45343-7

Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: Discovering splice junctions with

RNA-Seq. *Bioinformatics* **25**. doi:10.1093/bioinformatics/btp120

Tsuruoka T, Tamura A, Miyata A, Takei T, Inouye S, Matsuhashi M. 1985. Second

Lytic Target of Beta-Lactam Compounds That Have A Terminal d-Amino Acid

Residue. *Eur J Biochem* **151**. doi:10.1111/j.1432-1033.1985.tb09090.x

Vilchèze C, Jacobs WR. 2007. Isolation and analysis of Mycobacterium tuberculosis

mycolic acids. *Curr Protoc Microbiol* **Chapter 10**.

doi:10.1002/9780471729259.mc10a03s05

Wang M, Guo X, Yang X, Zhang B, Ren J, Liu A, Ran Y, Yan B, Chen F, Guddat LW,

Hu J, Li J, Rao Z. 2019. Mycobacterial dynamin-like protein IniA mediates

membrane fission. *Nat Commun* **10**. doi:10.1038/s41467-019-11860-z

White DW, Elliott SR, Odean E, Bemis LT, Tischler AD. 2018. Mycobacterium

tuberculosis Pst/SenX3-RegX3 Regulates Membrane Vesicle Production

Independently of ESX-5 Activity. *MBio* **9**. doi:10.1128/mBio.00778-18

Williams EP, Lee JH, Bishai WR, Colantuoni C, Karakousis PC. 2007. Mycobacterium

tuberculosis SigF regulates genes encoding cell wall-associated proteins and

directly regulates the transcriptional regulatory gene phoY1. *J Bacteriol* **189**.

doi:10.1128/JB.00201-07

Wiśniewski JR, Zougman A, Nagaraj N, Mann M. 2009. Universal sample preparation

method for proteome analysis. *Nat Methods* **6**. doi:10.1038/nmeth.1322

Zhang X, Smits AH, Tilburg GBA Van, Ovaa H, Huber W, Vermeulen M. 2018.

Proteome-wide identification of ubiquitin interactions using UbIA-MS. *Nat Protoc*

13:530–550. doi:10.1038/nprot.2017.147

Ziegenbalg A, Prados-Rosales R, Jenny-Avital ER, Kim RS, Casadevall A, Achkar JM.

2013. Immunogenicity of mycobacterial vesicles in humans: identification of a new

tuberculosis antibody biomarker. *Tuberc* **93**:448–455.

doi:10.1016/j.tube.2013.03.001

Zuber B, Haenni M, Ribeiro T, Minnig K, Lopes F, Moreillon P, Dubochet J. 2006.

Granular layer in the periplasmic space of gram-positive bacteria and fine

structures of Enterococcus gallinarum and Streptococcus gordonii septa revealed

by cryo-electron microscopy of vitreous sections. *J Bacteriol* **188**.

doi:10.1128/JB.00391-06

1600

1601

1602

1603

Figure Legends

Figure 1. Ultrastructural changes in the cell envelope of *virR*-deficient mutant associated with increased vesiculation. (**A**) Cryo-electron micrographs of indicated *Mtb* strains grown in high iron MM. Closed line rectangles were used to calculate grey value profiles of cell envelopes using ImageJ. The dashed line insets within the main micrograph were magnified to show a detailed view of the cell surface. Scale bars are 100 nm in main micrographs and 50 nm in the insets. (**B**) Density profiles based on grey values of the cross sections marked by solid line rectangles in A. (**C**) Mean values and standard errors of distances in nm between main cell envelope layers measured in A and C. *P < 0.05 after applying a Tukey´s multiple comparison test. The number of cells analyzed varied from n = 20 (WT), n = 15 (*virR^mut^*) and n = 23 (*virR^mut^*-C). CM, cytoplasmic membrane; OM, outer membrane; L1, layer 1; L2, layer 2.

Figure 2. Enhanced permeability and vesiculation are linked in the absence of LytR_C solo domain proteins in *Mtb*. (A) Uptake of ethidium bromide (EthBr) in the indicated strains, as measured by fluorescence at 590 nm for 65 min. Data are mean and standard deviation of three biological replicates. **(B)** Sensitivity of the indicated strains to different concentrations of vancomycin as measured by monitoring optical density at 580 nm (OD580) after 7 days. Data are mean and standard deviations of three biological replicated. Errors bars not shown are smaller than symbols. **(C)** Time course of the incorporation of FDAAs on the indicated strains in 7H9 as measured by fluorescence. Cells were labeled with HADA for 48 h, washed in 7H9 medium and labeled with FDL for 5 h. Then, cells were fixed and imaged under a fluorescence microscope. Representative images of WT, *virR^mut^* and *virR^mut^*-Comp strains in 7H9. Data are mean and standard deviations of three biological replicates. *** means *P*<0.001 and **** means *P*<0.0001 after applying a One-way ANOVA analysis. **(D)** immunoblot analysis of cell lysates of two independent conditional mutants (c1 and c2) for *virR* and *cei* for the presence of VirR and Cei proteins in cultures incubated with and without anhydrotetracycline (ATc). **(E)** Nanoparticle tracking analysis (NTA) of EV preparations derived from *virR* and *cei* conditional mutants (c2 (*virR*) and c1 (*cei*)) obtained from cultures with and without ATc showing number of particles per cm³ determined by Zeta View NTA in three independent EV preparations. Data are presented as mean ± SEM.

Figure 3. Transcriptional profiling of *virR^mut^* showcases systemic alterations in cell wall architecture and metabolism. (A) Experiment design for RNA-seq analyses. RNA was extracted from whole cell extracts from 4 replicate cultures of each strain (WT,

virR^mut, and *virR^mut* -Comp), and sequenced. **(B)** PCA plot of the transcriptomics data.
The loadings from the two first principal components are plotted. **(C)** Number of
differentially expressed genes between *virR^mut* and either WT (left) or *virR^mut*-Comp
(right). **(D)** Enrichment profiles between sets of genes found up and down-regulated for
the contrasts *virR^mut* vs WT and *virR^mut* vs *virR^mut*-Comp. While genes that are
differentially expressed in the same direction in both contrasts are mutually enriched,
genes whose differential expression goes in different directions in both comparisons are
mutually depleted (two-sided Fisher exact test p<0.05 in all four cases). **(E)** Gene
Ontology (GO) enrichments of genes that are 1) simultaneously downregulated in *virR^mut*
with respect to both WT and v *virR^mut*-Comp (left) and 2) simultaneously upregulated
(right). GO terms selected for testing included biological process and cell compartment
labels containing N > 10 genes, between levels 4-6 of the ontology tree. In this
representation, each node in the networks represents a significantly enriched term (terms
selected for visualization with FDR=10% and enrichment odds-ratio>2), nodes' size is
proportional to the enrichment significance, and links between terms are added wherever
the sets of genes contributing to the enrichments in a connected pair show an
intersection larger than 50% of the smallest of the gene sets involved in the pair (see
Methods for further details, and supplementary file S2 for an extended list of all the
enrichments found).

Figure 4. VirR-dependent genes intersect the regulons of key transcriptional
regulators of the responses to stress, dormancy, and cell wall remodeling. (A)
Transcription factors controlling regulons significantly enriched among genes that are 1)
simultaneously downregulated in *virR^mut* with respect to both WT and *virR^mut* -Comp
(bottom, 9 TFs) and 2) simultaneously upregulated (top, 1 TF). **(B)** Differential expression
patterns for enriched TFs whose expression is VirR-dependent. **(C)** Regulatory
subnetwork including the genes under transcriptional control of the enriched TFs
highlighted in figure 4A that were found to be VirR-dependent. In this visualization, color
captures the logFC for the contrast *virR^mut*-WT, and the size of its FDR.

1669

Figure 5. Cell permeability is a major driver of vesicle production in *Mtb*. (A) Time
course of the uptake of ethidium bromide (EthBr) by *Mtb* grown in the indicated
conditions, as measured by fluorescence at 590=nm for 65 min. Data are mean and
standard deviation of three biological replicates. **(B)** Dot-blot analysis of released EVs in
supernatants from *Mtb* cultures submitted to the indicated conditions. The loading
volume was normalized according to CFUs. The graph below represents the
quantification of the dotblot using ImageJ. The integrated cell density was normalized to
WT values. Data are mean and standard errors from the three biological replicates; ****

denotes P<0.0001, *** denotes P=0.0028 after applying a Tuke's multiple comparison

test. **(C)** Enrichment odds ratios for Fisher exact tests comparing sets of genes up, and

down regulated, respectively, in 1) cholesterol-supplemented medium vs. glucose-

supplemented medium (Pawełczyk et al., 2021), GEO accession GSE175812); and 2)

virR^{mut} vs WT as well as *virR^{mut}* vs *virR^{mut}*-Comp. **(D)** Euler diagram showing the

intersection between A) genes upregulated by cholesterol, B) genes down-regulated in

virR^{mut}, and C) the DevR regulon. **(E)** Expression patterns in response to TRZ (1h) and

in *virR^{mut}* for genes involved in metal signaling, metabolism and homeostasis.

1686

Figure 6. VirR regulates the protein content of EVs in *Mtb*. (A). Experiment design

for proteomics analyses. label-free mass spectrometry data was retrieved for replicates

of H37RV (WT), *virR^{mut}* and *virR^{mut}*-Comp. For each strain, data corresponding to

proteins extracted from either whole-cell extracts (WC) and EVs was produced, totaling

12 samples (2 strains times 2 cell compartments times 3 replicates). **(B)** PCA plot of the

proteomics data. The loadings from the two first principal components, adding up to

50.5% of variance explained, are plotted. **(C)** Expression patterns of VirR in the

proteomics dataset across cell compartments and strains. As expected, while *virR^{mut}*

shows significantly lower levels of VirR protein, the complemented strain does not just

restore WT levels of expression, but leads to a strong overexpression of VirR, both in

the lysates and the EVs, **(D)** Bar plots of differentially expressed proteins found between

strains in EVs (up: *virR^{mut}* vs WT; down *virR^{mut}* vs *virR^{mut}*-Comp at 5% FDR). **(E)**

Enrichment profiles between sets of proteins found up and down-regulated for the

contrasts *virR^{mut}* vs WT and *virR^{mut}* vs *virR^{mut}*-Comp. **(F)** Gene Ontology enrichments of

the downregulated proteins in the VirR mutant strain (left) and the upregulated ones

(Right), at the EV location. GO terms selected for testing included biological process and

cell compartment labels containing N > 3 genes, between levels 4-6 of the ontology tree.

As in **figure 3E**, terms selected for visualization with FDR=10% and enrichment odds-

ratio>2, nodes' size is proportional to the enrichment significance, and links between

terms are added wherever the sets of proteins contributing to the enrichments in a

connected pair shows an intersection larger than 50% of the smallest of the gene sets

involved in the pair (see Methods for further details, and supplementary file S5 for an

extended list of all the enrichments found). **(G)** Expression patterns of bacterioferritin

proteins in the EVs of WT and *virR^{mut}*. **(H)** Expression patterns of proteins contributing to

enrichments of terms related to mycolic acids biosynthesis. **(I)** Bottom left:

Concentrations of key metabolites in the propionyl-coa detoxification routes. Right:

Metabolic pathway of the propionyl-coa detoxification route through the methyl-malonyl

route, along with the differential expression statistics between the WT and *virR^{mut}* strains

observed at transcriptomic and proteomic levels for the main enzymes involved. The

framed plot represents the quantification of selected metabolites by mass spectrometry.

1717

Figure 7. Lipidomic analysis of the cell envelope and EVs from *virR^mut^*. (A)

Comparative total lipid analysis of the indicated strains by nano-LC-MS. Lipid species

with an abundance higher than 0.5% are shown. Data indicates mean and standard error

from three biological replicates. **(B)** Analysis of polar lipids of the indicated strains by

bidimensional (2D)-layer chromatography (TLC) (2D-TLC). Phosphatidyl-myo-inositol

dimannosides (Ac$_1$PIM$_2$ and Ac$_2$PIM$_2$, respectively); phosphatidyl-myo-inositol

hexamannosides (Ac$_x$PIM$_6$); phosphatidylethanolamine (PE); diphosphatidylglycerol

(DPG). Phospholipid (P). **(C)** Analysis of mycolic acid species in the indicated strains by

TLC. **(D)** Analysis of PIDMs and TAGs by TLC. **(E)** Analysis of apolar lipids of the

indicated strains by TLC. Diacylglycerol (DAG); free fatty acid (FF); free mycolic acid

(FMA). **(F)** Analysis of apolar lipids of MEVs isolated from the indicated strains.

Triacylglycerols (TAGs); phthiocerol dimycocerosates (PDIMs) **(G)** Analysis of polar

lipids of MEVs isolated from the indicated strains. Phosphatidyl-myo-inositol

dimannosides (Ac$_1$PIM$_2$ and Ac$_2$PIM$_2$, respectively); phosphatidyl-myo-inositol

hexamannosides (Ac$_x$PIM$_6$); diphosphatidylglycerol (DPG).

1733

Figure 8. Chemical and nanometric analysis of isolated cell walls shows higher

amounts of PG in the absence of VirR. (A) Glycosyl composition analysis of isolated

cell walls from the indicated strains. Acid hydrolysis prior to GC-MS analysis of isolated

cell walls was performed to determine the total amounts of galactose (G), arabinose (A),

rhamnose (R), and N-acetylglucosamine (NAGc). The ratio between different sugars is

shown to indicate: (i) the relative levels of arabinogalactan (A/G); (ii) the length of the AG

polysaccharide chain (G/R); (iii) the amount of PG (NAGc/Rham). Individual

measurements are shown from three biological replicates. Data are mean and standard

errors. **P < 0.01 after applying a Tukey´s multiple comparison test. **(B)** Quantification

of diaminopimelic acid (DAP) on isolated cell walls from the indicated strains. Data are

mean and standard errors from three biological replicates; **** denotes P<0.0001, *

denotes P$=0.0119$ after applying a Tukey´s multiple comparison test. **(C)** Atomic force

microscopy high resolution images taken in liquid from the external surface of purified

peptidoglycan from different samples, left to right: WT, *virR^mut^*, *virR^mut^-C*, the structure of

external peptidoglycan layer shows a mesh with pores of different sizes, **(D)** Graph

showing the pore diameter from n = 8 AFM images similar to **(C)** (each point represent

an image) of different samples; black triangles: WT, blue circles: *virR^mut^*, red squares:

virR^mut^-C. The pore diameter was calculated from the binary slice from each image

containing the greatest number of pores, according to (Pasquina-Lemonche, 2024). **(E)**
The number of pores per surface area from different samples; white triangles: WT, red
circles: *virR^{mut}*, blue squares: *virR^{mut}*-C, this was calculated from the slice containing the
maximum number of pores where the pore size was analyzed, following the method from
(Pasquina-Lemonche, 2024). **(F)** Graph showing the peptidoglycan thickness manually
measured using the 1D statistic tools from the open-source program Gwyddion (Nečas
and Klapetek, 2012) from AFM images in air containing several fragments of cell wall,
each point represents a different peptidoglycan fragment from a different cell. There were
three samples analyzed: black WT n=23 peptidoglycan fragments, blue *virR^{mut}* n=36 and
red *virR^{mut}*-C n=25. Statistics: D) Using a two-tailed t test with Welch's correction the
statistical comparison are: (***) pWT- *virR^{mut}* = 4.1 10^{-4} with t = 6.1, df = 7.5, (**) p *virR^{mut}*
- *virR^{mut}*-C = 0.007 with t = 3.2, df = 13.2, E) Using a two-tailed t test with Welch's
correction the statistical comparison are: (****) pWT- *virR^{mut}* = 6.6 10-10 with t = 16.3,
df = 12.7, (**) p *virR^{mut}* - *virR^{mut}* -C = 0.01 with t = 3.0, df = 13.2., F) Using a two-tailed t
test with Welch's correction the statistical comparison are: (****) pWT- *virR^{mut}* = 2.8 10-
18 with t = 17.8, df = 32.3, (****) *virR^{mut}* - *virR^{mut}* -C = 3.9 10-17 with t = 16.4, df = 31.8.
(G) List of muropeptides identified in the cell walls of Mtb. Features including retention
time (RT), observed and expected mass as well as the best match for each peak are
shown. Those deacetylated muropeptides are indicated in red. **(H)** Relative abundance
of deacetylated muropeptides in the indicated strains.
1772
Figure 9. VirR interacts with Lcp1. (A) Blue-native PAGE analysis of recombinant VirR
and Lcp1. The molecular mass of markers in kDa are indicated on the left side of the gel.
The size of the multimers of VirR and Lcp1 are indicated on the left side of the gel. **(B)**
Upper panel. Representative image of a flotation assay of recombinant VirR alone or in
combination with Lcp1 performed on an idioxanol density gradient. The presence of VirR
was detected in each fraction by dot-blot using murine polyclonal antibodies raised
against VirR. Numbers indicate collected fractions from top (1) to bottom (11). *Lower*
panel. Quantification of the dot-blot by measuring the pixel intensity of the dots using
ImageJ. Data are mean and standard error of three independent experiments. **(C)**
Bipartite split-GFP experiment using *M. smegmatis* D2 expressing plasmids depicted in
(**Fig S11A**), including CtpC_{Nterm}, VirR, VirRsol (Δ1-41), Rv3484, Rv3267 (Lcp1), Rv0822,
Rv3484_{LytR_C}, Rv3267_{LytR_C} and Rv0822_{LytR_C}. Bacteria were grown in complete 7H9 and
fluorescence was recorded by FACS. Pooled data obtained from four to six independent
cultures of each strain are shown. Statistical analysis was performed using non-
parametric two-sided Mann-Whitney test. ns: not significant. * and ** refer to p-values
<0.05 or 0.01, respectively.

1789
1790 **Supplementary Tables**

1791
1792 **Supplementary Table S1.** Plasmids and primers used in this study.

Construction / DNA fragment used	Template	Primers	Sequence 5'-3'
pGMCS-P1-*virR*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)			
P1	pGMCS-TetR-P1-*pacL1*$_{Flag}$- P$_{pacL1}$-*ctpC*$_{His}$ (Boudehen *et al.* 2022)	Inf-B2-P1-Fw	CAAAGTGGAGCGATACTCGACATCGGATCGTCGGCACCGTCAC
		Inf-P1-Left	GAAATGATGTATGCCGTGCTGGTC
pGMCS backbone	pGMCS-P$_{pacL}$-*pacL1*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10) (Boudehen *et al.* 2022)	ctpC-B2-Left	GTCGAGTATCGCTCCACTTTG
		Rv3269-P1-Fus-Fw	CAGCACGGCATACATCATTTCGGAAGAAAGGTACAGGCAATGG
pGMCS-P1-*virR*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)			
virR	*M. tuberculosis* H37Rv chromosomal DNA	If-AmL1-VirR	AAGGTACAGGCAATGgtgctggttacagtgggctcg
		If-Lk2-VirR-Rv	ACCGGATCCTTCCAGgccggtcaccacgacgatg
pGMCS backbone	pGMCS-P1-*pacL1*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Lk2-Right	CTGGAAGGATCCGGTGGCGG
		pGMC 3269 Am Left	CATTGCCTGTACCTTTCTTCC
pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)			
virRsol	*M. tuberculosis* H37Rv chromosomal DNA	If-AmL1-VirRsol	AAGGTACAGGCAATGctgggttcgtctccgaactccg
		If-Lk2-VirR-Rv	ACCGGATCCTTCCAGgccggtcaccacgacgatg
pGMCS backbone	pGMCS-P1-*pacL1*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Lk2-Right	CTGGAAGGATCCGGTGGCGG
		pGMC 3269 Am Left	CATTGCCTGTACCTTTCTTCC
pGMCS-P1-*virR*-Lk2-GFP11-*virR*-Lk15-GFP(1-10)			
virR	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-VirR	GAGCGCCTCGCCATGgtgctggttacagtgggctcg
		If-L15-VirR-Rv	AGACAGCCCTGGACCgccggtcaccacgacgatg
pGMCS backbone	pGMCS-P1-*virR*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*virRsol*-Lk15-GFP(1-10)			
virRsol	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-VirRsol	GAGCGCCTCGCCATGctgggttcgtctccgaactccg
		If-L15-VirR-Rv	AGACAGCCCTGGACCgccggtcaccacgacgatg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv0822c*(LytR_C)-Lk15-GFP(1-10)			
Rv0822c(LytR_C)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-0822cD	GAGCGCCTCGCCATGgagattcagcaccagcaggttacgacg
		If-L15-0822cD-Rv	AGACAGCCCTGGACCgcggcttatctgcacgctgacg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3267*(LytR_C)-Lk15-GFP(1-10)			
Rv3267(LytR_C)		If-AmMBD-3267D	GAGCGCCTCGCCATGcccgccaagaccacggcc

	M. tuberculosis H37Rv chromosomal DNA	If-L15-3267D-Rv	AGACAGCCCTGGACCcagcccggagcccggac
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
colspan="4"	pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3484*(LytR_C)-Lk15-GFP(1-10)
Rv3484(LytR_C)	M. tuberculosis H37Rv chromosomal DNA	If-AmMBD-3484D	GAGCGCCTCGCCATGgcggccgccaaacctaaccc
		If-L15-3484D-Rv	AGACAGCCCTGGACCggtatccaccgtcacacggatg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
colspan="4"	pGMCS-P1-*virRsol*-Lk2-GFP11-*rv0822c*(Cterm)-Lk15-GFP(1-10)
Rv0822c(Cterm)	M. tuberculosis H37Rv chromosomal DNA	If-AmMBD-0822cD	GAGCGCCTCGCCATGgagattcagcaccagcaggttacgacg
		If-L15-0822c-Rv2	AGACAGCCCTGGACCctcgcaggtggtgtcggcg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
colspan="4"	pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3267*(Cterm)-Lk15-GFP(1-10)
Rv0822c(Cterm)	M. tuberculosis H37Rv chromosomal DNA	If-AmMBD-3267D	GAGCGCCTCGCCATGgagattcagcaccagcaggttacgacg
		If-L15-3267-Rv2	AGACAGCCCTGGACCgttgatgcactccggcgcgtc
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
colspan="4"	pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3484*(Cterm)-Lk15-GFP(1-10)
Rv0822c(Cterm)	M. tuberculosis H37Rv chromosomal DNA	If-AmMBD-3267D	GAGCGCCTCGCCATGgagattcagcaccagcaggttacgacg
		If-L15-3484-Rv2	AGACAGCCCTGGACCgttcacgcagggcacgccg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
colspan="4"	pGMCS-P1-*virRsol*-Lk2-GFP11-*rv0822c*-Lk15-GFP(1-10)
Rv0822c	M. tuberculosis H37Rv chromosomal DNA	If-AmMBD-0822c	GAGCGCCTCGCCATGagtgacggcgagagcg
		If-L15-0822c-Rv	AGACAGCCCTGGACCctactcgcaggtggtgtcgg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
colspan="4"	pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3267*-Lk15-GFP(1-10)
Rv3267(Cterm)	M. tuberculosis H37Rv chromosomal DNA	If-AmMBD-3267	GAGCGCCTCGCCATGatgtctgcgcaacgtgtggttc
		If-L15-3267-Rv	AGACAGCCCTGGACCtcagttgatgcactccggcgc
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
colspan="4"	pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3484*-Lk15-GFP(1-10)
Rv3484(Cterm)	M. tuberculosis H37Rv chromosomal DNA	If-AmMBD-3484	GAGCGCCTCGCCATGgcgcgttctgagggc
		If-L15-3484-Rv	AGACAGCCCTGGACCctagttcacgcagggcacgc
		Flex15-Right	GGTCCAGGGCTGTCTGGCC

pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	ctpC-Am-left2	CATGGCGAGGCGCTCAAG
CRISPRi mutants			
virR (158.1) c2	Rv0431	Rv0431crispr 1 F	GGGAGAGCAGCAGGAAGACGACGC
		Rv0431crispr 1 R	AAACGCGTCGTCTTCCTGCTGCTC
virR (42.2) c1	Rv0431	Rv0431crispr 2 F	GGGAACGAACCCAGTGCCTGCCAGA
		Rv0431crispr 2 R	AAACTCTGGCAGGCACTGGGTTCGT
cei (145.2) c1	Rv2700	Rv2700crispr 1 F	GGGAATCACGCAAGTCACCACCAC
		Rv2700crispr 1 R	AAACGTGGTGGTGACTTGCGTGAT
cei (47.3) c2	Rv2700	Rv2700crispr 2 F	GGGAGTCGGTGCCGAGCGCAAGGTCG
		Rv2700crispr 2 R	AAACGACCTTGCGCTCGGCACCGAC
VirR-Venus primers			
VirR-F			TCACTGGGGTGCTGGTTACAGTGGGCTC
VirR-R			ATAGGATCGCCGGTCACCACGACGAT
pMV-F			TGACCGGCGATCCTATGGTGAGCAAGGGC
pMV-R			ACCAGCACCCCAGTGAAAAGTTCTTCTCCTTTACTC

1793

1794 **Supplementary Table 2**. Gradient mobile phase composition for ion-pairing LC-MS/MS
1795 for the quantification of acetyl-CoA, methyl citrate and methylmalonyl-CoA.

Time (min)	Mobile Phase (%)	
	A	B
1	100	0
2	55	45
7	50	50
7.5	5	95
8	5	95
8.5	100	0
10	End	

1796 A: water-methanol 07:3 (v/v) + 10mM tributylamine + 3mM acetic acid; B: pure methanol
1797

1798 **Supplementary Table 3**. Gradient mobile phase composition for ion-pairing LC-MS/MS
1799 for the quantification of pyruvate.

Time (min)	Mobile Phase (%)	
	A	B
1	5	95
6	95	5
7	95	5
8	5	95
10	End	

1800 A: pure water; B: pure methanol
1801

1802 **Supplementary Table 4**. Quantifier MRM transitions and collision energies used for
1803 acety-CoA, methyl citrate, methylmalonyl-CoA and pyruvate.

Metabolite	Precursor ion [M-H]⁻ (m/z)	Product ion (m/z)	Collision energy (eV)

Acetyl-CoA	808.00	407.95	41
Methyl citrate	204.90	125.10	13
Methylmalonyl-CoA	432.60	410.50	10
Pyruvate	87.20	43.10	13

1804
1805
1806
1807

Supplementary figures

Fig S1. Diversity of *Mtb* LCP proteins. (**A**) Domain structure of LCP proteins in *Mtb*. Proteins are organized according to the presence of (i) both LCP (green) and LytR_C (slight orange) terminal domain (Rv3267, Rv3484 and Rv0822); (ii) the solo LytR_C domain (VirR and Rv2700); and the solo LCP domain (Rv3840). Additional information is provided for *Mtb* mutants in the indicated genes. (**B**) Phylogenetic analysis of LCP proteins in Mycobacteria (*M. tuberculosis, M. smegmatis, M. marinum and M. leprae*). The unrooted tree includes seven separate clusters for proteins with both LCP and LytR_C terminal domains (CpsA1, CpsA2 and CpsA3); with the solo LytR_C terminal domain (Solo_LytRC1 and Solo_LytRC2); with the solo LCP domain; and a separate cluster including the characterized LCP proteins from *S. pneumonia* and *S. aureus*. Note the clustering of one of the *M. smegmatis* LCP proteins with this group. VirR is highlighted in blue.

Fig S2. The enhanced vesiculation phenotype of *virR^{mut}* occurs in the absence of cell lysis. **(A)** Nanoparticle tracking analysis (NTA) of EV preparations derived from WT and the indicated strains showing number of particles per cm^3 determined by Zeta View NTA in three independent EV preparations. Data are presented as mean ± SEM. *P ≤ 0.05 (One-way ANOVA with Tukey's multiple comparisons test). **(B)** Cell lysis control of *virR^{mut}*. Shown are immunoblots of cell lysates and culture supernatants (SN) for the indicated strains. The cytoplasmic protein IdeR was detected in cell lysates but not in culture supernatants confirming cell integrity, while the secreted protein Ag85b was readily detected in the culture supernatants. **(C)** Cell lysis control of *Mtb* submitted to conditions that alter permeability. Arrows indicate the size of the proteins of interest detected by immunoblot. **(D)** Alternatively, cell viability was tested by serial dilution and plating onto 7H10 medium. Cultures from B and C were submitted to serial dilutions (10^{-1} to 10^{-5}) and plated onto 7H10 after reaching an OD_{600nm} of 0.2.

Fig S3. H_2O_2 sensitivity assay. The indicated strains were initially grown on 7H9 medium supplemented with AND to an OD of 0.5. Washed cells were then transferred to

1838 the same medium supplemented with and without 13mM H_2O_2. After 8 h, cells were

1839 diluted down as indicated and spotted onto 7H10 solid medium.

1840

1841 **Fig S4. Expression patterns of permeability-dependent genes upregulated in**

1842 **cultures exposed to cholesterol and simultaneously downregulated in $virR^{mut}$.**

1843

Fig S5. Behaviour of $virR^{mut}$ upon conditions that modulate permeability. (A)

Growth curves of $virR^{mut}$ in MM supplemented with either glycerol, cholesterol or glycerol

plus TRZ at the indicated concentrations. **(B)** Dotblot analysis of vesiculation levels of

$virR^{mut}$ on the indicated culture media. a, b and c denote biological replicates.

Quantification of the dotblot was done by measuring the integrated density of each dot

using ImageJ. Data are mean and standard deviations. * p=0.021 after applying a Tuke's

multiple comparison test. **(C)** Time course of the uptake of ethidium bromide (EthBr) by

$virR^{mut}$ grown in the indicated conditions, as measured by fluorescence at 590=nm for 65

min. Data are mean and standard deviation of three biological replicates.

1853

1854 **Fig S6. Expression patterns of ribosomal proteins in the EVs from WT, $virR^{mut}$**

1855 **and $virR^{mut}$ -Comp strains. Proteins shown are downregulated in the vesicles of**

1856 **$virR^{mut}$ with respect to both VirR expressing strains.**

1857

1858 **Fig S7. Expression patterns in the EVs from WT, $virR^{mut}$ and $virR^{mut}$ -Comp strains**

1859 **of proteins differentially expressed in $virR^{mut}$ that contribute to enrichments of**

1860 **terms related to metabolism of: (A) nucleotides (B) amino-acids (C)**

1861 **oligosaccharides (D) glycogen.**

1862

Fig S8. TLC analysis of strains grown in the presence of cholesterol or TRZ.

Analysis of polar lipids, including SL1 and DATs lipids of the indicated strains by TLC.

Sulfolipid 1 (SL1); Glucose monomycolate (GMM); Trehalose monomycolate (TMM).

Phosphatidyl-myo-inositol dimannosides (Ac_1PIM_2 and Ac_2PIM_2, respectively);

phosphatidyl-myo-inositol hexamannosides (Ac_xPIM_6); diphosphatidylglycerol (DPG);

phospholipid (P); phosphatidylethanolamine (PE).

1869

Fig S9. Additional AFM data. (A-C) (i) Atomic force microscopy images taken in liquid

from a big area showing the sample heterogeneity overview with various peptidoglycan

fragments of the following samples: **(A)** WT, **(B)** $virR^{mut}$, **(C)**, $virR^{mut}$-C. **(ii)** Atomic force

microscopy images zoom in one individual peptidoglycan fragment corresponding to one

cell from different samples: **(A)** WT, **(B)** $virR^{mut}$ showing the background (Backg.), the

external surface (Ext) the internal (Int), **(C)**, *virR^{mut}*-C, white square showing the size of
the zoom images shown in Figure 5 from the external surface of the cells. **(D)** Curve of
grey points showing 255 slices of one AFM image taken using a FIJI macro, in the x axis
and the total number of pores in the y axis, the red line is a Gaussian fitted to the curve
and the maximum of the Gaussian is marked with a dash red line showing slice 140 in
this example corresponding to the slice of the image with the maximum number of pores.
(E) Binary slice number 140 from the original AFM image showing the pores in white and
the material represented in black. **(F)** The pores from 'E' were analyzed using the FIJI
macros where each pore area was measured and plotted in the x axis against the
cumulative fraction of total pore area in the y axis as shown in the curve by blue dots.
The red dot marks the most likely pore area which corresponds to the half of the
cumulative fraction of the total pore area, this area was then used to obtain the pore
diameter of the image. The same was repeated with the n = 8 images for each sample
to compose the graphs 'D' and 'E' from Figure 7.

1889

1890 **Fig S10**. **Sensitivity to lysozyme of *virR^{mut}*. (A)** Analysis of the minimum inhibitory
1891 concentration (MIC) of lysozyme of the indicated strains by the resazurin method. **(B)**
1892 Growth of WT and *virR^{mut}* treated with 0 or 50 µg/ml lysozyme (Lys). Data represents the
1893 mean ± SEM of three independent cultures.

1894

1895 **Fig S11. Localization of VirR by fluorescence microscopy**. Fluorescence images of
1896 a *Mtb* strain expressing Venus-tagged VirR.

1897

Fig S12. Split-GFP strategy. (A) *Upper panel*. Scheme of the plasmid used to test VirR
interactions in vivo. *Lower panel*. Scheme of the potential molecular interaction between
VirR and Lcp1 using split-GFP. **(B)** Gating strategy to analyze split-GFP flow cytometry
data. Bacteria were first sorted based on size (FSC) and structure (SSC) to exclude
debris (1). Then, aggregates were excluded based on FSC and SSC width (W) and
eighth (H) (2,3). GFP signal was then analyzed *vs*. SSC (4) or mCherry, when relevant
(5).

1905

1906 **Fig S13. Quantitative RT PCR of *virR* downstream genes Rv0432 and Rv0433.**

1907

1908 **Supplementary file S1**: FileS1_RNA_differential_expression.xlsx: Results of the
1909 differential expression analyses for the RNA-seq dataset.

1910

1911 **Supplementary file S2**: FileS2_GOenrichments_RNA.xlsx: Gene ontology enrichment

1912 analyses conducted on sets of genes up (N=399), and down-regulated (N=502) in the

1913 *virR^{mut}* strain. Rows highlighted correspond to terms selected for graphical

1914 representation (figure 3E, OR>2 & FDR<0.1)

1915

1916 **Supplementary file S3**: FileS3_TRN_regulon_enrichments.xlsx: Enrichment analyses

1917 for the Transcription factor regulons reported in (Minch et al. 2015).

1918

1919 **Supplementary file S4**: FileS4_Proteins_differential_expression.xlsx: Results of the

1920 differential expression analyses for the LF-MS dataset.

1921

1922 **Supplementary file S5**: FileS5_GOenrichments_proteins.xlsx: Gene ontology

1923 enrichment analyses conducted on sets of protein up (N=117), and down-regulated

1924 (N=108) in the extracellular vesicles of the *virR^{mut}* strain. Rows highlighted correspond

1925 to terms selected for graphical representation (figure 5F, OR>2 & FDR<0.1).

1926

A

WT   virR^mut   virR^mut-C

B

C

	CM	CM-OM	CM-L1	OM-L2	L1-L2	OM
WT	2.7±0.2	44.09±3.1	11.2±0.4	13.34±0.9	10.46±1.4	5.7±0.4
virR^mut	5.4±0.7*	62.7±3.2*	31.36±3.2*	15.1±2.2	6.8±0.9*	6.1±0.3
***virR^mut*-C**	2.5±0.5	41.6±0.7	12.4±1.4	10.03±1.1	12.6±1.3	5±0.2

Figure 1

Figure 1. Ultrastructural changes in the cell envelope of *virR*-deficient mutant associated with increased vesiculation. (A) Cryo-electron micrographs of indicated Mtb strains grown in high iron MM. Closed line rectangles were used to calculate grey value profiles of cell envelopes using ImageJ. The dashed line insets within the main micrograph were magnified to show a detailed view of the cell surface. Scale bars are 100 nm in main micrographs and 50 nm in the insets. (B) Density profiles based on grey values of the cross sections marked by solid line rectangles in A. (C) Mean values and standard errors of distances in nm between main cell envelope layers measured in A and C. *P < 0.05 after applying a Tukey´s multiple comparison test. The number of cells analyzed varied from n = 20 (WT), n = 15 (virRmut) and n = 23 (virRmut-C). CM, cytoplasmic membrane; OM, outer membrane; L1, layer 1; L2, layer 2.

Figure 2

Figure 2. Enhanced permeability and vesiculation are linked in the absence of LytR_C solo domain proteins in Mtb. (A) Uptake of ethidium bromide (EthBr) in the indicated strains, as measured by fluorescence at $590\,\text{nm}$ for 65 min. Data are mean and standard deviation of three biological replicates. (B) Sensitivity of the indicated strains to different concentrations of vancomycin as measured by monitoring optical density at 580 nm (OD580) after 7 days. Data are mean and standard deviations of three biological replicated. Errors bars not shown are smaller than symbols. (C) Time course of the incorporation of FDAAs on the indicated strains in both 7H9 and MM as measured by fluorescence. Representative images of both WT and virRmut strains in either 7H9 or MM at 8h after the addition of the FDAA. Data are mean and standard deviations of three biological replicates. The arrow indicates the time at which images were taken. Errors bar not shown are smaller than symbols. (D) immunoblot analysis of cell lysates of two independent conditional mutants (c1 and c2) for virR and cei for the presence of VirR and Cei proteins in cultures incubated with and without anhydrotetracycline (ATc). (E) Nanoparticle tracking analysis (NTA) of EV preparations derived from virR and cei conditional mutants (c2 (virR) and c1 (cei)) obtained from cultures with and without ATc showing number of particles per cm3 determined by Zeta View NTA in three independent EV preparations. Data are presented as mean $\pm\,\text{SEM}$.

Figure 3

Figure 3. Transcriptional profiling of *virR^mut^* showcases systemic alterations in cell wall architecture and metabolism. (A) Experiment design for RNA-seq analyses. RNA was extracted from whole cell extracts from 4 replicate cultures of each strain (WT, *virR^mut^*, and *virR^mut^*-Comp), and sequenced. **(B)** PCA plot of the transcriptomics data. The loadings from the two first principal components are plotted. **(C)** Number of differentially expressed genes between *virR^mut^* and either WT (left) or *virR^mut^*-Comp (right). **(D)** Enrichment profiles between sets of genes found up and down-regulated for the contrasts *virR^mut^* vs WT and *virR^mut^* vs *virR^mut^*-Comp. While genes that are differentially expressed in the same direction in both contrasts are mutually enriched, genes whose differential expression goes in different directions in both comparisons are mutually depleted (two-sided Fisher exact test $p<0.05$ in all four cases). **(E,F)** Gene Ontology (GO) enrichments of genes that are 1) simultaneously downregulated in *virR^mut^* with respect to both WT and *virR^mut^*-Comp **(E)** and 2) simultaneously upregulated **(F)**. GO terms selected for testing included biological process and cell compartment labels containing N > 10 genes, between levels 4-6 of the ontology tree. In this representation, each node in the networks represents a significantly enriched term (terms selected for visualization with FDR=10% and enrichment odds-ratio>2), nodes' size is proportional to the enrichment significance, and links between terms are added wherever the sets of genes contributing to the enrichments in a connected pair show an intersection larger than 50% of the smallest of the gene sets involved in the pair (see Methods for further details).

Figure 4

Figure 4. VirR-dependent genes intersect the regulons of key transcriptional regulators of the responses to stress, dormancy, and cell wall remodeling. (A) Transcription factors controlling regulons significantly enriched among genes that are 1) simultaneously downregulated in *virR^{mut}* with respect to both WT and *virR^{mut}*-Comp (bottom, 9 TFs) and 2) simultaneously upregulated (top, 1 TF). **(B)** Differential expression patterns for enriched TFs whose expression is VirR-dependent. **(C)** Regulatory subnetwork including the genes under transcriptional control of the enriched TFs highlighted in figure 4A that were found to be VirR-dependent. In this visualization, color captures the logFC for the contrast *virR^{mut}*-WT, and the size its FDR.

Figure 5

Figure 5. Cell permeability is a major driver of vesicle production in *Mtb*. (A) Time course of the uptake of ethidium bromide (EthBr) by *Mtb* grown in the indicated conditions, as measured by fluorescence at 590=nm for 65 min. Data are mean and standard deviation of three biological replicates. **(B)** Dot-blot analysis of released EVs in supernatants from *Mtb* cultures submitted to the indicated conditions. The loading volume was normalized according to CFUs. The graph below represents the quantification of the dotblot using ImageJ. The integrated cell density was normalized to WT values. Data are mean and standard errors from the three biological replicates; **** denotes P<0.0001, *** denotes P=0.0028 after applying a Tuke's multiple comparison test. **(C)** Enrichment odds ratios for Fisher exact tests comparing sets of genes up, and down regulated, respectively, in 1) cholesterol-supplemented medium vs. glucose-supplemented medium (Pawełczyk et al., 2021), GEO accession GSE175812; and 2) *virR^mut* vs WT as well as *virR^mut* vs *virR^mut*-Comp. **(D)** Euler diagram showing the intersection between A) genes upregulated by cholesterol, B) genes down-regulated in *virR^mut*, and C) the DevR regulon. **(E)** Expression patterns in response to TRZ (1h) and in *virRmut* for genes involved in metal signaling, metabolism and homeostasis.

Figure 6. VirR regulates the protein content of EVs in Mtb. (A). Experiment design for proteomics analyses. label-free mass spectrometry data was retrieved for replicates of H37RV (WT), *virR^mut* and *virR^mut*-Comp. For each strain, data corresponding to proteins extracted from either whole-cell extracts (WC) and EVs was produced, totaling 12 samples (2 strains times 2 cell compartments times 3 replicates). **(B)** PCA plot of the proteomics data. The loadings from the two first principal components, adding up to 50.5% of variance explained, are plotted. **(C)** Expression patterns of VirR in the proteomics dataset across cell compartments and strains. As expected, while *virR^mut* shows significantly lower levels of VirR protein, the complemented strain does not just restore WT levels of expression, but leads to a strong overexpression of VirR, both in the lysates and the EVs, **(D)** Bar plots of differentially expressed proteins found between strains in EVs (up: virRmut vs WT; down virRmut vs virRmut-Comp at 5% FDR). **(E)** Enrichment profiles between sets of proteins found up and down-regulated for the contrasts *virR^mut* vs WT and *virR^mut* vs *virR^mut*-Comp. **(F)** Gene Ontology enrichments of the downregulated proteins in the VirR mutant strain (left) and the upregulated ones (Right), at the EV location. GO terms selected for testing included biological process and cell compartment labels containing N > 3 genes, between levels 4-6 of the ontology tree. As in **figure 3E**, terms selected for visualization with FDR=10% and enrichment odds-ratio>2, nodes' size is proportional to the enrichment significance, and links between terms are added wherever the sets of proteins contributing to the enrichments in a connected pair shows an intersection larger than 50% of the smallest of the gene sets involved in the pair (see Methods for further details, and supplementary file S5 for an extended list of all the enrichments found). **(G)** Expression patterns of bacterioferritin proteins in the EVs of WT and *virR^mut*. **(H)** Expression patterns of proteins contributing to enrichments of terms related to mycolic acids biosynthesis.

Figure 6

Figure 7. Lipidomic analysis of cell envelope and EVs from virRmut. (A) Comparative total lipid analysis of the indicated strains by nano-LC-MS. Lipid species with an abundance higher than 0.5% are shown. Data indicates mean and standard error from three biological replicates. (B) Analysis of polar lipids of the indicated strains by bidimensional (2D)-layer chromatography (TLC) (2D-TLC). Phosphatidyl-myo-inositol dimannosides (Ac1PIM2 and Ac2PIM2, respectively); phosphatidyl-myo-inositol hexamannosides (AcxPIM6); phosphatidylethanolamine (PE); diphosphatidylglycerol (DPG). Phospholipid (P). (C) Analysis of mycolic acid species in the indicated strains by TLC. (D) Analysis of PIDMs and TAGs by TLC. (E) Analysis of apolar lipids of the indicated strains by TLC. Diacylglycerol (DAG); free fatty acid (FF); free mycolic acid (FMA). (F) Analysis of apolar lipids of MEVs isolated from the indicated strains. Triacylglycerols (TAGs); phthiocerol dimycocerosates (PDIMs) (G) Analysis of polar lipids of MEVs isolated from the indicated strains. Phosphatidyl-myo-inositol dimannosides (Ac1PIM2 and Ac2PIM2, respectively); phosphatidyl-myo-inositol hexamannosides (AcxPIM6); diphosphatidylglycerol (DPG).

The following table is from panel G:

Peak	R.T	Observed	Expected	Best Match
1	4.3	942.432	942.4155	M4
2	6.12	1752.80	1752.7379	D43 (-N-Ac)
3	6.59	1823.856	1823.7750	D44(-N-Ac)
4	6.62	1,823.862	1823.7750	D44 (-N-Ac)
5	6.87	1794.8	1794.7755	D43
6	7.1	1794.8	1794.7755	D34
7	7.76	1865.854	1865.8126	D44
8	8.82	1831.86	1831.748	D45N(-NH2,-2N-Ac)
9	9.7	2789.26	2789.2098	T444
10	10.56	1845,806	1774.7493	D44N (Anh)

Figure 7

Figure 7. Chemical and nanometric analysis of isolated cell walls shows higher amounts of PG in the absence of VirR. (A) Glycosyl composition analysis of isolated cell walls from the indicated strains. Acid hydrolysis prior to GC-MS analysis of isolated cell walls was performed to determine the total amounts of galactose (G), arabinose (A), rhamnose (R), and N-acetylglucosamine (NAGc). The ratio between different sugars is shown to indicate: (i) the relative levels of arabinogalactan (A/G); (ii) the length of the AG polysaccharide chain (G/R); (iii) the amount of PG (NAGc/Rham). Individual measurements are shown from three biological replicates. Data are mean and standard errors. **$P < 0.01$ after applying a Tukey´s multiple comparison test. (B) Quantification of diaminopimelic acid (DAP) on isolated cell walls from the indicated strains. Data are mean and standard errors from three biological replicates; **** denotes $P<0.0001$, * denotes $P=0.0119$ after applying a Tukey´s multiple comparison test. (C) Atomic force microscopy high resolution images taken in liquid from the external surface of purified peptidoglycan from different samples, left to right: WT, virRmut, virRmut-C, the structure of external peptidoglycan layer shows a mesh with pores of different sizes, (D) Graph showing the pore diameter from n = 8 AFM images similar to (C) (each point represent an image) of different samples; black triangles: WT, blue circles: virRmut, red squares: virRmut-C. The pore diameter was calculated from the binary slice from each image containing the greatest number of pores, according to (Pasquina-Lemonche, 2024). (E) The number of pores per surface area from different samples; white triangles: WT, red circles: virRmut, blue squares: virRmut-C, this was calculated from the slice containing the maximum number of pores where the pore size was analyzed, following the method from (Pasquina-Lemonche, 2024). (F) Graph showing the peptidoglycan thickness manually measured using the 1D statistic tools from the open-source program Gwyddion (Nečas and Klapetek, 2012) from AFM images in air containing several fragments of cell wall, each point represents a different peptidoglycan fragment from a different cell. There were three samples analyzed: black WT n=23 peptidoglycan fragments, blue virRmut n=36 and red virRmut-C n=25. Statistics: D) Using a two-tailed t test with Welch's correction the statistical comparison are: (***) pWT- virRmut = 4.1 10-4 with t = 6.1, df = 7.5, (**) p virRmut - virRmut-C = 0.007 with t = 3.2, df = 13.2, E) Using a two-tailed t test with Welch's correction the statistical comparison are: (****) pWT- virRmut = 6.6 10-10 with t = 16.3, df = 12.7, (**) p virRmut - virRmut -C = 0.01 with t = 3.0, df = 13.2., F) Using a two-tailed t test with Welch's correction the statistical comparison are: (****) pWT- virRmut = 2.8 10-18 with t = 17.8, df = 32.3, (****) virRmut - virRmut -C = 3.9 10-17 with t = 16.4, df = 31.8. (G) Analysis of deacetylated muropeptides. The table includes muropeptides detected by UPLC-UV. Muropeptides indicated in red are only detected in the PG of virRmut. (H) The relative abundance of deacetylated muropeptides was calculated by dividing the number of D-deacetylated muropeptides by the total number of muropeptides detected by UPLC-UV.

Figure 8

Figure 8. VirR interacts with Lcp1. (A) Blue-native PAGE analysis of recombinant VirR and Lcp1. The molecular mass of markers in kDa are indicated on the left side of the gel. The size of the multimers of VirR and Lcp1 are indicated on the left side of the gel. (B) Upper panel. Representative image of a flotation assay of recombinant VirR alone or in combination with Lcp1 performed on an idioxanol density gradient. The presence of VirR was detected in each fraction by dot-blot using murine polyclonal antibodies raised against VirR. Numbers indicate collected fractions from top (1) to bottom (11). Lower panel. Quantification of the dot-blot by measuring the pixel intensity of the dots using ImageJ. Data are mean and standard error of three independent experiments. (C) Bipartite split-GFP experiment using *M. smegmatis* Δ2 expressing plasmids depicted in (Supplementary Fig 9), including CtpCN, VirR, VirRsol(Δ1-41), Rv0822, Rv3267 (Lcp1), Rv3484, Rv3484LytR_C , Rv3267LytR_C and Rv0822 LytR_C . Bacteria were grown in complete 7H9 and fluorescence was recorded by FACS. Pooled data obtained from four to six independent cultures of each strain are shown. Statistical analysis was performed using non-parametric two-sided Mann-Whitney test. ns: not significant. * and ** refer to p-values <0.05 or 0.01, respectively.