Maintenance of cell wall remodeling and vesicle production are connected in *Mycobacterium tuberculosis*

Vivian Salgueiro[1]*, Jorge Bertol[2]*, Claude Gutierrez[3], Jose L. Serrano-Mestre[1], Noelia Ferrer-Luzon[2], Ainhoa Palacios[4], Laia Pasquina-Lemonche[5], Akbar Espalliat[6], Laura Lerma[1], Brian Weinrick[7], Jose L. Lavin[8], Felix Elortza[4], Mikel Azkalgorta[4], Alicia Prieto[9], Pilar Buendía-Nacarino[10], José L. Luque-García[10], Olivier Neyrolles[3], Felipe Cava[6], Jamie K. Hobbs[5], Joaquín Sanz[2]# and Rafael Prados-Rosales[1]#

[1]Department of Preventive Medicine and Public Health and Microbiology, Universidad Autónoma de Madrid, Madrid, Spain; [2]Institute for Bio-computation and Physics of Complex Systems BIFI, Department of Theoretical Physics, University of Zaragoza, Zaragoza, Spain; [3]Institut de Pharmacologie et de Biologie Structurale, IPBS, Université de Toulouse, CNRS, UPS, Toulouse, France; [4]CIC bioGUNE, , Basque Research and Technology Alliance (BRTA), Bizkaia Science and Technology Park, Derio, Spain; [5]Department of Physics and Astronomy, University of Sheffield, Sheffield, UK; [6]Department of Molecular Biology and Laboratory for Molecular Infection Medicine Sweden, Umeå Centre for Microbial Research, SciLifeLab, Umeå University, Umeå, Sweden; [7]Trudeau Institute, Saranac Lake, NY, USA; [8]Bioinformatics Unit, Neiker-Tecnalia, Derio, Spain; [9]Department of Microbial & Plan Biotechnology, Centro de Investigaciones Biológicas Margarita Salas, Spanish National Research Council (CSIC),Madrid, Spain; [10]Department of Analytical Chemistry, Universidad Complutense de Madrid, Madrid, Spain.

Key words: extracellular vesicles, *Mycobacterium tuberculosis*, vesiculogenesis, peptidoglycan, membrane, permeability.

#To whom correspondence should be addressed:

Dr. Rafael Prados-Rosales– rafael.prados@uam.es

Dr. Joaquín Sanz- jsanz@bifi.es

*Equal contribution

Running title: *Mycobacterium* vesicle biogenesis mechanism.

Abstract

Pathogenic and nonpathogenic mycobacteria secrete extracellular vesicles (EVs) under various conditions. EVs produced by *Mycobacterium tuberculosis* (*Mtb*) have raised significant interest for their potential in cell communication, nutrient acquisition, and immune evasion. However, the relevance of vesicle secretion during tuberculosis infection remains unknown due to the limited understanding of mycobacterial vesicle biogenesis. We have previously shown that a transposon mutant in the LCP-related gene *virR* (*virR^mut*) manifested a strong attenuated phenotype during experimental macrophage and murine infections, concomitant to enhanced vesicle release. In this study, we aimed to understand the role of VirR in the vesicle production process in *Mtb*. We employ genetic, transcriptional, proteomics, ultrastructural and biochemical methods to investigate the underlying processes explaining the enhanced vesiculogenesis phenomenon observed in the *virR^mut*. Our results establish that VirR is critical to sustain proper cell permeability via regulation of cell envelope remodeling possibly through the interaction with similar cell envelope proteins, which control the link between peptidoglycan and arabinogalactan. These findings advance our understanding of mycobacterial extracellular vesicle biogenesis and suggest that these set of proteins could be attractive targets for therapeutic intervention.

Introduction

Mycobacterium tuberculosis (*Mtb*) is one of the most successful human pathogens with the capacity to persist within the host establishing long-lasting infections. *Mtb* possesses multiple highly evolved mechanisms to manipulate host cellular machinery and evade the host immune system. Similar to other intracellular pathogens, *Mtb* releases proteins and lipids, that interfere with host cellular processes, via canonical or specialized secretion systems (Majlessi et al., 2015). How mycobacteria confined within phagosomes delivers virulence factors to other cellular compartments and/or into the extracellular milieu to affect the physiology of neighboring cells is currently a matter of investigation. Understanding the physiology of this phenomenon during the infection process might be critical to develop alternative therapies against tuberculosis (TB).

Like most forms of life, *Mtb* uses an alternative antigen release process based on extracellular vesicles (EVs) (Prados-Rosales et al., 2011). Mycobacterial EVs (MEVs) are membrane walled spheres of 60-300 nm in diameter, released by live bacteria *in vitro* and *in vivo*. MEVs contain iron scavenging molecules, immunologically active complex lipids and lipoproteins, and classical virulence factors such as the *Mycobacterium ulcerans* toxin mycolactone (Lee et al., 2015; Marsollier et al., 2007; Prados-Rosales et al., 2011, 2014b). Several studies have clearly established a role of MEVs in immunomodulation and shown that MEVs deliver factors that impair macrophage effector functions, inhibit T cell activation, and modify the response of host cells to infection (Athman et al., 2015, 2017; Prados-Rosales et al., 2011, 2014b, 2014a; Rath et al., 2013). MEVs have also shown potential as biomarkers of *Mtb* infection (Schirmer et al., 2022; Ziegenbalg et al., 2013). Although the importance of MEVs has been recognized, hardly anything is known regarding both how such vesicles are exported across the mycobacterial cell wall and the molecular mechanisms underlying vesicle formation in mycobacteria. To date, several modulators of MEVs have been described. Our group identified two conditions leading to stimulation of vesicle production, namely: iron starvation (Prados-Rosales et al., 2014b) and the deletion of *virR* (Rv0431, vesiculogenesis and immune response Regulator) (Rath et al., 2013). In addition, it was shown that the Pst/SenX3-RegX3 signal transduction system regulates MEVs production independent from VirR (White et al., 2018), suggesting an alternative mechanism of vesicle production. We have recently demonstrated that both iron starvation and deletion of VirR in *Mtb*, two hypervesiculating conditions, trigger the induction of the isoniazid-induced gene operon *iniBAC* (Gupta et al., 2023). We determined that both IniA and IniC are dynamin-like proteins (DLP) presumably assisting MEV release through the fusion of the cell membrane (Gupta et al., 2023; Wang et al., 2019). Understanding how defects

in VirR lead to hypervesiculation through DLPs is important to provide mechanistic
explanation for the vesiculation process in mycobacteria.

Several lines of evidence support the notion that VirR might be involved in
maintaining cell wall integrity: (i) First, sequence analysis of the VirR protein indicates
that it contains a conserved LytR_C domain (**Fig S1**), which is usually found in
combination with another domain named LytR-Cps2A-Psr (LCP), and proteins carrying
both domains have been associated with the remodeling of the cell wall in Gram-positive
bacteria (Hubscher et al., 2008). *Mtb* has six genes encoding for LCP proteins, three with
both LCP and LytR_C domains (Rv3267, Rv3484 and Rv0822), two with a solo LytR_C
domain (VirR and Rv2700) and one protein with a solo LCP domain (Rv3840) (**Fig S1**).
The LCP family includes enzymes that transfer glycopolymers from membrane-linked
precursors to peptidoglycan (PG) or cell envelope proteins and are central to cell
envelope integrity (Kawai et al., 2011). In mycobacteria its was demonstrated that some
members of the (LCP) family of proteins are responsible for the linkage between
arabinogalactan (AG) and PG (Grzegorzewicz et al., 2016; Harrison et al., 2016).
Rv3267, Rv3484 and Rv0822 appear to have overlapping functions in cell wall assembly,
Rv3267 being the primary ligase (Grzegorzewicz et al., 2016). Interestingly, while
knocking down the gene encoding for the main AG-PG ligase (cg0847, *lcpA*) in
Corynebacterium glutamicum leads to the release of outer membrane material to the
extracellular medium (Baumgart et al., 2016), this phenotype is not observed in any
knockout strain in the *Mtb* orthologs (Grzegorzewicz et al., 2016), probably reflecting the
ability of the mutants to negatively regulate the biosynthesis of cell wall constituents in
response to a decrease in ligase activity; (ii) Second, we and others have determined
the increased susceptibility of the transposon *virR* mutant (*virR^mut^*) with respect to the
WT (Ballister et al., 2019; Rath et al., 2013), and the mutant in the *virR* ortholog, Rv2700
(cell envelope integrity, *cei*) to cell wall targeting drugs such as meropenem (Ballister et
al., 2019). Based on these results, we hypothesize that LytR_C solo domain proteins
including VirR and Cei Rv2700 have critical roles in the maintenance of cell wall
homeostasis and their absence provokes cell wall alterations leading to enhanced
vesiculation.

In this work we show that the absence of VirR leads to ultrastructural changes in
the cell envelope as observed by cryo-electron microscopy (cryo-EM). These changes
are compatible with enhanced permeability and an enlargement of the PG layer, as
measured by high resolution atomic force microscopy (AFM). We demonstrate that VirR
interacts with canonical LCP proteins and propose a model for such interaction. These
results indicate that VirR is a central scaffold at the cell envelope remodeling process

assisting the PG-AG linking process and that this function is important for vesicle
production in *Mtb*.
164
165
166 **Results**
167

High EV production linked to the lack of VirR is associated with an altered cell
envelope in *Mtb* in the absence of cell lysis. While Gram-negative bacteria can
release outer membrane vesicles directly into the extracellular environment from their
outermost compartment, Gram-positive bacteria and mycobacteria have thick cell
envelopes surrounding the cytoplasmic membrane that acts as a permeability barrier.
The mycobacterial cell envelope is composed of PG covalently linked to AG, which in
turn is decorated with exceptionally long-chain mycolic acids (MA). These MA, together
with intercalating glycolipids, form the unique mycobacterial outer membrane (OM)
(Kalscheuer et al., 2019). Previous analysis of mycobacterial vesicle associated lipids
showed predominantly polar lipids, consistent with the cytoplasmic membrane being the
likely origin of the vesicles (Prados-Rosales et al., 2011). This is also true in EV produced
by Fe-limited *Mtb* whose lipid content consists mainly of polar lipids and the lipidic
siderophore, mycobactin (Prados-Rosales et al., 2014b). Given that these vesicles must
traffic from their point of origin in the plasma membrane through the cell envelope, we
hypothesized that local remodeling of the cell envelope is necessary for EV release, and
that this process may be exacerbated in a *virR* mutant (*virR^{mut}*). To test this, we examined
the cell envelope structure of *virR^{mut}* grown in minimal medium (MM), as well as wild type
(WT) and *virR^{mut}* complemented (*virR^{mut}*-C) strains (Rath et al., 2013) (**Fig 1**). Using cryo-
electron microscopy (cryo-EM) on whole cells, which allows the study of mycobacterial
cell surface-associated compartments in a close-to-native state (Sani et al., 2010), we
could easily discern three layers comprising the mycobacterial cell envelope: i) the outer
membrane (OM); ii) an intermediate layer composed of two sublayers (L1 and L2,
previously assigned to the mycolate-PG-AG (mAGP) network (Sani et al., 2010)); and iii)
the cytoplasmic membrane (CM) (**Fig 1**). We observed a significant increase in the
thickness of the overall cell envelope (OM-CM) in the *virR^{mut}* strain compared to the WT.
This difference was largely explained by the expansion of the CM-L1 layer, which
appeared also more granulated, suggesting cell wall alterations closer to the cell
membrane and indicating aberrant production of either AG, PG or both.
Complementation of *virR* restored normal cell envelope structure (**Fig 1A-C**).
In Gram-negative bacteria increased vesiculation has been observed in
conditions associated with either loss or maintenance of periplasm integrity (McMillan

and Kuehn, 2021). Moreover, a novel mechanism of vesicle production common to both Gram-negative and Gram-positive bacteria has recently been described, in which a cell explosion phenomenon driven by bacteriophages leads to the formation of EVs in culture when bacteria are under genotoxic stress (Nagakubo et al., 2021). Consequently, we tested whether the enhanced vesiculation described in $virR^{mut}$ (Rath et al., 2013) could be linked to cell lysis events. To do this, we analyzed the presence of cytoplasmic IdeR, a transcription factor which is not present in EVs and is not detected in secreted proteins preparations, in culture supernatants of *Mtb* strains, assuming that its detection could be linked to a loss of envelope integrity (Gupta et al., 2023). We first confirmed that the $virR^{mut}$ strain manifested an enhanced vesiculation by measuring EVs using nanoparticle tracking analysis (NTA) (**Fig S2A**). We did not detect differences in size distribution of isolated EVs between strains. The same supernatants and cell lysates were used to detect the presence of IdeR by immunoblot using a specific polyclonal antibody (Pandey and Rodriguez, 2014). As a control for normal release of an extracellular protein, we used a monoclonal antibody against Ag85b (Prados-Rosales et al., 2017). We could not detect IdeR in the supernatant of WT and $virR^{mut}$ strain, but a faint band corresponding to IdeR could be observed in the SN of the *virR* complemented strain ($virR^{mut}$-Comp). These results suggest that the absence of *virR* does not lead to cell lysis but its overexpression may compromise cell envelope integrity (**Fig S2B**). We validated this approach by serially diluting WT, $virR^{mut}$ and $virR^{mut}$-Comp strains on 7H10 solid media. As shown in **Fig S2D**, we did not observe differences in the number of cells across the diluted spots between strains. Taken together, the observed changes in the cell envelope of high EV producing bacteria suggest that cell envelope alterations in response to lack of VirR and increased membrane vesicle biogenesis may be connected in *Mtb*.

The absence of LytR_C solo domain proteins in *Mtb* leads to enhanced cell envelope permeability concomitant to hypervesiculation. The collective cryoEM observations that indicated an enlargement of the compartment next to the cell membrane may be linked to a more permeable cell envelope. Supporting this notion is the recent finding that $virR^{mut}$ or the mutant in the *virR* ortholog *cei* (Rv2700) manifest an enhanced uptake of ethidium bromide (EtBr) relative to WT strain (Ballister et al., 2019). We could reproduce these results and add that $virR^{mut}$-Comp strain restores the permeability levels of the WT strain (**Fig 2A**). This enhanced permeability was concomitant to an increased sensitivity to different antitubercular drugs, including meropenem and vancomycin (Ballister et al., 2019). In agreement with these results, we could measure a significantly enhanced sensitivity of $virR^{mut}$ to the PG-targeting drug vancomycin relative to WT and $virR^{mut}$-Comp strains (**Fig 2B**). Interestingly, the

magnitude of the sensitivity to cell wall targeting drugs was either moderate (ethambutol)
or absent (isoniazid) in *virR^{mut}* (Ballister et al., 2019).
Furthermore, we observed enhanced incorporation of fluorescent D-amino acids
(FDAAs) in *virR^{mut}* relative to the WT and complemented strains (**Fig 2C**). These probes
can freely enter cells and may be incorporated into PG by at least three different
mechanisms, depending on the species: through the cytoplasmic steps of PG
biosynthesis and via two distinct transpeptidation reactions taking place in the periplasm
(Kuru et al., 2019). Consequently, the differential labeling observed in *virR^{mut}* relative to
WT strain may be a consequence of altered PG turnover in the mutant.
We then tested whether the enhanced permeability and vesiculation of *virR^{mut}* are also
connected in an *Mtb* mutant lacking *cei* (Ballister et al., 2019). To do this, we generated
cei conditional mutants using the CRISPR interference (CRISPRi) technology (Rock et
al., 2017). Further, we also generated *virR* CRISPRi conditional mutants to properly
compare both strains (**Fig 2D**). When these strains were cultured in the presence of
anhydrotetracycline (ATc) to induce the transcriptional silencing of target genes, we
could measure enhanced vesiculation by analyzing vesicle concentration in the
supernatant by NTA (**Fig 2E**), relative to strains cultured in the absence of ATc. Overall,
these results validate the enhanced permeability phenotype of *Mtb* mutants in the
LytR_C solo domain proteins VirR and Cei and establish that this subfamily of proteins
participates in the regulation of vesicle production in *Mtb*. Moreover, the higher
susceptibility to PG-cross-linking inhibitors in both mutants and the enhanced
incorporation of FDAAs in the *virR* mutant strain suggests defects in PG remodeling.
258
Transcriptional profiling of *virR^{mut}* showcases systemic alterations in cell wall
architecture and metabolism. To identify *virR*-dependent mediators of vesicle
production, we carried out a transcriptional profiling of *virR^{mut}* and compared it to those
from a WT and *virR^{mut}*-Comp strains. Previous efforts to define the transcriptional profile
of *virR^{mut}* showed no significant differences compared to WT (Rath et al., 2013).
However, those experiments were carried out with bacteria grown in rich 7H9 medium,
while here we use a defined high-Fe minimal medium (MM), which is typically used for
vesiculation experiments (Prados-Rosales et al., 2011). We extracted RNA from four
independent cultures (average optical densities (OD) of 0.40, 0.38 and 0.42 for WT,
virR^{mut}, and *virR^{mut}* -Comp, respectively) one day before vesicle isolation, and conducted
RNA-seq (**Fig 3A**). Principal component analysis (PCA) was then performed on the
transcriptomic data, after filtering out lowly expressed genes, as well as genes containing
outlier observations (N=3985 genes tested, see Methods), and revealed significant
differences between strains. The first principal component, explaining 49% of the

variance, separated $virR^{mut}$ strain samples from strains bearing functional copies of $virR$

(**Fig 3B**, WT and $virR^{mut}$-Comp vs $virR^{mut}$, one-tailed Mann-Whitney test p=2.02E-2).

Furthermore, the second principal component (28.3% variance) captured $virR$-

independent effects of mutagenesis on transcriptional profiles, as it separated WT

samples from $virR^{mut}$-Comp and $virR^{mut}$ (one-tailed Mann-Whitney test p=2.02E-2).

After PCA, we conducted differential expression analysis between strains to find 1808,

and 2230 genes differentially expressed in $virR^{mut}$ compared to either WT, or $virR^{mut}$-

Comp strains, respectively (**Supplementary file S1**, **Fig 3C**, Benjamini-Hochberg

FDR<0.05), representing 45.4% and 56.0% of all genes tested. Although transcriptional

profiles of the WT and $virR^{mut}$-Comp strains are also significantly diverse, with up to 2184

differentially expressed genes between them, the differences in expression between

$virR^{mut}$ when compared to either WT or $virR^{mut}$ -Comp are significantly correlated (r=0.63,

p<2.2e-16), and the sets of both upregulated and downregulated genes in $virR^{mut}$ vs any

of the other strains are mutually, directionally enriched (**Fig 3D**). Finally, we defined a

set of N=399 genes that are simultaneously upregulated in $virR^{mut}$ with respect to any of

the two strains containing functional copies of $virR$ (logFC>0 and FDR<0.05 for both

$virR^{mut}$ vs WT and $virR^{mut}$ vs $virR^{mut}$-Comp), as well as a set of N=502 genes

downregulated in $virR^{mut}$ vs both $virR$-containing strains. This way, the differential

expression signature found for these genes in the $virR^{mut}$ strain is most likely VirR-

dependent and not-induced by non-specific consequences of the mutagenesis

procedures.

Next, we looked for gene ontologies (GO) enriched among these $virR$-dependent genes,

stratified by the direction of the effects (399 up-regulated, and 502 down-regulated in

$virR^{mut}$ relative to WT and $virR^{mut}$-Comp alike). A total of 51 GO terms related with

biological processes and cell compartment appeared significantly enriched among genes

upregulated in the $virR^{mut}$ strain (FDR=10%), while 85 terms were enriched among

downregulated genes (FDR=10%, see Methods, and **supplementary file S2**). When

focusing on the most prominent of these enrichments (OR>2), the analyses highlighted

several groups of biological processes and cell compartment ontologies that appeared

altered in $virR^{mut}$ with respect to both WT and $virR^{mut}$-Comp strains (**Fig 3E**).

First, among genes that were expressed at lower levels in $virR^{mut}$, we found terms related

to several stresses associated with the intra-phagosomal environment (Schnappinger et

al., 2003), such as response to hypoxia (FDR=1.7E-5), heat (FDR = 2.2E-2) or response

to metal ions (FDR=3.4E-2), suggesting that responses to host-induced stress are

compromised in the mutant, as it is also suggested by the weaker, albeit significant

enrichments of responses to host immune response (OR=1.88, FDR=7.47e-02, not

highlighted in **Fig 3E**). Furthermore, the mutant strain showed a reduction in transcripts

related with secretion (FDR=7.6e-03), and, specifically, protein secretion by the type VII secretion system (FDR=5.2 e-03). Contributing to this enrichment, we found that the key antigen ESAT-6 was significantly down-regulated in $virR^{mut}$ with respect to both WT (log2FC=-0.17, FDR=4.1E-3) and $virR^{mut}$-Comp strains (log2FC=-0.19, FDR=2.6E-3), while $cfp10$ (Rv3874), was similarly less expressed in $virR^{mut}$ than in WT (log2FC=-0.24, FDR=2.6E-3).

Second, we found enriched terms related with lipid metabolism and transport both among the sets of genes up and downregulated in the $virR^{mut}$ strain. On the one hand, genes downregulated in $virR^{mut}$ appeared enriched in fatty acid biosynthetic processes (FDR=6.7E-2), suggesting that the regulation of both anabolic and catabolic pathways of lipids, key for bacterial survival and pathogenesis upon infection (Ghazaei, 2018), are, at least in part, VirR-dependent. Furthermore, the mutant strain also shows a significant induction of a number of GO terms related with cell wall composition (e.g. cell wall organization: FDR=3.9e-2), including terms associated with the biosynthesis key components of the mycobacterial cell wall such as PDIMs (DIM/DIP cell wall layer assembly enrichment FDR=9.5e-03) and importantly peptidoglycan (peptidoglycan-based cell wall biogenesis FDR=2.2e-2) (**Fig 3F**).

Third, we found a series of enrichments related with global regulation of gene expression (positive: FDR=4.4e-2, negative: FDR=5.4e-2), and RNA biosynthesis (positive: FDR=5.4e-2, negative: FDR=5.8e-2), among genes downregulated in $virR^{mut}$, suggesting a global alteration of the transcriptional landscape in the mutant.

Fourth, we found that metabolic reprogramming in the $virR^{mut}$ strain is not restricted to lipids but also involves amino acids (cellular amino acid metabolic processes FDR=2.1e-3), and carbohydrates (cellular carbohydrate biosynthetic process FDR=6.8e-3), enriched among genes upregulated in the mutant. Similarly, energy production also appears altered in $virR^{mut}$, with related GO terms enriched among downregulated genes, such as electron transfer activity, (FDR=4.2e-2), as well as terms enriched among upregulated genes in the mutant such as NADH dehydrogenase activity, (FDR=3.3e-3). All together, these results combined with the differences concerning lipid metabolism discussed above, suggest a global remodeling of metabolic networks in $virR^{mut}$.

$virR$-dependent genes intersect the regulons of key transcriptional regulators of responses to stress, dormancy, and cell wall remodeling. To shed light on the specific transcriptional circuits whose disruption in the $virR^{mut}$ strain may lead to the differential gene expression patterns described, we resorted to the regulon annotations previously reported (Minch et al., 2015). In this work, authors collected ChIP-seq data on binding events between a panel of transcription factors (TFs) and DNA to annotate

interactions between 143 regulators and the promoters of 7248 target genes (peaks

between -150 and +70 nucleotides from transcription staring sites). Capitalizing on these

data, we interrogated whether the sets of genes up- (N=399) and downregulated in the

$virR^{mut}$ strain (N=502) were significantly enriched among each of the regulons reported

by Minch et al. A total number of 10 regulons appeared enriched among either down (9)

or upregulated genes (1) in $virR^{mut}$ (**Supplementary file S3** and **Fig 4A**, highlighted TFs

at OR>2 FDR<0.1). Furthermore, the expression of 7 out of 10 of the TFs controlling

these regulons was itself found to be $virR$-dependent in our transcriptional data (**Fig 4B**).

Genes more highly expressed in the $virR^{mut}$ strain appeared significantly enriched among

Rv0339 targets (OR=9.1, FDR=7.5E-2, **Fig 4A**), named IniR because of its role as key

regulator of the isoniazid-induced operon $iniBAC$, a key element of the response to cell

wall biosynthesis inhibition induced by isoniazid (Boot et al., 2017) and linked to vesicle

biogenesis in Mtb (Gupta et al., 2023). Several additional transcription factors known to

regulate cell wall biosynthesis and homeostasis were also enriched among genes

downregulated in the $virR^{mut}$ strain. These include SigF (OR=5.8, FDR=8.1E-2), a sigma

factor controlling cell envelope organization (Williams et al., 2007), $mtrA$ (OR=2.2,

FDR=6.0E-2), a TF controlling peptidoglycan cleavage and alterations in the context of

regulation of cell growth and division, whose disruption has recently been shown to lead

to increased permeability (Peterson et al., 2023); as well as Rv0472c (OR=11.7,

FDR=3.5E-2), a transcription factor involved in the regulation of mycolate remodeling

pathways (Peterson et al., 2019), a process used by the pathogen upon phagocytosis to

evade macrophage-induced stress and enter dormancy. Interestingly, two genes under

the control of Rv0472c, that are key to enable this process, are the mycolate desaturases

$desA1$ and $desA2$ (Peterson et al., 2019), which appear downregulated in the $virR^{mut}$

strain with respect to WT (logFC=-0.4, -0.1; FDR=2.3E-11, 3.2E-2, for $desA1$, and

$desA2$, respectively), as well as with respect to $virR^{mut}$-Comp (logFC=-0.4, -0.4;

FDR=7.2E-12, 5.3E-21). Furthermore, Rv0472c has also been linked with the regulation

of key proteins in the response to several sources of stress in Mtb, such as the putative

nitroreductase Rv3131 (response to oxidative/nitrosative stress), as well as the

triacylglycerol synthase $tgs1$, responsible for the synthesis and accumulation of

triacylglycerol, a major energy source for the bacterium during dormancy (Chauhan and

Tyagi, 2009), both of which are strongly downregulated in the $virR^{mut}$ strain vs both WT:

(logFC=-1.2, -1.7; FDR=3.1E-19, 3.2E-14) and $virR^{mut}$-Comp (logFC=-0.4, -1.9;

FDR=3.9E-5, 1.4E-17).

The adjacent genes Rv3131 and $tgs1$ (Rv3030c) appear under the control of other key

regulators of the response to intracellular stress in Mtb and the transition to dormancy,

whose regulons appear as well enriched among genes downregulated in $virR^{mut}$ (**Fig

4C). These include *devR* (OR=2.7, FDR=1.9E-5), the canonical regulator of the response to hypoxia and transition to dormancy in *Mtb* (Park et al., 2003), as well as Rv1985c (OR=4.1, FDR=4.5E-11), a key regulator of the homeostasis of the threonine biosynthesis pathway, a metabolic route needed by the bacterium for persistence *in vivo* (Hasenoehrl et al., 2019). Three of the four remaining TFs whose regulons we found enriched among *virR^mut* downregulated genes have been similarly linked to the response to assorted sources of stress that are cardinal features of the phagosome environment, such as oxidative stress, response to toxic metals, and low pH (Schnappinger et al., 2003). These include the regulator *lsr2* (OR=2.3, FDR=5.1E-4), a global histone-like protein responsible from protecting the DNA physically from oxidative stress (Bartek et al., 2014), the zinc-uptake regulator Zur (*furB*) (Maciąg et al., 2007) (OR=8.8, FDR=4.5E-2), and *tcrX* (OR=2.6, FDR=5.0E-2), part of the recently characterized acid-sensing two-component system tcrXY (Stupar et al., 2024). Notably, we observed that removing catalase from culture medium prior to submitting *virR^mut* to a source of oxidative stress like H202 for 8 h, significantly affected its growth relative to WT and *virR^mut*-C strains (**Fig S3**). This result validates the transcriptomics data reflecting a defect in regulating oxidative stress in *virR^mut* and encourage to investigate its connection with vesicle production.

Cell envelope permeability is a major driver of MEV release. If enhanced cell permeability is a global consequence of *virR* disruption that leads to enhanced vesiculation, we hypothesized that other conditions either inducing or decreasing cell envelope permeability could lead to alterations in vesicle production in *Mtb*. To test this, we treated *Mtb* with sublethal concentrations of thioridazine (TRZ), a known enhancer of cell permeability (De Keijzer et al., 2016); and, alternatively, we cultured the bacillus with cholesterol as a sole carbon source, a condition which reduces mycobacterial cell permeability (Brzostek et al., 2009) (**Fig 5A**). The enrichment of the DevR regulon among the genes that are downregulated in *virR^mut* is also suggestive of the existence of a convergent regulatory route controlling both cell permeability and vesiculation under control of this transcription factor, whose activation has been described as a key part of the pathogen's transcriptomic response to cholesterol (Soto-Ramirez et al., 2017), concomitant to a reduction in cellular permeability (Brzostek et al., 2009). Importantly, neither cholesterol nor TRZ altered bacterial viability as measured by the absence of the release of the transcription factor IdeR, as an indication of cell lysis or by spotting serially diluted cultures on solid medium from the corresponding conditions (**Fig S2C,D**). We then measured vesicle levels in culture supernatants and found that *Mtb* significantly releases more EVs upon treatment with TRZ, and significantly reduces vesicle release

in the presence of cholesterol as a sole carbon source (**Fig 5B**), as measured by dot blot after normalization to colony forming units (CFUs). These results strongly suggest that cell envelope permeability determines the magnitude of the vesiculation phenomenon in *Mtb*. We next investigated whether common transcriptional features underpin the regulation of cell envelope permeability in the tested conditions. To do that, we compared the transcriptional response of *virR^mut^* with that of available datasets including *Mtb* cultured in the presence of 0.01% of cholesterol as a sole carbon source (Pawełczyk et al., 2021), and *Mtb* treated with 6 μg ml-1 of TRZ (Dutta et al., 2010). Experimentally, we retrieved the set of genes up and downregulated in *Mtb* grown in cholesterol vs glucose-rich culture media (Brzostek et al., 2009), and interrogated whether those genes were enriched among the sets of differentially expressed genes in *virR^mut^* vs WT, and *virR^mut^* vs *virR^mut^*-Comp reported in this study (**Fig 5C**). Interestingly, we found that those genes whose expression changed in the same direction in conditions associated with low, or high permeability, in both studies were mutually enriched, while the cross-comparisons between sets with opposite responses to low-permeability-associated treatments in both studies led to significant depletion statistics. These results validate the initial hypothesis according to which changes in permeability induced by these two treatments are linked to the activation of a common transcriptional program. The inspection of genes that are simultaneously upregulated in treatments associated with low permeability in both studies identified DevR as a putative regulator associated with the orchestration of this transcriptional footprint of low cell permeability. The two-component system DevS-DevR (also known as DosS-DosR) has a well-known role as a global regulator of the transition to dormancy, and its regulon is consistently upregulated in non-permeable dormant bacteria (Honaker et al., 2009).

Interestingly, authors found that low-permeability bacteria cultured in a cholesterol medium, albeit featuring a regular growing phenotype, activated the DosR regulon, suggesting that cell wall remodeling leading to a decrease in permeability linked with the induction of DosR can be triggered independently to the transcriptional regulatory programs leading to growth arrest and dormancy in *Mtb* (Pawełczyk et al., 2021). Importantly, the integration of the genes that are downregulated in *virR^mut^* validates this hypothesis (**Fig 5D and FigS4**), since not only the intersection between the DevR regulon and the set of genes downregulated in *virR^mut^* are significantly larger than expected by chance (see **Fig 4A**, OR=2.67, p=2.7E-7), but also the intersection between DevR regulon and the genes that are activated by cholesterol in Brzostek et al. data (OR=1.89, p=3.5e-5). 30 out of the 173 genes annotated in the DosR regulon described in (Minch et al., 2015) are simultaneously upregulated in conditions linked to low permeability in both studies.

A similar analytic approach for the comparison of the response of *virR^mut* and the

response to the drug TRZ, a treatment well-known for inducing an increase in

permeability in *Mtb*-treated cells (Dutta et al., 2010) (GSE16626), led to conceptually

similar results, with genes simultaneously downregulated in both TRZ and *virR^mut*

(OR=1.96, p=9.1E-4) being enriched. Interestingly, among the 22 genes showing the

largest, and most significant repression effects by both treatments (log2FC>0.25 &

FDR<0.05 simultaneously, for TRZ effects, *virR^mut* vs WT), we found 4 genes whose

functions are related with metal nutrients uptake and homeostasis, such as genes

encoding the bacterioferritin BrfB, the metal exporter CtpJ, the cupper regulator CsoR

and Moew, involved in Molibdenum-cofactors biosynthesis (**Fig 5E**). Taken together,

these analyses highlight the transcriptional parallels between *virR^mut* and both TRZ and

cholesterol conditions, which strongly suggests the existence of a common

transcriptional signature associated with the control of cell permeability.

Modulation of EV levels and permeability in *virR^mut* by cholesterol and TRZ. We

next wondered about the effect of culturing *virR^mut* on both cholesterol or TRZ could have

on cell growth, permeability and EV production. In the case of cholesterol, it has also

been shown to affect other aspects of physiology (redox, respiration, ATP), which can

directly affect permeability (Lu et al., 2017). We monitored *virR^mut* growth in MM

supplemented with either glycerol, cholesterol as a sole carbon source, or TRZ at 3 ug

ml^{-1} for 20 days. While cholesterol significantly enhanced the growth *virR^mut* after 5 days

relative to glycerol medium, supplementation of glycerol medium with TRZ restricted

growth during the whole time-course (**Fig S5A**). The study of cell permeability in the

same conditions indicated that the enhanced cell permeability observed in glycerol MM

was reduced when *virR^mut* when cultured with cholesterol as sole carbon source.

Conversely, the presence of TRZ increased cell permeability relative to the medium

containing solely glycerol (**Fig S5C**). As we have previously observed for the WT strain,

either condition (Chol or TRZ) also modified vesiculation levels in the mutant accordingly

(**Fig S5B**). These results strongly indicates that other aspects of mycobacterial

physiology besides permeability are also affected in the absence of VirR and may

contribute to the observed enhanced vesiculation.

virR* regulates the protein content of EVs in *Mtb. To capture the impact of

transcriptional adaptation to the proteome, and the degree of divergence between

transcriptional shifts and proteins changes observed in the absence of *virR*, we next

performed label-free peptide mass spectrometry (LF-MS) of whole cell lysates (WCLs)

of *virR^mut*, WT, and *virR^mut*-Comp strains. Additionally, we isolated EVs from axenic

cultures and obtained their proteomic composition to examine for potential VirR-dependent differences in protein cargo that may alter the functionality of EVs (**Fig 6A**). As a result, we obtained proteomic profiles across strains and cell compartments for a total of N=935 proteins. Upon PCA analysis of the proteomic data (**Fig 6B**), we found that the first principal component (PC1), explaining 36.9% of total variance, separated samples from WCLs and EVs in all strains (PC1 differs across cell compartments: one-tailed Mann Whitney test p=2.1e-05), while the second PC (PC2) (13.6% of total variance explained) captured strain effects on the proteomic footprint only among EVs (one tailed Mann Whitney t test for PC2 differences between *virR^mut* EVs and those from either WT or *virR^mut* -Comp strains (one-tailed Mann Whitney p=1.2 e-02). These results suggest that differences in protein abundances between EVs and WCLs are larger than differences found between strains, and that, concerning the differences across strains, these are more important among EVs. To verify this, we conducted differential protein expression analyses to find barely 27 proteins differentially expressed between *virR^mut* and any of the two *virR*-expressing strains, which represents 2.9% of the total of proteins under analyses (**supplementary file S4**). Despite the scarcity of significant differences in expression across strains in WCLs, some of the features most differentially expressed in *virR^mut* are, besides VirR itself (**Fig 6C**), proteins involved in key processes for bacterial survival and virulence, including key metabolic enzymes such as the Acetyl-coenzyme A synthetase Acs (logFC = -3.92 (FDR=4.5e-3) and -2.96 (FDR=5.2E-2) for the differences between *virR^mut* and WT, and between *virR^mut* and *virR^mut*-Comp, respectively), as well as FprA, (logFC= -3.12, FDR=4.5e-3 e-02 for *virR^mut*-WT and logFC= -2.16, FDR=5.9 e-02 for *virR^mut*- *virR^mut*-Comp), involved in the reduction of NADP+ into NADPH, both downregulated in *virR^mut*.

Conversely to the WCL dataset, we did find more frequent differences in protein expression across strains in EVs, with as many as 361, and 382 proteins differentially expressed in *virR^mut* EVs when compared to those from WT and *virR^mut*-Comp strains, respectively (**supplementary file S4**, **Fig 6D**). This represents 38.6 and 40.9% of all proteins tested, respectively. Similarly to what was observed at the level of RNA expression, differences in expression for the contrasts (*virR^mut*-WT) and (*virR^mut*- *virR^mut*-Comp) were directionally enriched (**Fig 6E**), thus motivating the definition of two sets of proteins that 1) were observed at higher levels in the EVs isolated from *virR^mut* than in those from both WT and *virR^mut* -Comp strains (N=117 *virR^mut* -upregulated proteins); or 2) were observed at lower levels in the EVs from *virR^mut* than in those from any of the other two strains (N=108 *virR^mut*-downregulated proteins). Interestingly, as many as 21 out of the 117 proteins that appeared enriched in *VirR^mut* EVs were also upregulated in *virR^mut* at the transcriptional level (One-sided Fisher exact test of enrichment OR=2.02,

p=7.0e-3) suggesting that at least a part of these differences in EVs' protein cargo are a consequence of differences in gene expression already observed at a transcriptional level. Notwithstanding that, proteins appearing at lower concentrations in $virR^{mut}$ EVs were not enriched among genes bearing significant transcriptional effects in the mutant. Regarding downregulated EV proteins in $virR^{mut}$ we observed robust enrichments in GO terms related with protein biosynthesis, including translation (FDR=9.8 e-22), ribosome (FDR=5.3 e-29), and ribosome biogenesis (FDR=4.1 e-27, **Fig 6F** and **supplementary file S5**). Consistent with these results, 44 ribosomal proteins (out of 49 detected) were found to be significantly less expressed in the $virR^{mut}$ strain relative to either WT or $virR^{mut}$-Comp strains (FDR<0.05 for at least one comparison), while no ribosomal protein appeared significantly upregulated in $virR^{mut}$. Furthermore, 30 of these proteins were simultaneously downregulated in $virR^{mut}$ when compared to both VirR-containing strains (**Fig S6**) implying a stark ribosomal depletion in the EVs in the absence of VirR. Furthermore, we found that the two bacterioferritin proteins BfrA and BfrB were simultaneously less abundant in $virR^{mut}$ than in both WT and $virR^{mut}$-Comp strains (logFC<-1.5 and FDR<7.3E-4, for differences $virR^{mut}$ vs WT and $virR^{mut}$ vs $virR^{mut}$-Comp in both genes, **Fig 6G**, and **supplementary file S4**), together contributing to the enrichment of the GO term ferroxidase activity (FDR=6.4 e-02, **Fig 6F**). This result is suggestive of a possible defect of $virR^{mut}$ EVs in iron sequestration (Mohammadzadeh et al., 2021; Prados-Rosales et al., 2014b). We also found a series of metabolic pathways enriched among proteins upregulated in the EVs of the $virR^{mut}$ strain, including enzymes involved in metabolism of nucleotides (e.g. purine nucleotide biosynthetic process FDR=2.0e-02, amino acids (alpha amino acids FDR=8.3e-04), as well as carbohydrates (hexose metabolic process FDR=6.3e-2), whose catabolism appears also enriched among proteins downregulated in the mutant (e.g. glycolytic process FDR=2.2e-2 and carbohydrate catabolic process FDR=4.7E-2). The simultaneous enrichments of these metabolic pathways observed both among upregulated and downregulated vesicular proteins (**Fig S7**), indicates a complex, VirR-dependent, rewiring of metabolic routes in the vesicles from the mutant strain.

Further, proteins upregulated in $virR^{mut}$ EVs (N=117) were enriched in terms related to mycolic acids metabolic processes (FDR=2.1E-2), and peptidoglycan-based cell-wall biogenesis (FDR=9.0 e-02). Proteins such as InhA (logFC=3.71, 2.26 FDR=2.20 e-05, and 1.4E-3 for comparisons $virR^{mut}$ vs WT and $virR^{mut}$ vs $virR^{mut}$-Comp, respectively), and KasB (logFC=1.83; 2.49 FDR=2.23 e-02; 4.4e-3), involved in mycolic acid metabolism were also upregulated in $virR^{mut}$ (**Fig 6H**).

The upregulation of these proteins, along with the enrichments at the transcriptional level of DIM/DIP biosynthetic pathways (upregulated in $virR^{mut}$), suggests a shift in the control

of the pool of propionyl-CoA, both at the level of its production, -as a byproduct of the catabolism of methyl-branched lipids, odd-chain-length fatty acids as well as cholesterol- and its detoxification by the bacterium through different routes, such as the methylmalonyl pathway, the glyoxylate cycle and the methylcitrate cycles (Lee et al., 2013). To shed light on these changes at the metabolic level, we conducted a metabolic profiling to quantify the concentration of metabolites that are central in these different detoxification routes including propionyl-CoA: acetyl-CoA, pyruvate, methylcitrate, and methylmalonyl-CoA (**Fig 6I**). Only methylmalonyl-CoA was found to be significantly repressed in $virR^{mut}$ (p=0.028). We also observed differences between $virR^{mut}$ and $virR^{mut}$ -Comp strains in methylcitrate. This result, along with the trend towards increased protein, and especially gene expression that is observed for genes involved in the biosynthesis of PDIMs from methylmalonyl (including phenolpthiocerol synthesis type-I polyketide synthases A-E, (ppsA-E, **Fig 6I**), among others), suggests that the methylmalonyl pool is depleted in the mutant strain because of an increased activity of the PDIM biosynthetic pathway. Furthermore, most of the subunits of the acetyl/propionyl-CoA carboxylase Acc (AccA1, AccA2, AccA3, AccD1, AccD5 appear repressed in $virR^{mut}$ EVs (**Fig 6I**), suggesting that the production of methylmalonyl Co-A from the degradation of propionyl CoA is inhibited within EVs of the mutant strain.

Taken together, the omics analyses align with the structural and functional differences observed in $virR^{mut}$. We reasoned that the increased expression of PDIMs biosynthetic genes and the reduced expression of genes related to protein synthesis may be triggered to compensate the enhanced permeability measured in $virR^{mut}$. In addition, this enhanced permeability leads to an increase in the release of EV, which seems to be depleted in ribosomes and iron storage proteins.

The absence of VirR leads to an aberrant enlargement of PG in *Mtb*. To elucidate the specific molecular defects that explain the periplasm enlargement, the enhanced permeability, and the antibiotic sensitivity phenotypes, we performed a biochemical analysis of WT and $virR^{mut}$ cell envelopes. First, we analyzed the total lipid profile by liquid chromatography-mass spectrometry (LC-MS) and observed significant differences in prevalent polar lipids linked to the cell membrane including, lysophosphatidic and phosphatidic acid, species of monoacylated and diacylated glycerol, phosphatidylethanolamines and phosphatidyl-myo-inositol mannoside (PIM) species, being more abundant in $virR^{mut}$ relative to WT and complementing strain. (**Fig 7A**). To validate these results, we ran thin layer chromatography (TLC) experiments on the same lipid extracts and found that the mutant carries more phospholipids in the membrane including more $AcPIM_2$ and PE (**Fig 7B**). Moreover, we could not measure major

differences in the level of mycolic acid species (**Fig 7C**), while significantly more PDIMs were detected in the mutant relative to the other strains (**Fig 7D**). This later result partially validates proteomics and metabolomic data and suggest that the methylmalonyl pool is depleted in the mutant strain because of an increased activity of the PDIM biosynthetic pathway. However, other lipids including SL1 or PAT/DAT that share a common precursor (propionyl- CoA) with PDIM may also explain the depletion of methylmalonyl. Notably, we observed a reduction in the level of these lipids (SL1 or PAT/DAT) in *virR^{mut}* cultured in glycerol relative to WT and complemented strains, suggesting that the excess of PDIM synthesis can take away methyl malonyl CoA from the biosynthesis of SL-1 and PAT/DAT in the absence of VirR (**Fig S8**).

When analyzing apolar lipids, we also observed increased levels of free mycolic acids (FMA) and free fatty acids (FFA) upon *virR* deletion (**Fig 7E**). The enhanced release of these apolar lipids has also been documented in a *C. glutamicum* mutant in Cg0847 (Baumgart et al., 2016), a canonical LCP protein. Importantly, we could not detect any of these lipids in EVs from *virR^{mut}* (**Fig 7F**), while major polar phospholipids could be resolved in EVs from all strains (**Fig 7G**), as previously reported (Prados-Rosales et al., 2011). This result indicates that *Mtb* releases free apolar lipids independently from membrane associated EVs.

Next, we performed a glycosyl composition analysis of isolated cell walls (including mycolyl-arabinogalactan-peptidoglycan complex) and determined that there were no significant differences between strains in AG content as well as in the ratio between arabinose and galactose (A/G) (**Fig 8A**). Similarly, no changes were observed in the galactan length as measured by the ratio between galactose (G) and rhamnose (R) G/R (Justen et al., 2020). We noticed that the ratio between N-acetylglucosamine (NAGc) and rhamnose (R) (NAGc/R) was significantly higher in *virR^{mut}* relative to WT, suggesting an increase in PG content in this strain (**Fig 8A**). The validation of these results was performed independently by measuring the amount of diaminopimelic acid (DAP) in isolated cell walls, as a surrogate of PG content, and determined that *virR^{mut}* carries two times more PG than WT or complemented strains (**Fig 8B**).

We next obtained pure PG preparations of each strain and performed an analysis by atomic force microscopy (AFM) (**Fig 8C and Fig S9**). This technique has been recently used to investigate the nanometric features of isolated Gram-positive bacterial cell walls (Pasquina-Lemonche et al., 2020). Semi-automated analysis of liquid AFM images from mycobacterial isolated PG samples revealed that: (i) the absence of VirR leads to a significant reduction in PG pore diameter (down to 9.2 ± 0.6 nm) relative to WT (16 ± 3.3 nm) and complemented strains (10 ± 0.8 nm) (**Fig 8C, D and Fig S9**); (ii) the number of pores is significantly higher in *virR^{mut}* compared to WT and complemented strain (**Fig**

8E and Fig S9); (iii) manual analysis of AFM images in ambient conditions revealed that

virR^{mut} PG is significantly thicker than that of the other strains of the study (**Fig 8F**). These

results show for the first time the nanometric architecture of isolated PG in mycobacteria

by AFM and suggest a role for VirR in regulating PG thickness and porosity. Strikingly,

we found that the PG enlargement observed in the *virR^{mut}* correlated with increased

susceptibility to the muramidase lysozyme (**Fig S10**), indicating modifications in PG

remodeling in *virR^{mut}*. In fact, a deeper MS analysis of PG showed a significant increase

in the abundance of deacetylated muropeptides in *virR^{mut}* relative WT and *virR^{mut}*-Comp

strain (**Fig 8G, H**). This PG feature has been linked to variability in lysozyme susceptibility

in other Gram positive bacterial species (Bera et al., 2005). Collectively, although the

structural changes at nanoscopic level of the peripheral areas of the cell observed by

AFM leads to a more dense and thick structure harder to penetrate, the chemical

changes in the peptidoglycan structure observed and biochemical assays suggest that

changes associated with alterations in VirR-regulated PG remodeling makes the wall

more permeable and support increased release of EV in *Mtb*.

VirR interacts with canonical LCP proteins. As mentioned above VirR is a LytR_C

solo domain protein with no catalytic LCP domain. Other members of the LCP protein

family like Rv3267 and Rv3287 are the main PG-AG ligases (Grzegorzewicz et al., 2016;

Harrison et al., 2016) and also carry a LytR_C domain. We have previously shown that

VirR interacts with itself (Rath et al., 2013). Consequently, we reasoned that if VirR

contributed to PG-AG ligation it could do it via interaction with canonical LCP proteins

through the LytR_C domain. Although VirR is one of the most abundant proteins found

in tuberculin preparations (Prasad et al., 2013), we determined that it localizes at sites

of nascent cell wall by fluorescence microscopy (**Fig S11**). Therefore, its location could

initially allow for interactions with other cell surface proteins like LCP. Both recombinant

VirR and the main AG-PG ligase Lcp1 (Rv3267) oligomerize in a blue-native PAGE gel

(**Fig 9A**). This observation suggests that they interact with proteins sharing similar

domain features. We next performed a flotation assay where we examined the behavior

of VirR on an iodixanol gradient in the presence or absence of Lcp1. Using a specific

antibody raised against VirR, we observed that the presence of Lcp1 displaces the

relative location of VirR toward upper fractions of the gradient, indicating a potential

interaction of both proteins and the formation of higher molecular weight complexes (**Fig

9B**). To investigate the interaction of virR and Lcp1, along with other Lcp proteins, in a

more physiological environment we used the bipartite splitGFP approach (Cabantous et

al., 2013) (**Fig 9C and Fig S12**), which has been recently utilized to demonstrate the co-

localization and interaction of the membrane-bound metal ATPase CtpC with the

chaperon PacL1 (Boudehen et al., 2022). We first recapitulated the VirR self-interaction

and added that such interaction implicates the LytR_C domain since no differences in

fluorescence were measured between full and truncated versions of VirR (VirRsol)

lacking the first 41 amino acids. As negative controls, we included interactions between

VirR and the cell surface associated metal ATPase CtpC (**Fig 9C**). When testing

interactions between VirR and other Lcp proteins (Rv0822c, Lcp1 (Rv3267) and Rv3284)

we could not measure any GFP signal when full Lcp proteins were used in the assay.

Conversely, a significant increase in fluorescence was measured for all combinations

when truncated Lcp proteins including either the C-terminal domain or specifically the

LytR_C domain were tested (**Fig 9C**). These results strongly suggest that VirR interacts

with Lcp proteins and that these interactions occur via LytR_C terminal domains.

DISCUSSION

We reported *virR* as the first gene implicated in the biogenesis of EVs in *Mtb* ten years

back (Rath et al., 2013). A transposon mutant in *virR* (*virR^{mut}*) overproduces MEVs and

this provokes augmentation of cytokine responses in mouse and human macrophages.

This mutant manifested an attenuated phenotype in experimental infections on

macrophages and mice (Beaulieu et al., 2010; Rath et al., 2013). Such studies

established that VirR has a role in virulence by negatively modulating the release of

MEVs. These studies provided evidence that vesiculogenesis in *Mtb* is genetically

regulated. Since then, conditions such as iron starvation (Prados-Rosales et al., 2014b)

and other genes, including the bacterial dynamin-like *iniAC* (Gupta et al., 2023), and the

Pst/SenX3-RegX3 signal transduction system (White et al., 2018) have been shown to

contribute to the biogenesis of MEVs. Interestingly, VirR, a LytR_C solo domain protein

(**Fig S1**), shares homology with Lcp proteins, which in *Mtb* are known to participate in

the enzymatic connection between AG and PG (Grzegorzewicz et al., 2016; Harrison et

al., 2016). This suggests VirR function might be linked to cell wall remodeling.

In this study, we investigated how VirR can control the magnitude of MEV release

in *Mtb*. We did this by studying the underlying enhanced vesiculogenesis phenomenon

observed in *virR^{mut}* using genetic, transcriptiomics, proteomics and ultrastructural and

biochemical methods. It was clear from our cryo-EM studies that *virR^{mut}* has an aberrant

cell wall as the layer above the cell membrane was significantly larger than that of WT

and complemented strains. The enlarged compartment showed higher granular texture,

a feature that has been previously associated to PG precursors in unrelated Gram-

positive bacteria (Zuber et al., 2006). This observation and previous studies showing the

increased sensitivity of *VirR^{mut}* to vancomycin (Ballister et al., 2019) strongly suggested

that the cell wall defect is connected to PG remodeling. The use of AFM on isolated PG

confirmed this and showed for the first time the nanostructure of PG in *Mtb*, providing information on PG thickness, pore size and number of pores. AFM indicated that *virR^{mut}* PG is thicker, has significantly more pores of small size. Considering the pore sizes measured across strains (ranging from 9-16 nm), we initially ruled out the notion that EVs (which are spheres of 50-300 nm in diameter) are released through the peripheral cell wall. Therefore, *Mtb* must have an alternative mechanism to release EVs, possible during the division process where the PG is most fragile, presumably due to defects in crosslinking, something that awaits further experimental verification. These events were not possible to visualize in AFM given that PG fragments from all the strains were lacking the poles or clearly defined division sites but could be the subject of further studies in the future. However, we cannot exclude that MEV plasticity could allow for crossing the cell wall via PG pores.

A recent study showed that *Corynebacterium glutamicum* can produce and release different types of MVs through different routes, including mycomembrane blebbing or bubbling cell death, depending on the stress at which *C. glutamicum* is exposed (Nagakubo et al., 2021). In the present study, we have shown that *virR^{mut}* overproduces EVs under normal growing conditions in the absence of cell lysis, according to our assay which measured cytoplasmic leaking of the transcription factor IdeR or serially diluted spots on solid medium. Overall, we show that EV production in *Mtb* under our growing conditions, is not linked to cell death. This fact largely excludes the bubbling cell death mechanism. Moreover, our lipidomic analysis now and before (Prados-Rosales et al., 2011) indicates the absence of mycomembrane lipids in isolated MEVs from all strains, ruling out that MEVs originate at the mycomembrane. These apparent disagreements with the *C. glutamicum* study (Nagakubo et al., 2021) might indicate that *Mtb* EV biogenesis may differ from that of *C glutamicum*, and other mycolic acid-containing bacteria with shorter fatty acids. Additional variables that may explain differences between studies is the growth media and the time at which vesicles are isolated (Gupta et al., 2023). Of note, *M. smegmatis* EVs were isolated at day 6 (Nagakubo et al., 2021), a time at which this fast-growing mycobacteria is already in stationary phase.

The transcriptional profile of *virR^{mut}* can provide a partial explanation for the attenuated phenotype of the mutant due to a downregulation of genes activated upon stresses associated to the intraphagosomal environment. It is particularly interesting to observe the specific reduction in transcript levels of the two most important effectors of the esx-1 secretion system. On the other hand, genes involved in PDIM synthesis were upregulated. It is possible that the specific upregulation of PDIM synthesis in absence of *virR* is part of compensatory response to a reduced permeability. We were surprised to

find no differences between protein profiles in whole cells between WT and *virR^{mut}*
strains, despite difference in transcriptomics. A biological interpretation of these results
would point to the existence of global compensatory mechanisms of post-transcriptional
regulation used by the bacteria to partially recover comparable proteomic profiles in
virR^{mut} to those observed in the WT.
Contrary to whole cell lysates, EV protein profiles between *virR^{mut}* and WT were
largely different. In this context, it is accepted that certain proteins and lipids are enriched
in bacterial EVs and that preferential inclusion of such antigens in MVs supports a
specific vesicle biogenesis mechanism. In Gram positive bacteria and mycobacteria both
membrane and cytosolic proteins are included in EVs since the origin of these structures
is linked to the cell membrane (Prados-Rosales et al., 2011) and, presumably, the
incorporation of intracellular proteins may occur during the process of vesicle formation.
We found a higher abundance of proteins related to PG and MA synthesis in *virR^{mut}* EVs
relative to WT. Therefore, these two processes seem to contribute to EV biogenesis in
Mtb. Additional data presented in this study, which includes ultrastructural and chemical
analyses confirm this. Both PG and MA synthesis are complex processes, including a
diverse set of genes. What parts of the biogenesis of these molecules are important for
the formation of EVs is a matter of future investigations.
Another finding of the study is the connection between permeability and the
magnitude of the vesiculation process in *Mtb*. We linked these two phenomena first, by
measuring enhanced vesiculation and permeability in the same strain (*virR^{mut}*) and
second, by modifying the permeability using chemical (via sublethal exposure to TRZ)
and nutritional supplements (via providing cholesterol as sole carbon source) in WT *Mtb*.
TRZ, a repurposed drug that belongs to the class of phenothiazines, is known to affect
respiration in *Mtb* (Amaral et al., 1996). It is also assumed that TRZ can inhibit efflux
pumps leading to an increased susceptibility to first line antitubercular drugs when tested
in combination (De Knegt et al., 2014). This provides an explanation for the synergistic
effects of such combinations. An independent explanation of such synergy may also
come from the increased permeability, as it has been recently shown by a quantitative
proteomics analysis of *Mtb* cells exposed to sublethal concentration of TRZ (De Keijzer
et al., 2016). When we grew *Mtb* with cholesterol as a sole carbon source, a condition
that slows mycobacterial growth but also reduces cell permeability via cholesterol
accumulation in the cell wall (Brzostek et al., 2009), we observed the opposite trend on
MEV biogenesis. We consider this finding of high relevance since *Mtb* permeability is
one the major concerns when developing potential antimycobacterial drugs.
Nevertheless, bacterial cell permeability is a complex property that it is controlled by
physiochemical, biological, and chemical processes such as stereochemistry,

lipophilicity, saturation and unsaturation, flexibility, viscosity, fluidity, pressure,
temperature and physiological conditions (Nagamani and Sastry, 2021). How these
variables change to allow the release of MEVs is not known.

This study also proposes a model VirR function whereby VirR may work as a
scaffold for canonical Lcp proteins, which in *Mtb* links PG to AG. Based on the
demonstration that VirR interacts with canonical Lcp proteins, we believe that the
absence of VirR deregulates the localization of canonical Lcp proteins. Although we
cannot rule out the contribution of other phenomena including alteration of the PG
molecule, VirR is a central scaffold in the cell wall remodeling process. Taking into
consideration the fact that $VirR^{mut}$ is attenuated in virulence on preclinical models of
infections and manifests an enhanced vesiculogenesis and permeability, VirR may
represent a novel and interesting drug target. We envision that targeting VirR may not
only make *Mtb* weaker and more permeable to other antitubercular drugs, but to
stimulate the immune system, via MEVs, to fight infection.

MATERIALS AND METHODS

Bacterial Strains, media and growth conditions. *Mtb* H37Rv (ATCC) and derivative
strains $virR^{mut}$ and $virR^{mut-}C$ (complemented strain) (Rath et al., 2013) were used in this
study. Bacteria were initially grown in Middlebrook 7H9 medium (7H9) supplemented
with 10% (v/v) oleic acid-albumin-dextrose-catalase (OADC) enrichment (Becton
Dickinson Microbiology Systems, Spark, MD, United States), 0.5% (v/v) glycerol and with
Tyloxapol 0.05% (v/v; Sigma-Aldrich, Burlington, MA, United States) prior to inoculation
in a minimal medium (MM) consisting of KH_2PO_4 1 g/l, Na_2HPO_4 2.5 g/l, asparagine
0.5 g/l, ferric ammonium citrate 50 mg/l, $CaCl_2$ 0.5 mg/l, $ZnSO_4$ 0.1 mg/l, without
Tyloxapol 0.05% (v/v), containing 0.7% (v/v) glycerol, pH 7.0.

Mtb was also cultured in MM in the presence of sublethal concentrations of
thirodazine (TRZ) of 6 µgml⁻¹ as previously described. We determined that this
concentration does not affect cell growth. *Mtb* was also cultured in MM with cholesterol
as a sole carbon source as previously described (Brzostek et al., 2009).

Mtb expressing fluorescent VirR was generated by cloning VirR into pMV261-
Venus using fast cloning approach (Li et al., 2011).

M. smegmatis mc²155 (ATCC 700084) and recombinants derived from this strain
were grown at 37°C in 7H9 medium (Difco) supplemented with 10% albumin-dextrose-
catalase (ADC, Difco) and 0.05% Tween-80 (Sigma-Aldrich), or on complete 7H11 solid
medium (Difco) supplemented with 10% OADC(Difco). When needed, streptomycin (25
µg ml⁻¹) was added to the culture media. *Escherichia coli* strains StellarTM (Takara bio,

San Jose, CA, United States) were grown at 37°C in LB broth (Difco) or on L-agar plates

(Difco) supplemented with streptomycin (25 μg ml^{-1}) when required.

***M. smegmatis* expression vectors for split GFP experiment.** Plasmids used for split

GFP experiments were constructed by modifying the plasmid pGMCS-PpacL1-pacL1-

Lk2-GFP11-ctpC(1-85)-Lk15-GFP(1-10) (Boudehen et al., 2022) (**Table S1**). Plasmids

pGMCS, conferring resistance to streptomycin, are shuttle vectors, episomal in *E. coli*

and integrative in mycobacteria through insertion at the attL5 mycobacteriophage

insertion site in the glyV tRNA gene. Lk2 (LEGSGGGGSGGGS) and Lk15

(GPGLSGLGGGGGSLG) are flexible linkers separating the protein domains of interest

from the last 11th b-sheet or the first ten b-sheets of GFP, respectively. First, the Zinc-

inducible PpacL1 promoter was replaced by P1, a strong constitutive promoter from *M.*

smegmatis (Ariyachaokun et al., 2020), resulting in pGMCS-P1-pacL1-Lk2-GFP11-

ctpC(1-85)-Lk15-GFP(1-10). Second, the pacL1 (Rv3269) sequence was replaced by

that of virR (encoding amino-acids 1 to 164) or virRsol (encoding amino-acids 42 to 164)

by In-Fusion HD cloning (Takara). PCR fragments were amplified using appropriate

primer pairs (**Table S1**) and Phusion High-Fidelity PCR Master Mix with GC Buffer

(Thermo Scientific, Walthman, MA, United States). Linear fragments were purified on

agarose gels and inserted into pGMCS backbone linearized by PCR amplification with

appropriate primers, using the In-Fusion HD Cloning Kit (Takara), following the

manufacturer's instructions. Finally, the ctpC sequence was replaced on the resulting

plasmids by the virR or virRsol sequences, or that encoding either the C-terminal part or

only the LytR_C domains of the Rv0822c, Rv3267 or Rv3484 proteins from *M.*

tuberculosis H37Rv. Plasmids were constructed by transformation into *E. coli* Stellar

recipient cells. All plasmids were verified by sequencing before introduction by

electroporation into *M. smegmatis* mc²155 strain.

H$_2$0$_2$ sensitivity assays. Bacterial strains were initially cultured in 7H9+ADN until they

reached an OD600nm of 0.5. Cells were washed in PBS and transferred to roller bottles

including the same medium and 13 mM H$_2$0$_2$. After 8 h, bacterial cultures were serially

diluted and spotted onto 7H10 solid medium.

Generation of the CRISPR mutants. To generate the conditional CRISPR mutants we

took advantage of the CRISPR interference (CRISPRi) technology (Rock et al., 2017).

Briefly, we designed sgRNA oligonucleotides containing the target Rv0431 (*virR*) and

Rv2700 (*cei*) genes sequences (**Table S1**) located immediately upstream of the PAM

sequence. The strengths of the PAM sequence were chosen following authors

recommendation to get high and intermediate gene repression, 158.1 or 42.2-fold
repression for *virR* and 145.2 and 47.3 for *cei*, respectively. The primers were annealed
and ligated into the gel purified BsmBI-digested CRISPRi vector (pIJR965 plasmid)
followed by transformation of competent *Mtb* cells via electroporation. Transcriptional
knockdown of genes were quantified by qRT-PCR; and protein levels were examined by
immunoblot using specific antibodies raised against VirR (Rath et al., 2013) and Cei
(Ballister et al., 2019) .

EV isolation and purification. *Mtb*-EV were prepared and purified as previously
described. Briefly, the culture filtrate of 1 L cultures of *Mtb* grown in low iron MM without
detergent for 14 days was processed for vesicle isolation by differential centrifugation. In
parallel a 2 ml culture in MM supplemented with 0.05% Tyloxapol to disperse bacterial
clumps was used to determine viability by plating culture dilutions onto 7H10 agar plates
and enumerating colony forming unit (CFU) at the time of culture filtrate collection.
The membranous pellet containing vesicles obtained after ultracentrifugation of the
culture filtrate at 100,000 × g for 2 h at 4°C, was resuspended in 1 ml sterile phosphate-
buffered saline (PBS) and overlaid with a series of Optiprep (Sigma-Aldrich) gradient
layers with concentrations ranging from 30% to 5% (w/v). The gradients were centrifuged
at 100,000 × g for 16 h. At the end 1 ml fractions were removed from the top, diluted to
20 ml with PBS and purified vesicles recovered by sedimentation at 100,000 × g for 1 h.
Vesicle pellets were suspended in PBS before analysis. Protein concentration was
measured by Bradford assay (Bio Rad, Hercules, CA, United States) according to
manufacturer instructions. For all conditions test from which EVs were isolated,
normalization was performed by referring EVs concentration to colony forming units
(CFUs).

Nano particle tracking analysis (NTA). NTA was conducted using ZetaView (Particle
Metrix, Winning am Ammersee, Germany). Instrument calibration was performed prior
to EV analysis using 102 nm polystyrene beads (Thermo Fisher Scientific), according to
manufacturer instructions. Measurements were performed using a 405 nm 68 mW laser
and CMOS camera by scanning 11 cell positions and capturing 60 frames per position
at 25 C with camera sensitivity 85, shutter speed 100, autofocus and automatic
scattering intensity. Samples were diluted in pre-filtered PBS to approximately 106-107
particles·ml^{-1} in Millipore DI water. Analysis was performed using ZetaView Software
version 8.05.12 SP1 with a minimum brightness of 30, maximum brightness of 255,
minimum area of 5, maximum area of 1000, and s minimum trace length 15. Triplicate
videos of each sample were taken in light scatter mode. Particle size and concentration

were analyzed using the built-in EMV protocol and plotted using Prism software, version

8.0.1 (GraphPad Inc., San Diego, CA, United States).

Electron-Microscopy. Cells were fixed with 2% glutaraldehyde in 0.1 M cacodylate at

room temperature for 24 h, and then incubated overnight in 4% formaldehyde, 1%

glutaraldehyde, and 0.1% PBS. For scanning microscopy, samples were then

dehydrated through a graded series of ethanol solutions before critical-point drying using

liquid carbon dioxide in a Toumisis Samdri 795 device and sputter-coating with gold-

palladium in a Denton Vacuum Desk-2 device. Samples were examined in a Zeiss Supra

Field Emission Scanning Electron Microscope (Carl Zeiss Microscopy, LLC, North

America), using an accelerating voltage of 5 KV.

For Cryo-EM, grids were prepared following standard procedures and observed

at liquid nitrogen temperatures in a JEM-2200FS/CR transmission electron microscope

(JEOL Europe, Croissy-sur-Seine, France) operated at 200 kV. An in-column omega

energy filter helped to record images with improved signal/noise ratio by zero-loss

filtering. The energy selecting slit width was set at 9 eV. Digital images were recorded on

an UltraScan4000 CCD camera under low-dose conditions at a magnification of 55,058

obtaining a final pixel size of 2.7 Å/pixel.

Density profiles were calculated along rectangular selections with the ImageJ

software (NIH, Bethesda, MD, United States). The number of cells analyzed varied from

n = 20 (WT), n = 15 ($virR^{mut}$) and n = 23 ($virR^{mut}$-C). The comparisons extracted from this

data is statistically relevant with t test with Welch's correction performed. For each cell

three representative measurements were obtained. Mean and standard error was then

calculated for measurements pooled for each strain.

Western Dot blot. Two μl of EV isolates and two-fold serial dilutions were spotted onto

a nitrocellulose membrane (Abcam, Cambridge, United Kingdom) and process for dot

blot using anti-EV polyclonal murine serum (Prados-Rosales et al., 2014a) as primary

antibody and goat anti-mouse-HRP conjugated as secondary antibody and ECL prime

Western Blotting Detection chemiluminescent substrate (GE Healthcare, Chicago, IL,

United States). The signal was visualized in a Chemidoc MP imaging system (Bio Rad)s).

RNA sequencing. *Mtb* strains were grown in MM supplemented with 0.05% Tyloxapol

at 37ºC until they reached an OD595nm of 0.3 and harvested by centrifugation. The cell

pellets were resuspended in 1 ml Qiagen RNA protect reagent (Qiagen, Venlo,

Netherlands) and incubated for 24 h at room temperature. Cells were disrupted by

mechanical lysis in a FastPrep-24 instrument (MP Biomedicals, Santa Ana, CA, United

States) in Lysing Matrix B tubes and RNA was purified with the Direct-zol RNA miniprep

kit (Zymo Research, Irvine, CA, United States). The quantity and quality of the RNAs

were evaluated using Qubit RNA HS Assay Kit (Thermo Fisher Scientific) and Agilent

RNA 6000 Nano Chips (Agilent Technologies, Santa Clara, CA, United States),

respectively. Sequencing libraries were prepared using the Ribo-Zero rRNA Removal Kit

(Gram-positive Bacteria) (Illumina Inc., San Diego, CA, United Stated) and the

TruSeqStranded mRNA library prep kit (Illumina Inc), following the Ribo-Zero rRNA

Removal kit Reference guide and the "TruSeqStranded mRNA Sample Preparation

Guide. Briefly, starting from 1 µg of total RNA, bacterial rRNA was removed and the

remaining RNA was cleaned up using AgencourtRNAClean XP beads (Beckman

Coulter). Purified RNA was fragmented and primed for cDNA synthesis. cDNA first strand

was synthesized with SuperScript-II Reverse Transcriptase (Thermo Fisher Scientific)

for 10 min at $25°C$, 15 min at $42°C$, 15 min at $70°C$ and pause at $4°C$. cDNA second strand

was synthesized with Illumina reagents at $16°C$ for 1 hr. Then, A-tailing and adaptor

ligation were performed. Libraries enrichment was achieved by PCR (30 sec at $98°C$; 15

cycles of 10 sec at $98°C$, 30 sec at $60°C$, 30 sec at $72°C$; 5 min at $72°C$ and pause at $4°C$).

Afterwards, libraries were visualized on an Agilent 2100 Bioanalyzer using Agilent High

Sensitivity DNA kit (Agilent Technologies) and quantified using Qubit dsDNA HS DNA

Kit (Thermo Fisher Scientific). Library sequencing was carried out on an Illumina

HiSeq2500 sequencer with 50 nucleotides single end reads and at least 20 million reads

per individual library were obtained.

RNA-sequencing data analysis. Quality Control of sequenced samples was performed

by FASTQC software (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/).

Reads were mapped against the *M. tuberculosis* H37Rv strain

(GCF_000195955.2_ASM19595v2) reference genome using Tophat (Trapnell et al.,

2009) with --bowtie1 option, to align 50 bp reads. The resulting BAM alignment files for

the samples were the input to Rsubread's (Liao et al., 2019) featureCounts function to

generate a table of raw counts required for the Differential Expression (DE) analysis.

Data preprocessing and differential expression analysis of the transcriptomic data was

performed using DeSeq2 (Love et al., 2014). Genes with a median of expression lower

than 1 CPM, and genes containing outlier observations were filtered before statistical

modeling, and the remaining N=4004 genes were used for differential expression

analysis. P-values were corrected for multiple testing using the R package, implementing

the Storey-Tibshirani FDR correction method (Storey et al., 2023). Genes with an

FDR<0.05 were deemed as differentially expressed. Data were deposited at Gene

expression Omnibus (GEO); GSE143996.

Label-free mass spectrometry analysis. Total cell proteins and MEV-associated proteins were submitted to LC-MS label-free analysis using a Synapt G2Si ESI Q-Mobility-TOF spectrometer (Waters) coupled online to a NanoAcquity nano-HPLC (Waters), equipped with a Waters BEH C18 nano-column (200mm x 75 um ID, 1.8um). Samples were incubated and digested following the filter-aided sample preparation (FASP) protocol (Wiśniewski et al., 2009) Trypsin was added to a trypsin:protein ratio of 1:50, and the mixture was incubated overnight at 37°C ,dried out in a RVC2 25 speedvac concentrator (Christ), and resuspended in 0.1% FA. Peptides were desalted and resuspended in 0.1% FA using C18 stage tips (Millipore). Digested samples (500 ng) were loaded onto the LC system and resolved using a 60 min gradient (5 to 60% ACN). Data was acquired in HDDA mode that enhances signal intensities using the ion mobility separation step. Protein identification and quantification were carried out using Progenesis LC-MS (version 2.0.5556.29015, Nonlinear Dynamics). One of the runs was used as the reference to which the precursor masses in all other samples were aligned to. Only features comprising charges of 2+ and 3+ were selected. The raw abundances of each feature were automatically normalized and logarithmized against the reference run. Samples were grouped in accordance with the comparison being performed, and an ANOVA analysis was performed. A peak list containing the information of all the features was generated and exported to the Mascot search engine (Matrix Science Ltd.). This file was searched against a Uniprot/Swissprot database, and the list of identified peptides was imported back to Progenesis LC-MS. Protein quantitation was performed based on the three most intense non-conflicting peptides (peptides occurring in only one protein), except for proteins with only two non-conflicting peptides.

Proteomics data analysis. Normalization and imputation of the proteomics data was done using the DEP package for R (Zhang et al., 2018). Normalization was done using the vsn method (Huber et al., 2002), which transforms data tackling heteroskedasticity. Imputation of missing values was performed using the bpca method (Oba et al., 2003), which leans on principal component analysis to impute missing values.

Data was analyzed according to a design where differences between strains ($virR^{mut}$ vs -WT, as well as $virR^{mut}$ vs $virR^{mut}$-Comp) were estimated independently within each cell compartment, (whole cell lysates or EVs). according to a nested design: Expression ~ Compartment + Strain:Compartment.

Differential expression analysis of the proteomics data was performed using limma (Ritchie et al., 2015) and p-values were corrected for multiple testing according to the

1012   Storey-Tibshirani method using the R package qvalue (Storey et al., 2003). Proteins with
1013   an FDR<0.05 were selected as differentially expressed.

1014

Gene ontology (GO) enrichments. Enrichment analyses of the sets of differentially
expressed genes and proteins were performed in R using the terms of the Gene
Ontology database for testing, version 11.02.2020. Gene sets corresponding to the
biological process and cellular component ontologies between levels 4 to 6 (N=2245)
were selected and filtered according to their size (minimum size =10 in the RNAseq
analyses; and =3 in the proteomics analyses). The selected sets were tested for
enrichment using Fisher's exact test among genes, or proteins consistently up, or down
regulated in $virR^{mut}$ with respect to both $virR$-expressing strains in each case. Multiple
testing correction was performed using the qvalue R package (Storey et al., 2023).
Terms with an FDR < 0.1 and an odds ratio > 2 were selected for visualization in figures
3E, and 6F, where node sizes was proportional to the significance of the enrichments. In
these visualizations, two terms were connected depending on the amount of sharing
between the groups of genes contributing to the enrichments in each of them, drawing a
link whenever the intersection between the groups of genes was larger than 50% of the
smallest of the two gene sets involved. Ontology term clusters, represented by colors,
were then assigned using the Louvain's algorithm for modularity optimization (Blondel et
al., 2008), as implemented in the R package igraph (Csardi and Nepusz, 2006).
Visualization of the resulting networks was done using the open-source software Gephi
(Bastian et al., 2009).

1034

Targeted metabolomics. LC-MS analysis was carried out on a Schimazdu LC/MS-8030
coupled with a triple quadrupole mass analyser (QqQ) provided with an electrospray
source in a negative ionization mode. Extracted metabolites were quantified in Multiple
Reaction Monitoring (MRM) mode. A stock standard solution prepared in 17% (v/v)
methanol in water and containing all metabolites: acetyl-CoA, methyl citrate,
methylmalonyl-CoA, pyruvate, was used as external standard. Metabolites were
identified by ion pairing liquid chromatography. Acetyl-CoA, methyl citrate and
methylmalonyl-CoA were separated by reverse phase with a poroshell 120 Phenyl-Hexyl
analytical column (2.1 x 50mm, 2.7 μm, Agilent) and a binary gradient (**Table S2**)
consisting of a mobile phase A with water-methanol 97:3 (v/v), 10 mM tributylamine and
3 mM acetic acid, and mobile phase B composed of pure methanol. Pyruvate was
separated by normal phase using a Kinetex HILIC analytical column (2.1 x 150 mm, 2.6
μm, Phenomenex) and a binary gradient (**Table S3**) consisting of pure water as mobile
phase A and pure methanol as mobile phase B.

The flow rate was maintained at 0.4 ml/min and the injection volume was 20 μl.

Nitrogen was used as nebulizing and drying gas at a flow rate of 1.5 and 15 ml/min,

respectively. The desolvation line (DL) temperature was set at 250°C and the ionization

voltage was fixed at 4.5 kV. Argon was used as collision gas to perform collision-induced

dissociation at a pressure of 230 kPa. The dwell time was set at 100 ms and the detection

was performed in MRM mode (detector voltage of 1.8 Kv), monitoring three transitions

for each compound. The transition with higher intensity was selected as quantitative

purposes (**Table S4**), while the other two were used for confirmation of the identity.

1057

Lipidomic analysis by UPLC-MS. The effect of virR on relative abundance of different

cell wall lipids was characterized as had been done previously for MEVs (Prados-

Rosales et al., 2014b). Briefly, cell wall lipids were extracted from triplicate 10 ml cultures

of wild type, $virR^{mut}$, and $virR^{mut}$ complemented ($virR^{mut}$-C) strains overnight into 3 ml 2:1

chloroform:methanol at room temperature, extracts were dried under nitrogen before

being dissolved in 0.75 ml LC-MS grade 2:1:1 isopropanol:acetonitrile:water. Extracts

were separated on a Waters ACQUITY BEH C18 column UPLC heated to 55ºC and

eluted at a flow rate of 0.4 ml/minute with 60:40 acetonitrile:water as mobile phase A and

90:10 isopropanol:acetonitrile as mobile phase B, both containing 10 mM ammonium

formate and 0.1% formic acid. Initial condition was 40%, B which increased to 43% at 2

minutes, 50% at 2.1 minutes, 54% at 12 minutes, 70% at 12.1 minutes, 90% at 18

minutes before returning to initial condition at 18.1 minutes to complete the run at 20

minutes. The coupled Waters Synapt G2 q-TOF MS was operated in positive ion

resolution mode with the following conditions: capillary voltage 2kV, cone voltage 30V,

desolvation gas temperature 550°C , gas flow 900L/hour, source temperature 120°C.

Mass spectra were acquired in centroid mode from 100-3000 m/z with scan times of 0.5

second. Leucine enkephaline was used as reference. Relative abundances of Mass-

retention time pairs were normalized to total ion current and lipid species were identified

using Mycomass (Layre et al., 2011) and Mtb LipidDB databases (Sartain et al., 2011).

1077

Lipidomic analysis by thin layer chromatography (TLC). Lipidomic analysis of both

whole cells and EVs by TLC was performed as previously described (Boldrin et al.,

2021). *Mtb* strains were grown at 37°C in standing Middlebrook 7H9 and when they

reached an OD595 of 0.2, cells were washed and transferred to MM. Cells were grown

until they reached an OD595 of 0.5. Further, cells were harvested by centrifugation at

3,500 rpm for 5 min, and pellets were heat-inactivated at 100°C for 1 h and used for lipid

extraction. Five milliliters 2:1 (vol/vol) chloroform-methanol was added to the cell pellet

and incubated at room temperature (RT) for 12 h with constant stirring. The organic

extract was separated by centrifugation (1,000 × g, 5 min, RT) and decantation. The obtained pellet was extracted with 5 ml 2:1 (vol/vol) chloroform-methanol for 1 h at RT with constant stirring and separated under the same conditions. The combined two organic extracts (total lipids) were dried under a stream of nitrogen gas at RT and saved for lipid analysis.

Polar lipids were separated by 2D-TLC by spotting 200 μg of total lipids on 60 F254 silica gel plates (Merck) using chloroform, methanol, water (50:30:6) as a mobile phase 1 (first dimension), and chloroform, acetic acid, methanol, and water (40:25:3:6) as mobile phase 2 (second dimension). Analysis of PDIMs and TAGs was performed in a 1D-TLC format. Two hundred μg of total lipids were spotted on the silica plates and separated using 9:1 petroleum ether-diethyl ether as mobile phase. Free mycolates were separated by 1D-TLC using chloroform/methanol/ammonium hydroxide (80:20:2) as mobile phase. Mycolic acids were extracted and separated from cells as previously described (Vilchèze and Jacobs, 2007). Sulfolipids and diacyltrehaloses were separated using 60:12:1 chloroform/methanol/water as mobile phase. EV-associated lipids were separated and visualized following the same procedures as for WCLs. Development of TLC plates was performed by repeating the process of pulverizing the staining solution and heating the plate at 100°C 4 times. The staining solution contained 10.5 ml 15% ethanolic solution of 1-naphthol, 6.5 ml 97% sulfuric acid, 40.5 ml ethanol, and 4 ml water.

PG isolation. To obtain mycobacterial cell walls, whole cell pellets were resuspended in PBS and boiled for 15 min. Boiled cells were transferred to 2 ml Lysing matrix tubes with 0.1 mm silica beads (M.P. Biomedical, Santa Ana, CA, United States). And mechanically disrupted in a FastPrep-24TM (M.P. Biomedical) giving 12 cycles at predetermined speed of 6 during 30 sec. Samples were allowed to chill on ice for 3 min between cycles. Cell breakage was monitored by optical microscopy (using Methylene blue as stain), if 95% of the cells had not broken perform additional FastPrep-24TM cycles were performed. Lysed cells were centrifuged at 170 × g for 30 sec to separate them from beads and the supernatant were transferred to a clean 1.5 ml tube. The suspensions were further digested with 10 µg of DNase and RNase/ml for 1 h at 4°C to obtain a cell wall-enriched fraction after centrifugation at 27,000 × g for 10 min. Cell walls were resuspended in PBS containing 2% sodium dodecyl sulfate (SDS) and the suspension was incubated for 1 h at 90°C with constant stirring and recentrifuged at 27,000 × g for 30 min, and the supernatant was discarded. This process was repeated twice. The resulting pellet was washed trice in distilled water, 80% acetone and acetone and lyophilized. Pellets were resuspended in 0.9 ml Tris-HCL (50 mM, pH 7) buffer, plus 0.1

ml of protease K stock solution. The mixture was incubated at 60°C for 90 min, to digest any remaining surface proteins attached to the CW and centrifuged again for 3 min at 20,000 × g and resuspend pellet in distilled water. The mixture was then heated at 90°C for 1 h before centrifugation at 27,000 × g for 30 min. The supernatant was discarded and washed twice with PBS and four times with deionized water to remove SDS. Pure PG was obtained from isolated cell walls. Briefly, pure cell walls were resuspended in 0.5% KOH in methanol and stirred at 37°C for 4 days to hydrolyze mycolic acids (MA). The mixture was centrifuged (27,000 × g 20 min) and the pellet was washed twice with methanol. MA were separated from cell walls using diethyl ether and recentrifuged at 27,000 × gfor 20 min. The supernatant contained MAMEs. The extraction process with diethyl ether was repeated twice and let dry. The resulting arabinogalactan (AG)-PG was digested with 0.2 N H_2SO_4 at 85°C for 30 min and neutralized with $BaCO_3$ to remove the arabinogalactan. Treated AG-PG was centrifuged at 27,000 × g for 20 min. Supernatant contained soluble AG. The resulting insoluble PG was washed four times by centrifugation in deionized water and stored at 4°C or room temperature.

Analysis of diaminopimelic acid. For DAP quantification (Tsuruoka et al., 1985), samples were hydrolyzed overnight in 6N HCl at 100°C in 1 ml capacity-crystal ampoules (Wheaton). Samples were dried and resuspended in water and mixed in a 1:1:1 ratio with pure acetic acid and ninhydrin reagent (250 mg ninhydrin, 4 ml phosphoric acid 0.6 M, and 6 ml pure acetic acid). Samples were measured at OD 434nm, and the amounts of DAP were calculated based on a standard curve with pure DAP. The data displays the average ± standard deviations n=3.

Glycosyl composition of isolated cell walls. 500 µg of purified cell walls were mixed with 10 µg of inositol and hydrolyzed with 0.1 M trifluoroacetic acid (TFA) for 2 h at 121°C. The monosaccharides released were then converted into alditol acetates by reduction with $NaBH_4$ and acetylation with acetic anhydride in pyridine. The products were analyzed by gas chromatography-mass spectrometry in a 7890A/5975C system from Agilent, using a DB-5ht (30 m x 0.25 mm x 0.1 µm) fused silica capillary column. The compounds were identified by comparing the retention times of the peaks with those recorded for standards analyzed under identical conditions. Quantification was carried out considering the area of the peaks and the response factors of each compound with respect to inositol.

Atomic force microscopy and data analysis. The dry pellets of purified peptidoglycan from the three strains were dissolved in pure water and further broken by tip sonication

at 2 mA for 30 sec, three cycles. Then 10 µl of the PG suspension was added to standard
AFM substrate made of mica coated with 0.01 % Poly-L-lysine by incubating the sample
during 90 min to achieve good coverage. The sample was further rinsed with water and
dried with Nitrogen flow. The AFM in ambient conditions (i.e. air) was performed using
Tapping mode in a Bruker FastScan Bio machine (Santa Barbara, CA, United States)
with Nunano SCOUT 350 - Silicon AFM probes (spring constant: 42 N/m, Resonance
frequency: 350 kHz). The images were taken using a free amplitude of 10 nm with set
point of 70-80 % of free amplitude (e.g. 7 nm). For AFM high resolution imaging in liquid
the images were acquired in Peak force Tapping mode in imaging buffer made of 10 mM
Tris and 150 mM KCl, pH: 8, with the FastScan Bio machine using Bruker Fastscan-D
AFM probes (spring constant: 0.1 N/m, Resonance frequency: 70 kHz) at the force range
of 1-3 nN peak force set point. The imaging parameters used are as follows; Scan rate:
1 Hz; Scan angle: 0 °; Peak force amplitude: 80-100 nm and with high pixel resolution
(512-880).

The thickness measurements were performed with standard Gwyddion 1D height
distribution tool (selecting an area containing background and a single leaflet of
peptidoglycan). The number of cells analysed varied from n = 23 (for Mtb WT), n = 36
(for $virR^{mut}$) and n = 25 (for $virR^{mut}$-C). The comparisons extracted from this data is
statistically relevant with t test with Welch's correction performed. The analysis of the
high-resolution images obtained in liquid was performed using the semi-automated
custom-made routine from (Pasquina-Lemonche, et al. "Pore AFM", 2024) where the
pore area, later converted in diameter, and the pore number were measured from the 2D
slice of each image containing the maximum number of pores (see **Fig. S8**). The number
of images similar to images shown in **Fig. 5** were n = 8 for all strains, analyzing both pore
size and pore number, the statistical comparisons were also performed using a t test with
Welch's correction.

1186

Blue native PAGE. Oligomerization of VirR and Lcp1 was analyzed by Blue native
PAGE (Invitrogen, Waltham, MA, United States). Recombinant VirR and Lcp1 proteins
were prepared in 1xLB Native PAGE including DDM at 0.5% final concentration. After
incubation at 4ºC for 30 min, samples were centrifuged at 20,000 × g for 30 min at 4°C.
The supernatant was recovered and mixed with G-250 additive following manufacturer
instructions (Invitrogen) and loaded onto a Novex® Bist-Tris 4-16% gradient gel and run
following manufacturer instructions. Gel was first fixed and then stained with Coomassie.

1194

Liposome flotation assay. Individual or combined VirR and Lcp1 were prepared at 2
µM in buffer A in a final volume of 50 µl and incubated for 30 min at 37°C. This volume

was mixed with 100 μl of 60% optiprep (Sigma-Aldrich) to a final concentration of 40%
and overlaid with 100 μl of subsequent optiprep solutions at 35%, 30%, 20% and 10%
(optiprep solutions were prepared in Buffer A). The samples were centrifugated at 70000
rpm for 2 h at 4°C in an Optima TLX Ultracentrifuge (Beckman Coulter, Indianapolis, IN
United States). Eleven fractions of 50 μl were recovered and examined by dot blot
(performed as previously described). Polyclonal antibodies raised against VirR and Lcp1
were used as primary antibodies. Quantification of dotblots was performed in ImageJ as
previously described (Schirmer et al., 2022). Briefly, integrated densities were calculated
after background subtraction, and referred to wild type values to obtain the relative
integrated intensities.

1207

1208 **Lysozyme and PG and AG susceptibility assays.** *Mtb* WT and *virR^{mut}* strains were
1209 grown in 7H9 media supplemented with ADC and Tween 80 to an OD 540 of 0.2 and
1210 then diluted to an OD540 of 0.1 with 7H9 containing 0 or 50 μgml^{-1} lysozyme (Sigma).
1211 The cultures were incubated with agitation at 37°C and OD540 was monitored daily.

1212

Fluorescence microscopy. The *Mtb* strain bearing the plasmid VirR-pMV261-Venus
was grown in MM to an OD595nm of 0.4. At this stage, cells were harvested, washed
three times in 1 × PBS (pH 7.4), resuspended in 1/3 – 1/6 volume of 4%
paraformaldehyde (PFA) and incubated for 2 h at 37°C. Finally, the cells were washed
twice in 1 × PBS and resuspended in 100–200 μl of PBS. Microscope slides were
covered with 5 μl poly-L-lysine (Sigma) and excess poly-L-lysine was removed away with
distilled water. A 20 – 40 μl aliquot was applied to pre-treated slides, allowed to air dry
and covered with antifade solution (Prolong Gold, Invitrogen) and visualized under the
fluorescence microscope using a 100x oil immersion objective on an Orca 12-ERG digital
CCD camera (Hamamatsu Photonics) mounted on a Nikon E600 microscope.

1223

FACS analysis for split-GFP experiments. *M. smegmatis* mc²155 transformed with the
pGMCS derivatives were inoculated in complete 7H9 medium. After overnight incubation
at 37°C, bacteria were collected and analyzed by fluorescence activated cell sorting
using a BD FACS LSRFortessa X20 flow cytometer. Flow cytometry data analysis was
performed using FlowJo software (Version 10; Becton, Dickinson and Company,
Ashland, OR, United States). The gating strategy is displayed in supplementary figure
10 (**Fig. S10**).

1231

1232 **Statistical analysis.** The statistical significance of the difference between experimental
1233 groups was determined by the two-tailed Student's test using PRISM 5.0. P values ≤

than 0.05 were considered significant. Statistical analysis of RNAseq is detailed in the corresponding section of Materials and Methods.

Data Availability. The Datasets produced in this study are available in the following database: RNA-seq data: Gene expression Omnibus, GSE143996 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE143996).

Funding Information. R.P.-R. acknowledges support by MINECO/FEDER EU contracts SAF2016–77433-R, PID2019-110240RB-I00, PID2022-136611OB-I00; and NIH-R01AI162821. CICbioGUNE thanks MINECO for the Severo Ochoa Excellence Accreditation (SEV-2016–0644). L.P.-L and J.K.H acknowledges support by the Wellcome Trust (212197/Z/19/Z) which funded part of this research. For the purpose of open access, the authors have applied and will apply a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. J.S. acknowledges support from the Spanish Ministry of Science and Innovation (MICINN) through grant PID2019-106859GA-I00 and Ramón y Cajal research grant RYC-2017-23560, as well as to the Government of Aragón, Spain, through grant B49-23R (NeuroBioSys). FC lab research is supported by the Swedish Research Council (VR2018-02823 and VR2018-05882), the Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, the Knut and Alice Wallenberg Foundation (KAW) grant KAW2012.0184 and the Kempe Foundation.

Acknowledgments. We are very grateful to Carl Nathan for sharing mutant *Mtb* strains. We thank G. Marcela Rodriguez for sharing the a-IdeR antibody. We thank Heran Darwin for sharing a-cei polyclonal antibody. We thank Emmanuelle Näser (Génotoul TRI-IPBS platform, Toulouse) for helping with flow cytometry. We thank Dr. Luke Alderwick (University of Birmingham) for the donation of the plasmid to recombinantly express Lcp1 protein. We thank M. Gutiérrez for assistance with figures. The authors have no conflict of interest to declare.

References

Amaral L, Kristiansen JE, Abebe LS, Millett W. 1996. Inhibition of the respiration of multi-drug resistant clinical isolates of Mycobacterium tuberculosis by thioridazine: Potential use for initial therapy of freshly diagnosed tuberculosis. *J Antimicrob Chemother* **38**. doi:10.1093/jac/38.6.1049

Ariyachaokun K, Grabowska AD, Gutierrez C, Neyrolles O. 2020. Multi-stress induction of the Mycobacterium tuberculosis MbcTA bactericidal toxin-antitoxin system.

Toxins (Basel) **12**. doi:10.3390/toxins12050329

Athman JJ, Sande OJ, Groft SG, Reba SM, Nagy N, Wearsch PA, Richardson ET, Rojas R, Boom WH, Shukla S, Harding C V. 2017. Mycobacterium tuberculosis Membrane Vesicles Inhibit T Cell Activation. *J Immunol* **198**:2028–2037. doi:10.4049/jimmunol.1601199

Athman JJ, Wang Y, McDonald DJ, Boom WH, Harding C V, Wearsch PA. 2015. Bacterial Membrane Vesicles Mediate the Release of Mycobacterium tuberculosis Lipoglycans and Lipoproteins from Infected Macrophages. *J Immunol* **195**:1044–1053. doi:10.4049/jimmunol.1402894

Ballister ER, Samanovic MI, Darwin KH. 2019. Mycobacterium tuberculosis rv2700 contributes to cell envelope integrity and virulence. *J Bacteriol* **201**. doi:10.1128/JB.00228-19

Bartek IL, Woolhiser LK, Baughn AD, Basaraba RJ, Jacobs WR, Lenaerts AJ, Voskuil MI. 2014. Mycobacterium tuberculosis Lsr2 is a global transcriptional regulator required for adaptation to changing oxygen levels and virulence. *MBio* **5**. doi:10.1128/mBio.01106-14

Bastian M, Heymann S, Jacomy M. 2009. Gephi: An Open Source Software for Exploring and Manipulating Networks. *Proc Int AAAI Conf Web Soc Media* **3**. doi:10.1609/icwsm.v3i1.13937

Baumgart M, Schubert K, Bramkamp M, Frunzke J. 2016. Impact of LytRCpsA-Psr proteins on cell wall biosynthesis in Corynebacterium glutamicum. *J Bacteriol* **198**. doi:10.1128/JB.00406-16

Beaulieu AM, Rath P, Imhof M, Siddall ME, Roberts J, Schnappinger D, Nathan CF. 2010. Genome-wide screen for Mycobacterium tuberculosis genes that regulate host immunity. *PLoS One* **5**:e15120. doi:10.1371/journal.pone.0015120

Bera A, Herbert S, Jakob A, Vollmer W, Götz F. 2005. Why are pathogenic staphylococci so lysozyme resistant? The peptidoglycan O-acetyltransferase OatA is the major determinant for lysozyme resistance of Staphylococcus aureus. *Mol Microbiol* **55**. doi:10.1111/j.1365-2958.2004.04446.x

Blondel VD, Guillaume J-L, Lambiotte R, Lefebre E. 2008. Fast unfolding of community hierarchies in large networks. *J Stat Mech Theory Exp* **10**.

Boldrin F, Anso I, Alebouyeh S, Sevilla IA, Geijo M, Garrido JM, Marina A, Mazzabò LC, Segafreddo G, Guerin ME, Manganelli R, Prados-Rosales R. 2021. The phosphatidyl-myo-inositol dimannoside acyltransferase pata is essential for mycobacterium tuberculosis growth in vitro and in vivo. *J Bacteriol* **203**. doi:10.1128/JB.00439-20

Boot M, van Winden VJC, Sparrius M, van de Weerd R, Speer A, Ummels R, Rustad T,

Sherman DR, Bitter W. 2017. Cell envelope stress in mycobacteria is regulated by

the novel signal transduction ATPase IniR in response to trehalose. *PLoS Genet*

13:e1007131. doi:10.1371/journal.pgen.1007131

Boudehen YM, Faucher M, Maréchal X, Miras R, Rech J, Rombouts Y, Sénèque O,

Wallat M, Demange P, Bouet JY, Saurel O, Catty P, Gutierrez C, Neyrolles O.

2022. Mycobacterial resistance to zinc poisoning requires assembly of P-ATPase-

containing membrane metal efflux platforms. *Nat Commun* **13**.

doi:10.1038/s41467-022-32085-7

Brzostek A, Pawelczyk J, Rumijowska-Galewicz A, Dziadek B, Dziadek J. 2009.

Mycobacterium tuberculosis is able to accumulate and utilize cholesterol. *J*

Bacteriol **191**. doi:10.1128/JB.00488-09

Cabantous S, Nguyen HB, Pedelacq JD, Koraïchi F, Chaudhary A, Ganguly K, Lockard

MA, Favre G, Terwilliger TC, Waldo GS. 2013. A new protein-protein interaction

sensor based on tripartite split-GFP association. *Sci Rep* **3**.

doi:10.1038/srep02854

Chauhan S, Tyagi JS. 2009. Powerful induction of divergent tgs1-Rv3131 genes in

Mycobacterium tuberculosis is mediated by DevR interaction with a high-affinity

site and an adjacent cryptic low-affinity site. *J Bacteriol* **191**.

doi:10.1128/JB.00310-09

Csardi G, Nepusz T. 2006. The igraph software package for complex network

research. *InterJournal Complex Syst* **Complex Sy**.

De Keijzer J, Mulder A, De Haas PEW, De Ru AH, Heerkens EM, Amaral L, Van

Soolingen D, Van Veelen PA. 2016. Thioridazine alters the cell-envelope

permeability of mycobacterium tuberculosis. *J Proteome Res* **15**.

doi:10.1021/acs.jproteome.5b01037

De Knegt GJ, Ten Kate MT, Van Soolingen D, Aarnoutse R, Boeree MJ, Bakker-

Woudenberg IAJM, De Steenwinkel JEM. 2014. Enhancement of in vitro activity of

tuberculosis drugs by addition of thioridazine is not reflected by improved in vivo

therapeutic efficacy. *Tuberculosis* **94**. doi:10.1016/j.tube.2014.09.002

Dutta NK, Mehra S, Kaushal D. 2010. A Mycobacterium tuberculosis sigma factor

network responds to cell-envelope damage by the promising anti-mycobacterial

thioridazine. *PLoS One* **5**. doi:10.1371/journal.pone.0010069

Ghazaei C. 2018. Mycobacterium tuberculosis and lipids: Insights into molecular

mechanisms from persistence to virulence. *J Res Med Sci*.

doi:10.4103/jrms.JRMS_904_17

Grzegorzewicz AE, de Sousa-d'Auria C, McNeil MR, Huc-Claustre E, Jones V, Petit C,

Angala SK, Zemanova J, Wang Q, Belardinelli JM, Gao Q, Ishizaki Y, Mikusova K,

Brennan PJ, Ronning DR, Chami M, Houssin C, Jackson M. 2016. Assembling of

the Mycobacterium tuberculosis Cell Wall Core. *J Biol Chem* **291**:18867–18879.

doi:10.1074/jbc.M116.739227

Gupta S, Bhagavathula, M Sharma V, Sharma N, Sharma N, Biswas A, Palacios A,

Salgueiro V, Lavín J, Dogra N, Salgame P, Prados-Rosales, R Rodríguez G.

2023. Dynamin-like proteins mediate extracellular vesicle secretion in

Mycobacterium tuberculosis. *EMBO Rep* **24**:e55593.

Harrison J, Lloyd G, Joe M, Lowary TL, Reynolds E, Walters-Morgan H, Bhatt A,

Lovering A, Besra GS, Alderwick LJ. 2016. Lcp1 is a phosphotransferase

responsible for ligating arabinogalactan to peptidoglycan in mycobacterium

tuberculosis. *MBio* **7**. doi:10.1128/mBio.00972-16

Hasenoehrl EJ, Rae Sajorda D, Berney-Meyer L, Johnson S, Tufariello JAM, Fuhrer T,

Cook GM, Jacobs WR, Berney M. 2019. Derailing the aspartate pathway of

Mycobacterium tuberculosis to eradicate persistent infection. *Nat Commun* **10**.

doi:10.1038/s41467-019-12224-3

Honaker RW, Leistikow RL, Bartek IL, Voskui MI. 2009. Unique roles of DosT and

DosS in DosR regulon induction and Mycobacterium tuberculosis dormancy. *Infect*

Immun **77**. doi:10.1128/IAI.01449-08

Huber W, Heydebreck A Von, Ultmann HS¨, Poustka A, Vingron M. 2002. Variance

stabilization applied to microarray data calibration and to the quantification of

differential expression. *BIOINFORMATICS*.

Hubscher J, Luthy L, Berger-Bachi B, Stutzmann Meier P. 2008. Phylogenetic

distribution and membrane topology of the LytR-CpsA-Psr protein family. *BMC*

Genomics **9**:617. doi:10.1186/1471-2164-9-617

Justen AM, Hodges HL, Kim LM, Sadecki PW, Porfirio S, Ultee E, Black I, Chung GS,

Briegel A, Azadi P, Kiessling LL. 2020. Polysaccharide length affects

mycobacterial cell shape and antibiotic susceptibility. *Sci Adv* **6**.

doi:10.1126/sciadv.aba4015

Kalscheuer R, Palacios A, Anso I, Cifuente J, Anguita J, Jacobs Jr. WR, Guerin ME,

Prados-Rosales R. 2019. The Mycobacterium tuberculosis capsule: a cell

structure with key implications in pathogenesis. *Biochem J* **476**:1995–2016.

doi:10.1042/BCJ20190324

Kawai Y, Marles-Wright J, Cleverley RM, Emmins R, Ishikawa S, Kuwano M, Heinz N,

Bui NK, Hoyland CN, Ogasawara N, Lewis RJ, Vollmer W, Daniel RA, Errington J.

2011. A widespread family of bacterial cell wall assembly proteins. *EMBO J*

30:4931–4941. doi:10.1038/emboj.2011.358

Kuru E, Radkov A, Meng X, Egan A, Alvarez L, Dowson A, Booher G, Breukink E,

Roper DI, Cava F, Vollmer W, Brun Y, Vannieuwenhze MS. 2019. Mechanisms of

1383   Incorporation for D -Amino Acid Probes That Target Peptidoglycan Biosynthesis.

1384   *ACS Chem Biol* **14**. doi:10.1021/acschembio.9b00664

1385   Layre E, Sweet L, Hong S, Madigan CA, Desjardins D, Young DC, Cheng TY, Annand

1386   JW, Kim K, Shamputa IC, McConnell MJ, Debono CA, Behar SM, Minnaard AJ,

1387   Murray M, Barry CE, Matsunaga I, Moody DB. 2011. A comparative lipidomics

1388   platform for chemotaxonomic analysis of mycobacterium tuberculosis. *Chem Biol*

1389   **18**. doi:10.1016/j.chembiol.2011.10.013

1390   Lee JE, Moon PG, Lee IK, Baek MC. 2015. Proteomic Analysis of Extracellular

1391   Vesicles Released by Adipocytes of Otsuka Long-Evans Tokushima Fatty

1392   (OLETF) Rats. *Protein J* **34**:220–235. doi:10.1007/s10930-015-9616-z

1393   Lee W, VanderVen BC, Fahey RJ, Russell DG. 2013. Intracellular Mycobacterium

1394   tuberculosis exploits host-derived fatty acids to limit metabolic stress. *J Biol Chem*

1395   **288**:6788–6800. doi:10.1074/jbc.M112.445056

1396   Li C, Wen A, Shen B, Lu J, Huang Y, Chang Y. 2011. FastCloning: A highly simplified,

1397   purification-free, sequence- and ligation-independent PCR cloning method. *BMC*

1398   *Biotechnol* **11**. doi:10.1186/1472-6750-11-92

1399   Liao Y, Smyth GK, Shi W. 2019. The R package Rsubread is easier, faster, cheaper

1400   and better for alignment and quantification of RNA sequencing reads. *Nucleic*

1401   *Acids Res* **47**. doi:10.1093/nar/gkz114

1402   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

1403   dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**. doi:10.1186/s13059-

1404   014-0550-8

1405   Lu R, Schaefer CM, Nesbitt NM, Kuper J, Kisker C, Sampson NS. 2017. Catabolism of

1406   the Cholesterol Side Chain in Mycobacterium tuberculosis Is Controlled by a

1407   Redox-Sensitive Thiol Switch. *ACS Infect Dis* **3**. doi:10.1021/acsinfecdis.7b00072

1408   Maciąg A, Dainese E, Rodriguez GM, Milano A, Provvedi R, Pasca MR, Smith I, Palù

1409   G, Riccardi G, Manganelli R. 2007. Global analysis of the Mycobacterium

1410   tuberculosis Zur (FurB) regulon. *J Bacteriol* **189**. doi:10.1128/JB.01190-06

1411   Majlessi L, Prados-Rosales R, Casadevall A, Brosch R. 2015. Release of

1412   mycobacterial antigens. *Immunol Rev* **264**:25–45. doi:10.1111/imr.12251

1413   Marsollier L, Brodin P, Jackson M, Kordulakova J, Tafelmeyer P, Carbonnelle E, Aubry

1414   J, Milon G, Legras P, Andre JP, Leroy C, Cottin J, Guillou MLJ, Reysset G, Cole

1415   ST, Korduláková J, Tafelmeyer P, Carbonnelle E, Aubry J, Milon G, Legras P,

1416   Saint André JP, Leroy C, Cottin J, Guillou MLJ, Reysset G, Cole ST. 2007. Impact

1417   of Mycobacterium ulcerans biofilm on transmissibility to ecological niches and

1418   Buruli ulcer pathogenesis. *PLoS Pathog* **3**:e62. doi:10.1371/journal.ppat.0030062

McMillan HM, Kuehn MJ. 2021. The extracellular vesicle generation paradox: a
    bacterial point of view. *EMBO J* **40**. doi:10.15252/embj.2021108174

Minch KJ, Rustad TR, Peterson EJR, Winkler J, Reiss DJ, Ma S, Hickey M, Brabant W,
    Morrison B, Turkarslan S, Mawhinney C, Galagan JE, Price ND, Baliga NS,
    Sherman DR. 2015. The DNA-binding network of Mycobacterium tuberculosis. *Nat
    Commun* **6**. doi:10.1038/ncomms6829

Mohammadzadeh R, Ghazvini K, Farsiani H, Soleimanpour S. 2021. Mycobacterium
    tuberculosis extracellular vesicles: exploitation for vaccine technology and
    diagnostic methods. *Crit Rev Microbiol*. doi:10.1080/1040841X.2020.1830749

Nagakubo T, Tahara YO, Miyata M, Nomura N, Toyofuku M. 2021. Mycolic acid-
    containing bacteria trigger distinct types of membrane vesicles through different
    routes. *iScience* **24**. doi:10.1016/j.isci.2020.102015

Nagamani S, Sastry GN. 2021. Mycobacterium tuberculosis cell wall permeability
    model generation using chemoinformatics and machine learning approaches. *ACS
    Omega* **6**. doi:10.1021/acsomega.1c01865

Nečas D, Klapetek P. 2012. Gwyddion: An open-source software for SPM data
    analysis. *Cent Eur J Phys*. doi:10.2478/s11534-011-0096-2

Oba S, Sato MA, Takemasa I, Monden M, Matsubara KI, Ishii S. 2003. A Bayesian
    missing value estimation method for gene expression profile data. *Bioinformatics*
    19:2088–2096. doi:10.1093/bioinformatics/btg287

Pandey R, Rodriguez GM. 2014. IdeR is required for iron homeostasis and virulence in
    Mycobacterium tuberculosis. *Mol Microbiol* **91**. doi:10.1111/mmi.12441

Park HD, Guinn KM, Harrell MI, Liao R, Voskuil MI, Tompa M, Schoolnik GK, Sherman
    DR. 2003. Rv3133c/dosR is a transcription factor that mediates the hypoxic
    response of Mycobacterium tuberculosis. *Mol Microbiol* **48**. doi:10.1046/j.1365-
    2958.2003.03474.x

Pasquina-Lemonche L, Burns J, Turner RD, Kumar S, Tank R, Mullin N, Wilson JS,
    Chakrabarti B, Bullough PA, Foster SJ, Hobbs JK. 2020. The architecture of the
    Gram-positive bacterial cell wall. *Nature* **582**. doi:10.1038/s41586-020-2236-6

Pawełczyk J, Brzostek A, Minias A, Płociński P, Rumijowska-Galewicz A, Strapagiel D,
    Zakrzewska-Czerwińska J, Dziadek J. 2021. Cholesterol-dependent transcriptome
    remodeling reveals new insight into the contribution of cholesterol to
    Mycobacterium tuberculosis pathogenesis. *Sci Rep* **11**. doi:10.1038/s41598-021-
    91812-0

Peterson EJ, Bailo R, Rothchild AC, Arrieta-Ortiz ML, Kaur A, Pan M, Mai D, Abidi AA,
    Cooper C, Aderem A, Bhatt A, Baliga NS. 2019. Path-seq identifies an essential
    mycolate remodeling program for mycobacterial host adaptation. *Mol Syst Biol* **15**.

doi:10.15252/msb.20188584

Peterson EJR, Brooks AN, Reiss DJ, Kaur A, Do J, Pan M, Wu WJ, Morrison R,

Srinivas V, Carter W, Arrieta-Ortiz ML, Ruiz RA, Bhatt A, Baliga NS. 2023. MtrA

modulates Mycobacterium tuberculosis cell division in host microenvironments to

mediate intrinsic resistance and drug tolerance. *Cell Rep* **42**.

doi:10.1016/j.celrep.2023.112875

Prados-Rosales R, Baena A, Martinez LR, Luque-Garcia J, Kalscheuer R,

Veeraraghavan U, Camara C, Nosanchuk JD, Besra GS, Chen B, Jimenez J,

Glatman-Freedman A, Jacobs Jr. WR, Porcelli SA, Casadevall A, Jacobs WR,

Porcelli SA, Casadevall A. 2011. Mycobacteria release active membrane vesicles

that modulate immune responses in a TLR2-dependent manner in mice. *J Clin*

Invest **121**:1471–1483. doi:10.1172/JCI44261

Prados-Rosales R, Carreno L, Cheng T, Blanc C, Weinrick B, Malek A, Lowary TL,

Baena A, Joe M, Bai Y, Kalscheuer R, Batista-Gonzalez A, Saavedra NA,

Sampedro L, Tomas J, Anguita J, Hung SC, Tripathi A, Xu J, Glatman-Freedman

A, Jacobs Jr. WR, Chan J, Porcelli SA, Achkar JM, Casadevall A. 2017.

Enhanced control of Mycobacterium tuberculosis extrapulmonary dissemination in

mice by an arabinomannan-protein conjugate vaccine. *PLoS Pathog*

13:e1006250. doi:10.1371/journal.ppat.1006250

Prados-Rosales R, Carreño LJ, Batista-Gonzalez A, Baena A, Venkataswamy MM, Xu

J, Yu X, Wallstrom G, Mitchell Magee D, LaBaer J, Achkar JM, Jacobs WR, Chan

J, Porcelli SA, Casadevall A. 2014a. Mycobacterial membrane vesicles

administered systemically in mice induce a protective immune response to surface

compartments of mycobacterium tuberculosis. *MBio* **5**:e01921-14.

doi:10.1128/mBio.01921-14

Prados-Rosales R, Weinrick BC, Pique DG, Jacobs Jr. WR, Casadevall A, Rodriguez

GM, Piqué DG, Jacobs WR, Casadevall A, Rodriguez GM. 2014b. Role for

Mycobacterium tuberculosis membrane vesicles in iron acquisition. *J Bacteriol*

196:1250–1256. doi:10.1128/JB.01090-13

Prasad TSK, Verma R, Kumar S, Nirujogi RS, Sathe GJ, Madugundu AK, Sharma J,

Puttamallesh VN, Ganjiwale A, Myneedu VP, Chatterjee A, Pandey A, Harsha HC,

Narayana J. 2013. Proteomic analysis of purified protein derivative of

mycobacterium tuberculosis. *Clin Proteomics* **10**. doi:10.1186/1559-0275-10-8

Rath P, Huang C, Wang T, Wang T, Li H, Prados-Rosales R, Elemento O, Casadevall

A, Nathan CF. 2013. Genetic regulation of vesiculogenesis and

immunomodulation in Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A*

110:E4790-7. doi:10.1073/pnas.1320118110

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. Limma powers
differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res **43**:e47. doi:10.1093/nar/gkv007

Rock JM, Hopkins FF, Chavez A, Diallo M, Chase MR, Gerrick ER, Pritchard JR,
Church GM, Rubin EJ, Sassetti CM, Schnappinger D, Fortune SM. 2017.
Programmable transcriptional repression in mycobacteria using an orthogonal
CRISPR interference platform. *Nat Microbiol* **2**:16274.
doi:10.1038/nmicrobiol.2016.274

Sani M, Houben ENG, Geurtsen J, Pierson J, De Punder K, Van Zon M, Wever B,
Piersma SR, Jiménez CR, Daffé M, Appelmelk BJ, Bitter W, Van Wel N Der,
Peters PJ. 2010. Direct visualization by Cryo-EM of the mycobacterial capsular
layer: A labile structure containing ESX-1-secreted proteins. *PLoS Pathog* **6**.
doi:10.1371/journal.ppat.1000794

Sartain MJ, Dick DL, Rithner CD, Crick DC, Belisle JT. 2011. Lipidomic analyses of
Mycobacterium tuberculosis based on accurate mass measurements and the
novel "Mtb LipidDB." *J Lipid Res* **52**. doi:10.1194/jlr.M010363

Schirmer S, Rauh L, Alebouyeh S, Delgado-Velandia M, Salgueiro VC, Lerma L,
Serrano-Mestre JL, Azkargorta M, Elortza F, Lavín JL, García MJ, Tórtola-
Fernández MT, Gola S, Prados-Rosales R. 2022. Immunogenicity of
Mycobacterial Extracellular Vesicles Isolated From Host-Related Conditions
Informs About Tuberculosis Disease Status. *Front Microbiol* **13**:907296.
doi:10.3389/fmicb.2022.907296

Schnappinger D, Ehrt S, Voskuil MI, Liu Y, Mangan JA, Monahan IM, Dolganov G,
Efron B, Butcher PD, Nathan C, Schoolnik GK. 2003. Transcriptional adaptation of
Mycobacterium tuberculosis within macrophages: Insights into the phagosomal
environment. *J Exp Med* **198**:693–704. doi:10.1084/jem.20030846

Soto-Ramirez MD, Aguilar-Ayala DA, Garcia-Morales L, Rodriguez-Peredo SM,
Badillo-Lopez C, Rios-Muñiz DE, Meza-Segura MA, Rivera-Morales GY, Leon-
Solis L, Cerna-Cortes JF, Rivera-Gutierrez S, Helguera-Repetto AC, Gonzalez-y-
Merchand JA. 2017. Cholesterol plays a larger role during Mycobacterium
tuberculosis in vitro dormancy and reactivation than previously suspected.
Tuberculosis **103**. doi:10.1016/j.tube.2016.12.004

Storey JD, Bass AJ, Dabney A, Robinson D. 2023. qvalue: Q-value estimation for false
discovery rate control.

Stupar M, Tan L, Kerr ED, De Voss CJ, Forde BM, Schulz BL, West NP. 2024. TcrXY
is an acid-sensing two-component transcriptional regulator of Mycobacterium
tuberculosis required for persistent infection. *Nat Commun* **15**.

doi:10.1038/s41467-024-45343-7

Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: Discovering splice junctions with

RNA-Seq. *Bioinformatics* **25**. doi:10.1093/bioinformatics/btp120

Tsuruoka T, Tamura A, Miyata A, Takei T, Inouye S, Matsuhashi M. 1985. Second

Lytic Target of Beta-Lactam Compounds That Have A Terminal d-Amino Acid

Residue. *Eur J Biochem* **151**. doi:10.1111/j.1432-1033.1985.tb09090.x

Vilchèze C, Jacobs WR. 2007. Isolation and analysis of Mycobacterium tuberculosis

mycolic acids. *Curr Protoc Microbiol* **Chapter 10**.

doi:10.1002/9780471729259.mc10a03s05

Wang M, Guo X, Yang X, Zhang B, Ren J, Liu A, Ran Y, Yan B, Chen F, Guddat LW,

Hu J, Li J, Rao Z. 2019. Mycobacterial dynamin-like protein IniA mediates

membrane fission. *Nat Commun* **10**. doi:10.1038/s41467-019-11860-z

White DW, Elliott SR, Odean E, Bemis LT, Tischler AD. 2018. Mycobacterium

tuberculosis Pst/SenX3-RegX3 Regulates Membrane Vesicle Production

Independently of ESX-5 Activity. *MBio* **9**. doi:10.1128/mBio.00778-18

Williams EP, Lee JH, Bishai WR, Colantuoni C, Karakousis PC. 2007. Mycobacterium

tuberculosis SigF regulates genes encoding cell wall-associated proteins and

directly regulates the transcriptional regulatory gene phoY1. *J Bacteriol* **189**.

doi:10.1128/JB.00201-07

Wiśniewski JR, Zougman A, Nagaraj N, Mann M. 2009. Universal sample preparation

method for proteome analysis. *Nat Methods* **6**. doi:10.1038/nmeth.1322

Zhang X, Smits AH, Tilburg GBA Van, Ovaa H, Huber W, Vermeulen M. 2018.

Proteome-wide identification of ubiquitin interactions using UbIA-MS. *Nat Protoc*

13:530–550. doi:10.1038/nprot.2017.147

Ziegenbalg A, Prados-Rosales R, Jenny-Avital ER, Kim RS, Casadevall A, Achkar JM.

2013. Immunogenicity of mycobacterial vesicles in humans: identification of a new

tuberculosis antibody biomarker. *Tuberc* **93**:448–455.

doi:10.1016/j.tube.2013.03.001

Zuber B, Haenni M, Ribeiro T, Minnig K, Lopes F, Moreillon P, Dubochet J. 2006.

Granular layer in the periplasmic space of gram-positive bacteria and fine

structures of Enterococcus gallinarum and Streptococcus gordonii septa revealed

by cryo-electron microscopy of vitreous sections. *J Bacteriol* **188**.

doi:10.1128/JB.00391-06

1563

1564

1565

1566

Figure Legends

Figure 1. Ultrastructural changes in the cell envelope of *virR*-deficient mutant associated with increased vesiculation. (**A**) Cryo-electron micrographs of indicated *Mtb* strains grown in high iron MM. Closed line rectangles were used to calculate grey value profiles of cell envelopes using ImageJ. The dashed line insets within the main micrograph were magnified to show a detailed view of the cell surface. Scale bars are 100 nm in main micrographs and 50 nm in the insets. (**B**) Density profiles based on grey values of the cross sections marked by solid line rectangles in A. (**C**) Mean values and standard errors of distances in nm between main cell envelope layers measured in A and C. *P < 0.05 after applying a Tukey´s multiple comparison test. The number of cells analyzed varied from n = 20 (WT), n = 15 (*virR^{mut}*) and n = 23 (*virR^{mut}*-C). CM, cytoplasmic membrane; OM, outer membrane; L1, layer 1; L2, layer 2.

Figure 2. Enhanced permeability and vesiculation are linked in the absence of LytR_C solo domain proteins in *Mtb*. (A) Uptake of ethidium bromide (EthBr) in the indicated strains, as measured by fluorescence at 590 nm for 65 min. Data are mean and standard deviation of three biological replicates. **(B)** Sensitivity of the indicated strains to different concentrations of vancomycin as measured by monitoring optical density at 580 nm (OD580) after 7 days. Data are mean and standard deviations of three biological replicated. Errors bars not shown are smaller than symbols. **(C)** Time course of the incorporation of FDAAs on the indicated strains in 7H9 as measured by fluorescence. Cells were labeled with HADA for 48 h, washed in 7H9 medium and labeled with FDL for 5 h. Ten, cell were fixed and imaged under a fluorescence microscope. Representative images of WT, *virR^{mut}* and *virR^{mut}*-Comp strains in 7H9. Data are mean and standard deviations of three biological replicates. *** means *P*<0.001 and **** means *P*<0.0001 after applying a One-way ANOVA analysis. The arrow indicates the time at which images were taken. Errors bar not shown are smaller than symbols. **(D)** immunoblot analysis of cell lysates of two independent conditional mutants (c1 and c2) for *virR* and *cei* for the presence of VirR and Cei proteins in cultures incubated with and without anhydrotetracycline (ATc). **(E)** Nanoparticle tracking analysis (NTA) of EV preparations derived from *virR* and *cei* conditional mutants (c2 (*virR*) and c1 (*cei*)) obtained from cultures with and without ATc showing number of particles per cm^3 determined by Zeta View NTA in three independent EV preparations. Data are presented as mean ± SEM.

Figure 3. Transcriptional profiling of *virR^{mut}* showcases systemic alterations in cell wall architecture and metabolism. (A) Experiment design for RNA-seq analyses. RNA

was extracted from whole cell extracts from 4 replicate cultures of each strain (WT,
virR^mut, and *virR^mut* -Comp), and sequenced. **(B)** PCA plot of the transcriptomics data.
The loadings from the two first principal components are plotted. **(C)** Number of
differentially expressed genes between *virR^mut* and either WT (left) or *virR^mut*-Comp
(right). **(D)** Enrichment profiles between sets of genes found up and down-regulated for
the contrasts *virR^mut* vs WT and *virR^mut* vs *virR^mut*-Comp. While genes that are
differentially expressed in the same direction in both contrasts are mutually enriched,
genes whose differential expression goes in different directions in both comparisons are
mutually depleted (two-sided Fisher exact test p<0.05 in all four cases). **(E)** Gene
Ontology (GO) enrichments of genes that are 1) simultaneously downregulated in *virR^mut*
with respect to both WT and v *virR^mut*-Comp (left) and 2) simultaneously upregulated
(right). GO terms selected for testing included biological process and cell compartment
labels containing N > 10 genes, between levels 4-6 of the ontology tree. In this
representation, each node in the networks represents a significantly enriched term (terms
selected for visualization with FDR=10% and enrichment odds-ratio>2), nodes' size is
proportional to the enrichment significance, and links between terms are added wherever
the sets of genes contributing to the enrichments in a connected pair show an
intersection larger than 50% of the smallest of the gene sets involved in the pair (see
Methods for further details, and supplementary file S2 for an extended list of all the
enrichments found).

Figure 4. VirR-dependent genes intersect the regulons of key transcriptional
regulators of the responses to stress, dormancy, and cell wall remodeling. (A)
Transcription factors controlling regulons significantly enriched among genes that are 1)
simultaneously downregulated in *virR^mut* with respect to both WT and *virR^mut* -Comp
(bottom, 9 TFs) and 2) simultaneously upregulated (top, 1 TF). **(B)** Differential expression
patterns for enriched TFs whose expression is VirR-dependent. **(C)** Regulatory
subnetwork including the genes under transcriptional control of the enriched TFs
highlighted in figure 4A that were found to be VirR-dependent. In this visualization, color
captures the logFC for the contrast *virR^mut* -WT, and the size of its FDR.

1633

Figure 5. Cell permeability is a major driver of vesicle production in *Mtb*. (A) Time
course of the uptake of ethidium bromide (EthBr) by *Mtb* grown in the indicated
conditions, as measured by fluorescence at 590=nm for 65 min. Data are mean and
standard deviation of three biological replicates. **(B)** Dot-blot analysis of released EVs in
supernatants from *Mtb* cultures submitted to the indicated conditions. The loading
volume was normalized according to CFUs. The graph below represents the
quantification of the dotblot using ImageJ. The integrated cell density was normalized to

WT values. Data are mean and standard errors from the three biological replicates; ****

denotes P<0.0001, *** denotes P=0.0028 after applying a Tuke's multiple comparison

test. **(C)** Enrichment odds ratios for Fisher exact tests comparing sets of genes up, and

down regulated, respectively, in 1) cholesterol-supplemented medium vs. glucose-

supplemented medium (Pawełczyk et al., 2021), GEO accession GSE175812); and 2)

virR^{mut} vs WT as well as *virR^{mut}* vs *virR^{mut}*-Comp. **(D)** Euler diagram showing the

intersection between A) genes upregulated by cholesterol, B) genes down-regulated in

virR^{mut}, and C) the DevR regulon. **(E)** Expression patterns in response to TRZ (1h) and

in *virR^{mut}* for genes involved in metal signaling, metabolism and homeostasis.

1650

Figure 6. VrR regulates the protein content of EVs in Mtb. (A). Experiment design for

proteomics analyses. label-free mass spectrometry data was retrieved for replicates of

H37RV (WT), *virR^{mut}* and *virR^{mut}*-Comp. For each strain, data corresponding to proteins

extracted from either whole-cell extracts (WC) and EVs was produced, totaling 12

samples (2 strains times 2 cell compartments times 3 replicates). **(B)** PCA plot of the

proteomics data. The loadings from the two first principal components, adding up to

50.5% of variance explained, are plotted. **(C)** Expression patterns of VirR in the

proteomics dataset across cell compartments and strains. As expected, while *virR^{mut}*

shows significantly lower levels of VirR protein, the complemented strain does not just

restore WT levels of expression, but leads to a strong overexpression of VirR, both in

the lysates and the EVs, **(D)** Bar plots of differentially expressed proteins found between

strains in EVs (up: *virR^{mut}* vs WT; down *virR^{mut}* vs *virR^{mut}*-Comp at 5% FDR). **(E)**

Enrichment profiles between sets of proteins found up and down-regulated for the

contrasts *virR^{mut}* vs WT and *virR^{mut}* vs *virR^{mut}*-Comp. **(F)** Gene Ontology enrichments of

the downregulated proteins in the VirR mutant strain (left) and the upregulated ones

(Right), at the EV location. GO terms selected for testing included biological process and

cell compartment labels containing N > 3 genes, between levels 4-6 of the ontology tree.

As in **figure 3E**, terms selected for visualization with FDR=10% and enrichment odds-

ratio>2, nodes' size is proportional to the enrichment significance, and links between

terms are added wherever the sets of proteins contributing to the enrichments in a

connected pair shows an intersection larger than 50% of the smallest of the gene sets

involved in the pair (see Methods for further details, and supplementary file S5 for an

extended list of all the enrichments found). **(G)** Expression patterns of bacterioferritin

proteins in the EVs of WT and *virR^{mut}*. **(H)** Expression patterns of proteins contributing to

enrichments of terms related to mycolic acids biosynthesis. **(I)** Bottom left:

Concentrations of key metabolites in the propionyl-coa detoxification routes. Right:

Metabolic pathway of the propionyl-coa detoxification route through the methyl-malonyl

route, along with the differential expression statistics between the WT and *virR^{mut}* strains

observed at transcriptomic and proteomic levels for the main enzymes involved. The

framed plot represents the quantification of selected metabolites by mass spectrometry.

1681

Figure 7. Lipidomic analysis of cell envelope and EVs from *virR^{mut}*. (A) Comparative

total lipid analysis of the indicated strains by nano-LC-MS. Lipid species with an

abundance higher than 0.5% are shown. Data indicates mean and standard error from

three biological replicates. **(B)** Analysis of polar lipids of the indicated strains by

bidimensional (2D)-layer chromatography (TLC) (2D-TLC). Phosphatidyl-myo-inositol

dimannosides (Ac_1PIM_2 and Ac_2PIM_2, respectively); phosphatidyl-myo-inositol

hexamannosides (Ac_xPIM_6); phosphatidylethanolamine (PE); diphosphatidylglycerol

(DPG). Phospholipid (P). **(C)** Analysis of mycolic acid species in the indicated strains by

TLC. **(D)** Analysis of PIDMs and TAGs by TLC. **(E)** Analysis of apolar lipids of the

indicated strains by TLC. Diacylglycerol (DAG); free fatty acid (FF); free mycolic acid

(FMA). **(F)** Analysis of apolar lipids of MEVs isolated from the indicated strains.

Triacylglycerols (TAGs); phthiocerol dimycocerosates (PDIMs) **(G)** Analysis of polar

lipids of MEVs isolated from the indicated strains. Phosphatidyl-myo-inositol

dimannosides (Ac_1PIM_2 and Ac_2PIM_2, respectively); phosphatidyl-myo-inositol

hexamannosides (Ac_xPIM_6); diphosphatidylglycerol (DPG).

1697

Figure 8. Chemical and nanometric analysis of isolated cell walls shows higher

amounts of PG in the absence of VirR. (A) Glycosyl composition analysis of isolated

cell walls from the indicated strains. Acid hydrolysis prior to GC-MS analysis of isolated

cell walls was performed to determine the total amounts of galactose (G), arabinose (A),

rhamnose (R), and N-acetylglucosamine (NAGc). The ratio between different sugars is

shown to indicate: (i) the relative levels of arabinogalactan (A/G); (ii) the length of the AG

polysaccharide chain (G/R); (iii) the amount of PG (NAGc/Rham). Individual

measurements are shown from three biological replicates. Data are mean and standard

errors. **P < 0.01 after applying a Tukey´s multiple comparison test. **(B)** Quantification

of diaminopimelic acid (DAP) on isolated cell walls from the indicated strains. Data are

mean and standard errors from three biological replicates; **** denotes P<0.0001, *

denotes P=0.0119 after applying a Tukey´s multiple comparison test. **(C)** Atomic force

microscopy high resolution images taken in liquid from the external surface of purified

peptidoglycan from different samples, left to right: WT, *virR^{mut}*, *virR^{mut}-C*, the structure of

external peptidoglycan layer shows a mesh with pores of different sizes, **(D)** Graph

showing the pore diameter from n = 8 AFM images similar to **(C)** (each point represent

an image) of different samples; black triangles: WT, blue circles: *virR^{mut}*, red squares:

virR^{mut}-C. The pore diameter was calculated from the binary slice from each image

containing the greatest number of pores, according to (Pasquina-Lemonche, 2024). **(E)**

The number of pores per surface area from different samples; white triangles: WT, red

circles: *virR^{mut}*, blue squares: *virR^{mut}*-C, this was calculated from the slice containing the

maximum number of pores where the pore size was analyzed, following the method from

(Pasquina-Lemonche, 2024). **(F)** Graph showing the peptidoglycan thickness manually

measured using the 1D statistic tools from the open-source program Gwyddion (Nečas

and Klapetek, 2012) from AFM images in air containing several fragments of cell wall,

each point represents a different peptidoglycan fragment from a different cell. There were

three samples analyzed: black WT n=23 peptidoglycan fragments, blue *virR^{mut}* n=36 and

red *virR^{mut}*-C n=25. Statistics: D) Using a two-tailed t test with Welch's correction the

statistical comparison are: (***) pWT- *virR^{mut}* = 4.1 10^{-4} with t = 6.1, df = 7.5, (**) p *virR^{mut}*

- *virR^{mut}*-C = 0.007 with t = 3.2, df = 13.2, E) Using a two-tailed t test with Welch's

correction the statistical comparison are: (****) pWT- *virR^{mut}* = 6.6 10-10 with t = 16.3,

df = 12.7, (**) p *virR^{mut}* - *virR^{mut}* -C = 0.01 with t = 3.0, df = 13.2., F) Using a two-tailed t

test with Welch's correction the statistical comparison are: (****) pWT- *virR^{mut}* = 2.8 10-

18 with t = 17.8, df = 32.3, (****) *virR^{mut}* - *virR^{mut}* -C = 3.9 10-17 with t = 16.4, df = 31.8.

(G) List of muropeptides identified in the cell walls of Mtb. Features including retention

time (RT), observed and expected mass as well as the best match for each peak are

shown. Those deacetylated muropeptides are indicated in red. **(H)** Relative abundance

of deacetylated muropeptides in the indicated strains.

1736

Figure 9. VirR interacts with Lcp1. (A) Blue-native PAGE analysis of recombinant VirR

and Lcp1. The molecular mass of markers in kDa are indicated on the left side of the gel.

The size of the multimers of VirR and Lcp1 are indicated on the left side of the gel. **(B)**

Upper panel. Representative image of a flotation assay of recombinant VirR alone or in

combination with Lcp1 performed on an idioxanol density gradient. The presence of VirR

was detected in each fraction by dot-blot using murine polyclonal antibodies raised

against VirR. Numbers indicate collected fractions from top (1) to bottom (11). *Lower*

panel. Quantification of the dot-blot by measuring the pixel intensity of the dots using

ImageJ. Data are mean and standard error of three independent experiments. **(C)**

Bipartite split-GFP experiment using *M. smegmatis* D2 expressing plasmids depicted in

(**Fig S11A**), including $CtpC_{Nterm}$, VirR, VirRsol (Δ1-41), Rv3484, Rv3267 (Lcp1), Rv0822,

$Rv3484_{LytR\_C}$, $Rv3267_{LytR\_C}$ and $Rv0822_{LytR\_C}$. Bacteria were grown in complete 7H9 and

fluorescence was recorded by FACS. Pooled data obtained from four to six independent

cultures of each strain are shown. Statistical analysis was performed using non-

parametric two-sided Mann-Whitney test. ns: not significant. * and ** refer to p-values

<0.05 or 0.01, respectively.

1753

1754 **Supplementary Tables**

1755

1756 **Supplementary Table S1.** Plasmids and primers used in this study.

Construction / DNA fragment used	Template	Primers	Sequence 5'-3'
pGMCS-P1-*virR*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)			
P1	pGMCS-TetR-P1-*pacL1*Flag- P*pacL1*-*ctpC*His (Boudehen *et al.* 2022)	Inf-B2-P1-Fw	CAAAGTGGAGCGATACTCGACATCGGATCGTCGGCACCGTCAC
		Inf-P1-Left	GAAATGATGTATGCCGTGCTGGTC
pGMCS backbone	pGMCS-P*pacL*-*pacL1*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10) (Boudehen *et al.* 2022)	ctpC-B2-Left	GTCGAGTATCGCTCCACTTTG
		Rv3269-P1-Fus-Fw	CAGCACGGCATACATCATTTCGGAAGAAAGGTACAGGCAATGG
pGMCS-P1-*virR*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)			
virR	*M. tuberculosis* H37Rv chromosomal DNA	If-AmL1-VirR	AAGGTACAGGCAATGgtgctggttacagtgggctcg
		If-Lk2-VirR-Rv	ACCGGATCCTTCCAGgccggtcaccacgacgatg
pGMCS backbone	pGMCS-P1-*pacL1*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Lk2-Right	CTGGAAGGATCCGGTGGCGG
		pGMC 3269 Am Left	CATTGCCTGTACCTTTCTTCC
pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)			
virRsol	*M. tuberculosis* H37Rv chromosomal DNA	If-AmL1-VirRsol	AAGGTACAGGCAATGctgggttcgtctccgaactccg
		If-Lk2-VirR-Rv	ACCGGATCCTTCCAGgccggtcaccacgacgatg
pGMCS backbone	pGMCS-P1-*pacL1*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Lk2-Right	CTGGAAGGATCCGGTGGCGG
		pGMC 3269 Am Left	CATTGCCTGTACCTTTCTTCC
pGMCS-P1-*virR*-Lk2-GFP11-*virR*-Lk15-GFP(1-10)			
virR	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-VirR	GAGCGCCTCGCCATGgtgctggttacagtgggctcg
		If-L15-VirR-Rv	AGACAGCCCTGGACCgccggtcaccacgacgatg
pGMCS backbone	pGMCS-P1-*virR*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*virRsol*-Lk15-GFP(1-10)			
virRsol	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-VirRsol	GAGCGCCTCGCCATGctgggttcgtctccgaactccg
		If-L15-VirR-Rv	AGACAGCCCTGGACCgccggtcaccacgacgatg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv0822c*(LytR_C)-Lk15-GFP(1-10)			
Rv0822c(LytR_C)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-0822cD	GAGCGCCTCGCCATGgagattcagcaccagcaggttacgacg
		If-L15-0822cD-Rv	AGACAGCCCTGGACCgcggcttatctgcacgctgacg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3267*(LytR_C)-Lk15-GFP(1-10)			

Rv3267(LytR_C)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-3267D	GAGCGCCTCGCCATGcccgccaagaccacggcc
		If-L15-3267D-Rv	AGACAGCCCTGGACCcagcccggagcccggac
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3484*(LytR_C)-Lk15-GFP(1-10)			
Rv3484(LytR_C)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-3484D	GAGCGCCTCGCCATGgcggccgccaaacctaaccc
		If-L15-3484D-Rv	AGACAGCCCTGGACCggtatccaccgtcacacggatg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv0822c*(Cterm)-Lk15-GFP(1-10)			
Rv0822c(Cterm)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-0822cD	GAGCGCCTCGCCATGgagattcagcaccagcaggttacgacg
		If-L15-0822c-Rv2	AGACAGCCCTGGACCctcgcaggtggtgtcggcg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3267*(Cterm)-Lk15-GFP(1-10)			
Rv0822c(Cterm)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-3267D	GAGCGCCTCGCCATGgagattcagcaccagcaggttacgacg
		If-L15-3267-Rv2	AGACAGCCCTGGACCgttgatgcactccggcgcgtc
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3484*(Cterm)-Lk15-GFP(1-10)			
Rv0822c(Cterm)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-3267D	GAGCGCCTCGCCATGgagattcagcaccagcaggttacgacg
		If-L15-3484-Rv2	AGACAGCCCTGGACCgttcacgcagggcacgccg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv0822c*-Lk15-GFP(1-10)			
Rv0822c	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-0822c	GAGCGCCTCGCCATGagtgacggcgagagcg
		If-L15-0822c-Rv	AGACAGCCCTGGACCctactcgcaggtggtgtcgg
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3267*-Lk15-GFP(1-10)			
Rv3267(Cterm)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-3267	GAGCGCCTCGCCATGatgtctgcgcaacgtgtggttc
		If-L15-3267-Rv	AGACAGCCCTGGACCtcagttgatgcactccggcgc
pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	Flex15-Right	GGTCCAGGGCTGTCTGGCC
		ctpC-Am-left2	CATGGCGAGGCGCTCAAG
pGMCS-P1-*virRsol*-Lk2-GFP11-*rv3484*-Lk15-GFP(1-10)			
Rv3484(Cterm)	*M. tuberculosis* H37Rv chromosomal DNA	If-AmMBD-3484	GAGCGCCTCGCCATGgcgcgttctgagggc
		If-L15-3484-Rv	AGACAGCCCTGGACCctagttcacgcagggcacgc
		Flex15-Right	GGTCCAGGGCTGTCTGGCC

pGMCS backbone	pGMCS-P1-*virRsol*-Lk2-GFP11-*ctpC*(1-85)-Lk15-GFP(1-10)	ctpC-Am-left2	CATGGCGAGGCGCTCAAG
CRISPRi mutants			
virR (158.1) c2	Rv0431	Rv0431crispr 1 F	GGGAGAGCAGCAGGAAGACGACGC
		Rv0431crispr 1 R	AAACGCGTCGTCTTCCTGCTGCTC
virR (42.2) c1	Rv0431	Rv0431crispr 2 F	GGGAACGAACCCAGTGCCTGCCAGA
		Rv0431crispr 2 R	AAACTCTGGCAGGCACTGGGTTCGT
cei (145.2) c1	Rv2700	Rv2700crispr 1 F	GGGAATCACGCAAGTCACCACCAC
		Rv2700crispr 1 R	AAACGTGGTGGTGACTTGCGTGAT
cei (47.3) c2	Rv2700	Rv2700crispr 2 F	GGGAGTCGGTGCCGAGCGCAAGGTCG
		Rv2700crispr 2 R	AAACGACCTTGCGCTCGGCACCGAC
VirR-Venus primers			
VirR-F			TCACTGGGGTGCTGGTTACAGTGGGCTC
VirR-R			ATAGGATCGCCGGTCACCACGACGAT
pMV-F			TGACCGGCGATCCTATGGTGAGCAAGGGC
pMV-R			ACCAGCACCCCAGTGAAAAGTTCTTCTCCTTTACTC

1757

1758 **Supplementary Table 2**. Gradient mobile phase composition for ion-pairing LC-MS/MS
1759 for the quantification of acetyl-CoA, methyl citrate and methylmalonyl-CoA.

Time (min)	Mobile Phase (%)	
	A	B
1	100	0
2	55	45
7	50	50
7.5	5	95
8	5	95
8.5	100	0
10	End	

1760 A: water-methanol 07:3 (v/v) + 10mM tributylamine + 3mM acetic acid; B: pure methanol
1761

1762 **Supplementary Table 3**. Gradient mobile phase composition for ion-pairing LC-MS/MS
1763 for the quantification of pyruvate.

Time (min)	Mobile Phase (%)	
	A	B
1	5	95
6	95	5
7	95	5
8	5	95
10	End	

1764 A: pure water; B: pure methanol
1765

1766 **Supplementary Table 4**. Quantifier MRM transitions and collision energies used for
1767 acety-CoA, methyl citrate, methylmalonyl-CoA and pyruvate.

Metabolite	Precursor ion [M-H]⁻ (m/z)	Product ion (m/z)	Collision energy (eV)

Acetyl-CoA	808.00	407.95	41
Methyl citrate	204.90	125.10	13
Methylmalonyl-CoA	432.60	410.50	10
Pyruvate	87.20	43.10	13

1768
1769
1770
1771

Supplementary figures

Fig S1. Diversity of *Mtb* LCP proteins. (**A**) Domain structure of LCP proteins in *Mtb*. Proteins are organized according to the presence of (i) both LCP (green) and LytR_C (slight orange) terminal domain (Rv3267, Rv3484 and Rv0822); (ii) the solo LytR_C domain (VirR and Rv2700); and the solo LCP domain (Rv3840). Additional information is provided for *Mtb* mutants in the indicated genes. (**B**) Phylogenetic analysis of LCP proteins in Mycobacteria (*M. tuberculosis, M. smegmatis, M. marinum and M. leprae*). The unrooted tree includes seven separate clusters for proteins with both LCP and LytR_C terminal domains (CpsA1, CpsA2 and CpsA3); with the solo LytR_C terminal domain (Solo_LytRC1 and Solo_LytRC2); with the solo LCP domain; and a separate cluster including the characterized LCP proteins from *S. pneumonia* and *S. aureus*. Note the clustering of one of the *M. smegmatis* LCP proteins with this group. VirR is highlighted in blue.

Fig S2. The enhanced vesiculation phenotype of *virR^mut^* occurs in the absence of cell lysis. (**A**) Nanoparticle tracking analysis (NTA) of EV preparations derived from WT and the indicated strains showing number of particles per cm^3 determined by Zeta View NTA in three independent EV preparations. Data are presented as mean ± SEM. *$P \leq 0.05$ (One-way ANOVA with Tukey's multiple comparisons test). (**B**) Cell lysis control of *virR^mut^*. Shown are immunoblots of cell lysates and culture supernatants (SN) for the indicated strains. The cytoplasmic protein IdeR was detected in cell lysates but not in culture supernatants confirming cell integrity, while the secreted protein Ag85b was readily detected in the culture supernatants. (**C**) Cell lysis control of *Mtb* submitted to conditions that alter permeability. Arrows indicate the size of the proteins of interest detected by immunoblot. (**D**) Alternatively, cell viability was tested by serial dilution and plating onto 7H10 medium. Cultures from B and C were submitted to serial dilutions (10^{-1} to 10^{-5}) and plated onto 7H10 after reaching an OD_{600nm} of 0.2.

Fig S3. H2O2 sensitivity assay. The indicated strains were initially grown on 7H9 medium supplemented with AND to an OD of 0.5. Washed cells were then transferred to

1802 the same medium supplemented with and without 13mM H202. After 8 h, cells were

1803 diluted down as indicated and spotted onto 7H10 solid medium.

1804 **Fig S4. Expression patterns of permeability-dependent genes upregulated in**

1805 **cultures exposed to cholesterol and simultaneously downregulated in *virR^{mut}*.**

1806

Fig S5. Behaviour of *virR^{mut}* upon conditions that modulate permeability. (A)

Growth curves of *virR^{mut}* in MM supplemented with either glycerol, cholesterol or glycerol

plus TRZ at the indicated concentrations. **(B)** Dotblot analysis of vesiculation levels of

virR^{mut} on the indicated culture media. a, b and c denote biological replicates.

Quantification of the dotblot was done by measuring the integrated density of each dot

using ImageJ. Data are mean and standard deviations. * p=0.021 after applying a Tuke's

multiple comparison test. **(C)** Time course of the uptake of ethidium bromide (EthBr) by

virR^{mut} grown in the indicated conditions, as measured by fluorescence at 590=nm for 65

min. Data are mean and standard deviation of three biological replicates.

1816

1817 **Fig S6. Expression patterns of ribosomal proteins in the EVs from WT, *virR^{mut}***

1818 **and *virR^{mut}* -Comp strains. Proteins shown are downregulated in the vesicles of**

1819 ***virR^{mut}* with respect to both VirR expressing strains.**

1820

1821 **Fig S7. Expression patterns in the EVs from WT, *virR^{mut}* and *virR^{mut}* -Comp strains**

1822 **of proteins differentially expressed in *virR^{mut}* that contribute to enrichments of**

1823 **terms related to metabolism of: (A) nucleotides (B) amino-acids (C)**

1824 **oligosaccharides (D) glycogen.**

1825

1826 **Fig S8. TLC analysis of strains grown in the presence of cholesterol or TRZ.**

1827 Analysis of polar lipids, including SL1 and DATs lipids of the indicated strains by TLC.

1828 Sulfolipid 1 (SL1); diacylglytrehaloses (DATs); Trehalose dimycolate (TDM).

1829

Fig S9. Additional AFM data. (A-C) (i) Atomic force microscopy images taken in liquid

from a big area showing the sample heterogeneity overview with various peptidoglycan

fragments of the following samples: **(A)** WT, **(B)** *virR^{mut}*, **(C)**, *virR^{mut}*-C. **(ii)** Atomic force

microscopy images zoom in one individual peptidoglycan fragment corresponding to one

cell from different samples: **(A)** WT, **(B)** *virR^{mut}* showing the background (Backg.), the

external surface (Ext) the internal (Int), **(C)**, *virR^{mut}*-C, white square showing the size of

the zoom images shown in Figure 5 from the external surface of the cells. **(D)** Curve of

grey points showing 255 slices of one AFM image taken using a FIJI macro, in the x axis

and the total number of pores in the y axis, the red line is a Gaussian fitted to the curve

and the maximum of the Gaussian is marked with a dash red line showing slice 140 in this example corresponding to the slice of the image with the maximum number of pores. **(E)** Binary slice number 140 from the original AFM image showing the pores in white and the material represented in black. **(F)** The pores from 'E' were analyzed using the FIJI macros where each pore area was measured and plotted in the x axis against the cumulative fraction of total pore area in the y axis as shown in the curve by blue dots. The red dot marks the most likely pore area which corresponds to the half of the cumulative fraction of the total pore area, this area was then used to obtain the pore diameter of the image. The same was repeated with the n = 8 images for each sample to compose the graphs 'D' and 'E' from Figure 7.

Fig S10. **Sensitivity to lysozyme of *virR^mut*.** **(A)** Analysis of the minimum inhibitory concentration (MIC) of lysozyme of the indicated strains by the resazurin method. **(B)** Growth of WT and *virR^mut* treated with 0 or 50 µg/ml lysozyme (Lys). Data represents the mean ± SEM of three independent cultures.

Fig S11. Localization of VirR by fluorescence microscopy. Fluorescence images of a *Mtb* strain expressing Venus-tagged VirR.

Fig S12. Split-GFP strategy. (A) *Upper panel*. Scheme of the plasmid used to test VirR interactions in vivo. *Lower panel*. Scheme of the potential molecular interaction between VirR and Lcp1 using split-GFP. **(B)** Gating strategy to analyze split-GFP flow cytometry data. Bacteria were first sorted based on size (FSC) and structure (SSC) to exclude debris (1). Then, aggregates were excluded based on FSC and SSC width (W) and eighth (H) (2,3). GFP signal was then analyzed *vs*. SSC (4) or mCherry, when relevant (5).

Supplementary file S1: FileS1_RNA_differential_expression.xlsx: Results of the differential expression analyses for the RNA-seq dataset.

Supplementary file S2: FileS2_GOenrichments_RNA.xlsx: Gene ontology enrichment analyses conducted on sets of genes up (N=399), and down-regulated (N=502) in the *virR^mut* strain. Rows highlighted correspond to terms selected for graphical representation (figure 3E, OR>2 & FDR<0.1)

Supplementary file S3: FileS3_TRN_regulon_enrichments.xlsx: Enrichment analyses for the Transcription factor regulons reported in (Minch et al. 2015).

1876

1877   **Supplementary file S4**: FileS4_Proteins_differential_expression.xlsx: Results of the

1878   differential expression analyses for the LF-MS dataset.

1879

1880   **Supplementary file S5**: FileS5_GOenrichments_proteins.xlsx: Gene ontology

1881   enrichment analyses conducted on sets of protein up (N=117), and down-regulated

1882   (N=108) in the extracellular vesicles of the $virR^{mut}$ strain. Rows highlighted correspond

1883   to terms selected for graphical representation (figure 5F, OR>2 & FDR<0.1).

1884

	CM	CM-OM	CM-L1	OM-L2	L1-L2	OM
WT	2.7±0.2	44.09±3.1	11.2±0.4	13.34±0.9	10.46±1.4	5.7±0.4
virR^mut	5.4±0.7*	62.7±3.2*	31.36±3.2*	15.1±2.2	6.8±0.9*	6.1±0.3
virR^mut-C	2.5±0.5	41.6±0.7	12.4±1.4	10.03±1.1	12.6±1.3	5±0.2

Figure 2

Peak	R.T	Observed	Expected	Best Match
1	4.3	942.432	942.4155	M4
2	6.12	1752.80	1752.7379	D43 (-N-Ac)
3	6.59	1823.856	1823.7750	D44(-N-Ac)
4	6.62	1,823.862	1823.7750	D44 (-N-Ac)
5	6.87	1794.8	1794.7755	D43
6	7.1	1794.8	1794.7755	D34
7	7.76	1865.854	1865.8126	D44
8	8.82	1831.86	1831.748	D45N(-NH2,-2N-Ac)
9	9.7	2789.26	2789.2098	T444
10	10.56	1845,806	1774.7493	D44N (Anh)

	Growth defect	Release material	Attenuated Virulence
Rv3267	Yes	No	No
Rv3484	No	No	Yes
Rv0822	No	No	No
Rv2700	Yes	?	Yes
VirR	-/+	+++	Yes
Rv3840	?	?	?

Fig S1

Fig S2

untreated

13 mM H$_2$O$_2$

10^{-1} 10^{-2} 10^{-3} 10^{-4} 10^{-1} 10^{-2} 10^{-3} 10^{-4}

WT

VirR^{mut}

VirR^{mut}-C

Fig S3

Fig S4

Fig S5

Fig S6

carbohydrate catabolic process

Standardized expression

- 1.0
- 0.5
- 0.0
- −0.5
- −1.0
- −1.5

WT *virR^{mut}*-C *virR^{mut}*

fatty acid metabolic process

Standardized expression

- 2
- 1
- 0
- −1

WT *virR^{mut}*-C *virR^{mut}*

Fig S7

A

GLY　　CHOL　　TRZ

WT　*VirR*^mut　*VirR*^mut_C　WT　*VirR*^mut　*VirR*^mut_C　WT　*VirR*^mut　*VirR*^mut_C

DPG
PE
P
Ac$_2$PIM$_2$

Ac$_x$PIM$_6$

O

B

GLY　　CHOL　　TRZ

WT　*VirR*^mut　*VirR*^mut_C　WT　*VirR*^mut　*VirR*^mut_C　WT　*VirR*^mut　*VirR*^mut_C

TDM

SL

DAT

PolG
O

FigS8

| A | WT | B | *virR*^{mut} | C | *virR*^{mut}-C |

Fig S9

Fig S10

5 µm

Fig S11

A

pGMCS-P1-VirR-Lk10-GFP11-Lcp-Lk15-GFP(1-10)

Int   AttL5   Strepto^R   Ori colE1

GFP11   GFP1-10   Full GFP

VirR   Lcp   VirR   Lcp

B

Fig S12