Challenges in estimating species age from phylogenetic trees

Carlos Calderón del Cid[1,2], Torsten Hauffe[2], Juan D. Carrillo[2], Rachel C. M. Warnock[3], Daniele Silvestro[2,4]

[1]Laboratório de Ecologia Espacial, Instituto de Biologia, Universidade Federal da Bahia, CEP 40170-110, Salvador, Bahia, Brazil

[2]Department of Biology, University of Fribourg, Switzerland and Swiss Institute of Bioinformatics, Fribourg, Switzerland

[3]GeoZentrum Nordbayern, Friedrich-Alexander Universität Erlangen-Nürnberg, Loewenichstrasse 28, 91054, Erlangen, Germany

[4]Department of Biological and Environmental Sciences and Gothenburg Global Biodiversity Centre, University of Gothenburg, Sweden

Abstract

Aim. Species age, the elapsed time since origination, can give an insight into how species longevity might influence eco-evolutionary dynamics and has been hypothesized to influence extinction risk. Traditionally, species ages have been measured in the fossil record. However, recently, numerous studies have attempted to estimate the ages of extant species from the branch lengths of time-calibrated phylogenies. This approach poses problems because phylogenetic trees contain direct information about species identity only at the tips and not along the branches. Here, we show that taxon sampling, extinction, and different assumptions about speciation modes can significantly alter the relationship between true species age and phylogenetic branch lengths, leading to high error rates. We find that these biases can lead to erroneous interpretations of eco-evolutionary patterns derived from the comparison between phylogenetic age and other traits, such as extinction risk.

Innovation. For bifurcating speciation, which is the default assumption in most analyses, we propose a probabilistic approach to improve the estimation of species ages, based on the properties of a birth-death process. We show that our model can reduce the error by one order of magnitude under cases of high extinction.

Main conclusion. Our results call for caution in interpreting the relationship between phylogenetic ages and eco-evolutionary traits, and show that, under some assumptions, it is possible to obtain better approximations of species age by combining information from branch lengths with the expectations of a birth-death process.

Key words: Evolutionary history, Extinction rates, Incomplete sampling, Simulations, Speciation modes, Phylogeny.

Introduction

The estimation of species age, or the elapsed time since species origin, is important to evaluate mechanisms that link species longevity with eco-evolutionary processes (Benton, 2013; Swenson, 2019). For instance, age-dependent extinction hypotheses test the relationship between species age and extinction probability, assessing whether extinction rates differ between young and old species (Balmford, 1996; Eldredge et al., 2005; Pearson, 1995). Likewise, species age could be a measure of colonization time, especially in island systems (Tanentzap *et al.* 2015) or during biotic invasions triggered by geological events, such as the formation of the Central American Isthmus for the Great American Interchange (Carrillo *et al.* 2015, 2020). Species age is measured in the fossil record through different statistical and probabilistic approaches, based mostly on taxa's stratigraphic duration (i.e., the time between the first and last appearance of a taxon in the fossil record) (Foote, 1996; Foote & Raup, 1996). Several of these approaches consider differences in fossil sampling and temporal resolution (Alroy et al., 2001; Silvestro et al., 2019). Species ages estimated from paleobiological data offer a reliable measure of species' temporal duration which can be used in macroevolutionary studies (Benton, 2016; Silvestro et al., 2020; Van Valen, 1973). More recently, several studies have used the length of terminal branches in time-calibrated phylogenies as a proxy for the age of extant species, an approximation that we hereafter refer to as "phylogenetic age" (Alzate et al., 2023; Davies et al., 2011; Gaston & Blackburn, 1997; Johnson et al., 2002; Pie & Caron, 2023; Sonne et al., 2022; Tanentzap et al., 2020; Verde Arregoitia et al., 2013). These phylogenetic ages have been used as the basis to test for links between species age and current extinction risks (Tanentzap et al., 2020; Verde Arregoitia et al., 2013) and to assess various correlations with evolutionary, biogeographical, and ecological patterns in living species (Alzate et al., 2023; Freer et al., 2022; Kennedy et al., 2022; Pie & Caron, 2023)

While several studies have used phylogenetic age at face value for species age (e.g., Johnson et al. 2002; Tanentzap et al. 2020; Verde Arregoitia et al. 2013), their potential deviation from the true species ages remains unclear. Specifically, we identify three non-mutually exclusive shortfalls that can lead to over- or underestimation of species ages. First, incomplete sampling of extant species, either due to incomplete species sampling or linked

to species still being unknown to science, can bias phylogenetic age estimation by artificially

increasing the length of terminal branches (Heath et al., 2008; Mynard et al., 2023).

Second, extinction events will mask branching events in phylogenetic trees of extant

species (Harvey et al., 1994; Nee & May, 1997). Even in phylogenetic trees that include

extinct taxa, the incompleteness of the fossil record will inevitably lead to missing lineages

and incorrect topologies. Unsampled extant and extinct species from the phylogeny results

in an inflation of the length of terminal branches leading to sampled species (i.e., the tips of

the tree), thus altering phylogenetic species ages. For instance, if the extinct species of the

Homo genus are not included in a phylogeny, the phylogenetic age of *Homo sapiens* is

approximately 10 million years, i.e., the age of the last common ancestor with its sister

species, the chimpanzee (Rivas-Gonzáles et al. 2023). This estimate exceeds the age of the

oldest known fossil of modern humans (i.e., *Homo sapiens*) by two orders of magnitude (Fig.

1; Callaway 2017).

The third shortfall, is that the tree alone does not contain information about the

underlying speciation mode and does not include species labels along its branches, such that

only the tips can be unequivocally assigned to a named species (Losos & Glor, 2003).

Alternative speciation modes have been discussed in the literature reflecting different

biological processes and species concepts, including bifurcating, budding, and anagenetic

speciation (Foote, 1996; Silvestro et al., 2018). These modes define the relationship between

the ancestral species and its descendants, thus contributing to determining species ages

(Rosenblum et al., 2012; Wagner, Erwin, & Anstey, 1995) (Fig. 2). Most phylogenetic trees

are depicted in the rectangular shape, where the two descending lineages split symmetrically

from an ancestral lineage, thus suggesting a bifurcating speciation mode where two new

species replace the ancestral lineage (Baum et al., 2005; Caetano & Quental, 2022). However,

the often-unstated assumption of virtually all birth-death processes used to model

phylogenetic branching times, is that speciation occurs as a budding process, with a

speciation event leading to a single new species and the survival of the parent species, even

though we cannot determine which descendant branch is the new species (Gernhard, 2008;

Nee et al., 1994; Stadler, 2013). Anagenetic speciation, in contrast, does not lead to a

branching event and is therefore not ordinarily visible on a phylogenetic tree.

All speciation modes may reflect plausible biological processes, and evidence for
each mode has been found in the fossil record (Foote, 1996; Simpson, 1984) and in extant
species (Skeels & Cardillo, 2019). Under bifurcating speciation mode, phylogenetic ages
equal species ages when there is no extinction, and all species are sampled. A phylogenetic
tree, under budding speciation, typically lacks information about which of the two descendent
branches is the new species after a speciation event (but see Aze *et al.* 2011). Thus, in the
absence of extinction, the phylogenetic age of one sister species will equal its species age
while the other will be older but without the possibility to identify which one is which.
Anagenetic speciation is not associated with a branching event but can be used to describe
different species or morphospecies (Emerson & Patiño, 2018) delimited by substantial
phenotypic change occurring along a lineage (Roopnarine et al., 1999) and will cause a higher
phylogenetic age than the genuine species age.

Here we use simulations to quantify the predictability of species age from
phylogenetic trees of extant taxa, under different diversification scenarios. Specifically, we
performed simulations where we know the true age of species to: 1) quantify the error in
phylogenetic ages under various scenarios combining different speciation modes with a range
of speciation and extinction rates and incomplete sampling; 2) examine whether this error
affects our ability to make qualitative decisions on the order of species ages; 3) explore
whether the signal of age-correlated extinction risk is preserved in the phylogenetic age of
species. Finally, we propose a new method to estimate species age more accurately under the
assumption of bifurcating speciation and complete extant species sampling, which are the
most widely used assumptions in eco-evolutionary studies (e.g., Alzate et al., 2023;
Tanentzap et al. 2020), and assess its ability to improve our interpretation of age-dependent
extinction risks.

Methods
Simulating species ages
We generated complete phylogenies of extant and extinct species under a stochastic
birth-death process using the package TreeSim 2.4 (Stadler, 2010) for the R 4.3.0 statistical
programing environment (R Core Team 2023). Then we mapped species on the complete
phylogenies using the R package FossilSim 2.3.1 (Barido-Sottani et al., 2019) under different

speciation modes, thus assigning species labels across all branches of the tree. We used the

labels assigned to terminal extant taxa to determine the true species ages. We then dropped

all extinct species from the tree and obtained the length of terminal branches, to quantify the

phylogenetic age of extant species. Finally, we rescaled all phylogenetic trees to a root age

of one, which ensures that the absolute errors in species ages are comparable in plots, and

compared the relative true and phylogenetic ages among different simulation scenarios.

Error in equating phylogenetic and species age

To explore whether there is a consistent over- or underestimation of species ages and

to quantify error in approximating species ages with phylogenetic ages, we simulated a range

of datasets with different speciation modes and diversification rates. First, we simulated 3

sets of 100 phylogenetic trees with 100 extant species based on speciation rates equal to 0.1,

0.5, and 1, combined with 100 extinction rates ranging from 0 to 0.99 in equal increments

(Beaulieu & O'Meara, 2016). Second, on each of these phylogenies we mapped species

according to different scenarios of speciation: (1) budding speciation, (2) bifurcating

speciation, (3) a combination of budding speciation and anagenetic speciation with the rate

of anagenesis set to half of the speciation rate, and (4) bifurcating speciation combined with

anagenetic speciation with the rate of anagenesis set to half of the speciation rate.

Across all trees, we obtained in total 120,000 extant species, 30,000 for each speciation

scenario. For each speciation mode and extinction fraction rate (defined as death/birth

Beaulieu & O'Meara, 2016), we calculated the mean absolute percentage error (MAPE)

across all species for each tree as measure of the deviation between the phylogenetic ages

from the true age:

$$\mathrm{MAPE} = \frac{100}{n} \times \sum_{i=1}^{n} \frac{|\nu_i - s_i|}{s_i}$$

,

where s is the true species age, v is the phylogenetic age, and n is the number of tips in the

tree (pruned of the extinct species).

Next, for each speciation mode, we plotted the MAPE against the simulated extinction

fraction.

Impact of age error in comparing species ages

To explore whether the error introduced by approximating species age with

phylogenetic age impacts our ability to make qualitative judgements, such as which of two

extant species is the younger one, we simulated 1,000 phylogenetic trees with values of

extinction fractions (extinction rate divided by speciation rate) of 0.9, 0.5, and 0, combined

with a fixed speciation rate of 1 (3000 trees). Second, on each of these phylogenies, we

mapped species according to budding and bifurcating speciation. Thus, we simulated 300,000

extant species for each speciation mode. Next, we calculated the proportion of cases where

the younger of two species, according to its phylogenetic age, is, in fact, the older one given

the true age of the two species. We performed this comparison from the perspective of an

empirical researcher that can only obtain the phylogenetic ages. We made two types of

comparisons for each phylogeny: (1) between the youngest and oldest species in the

phylogeny, and (2) between two randomly selected species.

Error in the phylogenetic age due to uniform incomplete sampling

We used the intermediate extinction scenario with an extinction fraction of 0.5 to

explore the error in equating phylogenetic and true age that is introduced by uniform

incomplete sampling under the scenarios of bifurcating and budding speciation. We specified

a fully sampled phylogeny, with 25% or 50% incomplete sampling. We simulated trees

conditioning on the total tip number, such that they included 100 sampled tips, i.e., after

dropping the unsampled ones (134 or 200 total tips). We calculated the MAPE for each tree

and compared the incomplete sampling scenarios for each speciation mode.

A probabilistic method to infer species age

Under the assumption of bifurcating speciation and complete sampling, the

phylogenetic age represents the upper boundary of plausible species ages and corresponds to

the true age in the absence of extinction. However, the true age could be younger if extinction

led to the disappearance of recent cladogenetic events from the phylogeny of extant species.

Given a phylogenic age v_i the probability that the true species age s_i is exactly v_i is conditional

on no other speciation or extinction event having occurred between v_i and the present. We

approximate the probability of no speciation or extinction for an arbitrary small time bin t

based on the probability that a lineage results in a single descendant, which is (Kendall, 1946):

$$p_1(t, \lambda, \mu) = \frac{(\lambda - \mu)^2 e^{-(\lambda - \mu)t}}{\left(\lambda - \mu \times e^{-(\lambda - \mu)t}\right)^2}$$

213 based on a birth-death process with time-homogenous speciation rate λ and extinction rate μ.

214 The probability that no event occurs over a time window v, *i.e.*, until the age of the observed

215 node i, is approximated as:

$$p_{\text{no event}}(\nu, \lambda, \mu) = p_1(t, \lambda, \mu)^{\nu/t}$$

217 In the absence of extinction ($\mu = 0$), the probability of the true species age s_i to be

218 equal to v_i is 1, because any speciation event following the node v_i would be observed in the

219 tree of extant species. Thus, we calculate the normalized probability of the speciation event

220 to occur at time v_i as:

$$P(s_i = \nu_i) = \frac{p_{\text{no event}}(\nu_i, \lambda, \mu)}{p_{\text{no event}}(\nu_i, \lambda, 0)}$$

221 (1)

222 We then compute the probability of a speciation time for any given time τ as:

$$P(s_i = \tau) = \frac{p_{\text{no event}}(\tau, \lambda, \mu)}{\int_{v_i}^0 p_{\text{no event}}(u, \lambda, \mu) du} \left(1 - P(s_i = \nu_i)\right) \text{ with } \tau \in \{\nu_i, 0\}$$

, (2)

where the first term quantifies the probability of the species age along a vector of

times, while the second term normalizes it after accounting for the probability that speciation

occurred exactly at the node. We use equations (1) and (2) to approximate a density

describing the probability of a species origination at the observed phylogenetic age and along

the branch connecting the node with the tip. As point estimates of the species age, we tested

whether the mean or the median of the distribution better approximate the true species age

compared to the phylogenetic age.

To evaluate the accuracy of our probabilistic species age estimator, we simulated

1,000 phylogenetic trees with extinction rates of 0.25, 0.15, and 0.05, and a fixed speciation

rate of 0.3. Using these phylogenies as inputs, we mapped species on each tree based on a

bifurcating speciation process. Then, we calculated the MAPE for each tree as a measure of

the deviation between the function's estimated and the phylogenetic ages from the true age.

Simulation of age-dependent species extinction risks

To evaluate the impact of the erroneous estimation of species age due to the extinction

shortfall on macroevolutionary analyses, we explored whether the assumed relationship

between species age and contemporary extinction risk (e.g., Johnson et al. 2002; Tanentzap

et al. 2020; Verde Arregoitia et al. 2013) is preserved in the phylogenetic ages. For this, we

binned the same number of extant species according to their age in five categories reflecting

the increase in extinction risk with age encapsulated by the IUCN categories: Least Concern

(LC), Near Threatened (NT), Vulnerable (VU), Endangered (EN), and Critically Endangered

(CR; International Union for the Conservation of Nature 2016). With this, we generated a

positive effect with older species being at higher extinction risk regarding the IUCN

categories, assuming bifurcating speciation.

Then, we quantified the share of the 1000 datasets where the order of the mean age

per IUCN category did not match with the simulated monotonic increase when utilizing (a)

phylogenetic ages, and (b) the probabilistic species age estimator. We also evaluated the

effect of nonrandom incomplete sampling (older species were less prone to be sampled than

younger ones) on the evaluation of species age-correlated extinction risks.

Results

Error in equating phylogenetic and species age

Under the assumption of bifurcating speciation and with no extinction events,

phylogenetic ages matched the true age of extant species (Fig. 3). At low extinction fractions

(< 0.25), 96% of the phylogenetic age estimations were congruent with the true age. At higher

extinction fractions (> 0.75), this was also the case for most species (73%). However, age

overestimation increased with extinction fraction and in some cases the phylogenetic age

erroneously suggested that the species is as old as the root age. While under bifurcating

speciation, the phylogenetic age never underestimated the true species age, both over- and

underestimation occurred in the case of budding speciation. Moreover, the proportion of

cases where the phylogenetic ages equal the species age was lower than in the bifurcating

scenario (Fig. 3). Overestimated ages were more frequent with high extinction while

underestimations occurred with low extinction, but in principle both happened under the

complete range of extinction rates (Fig. 3). Even at low extinction fractions, ~50% of
phylogenetic ages did not match the true ages.

In datasets simulated under a mixture of anagenetic and bifurcating speciation,
phylogenetic ages deviated more strongly from the true ages than under a pure bifurcating
process, given that anagenetic events are unobserved in the phylogeny (Fig. SM1). With a
low extinction fraction, phylogenetic ages were congruent with the true species ages in 78%
of the cases instead of 96%, and with high extinction the share decreases from 73% to 62%.
Datasets with mixed anagenetic and budding speciation, phylogenetic ages also deviated
more that under a pure budding process; with a low extinction fraction, phylogenetic ages
were congruent 40% instead of 50% of the time, and with high extinction the accuracy
decreased to 34%.

While a budding speciation mode led to a higher baseline error than bifurcation, the
latter showed a stronger increase with extinction (Fig. 4). Under both modes of speciation,
speciation rates did not have a substantial impact on error in age. For strictly bifurcating
speciation, there was no error in the absence of extinction, but the MAPE increased to up to
150% with extinction fractions exceeding 0.75. In contrast, under budding speciation the
MAPE was around 25% in the absence of extinction, increasing to 30-120% with extinction
fractions exceeding 0.75. In datasets that included anagenetic speciation, the MAPE reached
as high as 500% in some simulations (Fig. SM2).

Impact of age error on comparing species ages

For the combination of strictly bifurcating speciation and all extinction scenarios,
selecting the phylogenetically youngest and oldest species never resulted in a case where the
presumed older species has been in fact the younger of the two species according to their
simulated age (Fig. SM3a). Thus, for this speciation mode, the risk of a qualitative error when
comparing species at the extremes of the age range is minimal. However, the age ranking of
two randomly selected species was found to be incorrect in 6% and 8%, for intermediate and
high extinction, respectively (Fig. SM4a). Thus, qualitative errors in comparing species ages
are non-negligible under the assumption of bifurcating speciation.

In contrast, for budding speciation, the age rank of the oldest and youngest species was
erroneously determined in 2.2% of the simulations in the absence of extinction, increasing to
7.5% and 12.2% for intermediate and high extinction, respectively (Fig. SM3b). Thus, under
the assumption of budding speciation, there is a substantial risk of mistaking the oldest and
youngest species in the clade. The error in age ranking of two randomly selected species was
even higher, exceeding 25%, irrespective of the extinction level (Fig. SM4b).

Error on equating phylogenetic and species age given uniform incomplete sampling

Under bifurcating speciation, the MAPE for completely sampled trees was $25 \pm 20\%$;
for trees missing 25% of the extant species the error increased by 15 orders of magnitude;
and for trees missing 50 % of the extant species the error increased by 85 orders of magnitude
(Fig. 5a). Under budding speciation, the MAPE for completely sampled trees was 42 ± 15 %;
for trees missing 25 % of the extant species the error increased by 5 orders of magnitude; and
for trees missing 50% of the extant species the error increased by 22 orders of magnitude
(Fig. 5b).

Probabilistic species age estimation

With increasing extinction, our probabilistic estimation of species ages resulted in an
increasingly lower error compared with the phylogenetic age (Figs. 6, SM5). Under low
extinction the MAPE was slightly worse compared to the use of phylogenetic ages (ΔMAPE
$= 2.83 \pm 3.9$ % when using the mean of the estimated ages and 2.1 ± 1.3 % for the median
across estimates). While with increasing extinction rates the MAPE of the phylogenetic ages
increased up to 25% (Fig. SM5), the probabilistic estimation strongly reduced the error
compared with phylogenetic age (ΔMAPE = -27 ± 26 % for the mean estimated ages and -
32 ± 28 % for the median across estimates; Figs. 6, SM5).

Detecting age-dependent extinction risk

The use of phylogenetic age as an approximation of species age led to error rates of
1.3, 7.2, and 18.6 % in detecting the correlation between species ages and extinction risk for
scenarios with low, intermediate, and high extinction rates, respectively (Fig. 7). Thus, even
under intermediate extinction the true relationship between age and extinction risk was

wrongly estimated in a significant fraction of the simulations, and higher extinction rates led

to a further substantial drop in the reliability of this approach. In contrast, estimating species

ages based on our probabilistic method led to much lower error rates (3 to 4-fold) that

dropped to 1.3% and 4% with intermediate and high extinction, respectively. Under

incomplete taxon sampling, in which the sampling probability was negatively correlated with

species age, the error rates increased to 12.8%, and 49.5% for scenarios with 25%, and 50%

of missing extant species, respectively (Fig. SM6).

Discussion

339   The use of branch lengths of phylogenetic trees as an approximation of species ages

is becoming central to an increasing number of studies that use them to evaluate the

relationship among macro-ecological and evolutionary patterns and current extinction risks

(Pie & Caron, 2023; Sonne et al., 2022; Tanentzap et al., 2020; Verde Arregoitia et al., 2013).

Here we showed that this approximation leads to substantial errors and that its accuracy is

hampered by three shortfalls: unobserved extinction events, unknown speciation mode, and

incomplete sampling of extant species. The only instance in which phylogenetic ages

correctly predict species age is under the assumption of a bifurcating speciation process in

the absence of extinction and with all living species included in the phylogenetic tree. While

the prevalence of true speciation modes remains difficult to estimate (Bapst & Hopkins, 2017;

Silvestro et al., 2018; Wagner et al., 1995), the fossil record unequivocally shows that

extinction occurs across all clades in the tree of life (Bambach 2006; Benton 2023; Pimm et

al., 2014), and there is substantial evidence that many living species remain unknown to

science and are therefore absent from empirical phylogenetic trees (Blackwell 2001; Mora et

al., 2011). Thus, the scenario under which phylogenetic age correctly predicts species age is

very unlikely.

355   Some authors acknowledged the problems associated with measuring species age

from phylogenies (Swenson, 2019), and have proposed approaches to account for them. For

example, Sonne *et al.* (2022) determined young and old Andean hummingbirds by assessing

the sensitivity of their results to incomplete taxon sampling by generating 1000 trees with

randomly missing species. Pie & Caron (2023) accounted for taxonomic incompleteness by

pruning an additional 1 – 5% of species and evaluated if their conclusions changed and found

that they did not. Yet, the magnitude of the error associated with the direct use of the length

of phylogenetic branches as estimators of species ages remains under-appreciated, as shown

by the many studies implementing this approach.

We showed that the largest error in estimating species ages from phylogenetic trees

is linked with incomplete sampling of extant species. This is a problem that in principle can

be solved by extending the scope of the sampling to include all species in the phylogenetic

inference. Yet, despite the advances in the scalability of DNA sequencing, this remains

impractical for large clades, including some of the best sampled ones such as vertebrate

groups, in which many species still lack genetic data (Jetz & Pyron 2018; Tonini et al., 2016;

Upham et al., 2019). In addition, a substantial proportion of species might be missing from

the phylogenetic trees because they are still unknown to science, a problem often identified

as the Linnean shortfall (Diniz Filho et al., 2023; Hortal et al., 2015). The magnitude of the

Linnean shortfall is unknown, but available estimates show that it affects some clades

significantly more than others (Moura & Jetz 2021; Ondo et al., 2023), with the diversity of

highly diverse groups, such as insects and fungi, likely to be highly underestimated

(Blackwell 2001; Mora et al., 2011).

Our simulations showed that under some scenarios the accurate estimation of species

ages is essentially impossible from phylogenetic trees using current approaches. Under the

assumption of budding speciation, the error is high even without extinction and with

complete sampling, because phylogenetic trees are agnostic about parent and descendant

species following a branching event (Fig. 2, 4). Phylogenetic ages are, by construction,

identical for sister species, which is necessarily wrong within a budding speciation scenario.

Similarly, anagenetic speciation also leads to high error, which did not vary substantially

with extinction. However, anagenetic speciation might be impossible to quantify, except

perhaps in high resolution fossil time series (Aze et al., 2011), resulting in a general debate

on the use of the term anagenesis in evolutionary biology (Vaux et al., 2015) and

biogeography (Emerson & Patiño, 2018; Meiri et al., 2018). Thus, species age is

unidentifiable under the assumption of speciation modes that deviate from a strictly

bifurcating scenario.

The lowest error in species age estimation was observed under scenarios of
bifurcating speciation. This is the implicit assumption of most studies using approximations
of species ages (Alzate et al., 2023; Freer et al., 2022; Kennedy et al., 2022), even though it
is at odds with the assumption of all birth-death models used in the molecular clock analyses
that estimate the phylogenetic trees in the first place (Gernhard, 2008; Nee et al., 1994;
Stadler, 2013). Despite the lower error, our simulations showed that both extinction and
missing lineages can lead to a substantial decrease in accuracy (Fig. 4-5, SM3-4) that can
even lead to qualitative misinterpretations of general patterns such as age-dependent
extinction risks (Fig. 7). Given the large inaccuracy in phylogenetic age, the question is
whether this affects the inferences made from the relationship between species age and eco-
evolutionary variables, such as extinction risk, range size, or environmental variables (Gaston
& Blackburn 1997, Johnson *et al.* 2002, Tanentzap *et al.* 2015, Pie & Caron 2023).

We found our probabilistic approach to efficiently reduce the biases associated with
extinction. It substantially improved the accuracy of the estimation of species ages, leading
to much lower error rates even under scenarios of high extinction fractions. The performance
of our estimator is however contingent on the ability of birth-death models to correctly
estimate speciation and extinction rates from completely sampled phylogenies of extant
species. The accuracy of these methods has been shown to be high under several simulation
settings (Silvestro et al., 2011; Stadler 2011), but their accuracy has been questioned for
empirical datasets and under complex models with rate variation (Louca & Pennell 2020;
Rabosky 2010). One commonly observed pattern is the estimation of 0-extinction rate from
empirical phylogenies (Louca & Pennell 2021), which would make corrected ages identical
to the phylogenetic ages. Yet, the fossil record decisively shows that extinction and speciation
rates vary within the same order of magnitude across virtually all clades (Parry 2021). The
robustness of estimated extinction rates can be increased through the use of fossil data in the
analyses (Heath et al., 2014; Silvestro et al., 2018; Warnock et al., 2020). Alternatively, our
probabilistic approach could be applied across a range of plausible values of speciation and
extinction rates as a way to evaluate the robustness of conclusions drawn from the patterns
of estimated species ages.

Conclusion

421
This study provides a quantification of the deviations between true species age and
phylogenetic age due to incomplete taxon sampling, extinction, and unknown speciation
modes. We found that phylogenetic age is a good proxy of species age only in a rather
unlikely case in which 1) all species in a clade are known to science and included in the
phylogenetic tree, 2) speciation occurs as a strictly bifurcating process, and 3) there is no or
little extinction. Using simulations, we identified that incomplete taxon sampling and
budding and anagenetic speciation modes cause the highest mismatch between phylogenetic
age and true species age and cannot be accounted for in the absence of additional information,
for instance, derived from fossil data. In contrast, under a scenario of bifurcating speciation
and fully sampled phylogeny, we proposed a probabilistic approach based on the properties
of the birth-death process that can drastically improve the accuracy of the estimated species
ages by reducing the bias linked with extinction. We note that, even in this case, the
robustness of such estimates will be contingent on the accuracy of the estimated speciation
and extinction rates and of course of the underlying phylogenetic tree and dating of the
branching times. In the light of our findings, we caution against a direct use of branch lengths
as proxies for species ages. We recommend the application of our probabilistic approach to
correct species ages when assumptions are met. We hope our results will stimulate a
discussion about the use of phylogenetic trees in inferring species age and lead to a critical
evaluation of the robustness of inferences linking species age with traits, ecological variables,
and extinction risks.

Acknowledgements

CC received a Ph.D. scholarship from CAPES (88887.814725/2023-00) and an abroad
internship CAPES-Print scholarship (88887.682496/2022-00). D.S. and TH received funding
from the Swiss National Science Foundation (PCEFP3_187012). D.S. received funding the
Swedish Research Council (VR: 2019-04739), and the Swedish Foundation for Strategic
Environmental Research MISTRA within the framework of the research programme
BIOPATH (F 2022/1448). JDC was supported by the Swiss National Science Foundation
grant TMPFP2_209818.

Fig. 1. Discrepancy between species ages. Estimations based on the length of the terminal branch in a time-calibrated phylogeny (red) and the first appearance in the fossil record (green). Shark data (*Trianenodon obesus* and *Carcharhinus obscurus*) obtained from Brée et al. (2022). Mammals' phylogenetic data (except *Homo sapiens*) obtained from Upham et al. (2019). Mammals' fossil data (except *Homo sapiens*) obtained from Silvestro *et al.* (2018). *Homo sapiens* fossil and phylogenetic data obtained from Callaway (2017) and Rivas-Gonzáles et al. (2023), respectively.

Fig. 2. Impact of speciation mode and extinction on species age. For the same hypothetical time-calibrated phylogeny of extant species, the continuation of the same color indicates the same species, solid and dashed lines represent extant and extinct species, respectively, under different speciation modes and extinction scenarios. Numbers display the resulting age of the respective species in million years (myr).

478   **Fig. 3.** True age versus phylogenetic age at low and high extinction fraction for bifurcating

479   and budding speciation. Each point represents a species and both ages, true and phylogenetic,

480   are scaled to the root age of the correspondent phylogenetic tree.

494   **Fig. 4.** Error in equating phylogenetic age with true age. The error was quantified as mean

495   absolute percentage error (MAPE) between the true and phylogenetic ages across all species

496   for each tree simulated under bifurcating and budding speciation. Each dot represents one

497   replicate of the 300 trees for each speciation mode using different rates of speciation and

498   extinction fraction.

Fig. 5. Effect of incomplete taxon sampling on error in species ages. Error in equating the phylogenetic age with true species age for the fully sampled phylogeny, and 25% and 50% of missing extant species, in an intermediate extinction scenario. The error was quantified as mean absolute percentage error (MAPE) between the true phylogenetic ages across 100 species for each of 1000 trees for each missing species scenario simulated under bifurcating and budding speciation.

Fig. 6. Performance of the probabilistic age estimator. ΔMAPE for the three extinction scenarios (low, intermediate, and high; from left to right) was quantified as the difference of the mean absolute percentage error (MAPE) of the probabilistic estimator point estimates (mean and median) and the MAPE of the phylogenetic age. The MAPE was quantified as the difference between the true and point estimates (mean and median) of phylogenetic ages for 100 species across 1000 trees for each extinction scenario simulated under bifurcating speciation. The red dashed line represents no difference between the compared MAPEs, negative ΔMAPE values indicate an improvement in the accuracy of the probabilistic estimator over the phylogenetic age.

Fig. 7. Power to recover an age extinction-risk relationship. Simulated species ages under three extinction scenarios and assuming bifurcating speciation were binned into conservation status categories, which represents an increase in extinction risk by age (LC = Least Concern; NT = Near Threatened; VU = Vulnerable; EN = Endangered; CR = Critically Endangered). We used the phylogenetic age and the mean age obtained from our probabilistic corrective function to calculate the mean age per conservation status category and assess if every mean age increases in comparison with the previous category with lower extinction risk. The error rate is the percentage across all 1000 phylogenies where the relationship between the mean ages and the conservation status categories is not always increasing (shown by grey lines).

References

Alroy, J., Marshall, C. R., Bambach, R. K., Bezusko, K., Foote, M., Fürsich, et al., (2001).
  Effects of sampling standardization on estimates of phanerozoic marine
  diversification. *Proceedings of the National Academy of Sciences of the United States
  of America*, *98*(11), 6261–6266. doi: 10.1073/pnas.111144698

Alzate, A., Rozzi, R., Velasco, J. A., Robertson, D. R., Zizka, A., Tobias, J. A., ... &
  Onstein, R. E. (2023). The evolutionary age-range size relationship is modulated by
  insularity and dispersal in plants and animals. *bioRxiv*, 2023-11. doi:
  10.1101/2023.11.11.566377

Aze, T., Ezard, T. H. G., Purvis, A., Coxall, H. K., Stewart, D. R. M., Wade, B. S., &
  Pearson, P. N. (2011). A phylogeny of Cenozoic macroperforate planktonic
  foraminifera from fossil data. *Biological Reviews*, *86*(4), 900–927. doi:
  10.1111/j.1469-185X.2011.00178.x

Balmford, A. (1996). Extinction filters and current resilience: The significance of past
  selection pressures for conservation biology. *Trends in Ecology and Evolution*. doi:
  10.1016/0169-5347(96)10026-4

Bapst, D. W., & Hopkins, M. J. (2017). Comparing cal3 and other a posteriori time-scaling
  approaches in a case study with the pterocephaliid trilobites. *Paleobiology, 43*(1), 49-
  67. doi: 10.1017/pab.2016.34

Barido-Sottani, J., Pett, W., O'Reilly, J. E., & Warnock, R. C. M. (2019). FossilSim: An r
  package for simulating fossil occurrence data under mechanistic models of
  preservation and recovery. *Methods in Ecology and Evolution*, *10*(6), 835–840. doi:
  10.1111/2041-210X.13170

Barraclough, T. G., & Vogler, A. P. (2000). Detecting the geographical pattern of
  speciation from species-level phylogenies. *The American Naturalist*, *155*(4), 419–434.
  doi: 10.1086/303332

Barraclough, T. G., Vogler, A. P., & Harvey, P. H. (1998). Revealing the factors that
  promote speciation. *Philosophical Transactions of the Royal Society B: Biological
  Sciences*, *353*(1366), 241–249. doi: 10.1098/rstb.1998.0206

Bambach, R. K. (2006). Phanerozoic biodiversity mass extinctions. *Annu. Rev. Earth*

Planet. Sci., 34, 127-155. doi: 10.1146/annurev.earth.33.092203.122654

Baum, D. A., Smith, S. D. W., & Donovan, S. S. S. (2005). The tree-thinking challenge.

Science, 310(5750), 979–980. doi: 10.1126/science.1117727

Beaulieu, J. M., & O'Meara, B. C. (2016). Detecting hidden diversification shifts in models

of trait-dependent speciation and extinction. *Systematic Biology, 65*(4), 583–601. doi:

10.1093/sysbio/syw022

Benton, M. J. (2013). Origins of biodiversity. *Palaeontology*. doi: 10.1111/pala.12012

Benton, M. J. (2016). Origins of Biodiversity. *PLoS Biology, 14*(11), 1–7. doi:

10.1371/journal.pbio.2000724

Benton, M. J. (2023). Extinctions: How Life Survives, Adapts and Evolves. Thames &

Hudson.

Blackwell, M. (2011). The Fungi: 1, 2, 3… 5.1 million species?. *American journal of*

botany, 98(3), 426-438. doi: 10.3732/ajb.1000298

Brée, B., Condamine, F. L., & Guinot, G. (2022). Combining palaeontological and

neontological data shows a delayed diversification burst of carcharhiniform sharks

likely mediated by environmental change. *Scientific Reports, 12*(1), 21906. doi:

10.1038/s41598-022-26010-7

Caetano, D. S., & Quental, T. B. (2022). How important is budding speciaiton for

comparative studies? *BioRxiv*. doi: 10.1101/2022.05.24.493296

Callaway, E. (2017). Oldest Homo sapiens fossil claim rewrites our species' history.

Nature, 546, 289-293. doi: 10.1038/nature.2017.22114

Carrillo, J. D., Faurby, S., Silvestro, D., Zizka, A., Jaramillo, C., Bacon, C. D., & Antonelli,

A. (2020). Disproportionate extinction of South American mammals drove the

asymmetry of the Great American Biotic Interchange. *Proceedings of the National*

Academy of Sciences of the United States of America, 117(42), 26281–26287. doi:

10.1073/pnas.2009397117

Carrillo, J. D., Forasiepi, A., Jaramillo, C., Sánchez-villagra, M. R., & Richardson, J. E.

(2015). *Neotropical mammal diversity and the Great American Biotic Interchange :*

spatial and temporal variation in South America ' s fossil record. 5(January), 1–11.

doi: 10.3389/fgene.2014.00451

Chang, J., Rabosky, D. L., & Alfaro, M. E. (2020). Estimating Diversification Rates on

Incompletely Sampled Phylogenies: Theoretical Concerns and Practical Solutions.

Systematic Biology, *69*(3), 602–611. doi: 10.1093/sysbio/syz081

Davies, T. J., Smith, G. F., Bellstedt, D. U., Boatwright, J. S., Bytebier, B., Cowling, R. M.,

et al., (2011). Extinction risk and diversification are linked in a plant biodiversity

hotspot. *PLoS Biology*. 9(5), e1000620. doi: 10.1371/journal.pbio.1000620

Diniz Filho, J. A. F., Jardim, L., Guedes, J. J., Meyer, L., Stropp, J., Frateles, L. E. F., ... &

Hortal, J. (2023). Macroecological links between the Linnean, Wallacean, and

Darwinian shortfalls. *Frontiers of Biogeography*, 15(2). doi: 10.21425/F5FBG59566

Eldredge, N., Thompson, J. N., Brakefield, P. M., Gavrilets, S., Jablonski, D., Jackson, J.

B. C., et al., (2005). The dynamics of evolutionary stasis. *Paleobiology* 31(S2), 133-

145. doi: 10.1666/0094-8373(2005)031

Emerson, B. C., & Patiño, J. (2018). Anagenesis, Cladogenesis, and Speciation on Islands.

Trends in Ecology and Evolution, *33*(7), 488–491. doi: 10.1016/j.tree.2018.04.006

Fitzpatrick, B. M., & Turelli, M. (2006). The Geography of Mammalian Speciation: Mixed

Signals From Phylogenies and Range Maps. *Evolution*, *60*(3), 601–615. doi:

10.1111/j.0014-3820.2006.tb01140.x

Fjeldså, J., Bowie, R. C. K., & Rahbek, C. (2012). The role of mountain ranges in the

diversification of birds. *Annual Review of Ecology, Evolution, and Systematics*, *43*,

249–265. doi: 10.1146/annurev-ecolsys-102710-145113

Foote, M. (1996). *On the Probability of Ancestors in the Fossil Record*. 22(2), 141–151.

doi: 10.1017/S0094837300016146

Foote, M., & Raup, D. M. (1996). *Fossil Preservation and the Stratigraphic Ranges of*

Taxa. 22(2), 121–140. doi: 10.1017/S0094837300016134

Freer, J. J., Collins, R. A., Tarling, G. A., Collins, M. A., Partridge, J. C., & Genner, M. J.

(2022). *Global phylogeography of hyperdiverse lanternfishes indicates sympatric*

speciation in the deep sea. Global Ecology and Biogeography, 31(11), 2353-2367.

doi: 10.1111/geb.13586

Gaston, K. J., & Blackburn, T. M. (1997). Evolutionary age and risk of extinction in the

global avifauna. *Evolutionary Ecology,* 11, 557–565. doi: 10.1007/s10682-997-1511-4

Gernhard, T. (2008). The conditioned reconstructed process. *Journal of theoretical biology,*

253(4), 769-778. doi: 10.1016/j.jtbi.2008.04.005

Harvey, P. H., May, R. M., & Nee, S. (1994). Phylogenies without fossils. *Evolution*, *48*(3), 523–529. doi: 10.1111/j.1558-5646.1994.tb01341.x

Heath, T. A., Hedtke, S. M., & Hillis, D. M. (2008). Taxon sampling and the accuracy of phylogenetic analyses. *Journal of Systematics and Evolution*, *46*(3), 239–257. doi: 10.3724/SP.J.1002.2008.08016

Heath, T. A., Huelsenbeck, J. P., & Stadler, T. (2014). The fossilized birth–death process for coherent calibration of divergence-time estimates. *Proceedings of the National Academy of Sciences, 111*(29), E2957-E2966. doi: 10.1073/pnas.1319091111

Hortal, J., de Bello, F., Diniz-Filho, J. A. F., Lewinsohn, T. M., Lobo, J. M., & Ladle, R. J. (2015). Seven shortfalls that beset large-scale knowledge of biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 46, 523-549. doi: 10.1146/annurev-ecolsys-112414-054400

IUCN. (2016). IUCN Red List of Threatened Species. Version 2016-2. *Fourth Quarter*.

Jetz, W., & Pyron, R. A. (2018). The interplay of past diversification and evolutionary isolation with present imperilment across the amphibian tree of life. *Nature ecology & evolution, 2*(5), 850-858. doi: 10.1038/s41559-018-0515-5

Johnson, C. N., Delean, S., & Balmford, A. (2002). *Phylogeny and the selectivity of extinction in Australian marsupials*. 135–142. doi: 10.1017/S1367943002002196

Kendall, M. G. (1946). *The advanced theory of statistics* (2nd ed.). Charles Griffin adn Co., Ltd., London.

Kennedy, J. D., Marki, P. Z., Reeve, A. H., Blom, M. P., Prawiradilaga, D. M., Haryoko, T., ... & Jønsson, K. A. (2022). *Diversification and community assembly of the world's largest tropical island. Global Ecology and Biogeography, 31*(6), 1078-1089. doi: 10.1111/geb.13484

López-Martínez, A. M., Schonenberger, J., von Balthazar, M., González-Martínez, C. A., Ramírez-Barahona, S., Sauquet, H., & Magallón, S. (2023). Integrating Fossil Flowers into the Angiosperm Phylogeny Using Molecular and Morphological Evidence. *Systematic Biology*, *syad017*. doi: 10.1093/sysbio/syad017

Losos, J. B., & Glor, R. E. (2003). Phylogenetic comparative methods and the geography of speciation. *Trends in Ecology and Evolution*, *18*(5), 220–227. doi: 10.1016/S0169-5347(03)00037-5

Louca, S., & Pennell, M. W. (2020). Extant timetrees are consistent with a myriad of
diversification histories. *Nature, 580*(7804), 502-505. doi: 10.1038/s41586-020-2176-
1

Louca, S., & Pennell, M. W. (2021). Why extinction estimates from extant phylogenies are
so often zero. *Current Biology, 31*(14), 3168-3173. doi: 10.1016/j.cub.2021.04.066

Meier, R., & Willmann, R. (2000). A defense of the Hennigian Species Concept. In *Species
concepts and phylogenetic theory: a debate* (p. 167).

Meiri, S., Raia, P., & Santos, A. M. (2018). Anagenesis and cladogenesis are useful island
biogeography terms. *Trends in Ecology & Evolution, 33*(12), 895-896. doi:
10.1016/j.tree.2018.09.005

Mora, C., Tittensor, D. P., Adl, S., Simpson, A. G., & Worm, B. (2011). How many species
are there on Earth and in the ocean?. *PLoS biology, 9*(8), e1001127. doi:
10.1371/journal.pbio.1001127

Moura, M. R., & Jetz, W. (2021). Shortfalls and opportunities in terrestrial vertebrate
species discovery. Nature Ecology & Evolution, 5(5), 631-639. doi: 10.1038/s41559-
021-01411-5

Mynard, P., Algar, A., Lancaster, L., Bocedi, G., Fahri, F., Gubry-Rangin, C., et al., (2023).
Impact of Phylogenetic Tree Completeness and Misspecification of Sampling
Fractions on Trait Dependent Diversification Models. *Systematic Biology*, *72*(1), 106–
119. doi: 10.1093/sysbio/syad001

Nee, S., May, R. M., & Harvey, P. H. (1994). The reconstructed evolutionary process.
*Philosophical Transactions of the Royal Society of London. Series B: Biological
Sciences, 344*(1309), 305-311. doi: 10.1098/rstb.1994.0068

Nee, S., & May, R. M. (1997). Extinction and the Loss of Evolutionary History. *Science*,
278(5338), 692–694. doi: 10.1126/science.278.5338.692

Ondo, I., Dhanjal-Adams, K., Pironon, S., Silvestro, D., Deklerck, V., Grace, O., ... &
Antonelli, A. (2023). Plant diversity darkspots for global collection priorities. *bioRxiv*,
2023-09. doi: 10.1101/2023.09.12.557387

Parry, L. A. (2021). Evolution: No extinction? No way!. *Current Biology, 31*(14), R907-
R909. doi: 10.1016/j.cub.2021.05.044

Pearson, P. N. (1995). Investigating age-dependency of species extinction rates using

dynamic survivorship analysis. *Historical Biology*, 10(2), 119–136. doi: 10.1080/10292389509380516

Pie, M. R., & Caron, F. (2023). Substantial variation in species ages among vertebrate clades. *BioRxiv*, *06*. doi: 10.1101/2023.06.08.544238

Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N., … Sexton, J. O. (2014). The biodiversity of species and their rates of extinction, distribution, and protection. *Science*, *344*(6187). doi: 10.1126/science.1246752

R Core Team. (2019). R: A language and environment for statistical computing. Vienna, Austria: *R Foundation for Statistical Computing*. Retrieved from https://www.r-project.org/

Rabosky, D. L. (2010). Extinction rates should not be estimated from molecular phylogenies. *Evolution*, 64(6), 1816–1824. doi: 10.1111/j.1558-5646.2009.00926.x

Rivas-Gonzáles, I., Rousselle, M., Li, F., Zhou, L., Dutheil, J., Munch, K., et al., (2023). Pervasive incomplete lineage sorting illuminates speciation and selection in primates. *Science*, *380*(6648), eabn4409. doi: 10.1126/science.abn4409

Roopnarine, P. D., Byars, G., Fitzgerald, P., Paleobiology, S., & Winter, N. (1999). Anagenetic evolution, stratophenetic patterns, and random walk models. *Paleobiology, 25*(1), 41–57. doi: 10.1666/0094-8373(1999)0252.3.CO;2

Rosenblum, E. B., Sarver, B. A. J., Brown, J. W., Des Roches, S., Hardwick, K. M., Hether, T. D., et al., (2012). Goldilocks Meets Santa Rosalia: An Ephemeral Speciation Model Explains Patterns of Diversification Across Time Scales. *Evolutionary Biology*, *39*(2), 255–261. doi: 10.1007/s11692-012-9171-x

Silvestro, D., Schnitzler, J., & Zizka, G. (2011). A Bayesian framework to estimate diversification rates and their variation through time and space. *BMC evolutionary biology, 11*, 1-15. doi: 10.1186/1471-2148-11-311

Silvestro, D., Salamin, N., & Schnitzler, J. (2014). PyRate: a new program to estimate speciation and extinction rates from incomplete fossil data. *Methods in Ecology and Evolution2*, *5*(10), 1126–1131. doi: 10.1111/2041-210X.12263

Silvestro, D., Castiglione, S., Mondanaro, A., Serio, C., Melchionna, M., Piras, P., et al., (2020). A 450 million years long latitudinal gradient in age-dependent extinction. *Ecology Letters*, *23*(3), 439–446. doi: 10.1111/ele.13441

Silvestro, D., Salamin, N., Antonelli, A., & Meyer, X. (2019). Improved estimation of
macroevolutionary rates from fossil data using a Bayesian framework. *Paleobiology*,
45(4), 546–570. doi: 10.1017/pab.2019.23

Silvestro, D., Warnock, R. C. M., Gavryushkina, A., & Stadler, T. (2018). Closing the gap
between palaeontological and neontological speciation and extinction rate estimates.
Nature Communications, *9*(1). doi: 10.1038/s41467-018-07622-y

Simpson, George Gaylord. (1984). *Tempo and mode in evolution*. Columbia University
Press.

Skeels, A., & Cardillo, M. (2019). Reconstructing the geography of speciation from
contemporary biodiversity data. *American Naturalist*, *193*(2), 240–255. doi:
10.1086/701125

Slater, G. J., & Harmon, L. J. (2013). Unifying fossils and phylogenies for comparative
analyses of diversification and trait evolution. *Methods in Ecology and Evolution*,
4(8), 699-702. doi: 10.1111/2041-210X.12091

Sonne, J., Dalsgaard, B., Borregaard, M. K., Kennedy, J., Fjeldså, J., & Rahbek, C. (2022).
Biodiversity cradles and museums segregating within hotspots of endemism.
Proceedings of the Royal Society B: Biological Sciences, *289*(1981), 20221102. doi:
10.1098/rspb.2022.1102

Stadler, T. (2011). Mammalian phylogeny reveals recent diversification rate shifts.
Proceedings of the National Academy of Sciences, 108(15), 6187-6192. doi:
10.1073/pnas.1016876108

Stadler, T. (2013). Recovering speciation and extinction dynamics based on phylogenies.
Journal of evolutionary biology, 26(6), 1203-1219. doi: 10.1111/jeb.12139

Stadler T (2019). TreeSim: Simulating Phylogenetic Trees. R package version
2.4, https://CRAN.R-project.org/package=TreeSim

Swenson, N. G. (2019). *Phylogenetic ecology: A history, critique, and remodeling*.
University of Chicago Press.

Tanentzap, A. J., Brandt, A. J., Smissen, R. D., Heenan, P. B., Fukami, T., & Lee, W. G.
(2015). When do plant radiations influence community assembly? The importance of
historical contingency in the race for niche space. *New Phytologist*, *207*(2), 468–479.
doi: 10.1111/nph.13362

Tanentzap, A. J., Igea, J., Johnston, M. G., & Larcombe, M. J. (2020). Does evolutionary

history correlate with contemporary extinction risk by influencing range size

dynamics? *American Naturalist*, 195(3), 569–576. doi: 10.1086/707207

Thomas, G. H., Hartmann, K., Jetz, W., Joy, J. B., Mimoto, A., & Mooers, A. O. (2013).

PASTIS: An R package to facilitate phylogenetic assembly with soft taxonomic

inferences. *Methods in Ecology and Evolution*, *4*(11), 1011–1017. doi: 10.1111/2041-

210X.12117

Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W., & Pyron, R. A. (2016). Fully-

sampled phylogenies of squamates reveal evolutionary patterns in threat status.

Biological Conservation, 204, 23-31. doi: 10.1016/j.biocon.2016.03.039

Upham, N. S., Esselstyn, J. A., & Jetz, W. (2019). Inferring the mammal tree: Species-level

sets of phylogenies for questions in ecology, evolution, and conservation. In *PLoS

Biology*, 17(12), e3000494. doi: 10.1371/journal.pbio.3000494

Van Valen, L. (1973). A new evolutionary law. *Evolutionary Theory*. 1, 1–30.

Vaux, F., Trewick, S. A., & Morgan-Richards, M. (2016). Lineages, splits and divergence

challenge whether the terms anagenesis and cladogenesis are necessary. Biological

Journal of the Linnean Society, 117(2), 165-176. doi: 10.1111/bij.12665

Verde Arregoitia, L. D., Blomberg, S. P., & Fisher, D. O. (2013). Phylogenetic correlates

of extinction risk in mammals: Species in older lineages are not at greater risk.

Proceedings of the Royal Society B: Biological Sciences 280(1765), 20131092. doi:

10.1098/rspb.2013.1092

Wagner, P. J., Erwin, D. H., & Anstey, R. L. (1995). Phylogenetic patterns as tests of

speciation models. In *New approaches to speciaiton in the fossil record* (pp. 87–122).

New York: Columbia University Press.

Warnock, R. C., Heath, T. A., & Stadler, T. (2020). Assessing the impact of incomplete

species sampling on estimates of speciation and extinction rates. *Paleobiology, 46*(2),

137-157. doi: 10.1017/pab.2020.12

790