Seabird nutrient subsidies enrich mangrove ecosystems and are exported to nearby coastal

habitats

Jennifer Appoo,[1,2,5*] Nancy Bunbury,[2,3] Sébastien Jaquemet,[1] and Nicholas A.J. Graham[4]

[1]UMR ENTROPIE, Université de La Réunion, 97744 Saint Denis Cedex 9, La Réunion, France

[2]Seychelles Islands Foundation, Victoria, Mahé, Seychelles

[3]Centre for Ecology and Conservation, University of Exeter, Cornwall Campus, Penryn, UK

[4]Lancaster Environment Centre, Lancaster University, Lancaster, UK

[5]Lead contact

*Correspondence: jennifer.appoo@univ-reunion.fr

SUMMARY

Eutrophication by human-derived nutrient enrichment is a major threat to mangroves, impacting productivity, ecological functions, resilience and ecosystem services. Natural mangrove nutrient enrichment processes, however, remain largely uninvestigated. Mobile consumers such as seabirds are important vectors of cross-ecosystem nutrient subsidies to islands but how they influence mangrove ecosystems is poorly known. We assessed the contribution, uptake, cycling and transfer of nutrients from seabird colonies in remote mangrove systems free of human stressors. We found that nutrients from seabird guano enrich mangrove plants, reduce nutrient limitations, enhance mangrove invertebrate food webs and are exported to nearby coastal habitats through tidal flow. We show that seabird nutrient subsidies in mangroves can be substantial, improving the nutrient status and health of mangroves and adjacent coastal habitats. Conserving mobile consumers, such as seabirds, is therefore vital to preserve and enhance their role in mangrove productivity, resilience and provision of diverse functions and services.

Keywords: Aldabra Atoll, Functional connectivity, Stable isotopes, Indian Ocean, Invertebrates, Island ecosystems, *Rhizophora mucronata*

INTRODUCTION

Mangrove forests occupies the interface between land and sea, where it plays critical roles in sustaining biodiversity and maintaining ecological functions and services [1,2]. For example, mangrove habitat provides protection against coastal erosion and tidal surges [3]; provides essential food, protection and habitat to numerous taxa during part or all of their life cycles [4]; and sequesters large amounts of carbon, contributing substantially to climate change adaptation and mitigation [5]. The provisioning of these functions and services are dependent on the status and health of mangrove forests [6], which are increasingly threatened by human activities. Eutrophication, caused by excessive human-derived nutrient inputs is a major threat for coastal ecosystems, including mangroves [7]. Nutrient inputs from anthropogenic activities tend to be rich in nitrogen (N) but poor in phosphorus (P), and can reduce mangrove growth, cause death of pneumatophores and soil anoxia [8], and increase mortality of mangrove forests [9]. Although strong links have been documented between human-derived nutrient enrichment and mangrove ecosystem functioning, there has been far less attention on the impacts of mangrove nutrient enrichment by natural sources.

Mobile consumers play a key role in the movement of nutrients across ecosystem boundaries [10]. Seabirds comprise an important group of mobile consumers involved in the transport of nutrients from sea to land, by feeding in oceanic areas and depositing large amounts of guano in their colonies [11]. Enriched in both N and P compounds, seabird-derived nutrients enhance primary productivity around their colonies, and in coastal habitats such as coral reefs, resulting in increased growth rate of reef fish [12,13] and coral [14], as well as increased fish biomass [12]. Despite the widespread occurrence of seabirds nesting in mangroves [7], much less is known about the influence of seabird-derived nutrients in mangrove ecosystems. Previous studies documented positive relationships between seabird nutrient subsidies and mangrove productivity [15] and nutrient status [16–18], but these studies only focused on mangroves in northern and central America in proximity to urban centers. How seabird-derived nutrients influence mangrove forests in regions such as the extensive and mangrove-rich Indo-Pacific, and in the absence of human influence, is unknown. Additionally, no studies have explored the vertical transfer of seabird-derived nutrients within mangrove food webs. Furthermore, tides strongly mediate connectivity of mangroves with adjacent coastal habitats through the exchange of detritus, fauna and nutrients [19]. Consequently, seabird influence may extend beyond mangroves in the coastal seascape; however, the horizontal transfer of seabird-derived nutrients across mangrove

boundaries has not been studied. Documenting these relationships and linkages is increasingly

important given the accelerated declines in both mangrove habitats [20] and seabird populations [21,22]

due to numerous anthropogenic threats.

Here, we examined the contribution, uptake and transfer of seabird-derived nutrients in

mangrove habitats on Aldabra Atoll, in the Southern Seychelles, one of the largest mangrove-nesting

seabird colonies in the Indian Ocean. Aldabra hosts the largest area of mangroves in the Seychelles

archipelago, which supports the largest breeding populations of frigatebirds, *Fregata minor* and *F.*

ariel, in the Indian Ocean [23], as well as one of the largest breeding populations of red-footed boobies

Sula sula in the region. Frigatebirds and red-footed boobies nest exclusively in mangrove forests

around Aldabra's lagoon shores [24]. With only a small research station, human impact on Aldabra is

minimal, presenting ideal conditions to assess the effects of seabird subsidies in a relatively

undisturbed mangrove system. We aimed to answer the following questions: (1) What quantities of

marine-derived nutrients are contributed by mangrove-nesting seabirds? (2) Are seabird-derived

nutrients transferred to and assimilated by mangroves? (3) Do seabird-derived nutrients alleviate

nutrient limitation and reduce nutrient resorption efficiency of mangroves? (4) Are seabird-derived

nutrients trophically transferred to mangrove fauna? And (5) Are seabird-derived nutrients in

mangroves exported to adjacent coastal habitats?

We achieved this using biogeochemical assays of multiple ecosystem components, including

seabird guano, mangrove sediment, leaves, gastropods, crabs, adjacent macroalgae and seawater,

sampled at 10 sites with (n = 5) and without (n = 5) nesting seabirds. Using seabird demography

metrics, we find that seabird nutrient subsidies in mangroves can be substantial. Nutrient assays

reveal seabird-derived nutrients enrich mangroves and reduce their nutrient limitations, but do not

influence their resorption efficiency. Seabird-derived nutrients enrich mangrove invertebrates through

trophic transfer and are exported to nearby coastal habitats through tidal flow. By showing that

seabird-derived nutrients promote nutrient status and health of mangroves, our results increase

understanding about the implications of declines of seabird populations, and suggest that efforts to

maintain or restore their populations and breeding grounds should be prioritized.

RESULTS

Seabird nutrient contributions to mangroves are substantial

Red-footed booby guano contained 10.1 ± 2.92 %N and 10.2 ± 0.87 %P (Mean ± SD, n = 4) and frigatebird guano had 10.9 ± 2.06 %N and 10.3 ± 1.14 %P (Mean ± SD, n = 4). Guano nitrogen isotopic values (δ^{15}N) for red-footed booby were 10.8 ± 2.71 (Mean ± SD, n = 20) and for frigatebirds 12.9 ± 3.63 (Mean ± SD, n = 20). Based on seabird population size and breeding metrics, and guano nutrient concentrations (see experimental procedures), estimated total annual nutrient contributions were higher for red-footed boobies (28.6 N tonne.year^{-1}; 28.9 P tonne.year^{-1}) than for frigatebirds (12.8 N tonne.year^{-1}; 12.0 P tonne.year^{-1}).

Seabird-derived nutrients are transferred and assimilated by mangroves

We recorded mangrove forest metrics at all sites, resulting in a total of 923 surveyed trees, with *R. mucronata* making up 79% of individuals and being the dominant species at all surveyed sites (Table S1). δ^{15}N in *R. mucronata* leaves and mangrove sediment were higher at seabird sites compared to non-seabird sites (Figure 1D, 1E; Table S2). Nutrient levels in *R. mucronata* leaves were higher in the presence of nesting seabirds (Table S2), for both N (Mean ± SD: seabird = 1.02 ± 0.21%, non-seabird = 0.73 ± 0.09%, $P < 0.0001$) and P (seabird = 0.09 ± 0.02%, non-seabird = 0.07 ± 0.01%, $P = 0.01$; Figure 2A, B).

Seabird-derived nutrients reduce mangrove nutrient limitations but not resorption efficiency

Nutrient ratios, an indicator for nutrient limitation, were lower at seabird sites (C:N: seabird = 43.3 ± 8.64, non-seabird = 58.0 ± 6.95, $P < 0.0001$; and C:P: seabird = 524.4 ± 122.6, non-seabird = 643.2 ± 115.0, $P = 0.031$; Figure 2C, D). The overall average N:P ratio of mangroves was 11.7 ± 2.45. We estimated mangrove nutrient resorption efficiency as an indicator for mangrove nutrient cycling, using nutrient concentrations of green and senescent leaves (see experimental procedures). Nutrient resorption efficiency was similar between sites with and without seabirds (N resorption efficiency [RE-N]: seabird = 53.5 ± 13.1%, non-seabird = 59.5 ± 7.39%, $P = 0.15$; and P resorption efficiency [RE-P]: seabird = 50.1 ± 25.2%, non-seabird = 59.2 ± 10.9%, $P = 0.24$; Figure 2E, F; Table S2).

Seabird-derived nutrients are transferred to mangrove invertebrate food web

δ^{15}N were higher at seabird sites compared to non-seabird sites at all trophic levels, including for herbivorous gastropods (seabird = 13.5 ± 4.64%, non-seabird = 4.45 ± 3.42%, $P < 0.0001$; Figure

1C), herbivorous sesarmid crabs (seabird = 12.03 ± 2.16%, non-seabird = 6.79 ± 1.47%, $P < 0.001$;

Figure 1B) and omnivorous portunid crabs (seabird = 11.78 ± 1.09%, non-seabird = 8.32 ± 1.62%, $P <$

0.005; Figure 1A).

Seabird-derived nutrients are exported to nearby coastal habitats

Surface seawater in the lagoon had higher NOx (nitrate + nitrite) at seabird sites than non-seabird

sites only during outgoing tides (post-hoc tests: $P = 0.033$; Figure 3A; Table S3), whereas phosphate

concentrations were higher at seabird than non-seabird sites during both incoming and outgoing tides

(post-hoc tests: incoming: $P < 0.0001$, outgoing: $P = 0.0051$; Figure 3B; Table S3). Macroalgae

Halimeda spp. sampled on substrates adjacent to mangroves had higher $\delta^{15}N$ at seabird than non-

seabird sites (seabird = 9.63 ± 1.62, non-seabird = 5.98 ± 0.91, $P = 0.021$; Figure 3C).

DISCUSSION

The ecological functions and ecosystem services provided by mangroves are reliant on the status and

health of mangrove forests [6]. It is therefore important to understand mangrove responses to natural

processes that influence their status and health. Our study presents insights into the scale and extent

of influence of cross-ecosystem nutrient subsidies on the ecology and function of Indo-Pacific

mangroves and in the absence of human influence. Seabirds feed in open ocean and excrete

substantial quantities of nutrients in mangrove forests where they nest. Mangroves with nesting

seabirds were enriched in N and P and relieved of their nutrient limitations, although seabirds did not

influence their nutrient resorption efficiencies. Through trophic pathways, seabird-derived nutrients

were transferred to mangrove-associated invertebrate food webs. Furthermore, we documented the

export of seabird-derived nutrients from mangroves to nearby coastal habitats through tidal flow,

showing that the influence of nutrient contributions by mangrove-nesting seabirds extends beyond

mangrove boundaries. Our study highlights the important roles of mobile consumers such as seabirds

in maintaining functional connectivity between oceanic systems and coastal seascapes [25], and in

improving the nutrient status of coastal habitats (Figure 4) and isolated island ecosystems. Our study

adds more weight and nuance to the growing body of research on the influence of seabirds on highly

productive coastal ecosystems [26–28].

Oceanic atolls are particularly nutrient deficient as they are isolated from mainland nutrient

sources and surrounded by oligotrophic waters. On atolls with seabird colonies, seabirds represent

one of the main sources of marine-derived nutrient subsidies [29]. Breeding red-footed boobies and

frigatebirds delivered a combined total of 41.4 N tonne.year^{-1} and 40.9 P tonne.year^{-1} to Aldabra's

mangroves. This represents a minimum estimate since we did not account for non-breeding

individuals. Furthermore, due to the bi-annual breeding of red-footed boobies and protracted breeding

of frigatebirds [30], seabird-derived nutrient subsidies are delivered year-round. However, the majority of

seabirds on Aldabra nest in tall mangrove trees along the northern and eastern lagoon shores [24].

Accordingly, seabird nutrient contributions are disproportionately distributed, with Aldabra's southern

lagoon shores and terrestrial habitats deprived of seabird-derived nutrient subsidies. Additional

sources of marine subsidies on Aldabra include algal wrack from seagrass beds [31], nesting sea turtles

[32] and shorebirds [33], all of which contribute mainly to the sandy beach habitat around the atoll.

Upwellings may also provide nutrient input to Aldabra's coasts [34]. Quantifying these additional

sources and assessing their influence on the land- and seascape would provide a more holistic

overview of marine nutrient budgets and functioning of marine-subsidized ecosystems on the atoll.

The mangrove forests in our study were dominated by *R. mucronata*, a species widespread in

the Indo-Pacific region, and the only *Rhizophora* species occurring in the Western Indian Ocean [35,36].

The higher $\delta^{15}N$ concentration found in sediment and leaves at seabird sites confirms that seabird-

derived nutrients are transferred to mangroves. Foliar N and P concentrations of *R. mucronata* were

39% and 28% higher, respectively, at seabird sites, providing evidence of seabird-derived nutrient

uptake by mangroves. Our results corroborate documented increases in nutrients at seabird sites

relative to control sites for coastal *Rhizophora* species in Central and North America, for example *R.*

mangle (Belize: +37% N [37]; Honduras: +41% N and +20% P [18]; Mexico: +25 %N and +17 %P [16];

Florida: +33 %N [15]). Mangroves on Aldabra grow in a lagoonal low-lying carbonate environment and

do not receive large volumes of terrigenous sediment supply and nutrients compared to mangroves

growing in deltas or estuaries [38,39]. In this nutrient-limiting environment, our results show that seabirds

contribute essential macronutrients and improve the nutrient status of mangroves.

Responses of mangroves to nutrient enrichment are site- and species-specific, with the main

limiting nutrient of the site governing species nutrient requirement and tolerances [40]. Nutrient limitation

in mangroves is usually attributed to N or P limitation [41]. Foliar N:P ratios can be used to infer nutrient

limitation [42], with N:P ratios > 32 indicating P limitation in mangroves [7,43]. Foliar N:P ratio in our study

was 11.7 ± 2.45 (Mean ± SD), suggesting mangroves in our study are N-limited, which is typical for

fringe mangroves influenced by frequent tidal flushing [7,44]. N limitation at our study site is further

indicated by the equally low N:P ratios documented in previous analyses on mangrove soil [45] and

lagoonal sediment porewater on Aldabra [46]. At our seabird sites, uptake of seabird-derived nutrients

alleviated mangrove nutrient limitations of both N and P, observed by reductions in foliar C:N and C:P

ratios of *R. mucronata*. Nutrient enrichment in N-limited mangroves can generate multiple higher-

order effects. For example, in Florida, seabird-derived nutrients increased primary production, leaf

nutritional value and herbivory [15]. Similarly, experimental nutrient enrichment in N-limited mangroves

has demonstrated increases in shoot biomass and tree growth [40].

Resorption of nutrients prior to leaf fall, a process in which nutrients are resorbed from

senescent leaves and are directly available for continued plant growth, is a vital nutrient recycling and

conservation mechanism in mangroves [7,47]. When nutrients become available, mangroves reduce

their nutrient resorption efficiency [48,49]. Based on this, we expected reductions in resorption efficiency

at our seabird sites compared to non-seabird sites; however, no differences in RE-N or RE-P were

detected. Furthermore, given that our study site is N-limited, we would expect greater resorption of N

compared to P [47], however, average RE-N values were similar to average RE-P values at both

seabird and non-seabird sites. Seabird-derived nutrient enrichment has been documented to reduce

mangrove resorption efficiency in Mexico [16] but not in Belize [37]. Similar contrasting patterns are shown

in experimental nutrient enrichment studies [50,51], suggesting internal nutrient cycling by mangroves is

complex, regulated by the interacting effects between species-specific physiological capacity to

conserve nutrients [49], nutrient demand (through increased growth rate) and supply [7,51].

A consequence of nutrient enrichment is that it can reduce stability and resilience of

mangrove forests [9,52,53]. Nutrient enrichment causes plants to invest more in aboveground growth and

less in belowground biomass [54,55]. Furthermore, nutrient availability increases rates of microbial

decomposition of organic matter [48,56]. A forest with lower root biomass and higher decomposition

rates, is likely to have reduced accretion of sediment organic matter [57], resulting in reduced shoreline

stability [52] and increased probability of mangrove death when faced with environmental stressors or

events [9,53]. On this basis, seabird-derived nutrient enrichment has been linked to declines of

mangrove cays in Belize [37]. However, in contrast to Belize, some of the greatest increases in

mangrove extent on Aldabra over the past two decades coincided with seabird nesting locations [58].

Furthermore, mangroves on Aldabra are far from urban or agricultural centers compared to Belize,

and are therefore not influenced by additional anthropogenic nutrient sources. Natural nutrient

sources provide N and P in optimal ratios [59,60], generating contrasting responses in coastal habitat

structure and functions compared to anthropogenic sources [61]. In addition, seabirds on Aldabra nest

primarily in tall trees in the mangrove fringes, which receive constant tidal flushing, limiting over-

enrichment of nutrients [9]. Altogether, this suggests that seabird-derived nutrient enrichment is unlikely

to be causing declines in mangroves on Aldabra. Soil nutrient content is a main driver of mangrove

aboveground biomass on Aldabra [45]. Therefore, by fertilizing mangroves where they nest, seabirds

likely enhance mangrove productivity in their breeding areas, further promoting suitable nesting

habitat, likely creating a positive-feedback loop for seabird populations [10]. Indeed, both mangroves

and mangrove-nesting seabird populations on Aldabra have increased over the last few decades [23,58].

Multiple pathways of energy and nutrient flow exist in mangrove food webs due to the wide

range of available food sources, such as plankton, benthic microalgae, mangroves, macroalgae and

macrophytes [4,62]. By assessing the trophic transfer of seabird-derived nutrients, we found that

seabirds also enriched the mangrove invertebrate food web. We detected higher δ^{15}N at seabird sites

at all trophic levels, including baseline (sediments), primary producers (mangrove leaves), primary

consumers (gastropods and sesarmid crabs) and a secondary consumer (portunid crab). Invertebrate

food web enrichment by seabird guano has been documented in other coastal ecosystems such as

littoral habitats [63], fjords [64] and coastal ponds [65], but not previously in mangrove forests. Invertebrates

are key components of mangrove-associated macrofauna, playing substantial roles in nutrient cycling

of detritus [7]. Further research should explore additional pathways of seabird-derived nutrient flow in

mangroves, as well as how seabird nutrient enrichment affects invertebrate trophic structure,

interactions and ecological functions.

Given the absence of anthropogenic influence in Aldabra's lagoon, the higher levels of NOx

and phosphate in seawater adjacent to mangroves with seabird colonies can be attributed to seabird

guano inputs. Increased levels of seawater nutrients around seabird colonies in remote oligotrophic

locations confirms seabird-derived nutrients as an important source of marine productivity [66]. Indeed,

δ^{15}N levels of macroalgae *Halimeda* spp. growing adjacent to mangroves were also higher compared

to non-seabird sites. These results indicate horizontal transfer of seabird-derived nutrients, from

mangroves through tidally-mediated nutrient exchange. Nutrients can also be transferred to adjacent

coastal habitats (such as coral reefs or seagrass areas) via mangrove leaf litter, which plays a key

role in sustaining adjacent marine food webs [62]. Leaves with higher-nutrient content or nutritive value

are more rapidly broken down than less nutrient-rich leaves [15]. Seabird-derived nutrients from

mangroves may therefore extend to adjacent marine communities through trophic pathways, for

example via fish communities from nearby reefs or seagrass beds that utilize mangroves as nursery

or feeding habitat.

Mangroves on Aldabra support a wide range of biodiversity and ecosystem service benefits.

In addition to supporting large breeding populations of seabirds, Aldabra's mangroves sustain

threatened and regionally important populations of numerous marine species. Aldabra's mangroves

provide nursery habitat for one of the largest green turtle populations in the region [32], play a critical

role as feeding, breeding and nursery habitats for numerous bony fish and shark species [67]. Aldabra's

mangroves support the highest biomass of herbivorous fish and the highest abundance of sharks in

Seychelles [68,69]. Furthermore, Aldabra's mangroves comprise one of Seychelles' largest blue carbon

ecosystems [45]. By improving mangrove nutrient status and health, we show that seabirds nesting on

Aldabra boost biodiversity and ecosystem services.

Our study provides important insights to nutrient contribution, cycling and transfer in a system

with strong land-sea connectivity and without human stressors. We present critical data to start

unravelling the mechanisms and extent of nutrient connectivity in mangroves, linked to the cross-

ecosystem ocean-derived subsidies provided by seabirds to islands. Given the critical roles of mobile

consumers such as seabirds in maintaining ecosystem health and functions, conservation and

management actions should focus on restoring seabird populations and their breeding grounds. Such

efforts will present maximum benefits for people and biodiversity at multiple scales [70].

EXPERIMENTAL PROCEDURES

Resource availability

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the lead

contact, Jennifer Appoo jennifer.appoo@univ-reunion.fr

Materials availability

This study did not generate new unique materials.

Data and code availability

Data and code to replicate the findings presented will be made available on Dryad Digital Repository

upon acceptance.

Any additional information required for reanalyzing the data reported in this paper is available from the

lead contact upon request.

Study site

Aldabra Atoll (9°24′ S, 46°20′ E) is part of the Seychelles archipelago (Figure S1). The atoll comprises

four main islands (land area 15,500 ha, mean elevation 8 m asl), encircling a large shallow lagoon

(20,500 ha) and separated by deep (10–30 m depth) narrow channels [71,72]. Aldabra's mangroves

occur primarily along the lagoon shores, covering 1,720 ha [73]. Mangrove habitat on the northern and

south-eastern lagoon shores consists of tall trees forming closed canopies, while mangrove areas on

Aldabra's southern shores are smaller and scattered (see Table S1). Seven mangrove species occur,

with *Rhizophora mucronata*, *Bruguiera gymnorhiza*, *Ceriops tagal* and *Avicennia marina* being the

most common [74]. Mangrove sediments are mainly sand (53%) and loamy sand (26%) and with an

average salinity of 8.3 ± 3.1 (practical salinity scale) [58]. Aldabra is influenced by a semi-diurnal meso-

tidal regime (range: 2–3 m), resulting in strong tidal currents that drain approximately 75% of the

lagoon at low tide through the main channels [72,75]. Rainfall patterns on Aldabra show marked

seasonal variations influenced by monsoon winds. The majority of rain is concentrated between

November and April with a mean of 975 mm.year^{-1}, however rainfall does not influence primary

productivity of the mangroves [76].

On Aldabra, great and lesser frigatebirds, *Fregata minor* and *F. ariel* respectively, nest

together in four separate colonies, with a combined breeding population estimated at 16,534 pairs

(SIF, *unpubl. data*). The red-footed booby *Sula sula* nests together with or outside the frigatebird

colonies, totaling 36,720 pairs (SIF, *unpubl. data*). Great frigatebird breeding peaks in October, while

lesser frigatebirds and red-footed boobies have two breeding peaks, in June and October, and

February and August, respectively (SIF, *unpubl. data*). The distribution of mangrove-nesting seabirds

is not uniform, with seabirds found only in mangroves on the northern, eastern and south-eastern

lagoon shores [30]. Mangroves and seabirds on Aldabra are recognized globally, as a Ramsar Wetland

Site of International Importance and an Important Bird Area (BirdLife International), respectively.

Furthermore, Aldabra's mangroves and seabirds have been strictly protected since designation of the

atoll as a Strict Nature Reserve in 1981 and UNESCO World Heritage Site in 1982.

Experimental design

We investigated the effects of breeding seabirds on Aldabra's mangroves during the nesting period of

all three species (November 2022 to March 2023). Sampling was conducted at 10 sites comprising

five sites with no or low numbers of nesting seabirds (< 20 nests) and five sites with high numbers of

nesting seabirds (Figure S1). Sites were distributed along the lagoon shore around the atoll and

situated at least 1 km apart. All sampling sites were located on the seaward fringe of mangroves,

which is where the majority of seabirds roost and nests on Aldabra. Within each site, we conducted

mangrove measurements within eight quadrats of 5 x 5 m, separated by a minimum of 50 m.

Mangrove sampling was conducted at spring low tide when mangroves and intertidal areas were

exposed.

Seabird nutrient contributions

To determine the amount of nutrients deposited by mangrove-nesting seabirds, we collected

droppings around nests of red-footed boobies and frigatebirds. For the latter, droppings were sampled

around nests with big chicks. Given the difficulty of correctly identifying frigatebird species with chicks,

the samples were generalized to represent both great and lesser frigatebirds. Droppings were kept

cool in the field and then stored frozen in the laboratory until further processing. Total nitrogen (N)

content in seabird droppings was obtained by mineralization according to the Kjeldahl method [77]

(Büchi KjelMaster K-375, Switzerland). Total phosphorus (P) was determined by preliminary dry

combustion (600 °C), then dissolving the residual material in 6 M HCl. P concentration was obtained

by extraction using ammonium molybdate and ascorbic acid and measured with a continuous flow

colorimeter (Proxima, Alliance Instrument, USA). Nutrient analyses were conducted at the CIRAD

laboratory of agronomy in Reunion Island (France). Additional seabird droppings were collected for

isotope analysis (see below).

We used N and P concentrations in seabird droppings to estimate the total annual N and P

input from breeding seabirds using previously used methods [78,12]:

$$Nutrient_{gi} = Q_{gi} \times Dr_i \times \left[(Bd_i \times T_i \times f_i) + \left(\frac{P_i}{2} \times Bd_i \right) \right]$$

where $Nutrient_{gi}$, t.yr^{-1} is the annual input per nutrient type and species, Q_{gi}, mg.g^{-1} is the

concentration of N or P measured in seabird droppings for each species, Bd_i is number of breeding

adults for that species, T_i is the length of the breeding period (number of days from courtship to chick-

rearing, f_i is the proportion of time spent at the colony during breeding to account for absence due to

feeding forays and P_i is the productivity of the species (fledged chicks per breeding pair). Defecation

rate for red-footed boobies was obtained from Young et al., [79]. For frigatebirds, we estimated

defecation rates based on red-footed booby measurements, scaled allometrically to body size

following Staunton-Smith and Johnson [80], and using a median value of the adult body sizes of lesser

and great frigatebirds. Values for the length of the breeding period, proportion of time at the colony,

productivity and adult body sizes for each species were obtained from Riddick et al., [78]. Our estimates

do not account for nutrient contributions by non-breeding individuals since we lacked data on

numbers and diurnal movement patterns.

Mangrove forest structure

We assessed forest characteristics in each quadrat following methods previously employed on

Aldabra [45]. We recorded the species of all trees >2 m height and measured their diameter at breast

height (DBH, 1.3 m height) using a measuring tape, following guidelines for trees with anomalies such

as prop roots [81]. We counted the number of seedlings and used an inclinometer (Suunto PM-5/66 PC,

Vantaa, Finland) to measure tree height to the nearest ± 0.5 m. We applied the allometric equation by

Chave et al., [82] to determine aboveground biomass (ABG) following Constance et al., [45]. Aboveground

biomass (ABG, kg) per tree was obtained from the wood density (ρ, g.cm^{-3}), the DBH (D, cm), and

height (H, m).

$$ABG = 0.0509 \times \rho \times D^2 \times H$$

Mangrove foliar nutrient analyses

To test the effects of seabird subsidies on mangrove nutrients, we measured leaf nutrient concentrations of the dominant species *R. mucronata*. In each sampling quadrat, we collected six young fully expanded green leaves exposed to the sun and six yellow (senescent) leaves from at least three individual trees. The leaves were pooled per quadrat to form one composite green and yellow leaf sample. Leaves were rinsed with fresh water, oven-dried at 60°C for a minimum of 48 hrs and powdered using a ball mill. Total carbon (C) and N were determined using an elemental-analyser (Elementar Vario Micro Cube, Lancaster University, UK). Total P content was determined by the Olsen-Dabin method using 0.5 M sodium hydrogen carbonate solution with a continuous flow colorimeter (Proxima, Alliance Instrument, USA) at the CIRAD laboratory of agronomy in Reunion Island (France). Sub-samples were obtained for isotope analysis.

We derived foliar nutrient ratios (C:N, C:P mass basis) to explore nutrient limitations to mangrove growth [7,16]. As an indicator of nutrient cycling, we calculated the nutrient resorption efficiency of N (RE-N) and P (RE-P), which corresponds to the percentage of N or P recovered from senescing leaves prior to leaf fall [40,47], based on the equation by Chapin and Cleve [83]:

$$RE \% = \frac{N \text{ or } P \text{ green leaves} - N \text{ or } P \text{ senescent leaves}}{N \text{ or } P \text{ green leaves}} \times 100$$

Isotope sampling and analyses

To assess the transfer of seabird-derived nutrients in mangroves we conducted isotopic analyses on seabird droppings, mangrove sediment and green *R. mucronata* leaves. Surface sediments were sampled by inserting a PVC corer ($\varnothing$ = 22 mm) in the top 2 cm sediment layer at three random positions in each quadrat. To assess the trophic transfer of seabird-derived nutrients to mangrove fauna we conducted isotopic analyses on mangrove invertebrates comprising of gastropods and crabs occupying different trophic levels. As primary consumers, we collected herbivorous gastropods of the genus *Littoraria* from mangrove roots and trunk within each quadrat, and leaf-eating sesarmid crab *Sesarma leptosoma* on mangrove trees at each site. As secondary consumers, we collected omnivorous portunid crab *Thalamita crenata* on the intertidal mudflat adjacent to mangrove forests. We sampled foot muscle from gastropods, body muscle from sesarmid crabs and limb muscle from portunid crabs. We collected macroalgae on rocky outcrops of the intertidal mudflat and within 50 m of

the mangrove fringe for isotopic analyses to assess nutrient uptake in the adjacent habitat. We

selected macroalgae since they incorporate nutrients over a relatively long period (weeks to months),

constituting an ideal proxy for ambient nutrient conditions [84]. At each site, we randomly collected thalli

of the macroalgae *Halimeda* spp. The latter has proven a good proxy for seabird nutrients since it

responds positively to seabird subsidies [60].

Isotopic values of seabird-derived nutrients are enriched in nitrogen ($\delta^{15}N$) due to the high

trophic position of seabirds in the marine food chain and enrichment of $\delta^{15}N$ in seabird guano after

deposition [85]. $\delta^{15}N$ is therefore used to trace seabird-derived nutrients in recipient communities [27]. All

samples for isotopic analyses were dried at 60°C and ground into a homogeneous powder using a

ball mill. Isotopic ratios were measured using an Isoprime 100 Isotope Ratio Mass Spectrometer with

international standards IAEA 600, USGS 41 and CH6, at the stable isotope facility at Lancaster

University (Lancaster, UK). Accuracy based on internal standards was within 0.1 ‰ SD and selected

samples were run in triplicate to further ensure accuracy of readings.

Surface seawater nutrient concentrations

The semi-diurnal tidal fluctuations at Aldabra generate large and rapid changes in water levels,

causing strong tidal currents [72]. Most of the guano inputs by mangrove-nesting seabirds on Aldabra

are deposited directly in the water or are washed away during the subsequent high tide when

deposited at low tide [86]. To assess nutrient export from the seabird colonies, we sampled surface

seawater in 30-ml containers at each site on two occasions; on the incoming tide while the lagoon fills

and on the outgoing tide while the lagoon drains. Samples were collected in duplicate at each location

and occasion and treated at 80°C in a drying oven for a minimum of 2.5 hrs. Samples were stored at

room temperature in the dark until laboratory analysis. NO_x (nitrate NO_3^- + nitrite NO_2^-) and phosphate

(PO_4^{3-}) concentrations were determined using an AA3 auto-analyser (Seal Analytical) following the

method of Aminot and Kérouel [87]. Samples were processed at IRD laboratory in Plouzané (France)

and had an accuracy of 0.5 µmol.L^{-1} and 0.7 µmol.L^{-1}, for NO_x and PO_4^{3-}, respectively.

Data analyses

We assessed the influence of seabirds on nutrient parameters using univariate tests of differences

between seabird and non-seabird sites. We formulated linear mixed models (LMMs) for each nutrient

parameter (%N, %P, C:N, C:P, RE-N, RE-P) of mangrove leaves and $\delta^{15}N$ for mangrove leaves,

sediment, gastropod, crabs and macroalgae with seabird status (seabird/no seabird) as fixed effect

and site as random effect. For some parameters, variances were heterogeneous between the

predictor levels (i.e., higher within seabird than non-seabird sites), therefore we included the

'VarIdent' error structure on seabird status, which allows different variance of predictor levels. We

assessed residual spatial autocorrelation of each model by plotting residuals against location and

found no spatial autocorrelation in any of these models. This was confirmed by comparing models

with and without spatial weights matrix using Akaike's Information Criterion (AIC), and models without

had the lowest AIC values. Model diagnostics were performed by plotting residuals against fitted

values and explanatory variables to verify assumptions of homogeneity of variance and independence

following Zuur and Ieno [88].

We assessed differences in surface seawater nutrient concentrations separately for NOx and

phosphate. We used LMMs with seabird status, including an interaction with tidal regime (incoming/

outgoing) as fixed effects and site as random effect. These models included the 'VarIdent' error

structure on seabird status. Where significant differences were detected, we performed post-hoc tests

using the *emmeans* package [89]. Model residuals were assessed to verify model assumptions and to

confirm there were no spatial dependencies [88]. All models were formulated using the function 'lme'

from package *nlme* [90] and analyses were conducted on R version 4.3.0 [91].

SUPPLEMENTAL INFORMATION

Document S1: Figure S1 and Tables S1-S3.

Supplemental Information includes one figure and three tables and can be found online at:

ACKNOWLEDGMENTS

We thank Seychelles Islands Foundation staff at head office and on Aldabra Atoll for providing

administrative and logistical support during the fieldwork. Funding was provided by the African World

Heritage Fund through a Moses Mapesa Research grant to J.A, the PADI Foundation marine

research grant to J.A, and the Bertarelli Foundation under the Bertarelli Marine Science Program to

N.A.J.G. J.A was supported by a doctoral fellowship from the Reunion Island Regional Council.

Fieldwork was conducted with approval of the Seychelles Bureau of Standards under permit number

A0157 and the Seychelles Islands Foundation. Icons for Figure 4 were obtained from University of

Maryland Center for Environmental Science, Integration and Application Network

(https://ian.umces.edu/media-library/), Freepik (https://www.freepik.com/) and Birds Caribbean

(https://www.birdscaribbean.org/).

AUTHOR CONTRIBUTIONS

Conceptualization: J.A, N.B, N.A.J.G and S.J; Methodology: J.A and N.B; Data collection: J.A; Formal

analysis: J.A; Writing – Original Draft: J.A; Writing – Review and Editing: N.A.J.G, N.B, S.J, A.L and

M.L.C.

DECLARATION OF INTERETS

The authors declare no competing interest.

REFERENCES

1. Walters, B.B., Rönnbäck, P., Kovacs, J.M., Crona, B., Hussain, S.A., Badola, R., Primavera, J.H.,
Barbier, E., and Dahdouh-Guebas, F. (2008). Ethnobiology, socio-economics and management
of mangrove forests: A review. Aquat. Bot. *89*, 220–236. 10.1016/j.aquabot.2008.02.009.

2. Worthington, T.A., Andradi-Brown, D.A., Bhargava, R., Buelow, C., Bunting, P., Duncan, C.,
Fatoyinbo, L., Friess, D.A., Goldberg, L., Hilarides, L., et al. (2020). Harnessing Big Data to
Support the Conservation and Rehabilitation of Mangrove Forests Globally. One Earth 2, 429–
443. 10.1016/j.oneear.2020.04.018.

3. Ewel, K.C., Twilley, R.R., and Ong, J.E. (1998). Different kinds of mangrove forests provide
different goods and services. Glob. Ecol. Biogeogr. Lett 7, 83–94.

4. Nagelkerken, I., Blaber, S.J.M., Bouillon, S., Green, P., Haywood, M., Kirton, L.G., Meynecke, J.-
O., Pawlik, J., Penrose, H.M., Sasekumar, A., et al. (2008). The habitat function of mangroves for
terrestrial and marine fauna: A review. Aquat. Bot. *89*, 155–185. 10.1016/j.aquabot.2007.12.007.

5. Alongi, D.M. (2020). Global Significance of Mangrove Blue Carbon in Climate Change Mitigation.
Sci *2*, 67. 10.3390/sci2030067.

6. Salem, M.E., and Mercer, D.E. (2012). The Economic Value of Mangroves: A Meta-Analysis.
Sustainability *4*, 359–383. 10.3390/su4030359.

7. Reef, R., Feller, I.C., and Lovelock, C.E. (2010). Nutrition of mangroves. Tree Physiol. *30*, 1148–
1160. 10.1093/treephys/tpq048.

8. Mandura, A.S. (1997). A mangrove stand under sewage pollution stress: Red Sea. Mangroves
Salt Marshes, 255–262.

9. Lovelock, C.E., Ball, M.C., Martin, K.C., and C. Feller, I. (2009). Nutrient Enrichment Increases
Mortality of Mangroves. PLoS ONE *4*, e5600. 10.1371/journal.pone.0005600.

10. McInturf, A.G., Pollack, L., Yang, L.H., and Spiegel, O. (2019). Vectors with autonomy: what
distinguishes animal-mediated nutrient transport from abiotic vectors? Biol. Rev. *94*, 1761–1773.
10.1111/brv.12525.

11. Doughty, C.E., Roman, J., Faurby, S., Wolf, A., Haque, A., Bakker, E.S., Malhi, Y., Dunning, J.B.,
and Svenning, J.-C. (2016). Global nutrient transport in a world of giants. Proc. Natl. Acad. Sci.
113, 868–873. 10.1073/pnas.1502549112.

12. Graham, N.A.J., Wilson, S.K., Carr, P., Hoey, A.S., Jennings, S., and MacNeil, M.A. (2018).
Seabirds enhance coral reef productivity and functioning in the absence of invasive rats. Nature
559, 250–253. 10.1038/s41586-018-0202-3.

13. Benkwitt, C.E., Taylor, B.M., Meekan, M.G., and Graham, N.A.J. (2021). Natural nutrient
subsidies alter demographic rates in a functionally important coral-reef fish. Sci. Rep. *11*.
https://www.nature.com/articles/s41598-021-91884-y.

14. Savage, C. (2019). Seabird nutrients are assimilated by corals and enhance coral growth rates.
Sci. Rep. *9*, 4284. 10.1038/s41598-019-41030-6.

15. Onuf, C.P., Teal, J.M., and Valiela, I. (1977). Interactions of Nutrients, Plant Growth and
Herbivory in a Mangrove Ecosystem. Ecology *58*, 514–526. 10.2307/1939001.

16. Adame, M., Fry, B., Gamboa, J., and Herrera-Silveira, J. (2015). Nutrient subsidies delivered by
seabirds to mangrove islands. Mar. Ecol. Prog. Ser. *525*, 15–24. 10.3354/meps11197.

17. Irick, D.L., Gu, B., Li, Y.C., Inglett, P.W., Frederick, P.C., Ross, M.S., Wright, A.L., and Ewe,
S.M.L. (2015). Wading bird guano enrichment of soil nutrients in tree islands of the Florida
Everglades. Sci. Total Environ. *532*, 40–47. 10.1016/j.scitotenv.2015.05.097.

18. McFadden, T.N., Kauffman, J.B., and Bhomia, R.K. (2016). Effects of nesting waterbirds on
nutrient levels in mangroves, Gulf of Fonseca, Honduras. Wetl. Ecol. Manag. *24*, 217–229.
10.1007/s11273-016-9480-4.

19. Feller, I.C., Lovelock, C.E., Berger, U., McKee, K.L., Joye, S.B., and Ball, M.C. (2010).
Biocomplexity in Mangrove Ecosystems. Annu. Rev. Mar. Sci. *2*, 395–417.
10.1146/annurev.marine.010908.163809.

20. Friess, D.A., Rogers, K., Lovelock, C.E., Krauss, K.W., Hamilton, S.E., Lee, S.Y., Lucas, R.,
Primavera, J., Rajkaran, A., and Shi, S. (2019). The State of the World's Mangrove Forests: Past,
Present, and Future. Annu. Rev. Environ. Resour. *44*, 89–115. 10.1146/annurev-environ-101718-
033302.

21. Paleczny, M., Hammill, E., Karpouzi, V., and Pauly, D. (2015). Population Trend of the World's
Monitored Seabirds, 1950-2010. PLOS ONE *10*, e0129342. 10.1371/journal.pone.0129342.

22. Dias, M.P., Martin, R., Pearmain, E.J., Burfield, I.J., Small, C., Phillips, R.A., Yates, O., Lascelles,
B., Borboroglu, P.G., and Croxall, J.P. (2019). Threats to seabirds: A global assessment. Biol.
Conserv. *237*, 525–537. 10.1016/j.biocon.2019.06.033.

23. Šúr, M., Bunbury, N., and Van De Crommenacker, J. (2013). Frigatebirds on Aldabra Atoll:
population census, recommended monitoring protocol and sustainable tourism guidelines. Bird
Conserv. Int. *23*, 214–220. 10.1017/S0959270913000087.

24. Burger, A.E., and Betts, M. (2001). Monitoring populations of Red-footed Boobies Sula sula and
frigatebirds Fregata spp. breeding on Aldabra Atoll, Indian Ocean. Bull. Br. Ornithol. Club *121*,
236–246.

25. Buelow, C., and Sheaves, M. (2015). A birds-eye view of biological connectivity in mangrove
systems. Estuar. Coast. Shelf Sci. *152*, 33–43. 10.1016/j.ecss.2014.10.014.

26. Young, H.S., Hurrey, L., and Kolb, G.S. (2011). Effects of Seabird-Derived Nutrients on Aquatic
Systems. In Seabird Islands, C. P. H. Mulder, W. B. Anderson, D. R. Towns, and P. J.
Bellingham, eds. (Oxford University Press), pp. 242–260.

27. Kazama, K. (2020). Bottom-Up Effects on Coastal Marine Ecosystems Due to Nitrogen Input from
Seabird Feces. Ornithol. Sci. *18*, 117. 10.2326/osj.18.126.

28. Signa, G., Mazzola, A., and Vizzini, S. (2021). Seabird influence on ecological processes in
coastal marine ecosystems: An overlooked role? A critical review. Estuar. Coast. Shelf Sci. *250*,
107164. 10.1016/j.ecss.2020.107164.

29. Polis, G.A., and Hurd, S.D. (1996). Linking marine and terrestrial food webs: allochtonous input
from the ocean supports high secondary productivity on small islands and coastal land
communities.pdf. Am. Nat. *147*, 396–423.

30. Diamond, A.W. (1971). The Ecology of the Sea Birds of Aldabra. Philos. Trans. R. Soc. Lond. B.
Biol. Sci. *260*, 561–571.

31. Haupt, P. (2019). Reef fish associations with benthic habitats at a remote protected coral reef
ecosystem in the Western Indian Ocean, Aldabra Atoll, Seychelles. PhD thesis, Rhodes
University, South Africa.

32. Pritchard, A., Sanchez, C., Bunbury, N., Burt, A., Currie, J., Doak, N., Fleischer-Dogley, F.,
Metcalfe, K., Mortimer, J., Richards, H., et al. (2022). Green turtle population recovery at Aldabra
Atoll continues after 50 yr of protection. Endanger. Species Res. *47*, 205–215. 10.3354/esr01174.

33. Pistorius, P.A., and Taylor, F.E. (2008). Population status and conservation of Crab Plovers
Dromas ardeola on Aldabra, Indian Ocean. Wader Study Group Bull. *115*, 36–40.

34. Heywood, K.J., Barton, E.D., and Simpson, J.H. (1990). The effects of flow disturbance by an
oceanic island. J. Mar. Res. *48*, 55–73. 10.1357/002224090784984623.

35. Duke, N.C., Lo, E., and Sun, M. (2002). Global distribution and genetic discontinuities of
mangroves – emerging patterns in the evolution of Rhizophora. Trees *16*, 65–79.
10.1007/s00468-001-0141-7.

36. Triest, L., Van Der Stocken, T., De Ryck, D., Kochzius, M., Lorent, S., Ngeve, M., Ratsimbazafy,
H.A., Sierens, T., Van Der Ven, R., and Koedam, N. (2021). Expansion of the mangrove species
Rhizophora mucronata in the Western Indian Ocean launched contrasting genetic patterns. Sci.
Rep. *11*, 4987. 10.1038/s41598-021-84304-8.

37. Simpson, L.T., Canty, S.W.J., Cissell, J.R., Steinberg, M.K., Cherry, J.A., and Feller, I.C. (2021).
Bird rookery nutrient over-enrichment as a potential accelerant of mangrove cay decline in Belize.
Oecologia *197*, 771–784. 10.1007/s00442-021-05056-w.

38. Castañeda-Moya, E., Rivera-Monroy, V.H., Chambers, R.M., Zhao, X., Lamb-Wotton, L., Gorsky,
A., Gaiser, E.E., Troxler, T.G., Kominoski, J.S., and Hiatt, M. (2020). Hurricanes fertilize
mangrove forests in the Gulf of Mexico (Florida Everglades, USA). Proc. Natl. Acad. Sci. *117*,
4831–4841. 10.1073/pnas.1908597117.

39. Worthington, T.A., Zu Ermgassen, P.S.E., Friess, D.A., Krauss, K.W., Lovelock, C.E., Thorley, J.,
Tingey, R., Woodroffe, C.D., Bunting, P., Cormier, N., et al. (2020). A global biophysical typology

of mangroves and its relevance for ecosystem structure and deforestation. Sci. Rep. *10*, 14652.
10.1038/s41598-020-71194-5.

40. Feller, I.C., Lovelock, C.E., and McKee, K.L. (2007). Nutrient Addition Differentially Affects
Ecological Processes of Avicennia germinans in Nitrogen versus Phosphorus Limited Mangrove
Ecosystems. Ecosystems *10*, 347–359. 10.1007/s10021-007-9025-z.

41. Alongi, D.M. (2018). Impact of Global Change on Nutrient Dynamics in Mangrove Forests.
Forests *9*, 596. 10.3390/f9100596.

42. Güsewell, S. (2004). N : P ratios in terrestrial plants: variation and functional significance. New
Phytol. *164*, 243–266. 10.1111/j.1469-8137.2004.01192.x.

43. Lovelock, C.E., Feller, I.C., Ball, M.C., Ellis, J., and Sorrell, B. (2007). Testing the Growth Rate
vs. Geochemical Hypothesis for latitudinal variation in plant nutrients. Ecol. Lett. *10*, 1154–1163.
10.1111/j.1461-0248.2007.01112.x.

44. Boto, K., and Wellington, J. (1983). Phosphorus and nitrogen nutritional status of a northern
Australian mangrove forest. Mar. Ecol. Prog. Ser. *11*, 63–69. 10.3354/meps011063.

45. Constance, A., Oehri, J., Bunbury, N., Wiesenberg, G.L.B., Pennekamp, F., A'Bear, L., Fleischer-
Dogley, F., and Schaepman-Strub, G. (2022). Soil nutrient content and water level variation drive
mangrove forest aboveground biomass in the lagoonal ecosystem of Aldabra Atoll. Ecol. Indic.
143, 109292. 10.1016/j.ecolind.2022.109292.

46. Von Hoyningen-Huene, A.J.E., Schneider, D., Fussmann, D., Reimer, A., Arp, G., and Daniel, R.
(2022). DNA- and RNA-based bacterial communities and geochemical zonation under changing
sediment porewater dynamics on the Aldabra Atoll. Sci. Rep. *12*, 4257. 10.1038/s41598-022-
07980-0.

47. Lin, Y.-M., Liu, X.-W., Zhang, H., Fan, H.-Q., and Lin, G.-H. (2010). Nutrient conservation
strategies of a mangrove species Rhizophora stylosa under nutrient limitation. Plant Soil *326*,
469–479. 10.1007/s11104-009-0026-x.

48. Feller, I.C., Whigham, D.F., O'Neill, J.P., and Mckee, K.L. (1999). Effects of Nutrient Enrichment
on within-Stand Cycling in a Mangrove Forest.

49. Feller, I.C., Whigham, D.F., McKee, K.L., and Lovelock, C.E. (2003). Nitrogen limitation of growth
and nutrient dynamics in a disturbed mangrove forest, Indian River Lagoon, Florida. Oecologia
134, 405–414. 10.1007/s00442-002-1117-z.

50. Feller, I.C., Lovelock, C.E., and Piou, C. (2009). Growth and Nutrient Conservation in Rhizophora
mangle in Response to Fertilization along Latitudinal and Tidal Gradients. Smithson. Contrib.
Mar. Sci. *38*.

51. Wei, L., Kao, S.-J., and Liu, C. (2019). Mangrove species maintains constant nutrient resorption
efficiency under eutrophic conditions. J. Trop. Ecol. *36*, 36–38. 10.1017/S0266467419000336.

52. Deegan, L.A., Johnson, D.S., Warren, R.S., Peterson, B.J., Fleeger, J.W., Fagherazzi, S., and
Wollheim, W.M. (2012). Coastal eutrophication as a driver of salt marsh loss. Nature *490*, 388–
392. 10.1038/nature11533.

53. Feller, I.C., Dangremond, E.M., Devlin, D.J., Lovelock, C.E., Proffitt, C.E., and Rodriguez, W.
(2015). Nutrient enrichment intensifies hurricane impact in scrub mangrove ecosystems in the
Indian River Lagoon, Florida, USA. Ecology *96*, 2960–2972. 10.1890/14-1853.1.

54. Chapin, F.S. (1980). The Mineral Nutrition of Wild Plants. Annu. Rev. Ecol. Syst. *11*, 233–260.
10.1146/annurev.es.11.110180.001313.

55. Tilman, D. (1990). Constraints and Tradeoffs: Toward a Predictive Theory of Competition and
Succession. Oikos *58*, 3. 10.2307/3565355.

56. Huxham, M., Langat, J., Tamooh, F., Kennedy, H., Mencuccini, M., Skov, M.W., and Kairo, J.
(2010). Decomposition of mangrove roots: Effects of location, nutrients, species identity and mix
in a Kenyan forest. Estuar. Coast. Shelf Sci. *88*, 135–142. 10.1016/j.ecss.2010.03.021.

57. Hayes, M.A., Jesse, A., Tabet, B., Reef, R., Keuskamp, J.A., and Lovelock, C.E. (2017). The
contrasting effects of nutrient enrichment on growth, biomass allocation and decomposition of
plant tissue in coastal wetlands. Plant Soil *416*, 193–204. 10.1007/s11104-017-3206-0.

58. Constance, A., Haverkamp, P.J., Bunbury, N., and Schaepman-Strub, G. (2021). Extent change
of protected mangrove forest and its relation to wave power exposure on Aldabra Atoll. Glob.
Ecol. Conserv. *27*, e01564. 10.1016/j.gecco.2021.e01564.

59. Allgeier, J.E., Layman, C.A., Mumby, P.J., and Rosemond, A.D. (2014). Consistent nutrient
storage and supply mediated by diverse fish communities in coral reef ecosystems. Glob. Change
Biol. *20*, 2459–2472. 10.1111/gcb.12566.

60. Benkwitt, C.E., Wilson, S.K., and Graham, N.A.J. (2019). Seabird nutrient subsidies alter patterns
of algal abundance and fish biomass on coral reefs following a bleaching event. Glob. Change
Biol. *25*, 2619–2632. 10.1111/gcb.14643.

61. Allgeier, J.E., Layman, C.A., Montaña, C.G., Hensel, E., Appaldo, R., and Rosemond, A.D.
(2018). Anthropogenic versus fish-derived nutrient effects on seagrass community structure and
function. Ecology *99*, 1792–1801. 10.1002/ecy.2388.

62. Sandoval, L., Mancera-Pineda, J., Leal-Flórez, J., Blanco-Libreros, J., and Delgado-Huertas, A.
(2022). Mangrove carbon sustains artisanal fish and other estuarine consumers in a major
mangrove area of the southern Caribbean Sea. Mar. Ecol. Prog. Ser. *681*, 21–35.
10.3354/meps13910.

63. Gagnon, K., Rothäusler, E., Syrjänen, A., Yli-Renko, M., and Jormalainen, V. (2013). Seabird
Guano Fertilizes Baltic Sea Littoral Food Webs. PLoS ONE *8*, e61284.
10.1371/journal.pone.0061284.

64. Zmudczyńska-Skarbek, K., Balazy, P., and Kuklinski, P. (2015). An assessment of seabird
influence on Arctic coastal benthic communities. J. Mar. Syst. *144*, 48–56.
10.1016/j.jmarsys.2014.11.013.

65. Vizzini, S., Signa, G., and Mazzola, A. (2016). Guano-Derived Nutrient Subsidies Drive Food
Web Structure in Coastal Ponds. PLOS ONE *11*, e0151018. 10.1371/journal.pone.0151018.

66. Lorrain, A., Houlbrèque, F., Benzoni, F., Barjon, L., Tremblay-Boyer, L., Menkes, C., Gillikin, D.P.,
Payri, C., Jourdan, H., Boussarie, G., et al. (2017). Seabirds supply nitrogen to reef-building
corals on remote Pacific islets. Sci. Rep. *7*, 3721. 10.1038/s41598-017-03781-y.

67. Stevens, J.D. (1984). Life-history and ecology of sharks at Aldabra Atoll, Indian Ocean. Proc. R.
Soc. Lond. B Biol. Sci. *222*, 79–106. 10.1098/rspb.1984.0050.

68. Friedlander, A.M., Ballesteros, E., Beets, J., Brown, E.K., Fay, J.M., Haupt, P., Henning, B.,
Rose, P., and Sala, E. (2015). Biodiversity and ecosystem health of the Aldabra group, Southern
Seychelles.pdf.

69. Koester, A., Gordó–Vilaseca, C., Bunbury, N., Ferse, S.C.A., Ford, A., Haupt, P., A'Bear, L.,
Bielsa, M., Burt, A.J., Letori, J., et al. (2023). Impacts of coral bleaching on reef fish abundance,
biomass and assemblage structure at remote Aldabra Atoll, Seychelles: insights from two survey
methods. Front. Mar. Sci. *10*, 1230717. 10.3389/fmars.2023.1230717.

70. Sandin, S.A., Becker, P.A., Becker, C., Brown, K., Erazo, N.G., Figuerola, C., Fisher, R.N.,
Friedlander, A.M., Fukami, T., Graham, N.A.J., et al. (2022). Harnessing island–ocean
connections to maximize marine benefits of island conservation. Proc. Natl. Acad. Sci. *119*,
e2122354119. 10.1073/pnas.2122354119.

71. Stoddart, D.R., Taylor, J.D., Fosberg, F.R., and Farrow, G.E. (1971). Geomorphology of Aldabra
Atoll. Philos. Trans. R. Soc. Lond. B Biol. Sci. *260*, 31–65.

72. Hamylton, S., Hagan, A., Bunbury, N., Fleischer-Dogley, F., and Spencer, T. (2018). Mapping the
Lagoon at Aldabra Atoll, Western Indian Ocean. Atoll Res. Bull. *619*, 45–59.

73. Walton, R., Baxter, R., Bunbury, N., Hansen, D., Fleischer-Dogley, F., Greenwood, S., and
Schaepman-Strub, G. (2019). In the land of giants: habitat use and selection of the Aldabra giant
tortoise on Aldabra Atoll. Biodivers. Conserv. *28*, 3183–3198. 10.1007/s10531-019-01813-9.

74. Macnae, W. (1971). Mangroves on Aldabra. Philos. Trans. R. Soc. Lond. B Biol. Sci. *260*, 237–
247. 10.1098/rstb.1971.0013.

75. Farrow, G.E., and Brander, K.M. (1971). Tidal Studies on Aldabra. Philos. Trans. R. Soc. Lond. B.
Biol. Sci. *260*, 93–121.

76. Haverkamp, P.J., Shekeine, J., De Jong, R., Schaepman, M., Turnbull, L.A., Baxter, R., Hansen,
D., Bunbury, N., Fleischer-Dogley, F., and Schaepman-Strub, G. (2017). Giant tortoise habitats
under increasing drought conditions on Aldabra Atoll—Ecological indicators to monitor rainfall
anomalies and related vegetation activity. Ecol. Indic. *80*, 354–362.
10.1016/j.ecolind.2017.05.029.

77. Kjeldahl, J. (1883). Neue Methode zur Bestimmung des Stickstoffs in organischen Körpern.
Fresenius Z. Für Anal. Chem. *22*, 366–382. 10.1007/BF01338151.

78. Riddick, S.N., Dragosits, U., Blackall, T.D., Daunt, F., Wanless, S., and Sutton, M.A. (2012). The
global distribution of ammonia emissions from seabird colonies. Atmos. Environ. *55*, 319–327.
10.1016/j.atmosenv.2012.02.052.

79. Young, H.S., McCauley, D.J., Dunbar, R.B., and Dirzo, R. (2010). Plants cause ecosystem
nutrient depletion via the interruption of bird-derived spatial subsidies. Proc. Natl. Acad. Sci. *107*,
2072–2077. 10.1073/pnas.0914169107.

80. Staunton Smith, J., and Johnson, C. (1995). Nutrient inputs from seabirds and humans on a
populated coral cay. Mar. Ecol. Prog. Ser. *124*, 189–200. 10.3354/meps124189.

81. Kauffman, J.B., and Donato, D.C. (2012). Protocols for the measurement, monitoring and
reporting of structure, biomass and carbon stocks in mangrove forests (Center for International
Forestry Research (CIFOR)) 10.17528/cifor/003749.

82. Chave, J., Andalo, C., Brown, S., Cairns, M.A., Chambers, J.Q., Eamus, D., Fölster, H., Fromard,
F., Higuchi, N., Kira, T., et al. (2005). Tree allometry and improved estimation of carbon stocks
and balance in tropical forests. Oecologia *145*, 87–99. 10.1007/s00442-005-0100-x.

83. Chapin, F.S., and Cleve, K.V. (1989). Approaches to studying nutrient uptake, use and loss in
plants. In Plant physiological ecology: field methods and instrumentation. (Springer Netherlands),
p. 185–207.

84. Donovan, M.K., Adam, T.C., Shantz, A.A., Speare, K.E., Munsterman, K.S., Rice, M.M., Schmitt,
R.J., Holbrook, S.J., and Burkepile, D.E. (2020). Nitrogen pollution interacts with heat stress to
increase coral bleaching across the seascape. Proc. Natl. Acad. Sci. *117*, 5351–5357.
10.1073/pnas.1915395117.

85. Caut, S., Angulo, E., Pisanu, B., Ruffino, L., Faulquier, L., Lorvelec, O., Chapuis, J.-L., Pascal,
M., Vidal, E., and Courchamp, F. (2012). Seabird Modulations of Isotopic Nitrogen on Islands.
PLoS ONE *7*, e39125. 10.1371/journal.pone.0039125.

86. Diamond, A.W. (1979). Dynamic Ecology of Aldabran Seabird Communities. Philos. Trans. R.
Soc. Lond. B. Biol. Sci. *286*, 231–240.

87. Aminot, A., and Kérouel, R. (2007). Dosage automatique des nutriments dans les eaux marines :
méthodes en flux continu (Ifremer). Editions Quae.

88. Zuur, A.F., and Ieno, E.N. (2016). A protocol for conducting and presenting results of regression-
type analyses. Methods Ecol. Evol. *7*, 636–645. 10.1111/2041-210X.12577.

89. Lenth, R.V. (2023). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package
version 1.8.6.

90. Pinheiro, J., Bates, D., Debroy, S., Sarkar, D., and R Core Team (2023). Nlme: linear and
nonlinear mixed effects models. R package version 3.1-162.

91. R Core Team (2022). R: A language and environment for statistical language and environment for
statistical computing. (R Foundation for Statistical Computing).

 FIGURE LEGENDS

Figure 1. Nitrogen isotopic values in mangrove abiotic and biotic (trophic) components at

seabird and non-seabird sites

$\delta^{15}N$ values of portunid crabs *Thalamita crenata* (A), sesarmid crabs *Sesarma leptosoma* (B),

gastropods *Littoraria* spp. (C), mangrove leaves *Rhizophora mucronata* (D) and mangrove sediment

(E) at seabird and non-seabird breeding sites on Aldabra Atoll. Black points and error bars display

predicted means ± SD of linear mixed models and green and purple points display raw data. Error

bars not visible in some cases because of scaling. Trophic level of each biotic component is indicated

by the triangular icon in the top right of each plot.

Figure 2. Nutrient parameters in mangrove leaves at seabird and non-seabird sites

Nutrient concentrations, %N (A) and %P (B), nutrient ratios C:N (C) and C:P (D), and nutrient

resorption efficiencies RE-N (E) and RE-P (F) of mangrove leaves *Rhizophora mucronata* at seabird

and non-seabird breeding sites on Aldabra Atoll. Black points and error bars display predicted means

± SD of linear mixed models and green and purple points display raw data. Error bars not visible in

some cases because of scaling. RE-N: nitrogen resorption efficiency, RE-P: phosphorus resorption

efficiency.

Figure 3. Nutrient levels in abiotic and biotic components adjacent to mangroves at seabird

and non-seabird sites

Nutrient concentrations in surface seawater, NOx (A) and phosphate (B) during incoming and

outgoing tides, and $\delta^{15}N$ values of macroalgae *Halimeda* spp. (C) adjacent to mangroves, at seabird

and non-seabird breeding sites on Aldabra Atoll. Black points and error bars display predicted means

± SD of linear mixed models and green and purple points display raw data. NOx = nitrate + nitrite.

Figure 4. Summary figure illustrating the transfer of seabird-derived nutrients in mangrove

forests

Seabirds forage at sea and deposit nutrient-rich guano in their mangrove breeding colonies. Seabird-

derived nutrients enrich mangroves and associated invertebrate fauna, and are exported to adjacent

habitats by tidal flow. Nutrient enrichment are indicated by plus (+) signs.

FIGURES

Figure 1.

Figure 2.

Figure 3.

Figure 4.