1

Male germ cell-associated kinase (MAK) is required for axoneme formation during ciliogenesis in zebrafish photoreceptors

Hung-Ju Chiang[1], Yuko Nishiwaki[1], Wei-Chieh Chiang[1], and Ichiro Masai[1,2]

[1]Developmental Neurobiology Unit, Okinawa Institute of Science and Technology Graduate University, Tancha, Okinawa, Japan.

[2] Correspondence should be addressed (masai@oist.jp)

Keywords: zebrafish, photoreceptor degeneration, MAK, ciliopathy, ciliogenesis

Summary statement
Male germ cell-associated kinase (MAK) is a cilium-associated serine/threonine kinase that promotes axoneme development during ciliogenesis in zebrafish photoreceptors to ensure intracellular protein transport and photoreceptor survival.

Abstract

Vertebrate photoreceptors are highly specialized retinal neurons that have cilium-derived membrane organelles called outer segments (OS), which function as platforms for phototransduction. Male germ cell-associated kinase (MAK) is a cilium-associated serine/threonine kinase, and its genetic mutation causes photoreceptor degeneration in mice and retinitis pigmentosa in humans. However, the role of MAK in photoreceptors is not fully understood. Here, we report that zebrafish *mak* mutants show rapid photoreceptor degeneration during embryonic development. In *mak* mutants, both cone and rod photoreceptors completely lack OSs and undergo apoptosis. Interestingly, zebrafish *mak* mutants fail to generate axonemes during photoreceptor ciliogenesis, whereas basal bodies are specified. These data suggest that MAK contributes to axoneme development in zebrafish, in contrast to mouse *Mak* mutants, which have elongated photoreceptor axonemes. Furthermore, the kinase activity of MAK is critical in ciliary axoneme development and photoreceptor survival. Thus, MAK is required for ciliogenesis and OS formation in zebrafish photoreceptors to ensure intracellular protein transport and photoreceptor survival.

Introduction

The vertebrate photoreceptor is highly compartmentalized to form specialized structures related to phototransduction (Fain et al., 2010). The outer segment (OS) is a specialized cilium of the photoreceptor, in which multiple photoreceptive membrane discs are regularly stacked to accommodate phototransduction molecules such as rhodopsin and opsins (Bachmann-Gagescu and Neuhauss, 2019; Chen et al., 2021). The inner segment (IS) is a mitochondria-enriched region between the OS and the nucleus. Finally, photoreceptors have a specialized synaptic structure in the most basal region that mediates transmission of electric signals to horizontal cells and bipolar cells (Mercer and Thoreson, 2011).

The photoreceptor OS is a highly specialized primary cilium that consists of stacked membrane discs around a microtubule-based backbone called the axoneme (Bachmann-Gagescu and Neuhauss, 2019; Chen et al., 2021). The axoneme is anchored to the basal body in the IS of photoreceptors and extends apically through the connecting cilium. The connecting cilium bridges the IS and the OS, through which phototransduction molecules are transported from the ER to the Golgi and then into the OS. Importantly, the connecting cilium is equivalent to the ciliary transition zone in other cell types and functions as a gating system (Park and Leroux, 2022), through which OS resident proteins are transported to the OS by the intraflagellar transport (IFT) complex (Klena and Pigino, 2022) and BBSome, composed of eight Bardet-Biedl syndrome (BBS) proteins (Tian et al., 2023).

The process of ciliary development is called ciliogenesis, which consists primarily of three steps: centriole (basal body) docking to the apical plasma membrane, establishment of a transition zone, and ciliary axoneme extension (Breslow and Holland, 2019; Chen et al., 2021). The first step of ciliogenesis is removal of CP110 from a distal appendage of the mother centriole, which enables ARL13b and components of IFT-A and IFT-B to assemble at the apical surface of a distal appendage in a rab8-dependent manner (Breslow and Holland, 2019; Ge et al., 2022). The second step is formation of the transition zone by transport of MKS and NPHP module components (Park and Leroux, 2022). The third step is extension of the axoneme through IFT-mediated ciliary transport, which transports and adds α-/β-tubulin dimers to the plus end of the microtubule comprising the axoneme

(Brouhard and Rice, 2018; Ge et al., 2022; Nakayama and Katoh, 2018). Defects in

ciliogenesis cause ciliopathy, which is associated with various abnormalities during

organogenesis, including retinal dystrophies, cystic kidney diseases, skeletal dysplasia, and

polydactyly (Mill et al., 2023). Although many factors are reportedly involved in

photoreceptor ciliogenesis and ciliopathy (Cideciyan et al., 2007; den Hollander et al., 2006;

Gao et al., 2002; Gorden et al., 2008; Kakakhel et al., 2020; Rao et al., 2015; Zhu et al.,

2021), the underlying mechanisms are not fully understood.

Male germ cell-associated kinase (MAK) belongs to the MAK/ICK/MOK

serine/threonine kinase family (Chen et al., 2013). MAK was identified by cross-

hybridization with tyrosine kinase v-ros in rat testicular cells, so it was considered to be a

spermatogenesis regulator (Matsushime et al., 1990). MAK functions as a coactivator of

androgen receptors (ARs) to promote AR-mediated signaling, which is associated with

prostate tumorigenesis (Ma et al., 2006; Wang and Kung, 2012; Xia et al., 2002).

Furthermore, *Mak* transcripts are expressed in the photoreceptor layer in mouse retina

(Blackshaw et al., 2004) and MAK protein is localized in cilia of photoreceptors (Omori et

al., 2010). In *Mak* knockout mice, photoreceptors show abnormally elongated axonemes

and malformation of membrane discs in the OS, leading to photoreceptor degeneration

(Omori et al., 2010). In humans, *MAK* mutations cause retinitis pigmentosa (RP), in which

patients progressively lose their vision (Ozgul et al., 2011; Stone et al., 2011). However,

human patients carrying *MAK* mutations do not show elongation of the photoreceptor layer

(Stone et al., 2011), although fibroblasts derived from RP patients carrying *MAK* mutations

have elongated cilia (Tucker et al., 2022). Thus, it is important how *mak* mutations affect

ciliary regulation in photoreceptors among vertebrate species.

In this study, we found that zebrafish photoreceptors of *mak* mutants degenerate

during embryonic development. In *mak* mutants, both rods and cones undergo apoptosis,

although their degeneration processes differ. Interestingly, *mak* mutants fail to form

axonemes in photoreceptor cilia, whereas basal bodies are specified, which is the opposite

of mouse *Mak* knockout photoreceptors, with elongated ciliary axonemes. Furthermore,

both cones and rods completely lack OSs in *mak* mutants, leading to ectopic distribution of

opsins. So, MAK is essential to form axonemes and OSs in photoreceptors. Finally, MAK

kinase activity is critical for axoneme formation and photoreceptor survival. Thus, MAK is

essential for ciliogenesis, OS formation, and photoreceptor survival.

Results

Zebrafish *pday* mutants carry a mutation in *mak* gene, leading to photoreceptor degeneration

Zebrafish *payday* (*pday*) mutants display no visual response during the embryonic stage (Muto et al., 2005). We selected *pday* homozygous mutant embryos using an optokinetic response (OKR) assay and examined their phenotypes. At 6 days post-fertilization (dpf), *pday* mutants showed no apparent morphological defect (Fig. 1A), but did not survive beyond 10 dpf, suggesting a lethal mutation. However, under intensive feeding, less than 1% of *pday* mutants survived until 2 months post-fertilization (mpf), displaying scoliosis, cardiac edema, and smaller body size (Fig. 1A).

Next, we mapped the *pday* mutant locus with simple sequence-length polymorphism (SSLP) markers (Knapik et al., 1998; Shimoda et al., 1999). The *pday* mutation was mapped between two SSLP markers z23011 (1/186 miosis) and z13695 (3/186 miosis), so the mutant gene is restricted to a genomic region between 8.517 and 9.123 Mb on chromosome 24. In this genomic region, 10 genes including *mak* were annotated in the zebrafish genomic database (GRCz10, Ensemble release 80). Since *Mak* mutations cause photoreceptor degeneration in mice (Omori et al., 2010) and humans (Ozgul et al., 2011; Stone et al., 2011), and because the other 9 genes were not reported to be involved in retinal development, degeneration, and functions, we focused on the *mak* gene. First, we designed a new polymorphic marker, called Mak-N1, located around 20 bp upstream of the first exon of the *mak* gene. The recombination rate of Mak-N1 for *pday* mutation was 0 of 186 meiosis, suggesting that *mak* is a strong candidate. Next, we cloned *mak* cDNA prepared from wild-type embryos, and found that *mak* cDNA comprises 15 exons (Fig. 1B) and that there are two isoforms of *mak* mRNA. One has and the other lacks exon 14 due to alternative splicing. Then, we focused on the longest isoform, which has exon 14 and is annotated as the full-length isoform in the zebrafish genomic database (mak-201, GRCz11, Ensemble release 110). We compared amino acid sequences of this longest zebrafish MAK isoform with those of human and mouse MAK. Amino acid identity between zebrafish MAK and human/mouse MAK was more than 50% for full-length

protein and more than 86% for the N-terminal kinase domain (Fig. S1). We cloned *mak* cDNAs from *pday* mutant mRNA and genomic fragments covering coding exons from the *pday* mutant genome. Sequencing revealed that a non-sense mutation occurs in exon 4 of the *mak* gene in *pday* mutants, which causes a premature termination codon at 70R (Fig. 1B). These data suggest that the *pday* mutant gene encodes MAK.

Next, we examined retinal phenotypes of wild-type and *pday* mutant embryos from 3 to 6 dpf (Fig. 1C, 1C'). In wild-type retinas at 3 dpf, photoreceptors differentiate to form the outer nuclear layer (ONL). The OS is formed in the apical ONL (Fig. 1C'). In *pday* mutants, the ONL was formed; however, densely labeled nuclei were observed in the basal region of the ventro-central ONL (Fig. 1C, 1C'), and the OS was markedly reduced (Fig. 1C'). In wild-type photoreceptors at 4 dpf, the OS was elongated. However, in *pday* mutants, the ONL was decreased in thickness (Fig. 1C, 1C'), and no OS-like structure was observed (Fig. 1C'). In addition, there were pyknotic nuclei and bubble-like structures (Fig. 1C, 1C'). In wild-type embryos at 5 dpf, the ONL was divided into two subnuclear layers (Fig. 1C'), the apical and basal regions of which contain nuclei of blue/red/green cones and rods/UV cones, respectively (D'Orazi et al., 2020; Oel et al., 2020). However, in *pday* mutants, the dorso-central ONL drastically decreased in thickness (Fig. 1C) and contained flattened nuclei (Fig. 1C, 1C'). In wild-type embryos at 6 dpf, photoreceptors showed more mature shapes in the ONL. However, in *pday* mutants, most photoreceptor nuclei disappeared in the dorso-central retina (Fig. 1C), so there are very flat nuclei in the outermost region of the retina, which are difficult to define as the ONL or horizontal cells (Fig. 1C'). Interestingly, in *pday* mutants, ONL thickness was less affected in the retinal region more ventral to the optic nerve head, even at 6 dpf (Fig. 1C). Thus, sensitivity of photoreceptor degeneration differs between the dorso-central and the ventral retina. In addition, the inner nuclear layer (INL), the inner plexiform layer (IPL), and the retinal ganglion cell layer (RGCL) seemed to be intact in *pday* mutants from 3 to 6 dpf (Fig. 1C), suggesting photoreceptor-specific degeneration in *pday* mutants.

To confirm photoreceptor degeneration in mutant retinas at 6 dpf, we introduced a transgenic line *Tg[crx:eGFP]* that expresses eGFP in photoreceptor precursors and mature

rod and cone photoreceptors under control of the *cone-rod homeobox* (*crx*) promoter

(Suzuki et al., 2013) (Fig. 1D). In wild-type retinas, eGFP signals were detected in the

ONL at 6 dpf. However, in *pday* mutant retinas, eGFP signals disappeared in the dorso-

central retina (Fig. 1D). We measured thicknesses of eGFP-positive areas along the dorso-

ventral axis (Fig. S2A), and found that photoreceptor degeneration is more severe in the

dorso-central retina than in the ventral retina (Fig. 1E). Thus, photoreceptors undergo

degeneration in *pday* mutants.

Both rods and cones degenerate in *pday* mutants

In human *MAK*-associated RP patients, rods degenerate more prominently (Ozgul et al.,

2011; Stone et al., 2011). To evaluate rod and cone degeneration separately, we generated

double transgenic lines, *Tg[rho:NLS-eGFP; gnat2:NLS-tdTomato]*, to visualize rods and

cones with eGFP and tdTomato fluorescence, respectively, and introduced them into *pday*

mutants (Fig. 2, Fig. S3A). In wild-type retinas, eGFP expression in rods was strong and

dense in the ventral ONL but relatively weak and sparse in the dorso-central ONL at 3 dpf

(Fig. S3A). On the other hand, eGFP expression was very faint in *pday* mutant retinas

under the same conditions. However, when laser intensity was increased, eGFP expression

was detected (Fig. S3A), suggesting that rods are maintained in 3 dpf *pday* mutant retinas,

although the rhodopsin promoter does not effectively drive *eGFP* mRNA or else translation

of *eGFP* mRNA to eGFP protein was affected in *pday* mutants. Therefore, in later

experiments using *Tg[rho:NLS-eGFP]*, we applied higher laser intensity to *pday* mutant

scanning, and compared rod phenotypes between *pday* mutants and wild-type siblings.

Since photoreceptor degeneration is more severe in the dorso-central retina than

the ventral retina at 6 dpf (Fig. 1D, 1E), we visualized morphology of rods and cones using

Tg[rho:NLS-eGFP; gnat2:NLS-tdTomato] from 3 to 6 dpf (Fig. 2A, 2B), and focused on

the dorsal and ventral ONL regions (Fig. 2C-F). First, we examined rods (Fig. 2A). In wild-

type retinas at 3 dpf, rods were sparse in the dorsal ONL (Fig. 2C), but densely produced in

the ventral ONL (Fig. 2D). In *pday* mutants, many rods show very round nuclei in both the

dorsal and ventral ONL (Fig. 2C and 2D). Interestingly, eGFP signals were predominantly

localized in nuclei of wild-type rods because eGFP was tagged with a nuclear localization

sequence (NLS); however, eGFP signals were often excluded from the round nuclei of *pday*

mutant rods, suggesting apoptotic-like pyknotic nuclei of *pday* mutant rods. After 4 dpf, in

wild-type retinas, rods progressively increased in number and aligned along the dorsal ONL.

Their nuclei are positioned in the most basal region of the ONL, and rods extend processes

to form the OS in the apical ONL (Fig. 2C). On the other hand, rods are more densely

differentiated in the ventral ONL (Fig. 2D). However, in *pday* mutants at 4 dpf, rods almost

disappeared in both the dorsal and ventral ONL. Only a few rods remained, but displayed

pyknotic nuclei or were excluded from the ONL (Fig. 2C and 2D). In *pday* mutants at 5 and

6 dpf, rods disappeared in the ONL (Fig. 2C and 2D), except in the peripheral retina

adjacent to the ciliary marginal zone (CMZ) where new photoreceptors continued to be

produced by retinal stem cells. So, rods are eliminated soon after they are specified from

the CMZ. Evaluation of numbers of rod nuclei along the DV axis of the retina confirmed

that rods were progressively diminished in the central ONL of *pday* mutant retinas and

were completely eliminated, except in the CMZ area at 6 dpf (Fig. S3B).

207   Next, we examined cones (Fig. 2B). In wild-type retinas at 3 dpf, cones

differentiated uniformly in the dorsal ONL (Fig. 2E); however, cone density became

progressively sparse toward the CMZ region in the ventral ONL (Fig. 2F), a pattern

complementary to that of rods (Fig. 2D). In *pday* mutant retinas, cones normally

differentiated to form a spatial pattern similar to that of wild-type retinas (Fig. 2E, 2F).

However, there were tdTomato-negative rounded nuclei in both the dorsal and ventral ONL

(Fig. 2E and 2F), whose positions are identical to pyknotic-like rod nuclei (Fig. 2C and 2D).

Thus, rods lose their normal localization in the basal ONL in *pday* mutants at 3 dpf. After 4

dpf, in wild-type retinas, cones form two subnuclear layers in the dorso-central ONL (Fig.

2E). However, cone density was complementary to rod density and became progressively

sparse toward the CMZ region in the ventral ONL (Fig. 2F). In *pday* mutants at 4 dpf,

segregation of two cone subnuclear layers was incomplete in the dorso-central ONL (Fig.

2E). Furthermore, cones were sparse and shortened in the ventral ONL (Fig. 2F). In *pday*

mutants at 5 dpf, cones were shortened in the dorsal ONL (Fig. 2E). Interestingly, cone

density increased toward the CMZ in the ventral ONL of *pday* mutants (Fig. 2F), where

rods failed to be maintained (Fig. 2D). In *pday* mutants at 6 dpf, cones were extremely

flattened and there were gaps between them in the dorsal ONL (Fig. 2E), indicating that

cones had been eliminated. Furthermore, cones were similarly flattened in the ventro-

central ONL (Fig. 2F), whereas cones were uniformly produced and less flattened toward

the CMZ of the ventral ONL (Fig. 2F). Therefore, rods undergo acute apoptosis-like cell

death in *pday* mutants from 3 to 4 dpf, whereas cones undergo progressive shrinkage after 4

dpf and eventually disappear by 6 dpf in *pday* mutants (Fig. 2G), although cone

degeneration is less severe in the peripheral region of the ventral ONL.

Both rods and cones undergo apoptosis in *pday* mutants

Condensed rod nuclei in *pday* mutants at 3 dpf (Fig. 2C, 2D) are reminiscent of apoptosis,

in which chromatin condensation is a hallmark (Oberhammer et al., 1994). We performed

TdT-mediated dUTP-nick-end labeling (TUNEL) to *pday* mutant retinas at 3, 4, and 5 dpf

(Fig. 3A, 3A', Fig. S4A). In wild-type retinas, the number of TUNEL-positive cells was

continuously very low in the ONL (Fig. 3A, Fig. S4A). In *pday* mutant retinas at 3 dpf,

TUNEL signals were detected in the central and ventral ONL (Fig. 3A'), which is

consistent with our observation that rod pyknotic nuclei were detected at 3 dpf (Fig. 2C,

2D), suggesting that these early TUNEL-positive cells are likely to be rods. In *pday* mutant

retinas at 4 and 5 dpf, TUNEL signals increased in the dorsal and central ONL (Fig. 3A').

Interestingly, TUNEL-positive cells were positioned in a very thin central ONL of 5 dpf

pday mutant retinas (Fig. 3A'), suggesting that these late TUNEL-positive cells are likely

to be cones. TUNEL-positive cells were more numerous in *pday* mutant ONL than wild-

type ONL at 3 dpf, but the difference was not significant. However, the number of TUNEL-

positive cells was the highest in *pday* mutant ONL at 4 dpf and was still significantly

higher in *pday* mutant ONL at 5 dpf, compared with wild-type ONL (Fig. S4A). The

number of TUNEL signals in other retinal layers, INL, and RGCL, was not significantly

different between *pday* mutants and wild-type siblings at stages 3, 4, or 5 dpf (Fig. S4B,

S4C), suggesting that apoptosis is specific to photoreceptors.

Next, we examined whether cones undergo apoptosis in *pday* mutants. First, we

carried out double labeling of 5-dpf, wild-type, and *pday* mutant retinas with anti-zebrafish

rhodopsin (Vihtelic et al., 1999) and zpr1 (Larison and Bremiller, 1990) antibodies, which

label rod OSs and double-cone-type photoreceptors (red/green cones), respectively (Fig.

3B). Consistent with data of *Tg[rho:NLS-eGFP; gnat2:NLS-tdTomato]* (Fig. 2), rods were

only detected in the peripheral region of the dorsal and ventral ONL of *pday* mutants (Fig.

3B'). On the other hand, cones were observed in the peripheral region of the dorsal and

ventral ONL and as very flattened cells in the central ONL (Fig. 3B'). Next, we carried out

TUNEL of 5-dpf, wild-type and *pday* mutant retinas combined with zpr1 antibody labeling

(Fig. 3C). Many TUNEL-positive cells were observed in dorsal, central and ventral ONL of

pday mutants (Fig. 3C'). A significant fraction of these TUNEL-positive cells was

associated with zpr1 signals (Fig. 3C'). Third, we carried out TUNEL in 5-dpf, wild-type

and *pday* mutant retinas combined with anti-rhodopsin antibody labeling (Fig. 3D). Many

TUNEL-positive cells were observed in dorsal, central, and ventral ONL of *pday* mutants

(Fig. 3D'). However, rhodopsin signals were detected only in the CMZ of the dorsal and

ventral ONL (Fig. 3D'), but not in the central ONL. Thus, many TUNEL signals were not

associated with rhodopsin signals in the central ONL (Fig. 3D'). Since flattened

photoreceptors in the central ONL of 5 dpf *pday* mutants are cones (Fig. 2), cones undergo

apoptosis at or before 5 dpf in *pday* mutants.

To confirm that both rods and cones undergo apoptosis in *pday* mutants, we

combined a transgenic line *Tg[hsp:mCherry-Bcl2]*, which expresses an anti-apoptotic

protein Bcl2 (Nishiwaki and Masai, 2020), with *pday* mutants and evaluated cell survival

by labeling with zpr1 (Larison and Bremiller, 1990) and zpr3 (Gao et al., 2022) antibodies,

which visualize double cone photoreceptors and rhodopsin/green opsin, respectively.

Combined with quantitative analyses (Fig. S2A, S2B), we found that Bcl2 overexpression

significantly rescued survival of both cones and rods in *pday* mutants at 6 dpf (Fig. 3E, 3F;

Fig. S4D, S4E) even along the DV axis (Fig. S4F), suggesting that both cones and rods

undergo apoptosis in *pday* mutants.

Although rods and cones survived in *pday* mutants that overexpressed Bcl2, zpr3
signals fail to be localized in the OS, and spread in the plasma membrane throughout the
cell body, including the basal synaptic region (Fig. 3F'), suggesting that green opsin and
rhodopsin are mislocalized in Bcl2-overexpressing *pday* mutant photoreceptors.
Furthermore, photoreceptor nuclear shape differed between Bcl2-overexpressing *pday*
mutants and wild-type siblings (Fig. S4G). Next, we examined OKR in embryos produced
by crosses of *pday* heterozygous fish. On average, 25% of embryos were homozygous for
the *pday* mutation and did not show OKR. Bcl2 overexpression did not restore OKR of
pday mutants (Fig. S4H). Thus, MAK activity is primarily required for visual functions of
photoreceptors, disruption of which may subsequently cause Bax-mediated apoptosis.
288
***mak* is responsible for *pday* mutant phenotypes**
To validate whether the *mak* gene is responsible for *pday* mutant phenotypes, we generated
a zebrafish transgenic line *Tg[crx:eGFP-MAK],* which expresses the N-terminal eGFP-
tagged, full-length isoform of MAK under control of the *crx* promotor. A transgenic line
Tg[crx:eGFP] was used as a control. Labeling with zpr1 antibody revealed that
Tg[crx:eGFP-MAK] rescued cone photoreceptor degeneration in *pday* mutants at 6 dpf,
whereas *Tg[crx:eGFP]* did not (Fig. 4A, 4A', and 4B). Furthermore, we examined OKR in
embryos produced by crosses of *pday* heterozygous fish. On average, 25% of embryos were
homozygous for the *pday* mutation and did not show OKR. However, *Tg[crx:eGFP-MAK]*
restored OKR of *pday* mutants at 6 dpf, whereas *Tg[crx:eGFP]* did not (Fig. 4C). Thus,
mak is responsible for *pday* mutant phenotypes: photoreceptor degeneration, and visual
response defects.
301
302   **MAK is localized in the connecting cilium and promotes axoneme development during**
303   **ciliogenesis**
304   We examined *mak* mRNA expression during zebrafish embryonic development by whole-
305   mount *in situ* hybridization (Fig. 5A). *mak* mRNA is ubiquitously expressed from 1- to 4-
306   cell stages, indicating maternal mRNA expression. *mak* mRNA expression disappeared at

the 50%-epiboly stage, but was detected in Kupffer's vesicle, a ciliated embryonic organ that is important for left-right asymmetry patterning (Essner et al., 2005), at the tail-bud stage. At 24 hpf, *mak* mRNA is expressed in the central nervous system, and especially strongly expressed in the epiphysis and retina. At 48 hpf, *mak* mRNA is prominently expressed in the brain, the retina, and the pectoral fin bud. At 72 hpf, *mak* mRNA is highly expressed in the brain, including the optic tectum and the photoreceptor layer of the retina. We also confirmed that *mak* mRNA expression was markedly decreased in *pday* mutant embryos at 72 hpf, probably due to nonsense-mediated mRNA decay (Fig. S5A). Thus, *mak* mRNA is expressed in various cell types in zebrafish embryos, including Kupffer's vesicle and retinal and pineal photoreceptors, in which ciliation supports these cell functions (Essner et al., 2005; Liu et al., 2004; Sukumaran and Perkins, 2009; Zhai et al., 2014). In addition, zebrafish mutants with cilia defects tend to develop scoliosis (Gray et al., 2021; Latour et al., 2020; Wang et al., 2022), which was observed in *pday* mutants at 2 mpf (Fig. 1A). Thus, MAK regulates cilium-associated mechanisms in photoreceptors in zebrafish.

Next, we examined MAK protein localization in zebrafish photoreceptors by tracking the eGFP signal in *Tg[crx:eGFP-MAK]* retinas. eGFP-MAK is localized in a stem-like structure inside the OS of live photoreceptors (Fig. 5B). Next, we examined double labeling of *Tg[crx:eGFP-MAK]* retinas with anti-GFP antibody and anti-acetylated α-tubulin antibody. The latter mainly labels the ciliary axoneme. eGFP-MAK was co-localized to the ciliary axoneme visualized by an anti-acetylated α-tubulin antibody (Fig. 5C, Fig. S5B). Although overexpression of eGFP-MAK could lead to its localization to places where endogenous protein is not, it is likely that MAK is a ciliary protein in photoreceptors.

In zebrafish, photoreceptor precursors undergo one round of symmetric cell division to produce two daughter cones after 60 hpf, in which apically localized centrosomes start ciliogenesis (Zolessi et al., 2021). So, we examined the ciliary structure in *pday* mutant photoreceptors at 3 dpf. We used anti-γ tubulin and anti-acetylated α tubulin antibodies, which visualize the basal body and the axoneme of the cilium, respectively (Chen et al., 2021). Eyes shut (EYS) is a secreted extracellular matrix protein localized near at the

transition zone of photoreceptors and required for the ciliary pocket formation in zebrafish
(Yu et al., 2016). CEP290 (also known as NPHP6) is a large multidomain coiled coil
protein localized at the proximal part of the transition zone (Goncalves and Pelletier, 2017;
Park and Leroux, 2022). Genetic mutations of CEP290 are associated with Leber's
congenital amaurosis in humans (den Hollander et al., 2006) and photoreceptor
degeneration in zebrafish (Cardenas-Rodriguez et al., 2021). So, we used anti-EYS and
anti-CEP290 antibodies, which visualize transition zone-related areas. Labeling with anti-
acetylated α tubulin and anti-EYS antibodies revealed that EYS signals were normally
detected, but acetylated α tubulin signals were absent in *pday* mutant photoreceptors (Fig.
6A). Labeling with anti-γ tubulin and anti-EYS antibodies revealed that both γ tubulin and
EYS signals were normally associated in *pday* mutant photoreceptors (Fig. 6B). Next,
labeling with anti-γ tubulin and anti-CEP290 antibodies revealed that γ tubulin signals were
normal in *pday* mutant photoreceptors; however, CEP290 signals were drastically reduced
(Fig. 6C). These data indicate that the ciliary axoneme is absent and the transition zone is
also defective in *pday* mutants, whereas the basal body seems to be normal, suggesting that
the ciliary axoneme and the transition zone are the primary target of MAK. On the other
hand, normal EYS signals in *pday* mutant photoreceptors suggest that MAK is not involved
in EYS-mediated ciliary pocket formation.

The OS is a specialized cilium of photoreceptors (Bachmann-Gagescu and Neuhauss,
2019; Chen et al., 2021). Arl13b is a Joubert syndrome protein, which is localized in the
primary cilium and mutated in zebrafish *scorpion* (*sco*) mutants (Duldulao et al., 2009). In
addition, eGFP-tagged Arl13b visualizes cilia in zebrafish (Bachmann-Gagescu et al., 2011;
Borovina et al., 2010). We established two zebrafish transgenic lines, *Tg[gnat2:arl13b-
tdTomato; rho:arl13b-eGFP]*, and found that fluorescent protein-tagged Arl13b
overexpressed under the *gnat2* and *rho* promoters was localized more broadly within the
OSs of cones and rods in wild-type retinas (Fig. 6D). However, both Arl13b-tdTomato and
Arl13b-eGFP expression were absent in 3-dpf *pday* mutant photoreceptors. These data
suggest that OS formation is compromised in both rod and cone photoreceptors in *pday*
mutants.

Next, we applied electron microscopy (EM) to examine the ultrastructure of
photoreceptors at 3 dpf (Fig. 6E, 6F). In wild-type photoreceptors at 3 dpf, photoreceptors
show columnar shapes and their OSs started to form (Fig. 6E). On the other hand, OSs were
not observed in *pday* mutant photoreceptors (Fig. 6F). Higher magnification EM images
detected a basal body (a pair of centrioles) near the apical domain underneath the OS, as
well as a long extension from the basal body, which corresponds to the transition zone and
the axoneme, in wild-type photoreceptors (Fig. 6E'). However, a basal body and only a
short extension from the basal body, which might correspond to part of the transition zone,
were observed in *pday* mutant photoreceptors (Fig. 6F'). These data suggest that the OS
and the ciliary axoneme are absent in *pday* mutant photoreceptors, whereas the basal body
is normally formed. Thus, MAK regulates photoreceptor ciliogenesis by promoting
axonemal formation.

Phototransduction molecules fail to be transported to the OS in *pday* mutant
photoreceptors
Phototransduction molecules are normally transported to the OS through the cilium
(Athanasiou et al., 2018; Gulati and Palczewski, 2023; Karan et al., 2008; Wang and
Deretic, 2014). However, mislocalization of opsins leads to photoreceptor degeneration,
although detailed mechanisms are not fully understood (Alfinito and Townes-Anderson,
2002; Lopes et al., 2010; Rohrer et al., 2005). Therefore, we examined distributions of
opsins and rhodopsin in *pday* mutant photoreceptors using zpr3 (Gao et al., 2022) and 1D4
(Yin et al., 2012) antibodies, which label green opsin/rhodopsin and red opsin in zebrafish,
respectively. In 3 dpf wild-type photoreceptors, rhodopsin and red/green opsins were
normally localized in the OS (Fig. 6G, 6H). However, in *pday* mutants, these opsins were
mislocalized to the IS, plasma membrane, and the basal synaptic region (Fig. 6G, 6H). Thus,
transport of phototransduction molecules to the OS is compromised in *mak* mutants.

Kinase activity of MAK is essential for photoreceptor survival and ciliary axoneme
formation

To understand how MAK regulates photoreceptor survival and ciliary axoneme formation,
we focused on kinase-associated motifs (Wang and Kung, 2012) of MAK and conducted
site-directed mutagenesis to silence kinase activity (Fig. 7A). Like human and mouse MAK,
the kinase domain is located at the N-terminal region of zebrafish MAK (Fig. S1). In this
kinase domain, KR (32-33 aa) and TDY (157-159 aa) are required for kinase activity as an
ATP-binding motif and an autophosphorylation site, respectively (Wang and Kung, 2012).
We designed two kinase-dead mutants of zebrafish MAK, namely AR MAK and ADF
MAK, by replacing KR and TDY in wild-type zebrafish MAK with AR and ADF,
respectively (Fig. 7A). We also prepared DM MAK, which carries both AR and ADF
mutant motifs. *In vitro* kinase assay using MAK proteins purified from an *E.* coli.
expression system revealed that all three kinase-dead MAK mutants, AR MAK, ADF MAK,
and DM MAK, failed to phosphorylate a MAK mock substrate, myelin basic protein
(MBP), indicating that MAK kinase activity is silenced in either mutant, AR MAK, ADF
MAK, or DM MAK (Fig. 7B).

Next, we generated zebrafish transgenic lines, *Tg[crx:eGFP-AR MAK]*,
Tg[crx:eGFP-ADF MAK], and *Tg[crx:eGFP-DM MAK]*, by introducing kinase-dead
mutant motifs into the wild-type MAK-expressing construct, *Tg[crx:eGFP-WT MAK]*. First,
we examined subcellular localization of these three kinase-dead mutant MAK proteins in
zebrafish photoreceptors by labeling these transgenic wild-type retinas with anti-GFP
antibody (Fig. 7C). eGFP-WT MAK was localized in a stem-like pattern similar to the
axoneme of photoreceptor cilium. However, eGFP-AR MAK was localized in a spot-like
region near the proximal region of photoreceptor cilium, which seems to correspond to
either the basal body or the transition zone. Thus, ATP-binding activity is important for
proper localization of MAK in the axoneme of photoreceptor cilium. eGFP-ADF MAK was
likely to be localized in the axoneme of photoreceptor cilium, but its expression level was
low, suggesting that eGFP-ADF MAK may be unstable. eGFP-DM MAK showed low
expression level as well, which made it difficult to define its subcellular localization. Thus,
MAK kinase activity influences its localization in the axoneme of photoreceptor cilium.

Next, using zpr1 antibody, we examined whether MAK kinase activity is required for

survival of double-cone photoreceptors. We combined transgenic lines, *Tg[crx:eGFP-AR*

MAK], *Tg[crx:eGFP-ADF MAK]*, and *Tg[crx:eGFP-DM MAK]*, with the *pday* mutants. In

contrast to *Tg[crx:eGFP-WT MAK]*, all three kinase-dead mutant MAK transgenes failed to

rescue survival of double-cone photoreceptors at 6 dpf (Fig. 7D), suggesting that MAK

kinase activity is important for photoreceptor survival. Furthermore, we examined ciliary

axonemes in *pday* mutants by labeling *pday* mutants combined with *Tg[crx:eGFP-AR*

MAK], *Tg[crx:eGFP-ADF MAK]*, and *Tg[crx:eGFP-DM MAK]* with anti-acetylated α-

tubulin antibody. In contrast to *Tg[crx:eGFP-WT MAK]*, all three kinase-dead mutant

MAK transgenes failed to rescue axoneme formation at 6 dpf (Fig. 7E). Thus, MAK kinase

activity is critical for ciliary axoneme development and photoreceptor survival (Fig. 8).

Discussion

In humans and other animal models including zebrafish, many genetic mutations showing defects in photoreceptor ciliogenesis have been identified (Chen et al., 2021). In zebrafish mutants of transition zone components, selective transport of phototransduction molecules into the OS, is compromised, resulting in ectopic distribution of these OS-resident proteins outside the OS (Wang et al., 2022). However, the axoneme itself is formed in almost all of these transition-zone-defective mutants. In zebrafish mutants of BBS components, which cooperate with IFT to regulate intra-ciliary transport (Tian et al., 2023), OS non-resident proteins are mislocalized into the OS, but the axoneme and the OS are formed (Masek et al., 2022). Furthermore, in zebrafish *arl13b* mutants, with compromised cargo release in the OS, the axoneme and the OS are reduced in size, but still formed (Song et al., 2016). Thus, most ciliogenesis-defective mutants show formation of axoneme structure in photoreceptors. However, there are a few exceptions. In zebrafish mutants of an IFT-B component, IFT88 (also named *oval*), Traf3ip1 (also named *elipsa*), and a Kinesin II family protein Kif3a, the axoneme and the OS are totally absent in photoreceptors (Krock and Perkins, 2008; Omori et al., 2008; Sukumaran and Perkins, 2009; Zhu et al., 2021). Interestingly, in *ift88* and *traf3ip1* mutants, a single spot-like γ-tubulin or centrin signal is detected in the apical region of photoreceptors, similar to zebrafish *mak* mutants, suggesting that basal bodies are formed in both mutants (Omori et al., 2008; Sukumaran and Perkins, 2009; Tsujikawa and Malicki, 2004). Thus, it is likely that MAK cooperates with anterograde IFT complexes including IFT88 to promote axoneme formation during ciliogenesis.

Overexpression of wild-type MAK recovers formation of axonemes, leading to photoreceptor survival, in *mak* mutants. However, overexpression of kinase-mutant forms of MAK fail to rescue defects in axoneme and cell survival, so phosphorylation events on MAK kinase substrates are necessary for axoneme formation. Interestingly, GFP-tagged AR MAK is localized in a single spot area near the apical region of photoreceptors, which seems to correspond to the basal body or transition zone, whereas GFP-tagged, wild-type MAK is located along the axoneme. One possibility is that target proteins of MAK-mediated phosphorylation exist near the basal body or transition zone and move with MAK

along the axoneme after phosphorylation by MAK. Another possibility is that MAK interacts with adaptor proteins that are localized in the basal body or transition zone to support MAK substrate phosphorylation for ciliogenesis. At present, MAK phosphorylation substrates and adaptor proteins that facilitate MAK substrate phosphorylation are unknown in zebrafish; however, since the IFT complex is assembled and transported through the transition zone in the first step of ciliogenesis (Breslow and Holland, 2019), IFT complex components may be promising candidates to promote MAK substrate phosphorylation.

MAK proteins share a highly conserved amino acid sequence between zebrafish and mice. However, in contrast to zebrafish *mak* mutants, the axoneme is elongated in mouse *Mak* mutants (Omori et al., 2010). Why does *mak* gene knockout induce such opposite phenotypes in zebrafish and mice? From a phylogenetic point of view, MAK belongs to the MAK/ICK/MOK serine/threonine kinase family, all members of which share a highly conserved kinase domain (Chen et al., 2013; Miyata and Nishida, 1999). CILK1 (previously known as ICK) is a member of this kinase family and is now classified as a paralog of MAK. Indeed, human CILK1 shows a highly similar amino acid sequence in the kinase domain to that of zebrafish MAK (81.5% amino acid identity). Interestingly, in *Cilk1* mouse mutants, the ciliary axonemes of fibroblasts are shortened compared with those of wild-type controls (Chaya et al., 2014). Furthermore, mouse CILK1 interacts with IFT-B components, including IFT88 (Nakamura et al., 2020). Since a CILK1 homologous gene has not been identified in the zebrafish genomic database, zebrafish MAK and mouse CILK1 may share a common regulatory pathway for axoneme formation during ciliogenesis. It will be important to investigate whether zebrafish MAK cooperates with IFT-B to regulate axoneme formation during ciliogenesis.

Another interesting finding is that there is a difference in degeneration of rods and cones in *mak* mutants. In *mak* mutants, rods undergo acute apoptosis from 3 to 4 dpf, whereas cones undergo progressive shrinkage from 4 to 6 dpf. Interestingly, early pyknotic nuclei and later progressive shrinkage of the photoreceptor layer was also observed in zebrafish *ift88* and *traf3ip1* mutants, both of which show defects in axoneme formation (Bahadori et al., 2003; Doerre and Malicki, 2002; Sukumaran and Perkins, 2009; Wang et

al., 2022). Recently, it was reported that photoreceptor degeneration in ciliogenesis-
defective zebrafish *Tulp1* mutants is associated with pathological features of ferroptosis (Jia
et al., 2022), which is a new form of cell death linked to accumulation of lipid peroxides
(Yu et al., 2017). Although both rod and cone degeneration in *mak* mutants depend on
apoptosis, it is interesting to examine why cones show progressive shrinkage in the absence
of axoneme formation. Recent metabolic transcriptome analysis revealed that cellular
metabolism of rods and cones depends on aerobic glycolysis and oxidative phosphorylation,
respectively, but that cones shift toward glycolysis in the early pathological stage of
retinitis pigmentosa, suggesting an intrinsic difference of metabolic regulation between
rods and cones (Lee et al., 2023). It is interesting to examine how metabolic states regulated
differently in rods and cones in *mak* mutants.

Lastly, *mak* mutants show scoliosis and smaller body size in later development. A
comprehensive analysis of zebrafish transition zone mutants revealed phenotypic variations
between these ciliary-defective mutants (Wang et al., 2022). Among them, *cep290* mutants
show severe scoliosis in the adult stage, whereas *tmem216* mutants show smaller body size.
Indeed, CEP290 is markedly decreased in *mak* mutants, so it is possible that scoliosis and
smaller body size in *mak* mutants are caused by reduced activity of CEP290 and Tmem216,
respectively. Further study on the ciliary regulator network in *mak* mutants will reveal the
role of MAK in ciliopathy mechanism.

Materials and Methods

Ethics statement

Zebrafish were maintained at 28.5°C, on a 14h light/10h dark cycle, following standard

procedures (Westerfield, 2000). Collected embryos were cultured in E3 embryo medium (5

mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$) containing 0.003% 1-phenyl-

2-thiouera (PTU) to prevent pigmentation and 0.0002 % methylene blue to prevent fungal

growth. All experiments were performed on zebrafish embryos between the one-cell stage

and 6 dpf prior to sexual differentiation. Therefore, sexes of embryos could not be

determined. All zebrafish experiments were performed under the OIST Animal Care and

Use Program based on the Guide for the Care and Use of Laboratory Animals by the

National Research Council of the Nation Academies and approved by the Association for

Assessment and Accreditation of Laboratory Animal Care (AAALAC). The OIST

Institutional Animal Care and Use Committee approved all experimental protocols

(Protocol: ACUP-2023-016, ACUP -2023-017, ACUP -2023-018, ACUP -2023-019,

ACUP -2023-020).

Zebrafish strains and maintenance

Okinawa wild-type (oki) was used as a wild-type strain to maintain all mutant and

transgenic strains. The zebrafish mutant, *pday*[s351], was originally isolated by the Herwig

Baier lab (Muto et al., 2005). Mapping of the *pday*[s351] mutation was carried out in the

genetic background of WIK. Zebrafish transgenic lines, *Tg[gnat2:NLS-tdTomato]*[oki070] and

Tg[rho:NLS-eGFP][oki071], were used to visualize cone and rod nuclei, respectively.

Tg[hsp:mCherry-Bcl2][oki029] (Nishiwaki and Masai, 2020) was used to express mCherry-

tagged zebrafish Bcl2 under control of the zebrafish *heat-shock 70* (*hsp70*) promoter.

Tg[crx:eGFP][oki072], *Tg[crx:eGFP-WT MAK]*[oki073], *Tg[crx:eGFP-AR MAK]*[oki074] and

Tg[crx:eGFP-ADF MAK][oki075], and *Tg[crx:eGFP-DM MAK]*[oki076] were used to express

eGFP-tagged wild-type and kinase-deficient mutants MAK, respectively. *Tg[gnat2:arl13b-*

tdTomato][oki077] and *Tg[rho:arl13b-eGFP]*[oki078] were used to visualize cone and rod OSs,

respectively.

Mapping and cloning of the *pday* mutant gene

pday[+/-] fish maintained in the *oki* wild-type background were outcrossed with the *WIK* wild-type strain to generate F1 *pday*[+/-] fish with an *oki* and *WIK* trans-heterozygous wild-type background. These F1 *pday*[+/-] male and female fish were crossed to produce F2 embryos. F2 *pday*[-/-] embryos were selected by measuring OKR at 5 dpf and stored in methanol at -80°C. Genomic DNA was extracted from individual F2 *pday*[-/-] embryos. First, two DNA pools of 20 wild-type and 20 homozygous mutant embryos, respectively, were used for PCR-mediated amplification of SSLP markers (Knapik et al., 1998; Shimoda et al., 1999) to examine which chromosome hosts the *pday* mutation. Furthermore, 93 F2 *pday*[-/-] embryos (186 meiosis) were used to restrict the cytogenetic position of the *pday* mutation using SSLP markers mapped on the linked chromosome. We restricted the *pday* mutation within the genomic region flanked between two SSLP markers, z23011 and z13695, which were annotated at positions 8.517 and 9.123 Mb on chromosome 24 (zebrafish genomic database, GRCz10, Ensemble release 80). This region contained 10 genes (*tfap2a*, *tmem14ca*, *mak*, *gcm2*, *elovl2*, *BX546453*, *gnal*, *mppe1*, *CR318624*, and *dlgp1b*). Since, among these, *mak* is the only gene whose mutations cause inherited photoreceptor degeneration diseases, such as retinitis pigmentosa in humans (Ozgul et al., 2011; Stone et al., 2011), we designed a new SSLP marker, namely Mak-N1, which was located around 20 bp upstream of the first exon of the *mak* gene, and confirmed no recombination of Mak-N1 in all 93 F2 *pday*[-/-] embryos (186 meiosis). Sequence information on forward and reverse primers of z23011, z13695, and Mak-N1 is provided in the appendix.

To confirm that *pday* gene encodes MAK, full-length *mak* cDNA was amplified by PCR using mRNA prepared from wild-type and *pday*[s351] homozygous embryos at 6 dpf. However, only a partial cDNA fragment that corresponds to the C-terminal coding region covering from the middle of exons 7 to 15 was amplified by PCR from *pday* homozygous mutant mRNA. We sequenced the full-length cDNA of wild type and the C-terminal partial cDNA of *pday* homozygous mutants and found no amino acid change in the MAK coding region from exons 8 to 15 between wild-type and *pday*[-/-] cDNA. Next, we amplified six

DNA fragments from *the pday^-/-* genome, each of which covers six N-terminal coding

exons (the exon 2 - 7), respectively. Sequencing of these six DNA fragments revealed a

non-sense mutation in exon 4 of the *pday^-/-* genome. To confirm that the *pday* mutant gene

encodes MAK, we combined *pday* mutants with the transgenic line *Tg[crx:eGFP-WT*

MAK][oki073], and showed that expression of eGFP-tagged MAK in photoreceptor precursors

rescues photoreceptor degeneration in *pday* mutants.

575 Generation of DNA expression constructs and their transgenic lines

Tol2[crx: membrane targeted YFP (MYFP)] was used as a template to make the expression

construct *Tol2[crx: eGFP-MAK]. Tol2[crx: MYFP]* was constructed using the To2kit

multiple gateway-based construction system (Kwan et al., 2007), whose Tol2 vector

backbone contains the *cmcl2:CFP* transgenesis marker (Suzuki et al., 2013). *mak* cDNA

corresponding to a full-length isoform annotated on the zebrafish genomic database (mak-

201, GRCz11, Ensemble release 110), was amplified by PCR using a mRNA pool of 3-dpf,

wild-type embryos and was subcloned into pCR TOPO vector (Invitrogen). After the eGFP

cDNA fragment was tagged in-frame to the *mak* full-length cDNA at the N-terminus, the

MYFP region of *Tol2[crx: MYFP]* was replaced with the DNA fragment encoding eGFP-

tagged *mak* cDNA at the *BamHI* and *ClaI* sites to make the expression construct

pTol2[crx:eGFP-WT MAK]. The *pTol2[crx:eGFP-WT MAK]* was used as a template in

site-directed mutagenesis for making three kinase-dead MAK constructs, *Tg[crx:eGFP-AR*

MAK], Tg[crx:eGFP-ADF MAK] and *Tg[crx:eGFP-DM MAK].*

The *EF1α* promoter of the Tol2 transposon vector pT2AL200R150G (Urasaki et al.,

2006) was replaced with the zebrafish *gnat2* promoter (Iribarne et al., 2019) at the *XhoI* and

BamHI site to make *Tol2[gnat2:eGFP].* Zebrafish *arl13b* cDNA was amplified by PCR

using an mRNA pool prepared from 4-dpf, wild-type embryos and subcloned into the pCR

TOPO vector (Invitrogen). After a tdTomato cDNA fragment was tagged in-frame to the

arl13b cDNA at the C-terminus, a DNA fragment of tdTomato tagged *arl13b* cDNA was

further subcloned into the modified pT2AL200R150G carrying the *gnat2* promoter at the

BamHI and *ClaI* sites, to make the expression construct *pTol2[gnat2:arl13b-tdTomato].*

Similarly, the *EF1α* promoter of the Tol2 transposon vector pT2AL200R150G was

replaced with the zebrafish *rhodopsin* promoter (1.1kb upstream of the transcription unit) at

the *XhoI* and *BamHI* site. A DNA fragment of C-terminal eGFP-tagged *arl13b* cDNA was

prepared and subcloned into the modified pT2AL200R150G carrying the *rhodopsin*

promoter at the *BamHI* and *ClaI* sites, to make the expression construct *pTol2[rho:arl13b-*

eGFP].

The nuclear localization signal (NLS) was fused to cDNA encoding tdTomato or

eGFP at the N-terminus. DNA fragments of NLS-tdTomato and NLS-eGFP were

subcloned into the modified pT2AL200R150G carrying the zebrafish *gnat2* promoter and

the zebrafish *rhodopsin* promoter, respectively, at the *BamHI* and *ClaI* sites, to make the

expression constructs *pTol2[gnat2:NLS-tdTomato]* and *pTol2[rho:NLS-eGFP]*.

Next, these Tol2 expression construct plasmids and Tol2 transposase mRNA were

co-microinjected into one-cell-stage zebrafish fertilized eggs for generation of transgenic

lines, *Tg[crx:eGFP-WT MAK], Tg[crx:eGFP-AR MAK], Tg[crx:eGFP-ADF MAK],*

Tg[crx:eGFP-DM MAK], Tg[gnat2:arl13b-tdTomato], *Tg[rho:arl13b-eGFP],*

Tg[gnat2:NLS-tdTomato] and *Tg[rho:NLS-eGFP]*. Injected F0 fish expressing fluorescent

protein (eGFP or tdTomato) in the ONL were bred until adulthood and used to identify

founder fish, which produce fluorescent protein-expressing F1 progeny. These transgenic

F1 embryos were raised and used to produce F2 generation for establishment of the

transgenic strain.

Histology

For plastic sections, zebrafish embryos were anesthetized in 0.02% 3-aminobenzoic acid

ethyl ester (tricaine) and fixed in PBS (150mM NaCl, 10mM $PO4^{3-}$, pH 7.4) (Invitrogen)

with 4% paraformaldehyde (PFA) at 4^oC overnight. Samples were dehydrated with an

ethanol gradient before being embedded in JB4 resin (Polysciences). Samples were

sectioned at 5 µm with a microtome Rotatif, NM335E (MICROM International GmbH) and

stained with 0.1% toluidine blue for further analysis.

For TUNEL, samples were covered with the staining mixture at 37°C for 1 h

following instructions of the In Situ Cell Death Detection Kit (Roche, 11684795910). After

the TUNEL reaction, nuclei were stained with 60 mM PO_4 buffer (PB) (42.6 mM NaH_2PO_4,

17.4 mM Na_2HPO_4, pH 7.3) containing Hoechst 33342 (Fujifilm, 346-07951) at 1:1000,

followed by washing with PB with 0.1% TritonX-100 (0.1% PBTx) for 3 x 5 min. Samples

were mounted with Fluoromount (Diagnostic BioSystems, K024) for further analysis.

For immunohistochemistry, 3-6-dpf embryos were fixed with 4% PFA at room

temperature for 2 hours, and transferred to 30% sucrose in PB overnight before being

embedded and frozen in OCT compound. However, we used non PFA-fixed samples for

anti-GFP labeling shown in Fig. 6C, because the eGFP- MAK signals are stably detected

only in non-fixed tissues. Cryosections were prepared at 7 µm with a cryostat (Cryostar

NX70, Thermo Scientific.), and at 20 µm for imaging cilia components. Cryosections were

air-dried for at least 2 h, then rehydrated in 0.5% PBTx for 3 x 5 min. Rehydrated sections

were immersed in blocking solution (10% goat serum in 0.5% PBTx) for 1 h at room

temperature. Then primary antibody in blocking solution at an appropriate dilution (see

appendix) was applied, and sections were washed with 0.5% PBTx 3 x 10 min. Next,

primary antibody-labeled sections were treated with fluorophore-conjugated secondary

antibody at 1:500 and Hoechst 33342 at 1,000 (DOJINDO Laboratories, CAS23491-52-3)

in blocking solution for 1 h at room temperature, or 50 nM Sytox-Green (Molecular Probes)

in 0.1% PBTx 3 x 10 min after secondary antibody, and then washed with 0.1% PBTx 3 x

10 min. Sections were finally mounted with Fluoromount. Confocal images were scanned

with a confocal laser-scanning microscope, LSM780 (Carl Zeiss) or Fluoview FV3000

(Olympus, Evident). Information on primary and secondary antibodies is provided in the

appendix.

For EM analysis, 3-dpf embryos were fixed in PB with fixative (2.5%

glutaraldehyde, 1% paraformaldehyde, 3% sucrose, and 30 mM HEPES) on ice for 2 h,

followed by washing with PB with 3% sucrose and 30 mM HEPES 3 x 5 min. Embryos

were postfixed in 60 mM PB with 1% OsO_4 for 1 h on ice, followed by washing with Milli-

Q water three times. Samples were dehydrated with a gradient series of ethanol that was

replaced with acetone. After treatment with a gradient series of propylene oxide (PO),

epoxy resin EPON812 (Nisshin-EM, 3402) was applied to the samples before they were

embedded in resin and solidified for 2 days at 60^{o}C. Ultra-thin sections at 50 nm were

prepared with a Leica UC7 ultra-microtome and processed with Reynolds' stain (Reynolds,

1963). Images were observed and captured using a JOEL JEM-1230R transmission EM.

Bcl2 overexpression in *pday* mutant retinas

Transgenic line *Tg[hsp:mCherry-Bcl2]* (Nishiwaki and Masai, 2020) was combined with

pday mutants. Heat shock treatment was carried out by incubation of embryos at 39^{o}C for 1

h every 12 h after 1 dpf until 6 dpf.

In situ **hybridization**

Whole-mount *in situ* hybridization was carried out using a published protocol (Thisse and

Thisse, 2008). A DNA fragment containing *mak* full-length cDNA combined with the T7

RNA polymerase promoter, which is located at the 3' end of the *mak* full-length cDNA,

was amplified by PCR. Digoxigenin (DIG)-labeled anti-sense *mak* RNA probe was

generated by *in vitro* transcription from this T7 promoter: *mak* cDNA fragment as the

template using a DIG RNA labeling kit (Roche, 11175025910). Zebrafish embryos from

the 1-cell stage to 72 hpf were fixed with 4% PFA at 4^{o}C overnight, and were stored in

100% methanol at -70^{o}C. Samples were rehydrated from methanol to PBS, and then treated

with proteinase K (10 µg/mL) to increase permeability. After refixing with 4% PFA for 20

min, samples were soaked with prehybridization buffer at 65^{o}C for 2 h, then incubated with

hybridization buffer containing *mak* antisense RNA probe at 65^{o}C overnight. Samples were

washed with 2xSSC with 0.1 % Tween20 for 5 min, room temperature, and then 0.2xSSC

with 0.1 % Tween20 3 x 30 min at 65^{o}C. Washed samples were transferred to maleic acid

buffer with 0.1% Tween20 (MABT), and then to MABT with 2% blocking reagent (Roche,

11096176001) at 4^{o}C overnight. Samples were incubated with anti-DIG-AP antibody

(Roche, 11093274910) at 1:6,000 in MABT for 1 h at room temperature, washed with

MABT 3 x 15 min, and then transferred to Tris-HCl buffer (pH 9.5) for 15 min. For the

alkaline phosphatase (AP) reaction, samples were transferred to BM-Purple (Roche,

11442074001) until a blue signal developed, and were then fixed with 4% PFA overnight at

4^{o}C. These samples were mounted in 70% glycerol-PBS and stored at 4^{o}C for long-term

preservation. Images of the embryos were observed with a SteREO Discovery V12

dissection microscope (Carl Zeiss).

Live imaging

For live imaging using confocal LSM, embryos were anesthetized in 0.02% tricaine and

mounted on glass slides in appropriate positions and glued with 3% methylcellulose. An

upright confocal LSM, FV3000 (Olympus, Evident) was used for scanning fluorescent

signals in the photoreceptor OS of *Tg[crx:GFP-MAK]*.

In vitro **kinase assay**

The wild-type zebrafish *mak* cDNA fragment was introduced into pGEX6p3 (GE Health

Care Life Science). Kinase-dead *mak* cDNA fragments were generated from pGEX6p3-

zMAK(wt) with primers for site-directed mutagenesis. Plasmids were transformed into *E.*

coli strain BL21-competent cells (TaKaRa Bio, TKR9126). Colonies were inoculated into

LB broth for starter cultures incubated at 37^{o}C overnight, followed by amplification

cultures containing 1% of the starters. Amplification cultures were cooled on ice once the

O.D. reached 0.6 and IPTG was added (TaKaRa Bio, TKR9030) to a final concentration of

1 mM, and incubated at 20^{o}C overnight. Cultures were pelleted, washed with cold PBS, and

sonicated in 1% PBS TritonX-100 (PBSTx) with 1x protease inhibitor (Roche,

04693132001). Debris was then pelleted, and supernatant fractions were subjected to

Glutathione Sepharose 4B (Cyvita, 17-0756-01) for 1.5 h at 4^{o}C. After being washed with

cold PBS 3x, Sepharose beads were incubated with PreScission protease (Cyvita, 27084301)

in the solution at 4^{o}C to release MAK from the GST tag. Digested products were pelleted,

and supernatant fractions were used for *in vitro* kinase assays.

Each reaction for *in vitro* kinase assays is composed of 50 mM Tris-HCl (pH 7.0),

10 mM MgCl$_2$, 5 mM ATP (Thermo Scientific, R0441), 1 mM DTT (Thermo Scientific,

20290), 5 µg dephosphorylated myelin basic protein (Merck, 13-110), and purified MAK

protein. Reactions were incubated 1 h at 30°C. Samples were mixed with SDS sample dye

and incubated for 20 min at 75°C to stop the reactions. Samples were separated by SDS-

PAGE and transferred onto PVDF membranes for later analyses. Membranes were blocked

with 5% skim milk for 1 h at room temperature followed by probing with primary

antibodies and HRP-conjugated secondary antibodies. ECL substrate (Thermo Scientific,

34577) was applied to visualize signals using a iBright FL1500 imager (Invitrogen).

Data quantification and statistical analysis

To qualify the thickness of photoreceptor layers, photoreceptor layers in the images were

straightened with the straighten tool of ImageJ (Fig. S2A). A line dividing the

photoreceptor layer into 10 parts was generated with Adobe Illustrator. Photoreceptor layer

thickness at each decile was measured with ImageJ by following the conversion ratio of

pixels to the actual length of the scale bar. To qualify photoreceptor numbers,

photoreceptor layers in the images were straightened with ImageJ, and a reference line

marking every 1/10 points was drawn with Adobe Illustrator. Cell numbers were manually

counted. TUNEL signals in retinas were manually counted. The zpr1-positive area in the

total retinal was calculated as previously described (Fig. S2B) (Nishiwaki et al., 2013). All

statistical analyses were performed with GraphPad Prism 9.2.1. Significance levels,

numbers of samples, and details of analyses are indicated in each figure and figure legend.

Acknowledgments

We are grateful to Herwig Baier and Akira Muto for providing the *pday* mutant line. We also thank Tadashi Yamamoto for providing the vector, pGEX6p3. We thank previous lab members, Brandy Lee Denis and Maria Iribarne for supporting the experiment to clone *pday* mutant gene, Shohei Suzuki for cloning of the zebrafish *rhodopsin* promoter, Sachihiro Suzuki for cloning of the zebrafish *gnat2* promoter, Yutaka Kojima, Ayano Harata and Hiroshi Izumi for supporting histological analyses of *pday* mutant phenotypes. We are grateful to the Imaging Section and the sequencing section of the Research Support Division of OIST. We thank Steven D. Aird for critical reading and editing of the manuscript.

Data availability

N/A

Author contributions statement

Conceptualization: H-JC, IM; Methodology: H-JC, YN, IM; Software: H-JC, IM; Validation: H-JC, YN, W-CC, IM; Formal analysis: H-JC, YN, IM; Investigation: H-JC, YN, W-CC, IM; Resources: H-JC, YN, W-CC, IM; Data curation: H-JC, YN, W-CC, IM; Writing - original draft: H-JC, IM; Writing - review & editing: H-JC, IM; Visualization: H-JC, YN, W-CC, IM; Supervision: IM; Project administration: H-JC, YN, IM; Funding acquisition: IM.

Funding

This work was funded by a grant from the Okinawa Institute of Science and Technology Graduate University to IM.

Declaration of interests

The authors declare no competing interests.

Figure Legends

Fig. 1: Zebrafish *mak* mutants, *pday,* show photoreceptor degeneration.

(A) *pday* mutants at 6 dpf and 2 mpf. At 6 dpf, *pday* mutants show no morphological
differences from wild-type siblings. At 2 mpf, *pday* mutants have smaller bodies with
less-developed fins, scoliosis, and cardiac edema (open arrowhead).

(B) The zebrafish *mak* gene consists of 15 exons and encodes 633 amino acids with a
serine/threonine kinase domain. In *pday* mutants, a non-sense mutation (R70stop)
results in a truncated protein lacking most of kinase domain.

(C) Wild-type and *pday* mutant retinas. The ONL in *pday* mutants shows dense nuclear
granules (yellow arrowheads), bubble-like structures (red arrowheads) and reduced
outer segments (yellow lines). In *pday* mutants, the dorso-central ONL progressively
decreases in thickness after 4 dpf, becomes extremely flat at 5 dpf and disappears at 6
dpf (red dotted lines).

(C') Higher-magnification of the central ONL. The OS is drastically reduced in *pday*
mutants (red lines). Dense nuclear granules (yellow arrowheads), bubble-like structures
(red arrowheads), and the progressive reduction of ONL thickness (yellow lines) are
observed in *pday* mutants. HC, horizontal cells; PR, photoreceptors.

(D) Wild-type and *pday* mutant retinas carrying the transgenic line *Tg[crx:eGFP]*. Nuclei
were counterstained with Hoechst 33342 (blue). crx:eGFP (green) expression is more
severely decreased in the dorsal retina (red dotted line) than in the ventral retina of
pday mutants.

(E) Thickness of eGFP-positive photoreceptors along the DV axis of the retina. The
thickness is more severely decreased in the dorsal retina than in the ventral retina of
pday mutants. Color bars and lines indicate mean ± SD. Statistical difference is
evaluated with 2-way ANOVA and Sidak's multiple comparison tests, n=3 for each
point. ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Scale bars: 1 mm (A), 50 µm (C), 10 µm (C'), 40 µm (D).

Fig. 2: Rods undergo apoptotic-like degeneration in *pday* mutants, whereas cones progressively shrink.

(A-B) Wild-type and *pday* mutant retinas carrying the transgenic lines *Tg[rho:NLS-eGFP]* and *Tg[gnat2:NLS-tdTomato]*, which visualize rods (green) and cones (red), respectively. Nuclei were counterstained with Hoechst 33342 (blue). (A) Only green and blue channel. (B) Only red and blue channel. Squares indicate the dorsal and ventral ONL, shown in (C–F).

(C) The dorsal ONL of wild-type and *pday* mutant retinas with *Tg[rho:NLS-eGFP]* (green). In *pday* mutants, rods have condensed, round nuclei (white arrowheads) at 3 dpf. In *pday* mutants, rods are drastically reduced in number at 4 dpf and disappear at 5 and 6 dpf (asterisks), except in the peripheral region near the dorsal CMZ.

(D) The ventral ONL of wild-type and *pday* mutant retinas with *Tg[rho:NLS-eGFP]* (green). In *pday* mutants, rods have condensed, round nuclei (white arrowheads) at 3 and 4 dpf. In *pday* mutants, rods disappear at 5 and 6 dpf (asterisks), except in the peripheral region near the ventral CMZ.

(E) The dorsal ONL of wild-type and *pday* mutant retinas with *Tg[gnat2:NLS-tdTomato]* (red). In wild type, cones form monolayer at 3 dpf and two subnuclear layers after 4 dpf. In *pday* mutants, tdTomato-negative nuclei are observed, whose position is identical to that of condensed rod nuclei (white arrowheads). From 4 to 5 dpf, cones shrink progressively and become flattened (yellow line). At 6 dpf, cones are extremely flattened and disappear, leaving gaps (yellow arrowheads), except in the peripheral region near the dorsal CMZ.

(F) The ventral ONL of wild-type and *pday* mutant retinas with *Tg[gnat2:NLS-tdTomato]* (red). In wild type, cones differentiate at 3 dpf, but progressively disappear toward the periphery of the ventral ONL, where rods densely differentiate. In *pday* mutants, ONL thickness is progressively decreased after 4 dpf. Interestingly, after 5 dpf, the cone differentiating area is expanded toward the periphery of the ventral ONL (asterisks), where rods disappear.

(G) Cone and rod degeneration process in *pday* mutants. Rods undergo acute degeneration

with apoptotic-like pyknotic nuclei during 3-4 dpf, whereas cones undergo progressive

shrinkage of cell volume during 4-6 dpf and disappear by 6 dpf.

Scale bars: 40 µm (A, B).

Fig. 3: Rods and cones undergo apoptosis in *pday* mutants.

(A) TUNEL (red) of wild-type and *pday* mutant retinas. Nuclei are counterstained with

Sytox-Green (green).

(A') Higher-magnification of dorsal, central, and ventral ONL of *pday* mutant retinas,

which are indicated by white squares in (A). White arrowheads indicate TUNEL

signals (red).

(B) Double labeling of 5-dpf wild-type and *pday* mutant retinas with zpr1 (red) and anti-

rhodopsin (green) antibodies.

(B') Higher-magnification of dorsal, central, and ventral ONL of *pday* mutant retinas,

which are indicated by white squares in (B). Both rhodopsin (yellow arrowheads) and

zpr1 signals are observed in the peripheral region of the dorsal and ventral ONL. In

addition, zpr1-positive flattened cells are located in the central ONL (white arrow).

(C) TUNEL (green) of 5-dpf wild-type and *pday* mutant retinas combined with zpr1

antibody labeling (red).

(C') Higher-magnification of dorsal, central, and ventral ONL of *pday* mutant retinas,

indicated by white squares in (C). Many TUNEL signals are associated with zpr1

signals (white arrows); however, a few are not associated with zpr1 (white arrowheads).

(D) TUNEL (red) of 5-dpf wild-type and *pday* mutant retinas combined with anti-rhodopsin

antibody labeling (green).

(D') Higher-magnification of dorsal, central, and ventral ONL of *pday* mutant retinas,

indicated by white squares in (D). Rhodopsin signals are detected in the peripheral

region of dorsal and ventral ONL (yellow arrowheads). TUNEL signals associated with

rhodopsin signals are observed only in the peripheral region of the dorsal and ventral

ONL (white arrowheads). TUNEL signals are not associated with rhodopsin signals in

central ONL and non-peripheral regions of the dorsal and ventral ONL (white arrows).

(E, F) Wild-type and *pday* mutant retinas with and without the transgenic line *Tg[hsp:*

mCherry-Bcl2], labelled with zpr1 (E) and zpr3 (F) antibodies. Nuclei were

counterstained with Hoechst 33342 (blue). Both zpr1 and zpr3 signals in the ONL are

recovered in *pday* mutant retinas with *Tg[hsp: mCherry-Bcl2]* (white arrowheads).

(F') Higher-magnification of the dorsal ONL indicated by white squares, shown in (F).

Scale bars: 50 µm (A-D), 40 µm (E, F)

Fig. 4: Overexpression of MAK restores photoreceptor survival and visual function of

***pday* mutants.**

(A) Labeling of wild-type and *pday* mutant retinas with zpr1 (red) and anti-eGFP (green)

antibodies carrying transgenic line *Tg[crx:eGFP]* or *Tg[crx:eGFP-MAK]*. Nuclei were

counterstained with Hoechst 33342 (blue). (A') Only red channel. zpr1 signals are

markedly decreased in the dorso-central region of *pday* mutant retinas carrying

Tg[crx:eGFP] (white dotted lines), but maintained in *pday* mutant retinas carrying

Tg[crx:eGFP-MAK].

(B) The fraction of zpr1-positive area in the total retinal area in wild-type and *pday* mutants

with *Tg[crx:eGFP]* or *Tg[crx:eGFP-MAK]*. The fraction is markedly reduced in *pday*

mutants with *Tg[crx:eGFP]* (red bar), but recovered in *pday* mutants with

Tg[crx:eGFP-MAK] (purple bar), equivalent to wild-type with *Tg[crx:eGFP]* (blue

bar) or *Tg[crx:eGFP-MAK]* (green bar). Bars and lines indicate mean ± SD. Statistical

significance was evaluated with ordinary one-way ANOVA and Turkey's multiple

comparison tests; $*p<0.0332$, $****p<0.0002$, ns: not significant.

(C) The fraction of OKR$^+$ (blue) and OKR$^-$ (red) embryos relative to total progeny

produced by crosses of *pday*$^{+/-}$ parent fish, *pday*$^{+/-}$ parent fish carrying *Tg[crx:eGFP]*,

and *pday*$^{+/-}$ parent fish carrying *Tg[crx:eGFP-MAK]*. 57 of 244 (23.3%) embryos

produced by three independent crosses of *pday*$^{+/-}$ parent fish were negative for OKR.

However, 100% of total 257 embryos produced by three independent crosses of *pday*$^{+/-}$

parent fish carrying *Tg[crx:eGFP-MAK]* were positive for OKR. 68 of 266 (25.5%)

embryos produced by three independent crosses of *pday*$^{+/-}$ parent fish carrying

Tg[crx:eGFP] were negative for OKR. Bars and lines indicate mean ± SD. Statistical

significance was evaluated with 2-way ANOVA and Turkey's multiple comparison

tests. *** $p < 0.0002$, ns: not significant.

Scale bars: 20 µm (A, A')

Fig. 5: MAK is localized in axonemes of photoreceptor cilia.

(A) Whole-mount *in situ* hybridization of zebrafish embryos with *mak* RNA probe. *mak*

mRNA expression is observed in Kupffer's vesicles (red open arrowhead) at the tail-

bud stage, in retina (arrows) and epiphysis (red open arrowhead) at 24 hpf, and in retina,

brain and fin (arrows) at 48 hpf. Retinal expression is restricted to the photoreceptor

layer (yellow arrowheads) at 72 hpf. kv, Kuffper's vesicle; re, retina; epi, epiphysis; br,

brain; fin, pectoral fin; Tc, optic tectum; PR, photoreceptor layer.

(B) Confocal live image of *Tg[crx: eGFP-MAK]* transgenic retina at 5 dpf. eGFP-MAK

(green) is localized in the OS of photoreceptors (yellow arrowheads).

(C) Labeling of 5-dpf, *Tg[crx: eGFP-MAK]* transgenic retina with anti-acetylated α-tubulin

(red) and anti-GFP (green) antibodies. Nuclei were counterstained with Hoechst 33342

(blue). The right panel indicates higher magnification of the square shown in the left

panel. eGFP-MAK is localized in the axoneme (white open arrowheads, right panel).

Scale bars: 200 µm (A), 5 µm (B, C left), and 1 µm (C, right).

Fig. 6: The axoneme is absent in *pday* mutant photoreceptors.

(A) Wild-type and *pday* mutant retinas with anti-acetylated α-tubulin (red) and anti-EYS

(green) antibodies. Nuclei were counterstained with Hoechst 33342 (blue). Bottom

panels are higher magnification of the ONL shown in top panels.

(B) Wild-type and *pday* mutant retinas with anti-γ-tubulin (red) and anti-EYS (green)

antibodies. Nuclei were counterstained with Hoechst 33342 (blue). Bottom panels are

higher magnification of the ONL shown in top panels.

(C) Wild-type and *pday* mutant retinas with anti-γ-tubulin (red) and anti-CEP290 (green)
antibodies. Nuclei were counterstained with Hoechst 33342 (blue). Bottom panels
show higher magnification views of the ONL seen in the top panels.

(D) Wild-type and *pday* mutant retinas combined with the transgenic lines, *Tg[gnat2:*
arl13b-tdTomato] (red) and *Tg[rho: ail13b-eGFP]* (green). Nuclei were counterstained
with Hoechst 33342 (blue). Bottom panels are higher magnification of the ONL.

(E, F) EM analysis of wild-type (E) and *pday* mutant (F) photoreceptors at 3 dpf. Bottom
left panels indicate higher-magnification (3000x) of apical region of photoreceptors in
wild-type siblings (E') and *pday* mutants (F'). Bottom right panels are enlarged images
of the square region shown in the bottom left panels. Red asterisks indicate the OS.
Red and yellow arrowheads indicate basal body and ciliary axoneme, respectively.
Orange arrowheads indicate short ciliary extension from the basal body observed in
pday mutants.

(G, H) Labeling of wild-type and *pday* mutant retinas with zpr3 antibody (G) and 1D4
antibody (H). Nuclei were counterstained with Hoechst 33342 (blue). Bottom panels
are higher magnification of the ONL. Arrowheads indicate ectopic distribution of zpr3
and 1D4 signals in plasma membrane including synaptic area in *pday* mutants.

Scale bars: 5 µm (top panels of A, B, C; bottom panels of D, G, H), 1 µm (bottom panels of
A, B, C; E, E', F, F'), and 20 µm (top panels of D, G, H)

Fig. 7: MAK kinase activity is required for MAK protein localization in the ciliary
axoneme, axoneme formation, and photoreceptor maintenance.

(A) A scheme of generation of three types of MAK kinase-dead mutants: AR MAK, ADF
MAK and DM MAK. Zebrafish MAK has a catalytic domain, which contains an ATP
binding motif, KR, and autophosphorylation site, TDY. AR MAK and ADF MAK were
generated by conversion of KR and TDY into AR and ADF, respectively. DM MAK
has both AR and ADF mutations.

(B) *In vitro* kinase assay of MAK using a substrate, MBP. Only wild-type MAK
phosphorylates MBP.

(C) Labeling of 6-dpf, wild-type ONL carrying the transgenic line, *Tg[crx:eGFP],*

Tg[crx:eGFP-WT MAK], Tg[crx:eGFP-AR MAK], Tg[crx:eGFP-ADF MAK], or

Tg[crx:eGFP-DM MAK], with anti-GFP antibody (green). Nuclei were counterstained

with Hoechst 33342 (blue).

(D) Labeling of 6-dpf, wild-type and *pday* mutant retinas carrying a transgenic line,

Tg[crx:eGFP], Tg[crx:eGFP-WT MAK], Tg[crx:eGFP-AR MAK], Tg[crx:eGFP-ADF

MAK], or *Tg[crx:eGFP-DM MAK]*, with zpr1 (red) and anti-GFP (green) antibodies.

Nuclei were counterstained with Hoechst 33342 (blue). zpr1-positive photoreceptors are

degenerated in dorso-central retinas of *pday* mutants carrying *Tg[crx:eGFP]* (white

dotted line). However, photoreceptor degeneration is rescued only in *pday* mutants

carrying *Tg[crx:eGFP-WT MAK]*, but not in *pday* mutants carrying kinase-dead mutant

transgenes (white dotted line).

(E) Labeling of 3-dpf, wild-type and *pday* mutant ONL carrying the transgenic line,

Tg[crx:eGFP], Tg[crx:eGFP-WT MAK], Tg[crx:eGFP-AR MAK], Tg[crx:eGFP-ADF

MAK], or *Tg[crx:eGFP-DM MAK]*, with anti-acetylated α-tubulin antibody (red).

Nuclei were counterstained with Hoechst 33342 (blue). Axoneme formation defects are

rescued only in *pday* mutants carrying *Tg[crx:eGFP-WT MAK]* (arrowheads), but not in

pday mutants carrying kinase-dead mutant transgenes (asterisks).

Scale bars: 5 µm (C, E) and 40 µm (D).

Fig. 8: MAK is essential for ciliary axoneme formation in zebrafish photoreceptor

ciliogenesis.

Phenotypes of zebrafish *mak* mutant photoreceptors. In the absence of MAK, the axoneme

(orange) does not form and the transition zone (green) is affected, resulting in failure to

form the OS and to transport phototransduction molecules into the OS.

References

Alfinito, P. D. and Townes-Anderson, E. (2002). Activation of mislocalized opsin kills

rod cells: a novel mechanism for rod cell death in retinal disease. *Proc Natl Acad Sci U S A*

99, 5655-5660.

Athanasiou, D., Aguila, M., Bellingham, J., Li, W., McCulley, C., Reeves, P. J. and

Cheetham, M. E. (2018). The molecular and cellular basis of rhodopsin retinitis

pigmentosa reveals potential strategies for therapy. *Prog Retin Eye Res* **62**, 1-23.

Bachmann-Gagescu, R. and Neuhauss, S. C. (2019). The photoreceptor cilium and its

diseases. *Curr Opin Genet Dev* **56**, 22-33.

Bachmann-Gagescu, R., Phelps, I. G., Stearns, G., Link, B. A., Brockerhoff, S. E.,

Moens, C. B. and Doherty, D. (2011). The ciliopathy gene cc2d2a controls zebrafish

photoreceptor outer segment development through a role in Rab8-dependent vesicle

trafficking. *Hum Mol Genet* **20**, 4041-4055.

Bahadori, R., Huber, M., Rinner, O., Seeliger, M. W., Geiger-Rudolph, S., Geisler, R.

and Neuhauss, S. C. (2003). Retinal function and morphology in two zebrafish models of

oculo-renal syndromes. *Eur J Neurosci* **18**, 1377-1386.

Blackshaw, S., Harpavat, S., Trimarchi, J., Cai, L., Huang, H., Kuo, W. P., Weber, G.,

Lee, K., Fraioli, R. E., Cho, S. H., et al. (2004). Genomic analysis of mouse retinal

development. *PLoS Biol* **2**, E247.

Borovina, A., Superina, S., Voskas, D. and Ciruna, B. (2010). Vangl2 directs the

posterior tilting and asymmetric localization of motile primary cilia. *Nat Cell Biol* **12**, 407-

412.

Breslow, D. K. and Holland, A. J. (2019). Mechanism and Regulation of Centriole and

Cilium Biogenesis. *Annu Rev Biochem* **88**, 691-724.

Brouhard, G. J. and Rice, L. M. (2018). Microtubule dynamics: an interplay of
biochemistry and mechanics. *Nat Rev Mol Cell Biol* **19**, 451-463.

Cardenas-Rodriguez, M., Austin-Tse, C., Bergboer, J. G. M., Molinari, E., Sugano, Y.,
Bachmann-Gagescu, R., Sayer, J. A. and Drummond, I. A. (2021). Genetic
compensation for cilia defects in cep290 mutants by upregulation of cilia-associated small
GTPases. *J Cell Sci* **134**.

Chaya, T., Omori, Y., Kuwahara, R. and Furukawa, T. (2014). ICK is essential for cell
type-specific ciliogenesis and the regulation of ciliary transport. *EMBO J* **33**, 1227-1242.

Chen, H. Y., Kelley, R. A., Li, T. and Swaroop, A. (2021). Primary cilia biogenesis and
associated retinal ciliopathies. *Semin Cell Dev Biol* **110**, 70-88.

Chen, T., Wu, D., Moskaluk, C. A. and Fu, Z. (2013). Distinct expression patterns of
ICK/MAK/MOK protein kinases in the intestine implicate functional diversity. *PLoS One* **8**,
e79359.

Cideciyan, A. V., Aleman, T. S., Jacobson, S. G., Khanna, H., Sumaroka, A., Aguirre,
G. K., Schwartz, S. B., Windsor, E. A., He, S., Chang, B., et al. (2007). Centrosomal-
ciliary gene CEP290/NPHP6 mutations result in blindness with unexpected sparing of
photoreceptors and visual brain: implications for therapy of Leber congenital amaurosis.
Hum Mutat **28**, 1074-1083.

D'Orazi, F. D., Suzuki, S. C., Darling, N., Wong, R. O. and Yoshimatsu, T. (2020).
Conditional and biased regeneration of cone photoreceptor types in the zebrafish retina. *J*
Comp Neurol **528**, 2816-2830.

den Hollander, A. I., Koenekoop, R. K., Yzer, S., Lopez, I., Arends, M. L., Voesenek,
K. E., Zonneveld, M. N., Strom, T. M., Meitinger, T., Brunner, H. G., et al. (2006).
Mutations in the CEP290 (NPHP6) gene are a frequent cause of Leber congenital
amaurosis. *Am J Hum Genet* **79**, 556-561.

Doerre, G. and Malicki, J. (2002). Genetic analysis of photoreceptor cell development in
the zebrafish retina. *Mech Dev* **110**, 125-138.

Duldulao, N. A., Lee, S. and Sun, Z. (2009). Cilia localization is essential for in vivo
functions of the Joubert syndrome protein Arl13b/Scorpion. *Development* **136**, 4033-4042.

Essner, J. J., Amack, J. D., Nyholm, M. K., Harris, E. B. and Yost, H. J. (2005).
Kupffer's vesicle is a ciliated organ of asymmetry in the zebrafish embryo that initiates left-
right development of the brain, heart and gut. *Development* **132**, 1247-1260.

Fain, G. L., Hardie, R. and Laughlin, S. B. (2010). Phototransduction and the evolution
of photoreceptors. *Curr Biol* **20**, R114-124.

Gao, J., Cheon, K., Nusinowitz, S., Liu, Q., Bei, D., Atkins, K., Azimi, A., Daiger, S. P.,
Farber, D. B., Heckenlively, J. R., et al. (2002). Progressive photoreceptor degeneration,
outer segment dysplasia, and rhodopsin mislocalization in mice with targeted disruption of
the retinitis pigmentosa-1 (Rp1) gene. *Proc Natl Acad Sci U S A* **99**, 5698-5703.

Gao, P., Qin, Y., Qu, Z., Huang, Y., Liu, X., Li, J., Liu, F. and Liu, M. (2022). The Zpr-
3 antibody recognizes the 320-354 region of Rho and labels both rods and green cones in
zebrafish. *biorxiv*.

Ge, R., Cao, M., Chen, M., Liu, M. and Xie, S. (2022). Cytoskeletal networks in primary
cilia: Current knowledge and perspectives. *J Cell Physiol* **237**, 3975-3983.

Goncalves, J. and Pelletier, L. (2017). The Ciliary Transition Zone: Finding the Pieces
and Assembling the Gate. *Mol Cells* **40**, 243-253.

Gorden, N. T., Arts, H. H., Parisi, M. A., Coene, K. L., Letteboer, S. J., van Beersum,
S. E., Mans, D. A., Hikida, A., Eckert, M., Knutzen, D., et al. (2008). CC2D2A is
mutated in Joubert syndrome and interacts with the ciliopathy-associated basal body protein
CEP290. *Am J Hum Genet* **83**, 559-571.

Gray, R. S., Gonzalez, R., Ackerman, S. D., Minowa, R., Griest, J. F., Bayrak, M. N., Troutwine, B., Canter, S., Monk, K. R., Sepich, D. S., et al. (2021). Postembryonic screen for mutations affecting spine development in zebrafish. *Dev Biol* **471**, 18-33.

Gulati, S. and Palczewski, K. (2023). Structural view of G protein-coupled receptor signaling in the retinal rod outer segment. *Trends Biochem Sci* **48**, 172-186.

Iribarne, M., Hyde, D. R. and Masai, I. (2019). TNFalpha Induces Muller Glia to Transition From Non-proliferative Gliosis to a Regenerative Response in Mutant Zebrafish Presenting Chronic Photoreceptor Degeneration. *Front Cell Dev Biol* **7**, 296.

Jia, D., Gao, P., Lv, Y., Huang, Y., Reilly, J., Sun, K., Han, Y., Hu, H., Chen, X., Zhang, Z., et al. (2022). Tulp1 deficiency causes early-onset retinal degeneration through affecting ciliogenesis and activating ferroptosis in zebrafish. *Cell Death Dis* **13**, 962.

Kakakhel, M., Tebbe, L., Makia, M. S., Conley, S. M., Sherry, D. M., Al-Ubaidi, M. R. and Naash, M. I. (2020). Syntaxin 3 is essential for photoreceptor outer segment protein trafficking and survival. *Proc Natl Acad Sci U S A* **117**, 20615-20624.

Karan, S., Zhang, H., Li, S., Frederick, J. M. and Baehr, W. (2008). A model for transport of membrane-associated phototransduction polypeptides in rod and cone photoreceptor inner segments. *Vision Res* **48**, 442-452.

Klena, N. and Pigino, G. (2022). Structural Biology of Cilia and Intraflagellar Transport. *Annu Rev Cell Dev Biol* **38**, 103-123.

Knapik, E. W., Goodman, A., Ekker, M., Chevrette, M., Delgado, J., Neuhauss, S., Shimoda, N., Driever, W., Fishman, M. C. and Jacob, H. J. (1998). A microsatellite genetic linkage map for zebrafish (Danio rerio). *Nat Genet* **18**, 338-343.

Krock, B. L. and Perkins, B. D. (2008). The intraflagellar transport protein IFT57 is required for cilia maintenance and regulates IFT-particle-kinesin-II dissociation in vertebrate photoreceptors. *J Cell Sci* **121**, 1907-1915.

Kwan, K. M., Fujimoto, E., Grabher, C., Mangum, B. D., Hardy, M. E., Campbell, D.
S., Parant, J. M., Yost, H. J., Kanki, J. P. and Chien, C. B. (2007). The Tol2kit: a
multisite gateway-based construction kit for Tol2 transposon transgenesis constructs. *Dev*
Dyn **236**, 3088-3099.

Larison, K. D. and Bremiller, R. (1990). Early onset of phenotype and cell patterning in
the embryonic zebrafish retina. *Development* **109**, 567-576.

Latour, B. L., Van De Weghe, J. C., Rusterholz, T. D., Letteboer, S. J., Gomez, A.,
Shaheen, R., Gesemann, M., Karamzade, A., Asadollahi, M., Barroso-Gil, M., et al.
(2020). Dysfunction of the ciliary ARMC9/TOGARAM1 protein module causes Joubert
syndrome. *J Clin Invest* **130**, 4423-4439.

Lee, S. J., Emery, D., Vukmanic, E., Wang, Y., Lu, X., Wang, W., Fortuny, E., James,
R., Kaplan, H. J., Liu, Y., et al. (2023). Metabolic transcriptomics dictate responses of
cone photoreceptors to retinitis pigmentosa. *Cell Rep* **42**, 113054.

Liu, Q., Zuo, J. and Pierce, E. A. (2004). The retinitis pigmentosa 1 protein is a
photoreceptor microtubule-associated protein. *J Neurosci* **24**, 6427-6436.

Lopes, V. S., Jimeno, D., Khanobdee, K., Song, X., Chen, B., Nusinowitz, S. and
Williams, D. S. (2010). Dysfunction of heterotrimeric kinesin-2 in rod photoreceptor cells
and the role of opsin mislocalization in rapid cell death. *Mol Biol Cell* **21**, 4076-4088.

Ma, A. H., Xia, L., Desai, S. J., Boucher, D. L., Guan, Y., Shih, H. M., Shi, X. B.,
deVere White, R. W., Chen, H. W., Tepper, C. G., et al. (2006). Male germ cell-
associated kinase, a male-specific kinase regulated by androgen, is a coactivator of
androgen receptor in prostate cancer cells. *Cancer Res* **66**, 8439-8447.

Masek, M., Etard, C., Hofmann, C., Hulsmeier, A. J., Zang, J., Takamiya, M.,
Gesemann, M., Neuhauss, S. C. F., Hornemann, T., Strahle, U., et al. (2022). Loss of

the Bardet-Biedl protein Bbs1 alters photoreceptor outer segment protein and lipid
composition. *Nat Commun* **13**, 1282.

Matsushime, H., Jinno, A., Takagi, N. and Shibuya, M. (1990). A novel mammalian
protein kinase gene (mak) is highly expressed in testicular germ cells at and after meiosis.
Molecular and cellular biology **10**, 2261-2268.

Mercer, A. J. and Thoreson, W. B. (2011). The dynamic architecture of photoreceptor
ribbon synapses: cytoskeletal, extracellular matrix, and intramembrane proteins. *Vis*
Neurosci **28**, 453-471.

Mill, P., Christensen, S. T. and Pedersen, L. B. (2023). Primary cilia as dynamic and
diverse signalling hubs in development and disease. *Nat Rev Genet* **24**, 421-441.

Miyata, Y. and Nishida, E. (1999). Distantly related cousins of MAP kinase: biochemical
properties and possible physiological functions. *Biochem Biophys Res Commun* **266**, 291-
295.

Muto, A., Orger, M. B., Wehman, A. M., Smear, M. C., Kay, J. N., Page-McCaw, P. S.,
Gahtan, E., Xiao, T., Nevin, L. M., Gosse, N. J., et al. (2005). Forward genetic analysis
of visual behavior in zebrafish. *PLoS Genet* **1**, e66.

Nakamura, K., Noguchi, T., Takahara, M., Omori, Y., Furukawa, T., Katoh, Y. and
Nakayama, K. (2020). Anterograde trafficking of ciliary MAP kinase-like ICK/CILK1 by
the intraflagellar transport machinery is required for intraciliary retrograde protein
trafficking. *J Biol Chem* **295**, 13363-13376.

Nakayama, K. and Katoh, Y. (2018). Ciliary protein trafficking mediated by IFT and
BBSome complexes with the aid of kinesin-2 and dynein-2 motors. *J Biochem* **163**, 155-
164.

Nishiwaki, Y. and Masai, I. (2020). beta-SNAP activity in the outer segment growth

period is critical for preventing BNip1-dependent apoptosis in zebrafish photoreceptors. *Sci*

Rep **10**, 17379.

Nishiwaki, Y., Yoshizawa, A., Kojima, Y., Oguri, E., Nakamura, S., Suzuki, S., Yuasa-

Kawada, J., Kinoshita-Kawada, M., Mochizuki, T. and Masai, I. (2013). The BH3-only

SNARE BNip1 mediates photoreceptor apoptosis in response to vesicular fusion defects.

Dev Cell **25**, 374-387.

Oberhammer, F. A., Hochegger, K., Froschl, G., Tiefenbacher, R. and Pavelka, M.

(1994). Chromatin condensation during apoptosis is accompanied by degradation of lamin

A+B, without enhanced activation of cdc2 kinase. *J Cell Biol* **126**, 827-837.

Oel, A. P., Neil, G. J., Dong, E. M., Balay, S. D., Collett, K. and Allison, W. T. (2020).

Nrl Is Dispensable for Specification of Rod Photoreceptors in Adult Zebrafish Despite Its

Deeply Conserved Requirement Earlier in Ontogeny. *iScience* **23**, 101805.

Omori, Y., Chaya, T., Katoh, K., Kajimura, N., Sato, S., Muraoka, K., Ueno, S.,

Koyasu, T., Kondo, M. and Furukawa, T. (2010). Negative regulation of ciliary length

by ciliary male germ cell-associated kinase (Mak) is required for retinal photoreceptor

survival. *Proc Natl Acad Sci U S A* **107**, 22671-22676.

Omori, Y., Zhao, C., Saras, A., Mukhopadhyay, S., Kim, W., Furukawa, T., Sengupta,

P., Veraksa, A. and Malicki, J. (2008). Elipsa is an early determinant of ciliogenesis that

links the IFT particle to membrane-associated small GTPase Rab8. *Nat Cell Biol* **10**, 437-

444.

Ozgul, R. K., Siemiatkowska, A. M., Yucel, D., Myers, C. A., Collin, R. W., Zonneveld,

M. N., Beryozkin, A., Banin, E., Hoyng, C. B., van den Born, L. I., et al. (2011). Exome

sequencing and cis-regulatory mapping identify mutations in MAK, a gene encoding a

regulator of ciliary length, as a cause of retinitis pigmentosa. *Am J Hum Genet* **89**, 253-264.

Park, K. and Leroux, M. R. (2022). Composition, organization and mechanisms of the

transition zone, a gate for the cilium. *EMBO Rep* **23**, e55420.

Rao, K. N., Li, L., Anand, M. and Khanna, H. (2015). Ablation of retinal ciliopathy

protein RPGR results in altered photoreceptor ciliary composition. *Sci Rep* **5**, 11137.

Reynolds, E. (1963). The use of lead citrate at high pH as an electron-opaque stain in

electron microscopy. *The Journal of cell biology* **17**, 208-212.

Rohrer, B., Lohr, H. R., Humphries, P., Redmond, T. M., Seeliger, M. W. and Crouch,

R. K. (2005). Cone opsin mislocalization in Rpe65-/- mice: a defect that can be corrected

by 11-cis retinal. *Invest Ophthalmol Vis Sci* **46**, 3876-3882.

Shimoda, N., Knapik, E. W., Ziniti, J., Sim, C., Yamada, E., Kaplan, S., Jackson, D.,

de Sauvage, F., Jacob, H. and Fishman, M. C. (1999). Zebrafish genetic map with 2000

microsatellite markers. *Genomics* **58**, 219-232.

Song, P., Dudinsky, L., Fogerty, J., Gaivin, R. and Perkins, B. D. (2016). Arl13b

Interacts With Vangl2 to Regulate Cilia and Photoreceptor Outer Segment Length in

Zebrafish. *Invest Ophthalmol Vis Sci* **57**, 4517-4526.

Stone, E. M., Luo, X., Héon, E., Lam, B. L., Weleber, R. G., Halder, J. A., Affatigato,

L. M., Goldberg, J. B., Sumaroka, A., Schwartz, S. B., et al. (2011). Autosomal

Recessive Retinitis Pigmentosa Caused by Mutations in theMAKGene. *Investigative*

Opthalmology & Visual Science **52**.

Sukumaran, S. and Perkins, B. D. (2009). Early defects in photoreceptor outer segment

morphogenesis in zebrafish ift57, ift88 and ift172 Intraflagellar Transport mutants. *Vision*

Res **49**, 479-489.

Suzuki, S. C., Bleckert, A., Williams, P. R., Takechi, M., Kawamura, S. and Wong, R.

O. (2013). Cone photoreceptor types in zebrafish are generated by symmetric terminal

divisions of dedicated precursors. *Proc Natl Acad Sci U S A* **110**, 15109-15114.

Thisse, C. and Thisse, B. (2008). High-resolution in situ hybridization to whole-mount

zebrafish embryos. *Nat Protoc* **3**, 59-69.

Tian, X., Zhao, H. and Zhou, J. (2023). Organization, functions, and mechanisms of the

BBSome in development, ciliopathies, and beyond. *Elife* **12**.

Tsujikawa, M. and Malicki, J. (2004). Intraflagellar transport genes are essential for

differentiation and survival of vertebrate sensory neurons. *Neuron* **42**, 703-716.

Tucker, B. A., Burnight, E. R., Cranston, C. M., Ulferts, M. J., Luse, M. A., Westfall,

T., Scott, C. A., Marsden, A., Gibson-Corley, K., Wiley, L. A., et al. (2022).

Development and biological characterization of a clinical gene transfer vector for the

treatment of MAK-associated retinitis pigmentosa. *Gene Ther* **29**, 259-288.

Urasaki, A., Morvan, G. and Kawakami, K. (2006). Functional dissection of the Tol2

transposable element identified the minimal cis-sequence and a highly repetitive sequence

in the subterminal region essential for transposition. *Genetics* **174**, 639-649.

Vihtelic, T. S., Doro, C. J. and Hyde, D. R. (1999). Cloning and characterization of six

zebrafish photoreceptor opsin cDNAs and immunolocalization of their corresponding

proteins. *Vis Neurosci* **16**, 571-585.

Wang, J. and Deretic, D. (2014). Molecular complexes that direct rhodopsin transport to

primary cilia. *Prog Retin Eye Res* **38**, 1-19.

Wang, J., Thomas, H. R., Thompson, R. G., Waldrep, S. C., Fogerty, J., Song, P., Li,

Z., Ma, Y., Santra, P., Hoover, J. D., et al. (2022). Variable phenotypes and penetrance

between and within different zebrafish ciliary transition zone mutants. *Dis Model Mech* **15**.

Wang, L. Y. and Kung, H. J. (2012). Male germ cell-associated kinase is overexpressed

in prostate cancer cells and causes mitotic defects via deregulation of APC/CCDH1.

Oncogene **31**, 2907-2918.

Westerfield, M. (2000). *The zebrafish book. A guide for the laboratory use of zebrafish*

(Danio rerio). (4th edn): Univ. of Oregon Press, Eugene.

Xia, L., Robinson, D., Ma, A. H., Chen, H. C., Wu, F., Qiu, Y. and Kung, H. J. (2002).

Identification of human male germ cell-associated kinase, a kinase transcriptionally

activated by androgen in prostate cancer cells. *J Biol Chem* **277**, 35422-35433.

Yin, J., Brocher, J., Linder, B., Hirmer, A., Sundaramurthi, H., Fischer, U. and

Winkler, C. (2012). The 1D4 antibody labels outer segments of long double cone but not

rod photoreceptors in zebrafish. *Invest Ophthalmol Vis Sci* **53**, 4943-4951.

Yu, H., Guo, P., Xie, X., Wang, Y. and Chen, G. (2017). Ferroptosis, a new form of cell

death, and its relationships with tumourous diseases. *J Cell Mol Med* **21**, 648-657.

Yu, M., Liu, Y., Li, J., Natale, B. N., Cao, S., Wang, D., Amack, J. D. and Hu, H.

(2016). Eyes shut homolog is required for maintaining the ciliary pocket and survival of

photoreceptors in zebrafish. *Biol Open* **5**, 1662-1673.

Zhai, G., Gu, Q., He, J., Lou, Q., Chen, X., Jin, X., Bi, E. and Yin, Z. (2014). Sept6 is

required for ciliogenesis in Kupffer's vesicle, the pronephros, and the neural tube during

early embryonic development. *Mol Cell Biol* **34**, 1310-1321.

Zhu, P., Xu, J., Wang, Y. and Zhao, C. (2021). Loss of Ift74 Leads to Slow

Photoreceptor Degeneration and Ciliogenesis Defects in Zebrafish. *Int J Mol Sci* **22**.

Zolessi, F. R., Berois, N., Brauer, M. M. and Castillo, E. (2021). Building the embryo of

Developmental Biology in Uruguay. *Int J Dev Biol* **65**, 71-76.

1197

1198
1199

1200   **Appendix**

1201   **Antibodies**

NAME	SPECIES	SOURCE	USAGE	REMARK
Zpr-1	Mouse mAb	ZIRC	IHC: 1:100	
Zpr-3	Mouse mAb	ZIRC	IHC: 1:100	recognize zebrafish green opsin and rhodopsin (Gao et al., 2022)
Anti-Rhodopsin	Mouse mAb, clone 1D4	Abcam, ab5417	IHC: 1:100	recognize zebrafish red opsin (Yin et al., 2012)
Anti-GFP	Rabbit pAb	Invitrogen, A11122	IHC: 1:100	
Anti-acetylated Tubulin	Mouse mAb, clone 6-11B-1	Sigma Aldrich, T7451	IHC: 1:250	
Anti-EYS	Rabbit pAb	Novus Bio, NBP1-90038	IHC: 1:300	pH 6.0 HIAR
Anti-gamma Tubulin	Mouse mAb, clone GTU88	Sigma Aldrich, T6557	IHC: 1:100	pH 6.0 HIAR
Anti-MAK	Rabbit pAb	Scrum Inc.	WB: 1:1000	custom ab
Anti-MBP	Mouse mAb, clone SKB3	Merck, 05-675	WB: 1:5000	
Anti-phospho-MBP	Mouse mAb, clone P12	Merck, 05-429	WB: 1:10000	
Anti-GST	Mouse mAb, clone B-14	Santa Cruz, sc-138	WB: 1:10000	
Anti-zebrafish rhodopsin	Rabbit pAb	ZFIN: ZDB-ATB-09058-5	IHC: 1:5000	Ref. Vihtelic and Hyde, 1999 (DOI: 10.1017/s0952523899163168)
Anti-CEP290	Rabbit pAb	Abcam, ab84870	IHC: 1:100	

1202

1203   **Primers**

NAME	SEQUENCE (5' TO 3')	REMARK
z23011_Fw	TTGGGGAAAAAGTCTGCTTG	SSLP BSA
z23011_Rev	TCCAACATGTAAAAACCTGACAA	SSLP BSA
mak-N1_Fw	GTGCAAGGTGTGAATTAGAC	SSLP BSA
mak-N1_Rev	AACTTCCCGACTTCTGTAAC	SSLP BSA
mak_Fw3	ACAGTGGGATGGAGTGGCATCCCTT	mak cDNA cloning
mak_Rev3	TTTCCGAGCGTTGAAAGTAGTCC	mak cDNA cloning
mak-I4_Fw	TGATGTTCACCAGCCACACT	pday site sequence
mak-I3_Rev	TCCCAGTTTTGATGTTGCAT	pday site sequence
mak-E5_Fw	CACAAATGCCAAACCAGAGA	pday site sequence
mak-E4_Rev	TGGTGAAGCTAAAGGAGGTGA	pday site

		sequence
MAK-AR_Fw	AGTGGCCATCgcgAGAATGAAGAG	AR MAK mutagenesis
MAK-AR_Rev	CTCTTCATTCTcgcGATGGCCACT	AR MAK mutagenesis
MAK-ADF_Fw	cttcGTCTCAACACGATGGTATCGAG	ADF MAK mutagenesis
MAK-ADF_Rev	tctgcGTAAGGTGGCCGAGAGCG	ADF MAK mutagenesis
arl13b_Fw	GGAAATGTTCAATCTGATGGCGAACTG	arl13b cDNA cloning
arl13b_Rev	GTGTTAGGAAATGACATCATGAGTATCGC	arl13b cDNA cloning
T3_mak_Fw	CTAATTAACCCTCACTAAAGGATGAACCGTTACACGACACTCAAG	mak in situ probe
mak_T7_Rev	CTTAATACGACTCACTATAGGGTTAACGATGGCTTCCATATT	mak in situ probe
Z13695_Fw	TGTCGTAGGACATGGGGATT	BSA SSLP
Z13695_Rev	TCGCTGCTTCGTTCTCACTA	BSA SSLP

1204
1205

Figure 1

Figure 2

Figure 3

Figure 4

A

				tail-bud
1-cell	2-cell	4-cell	50%-epi	lateral / ventral

24 hpf — epi, re
48 hpf — re
72 hpf — re, Tc

24 hpf — epi, re
48 hpf — br, re, fin
72 hpf — PR, Tc

B

Photoreceptors
OS
IS

5 dpf
Tg[crx:eGFP-MAK]
ONL
INL

C

anti-acetylated α-tubulin / anti-GFP

5 dpf
Tg[crx:eGFP-MAK]

Figure 5

Figure 6

Figure 7

Figure 8