Title

High radiosensitivity in Norway spruce (*Picea abies*) is rendered by less comprehensive

mobilisation of protection, repair and stress responses compared to the radiotolerant *A.*

thaliana

Payel Bhattacharjee[1, 2*], Dajana Blagojevic[1, 2+], YeonKyeong Lee[1, 2, 3+], Gareth B Gillard[4],

Lars Grønvold[4], Torgeir Rhoden Hvidsten[5], Simen Rød Sandve[4], Ole Christian Lind[2, 6], Brit

Salbu[2, 6], Dag Anders Brede[2, 6], Jorunn E. Olsen[1, 2]

[1] Department of Plant Sciences, Faculty of Biosciences, Norwegian University of Life

Sciences, Ås, Norway

[2] Centre for Environmental Radioactivity (CERAD CoE), Norwegian University of Life

Sciences, Ås, Norway

[3] Department of Plant Biotechnology, Korea University Graduate School, 145 Anam-ro,

Seongbuk-ku, Seoul, Republic of Korea

[4] Department of Animal and Aquacultural Sciences, Faculty of Biosciences, Norwegian

University of Life Sciences, Ås, Norway

[5] Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life

Sciences, Ås, Norway

[6] Department of Environmental Sciences, Faculty of Environmental Sciences and Natural

Resource Management, Norwegian University of Life Sciences, Ås, Norway

[+] These authors contributed equally.

*Corresponding author at: Department of Plant Sciences, Faculty of Biosciences, Norwegian

University of Life Sciences, P.O. Box 5003, 1432 Ås, Norway

E-mail address: payel.bhattachrjee@nmbu.no

Abstract

Following nuclear events, risk assessment and protection of plant communities in

contaminated ecosystems require in-depth understanding of radiosensitivity of plants.

However, the physiological and molecular factors defining differential sensitivity to chronic

ionising radiation exposure are poorly understood. In this study, we compared early

molecular events associated with protection, repair, and stress responses as well as

phenotypic and cellular effects in gamma-irradiated seedlings of the radiosensitive conifer

Norway spruce (*Picea abies*) and the radiotolerant herbaceous *Arabidopsis thaliana*. After

48-h of irradiation, Norway spruce showed reduced growth at 290 mGy h^{-1} and severe

organelle damage at $\geq$ 1 mGy h^{-1} whereas *A. thaliana* showed unaffected development, minor

organelle damage at $\geq$ 100 mGy h^{-1} only and significantly less DNA damage at all dose rates.

Comparative transcriptomics revealed that Norway spruce mobilized transcription of DNA

damage repair and antioxidant genes at $\geq$ 40 mGy h^{-1} only while *A. thaliana* showed massive

activation of genes related to DNA damage repair, antioxidants, and other stress responses as

well as growth-promoting hormones and cell wall components at $\geq$ 1 mGy h^{-1}. Adverse

effects on chloroplasts and mitochondria from low dose rates on and comprehensive

downregulation of photosynthetic genes and activation of respiration genes at $\geq$ 40 mGy h^{-1}

in Norway spruce but not in *A. thaliana* may reflect the higher energy demand in Norway

spruce to simultaneously maintain its far larger genome and engage protection and repair

systems. Hence, the absence of transcriptional response at lower gamma doses and activation

of repair and protection at high dose rates only, when accumulated damage is high, is

consistent with the high radiosensitivity of Norway spruce. Conversely, the more massive

transcriptional activation of crucial repair and protection pathways even at low dose rates

complies with the high radiotolerance of *A. thaliana*.

Key words

Radiosensitivity in plants

Organelle damage

Antioxidant protection

DNA damage repair

Comparative transcriptomics

1. Introduction

Plants are frequently exposed to environmental stress and have therefore evolved stress-

management strategies. Stress tolerance capacity in general is highly variable and a defining

factor for species niche and habitat preference. Ionising radiation is a stressor that can induce

harmful effects to organisms due to damage to biological macromolecules such as DNA,

proteins, and lipids (Caplin and Wiley 2018). Field experiments and surveys of radionuclide-

contaminated plant communities and ecosystems have shown that species that are sensitive to

chronic ionising radiation may succumb and be replaced by more tolerant species (Amiro and

Sheppard 1994; Caplin and Willey 2018). Laboratory studies have confirmed that

radiosensitivity of different plant species ranges over orders of magnitude (Caplin and Wiley

2018; Blagojevic et al., 2019a). However, the underlying mechanisms conferring tolerance to

ionising radiation are not well understood. Such knowledge is needed to improve our ability

to predict environmental consequences of nuclear accidents and release of radionuclides

leading to exposure to high levels of ionizing radiation.

The average natural background level of ionising radiation is estimated to 0.3 µGy h^{-1}, while

15 µGy h^{-1} has been measured in areas with naturally elevated radionuclide levels in bedrock,

sand and/or sediments (Caplin and Willey 2018). The nuclear power plant accident in

Chernobyl in April 1986 as well as other accidents and experiments have shown that acute

high doses of 10-1000 Gy can be lethal for plants and that radiosensitivity differs between

species (UNSCEAR 1996). In response to sub-lethal chronic radiation, adverse effects such

as growth inhibition, reduced apical dominance (increased branching) and leaf chlorosis have

been observed, particularly in conifers (Caplin and Wiley 2018; (Kashparova et al., 2020). A

dose rate of 10 mGy d^{-1} (417 µGy h^{-1}) has been suggested as a threshold value in the sense

that lower dose rates have "no detrimental effects for populations of terrestrial plants in the

field" (International Atomic Energy Agency [IAEA]; Caplin and Willey 2018; UNSCEAR).

However, this does not appear to hold true for boreal conifer forests. In a long-term field

irradiation experiment in Canada persistent effects on forests including conifer trees were

observed at chronic dose rates ≥100 µGy h^{-1} (Amiro and Sheppard 1994). In addition to the

differential radiosensitivity between species, effects of ionising radiation depend on dose rate,

duration (acute or chronic) and thereby total dose, life stages and the co-occurrence of other

contaminants as well as environmental conditions (Garnier-Laplace et al. 2013). A range of

plant species, including *Arabidopsis thaliana* and rice, have shown high tolerance to ionising

radiation compared to conifers (Blagojevic et al., 2019; Vanhoudt et al., 2014; Choi et al.,

2021; Hase et al., 2020; *International Atomic Energy Agency, the Annual Report for 1992*).

The sensitivity of plant species to acute irradiation has been suggested to be attributed to the

efficiency of DNA repair systems and genome organisation and size (De Micco et al., 2011).

A. thaliana has a small genome of 135 Mb, whereas conifers like Norway spruce and Scots

pine have giant genomes of 20 and 23 Gb (Nystedt et al., 2013; Zimin et al., 2017). On the

other hand, the sensitivity to chronic irradiation appears to be regulated by delicate

mechanisms on different levels of biological organisation (Kim et al., 2011; Caplin and

Wiley 2018; Kim et al., 2019). However, factors that define sensitivity to chronic exposure

are poorly understood and apparently have functional rather than structural origin, depending

on the molecular architecture of stress-responsive pathways of different plant species.

Ionising radiation can result in damage to proteins, lipids and nucleic acids both directly and

indirectly via the formation of free radicals and subsequent reactive oxygen species (ROS)

(Caplin and Willey 2018). Thus, a significant role in plant reactions to chronic radiation is

attributed to high levels of ROS conferring oxidative stress. Plants possess potent antioxidant

systems involving enzymes such as catalases (CAT), superoxide dismutase (SOD) and

peroxidases (PER) as well as metabolites such as ascorbate (AsA), phenolic compounds and

glutathione (GSH) that can neutralise ROS and thereby prevent oxidative damage

(Dumanović et al. 2020). Studies of specific plant species including *A. thaliana*, have

revealed induction of antioxidant-related genes, including *SOD* and *ASCORBATE*

PEROXIDASE (*APX*) following ionising radiation exposure but there is limited information

about antioxidant homeostasis in conifers (Vandenhove et al., 2009; Vanhoudt et al., 2014;

van de Walle et al., 2016; Kim et al., 2011).

DNA strand breaks induced by ROS or directly by the ionising radiation trigger the activation

of DNA damage repair (DDR) pathways (Kim et al., 2019). For instance, a range of

homologous-recombination (HR) repair-mediated DDR genes known to be activated by the

plant-specific transcription factor *SUPRESSOR OF GAMMA RESPONSE 1 (SOG1)* were

significantly upregulated by gamma irradiation at 100 Gy in *A. thaliana* (Bourbousse,

Vegesna, and Law 2018; Yoshiyama, Sakaguchi, and Kimura 2013). The initiation of the

DDR is closely interlinked with the stalling of the cell cycle through the activation of cell

cycle checkpoints by regulating the cyclin (CYC)-cyclin-dependent kinase (CDK)-complexes (Qi and Zhang 2019). This can further lead to DNA damage-induced endoreduplication (Nisa et al., 2019). In addition, transcriptional recruitment of DDR proteins in response to gamma radiation requires chromatin remodelers involved in DNA methylation and histone 2A (H2A) modification (Choi et al., 2019; Kim 2019; Kim et al., 2021). In that context, increased global methylation in Scots pine plants from the Chernobyl exclusion zone has been reported (Kovalchuk et al., 2003). However, there is a lack of knowledge related to molecular events associated with DDR and other relevant pathways such as epigenetic modifications in conifers exposed to ionising radiation.

Although transcriptional regulation in response to high doses of ionising radiation in *A. thaliana* and a few other species has been explored (Choi et al., 2021; Vanhoudt et al., 2014), information on the molecular mechanisms leading to the differential radiosensitivity among plant species is yet to be clarified. Knowledge about mechanisms and factors that determine the radiosensitivity of plant species is of vital importance both with respect to risk assessment and protection of ecosystems from ionising radiation hazard. Hence, by employing gamma irradiation under standardised laboratory conditions, RNA-sequencing (RNA seq), concomitant analyses of growth, organelle, and DNA damage, as well as the dynamics of DDR gene expression, this study aimed to provide an overview of mechanisms involved in plant radiosensitivity. In this respect, early physiological, cellular, and molecular effects were compared in the radiosensitive conifer Norway spruce and the radio-tolerant *A. thaliana* to provide new insight regarding observed effects in plant communities growing in areas contaminated with radionuclides.

2. Materials and methods

2.1. Plant materials and pre-growing conditions

Seeds of the Landsberg erecta (L*er*) accession of *A. thaliana thaliana* (L) Heyhn, and

Norway spruce (*Picea abies* L. H. Karst) from the provenance CØ1 from Halden, Norway

(59°N latitude, seed lot 98063, Skogfrøverket, Hamar, Norway (skogfroverket.no) were

surface-sterilised and sowed according to the protocol in (Blagojevic, et al. 2019 a, b). The

seeds were germinated for 6 days at 20 °C under a 16 h photoperiod with a photosynthetic

photon flux density (PPFD) of 30 μmol m^{-2} s^{-1} at 400-700 nm (TL-D 58W/840 lamps,

Philips, Eindhoven, The Netherlands), as measured at the top of the petri dishes with a Li-Cor

Quantum/Radiometer/Photometer (model LI-250, LI-COR, Lincoln, NE, USA).

2.2.   Gamma irradiation of seedlings and growing conditions during the exposure

Six days old seedlings of *A. thaliana* and Norway spruce were exposed for 48 h using the

FIGARO low dose gamma irradiation facility (^{60}Co; 1173.2 and 1332.5 keV γ-rays) at the

Norwegian University of Life Sciences (Lind et al., 2019). Petri dishes (3.5 cm diameter)

with seedlings were placed at different distances from the gamma source to obtain the

average dose rates to water 1, 10, 20, 40, 100 and 290 mGy h^{-1} (Table 1). The exposure

experiments were repeated three times to obtain enough material for all the analyses. For one

out three repeated experiments, the highest dose rate was 250 mGy h^{-1} as 290 mGy h^{-1} could

not be obtained due to the decay of ^{60}Co in the source (Table S1). The dose rates to water and

dose rate intervals (in front and back of the petri dishes) were calculated from the air kerma

rates according to (Hansen et al., 2019) as previously described (Blagojevic et al. 2019 a, b).

The total doses were calculated from the estimated absorbed dose rates to water, multiplied

by the exposure time (48 h). During the exposure the room temperature was 20 ˚C ± 1 °C and

the photoperiod 12 h with an irradiance of 55 μmol m^{-2} s^{-1} from high pressure metal halide

lamps (HPI-T Plus 250W lamps, Philips, Eindhoven, The Netherlands). The red:far red

(R:FR) ratio was measured to 3.5 using a 660/730 nm Skye sensor (Skye Instruments, Powys,

Wales, UK).

2.3. Post-irradiation growing conditions

Since phenotypic effects of gamma irradiation may take some time to be manifested, the

post-irradiation effects of the 48 h of gamma exposure were recorded after transfer of

seedlings to pots filled with S-soil (Hasselfors Garden AS, Örebro, Sweden), one plant per

pot. All plants were kept in growth chambers (manufactured by Norwegian University of Life

Sciences) with temperature, relative air humidity (RH), R:FR ratio, and photoperiod

maintained as previously described (Blagojevic, et al. 2019).

2.4. Growth parameter recordings

At the end of the gamma irradiation, seedlings from both species were placed between two

transparent plastic sheets with a millimetre paper on top and scanned. The shoot and root

lengths of Norway spruce and the lengths of the hypocotyl and cotyledons in *A. thaliana* were

measured on the scanned seedlings using Image J (US National Institutes of Health,

Bethesda, MD, USA; imagej.nih.gov). Per gamma dose rate, 29-40 and 28-35 seedlings of *A.*

thaliana and Norway spruce respectively, were measured.

Post-irradiation, the number of rosette leaves as well as the number of plants with floral buds,

inflorescence and open flower buds were recorded in 23-24 *A. thaliana* plants per gamma

dose rate in time courses up to 75 days post-irradiation (dpi), and percentages of plants with

floral buds, inflorescence and open flowers were calculated. Leaf width and leaf length were

also measured at 34 and 41 dpi and height of the inflorescence at day 64. In Norway spruce,

shoot elongation, the number of needles and plant diameter were recorded in a time course of

59 dpi in 11-22 plants per dose rate. The plant diameter was calculated by averaging two

perpendicular measurements from needle tip to needle tip across the plant at the shoot apex.

Also, plant height was measured from the pot edge to the shoot apical meristem and the

cumulative shoot elongation was calculated.

2.5. Histological and cytological studies of shoot tips

Histological studies of shoot apical meristems at the end of the gamma irradiation were

performed according to (Lee et al. 2017). Shoot tips of Norway spruce and *A. thaliana* were

then fixed in 4 % paraformaldehyde and 0.025 % glutaraldehyde in phosphate buffered saline

(PBS, pH 7.0), vacuum infiltrated for 1 h at room temperature and stored at 4 °C overnight.

The fixed tissues were washed in PBS solution for 30 min and dehydrated in a graded ethanol

solution. After dehydration, approximately 3 mm of the shoot tips of each of 5 plants per

treatment were infiltrated and embedded in LR White (London Resin Company, London,

UK). After polymerisation, 1 µm sections were made using an Ultracut Leica EM UC6

microtome (Leica, Mannheim, Germany), and stained with toluidine blue O to visualise the

cells. The stained sections were inspected using a light microscope with bright field optics

(DM6B, Leica).

For cytological studies, LR white-embedded samples were sectioned into 70 nm ultrathin

sections, using an Ultracut microtome (Leica EMUC6) and stained with uranyl acetate and

lead citrate before examining by transmission electron microscopy (TEM) (FEI Morgagni

268, United States).

2.6. Comet assay for analysis of DNA damage

Single and double-strand DNA breaks at the end of the gamma irradiation and 77 dpi were

quantified using the COMET assay (Gichner 2003) with some modifications (Blagojevic, et

al. 2019). Briefly, this included isolation of cell nuclei and electrophoresis under alkaline pH,

resulting in a COMET-like appearance of damaged DNA due to movement out of the nuclei.

This was followed by quantification of the DNA in the COMET tail using fluorescence

microscopy. Three biological replicates per dose rate, each consisting of 3-4 mm of shoot tips

from 3 plants of Norway spruce and 5 plants of *A. thaliana*, were used to assess DNA

damage. For each of the three biological replicates per dose rate, totally about 180-200 cell

nuclei were scored respectively, with 60-70 nuclei scored in each of 3 technical replicates

(gels) and the overall median values were calculated.

2.7. RNA isolation for RNA sequencing

At the termination of the 48-h gamma irradiation, four biological replicates per gamma dose

rate (0, 1, 10, 40, 100 (both species) and 290 mGy h^{-1} (*A. thaliana* only due to lack of growth

inhibition and lack of/less organelle and DNA damage at lower dose rates), each containing

shoots from 20-30 and four seedlings for *A. thaliana* and Norway spruce, respectively, were

harvested in liquid nitrogen. For *A. thaliana*, total RNA was extracted using the RNeasy Plant

Mini Kit (QIAGEN Inc., Valencia, CA) following the manufacturer's protocol. For Norway

spruce, total RNA was extracted using the Masterpure Complete DNA and RNA Purification

Kit (Epicenter, Madison, WI, USA) following the manufacturer's protocol, except that 3 µl

beta-mercaptoethanol per sample replaced 1,4-dithiothreitol (DTT) and 0.5%

polyvinylpyrrolidone (PVP, mw 360 000, Sigma Aldrich, Steinheim, Germany) was added to

the extraction buffer. The purified RNA was quantified using a NanoDrop ND-1000

Spectrophotometer (Thermo Scientific, Wilmington, DE, USA) and the quality assessed with

an Agilent 2100 Bioanalyzer with an RNA 144000 Nano Kit (Agilent technologies, Palo

Alto, CA, USA).

2.8. RNA sequencing

The transcriptomes of the shoots were sequenced on an Illumina HiSeq4000 platform at the

Norwegian Sequencing Centre (Oslo, Norway) using the Strand specific 20xTruseq RNA

library preparation (paired end, read length 150 bp) as recommended by the manufacturer

(Illumina, San Diego, CA, USA). All samples were sequenced in two repeated lanes on the

HiSeq4000. The raw sequence data were submitted to the ArrayExpress under the accession

number E-MTAB-8081 for Norway spruce and E-MTAB-11120 for *A. thaliana.*

2.9. Differential gene expression analyses and functional term enrichment

The transcript abundances were estimated using the Salmon program (Patro et al. 2017)

comparing each rate with either the Norway spruce genome (*Picea abies* 1.0 assembly,

plantgenie.org) (Nystedt, et al., 2013) or *A. thaliana* genome (TAIR10, arabidopsis.org)

(Berardini et al. 2015). Gene expression was compared between the samples at different

gamma radiation dose rates and the unexposed control samples using the DESeq2 (Love et

al., 2014). Genes were classified as differentially expressed genes (DEGs) if the False

Discovery Rate (Benjamini & Hochberg correction) adjusted p-values were < 0.05.

Gene Ontology (GO) annotations of Norway spruce genes, as well as spruce to *A. thaliana*

orthologs by best sequence matches, were downloaded from plantgenie.org. The *A. thaliana*

orthologs were used to associate the Norway spruce genes with KEGG (Kyoto Encyclopaedia

of Genes and Genomes) pathways. Enrichment of GO or KEGG terms in sets of DEGs was

found using topGO (version 2.34.0) or kegga from the limma package (version 3.38.2),

respectively.

2.10. RT-qPCR analysis

Based on the significant regulation in one or both species in response to 48 h gamma

irradiation, 11 DNA repair and cell cycle control genes were selected for further RT-qPCR

validation and analysis of their regulation after 12, 24 and 48 h of gamma irradiation at 1, 10,

20, 40 and 100 mGy h^{-1} (Table S2 and S3). Four biological replicates per dose rate, each

consisting of shoots from eight and 30 seedlings for Norway spruce and *A. thaliana*,

respectively, were harvested and RNA was isolated as described above. cDNA was

synthesised from 1250 ng RNA for *A. thaliana* and 2500 ng RNA from Norway spruce in 20

µl reactions using SuperScript™ IV VILO™ cDNA Synthesis Kit (Invitrogen, Carlsbad, CA,

US) following the manufacturer's protocol, and no enzyme controls (-RT) were made for

each sample.

Gene sequences were retrieved from NCBI (ncbi.nlm.nih.gov/gene) and TAIR

(arabidopsis.org) for *A. thaliana* and from plantgenie.org for Norway spruce and exon-exon-

spanning primers were designed to avoid non-specific amplification from genomic DNA

contamination (if any) in the RNA samples, using the online tools Oligoanalyzer

(eu.idtdna.com/calc/analyzer) and Primer Blast (ncbi.nlm.nih.gov/tools/primer-blast) with

default parameters. Amplifications were performed in a 7500 Fast Real-time PCR system

(Applied Biosystems, Foster city, USA) using 20 µl of Platinum Quantitative PCR Supermix-

UDG and SYBRGreen (Thermo Fisher, Carlsbad, CA, US), using standard protocol.

To compare the relative transcript levels of the selected genes in the samples exposed to

different gamma radiation dose rates at different time points with the unexposed controls (0

mGy h^{-1}), the comparative $\Delta\Delta CT$ method was used (Livak and Schmittgen 2001). The

expression of the target genes was normalised by the mean expression of the three reference

genes, *ACTIN* (*ACT*), *ELONGATION FACTOR 1α* (*EF-1α*) and *UBIQUITIN*.

2.11. Statistical analysis for growth recordings, DNA damage and qPCR results

DNA damage, hypocotyl, cotyledon and shoot length, as well as post-irradiation growth

parameters at the final sampling time (34 or 41 dpi for *A. thaliana* and 59 dpi for Norway

spruce) were tested for normality and homoscedasticity by the Ryan-Joiner and Levene tests,

respectively. One-way ANOVA tests (glm model in Minitab 21.1, Minitab Inc, PA, USA)

were then used to test for differences in these response variables between plants exposed to

48 h of gamma radiation versus control plants. Due to the lack of normality and

homoscedasticity of the non-transformed RT-qPCR gene expression data, log-transformed data was used in two-way ANOVA for individual species (dose rate and gamma-irradiation duration as factors) and three-way ANOVA for comparison of species (plant species, dose rate and gamma-irradiation duration as factors). Tukey`s post hoc test was used to test for differences between means ($p \leq 0.05$).

3. Results

3.1. Gamma irradiation for 48 h affects growth negatively in Norway spruce but not A. thaliana

In *A. thaliana*, no significant effect of the gamma irradiation on hypocotyl and cotyledon lengths was identified (Fig. 1A and B). At 41 dpi, no significant dose response-relationship was observed for the length, width, or the number of leaves (Fig. S1A, B and C). Although there was slight delay in flowering post-irradiation at 250-290 mGy h^{-1}, eventually all plants flowered (63 dpi) and showed very similar overall morphology, flowering, and silique development (77 dpi) (Fig. S1D, E and F and Fig. 1C). In Norway spruce, the shoot length was significantly reduced (27 %) after 48 h of gamma irradiation at 290 mGy h^{-1} only, but the root length was not affected (Fig. 1D and E). At 59 dpi, the shoot diameter and number of needles were significantly reduced by 5 % and 12 %, in 100 mGy h^{-1} and 9 % and 35 % at 290 mGy h^{-1}, but no such effects were detected at 77 dpi (Fig. S2A and Fig. 1F, G and H). Hence, there was temporary growth impairment in Norway spruce after exposure, while development in *A. thaliana* remained unaffected.

3.2. More severe gamma radiation-induced organelle damage in Norway spruce than in A. thaliana

Normal dome-shaped shoot apical meristems with compact cells were observed in both species after 48 h of gamma irradiation (Fig. 2). In *A. thaliana*, slightly damaged

mitochondria were observed at 100-290 mGy h^{-1}, but the chloroplasts and cell nuclei

appeared normal at all dose rates compared to the unexposed control except that a few cells

showed less dense nuclei morphology at ≥ 1 mGy h^{-1} (Fig. 3A, B and C). In comparison, in

Norway spruce, disrupted nuclear and mitochondrial membranes were observed at ≥ 1 mGy

h^{-1}, as well as swollen chloroplasts and disrupted thylakoid membranes at ≥ 40 mGy h^{-1} (Fig.

3D, E and F).

3.3. Higher levels of DNA damage in gamma-irradiated Norway spruce than in A. thaliana

In *A. thaliana*, the Comet assay showed low but significant damage with ≤ 5 % tail DNA at

1-100 mGy h^{-1} and 12 % tail DNA at 290 mGy h^{-1} (Fig. 4A). In comparison, DNA damage

was consistently 2- to 8-fold higher in Norway spruce with 10 %, 12 %, 15 %, 17.5% and 40

% DNA in the Comet tail after the 48-h of irradiation at 1, 10, 40, 100 and 290 mGy h^{-1},

respectively (Fig. 4B). At 77 dpi, the DNA damage was substantially reduced but still

observed in *A. thaliana* at ≥ 10 mGy h^{-1} with ≤ 3% tail DNA and in Norway spruce seedlings

at ≥ 1 mGy h^{-1}, with ≤ 5% tail DNA at all dose rates (Fig. 4 A and B).

3.4 Profound transcriptome differences between gamma-irradiated Norway spruce and A.

thaliana

Comparative RNA-seq analysis of gamma-irradiated shoot samples (1, 10, 40, 100 mGy h^{-1}

both species, 290 mGy h^{-1} *A. thaliana* only), revealed extensive differences between the

species. *A. thaliana* expressed a total of 26,643 genes, of which 11,592 were differentially

expressed genes (DEGs) between any dose rate and the unexposed control (43.5 %) in one or

multiple gamma dose rates (Fig. 5A). Norway spruce expressed a total of 66,069 genes, of

which 5326 were DEGs (8.1%, Fig. 5C). In *A. thaliana*, a high number of DEGs was

observed already from 1 mGy h^{-1} (3442 DEGs), with the higher dose rates having ~4400-

5500 DEGs (Fig. 5A and B, Fig. S3A). At 100-290 mGy h^{-1}, there was a higher proportion of

upregulated genes with more than 2-fold change as compared to the lower dose rates (Fig.

5A, Fig. S3B). By contrast, in Norway spruce, very few DEGs were observed at 1 and 10

mGy h^{-1} (38 and 43 DEGs), while 822 and 5047 DEGs were observed at 40 and 100 mGy h^{-1},

respectively (Fig. 5C and D). A principal component analysis (PCA) showed that *A. thaliana*

samples exposed to 100 and 290 mGy h^{-1} of gamma radiation clustered separately from the

low dose rate groups (1-40 mGy h^{-1}) and samples without radiation (0 mGy h^{-1}), reflecting

overall differences in both high and low dose rate groups (data not shown). In Norway

spruce, samples exposed to 100, and to some extent 40 mGy h^{-1}, clustered separately from

other groups, reflecting the overall differences of doses higher than 40 mGy h^{-1} in Norway

spruce (Fig. S3D). The increase in DEGs from 40 to 100 mGy h^{-1} was much more

pronounced in Norway spruce than in *A. thaliana* (Fig. 5A vs C). The largest overlap of *A.*

thaliana DEGs was between 100 and 290 mGy h^{-1}, and between 1, 10 and 40 mGy h^{-1} (Fig.

5B). Upregulation at higher dose rates had more overlap than downregulation. A good

proportion of Norway spruce DEGs at 40 and 100 mGy h^{-1} overlapped, showing some

consistency in the response even with a large difference in total numbers of DEGs, including

many unique ones (Fig. 5D).

3.5. Distinctive pathway enrichment in Norway spruce and A. thaliana at different dose rates

GO and KEGG pathway analyses on sets of DEGs revealed a gamma dose rate-dependent

enrichment of different pathways. In *A. thaliana*, DNA replication, DNA repair, cell cycle,

endoreduplication, and endoplasmic reticulum (ER) stress were enriched at 1-40 mGy h^{-1}

(Fig. S4 and S5). Hormone biosynthesis and signalling pathways showed considerable

enrichment along with stress response at 100-290 mGy h^{-1} (Fig. S4 and S5). Oxidative stress,

defence response, growth and development, protein synthesis and catabolism as well as

mitogen-activated protein kinase (MAPK)-mediated signalling were significantly regulated at

all dose rates (Fig. S4 and S5). In Norway spruce, pathways such as DNA replication, DNA

repair, cell cycle, endoreduplication, unfolded protein response (UPR) and endoplasmic

reticulum (ER) stress, response to gamma, energy metabolism and growth and development

were all enriched at $\geq$ 40 mGy h^{-1} (Fig. S6 and S7). Photosynthesis and protein biosynthesis

and metabolism were enriched in downregulated DEGs at 100 mGy h^{-1} (Fig. S6 and S7) in

Norway spruce only (Fig. S6 and S7).

In *A. thaliana* KEGG pathway analysis revealed that the DNA damage repair (DDR)

pathways such as HR, mismatch repair (MMR), non-homologous end-joining (NHEJ) and

base excision repair (BER) were regulated at 10-40 mGy h^{-1}, and BER also at 290 mGy h^{-1}

(Fig. S5). In Norway spruce, NER, MMR, HR and NHEJ pathways were significantly

regulated at 40 and 100 mGy h^{-1} and BER at 40 mGy h^{-1} only (Fig. S7).

3.6. Distinct differences in stress response pathways between Norway spruce and A. thaliana

The specific DEGs in DDR, antioxidant protection and other stress response pathways were

further analysed to compare *A. thaliana* and Norway spruce response after the exposure to 48

h of gamma radiation.

3.6.1. DNA damage repair

In *A. thaliana*, most DEGs in the DDR pathways showed a 30-70% change in expression

(Table S3). Genes involved in DNA double strand break (DSB) repair via HR, including

RAD51, *SMC6B*, *PROLIFERATING CELL NUCLEAR ANTIGEN 2* (*PCNA2*) and *BRCA2B*,

showed significant upregulation at all dose rates in the 10-290 mGy h^{-1} range. In Norway

spruce, most DDR DEGs at 40-100 mGy h^{-1} had at least a 60% change in expression (Table

S3), and genes involved in DSB repair via HR; *SRS2*, *RAD52*, *RPA1A*, *MCM8, XRCC2* and

XRCC3, were all upregulated at 100 mGy h^{-1}. A few other genes such as *PARP1* and *PARP2*

were significantly upregulated at the highest dose rates in both species. Some DEGs showed

opposite changes in expression between species; *MRE11*, *RPA2A/ROR1* and *BRCA1* in the

HR pathway, and *MSH7* in the MMR pathway. Unique DEGs to Norway spruce included

upregulation of *KU80* in NHEJ at 40-100 mGy h^{-1} and *SOG1* at 100 mGy h^{-1}. Unique to *A.*

thaliana was upregulation of *ATM* at 100 mGy h^{-1}.

3.6.2.   Cell cycle and endoreduplication

In *A. thaliana*, most cell cycle-related DEGs including *CYCBs* and *CDKBs*, were observed at

1-40 mGy h^{-1} but not 100-290 mGy h^{-1} (Table S4). *CYCB1* was significantly upregulated at

all dose rates $\leq$ 40 mGy h^{-1}, along with many other *CYCBs*. In Norway spruce, no such

effects on B type cyclins were observed and most of the cell cycle genes showed changes in

expression at $\geq$ 40 mGy h^{-1} only. Anaphase-promoting factors *APC6* and *13*, chromatin

assembly factor *FAS2*, and the cyclin-dependent kinases regulatory subunit *CKS1* showed

upregulation at $\geq$ 40 mGy h^{-1}. Positive regulators of the cell cycle such as D type cyclins

(e.g., *CYCD1*) were significantly upregulated in *A. thaliana* at $\leq$ 40 mGy h^{-1} but

downregulated in Norway spruce at 100 mGy h^{-1}. *RBR1* was upregulated in both species.

E2Fb was upregulated at 40 mGy h^{-1} in Norway spruce only. An important G2/M checkpoint

regulator, nuclear kinase *WEE1*, was upregulated at $\leq$ 40 mGy h^{-1} in *A. thaliana*, but at $\geq$ 40

mGy h^{-1} in Norway spruce. Most of the cell cycle-related genes, such as *CYCs* and *CDKs*, are

also involved in the endocycle and the endoreduplication-related biological pathway was

enriched in *A. thaliana* at $\leq$ 40 mGy h^{-1} but very few such DEGs were observed in Norway

spruce (Table S5).

3.6.3.   Chromatin remodelers and genes in epigenetic pathways

Significant activation of genes involved in the pathway associated with de-condensation of

the nucleosome structure and activation of DDR pathways, was predominantly observed in

Norway spruce at 40-100 mGy h^{-1} (Table S6). *In A. thaliana*, a few such genes were

upregulated at $\leq$ 40 mGy h^{-1} only, and a few were downregulated at higher dose rates. In this

species, the DNA helicase *DDM1* was upregulated in *A. thaliana* at $\leq$ 40 mGy h^{-1} and DNA

methyltransferase *CMT3,* histone 2A variant *HTA8* and a component of the core chromatin

remodelling complex *ARP6* were significantly downregulated at 100-290 mGy h^{-1} (Table S6).

In Norway spruce, histone deacetylase *HOS15* genes involved in the regulation of DNA

methylation, such as *DMT1*, *IDN2* and *SUVH4*, as well as the histone 2 A variant *H2AV*,

showed significant upregulation at 40-100 mGy h^{-1}.

3.6.4.   Hormone biosynthesis and signalling

A. thaliana genes involved in biosynthesis and signalling of main growth-promoting

hormones such as auxin, cytokinin, auxin, cytokinin, gibberellin (GA) and brassinosteroid, as

well as important stress-response-related hormones such as abscisic acid (ABA) and ethylene,

were significantly regulated in response to the gamma irradiation (Table S7) especially at

higher dose rates. The auxin-responsive MYB transcription factors *AtMYB93* and *MYB61*, the

auxin-responsive genes *IAA13, IAA14, IAA27, SAUR10* and *GH3.6*, the auxin biosynthesis

gene *YUC9* as well as the auxin efflux carriers and transporters involved in polar auxin

transport, *PIN2*, *PIN3*, *AUX1* and several ABC transporters, were significantly upregulated in

this species (Table S6) but downregulated in Norway spruce. Although the auxin-

homeostasis-related gene *IAMT1*, and the auxin responsive genes *SAR1* and *SAUR76* showed

increased expression at 40-100 mGy h^{-1} in Norway spruce, several auxin signalling genes

including *ARF6, TIR1, AIR9* and *IAA26* were significantly downregulated (Table S7). In *A.*

thaliana, cytokinin biosynthesis and metabolism as well as cytokinin signalling genes such as

LOG7, AHP1 and SOT8, cytokinin-responsive genes *ARR2* and *ARR3*, and cytokinin

metabolism genes *CKX5*, were upregulated at 40-290 mGy h^{-1} (Table S7). Contrastingly none

of these genes showed significant regulation in Norway spruce. Similarly, the gibberellin

(GA) biosynthesis genes *GA2* (*ent-kaurene synthase 1*) and *GA 20-OXIDASE 2* (*GA20OX2),*

and the brassinosteroid biosynthesis and homeostasis genes *CYP90C1, CYP708A1,* and

CYP85A1, were upregulated in *A. thaliana* but not Norway spruce.

In general, ABA biosynthesis and signalling genes did not show any gradual up- or

downregulation in *A. thaliana* (Table S7). However, a few were downregulated, including the

negative regulator of ABA biosynthesis *PUB44*, and the ABA signalling genes *PYL4*, *CRK29*

and *ATHB-6*. Many stress- and ABA-responsive genes including *RCI2A*, *LTP3*, and

BFRUCT3 showed upregulation in 100-290 mGy h^{-1}. Some of the ABA-responsive genes

involved in an ABA-mediated signalling pathway such as *ASPG1*, *PPCK1*, and *RD22*

showed consistent increasing expression from 10 mGy h^{-1} to 290 mGy h^{-1}. Among the genes

regulated in both species, the ABA signalling gene *CIPK3* showed downregulation in all dose

rates in *A. thaliana*, and at 100 mGy h^{-1} in Norway spruce, and *ADH1* were upregulated in

both species at higher dose rates. In *A. thaliana*, ethylene biosynthesis and signalling genes

were mostly downregulated at lower dose rates with few exceptions (Table S6). Ethylene-

responsive genes such as *CTR1, RAP2-2, RAP2-12*, and *ERFs* as well as the ethylene

biosynthesis genes *EOL1* and *ACS5* showed significant regulation in *A. thaliana*, but not in

Norway spruce.

3.6.5. Genes in antioxidant metabolism pathways

Significant overall upregulation of antioxidant biosynthesis and signalling genes was

observed in *A. thaliana* even from 1 mGy h^{-1}. By contrast, in Norway spruce, most of the

antioxidant genes were not significantly affected below 100 mGy h^{-1} (Table S8). The key

flavonoid and anthocyanin biosynthesis genes *CHS* and *DIHYDROFLAVONOL*

REDUCTASE (*DFRA*) were upregulated in *A. thaliana* from 1 mGy h^{-1}. but were

significantly downregulated in Norway spruce at 100 mGy h^{-1}. Flavanol synthase (*FLS1*) was

upregulated in *A. thaliana* at $\leq$ 10 mGy h^{-1} and Norway spruce at 100 mGy h^{-1}. Many

PEROXIDASE (*PER*) genes including *PER3* and *PER 11* were regulated in both species but

from lower dose rates in *A. thaliana* than Norway spruce. *GLUTATHIONE-S-TRANFERASE*

(*GSTU/GSTF*) genes were upregulated in both species.

3.6.6. Cell wall component genes

In *A. thaliana* genes involved in the biosynthesis of cell wall components, cell wall

maintenance and fortification, as well as membrane receptors were significantly upregulated

at 100-290 mGy h^{-1} with large fold-changes. In Norway spruce, only very few genes related

to cell wall biogenesis and maintenance showed regulation, i.e., at 40 and 100 mGy h^{-1} (Table

S9). In *A. thaliana*, *PECTIN METHYL TRANSFERASE* (*PMEPCRF*) and *REQUIRES HIGH*

BORON (*BOR2*) involved in cellulose and pectin metabolism, *HHT1* involved in pectin and

suberin biosynthesis, the cell wall organisation and biosynthesis genes *FATTY ALCOHOL*

CAFFEOYL-COA CAFFEOYL TRANSFERASE (*FACT*) and *CYP86B1*, were upregulated at

100-290 mGy h^{-1}. The endoglucanases *GH9C2* and *GH9B13* involved in cell wall

organisation were upregulated in all dose rates. Although none of those genes showed any

significant changes at any dose rate in Norway spruce, a few other genes such as the

EXPANSINs 4 and 8, CELLULOSE SYNTHASE 5 (CESA5), and *XYLOGLUCAN*

ENDOTRANSGLUCOSYLASE/HYDROLASES XTH 5 and *9* were significantly regulated in

both species.

3.6.7. Photosynthesis and energy metabolism

Photosynthesis was massively downregulated in Norway spruce while almost no significant

changes were observed in *A. thaliana*. Photosystem (PS) I and II-related genes such as *PS I*

*REACTION CENTRE SUBUNIT*s *PSAG, PSAK* and *PSAN*, *PHOTOSYSTEM 1 LIGHT*

HARVESTING COMPLEX 1 (*LHCA1*), PS II assembly factor *HIGH CHLOROPHYLL*

FLUORESCENCE 36 (*HCF136*), essential enzymes such as *NADPH-*

PROTOCHLOROPHYLLIDE OXIDOREDUCTASE (*PORA*), and *FRUCTOSE-1,6-*

BISPHOSPHATASE (*CYFBP*), chloroplast binding proteins and membrane efflux carriers,

were downregulated in Norway spruce at 100 mGy h^{-1} (Table S10). In *A. thaliana,* only a

few genes were regulated, such as upregulation of the PS I and II reaction centre components

PSAB and *PSBB*, and *NAD(P)H-QUININE OXIDOREDUCTASE SUBUNIT 3* (*NDHC*)

(Table S10).

Similarly, energy metabolism was far more affected in Norway spruce than *A. thaliana*

(Table S11). Norway spruce particularly showed upregulation at $\geq$ 40 mGy h^{-1} of a wide

range of genes involved in glycolysis, TCA cycle and gluconeogenesis, such as *PYRUVATE*

KINASE,   GLYCERALDEHYDE-3-PHOSPHATE   DEHYDROGENASE   (GAPC1),   6-

PHOSPHOFRUCTOKINASE 4 (PFK4), TRIOSEPHOSPHATE ISOMERASE (CTIMC),

MALATE DEHYDROGENASE (MDH1), SUCCINATE DEHYDROGENASE (SDH2-1), and

several homologs of *PHOSPHOENOLPYRUVAT CABOXKINASE* (*PCKA*) and *PFK2*.

Additionally, the *CYTOCHROME C OXIDASE 6B1* (*COX6B-1*) and *CYTOCHROME C1*

(*CYTC-1*) genes involved in mitochondrial electron transport were significantly upregulated

at 100 mGy h^{-1}. *ALCOHOL DEHYDROGENASE 1* (*ADH1*) was significantly upregulated in

both species from 100 mGy h^{-1}.

3.7.   Differential dynamic regulation of DNA repair genes in A. thaliana and Norway

spruce during gamma irradiation

To understand the dynamics of DDR and cell-cycle-related gene expression during radiation

stress, both species were exposed to 12, 24 and 48 h of gamma radiation and the transcript

levels of selected genes analysed by RT-qPCR. Consistent with the RNA-seq data,

significantly increased transcript levels of *WEE1, PARP2, RAD51* and *CYCB1.2* were

observed after 48 h in *A. thaliana*, by 5-10-fold as compared to the unexposed controls (Fig.

6). However, after 12 and 24 h exposure no significant effect of the gamma irradiation on

their expression was observed. By contrast, in Norway spruce the strongest response was

generally observed after 24 h of gamma irradiation (Fig. 7). The transcript levels of *WEE1*

increased significantly at 20-100 mGy h^{-1} after 24 h only and those of *SOG1* increased at 40-

100 mGy h^{-1} after 24 h, while *PARP2* expression increased significantly by 10-30-fold at 40-

100 mGy h^{-1} after 24 and 48 h. There were no significant effects on expression at 12 h.

4.   Discussion

To improve the knowledge base for risk assessment and protection of plant communities in

radionuclide-contaminated ecosystems against ionising radiation, the current comparative

study has focussed on underlying mechanisms with respect to differential radiosensitivity in

the highly radio-sensitive conifer Norway spruce and the radio-tolerant *A. thaliana*. Through

exposure of seedlings to gamma radiation under standardised conditions and comparative

assessment of phenotypic, cellular, and genotoxic effects, combined with RNA-seq and qPCR

analysis, we present a novel in-depth understanding of the differences between the two

species in mechanisms involved in defence and repair as well as adverse effects.

Previously, we have shown dose rate-dependent impaired development in Norway spruce

seedlings exposed to gamma radiation for 144 h, with significant growth inhibition at $\geq$ 40

mGy h^{-1} and progressively developing after-effects eventually causing mortality (Dajana

Blagojevic et al. 2019). By contrast, *A. thaliana* showed no significant adverse effects at dose

rates $\leq$ 540 mGy h^{-1} (Blagojevic et al., 2019a). Norway spruce recovers better from inhibited

growth 77 dpi after 48 h exposure to 290 mGy h^{-1} (Fig. 1 D and F-H) compared to the

previous situation (after 144 h exposure to 40 mGy h $^{-1}$, (Blagojevic, et al. 2019) despite the

total dose after 48 h being higher (13,920 mGy) than after 144 h (5,760 mGy), emphasising

the severity of chronic exposure. Consistent with previous findings, *A. thaliana* did not show

any significant growth or developmental effects in this study (Fig. 1A, B, and C). The RNA-

seq analysis revealed, however, profound differences between the two species in response to

gamma irradiation (Fig. 5, S4, S5). The fact that *A. thaliana* mounts a gene regulatory

response at lower dose rates than Norway spruce signified that *A. thaliana* was more poised

towards a rapid response to inflicted radiation stress.

The energy deposited at a specific gamma radiation dose rate in the two exposed plant

species should be similar, with ionization and excitation of electrons occurring uniformly

throughout the cells. Thus, induced damage should be randomly distributed in biomolecules

such as DNA and relative to the DNA content (genome size) (Einset and Collins 2018). Our

previous work as well as this study showed less DNA damage in *A. thaliana* than in Norway

spruce and Scots pine, both at the end of the irradiation and during the post-irradiation period

(Blagojevic et al., 2019a, b; Fig. 4). We therefore hypothesised that *A. thaliana* has more

effective DNA repair and/or protection systems. Yet, the DNA damage at 77 dpi in both

species (Fig. 4), albeit at lower level than at the end of the 48-h irradiation, demonstrates that

some DNA damage or genomic instability is not critical to survival or development.

Although genomic instability may result into detrimental growth effects (Mothersill and

Seymour 1998), the lack of clear negative effects on the vegetative growth post-irradiation in

A. thaliana in contrast to Norway spruce, and only a slight delay in reproductive development

after 144 h of gamma irradiation at high dose rates in *A. thaliana*, (Blagojevic et al., 2019a)

confirms better tolerance compared to conifers. This is also supported by the mortality

observed in Norway spruce and Scots pine at dose rates > 40 mGy h^{-1}, but not in *A. thaliana*

even after exposure to 400-540 mGy h^{-1} (Blagojevic et al., 2019a). No mortality after 48 h of

gamma irradiation in both species in this study again points to a certain tolerance to damage.

The comparative RNA-seq analysis revealed that both *A. thaliana* and Norway spruce

initiated a plethora of DDR pathways to repair the damage to avoid the accumulation of

harmful mutations and maintain genome integrity. However, the response of the two species

differed in two main aspects (Fig. S9). Firstly, the upregulation of DDR-related genes at

lower doses dose rates in *A. thaliana* than in Norway spruce suggested that the former is far

more responsive towards lower levels of DNA damage. Secondly, the prominent DNA repair

pathways in *A. thaliana* were mostly HR, and to some extent MMR, whereas in Norway

spruce, in addition to HR, several other pathways such as NHEJ, MMR and DNA transleison

synthesis were activated. Hence, it is tempting to speculate that Norway spruce needed to

engage a wider range of DDR because of the significantly higher levels of DNA damage

compared to *A. thaliana*, possibly as a consequence of less antioxidant protection and a larger

genome.

One of the key contributor genes in gamma radiation-induced DDR in plants, *SOG1*, which

activates several DDR pathways and cell cycle checkpoints (Kim et al., 2019), was not

affected at any time points or dose rates tested in the gene expression analysis (Table S3 and

Fig. 6). This might indicate that induction of *SOG1* gene expression is only modulated in

higher total doses or in acute radiation that generate high level of DNA damage (Bourbousse,

Vegesna, and Law 2018). However, no change in *SOG1* expression after 48 h of gamma

irradiation does not necessarily indicate the lack of SOG1 activity as that could rather depend

on the post-translational modification by ATM- or ATR-kinases (Bourbousse et al., 2018;

Yoshiyama et al., 2013). The involvement of *SOG1* in *A. thaliana* is indeed supported by the

activation of many primary target genes of SOG1 (i.e., *RAD51, BRCA1, PARP1* and *2,*

GMI1) in response to the 48 h of gamma irradiation. By contrast, significant upregulation of

SOG1 in Norway spruce after 24 h at $\geq$ 40 mGy h^{-1} and 48 h at 100 mGy h^{-1} (Fig. 7) but no

significant regulation of ATM (Table S3)3), might imply that elevated *SOG1* expression was

triggered in Norway spruce possibly due to higher levels of accumulated DNA damage.

Moreover, upregulation of *WEE1,* a direct target of SOG1 and key regulator of CDK-

inhibition and stress-induced cell cycle arrest (Kevei et al., 2011; De Schutter et al., 2007),

coincided with *SOG1* expression in Norway spruce, strongly suggesting SOG1 involvement

in gamma-induced DNA damage repair and cell cycle arrest (Pedroza-Garcia et al., 2022;

Bourbousse et al., 2018).

Transcriptional activation of numerous B type *CYC-* and *CDKB*-genes in *A. thaliana* after

48 h at 1-40 mGy h^{-1} (Table S4) might indicate gamma radiation stress-induced cell cycle

arrest and subsequent DNA damage repair through HR by maintaining higher cyclin

threshold (Endo et al. 2006). This further supports the previous notion that enhanced CDKB1

activity is involved in ionising radiation-induced cell cycle arrest leading to HR repair in

damaged mitotically active cells (Weimer et al., 2016). On the contrary, there were few

regulations of B-type *CYC* or *CDK* genes in Norway spruce (Table S4), indicating no

significant involvement of such genes in cell cycle arrest. Furthermore, upregulation in *A.*

thaliana of the D-type cyclins responsible for cell division progress by repressing the *RBR1-*

E2F pathway along with *RBR1* and *E2FB* (Desvoyes and Gutierrez 2020), indicates further

repression of cell cycle progression in higher dose rates in contrast to Norway spruce. Hence,

the regulation of the key contributors in cell cycle progression in both species, albeit to a

larger degree in *A. thaliana*, suggests that the cell cycle is one of the most affected biological

pathways in the applied gamma radiation stress. Most of the CYCs and CDKs regulated in *A.*

thaliana in our study are also involved in the regulation of the endocycle (Table S5).

Although very few genes related to endocycle were regulated in Norway spruce, the observed

upregulation of *CKS1* and *APC6* might play a role in endoreduplication (Heyman and De

Veylder 2012; De Veylder et al., 2001). However, stress-indued endoreduplication is less

common in gymnosperms than angiosperms (De Veylder et al., 2001). A preliminary flow

cytometry study did not reveal any effect of 48 h gamma radiation at 100 mGy h^{-1} on the

endoreduplication status in *A. thaliana* and Norway spruce (results not shown), suggesting

that the applied dose was not sufficient to induce the endocycle. Increased endoreduplication

was previously shown in *A. thaliana* at 1 Gy min^{-1} (150 Gy totally) and *Lemna minor* at 1500

mGy h^{-1} for 7 days, but not at lower dose rates (Adachi et al., 2009; Van Hoeck et al., 2015).

DDR is closely related to epigenetic regulation and chromatin structure must be decondensed

for the DDR proteins to reach the damaged sites to carry out efficient repair (Choi et al.,

2019; Kim et al., 2019). Although upregulation of epigenetically-related genes were shown

previously in *A. thaliana* exposed to ionising radiation (Sidler et al., 2015), very few such

genes were regulated in *A. thaliana* in our study. By contrast, the upregulation of *H2A*

variants, DNA methylation and deacetylation-related genes particularly in Norway spruce

(Table S6, Fig. S6), clearly indicate chromatin modification at $\geq$ 40 mGy h^{-1} in this species.

The strong enrichment of DNA methylation regulation is consistent with the previously

reported DNA hypermethylation in *P. sylvestris* trees exposed to high doses of ionising

radiation in the Chernobyl area, a mechanism suggested to stabilise the genome (Kovalchuk

et al., 2003). Despite the genome of Norway spruce being 150 times bigger than that of *A.*

thaliana, it only contains 2.5 times more genes (Nystedt et al., 2013), and has large

heterochromatin regions which requires the activation of chromatin remodelers for

decondensation of chromatin regions for efficient DDR (Fig. S9). Moreover, gamma

radiation-induced damage to heterochromatin regions does not initiate rapid activation of

DDR pathways (Falk et al., 2008). It is thus plausible that Norway spruce requires a higher

level of DNA damage for the activation of DDR and chromosome remodelers only when the

damage was significantly high.

Our previous study with gamma irradiation for 144 or 360 h showed that the apical meristems

of the two species were affected differently with adverse effects in Norway spruce only

(Blagojevic et al., 2019). In this study the 48-h exposure did not affect the structural integrity

of the apical meristems (Fig. 2). We therefore proceeded to investigate the effect of the

radiation exposure on organelles. TEM image analysis revealed that integrity of thylakoid

and nuclear membranes was compromised in Norway spruce, but not in *A. thaliana* (Fig. 3).

Notably, in both species the mitochondria appeared to be the most vulnerable organelle.

Furthermore, the fact that disrupted mitochondria were evident from 1 mGy h^{-1} in Norway

spruce, while *A. thaliana* showed no damage below 100 mGy h^{-1}, indicating a major

difference in their capacity to maintain cell homeostasis.

Previous studies of gamma-irradiated plant species such as *A. thaliana* and *L. minor* have

demonstrated increased activities of ROS-scavenging enzymes such as SOD and APX

(Vanhoudt, et al. 2014; Van Hoeck et al. 2017; Xie et al. 2019), but such information was

lacking in conifers. Consistent with the notion that species-specific antioxidant response

determines oxidative stress-induced damage, less activation of such genes occurred in

Norway spruce < 40 mGy h^{-1}, while *A. thaliana* showed comprehensive modulation of

enzymatic and non-enzymatic antioxidant-related genes (*PERs, CHSs, GSTs*, anthocyanin

and flavanol synthesis genes) (Table S8). This could at least partly explain the massive

damage to mitochondria, chloroplasts, and nuclear membranes in Norway spruce in contrast

to only slight mitochondrial damage in *A. thaliana* (Fig. 2 and 3). These observations imply

that *A. thaliana* could have a higher capacity to ameliorate cellular ROS generated by gamma

irradiation, thus preventing damage to organelles and indirect DNA damage. Moreover,

antioxidants such as *PERs* can modulate cell wall composition during stress (Díaz-Tielas et

al. 2012). Cellulose and pectin were previously shown to be damaged by very high doses of

gamma radiation (1-2.5 kGy) (Kovács and Keresztes 2002). Hence, in contrast to Norway

spruce which was negatively affected, the radical upregulation of cell wall component genes

even at lower gamma dose rates in *A. thaliana* (Table S9) might suggest efficient

replacement of damaged cell wall components and cell wall fortification in *A. thaliana* (Le

Gall et al. 2015).

Antioxidants such as *GST*s regulate auxin-responsive genes which modulate plant growth in

presence of ABA to induce tolerance towards abiotic stress (Sylvestre-Gonon et al. 2019;

Jiang et al. 2010). Thus, activation of the *GST*s in Norway spruce might have affected plant

growth in addition to counteracting oxidants. This is consistent with the observed growth

retardation after 48 h irradiation (290 mGy h^{-1}) and during the post-irradiation period (100-

290 mGy h^{-1}) in Norway spruce only (Fig. 1 and Fig. S1), and the downregulation or no

regulation of growth- promoting hormones in Norway spruce (Table S7). By contrast, the

gamma-irradiation-induced upregulation of auxin and cytokinin-related genes (biosynthesis,

metabolism and signalling as well as auxin transport) in *A. thaliana* complies with the

maintenance of meristem function to sustain normal vegetative growth. These results are in

line with the previously demonstrated activation of such genes in response to abiotic stress,

indicating the importance of growth-hormone homeostasis under stress (Verma, Ravindran,

and Kumar 2016). Similar to previous reports on the induction of ABA biosynthesis and

signalling genes triggered by ROS in gamma-irradiated *A. thaliana* (Vishwakarma et al.

2017), upregulation of many such ABA-responsive genes were observed in *A. thaliana* in our

study even though the total doses applied were much lower. Also, upregulation of the

ethylene-responsive oxidative stress-pathway (multiple *ERFs*, *ACSs* and *MPK6*) and SA-

responsive genes (*WRKY* and *MYB* transcription factors) in *A. thaliana*, but not Norway

spruce (Table S7), indicate less induction of hormone response to gamma stress in the latter

and is consistent with the greater tolerance of *A. thaliana* to gamma radiation. Hence, in *A.*

thaliana better homeostasis of antioxidants, cell wall biosynthesis/fortification and hormone

signalling might contribute to the tolerance to the gamma irradiation. Conversely, less

transcriptional activation of the same protective pathways in Norway spruce are likely to

have resulted in adverse effects on overall growth and higher susceptibility to radiation-

induced oxidative damage.

The RNA-seq analysis also provided evidence that energy metabolism and photosynthesis

in Norway spruce were strongly affected by gamma irradiation in Norway spruce, but only

marginally in *A. thaliana*. In Norway spruce, the reduced expression of photosynthesis-

related genes coincided with the reduced growth and chloroplast impairment at higher dose

rates ($\geq$ 40 mGy h^{-1}). Concurringly, the strong upregulation of energy metabolism-related

genes (glycolysis, Krebs cycle, mitochondrial electron transport-related genes) (Table S11) in

this species may support the hypothesis that there is a high ATP requirement for damage

repair including maintenance of the large genome, and overall growth. Furthermore, the fact

that energy metabolism genes were upregulated, may indicate a compensatory effect to

counteract the damage to mitochondria. Hence, it appears that radiation induces energy

deficiency, which in turn requires increased energy expenditure on damage repair at the

expense of growth and is reflected by the downregulation of growth-promoting hormones in

Norway spruce at higher dose rates. However, the upregulation of energy-metabolism-genes

might additionally indicate that the damage to mitochondria at high dose rates could have led

to inefficient energy production and thus a need for the activation of respiratory genes to

meet the energy demands.

5.   Conclusions

This study demonstrates that compared to the radiosensitive Norway spruce, radiotolerant

A. thaliana mobilises more comprehensive gamma radiation-induced defence, repair, and

homeostasis-maintaining stress responses, even at very low dose rates. Despite significant

adverse effects in Norway spruce related to DNA and organelle damage and growth at low

dose rates, transcriptional activation of DNA damage repair was evident at dose rates higher

dose than 40-100 mGy h^{-1}. Furthermore, in Norway spruce, mobilization of antioxidants, as

well as modulation of growth-regulatory and stress-related hormones at lower dose was less

efficient compared to *A. thaliana*. Thus, the short-lived *A. thaliana* with quick generation-

time, and small genome with less energy requirement for maintenance, showed better

plasticity in response to radiation stress compared to the long-lived Norway spruce with

lengthy generation time and gigantic genome. These observations expand our understanding

of differential radiosensitivity in plants, and thus improves the knowledge base for risk

assessment and protection of ecosystems from ionising radiation exposure. In the context of

environmental impact, the findings strongly support that under severe ecosystem exposure

situations, radiosensitive conifers require special attention with respect to remediating

measures.

Acknowledgments

This study has been funded by the Research Council of Norway through its Centre of

Excellence (CoE) funding scheme (Project No. 223268/F50) and the Norwegian University

of Life Science (NMBU). Sincere thanks to Marit Siira and Svein Andre Kolltveit for skillful

assistance in plant growing and growth measurements and Tone Ingeborg Melby and Linda

Ripel for skillful work with qPCR analyses of gene expression.

CRediT authorship contribution statement

Payel Bhattacharjee: Conceptualization, Data curation, Formal analysis, Investigation,

Methodology, Software, Validation, Visualization, Writing - original draft, Writing - review

& editing; Dajana Blagojevic: Conceptualization, Data curation, Formal analysis,

Investigation, Methodology, Writing – review and editing; YeonKyeong Lee: Methodology,

Visualization, Formal analysis, Writing – review and editing; Gareth B Gillard: Data

curation, Formal analysis, Visualization, Writing - review & editing; Lars Grønvold, Torgeir

R Hvidsten, Simen R Sandve: Data curation, Formal analysis, Writing – review and editing;

Ole C Lind: Conceptualization , Writing – review and editing; Brit Salbu : Resources;

Funding acquisition, Writing – review and editing; Dag A Brede: Investigation,

Methodology, Writing - review & editing, Jorunn E Olsen: Conceptualization, Funding

acquisition, Investigation, Methodology, Project administration, Resources, Supervision,

Validation, Visualization, Writing - original draft, Writing - review & editing.

References

743 Berardini, T. Z., L. Reiser, D. Li, Y. Mezheritsky, R. Muller, E. Strait, and E. Huala. 2015. "The
744   Arabidopsis Information Resource: Making and Mining The "Gold Standard" Annotated
745   Reference Plant Genome." *Genesis* 53, no. 8 (Aug): 474-85.
746   https://dx.doi.org/10.1002/dvg.22877.

747
748 Blagojevic, D., Y. Lee, D. A. Brede, O. C. Lind, I. Yakovlev, K. A. Solhaug, C. G. Fossdal, B. Salbu, and J.
749   E. Olsen. 2019. "Comparative Sensitivity to Gamma Radiation at the Organismal, Cell and
750   Dna Level in Young Plants of Norway Spruce, Scots Pine and Arabidopsis Thaliana." *Planta*
751   (Aug). https://dx.doi.org/10.1007/s00425-019-03250-y.

752
753

754
755 Blagojevic Dajana, Yeonkyeong Lee, Dag Brede, Ole Christian Lind, Brit Salbu, Knut Asbjørn Solhaug,
756   Line Nybakken, and Jorunn Olsen.. "Comparative Toxicity Assessment of Ionizing Radiation in
757   Scots Pine, Norway Spruce and *Arabidopsis Thaliana*." *Planta* (2019) 250:1567–1590.
758   https://doi.org/10.1007/s00425-019-03250-y

759
760 Bourbousse, C., N. Vegesna, and J. A. Law. 2018. "Sog1 Activator and Myb3r Repressors Regulate a
761   Complex Dna Damage Network In." *Proc Natl Acad Sci U S A* 115, no. 52 (12 26): E12453-
762   E12462. https://dx.doi.org/10.1073/pnas.1810582115.

763
764 Caplin, N., and N. Willey. 2018. "Ionizing Radiation, Higher Plants, and Radioprotection: From Acute
765   High Doses to Chronic Low Doses." *Front Plant Sci* 9: 847.
766   https://dx.doi.org/10.3389/fpls.2018.00847.

767
768 Choi, H. I., S. M. Han, Y. D. Jo, M. J. Hong, S. H. Kim, and J. B. Kim. 2021. "Effects of Acute and Chronic
769   Gamma Irradiation on the Cell Biology and Physiology of Rice Plants." *Plants (Basel)* 10, no. 3
770   (Feb 25). https://dx.doi.org/10.3390/plants10030439.

771
Choi, S. H., T. H. Ryu, J. I. Kim, S. Lee, S. S. Lee, and J. H. Kim. 2019. "Mutation in Ddm1 Inhibits the
Homology Directed Repair of Double Strand Breaks." *PLoS One* 14, no. 2: e0211878.
https://dx.doi.org/10.1371/journal.pone.0211878.

De Schutter, K., J. Joubès, T. Cools, A. Verkest, F. Corellou, E. Babiychuk, E. Van Der Schueren, T.
Beeckman, S. Kushnir, D. Inzé, and L. De Veylder. 2007. "Arabidopsis Wee1 Kinase Controls
Cell Cycle Arrest in Response to Activation of the Dna Integrity Checkpoint." *Plant Cell* 19,
no. 1 (Jan): 211-25. https://dx.doi.org/10.1105/tpc.106.045047.

De Veylder, L., G. T. Beemster, T. Beeckman, and D. Inzé. 2001. "Cks1at Overexpression in
Arabidopsis Thaliana Inhibits Growth by Reducing Meristem Size and Inhibiting Cell-Cycle
Progression." *Plant J* 25, no. 6 (Mar): 617-26. https://dx.doi.org/10.1046/j.1365-
313x.2001.00996.x.

Desvoyes, B., and C. Gutierrez. 2020. "Roles of Plant Retinoblastoma Protein: Cell Cycle and Beyond."
EMBO J 39, no. 19 (10 01): e105802. https://dx.doi.org/10.15252/embj.2020105802.

Dumanović, J., E. Nepovimova, M. Natić, K. Kuča, and V. Jaćević. 2020. "The Significance of Reactive
Oxygen Species and Antioxidant Defense System in Plants: A Concise Overview." *Front Plant
Sci* 11: 552969. https://dx.doi.org/10.3389/fpls.2020.552969.

Díaz-Tielas, C., E. Graña, M. J. Reigosa, and A. M. Sánchez-Moreiras. 2012. "The Role of Peroxidases
on the Mode of Action of Chalcone in Arabidopsis Roots." *Plant Signal Behav* 7, no. 10 (Oct
01): 1274-6. https://dx.doi.org/10.4161/psb.21594.

Einset, J., and A. R. Collins. 2018. "Genome Size and Sensitivity to Dna Damage by X-Rays-Plant
Comets Tell the Story." *Mutagenesis* 33, no. 1 (Feb 24): 49-51.
https://dx.doi.org/10.1093/mutage/gex029.

Endo, M., Y. Ishikawa, K. Osakabe, S. Nakayama, H. Kaya, T. Araki, K. Shibahara, K. Abe, H. Ichikawa,
L. Valentine, B. Hohn, and S. Toki. 2006. "Increased Frequency of Homologous
Recombination and T-Dna Integration in Arabidopsis Caf-1 Mutants." *EMBO J* 25, no. 23 (Nov
29): 5579-90. https://dx.doi.org/10.1038/sj.emboj.7601434.

Falk, M., E. Lukásová, and S. Kozubek. 2008. "Chromatin Structure Influences the Sensitivity of Dna
to Gamma-Radiation." *Biochim Biophys Acta* 1783, no. 12 (Dec): 2398-414.
https://dx.doi.org/10.1016/j.bbamcr.2008.07.010.

Garnier-Laplace, J., S. Geras'kin, C. Della-Vedova, K. Beaugelin-Seiller, T. G. Hinton, A. Real, and A.
Oudalova. 2013. "Are Radiosensitivity Data Derived from Natural Field Conditions Consistent
with Data from Controlled Exposures? A Case Study of Chernobyl Wildlife Chronically
Exposed to Low Dose Rates." *J Environ Radioact* 121 (Jul): 12-21.
https://dx.doi.org/10.1016/j.jenvrad.2012.01.013.

Gichner, T. 2003. "Dna Damage Induced by Indirect and Direct Acting Mutagens in Catalase-Deficient
Transgenic Tobacco. Cellular and Acellular Comet Assays." *Mutat Res* 535, no. 2 (Mar 03):
187-93. https://dx.doi.org/10.1016/s1383-5718(02)00320-0.

Hansen, E. L., O. C. Lind, D. H. Oughton, and B. Salbu. 2019. "A Framework for Exposure
821   Characterization and Gamma Dosimetry at the Nmbu Figaro Irradiation Facility." *Int J Radiat*
822   *Biol* 95, no. 1 (01): 82-89. https://dx.doi.org/10.1080/09553002.2018.1539878.

824   Hase, Y., K. Satoh, H. Seito, and Y. Oono. 2020. "Genetic Consequences of Acute/Chronic Gamma and
825   Carbon Ion Irradiation Of." *Front Plant Sci* 11: 336.
826   https://dx.doi.org/10.3389/fpls.2020.00336.

828   Heyman, J., and L. De Veylder. 2012. "The Anaphase-Promoting Complex/Cyclosome in Control of
829   Plant Development." *Mol Plant* 5, no. 6 (Nov): 1182-94.
830   https://dx.doi.org/10.1093/mp/sss094.

832   *International Atomic Energy Agency, the Annual Report for 1992*.

834   Jiang, H. W., M. J. Liu, I. C. Chen, C. H. Huang, L. Y. Chao, and H. L. Hsieh. 2010. "A Glutathione S-
835   Transferase Regulated by Light and Hormones Participates in the Modulation of Arabidopsis
836   Seedling Development." *Plant Physiol* 154, no. 4 (Dec): 1646-58.
837   https://dx.doi.org/10.1104/pp.110.159152.

839   Kevei, Z., M. Baloban, O. Da Ines, H. Tiricz, A. Kroll, K. Regulski, P. Mergaert, and E. Kondorosi. 2011.
840   "Conserved Cdc20 Cell Cycle Functions Are Carried out by Two of the Five Isoforms in
841   Arabidopsis Thaliana." *PLoS One* 6, no. 6: e20618.
842   https://dx.doi.org/10.1371/journal.pone.0020618.

844   Kim, D. S., J. B. Kim, E. J. Goh, W. J. Kim, S. H. Kim, Y. W. Seo, C. S. Jang, and S. Y. Kang. 2011.
845   "Antioxidant Response of Arabidopsis Plants to Gamma Irradiation: Genome-Wide
846   Expression Profiling of the Ros Scavenging and Signal Transduction Pathways." *J Plant Physiol*
847   168, no. 16 (Nov 01): 1960-71. https://dx.doi.org/10.1016/j.jplph.2011.05.008.

849   Kim, J. H. 2019. "Chromatin Remodeling and Epigenetic Regulation in Plant Dna Damage Repair." *Int
850   J Mol Sci* 20, no. 17 (Aug). https://dx.doi.org/10.3390/ijms20174093.

852   Kim, J. H., K. Hwangbo, E. Lee, S. K. Dubey, M. S. Chung, B. Y. Chung, and S. Lee. 2021. "Application of
853   Gamma Ray-Responsive Genes for Transcriptome-Based Phytodosimetry in Rice." *Plants
854   (Basel)* 10, no. 5 (May 13). https://dx.doi.org/10.3390/plants10050968.

856   Kim, J. H., T. H. Ryu, S. S. Lee, S. Lee, and B. Y. Chung. 2019. "Ionizing Radiation Manifesting Dna
857   Damage Response in Plants: An Overview of Dna Damage Signaling and Repair Mechanisms
858   in Plants." *Plant Sci* 278 (Jan): 44-53. https://dx.doi.org/10.1016/j.plantsci.2018.10.013.

860   Kovalchuk, O., P. Burke, A. Arkhipov, N. Kuchma, S. J. James, I. Kovalchuk, and I. Pogribny. 2003.
861   "Genome Hypermethylation in Pinus Silvestris of Chernobyl--a Mechanism for Radiation
862   Adaptation?" *Mutat Res* 529, no. 1-2 (Aug 28): 13-20. https://dx.doi.org/10.1016/s0027-
863   5107(03)00103-9.

Kovács, E., and A. Keresztes. 2002. "Effect of Gamma and Uv-B/C Radiation on Plant Cells." *Micron*
33, no. 2: 199-210. https://dx.doi.org/10.1016/s0968-4328(01)00012-9.

Le Gall, H., F. Philippe, J. M. Domon, F. Gillet, J. Pelloux, and C. Rayon. 2015. "Cell Wall Metabolism in
Response to Abiotic Stress." *Plants (Basel)* 4, no. 1 (Feb 16): 112-66.
https://dx.doi.org/10.3390/plants4010112.

Lee, Y., C. Karunakaran, R. Lahlali, X. Liu, K. K. Tanino, and J. E. Olsen. 2017. "Photoperiodic
Regulation of Growth-Dormancy Cycling through Induction of Multiple Bud-Shoot Barriers
Preventing Water Transport into the Winter Buds of Norway Spruce." *Front Plant Sci* 8: 2109.
https://dx.doi.org/10.3389/fpls.2017.02109.

Lind, O. C., D. Helen Oughton, and B. Salbu. 2019. "The Nmbu Figaro Low Dose Irradiation Facility."
Int J Radiat Biol 95, no. 1 (01): 76-81. https://dx.doi.org/10.1080/09553002.2018.1516906.

Livak, K. J., and T. D. Schmittgen. 2001. "Analysis of Relative Gene Expression Data Using Real-Time
Quantitative Pcr and the 2(-Delta Delta C(T)) Method." *Methods* 25, no. 4 (Dec): 402-8.
https://dx.doi.org/10.1006/meth.2001.1262.

Mothersill, C., and C. B. Seymour. 1998. "Mechanisms and Implications of Genomic Instability and
Other Delayed Effects of Ionizing Radiation Exposure." *Mutagenesis* 13, no. 5 (Sep): 421-6.
https://dx.doi.org/10.1093/mutage/13.5.421.

Nisa, M. U., Y. Huang, M. Benhamed, and C. Raynaud. 2019. "The Plant Dna Damage Response:
Signaling Pathways Leading to Growth Inhibition and Putative Role in Response to Stress
Conditions." *Front Plant Sci* 10: 653. https://dx.doi.org/10.3389/fpls.2019.00653.

Nystedt, B., N. R. Street, A. Wetterbom, A. Zuccolo, Y. C. Lin, D. G. Scofield, F. Vezzi, N. Delhomme, S.
Giacomello, A. Alexeyenko, R. Vicedomini, K. Sahlin, E. Sherwood, M. Elfstrand, L. Gramzow,
K. Holmberg, J. Hällman, O. Keech, L. Klasson, M. Koriabine, M. Kucukoglu, M. Käller, J.
Luthman, F. Lysholm, T. Niittylä, A. Olson, N. Rilakovic, C. Ritland, J. A. Rosselló, J. Sena, T.
Svensson, C. Talavera-López, G. Theißen, H. Tuominen, K. Vanneste, Z. Q. Wu, B. Zhang, P.
Zerbe, L. Arvestad, R. Bhalerao, J. Bohlmann, J. Bousquet, R. Garcia Gil, T. R. Hvidsten, P. de
Jong, J. MacKay, M. Morgante, K. Ritland, B. Sundberg, S. L. Thompson, Y. Van de Peer, B.
Andersson, O. Nilsson, P. K. Ingvarsson, J. Lundeberg, and S. Jansson. 2013. "The Norway
Spruce Genome Sequence and Conifer Genome Evolution." *Nature* 497, no. 7451 (May):
579-84. https://dx.doi.org/10.1038/nature12211.

Patro, R., G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford. 2017. "Salmon Provides Fast and Bias-
Aware Quantification of Transcript Expression." *Nat Methods* 14, no. 4 (Apr): 417-419.
https://dx.doi.org/10.1038/nmeth.4197.

Pedroza-Garcia, J. A., Y. Xiang, and L. De Veylder. 2022. "Cell Cycle Checkpoint Control in Response
to Dna Damage by Environmental Stresses." *Plant J* 109, no. 3 (02): 490-507.
https://dx.doi.org/10.1111/tpj.15567.

Qi, F., and F. Zhang. 2019. "Cell Cycle Regulation in the Plant Response to Stress." *Front Plant Sci* 10:
1765. https://dx.doi.org/10.3389/fpls.2019.01765.

913
914   Sheppard, B.D. Amiro and S.C. 1994. *Effects of Ionizing Radiation on the Boreal Forest: Canada's Fig
915   Experiment, with Implications for Radionuclides*. Vol. 157: The Science of the Total
916   Environment.

917
918   Sidler, C., D. Li, O. Kovalchuk, and I. Kovalchuk. 2015. "Development-Dependent Expression of Dna
919   Repair Genes and Epigenetic Regulators in Arabidopsis Plants Exposed to Ionizing Radiation."
920   *Radiat Res* 183, no. 2 (Feb): 219-32. https://dx.doi.org/10.1667/RR13840.1.

921
922   Sylvestre-Gonon, E., S. R. Law, M. Schwartz, K. Robe, O. Keech, C. Didierjean, C. Dubos, N. Rouhier,
923   and A. Hecker. 2019. "Functional, Structural and Biochemical Features of Plant Serinyl-
924   Glutathione Transferases." *Front Plant Sci* 10: 608.
925   https://dx.doi.org/10.3389/fpls.2019.00608.

926
927   UNSCEAR. *Sources and Effects*

928   *of Ionizing Radiation*: United Nations Scientific Committee on the Effects of Atomic Radiation
929   UNSCEAR 1996 Report to the General Assembly, with Scientific Annexes.

930
931   van de Walle, J., N. Horemans, E. Saenen, M. Van Hees, J. Wannijn, R. Nauts, A. van Gompel, J.
932   Vangronsveld, H. Vandenhove, and A. Cuypers. 2016. "Arabidopsis Plants Exposed to
933   Gamma Radiation in Two Successive Generations Show a Different Oxidative Stress
934   Response." *J Environ Radioact* 165 (Dec): 270-279.
935   https://dx.doi.org/10.1016/j.jenvrad.2016.10.014.

936
937   Van Hoeck, A., N. Horemans, R. Nauts, M. Van Hees, H. Vandenhove, and R. Blust. 2017. "Lemna
938   Minor Plants Chronically Exposed to Ionising Radiation: Rna-Seq Analysis Indicates a Dose
939   Rate Dependent Shift from Acclimation to Survival Strategies." *Plant Sci* 257 (Apr): 84-95.
940   https://dx.doi.org/10.1016/j.plantsci.2017.01.010.

941
942   Vandenhove, H., N. Vanhoudt, J. Wannijn, M . Van Hees, and A . Cuypers. 2009. *Effect of Low-Dose
943   Chronic Gamma Exposure on Growth and*

944   *Oxidative Stress Related Responses in Arabidopsis Thaliana*. Vol. 44: Radioprotection.

945
946   Vanhoudt, N., N. Horemans, J. Wannijn, R. Nauts, M. Van Hees, and H. Vandenhove. 2014. "Primary
947   Stress Responses in Arabidopsis Thaliana Exposed to Gamma Radiation." *J Environ Radioact*
948   129 (Mar): 1-6. https://dx.doi.org/10.1016/j.jenvrad.2013.11.011.

949
950   Verma, V., P. Ravindran, and P. P. Kumar. 2016. "Plant Hormone-Mediated Regulation of Stress
951   Responses." *BMC Plant Biol* 16 (Apr 14): 86. https://dx.doi.org/10.1186/s12870-016-0771-y.

952
Vishwakarma, K., N. Upadhyay, N. Kumar, G. Yadav, J. Singh, R. K. Mishra, V. Kumar, R. Verma, R. G.
Upadhyay, M. Pandey, and S. Sharma. 2017. "Abscisic Acid Signaling and Abiotic Stress

Tolerance in Plants: A Review on Current Knowledge and Future Prospects." *Front Plant Sci*
8: 161. https://dx.doi.org/10.3389/fpls.2017.00161.

Weimer, A. K., S. Biedermann, H. Harashima, F. Roodbarkelari, N. Takahashi, J. Foreman, Y. Guan, G.
Pochon, M. Heese, D. Van Damme, K. Sugimoto, C. Koncz, P. Doerner, M. Umeda, and A.
Schnittger. 2016. "The Plant-Specific Cdkb1-Cycb1 Complex Mediates Homologous
Recombination Repair in Arabidopsis." *EMBO J* 35, no. 19 (10): 2068-2086.
https://dx.doi.org/10.15252/embj.201593083.

Xie, L., K. A. Solhaug, Y. Song, D. A. Brede, O. C. Lind, B. Salbu, and K. E. Tollefsen. 2019. "Modes of
Action and Adverse Effects of Gamma Radiation in an Aquatic Macrophyte Lemna Minor."
Sci Total Environ 680 (Aug 25): 23-34. https://dx.doi.org/10.1016/j.scitotenv.2019.05.016.

Yoshiyama, K. O., J. Kobayashi, N. Ogita, M. Ueda, S. Kimura, H. Maki, and M. Umeda. 2013. "Atm-
Mediated Phosphorylation of Sog1 Is Essential for the Dna Damage Response in
Arabidopsis." *EMBO Rep* 14, no. 9 (Sep): 817-22.
https://dx.doi.org/10.1038/embor.2013.112.

Yoshiyama, K. O., K. Sakaguchi, and S. Kimura. 2013. "Dna Damage Response in Plants: Conserved
and Variable Response Compared to Animals." *Biology (Basel)* 2, no. 4 (Nov): 1338-56.
https://dx.doi.org/10.3390/biology2041338.

Zimin, A. V., K. A. Stevens, M. W. Crepeau, D. Puiu, J. L. Wegrzyn, J. A. Yorke, C. H. Langley, D. B.
Neale, and S. L. Salzberg. 2017. "An Improved Assembly of the Loblolly Pine Mega-Genome
Using Long-Read Single-Molecule Sequencing." *Gigascience* 6, no. 1 (01 01): 1-4.
https://dx.doi.org/10.1093/gigascience/giw016.

Fig. 1   *Bhattacharjee et al. 2023*

A. thaliana	*P. abies*	Dose rates (mGy h^{-1})

Fig. 2   *Bhattacharjee et al. 2023*

Dose rate (mGy h⁻¹)

Fig. 3   *Bhattacharjee et al. 2023*

Fig. 4   *Bhattacharjee et al. 2023*

Fig. 5   *Bhattacharjee et al. 2023*

Fig. 6   *Bhattacharjee et al. 2023*

Fig. 7   *Bhattacharjee et al. 2023*

Figure legends

Fig. 1. Effect of 48 h of gamma irradiation from day 7 after sowing on growth in *A. thaliana* and Norway spruce (*P. abies*). A. Hypocotyl and B. cotyledon length in *A. thaliana* (n = 29-40) after the 48-h irradiation and C. morphology 77 days post-irradiation (dpi). D. Shoot and E. root length of Norway spruce (n = 28-35) after the 48-h irradiation and F. shoot diameter and G. number of needles in time courses (n = 11-22) and F. morphology 77 dpi. Scale bars: 3 cm. Means ± standard errors are shown. Different letters within each figure denotes significant difference (p ≤ 0.05) based on analysis of variance followed by Tukey`s test.

Fig. 2. Light micrographs of shoot apical meristems after 48 h of gamma irradiation from day 7 after sowing in *A. thaliana* and Norway spruce (*P. abies*). Five plants were investigated per treatment. Scale bars: 25 µm.

Fig. 3. Transmission electron micrographs showing cellular organelles after 48 h of gamma irradiation from day 7 after sowing in A – C *A. thaliana* and D - E Norway spruce (*P. abies*). nu: nucleus, no: nucleoli, mt: mitochondrion, ch: chloroplast, sb: starch body. Arrows indicate nuclear membrane. Arrowheads indicate disrupted mitochondria. Five plants were investigated per treatment. Scale bars: Scale nu: 1 µm, ch: 500 nm, mt: 500 nm.

Fig. 4. Effect of 48 h of gamma irradiation from day 7 after sowing on DNA damage detected by the COMET assay in A. *A. thaliana* and B. Norway spruce after the 48-h irradiation and day 77 post-irradiation. The line in each box represents the median values of 3 samples with 200 nuclei scored in each. Lower and upper box boundaries = 25 and 75 percentiles, error bars = 10 and 90 percentiles with data points outside these shown as dots. Different letters within each figure indicate significant difference (p ≤ 0.05) based on analysis of variance followed by Tukey's test.

Fig. 5. Number of differentially expressed genes (DEGs) and their $\log_2$ fold change after 48 h of gamma irradiation from day 7 after sowing relative to unexposed controls of A-B. *A. thaliana* and C-D Norway spruce, with false discovery rate (FDR) < 0.05 (n = 4). Overlap of upregulated (red numbers) and downregulated (blue numbers) genes at different dose rates in (b) *A. thaliana* and (d) Norway spruce.

Fig. 6. Relative transcript levels of DNA repair and cell cycle genes after 12, 24 and 48 h of gamma irradiation from day 7 after sowing in shoots of *A. thaliana*. Transcript levels were normalized against *ACTIN*, *ELONGATION FACTOR 1 α* and *UBIQUITIN 10* and shown relative to the unexposed controls at 12 h. The results are mean ± SE (n = 3-4 with 3-4 technical replicates). Different letters within each gene indicate significant difference (p ≤ 0.05) based on analysis of variance followed by Tukey's post-hoc test.

Fig. 7. Relative transcript level of DNA repair and cell cycle genes after 12, 24 and 48 h of gamma irradiation from day 7 after sowing in shoots of Norway spruce. Transcript levels were normalized against *ACTIN*, *ELONGATION FACTOR 1 α* and *UBIQUITIN 10* and shown relative to the unexposed controls at 12 h. The results are mean $\pm$SE (n = 3-4 with 3-4 technical replicates). Different letters within each gene indicate significant differences (p $\leq$ 0.05) based on analysis of variance followed by Tukey's post-hoc test.

Table 1. The gamma radiation dose rates and total doses applied in experiments with 48 h exposure of seedlings of *Arabidopsis thaliana* and Norway spruce from day 7 after sowing, using a ^{60}Co source.

Dose rate (mGy h^{-1})	Dose rate interval (mGy h^{-1})		Total dose (Gy) after 48 h exposure	Total dose interval (Gy) after 48 h exposure)	
	Minimum	Maximum		Minimum	Maximum
290	255	325	13.9	12.2	15.6
100	88	112	4.8	4.3	5.4
40	35	45	1.9	1.69	2.15
10	8,8	11,2	0.48	0.41	0.55
1	0,88	1,12	0.048	0.041	0.055
0	-	-	0	-	-