High radiosensitivity in the conifer Norway spruce (*Picea abies*) due to less

comprehensive mobilisation of protection and repair responses compared

to the radiotolerant *Arabidopsis thaliana*

Payel Bhattacharjee[1,2], Dajana Blagojevic[1,2+], YeonKyeong Lee[1,2,3+], Gareth B Gillard[4], Lars

Grønvold[4], Torgeir Rhoden Hvidsten[5], Simen Rød Sandve[4], Ole Christian Lind[2,6], Brit Salbu

[2,6], Dag Anders Brede[2,6], Jorunn E. Olsen[1,2]

[1]Department of Plant Sciences, Faculty of Biosciences, Norwegian University of Life

Sciences, Ås, Norway

[2]Centre for Environmental Radioactivity (CERAD CoE), Norwegian University of Life

Sciences, Ås, Norway

[3]Department of Plant Biotechnology, Korea University Graduate School, 145 Anam-ro,

Seongbuk-ku, Seoul, Republic of Korea

[4]Department of Animal and Aquacultural Sciences, Faculty of Biosciences, Norwegian

University of Life Sciences, Ås, Norway

[5]Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life

Sciences, Ås, Norway

[6]Department of Environmental Sciences, Faculty of Environmental Sciences and Natural

Resource Management, Norwegian University of Life Sciences, Ås, Norway

[+] These authors contributed equally.

Authors for correspondence: Payel Bhattacharjee, Jorunn E. Olsen

Email address: payel.bhattacharjee@nmbu.no

E-mail address: jorunn.olsen@nmbu.no

Abstract

Risk assessment and protection of plant communities in contaminated ecosystems require in-depth understanding of differential sensitivity to chronic ionising radiation in plants. However, the contributing molecular factors to differential radiosensitivity among plant species are poorly understood. To shed light on this, we compared early events associated with protection, repair, and stress responses in gamma-irradiated (1-290 mGy h^{-1}) seedlings of the radiosensitive conifer Norway spruce (*Picea abies*) and the radiotolerant *Arabidopsis thaliana*, by analysing growth, organelle and DNA damage, transcriptomes and the dynamics of antioxidant activities and expression of relevant genes. After 48 h of gamma radiation exposure, Norway spruce showed significantly reduced growth at 100-290 mGy h^{-1} and organelle damage, especially in mitochondria, at ≥ 1 mGy h^{-1} whereas *A. thaliana* showed normal vegetative growth at all dose rates, transiently delayed reproductive development at 290 mGy h^{-1} only, minor organelle damage only at ≥ 100 mGy h^{-1} and significantly less DNA damage than in Norway spruce at all dose rates. Comparative transcriptomics revealed that *A. thaliana* showed massive activation of genes related to DDR, antioxidants, and other stress responses at ≥ 1 mGy h^{-1} while Norway spruce mobilized transcription of such pathways only at ≥ 40 mGy h^{-1}. The transcriptional activation of repair and protection responses at higher gamma dose-rates only and its absence in lower dose-rates, corelates with high radiosensitivity of Norway spruce, compared to the massive transcriptional activation from low dose-rates in the radiotolerant *A. thaliana*.

1. Introduction

Stress tolerance capacity in plants in general is highly variable and a defining factor for species niche and habitat preference. Ionising radiation is a stressor that can induce harmful effects to organisms due to damage to biological macromolecules such as DNA, proteins and lipids (Caplin and Wiley, 2018). Field experiments and surveys of radionuclide-contaminated plant communities and ecosystems have shown that species that are sensitive to chronic ionising radiation may succumb and be replaced by more tolerant species (Amiro and Sheppard, 1994; Caplin and Willey, 2018). Laboratory studies have confirmed that radiosensitivity of different plant species ranges over orders of magnitude (Caplin and Wiley, 2018; Blagojevic et al., 2019a and b). However, the underlying mechanisms conferring tolerance to ionising radiation are not well understood. Such knowledge is needed to improve our ability to predict environmental consequences of nuclear accidents and release of radionuclides leading to exposure to high levels of ionising radiation.

The average natural background level of ionising radiation is estimated to 0.29 μGy h^{-1}, while up to 15 μGy h^{-1} has been measured in areas with naturally elevated radionuclide levels (Caplin and Willey, 2018). The nuclear power plant accident in Chernobyl in April 1986 as well as other accidents and experiments have shown that acute high doses (10-1000 Gy) can be lethal for plants and that radiosensitivity differs between species (UNSCEAR, 1996). In response to sub-lethal chronic radiation, adverse effects such as growth inhibition, reduced apical dominance (increased branching) and leaf chlorosis have been observed, particularly in conifers (Caplin and Wiley, 2018; Kashparova et al., 2020). A dose rate of 10 mGy d^{-1} (417 μGy h^{-1}) has been suggested as a threshold value in the sense that lower dose rates have "no detrimental effects for populations of terrestrial plants in the field" (International Atomic Energy Agency [IAEA], 1992; Caplin and Willey, 2018; UNSCEAR, 1996). However, this does not appear to hold true for boreal conifer forests. In a long-term field irradiation experiment in Canada persistent effects on forests including conifer trees were observed at chronic dose rates $\geq$ 100 μGy h^{-1} (Amiro and Sheppard, 1994). In addition to the differential radiosensitivity between species (Blagojevic et al., 2019a; Vanhoudt et al., 2014; Choi et al., 2021; IAEA, 1992 ; Blagojevic et al., 2019b), effects of ionising radiation depend on dose rate, duration (acute or chronic), total dose, life stages and the co-occurrence of other contaminants and environmental conditions (Garnier-Laplace et al., 2013).

The sensitivity of plant species to acute irradiation has been suggested to be attributed to the

efficiency of DNA repair systems and genome organisation and size (De Micco et al., 2011).

A. thaliana has a small genome of 135 Mb, whereas conifers like Norway spruce and Scots

pine have giant genomes of 20 and 23 Gb (Nystedt et al., 2013; Zimin et al., 2017). On the

other hand, the sensitivity to chronic irradiation appears to be regulated by delicate

mechanisms on different levels of biological organisation (Caplin and Willey, 2018; Kim et

al., 2019). However, factors that define sensitivity to chronic exposure are poorly understood

and apparently have functional rather than structural origin, depending on the molecular

architecture of stress-responsive pathways of different plant species.

Ionising radiation can cause damage both directly and indirectly via the formation of free

radicals and subsequent reactive oxygen species (ROS) (Caplin and Willey, 2018). As one of

the key forces of adaptive responses, non-lethal levels of ROS can modulate cellular

responses related to redox homeostasis and initiate retrograde signalling between organelles,

such as mitochondria, chloroplasts, and the endoplasmic reticulum, and the cell nucleus,

thereby affecting gene expression (Volkova et al., 2022). Plants possess antioxidant systems

involving enzymes such as catalases (CAT), superoxide dismutase (SOD) and peroxidases

(PER) as well as metabolites such as ascorbate, phenolic compounds and glutathione (GSH)

that can neutralise ROS and thereby prevent oxidative damage (Dumanović et al., 2020).

Studies of specific plant species including *A. thaliana*, have revealed induction of

antioxidant-related genes, including *SOD* and *ASCORBATE PEROXIDASE* (*APX*) following

ionising radiation exposure but there is limited information about antioxidant homeostasis in

conifers (van de Walle et al., 2016; Kim et al., 2011).

DNA strand breaks induced by ROS or directly by ionising radiation trigger the activation of

DNA damage repair (DDR) pathways (Kim et al., 2019). A range of homologous-

recombination (HR) repair-mediated DDR genes known to be activated by the plant-specific

transcription factor *SUPRESSOR OF GAMMA RESPONSE 1 (SOG1)* were significantly

upregulated by gamma irradiation at 100 Gy in *A. thaliana* (Bourbousse et al., 2018). The

initiation of the DDR is closely interlinked with cell cycle stalling through the activation of

cell cycle checkpoints by regulating cyclin (CYC)-cyclin-dependent kinase (CDK)-

complexes (Qi and Zhang, 2019), further leading to DNA damage-induced endoreduplication

(Nisa et al. 2019). In addition, transcriptional recruitment of DDR proteins in response to

gamma radiation requires chromatin remodelers involved in DNA methylation and histone

2A (H2A) modification (Choi et al., 2019; ; Kim et al., 2021). However, there is a lack of

knowledge related to molecular events associated with DDR and other relevant pathways in
conifers exposed to ionising radiation.

Knowledge of the factors determining the radiosensitivity of plant species is crucial for risk
assessment and the protection of ecosystems from ionising radiation hazard. Although
transcriptional regulation in response to very high doses of ionising radiation has been
investigated, albeit to a limited extent, in *A. thaliana* and a few other species (Choi et al.,
2021; Vanhoudt et al., 2014), the mechanisms leading to the differential radiosensitivity
among plant species are poorly understood. Hence, this study aimed to provide an overview
of mechanisms involved in plant radiosensitivity by employing gamma irradiation under
standardised conditions and concomitant analyses of growth, organelle- and DNA damage as
well as transcriptomics and dynamics of antioxidant activities and expression of repair- and
protection-related genes. In this respect, early physiological, cellular, and molecular effects
were compared in the radiosensitive conifer Norway spruce and the radio-tolerant *A. thaliana*
to provide novel insights into the observed effects in plant communities growing in areas
contaminated with radionuclides.

2. Materials and methods

2.1. Plant materials and pre-growing conditions

Seeds of the Landsberg erecta (L*er*) accession of *A. thaliana* (L) Heyhn, and Norway spruce
(*Picea abies* L. H. Karst) from the provenance CØ1 from Halden, Norway (59°N latitude,
Skogfrøverket, Hamar, Norway) were surface-sterilised and sowed according to the protocol
in (Blagojevic et al. 2019a). The seeds were germinated for 6 days at 20°C under a 16 h
photoperiod with a photosynthetic photon flux density (PPFD) of 30 μmol m^{-2} s^{-1} at 400-700
nm (TL-D 58W/840 lamps, Philips, Eindhoven, The Netherlands), as measured at the top of
the Petri dishes with a Li-Cor Quantum/Radiometer/Photometer (LI-250, LI-COR, USA).

2.2. Gamma irradiation of seedlings and growing conditions during the exposure

Six days old seedlings of *A. thaliana* and Norway spruce were exposed for 48 h using the
FIGARO gamma irradiation facility (^{60}Co; 1173.2 and 1332.5 keV γ-rays) at the Norwegian
University of Life Sciences (Lind et al, 2019). Petri dishes (3.5 cm diameter) with seedlings
were placed at different distances from the gamma source to obtain the average dose rates to
water 1, 10, 20, 40, 100 and 290 mGy h^{-1} (Table 1). These 48 h- irradiation experiments were
repeated three times (Table S1. For one of these experiments, the highest dose rate was 250

mGy h^{-1} as 290 mGy h^{-1} could not be obtained due to the decay of the ^{60}Co source. In

additional experiments aiming at studies of dynamic effects early during gamma irradiation

on gene expression and antioxidant contents, seedlings of both species were exposed to 0, 1,

10, 20, 40 and 100 mGy h^{-1} for 12, 24 or 48 h, with 2 repeated experiments for each duration

(Table S1). The dose rates to water and dose rate intervals (in front and back of the petri

dishes) were calculated from the air kerma rates according to (Hansen et al., 2019) as

previously described (Blagojevic et al., 2019a). The total doses were calculated by

multiplying the absorbed dose rates to water and exposure duration. During the exposure the

room temperature was 20˚C and the photoperiod 12 h with an irradiance of 55 μmol m^{-2} s^{-1}

from high pressure metal halide lamps (HPI-T Plus 250W lamps, Philips, The Netherlands).

The red:far red (R:FR) ratio was measured to 3.5 using a 660/730 nm Skye sensor (Skye

Instruments, UK).

2.3. Post-irradiation growing conditions

171 The post-irradiation effects of the 48 h of gamma exposure were recorded after transfer of

172 seedlings to pots filled with S-soil (Hasselfors Garden AS, Örebro, Sweden), one plant per

173 pot. All plants were kept in growth chambers (manufactured by Norwegian University of Life

174 Sciences) with temperature, relative air humidity, R:FR ratio, and photoperiod maintained as

175 previously described (Blagojevic et al., 2019a).

2.4. Growth parameter recordings

At the end of the gamma irradiation, seedlings from both species were placed between two

transparent plastic sheets with a millimetre paper on top and scanned. The shoot and root

lengths of Norway spruce and the lengths of the hypocotyl and cotyledons in *A. thaliana* were

measured using Image J (imagej.nih.gov) with 28-35 and 29-40 seedlings per gamma dose

rate for the respective species.

Post-irradiation, the number of rosette leaves as well as the number of plants with floral buds,

inflorescence and open flower buds were recorded in 23-24 *A. thaliana* plants per gamma

dose rate in time courses up to 75 days post-irradiation (dpi), and percentages of plants with

floral buds, inflorescence and open flowers were calculated. Leaf width and leaf length were

measured at 34 and 41 and height of the inflorescence at 64 dpi. In Norway spruce, shoot

elongation, the number of needles and shoot diameter were recorded in a time course of 59

dpi in 11-22 plants per dose rate. The shoot diameter was calculated by averaging two

perpendicular measurements from needle tip to needle tip across the plant at the shoot apex.

Also, plant height was measured from the pot edge to the shoot apical meristem and the
cumulative shoot elongation was calculated.

2.5. Histological and cytological studies of shoot tips

Histological studies of shoot apical meristems (SAMs) at the end of the gamma irradiation
were performed according to (Lee et al., 2017). Shoot tips of Norway spruce and *A. thaliana*
were then fixed in 4 % paraformaldehyde and 0.025 % glutaraldehyde in phosphate buffered
saline (PBS, pH 7.0), vacuum infiltrated at room temperature and stored at 4°C. The fixed
tissues were washed in PBS solution and dehydrated in a graded ethanol solution.
Approximately 3 mm of the shoot tips of each of 5 plants per species per treatment were
infiltrated and embedded in LR White (London Resin Company, UK). After polymerisation,
1 μm sections were made using an Ultracut Leica EM UC6 microtome (Leica, Germany),
stained with toluidine blue O and inspected using a light microscope with bright field optics
(DM6B, Leica).

For cytological studies, LR white-embedded samples from 5 plants per species per treatment
were sectioned into 70 nm ultrathin sections, (Leica EMUC6), stained with uranyl acetate and
lead citrate before examining by transmission electron microscopy (TEM) (FEI Morgagni
268, United States).

2.6. Comet assay for analysis of DNA damage

Single and double-strand DNA breaks at the end of the gamma irradiation and 77 dpi were
quantified using the comet assay (Gichner, 2003) with some modifications et al. 2019a).
Briefly, this included isolation of cell nuclei and electrophoresis under alkaline pH, resulting
in a comet-like appearance of damaged DNA due to movement out of the nuclei. This was
followed by quantification of the DNA in the comet tail using fluorescence microscopy.
Three biological replicates per dose rate, each consisting of 3-4 mm of shoot tips from 3
plants of Norway spruce or 5 plants of *A. thaliana*, were used to assess DNA damage. For
each sample, about 180-200 cell nuclei were scored, with 60-70 nuclei scored in each of 3
technical replicates (gels) and the median value was calculated.

2.7. RNA isolation for RNA sequencing

At the termination of the 48-h gamma irradiation, four biological replicates per gamma dose
rate (0, 1, 10, 40, 100 (both species) and 290 mGy h^{-1} (*A. thaliana* only due to lack of growth
inhibition, organelle and DNA damage at lower dose rates)) were harvested in liquid

nitrogen. For *A. thaliana* (20-30 shoots per dose rate), total RNA was extracted using the

RNeasy Plant Mini Kit (QIAGEN Inc., Valencia, CA) following the manufacturer's protocol.

For Norway spruce (four shoots per dose rate), total RNA was extracted using the Masterpure

Complete DNA and RNA Purification Kit (Epicenter, Madison, USA) following the

manufacturer's protocol, except that 3 µl beta-mercaptoethanol per sample replaced 1,4-

dithiothreitol (DTT) and 0.5% polyvinylpyrrolidone (PVP, mw 360 000, Sigma Aldrich,

Germany) was added to the extraction buffer. The purified RNA was quantified using a

NanoDrop ND-1000 Spectrophotometer (Thermo Scientific, USA) and the quality assessed

with an Agilent 2100 Bioanalyzer with an RNA 144000 Nano Kit (Agilent technologies,

USA).

2.8. RNA sequencing

The transcriptomes of the shoots were sequenced with two repeated lanes per sample, on an

Illumina HiSeq4000 platform at the Norwegian Sequencing Centre (Oslo, Norway) using the

Strand specific 20xTruseq RNA library preparation (paired end, read length 150 bp) as

recommended by the manufacturer (Illumina, San Diego, CA, USA). The raw sequence data

were submitted to the ArrayExpress under the accession number E-MTAB-8081 for Norway

spruce and E-MTAB-11120 for *A. thaliana*.

2.9. Differential gene expression analyses and functional term enrichment

The transcript abundances were estimated using the Salmon program (Patro et al., 2017),

comparing with the genomes of either Norway spruce (*Picea abies* 1.0 assembly,

plantgenie.org) (Nystedt et al., 2013) or *A. thaliana* (TAIR10, arabidopsis.org) (Berardini et

al. 2015). Gene expression was compared between the samples at different gamma radiation

dose rates and the unexposed control samples using the DESeq2 (Love et al., 2014). Genes

were classified as differentially expressed genes (DEGs) if the False Discovery Rate

(Benjamini and Hochberg, correction) adjusted p-values were < 0.05.

Gene Ontology (GO) annotations of Norway spruce genes, as well as spruce to *A. thaliana*

orthologs by best sequence matches, were downloaded from plantgenie.org. The *A. thaliana*

orthologs were used to associate the Norway spruce genes with KEGG pathways. Enrichment

of GO or KEGG terms in sets of DEGs was found using topGO (version 2.34.0) or kegga

from the limma package (version 3.38.2), respectively.

2.10. RT-qPCR analysis of dynamics in gene expression under ionising radiation

Based on the significant regulation in one or both species in response to 48 h gamma

irradiation, 11 DNA repair- and cell cycle control genes as well as 4 antioxidant- and 2

retrograde signalling-related genes were selected for RT-qPCR analysis of their dynamic

regulation after 12, 24 and 48 h of gamma irradiation at 0, 1, 10, 20, 40 and 100 mGy h^{-1}

(Table S2). RNA was isolated as described above from four biological replicates per dose

rate, each consisting of shoots from eight and 20-30 seedlings for Norway spruce and *A.*

thaliana, respectively. cDNA was synthesised using SuperScript™ IV VILO™ cDNA

Synthesis Kit (Invitrogen, Carlsbad, CA, US) following the manufacturer's protocol, and no

enzyme controls (-RT) were made for each sample.

Gene sequences were retrieved from NCBI (ncbi.nlm.nih.gov/gene) and TAIR

(arabidopsis.org) for *A. thaliana* and plantgenie.org for Norway spruce and exon-exon-

spanning primers were designed to avoid non-specific amplification from any genomic DNA

contamination in the RNA samples, using the online tools Oligoanalyzer

(eu.idtdna.com/calc/analyzer) and Primer Blast (ncbi.nlm.nih.gov/tools/primer-blast) with

default parameters. Amplifications were performed in a 7500 Fast Real-time PCR system

(Applied Biosystems, Foster city, USA) using Platinum Quantitative PCR Supermix-UDG

and SYBRGreen (Thermo Fisher, Carlsbad, CA, US), using standard protocol.

To compare the relative transcript levels of the selected genes in the samples exposed to

different gamma radiation dose rates at different time points with the unexposed controls (0

mGy h^{-1}), the comparative ΔΔCT method was used (Livak and Schmittgen, 2001). The

expression of the target genes was normalised by the mean expression of the three reference

genes, *ACTIN* (*ACT*), *ELONGATION FACTOR 1α* (*EF-1α*) for both species and *UBIQUITIN*

and *α-Tubulin* for *A. thaliana* and Norway spruce respectively.

2.11. Antioxidant assays

Shoot samples for analyses of PER and CAT concentration, SOD activity and flavonoid

content in seedling shoots of both plant species were flash-frozen in liquid nitrogen after 12,

24 and 48 h of gamma irradiation at 0, 1, 10, 20, 40 and 100 mGy h^{-1}. Samples of 50 mg -100

mg were homogenized maintaining freezing conditions with stainless steel beads in a tissue

homogenizer (Retsch, Haan, Düsseldorf, Germany). The samples were then mixed with

appropriate buffers according to the protocol for each of the different antioxidant kits used

(OxiSelect, Cell Biolabs. Inc., San Diego, CA, USA). The mixtures were vortexed and

subsequently centrifuged to precipitate cellular debris, and the clear lysates were collected for

the assays. The assays were performed in room temperature following the instructions in the manufacturers' protocol for each antioxidant. Colorimetric analyses were done in a plate reader (Microplate reader, Biochrom (ASYS UVB340)). For each antioxidant, three (peroxidase, catalase, SOD) or four (flavonoid) biological replicates, each consisting of materials from 20-30 shoots for *A. thaliana* and 4 shoots for Norway spruce, were analysed per dose rate and gamma irradiation duration with 2-3 technical replicates of each.

2.12. Statistical analysis for growth recordings, DNA damage, qPCR and antioxidant assay

DNA damage, hypocotyl, cotyledon and shoot length, as well as post-irradiation growth parameters at the final sampling time (34, 41 or 64 dpi for *A. thaliana* and 59 dpi for Norway spruce) were tested for normality and homoscedasticity by the Ryan-Joiner and Levene tests, respectively. One-way ANOVA tests (glm model in Minitab 21.1, Minitab Inc, PA, USA) were then used to test for differences in these response variables between plants exposed to 48 h of gamma radiation versus unexposed control plants. Due to the lack of normality and homoscedasticity of the non-transformed RT-qPCR gene expression and antioxidant analyses data, log-transformed values were used in two-way ANOVA for individual species (dose rate and gamma-irradiation duration as factors) and three-way ANOVA for comparison of species (plant species, dose rate and gamma-irradiation duration as factors). Tukey`s post hoc test was used to test for differences between means ($p \leq 0.05$).

3. Results

3.1. Gamma irradiation for 48 h affects growth more negatively in Norway spruce but not A. thaliana

In *A. thaliana*, no significant effect of the 48-h of gamma irradiation on hypocotyl and cotyledon lengths was identified (Fig. 1A, B). Also, the number of rosette leaves developing post-irradiation was not significantly affected (Fig. 1C). Only a slight, significant reduction in the width and length of rosette leaves was observed at the highest dose rate compared to the unexposed control at 34 and 41 dpi, but apart from this, there was no clear dose response-relationship (Fig. S1A, B). Although post-irradiation flowering was slightly delayed at 250-290 mGy h^{-1}, eventually all plants flowered (63 dpi) and showed very similar overall morphology, flowering, and silique development (77 dpi) (Fig. 1C, S1D, E, F, G). In Norway spruce, the shoot length was significantly reduced (27%) after 48 h of gamma irradiation at 290 mGy h^{-1}, but the root length was not affected (Fig. 1D, E). At 59 dpi, the shoot diameter and number of needles were significantly reduced by 5% and 12%, at 100 mGy h^{-1} and 9%

316 and 35% at 290 mGy h^{-1} whereas the cumulative growth did not differ significantly between
317 the gamma dose rates but (Fig. 1F, G, H, Fig. S2). However, the growth inhibition effects
318 was no longer as evident at 77 dpi (Fig. 1H).

3.2. More severe gamma radiation-induced organelle damage in Norway spruce than A.
thaliana

Normal dome-shaped shoot apical meristems with compact cells were observed in both
species after 48 h of gamma irradiation (Fig. 2). In *A. thaliana*, slightly damaged
mitochondria were observed at 100-290 mGy h^{-1}, but the chloroplasts and cell nuclei
appeared normal at all dose rates compared to the unexposed control except that a few cells
showed less dense nuclei at ≥ 1 mGy h^{-1} (Fig. 3A, B, C). In comparison, in Norway spruce,
disrupted nuclear and mitochondrial membranes were observed at ≥ 1 mGy h^{-1}, as well as
swollen chloroplasts with large starch bodies and disrupted thylakoid membranes at ≥ 40
mGy h^{-1} (Fig. 3D, E, F).

3.3 Higher levels of DNA damage in gamma-irradiated Norway spruce than A. thaliana

In *A. thaliana*, the Comet assay showed low but significant damage with ≤ 5% tail DNA at 1-
100 mGy h^{-1} and 12% tail DNA at 290 mGy h^{-1} (Fig. 4A). In comparison, DNA damage was
consistently 2-8-fold higher in Norway spruce with 10%, 12%, 15%, 17.5% and 40% DNA in
the Comet tail after the 48-h of irradiation at 1, 10, 40, 100 and 290 mGy h^{-1}, respectively
(Fig. 4B). At 77 dpi, the DNA damage was substantially reduced but still observed in *A.*
thaliana at ≥ 10 mGy h^{-1} with ≤ 3% tail DNA and in Norway spruce seedlings at ≥ 1 mGy h^{-1},
with ≤ 5% tail DNA at all dose rates (Fig. 4A, B).

3.4. Profound transcriptome differences between gamma-irradiated Norway spruce and A.
thaliana

Comparative RNA-seq analysis of gamma-irradiated shoot samples (1-100 mGy h^{-1} both
species, 290 mGy h^{-1} *A. thaliana* only), revealed extensive differences between the species.
A. thaliana expressed a total of 26,643 genes, of which 11,592 were DEGs (43.5%) between
any dose rate and the unexposed control in one or multiple gamma dose rates (Fig. 5A, upper
panel). Norway spruce expressed a total of predicted 66,069 genes, of which 5326 were
DEGs (8.1%, Fig. 5A, lower panel). In *A. thaliana*, a high number of DEGs was observed
already from 1 mGy h^{-1} (3442 DEGs), with the higher dose rates having ~4400-5500 DEGs
(Fig. 5A, ,). At 100-290 mGy h^{-1}, there was a higher proportion of upregulated genes with

more than 2-fold change as compared to the lower dose rates (Fig. 5A,). By contrast, in
Norway spruce, very few DEGs were observed at 1-10 mGy h^{-1} (38 and 43 DEGs), while 822
and 5047 DEGs were observed at 40-100 mGy h^{-1}, respectively (Fig. 5B, C). The increase in
DEGs from 40 to 100 mGy h^{-1} was much more pronounced in Norway spruce than in *A.*
thaliana (Fig. 5A,). The largest overlap of *A. thaliana* DEGs was between 100 and 290 mGy
h^{-1}, and between 1, 10 and 40 mGy h^{-1} (Fig. 5B). A good proportion of Norway spruce DEGs
at 40 and 100 mGy h^{-1} overlapped, showing some consistency in the response even with a
large difference in total numbers of DEGs, including many unique ones (Fig. 5C).

3.5. Distinctive pathway enrichment in Norway spruce and A. thaliana at different dose rates

GO and KEGG pathway analyses on sets of DEGs revealed a gamma dose rate-dependent
enrichment of different pathways. In *A. thaliana*, DNA replication, DNA repair, cell cycle,
endoreduplication, and endoplasmic reticulum (ER) stress were enriched at 1-40 mGy h^{-1}
(Fig. S3, S4). Hormone biosynthesis and signalling pathways showed considerable
enrichment along with stress response at 100-290 mGy h^{-1} (Fig. S3, S4). Oxidative stress,
defence response, growth and development, protein synthesis and catabolism as well as
mitogen-activated protein kinase (MAPK)-mediated signalling were significantly regulated at
all dose rates (Fig. S3, S4). In Norway spruce, pathways such as DNA replication, DNA
repair, cell cycle, endoreduplication, unfolded protein response (UPR) and ER stress,
response to gamma, energy metabolism and growth and development were all enriched at ≥
40 mGy h^{-1} (Fig. S5, S6). Photosynthesis and protein biosynthesis and metabolism were
enriched in downregulated DEGs at 100 mGy h^{-1} (Fig. S5, S6) in Norway spruce only.

In *A. thaliana* KEGG pathway analysis revealed that the DDR pathways such as HR,
mismatch repair (MMR), non-homologous end-joining (NHEJ) and base excision repair
(BER) were regulated at 10-40 mGy h^{-1}, and BER also at 290 mGy h^{-1} (Fig. S4). In Norway
spruce, NER, MMR, HR and NHEJ pathways were significantly regulated at 40-100 mGy h^{-1}
and BER at 40 mGy h^{-1} only (Fig. S6).

3.6. Distinct differences in stress response pathways between Norway spruce and A. thaliana

The specific DEGs in DDR, antioxidant protection and other stress response pathways were
further analysed to compare *A. thaliana* and Norway spruce response after the 48-h of gamma
irradiation.

3.6.1. DNA damage repair

In *A. thaliana*, most DEGs in the DDR pathways showed a 30-70% change in expression

(Table S3). Genes involved in DNA double strand break (DSB) repair via HR, including

RAD51, *SMC6B* and *BRCA2B*, showed significant upregulation at all dose rates in the 10-290

mGy h^{-1} range. In Norway spruce, most DDR DEGs at 40-100 mGy h^{-1} had at least a 60%

change in expression (Table S3), and genes involved in DSB repair via HR; *SRS2*, *RAD52*,

RPA1A, *MCM8*, *XRCC2* and *XRCC3*, were all upregulated at 100 mGy h^{-1}. A few other genes

such as *PARP1* and *PARP2* were significantly upregulated at the highest dose rates in both

species. Some DEGs showed opposite changes in expression between species; *MRE11*,

RPA2A/ROR1 and *BRCA1* in the HR pathway, and *MSH7* in the MMR pathway. Unique

DEGs to Norway spruce included upregulation of *KU80* at 40-100 mGy h^{-1} and *SOG1* at 100

mGy h^{-1}. Unique to *A. thaliana* was upregulation of *ATM* at 100 mGy h^{-1}.

3.6.2. Cell cycle and endoreduplication

In *A. thaliana*, most cell cycle-related DEGs including *CYCB*s and *CDKB*s, were observed at

1-40 mGy h^{-1} but not 100-290 mGy h^{-1} (Table S4). In Norway spruce, no such effects on B

type cyclins were observed and most of the cell cycle genes showed changes in expression at

≥ 40 mGy h^{-1}. Anaphase-promoting factors *APC6* and *13* and the cyclin-dependent kinases

regulatory subunit *CKS1* showed upregulation at ≥ 40 mGy h^{-1}. Positive regulators of the cell

cycle such as D type cyclins (e.g., *CYCD1*) were significantly upregulated in *A. thaliana* at ≤

40 mGy h^{-1} but downregulated in Norway spruce at 100 mGy h^{-1}. *RBR1* was upregulated in

both species but at 100 mGy h^{-1} only in Norway spruce and at most dose rates in *A. thaliana*.

E2FB was upregulated at 40 mGy h^{-1} in Norway spruce only. *WEE1,* an important G2/M

checkpoint regulator, was upregulated at ≤ 40 mGy h^{-1} in *A. thaliana*, but at ≥ 40 mGy h^{-1} in

Norway spruce. Most of the cell cycle-related genes, such as *CYC*s and *CDKs* that are also

involved in the endocycle, and the endoreduplication-related biological pathway, were

enriched in *A. thaliana* at ≤ 40 mGy h^{-1} but fewer such DEGs were observed in Norway

spruce (Table S4, S5). A few endoreduplication genes such as *APC5* and *APC6* were

upregulated in Norway spruce at 100 mGy h^{-1} but not in *A. thaliana* (Table S5).

3.6.3. Chromatin remodelers and genes in epigenetic pathways

Significant activation of genes involved in the pathway associated with de-condensation of

the nucleosome structure and activation of DDR pathways, was observed in Norway spruce at

40-100 mGy h^{-1} (Table S6). By contrast, in *A. thaliana*, a number of such genes were

downregulated at ≥ 100 mGy h^{-1}, and several were upregulated at ≤ 40 mGy h^{-1} only (Table

S6). The DNA helicase *DDM1* was upregulated in *A. thaliana* at ≤ 40 mGy h^{-1} and DNA

methyltransferase *CMT3,* histone 2A variant *HTA8* and a component of the core chromatin

remodelling complex *ARP6* were significantly downregulated at 100-290 mGy h^{-1} (Table S4,

S6). In Norway spruce, histone deacetylase *HOS15* and genes involved in the regulation of

DNA methylation, such as *DMT1*, *IDN2* and *SUVH4*, as well as the histone 2 A variant

H2AV, showed significant upregulation at 100 mGy h^{-1}.

3.6.4. Hormone biosynthesis and signalling

A. thaliana genes involved in biosynthesis and signalling of main growth-promoting

hormones auxin, cytokinin, cytokinin, gibberellin (GA) and brassinosteroid, as well as

important stress-response-related hormones such as abscisic acid (ABA) and ethylene, were

significantly regulated in response to the gamma irradiation (Table S7) especially at higher

dose rates. The auxin-responsive MYB transcription factors *MYB93* and *MYB61*, the auxin-

responsive genes *IAA13, IAA14*, *IAA27, SAUR10* and *GH3.6*, the auxin biosynthesis gene

YUC9 as well as the auxin efflux carriers and transporters involved in polar auxin transport,

PIN2, *PIN3*, *AUX1* and several ABC transporters, were significantly upregulated in this

species (Table S7) but downregulated in Norway spruce. Although the auxin-homeostasis-

related gene *IAMT1*, and the auxin responsive genes *SAR1* and *SAUR76* showed increased

expression at 40-100 mGy h^{-1} in Norway spruce, several auxin signalling genes including

ARF6, TIR1, AIR9 and *IAA26* were significantly downregulated (Table S7). In *A. thaliana*,

cytokinin biosynthesis, metabolism as well as cytokinin signalling genes such as *CKX5*,

LOG7, AHP1, SOT8, ARR2 and *ARR3*, were upregulated at 40-290 mGy h^{-1} (Table S7).

Contrastingly, none of these genes showed significant regulation in Norway spruce.

Similarly, the GA biosynthesis genes *GA2* (*ent-kaurene synthase 1*) and *GA 20-OXIDASE 2*

(*GA20OX2),* and the brassinosteroid biosynthesis and homeostasis genes *CYP90C1* and

CYP85A1, were upregulated in *A. thaliana* but not in Norway spruce.

In general, ABA biosynthesis and signalling genes did not show any gradual up- or

downregulation in *A. thaliana* (Table S7). However, the negative regulator of ABA

biosynthesis *PUB44*, and the ABA signalling genes *PYL4, CRK29* and *ATHB-6* were

downregulated at < 40 mGy h^{-1}. Many stress- and ABA-responsive genes such as *ASPG1,*

PPCK1, and *RD22* showed consistent increasing expression from 10-290 mGy h^{-1}. Among

the genes regulated in both species, the ABA signalling gene *CIPK3* showed downregulation

in all dose rates in *A. thaliana*, and at 100 mGy h^{-1} in Norway spruce, and *ADH1* were

upregulated in both species at higher dose rates. In *A. thaliana*, ethylene biosynthesis and signalling genes were mostly downregulated at lower dose rates with few exceptions (Table S7). Ethylene-responsive genes such as *CTR1, RAP2-2, RAP2-12,* and *ERFs* as well as the ethylene biosynthesis genes *EOL1* and *ACS5* showed significant regulation in *A. thaliana*, but not in Norway spruce.

3.6.5. Genes in antioxidant metabolism pathways

Significant overall upregulation of antioxidant biosynthesis and signalling genes was observed in *A. thaliana* even from 1 mGy h^{-1}. By contrast, in Norway spruce, most of the antioxidant genes were not significantly affected below 100 mGy h^{-1} (Table S8). The key flavonoid and anthocyanin biosynthesis genes *CHS* and *DFRA* were upregulated in *A. thaliana* from 1 mGy h^{-1} but significantly downregulated in Norway spruce at 100 mGy h^{-1}. A flavanol synthase gene (*FLS1*) was upregulated in *A. thaliana* at ≤ 10 mGy h^{-1} and 290 mGy h^{-1} and Norway spruce at 100 mGy h^{-1}. The *PER* gene *PER3* was regulated in both species but from lower dose rates in *A. thaliana* than Norway spruce. *GLUTATHIONE-S-TRANFERASEs* (*GSTU/GSTF*) were upregulated in both species.

3.6.6. Cell wall component genes

In *A. thaliana* genes involved in the biosynthesis of cell wall components, cell wall maintenance and fortification, as well as membrane receptors were significantly upregulated at 100-290 mGy h^{-1}, commonly with large fold-changes. In Norway spruce, only very few genes related to cell wall biogenesis and maintenance showed regulation, i.e., at 40-100 mGy h^{-1} (Table S9). In *A. thaliana*, *BOR2* involved in cellulose and pectin metabolism, *HHT1* involved in pectin and suberin biosynthesis, the cell wall organisation and biosynthesis genes *FACT* and *CYP86B1*, were upregulated at 100-290 mGy h^{-1}. The endoglucanase *GH9C2* involved in cell wall organisation were upregulated in all dose rates. None of these genes showed any significant changes at any dose rate in Norway spruce. A few other genes involved in biogenesis and metabolism of expansins, cellulose and xyloglucan such as the *EXP4* and *8* and *CESA5* were significantly regulated in both species with downregulation in Norway spruce and upregulation in *A. thaliana*, whereas *XTH5* was upregulated in Norway spruce at 100 mG h^{-1}

3.6.7. Photosynthesis and energy metabolism

Photosynthesis was massively downregulated in Norway spruce but not in *A. thaliana*. Photosystem (PS) I and II-related genes such as PS I reaction centre subunits *PSAG, PSAK* and *PSAN*, PS 1 light harvesting complex *LHCA1*, PS II assembly factor *HCF136*, essential enzymes such as NADPH-protochlorophyllide oxidoreductase *PORA*, and Fructose-1,6-bisphosphatase *CYFBP*, chloroplast binding proteins and membrane efflux carriers, were downregulated in Norway spruce at 100 mGy h^{-1} (Table S10). In *A. thaliana*, many genes were upregulated, such as the PS I and II reaction centre components *PSAB* and *PSBB*, and *NDHC* (Table S10).

Similarly, energy metabolism was far more affected in Norway spruce than *A. thaliana* (Table S11). Norway spruce particularly showed upregulation at $\geq$ 40 mGy h^{-1} of a wide range of genes involved in glycolysis, TCA cycle and gluconeogenesis, such as, Glyceraldehyde-3-phosphate dehydrogenase *GAPC1*, *MALATE DEHYDROGENASE* (*MDH1*), *SUCCINATE DEHYDROGENASE (SDH2-1)*, and several homologs of phosphoenolpyruvate carboxykinase *PCKA* and *PFK2*. Additionally, the *CYTOCHROME C1* (*CYTC-1*) and *CYTC OXIDASE 6B1* (*COX6B-1*) genes involved in mitochondrial electron transport were significantly upregulated at 100 mGy h^{-1}. *ALCOHOL DEHYDROGENASE 1* (*ADH1*) was significantly upregulated in both species from 100 mGy h^{-1}.

3.7. Dynamic regulation of repair and protection-related genes in A. thaliana and Norway spruce during gamma irradiation

To evaluate the dynamics of gene expression during radiation stress, both species were exposed to 12, 24 and 48 h of gamma radiation and the transcript levels of selected genes analysed by RT-qPCR (Fig 6, Fig. S7, S8). Consistent with the RNA-seq data, significantly increased transcript levels of the DDR-related *WEE1, PARP2, RAD51, BRCA1* and the cell cycle-related *CYCB1.2* were observed after 48 h in *A. thaliana*, by 5-10-fold as compared to the unexposed controls (Fig. 6, Fig S7). However, after 12 and 24 h exposure no significant effect of the gamma irradiation on their expression was observed. Also, the expression of *SOG1*, *BARD1*, *KU80*, *MSH7*, *PCNA* and *CYCB1.1* did not differ between the controls and any of the gamma dose rates. Of these DDR- and cell cycle-related genes in Norway spruce, the transcript levels of *WEE1* were significantly higher at 20-100 mGy h^{-1} after 24 h only compared to the unexposed controls and those of *SOG1* increased at 40-100 mGy h^{-1} after 24 h, while *PARP2* expression increased significantly by 10-30-fold at 40-100 mGy h^{-1} after 24 and 48 h. There were no significant effects on expression at 12 h.

Out of the four antioxidant genes analysed (*DFRA1* and *CHS1* (phenolic/flavonoid
biosynthesis)*, GSTU17* and *PER52), GSTU17* showed significant downregulation in *A.*
thaliana at 48 h at all dose rates except 10 mGy h^{-1}, but no such regulation was observed for
Norway spruce (Fig. 6, Fig. S8). Additionally, *CHS1* showed significant upregulation in *A.*
thaliana at 12 and 24 h after exposure to 1 and 100 mGy h^{-1} (Fig. S8). Moreover, amongst the
two analysed marker genes of retrograde signalling, *AOX1* (alternative oxidase) and *WRKY40*
(a transcription factor)*, WRKY40* showed small but significant downregulation and 100 mGy
h^{-1} in 12 and 24 h of radiation in *A. thaliana* (Fig. S8)*.* However, no significant regulation of
these genes was observed in Norway spruce.

3.8. Antioxidants in A. thaliana and Norway spruce during gamma irradiation

To evaluate antioxidant dynamics, the activities/concentrations of CAT, PER, SOD, and
flavonoids were investigated in *A. thaliana* and Norway spruce seedlings exposed to 12, 24
and 48 h of gamma radiation (Fig. 7). Although the concentrations of PER and CAT differed
significantly between the two species with *A. thaliana* being highest in PER and Norway
spruce in CAT, there were generally no significant effects of the gamma irradiation.
However, compared to the unexposed controls, the SOD activity in Norway spruce showed
significant increase after 48 h at 40-100 mGy h^{-1}. There was no effect of gamma radiation
after 12 and 24 h radiation in Norway spruce and no significant regulation of SOD in *A.*
thaliana. Flavonoid concentrations showed significant downregulation with time in *A.*
thaliana regardless of gamma irradiation or not. In Norway spruce, no such effect was
observed.

4. Discussion

To improve the knowledge base for risk assessment and protection of plant communities in
radionuclide-contaminated ecosystems against ionising radiation, the current comparative
study has focussed on underlying mechanisms with respect to differential radiosensitivity in
the highly radiosensitive conifer Norway spruce and the radiotolerant *A. thaliana*. Through
exposure of seedlings to gamma radiation under standardised conditions and comparative
assessment of phenotypic, cellular, and genotoxic effects, combined with RNA-seq and
studies of dynamics of antioxidant contents and expression of relevant genes, we present
novel understanding of differences between the two species in the mechanisms involved in
protection and repair.

Previously, we have shown dose rate-dependent impaired development in Norway spruce

seedlings exposed to gamma radiation for 144 h, with significant growth inhibition at $\geq$ 40

mGy h^{-1} and progressively developing after-effects eventually causing mortality (Blagojevic

et al., 2019a). By contrast, *A. thaliana* showed no significant adverse effects at dose rates $\leq$

540 mGy h^{-1} except a transient delay in reproductive development at the highest dose rates

(Blagojevic et al., 2019a). Consistent with the previous findings, in this study *A. thaliana* did

not show any clear negative effect on growth after 48 h irradiation up to 290 mGy h^{-1} except

a slight transient delay in floral development, whereas Norway spruce showed distinct growth

inhibition (Fig. 1, S1). In agreement with this, the RNA-seq analysis revealed profound

differences between the two species in response to gamma irradiation (Fig. 5, S3-S6). The

fact that *A. thaliana* mounts a transcriptomic response at lower dose rates than Norway

spruce signifies that it is more poised towards rapid response to inflicted radiation stress.

The energy deposited at a specific gamma radiation dose rate in the two plant species should

be similar, with ionisation and excitation of electrons occurring uniformly throughout the

cells. Thus, induced damage should also be randomly distributed in biomolecules such as

DNA and relative to the DNA content (genome size) (Einset and Collins, 2018). Consistent

with a previous report (Blagojevicet al., 2019a) this study showed less DNA damage in *A.*

thaliana than in Norway spruce both at the end of the irradiation and post-irradiation period

(Fig. 4). We therefore hypothesised that *A. thaliana* has more effective DNA repair and/or

protection systems. Yet, the DNA damage at 77 dpi in both species (Fig. 4), albeit at

considerably lower level than at the end of the 48-h irradiation, demonstrates that some DNA

damage or genomic instability is not critical to survival or development. Although genomic

instability may result into detrimental growth effects (Mothersill and Seymour, 1998), no

clear negative effects on the vegetative growth post-irradiation in *A. thaliana* even after

exposure to 400-540 mGy h^{-1} in contrast to the severe growth inhibition and mortality

observed in Norway spruce and Scots pine at dose rates $\geq$ 40 100 mGy h^{-1}, (Blagojevic et al.,

2019a) confirm better tolerance in *A. thaliana* compared to conifers. In our current study, no

mortality after 48 h of gamma irradiation in either species again points to a certain tolerance

to damage in both species.

The comparative RNA-seq analysis revealed that both *A. thaliana* and Norway spruce

initiated a plethora of DDR pathways to avoid harmful mutations and maintain genome

integrity. However, the response of the two species differed in two main aspects (Fig. S9;

Table S3). Firstly, the upregulation of DDR-related genes at lower doses dose rates in *A.*

thaliana than in Norway spruce suggests that the former is far more responsive towards lower

levels of DNA damage. Secondly, the prominent DNA repair pathways in *A. thaliana* were

mostly HR, and to some extent MMR, whereas in Norway spruce, in addition to HR, several

other pathways such as NHEJ, MMR and DNA transleison synthesis were activated. Hence,

it is tempting to speculate that Norway spruce needed to engage a wider range of DDR

because of the significantly higher levels of DNA damage compared to *A. thaliana*, possibly

because of less antioxidant protection and a larger genome.

One of the key contributor genes in gamma radiation-induced DDR in plants, *SOG1*, which

activates several DDR pathways and cell cycle checkpoints (Kim et al., 2019), was not

upregulated at any time points or dose rates in *A. thaliana* (Fig. 6, Table S3). This might

indicate that induction of *SOG1* expression is only modulated in higher total doses or acute

radiation that generate very high level of DNA damage (Bourbousse et al., 2018). However,

no change in *SOG1* expression after 48 h of gamma irradiation does not necessarily indicate

the lack of SOG1 activity as that could rather depend on the post-translational modification

by ATM- or ATR-kinases (Bourbousse et al., 2013). The involvement of *SOG1* in *A. thaliana*

is indeed supported by the activation of many primary target genes of SOG1 (i.e., *RAD51,*

BRCA1, PARP1 and *2, GMI1*) in response to the 48 h of gamma irradiation. By contrast,

significant upregulation of *SOG1* in Norway spruce after 24 h at ≥ 40 mGy h^{-1} and 48 h at

100 mGy h^{-1} (Fig. 7, Table S3) but no significant regulation of *ATM* (Table S3), might imply

that elevated *SOG1* expression was triggered in Norway spruce possibly due to higher levels

of accumulated DNA damage. Moreover, upregulation of *WEE1,* a direct target of SOG1 and

key regulator of CDK-inhibition and stress-induced cell cycle arrest (Kevei et al., 2011; De

Schutter et al., 2007) (Fig. 6, Table S4), coincided with *SOG1* expression in Norway spruce,

strongly suggesting SOG1 involvement in gamma-induced DDR and cell cycle arrest

(Pedroza-Garcia et al., 2022; Bourbousse et al., 2018).

Transcriptional activation of numerous B-type *CYC*- and *CDKB*-genes in *A. thaliana* after 48

h at 1-40 mGy h^{-1} (Fig. 6, Table S4) might indicate gamma radiation stress-induced cell cycle

arrest and subsequent DDR through HR by maintaining higher cyclin threshold (Endo et al.,

2006). This further supports the previous notion that enhanced CDKB1 activity is involved in

ionising radiation-induced cell cycle arrest leading to HR-repair in damaged mitotically

active cells (Weimer et al., 2016). On the contrary, there were fewer regulations and

downregulation only of B-type *CYC* or *CDK* genes in Norway spruce (Table S4), indicating

no significant involvement of such genes in such cell cycle arrest. The regulation of the key

contributors in cell cycle progression in both species, albeit to a larger degree in *A. thaliana*,
suggests that the cell cycle is one of the most affected biological pathways in the applied
gamma radiation. Most of the CYCs and CDKs regulated in *A. thaliana* in our study are also
involved in the regulation of the endocycle. However, stress-indued endoreduplication is less
common in gymnosperms than angiosperms (De Veylder et al., 2001) and fewer genes
related to the endocycle were upregulated in Norway spruce compared to *A. thaliana* (Table
S4, S5). Increased endoreduplication was previously shown in *A. thaliana* at 1 Gy min^{-1} (150
Gy totally) and *Lemna minor* at 1500 mGy h^{-1} for 7 days, but not at lower dose rates (Adachi
et al., 2009; Van Hoeck et al., 2015). Our preliminary study did not reveal any effect of 48 h
gamma radiation at 100 mGy h^{-1} on endoreduplication in *A. thaliana* and Norway spruce
(flow cytometry results not shown), suggesting that the applied dose was not sufficient to
induce the endocycle.

DDR is closely related to epigenetic regulation and chromatin structure must be decondensed
for the DDR proteins to carry out efficient repair (Choi et al., 2019; Kim et al., 2019).
Chromatin remodelers were previously shown to be upregulated in *A. thaliana* at very high
dose rates of ionising radiation (X-rays at 5.25 Gy h^{-1} with total doses of 10-100 Gy) (Sidler
et al., 2015). However, in our study upregulation of such genes in *A. thaliana* occurred at $\leq$
40 mGy h^{-1} only. By contrast, the upregulation of *H2A* variants, DNA methylation and
deacetylation-related genes in Norway spruce at $\geq$ 40 mGy h^{-1} only (Table S5, Fig. S5)
clearly indicates chromatin modification under high radiation stress in this species. The
enrichment of DNA methylation regulation is consistent with the previously reported DNA
hypermethylation in *P. sylvestris* trees exposed to high doses of ionising radiation in the
Chernobyl area, a mechanism suggested to stabilise the genome (Kovalchuk et al., 2003).
Despite the genome of Norway spruce being 150 times bigger than that of *A. thaliana*, it only
contains 2.5 times more predicted genes (Nystedt et al. 2013), and has large heterochromatin
regions which requires the activation of chromatin remodelers for decondensation for
efficient DDR (Fig. S9). Moreover, gamma radiation-induced damage to heterochromatin
regions does not initiate rapid activation of DDR pathways (Falk et al., 2008). It is thus
plausible that Norway spruce requires significantly more DNA damage for the activation of
DDR and chromosome remodelers.

The 48-h exposure did not affect the structural integrity of the SAMs (Fig. 2) in contrast to
the previous studies where 144 or 360-h gamma irradiation resulted in adverse effects in
SAMs of Norway spruce only (Blagojevic et al., 2019a). However, TEM image analysis

revealed that integrity of thylakoid and nuclear membranes was compromised in Norway

spruce, but not in *A. thaliana* (Fig. 3). Notably, in both species the mitochondria appeared to

be the most vulnerable organelle. Furthermore, the fact that disrupted mitochondria were

evident from 1 mGy h^{-1} in Norway spruce, while *A. thaliana* showed no damage below 100

mGy h^{-1}, indicates a major difference in their capacity to maintain cell homeostasis.

Former studies have demonstrated activation of ROS-scavenging enzymes such as SOD and

APX in gamma-irradiated plant species such as *A. thaliana* and *L. minor* (Vanhoudt et al.,

2014; Xie et al., 2019) and flavonoids in *Onobrychis viciifolia* and *Glycine max* (Katiyar et

al., 2022). Lack of activation of antioxidant-related genes in Norway spruce at ≤ 40 mGy h^{-1}

but comprehensive upregulation of such genes (*PERs, CHSs, GSTUs*, anthocyanin and

flavanol synthesis genes) at all dose rates in *A. thaliana* (Table S8), may at least partly

explain the damage to mitochondria, chloroplasts, and nuclear membranes at all dose rates in

Norway spruce in contrast to only slight mitochondrial damage at high dose rates in *A.*

thaliana (Fig. 3). Despite the observed organelle damage, the lack of significant upregulation

of two marker genes for retrograde signalling, *AOX1* and the transcription factor *WRKY40,* at

all dose rates in both species (Fig S8), may possibly suggest that the duration or levels of the

gamma irradiation were insufficient to activate retrograde signalling.

In the time-kinetic analyses of antioxidants (PER, CAT, flavonoids, SOD) in this study, only

SOD in Norway spruce exhibited increased activity in response to gamma irradiation (after

48 h at 40 and 100 mGy h^{-1}; Fig. 7). This suggests that the dynamics of the formation of these

antioxidants in response to radiation stress may vary among different species. Additionally,

increased biosynthesis of antioxidants may require a longer period of gamma irradiation. At

least the lack of increase in flavonoids in Norway spruce and *A. thaliana* in this study (Fig. 7)

and in Scots pine after 144 h of gamma irradiation (Blagojevic et al., 2019b) but elevated

flavonoid levels in Scots pine under chronic high radiation in the Chernobyl area (Nybakken

et al., 2023) are consistent with requirement of long-term irradiation. It is also possible that

assays of total antioxidant content/activity might not detect nuances of antioxidant

homeostasis provided by individual members of the antioxidant families, including the

antioxidant status in different cellular compartments.

Moreover, antioxidants such as PERs can modulate cell wall composition during stress (Díaz-

Tielas et al., 2012). The cell wall components cellulose and pectin were previously shown to

be damaged by very high doses of gamma radiation (1-2.5 kGy) (Kovács and Keresztes,

2002). Hence, in contrast to Norway spruce, the radical upregulation of cell wall component
genes even at lower gamma dose rates in *A. thaliana* (Table S9) might suggest efficient
replacement of damaged cell wall components and cell wall fortification in *A. thaliana* (Le
Gall et al., 2015).

Antioxidants such as *GST*s may also downregulate auxin-responsive genes (Sylvestre-Gonon
et al., 2019; Jiang et al., 2010). The upregulation of *GSTs* and growth retardation in Norway
spruce at higher dose rates as well as downregulation or no regulation of growth-promoting
hormone pathways are consistent with this notion (Fig.1, S1, Table S7). By contrast, the
gamma-irradiation-induced upregulation of auxin and cytokinin-related genes (biosynthesis,
metabolism, signalling, auxin transport) in *A. thaliana* complies with the maintenance of
meristem function to sustain normal growth. These results are in line with the previously
demonstrated activation of such genes in response to abiotic stress, indicating the importance
of growth-hormone homeostasis under stress (Verma et al., 2016). Similar to previous reports
on the induction of ABA biosynthesis and signalling genes triggered by ROS in gamma-
irradiated *A. thaliana* (Vishwakarma et al., 2017), upregulation of many such ABA-
responsive genes were observed in *A. thaliana* in our study, although the total doses applied
were much lower. Also, upregulation of the ethylene-responsive oxidative stress-pathway
(*ERFs*, *ACS5*) and SA-responsive genes (*MYBs*) in *A. thaliana*, but not Norway spruce (Table
S7), indicate less induction of hormone response to gamma-stress in the latter and is
consistent with the greater tolerance of *A. thaliana* to gamma radiation. Hence, in *A. thaliana*
better antioxidant homeostasis, cell wall biosynthesis/fortification and hormone signalling
might contribute to the high tolerance to the gamma irradiation. Conversely, less
transcriptional activation of the same protective pathways in Norway spruce possibly have
resulted in adverse effects on overall growth and higher susceptibility to radiation-induced
oxidative damage.

The RNA-seq analysis provided evidence that energy metabolism and photosynthesis in
Norway spruce were strongly affected by gamma irradiation in Norway spruce, but only
marginally in *A. thaliana*. In Norway spruce, the reduced expression of photosynthesis-
related genes coincided with the reduced growth and chloroplast impairment at higher dose
rates ($\geq$ 40 mGy h^{-1}) (Table S10). Concurringly, the strong upregulation of energy
metabolism-related genes (glycolysis, Krebs cycle, mitochondrial electron transport-related
genes) (Table S11) in this species may support the hypothesis that there is a high ATP
requirement for damage repair including maintenance of the large genome. Hence, it appears

that radiation induces energy deficiency, which in turn requires increased energy expenditure
on damage repair at the expense of growth and is reflected by the downregulation of growth-
promoting hormones in Norway spruce at higher dose rates. However, the upregulation of
energy metabolism genes might additionally indicate that the damage to mitochondria at high
dose rates could have led to inefficient energy production and thus a need for the activation of
respiratory genes to meet the energy demands.

5. Conclusions

This study demonstrates that compared to the radiosensitive Norway spruce, the radiotolerant
A. thaliana mobilises more comprehensive gamma radiation-induced defence, repair, and
homeostasis-maintaining stress responses, even at low dose rates given for 48 h. Despite
significant adverse effects in Norway spruce related to DNA and organelle damage and
growth at low dose rates, transcriptional activation of DDR was evident at 40-100 mGy h^{-1}
only. Furthermore, in Norway spruce, mobilisation of antioxidant genes, as well as
modulation of growth-regulatory and stress-related hormones at lower dose rates were less
efficient than in *A. thaliana*. Thus, the short-lived *A. thaliana* with short generation-time, and
small genome with less energy requirement for maintenance, shows better plasticity in
response to radiation stress compared to the long-lived Norway spruce with a lengthy
generation time and gigantic genome. These observations expand our understanding of
differential radiosensitivity in plants and thus, improves the knowledge base for risk
assessment and protection of ecosystems from ionising radiation exposure. In the context of
environmental impact, the findings confirm that under severe ecosystem exposure situations,
radiosensitive conifers require special attention with respect to remediating measures.

Acknowledgments

This study has been funded by the Research Council of Norway through its Centre of
Excellence (CoE) funding scheme (Project No. 223268/F50) and the Norwegian University
of Life Science (NMBU). Sincere thanks to Marit Siira and Svein Andre Kolltveit for skilful
assistance in plant growing and growth measurements and Tone Ingeborg Melby and Linda
Ripel for skilful work with qPCR analyses of gene expression.

Competing interests

None declared

Author contributions

Payel Bhattacharjee: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Validation, Visualization, Writing - original draft, Writing - review & editing; Dajana Blagojevic: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing – review and editing; YeonKyeong Lee: Methodology, Visualization, Formal analysis, Writing – review and editing; Gareth B Gillard: Data curation, Formal analysis, Visualization, Writing - review & editing; Lars Grønvold, Torgeir R Hvidsten, Simen R Sandve: Data curation, Formal analysis, Writing – review and editing; Ole C Lind: Conceptualization , Writing – review and editing; Brit Salbu : Resources; Funding acquisition, Writing – review and editing; Dag A Brede: Investigation, Methodology, Writing - review & editing, Jorunn E Olsen: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Validation, Visualization, Writing - original draft, Writing - review & editing.

Data availability

The raw RNA sequence data are available in ArrayExpress under the accession number E-MTAB-8081 for Norway spruce and E-MTAB-11120 for *A. thaliana*. Other data that support the findings of this study are available within the article, in the Supporting Information of this article and in online repositories. The names of the online repository and the link of the identifiers will be available in the Norwegian University of Life Sciences archive once the submission is approved.

References

Berardini, T. Z., L. Reiser, D. Li, Y. Mezheritsky, R. Muller, E. Strait, and E. Huala. 2015. "The Arabidopsis information resource: Making and mining the "gold standard" annotated reference plant genome." *Genesis* 53 (8): 474-85. https://doi.org/10.1002/dvg.22877. https://www.ncbi.nlm.nih.gov/pubmed/26201819.

Blagojevic, D., Y. Lee, D. A. Brede, O. C. Lind, I. Yakovlev, K. A. Solhaug, C. G. Fossdal, B. Salbu, and J. E. Olsen. 2019a. "Comparative sensitivity to gamma radiation at the organismal, cell and DNA level in young plants of Norway spruce, Scots pine and Arabidopsis thaliana." *Planta*. https://doi.org/10.1007/s00425-019-03250-y. https://www.ncbi.nlm.nih.gov/pubmed/31372744.

Blagojevic, Dajana, YeonKyeong Lee, Li Xie, Dag A. Brede, Line Nybakken, Ole Christian Lind, Knut Erik Tollefsen, Brit Knut Salbu, Asbjørn Solhaug, and Jorunn E. Olsen. 2019b. No evidence of a protective or cumulative negative effect of UV-B on growth inhibition induced by gamma

764  radiation in Scots pine (Pinus sylvestris) seedlings. *Photochemical & Photobiological*
765  *Sciences*.
766 Bourbousse, C., N. Vegesna, and J. A. Law. 2018. "SOG1 activator and MYB3R repressors regulate a
767  complex DNA damage network in." *Proc Natl Acad Sci U S A* 115 (52): E12453-E12462.
768  https://doi.org/10.1073/pnas.1810582115.
769  https://www.ncbi.nlm.nih.gov/pubmed/30541889.
770 Caplin, N., and N. Willey. 2018. "Ionizing Radiation, Higher Plants, and Radioprotection: From Acute
771  High Doses to Chronic Low Doses." *Front Plant Sci* 9: 847.
772  https://doi.org/10.3389/fpls.2018.00847.
773  https://www.ncbi.nlm.nih.gov/pubmed/29997637.
774 Choi, H. I., S. M. Han, Y. D. Jo, M. J. Hong, S. H. Kim, and J. B. Kim. 2021. "Effects of Acute and Chronic
775  Gamma Irradiation on the Cell Biology and Physiology of Rice Plants." *Plants (Basel)* 10 (3).
776  https://doi.org/10.3390/plants10030439.
777  https://www.ncbi.nlm.nih.gov/pubmed/33669039.
778 Choi, S. H., T. H. Ryu, J. I. Kim, S. Lee, S. S. Lee, and J. H. Kim. 2019. "Mutation in DDM1 inhibits the
779  homology directed repair of double strand breaks." *PLoS One* 14 (2): e0211878.
780  https://doi.org/10.1371/journal.pone.0211878.
781  https://www.ncbi.nlm.nih.gov/pubmed/30742642.
782 De Schutter, K., J. Joubès, T. Cools, A. Verkest, F. Corellou, E. Babiychuk, E. Van Der Schueren, T.
783  Beeckman, S. Kushnir, D. Inzé, and L. De Veylder. 2007. "Arabidopsis WEE1 kinase controls
784  cell cycle arrest in response to activation of the DNA integrity checkpoint." *Plant Cell* 19 (1):
785  211-25. https://doi.org/10.1105/tpc.106.045047.
786  https://www.ncbi.nlm.nih.gov/pubmed/17209125.
787 De Veylder, L., G. T. Beemster, T. Beeckman, and D. Inzé. 2001. "CKS1At overexpression in
788  Arabidopsis thaliana inhibits growth by reducing meristem size and inhibiting cell-cycle
789  progression." *Plant J* 25 (6): 617-26. https://doi.org/10.1046/j.1365-313x.2001.00996.x.
790  https://www.ncbi.nlm.nih.gov/pubmed/11319029.
791 Dumanović, J., E. Nepovimova, M. Natić, K. Kuča, and V. Jaćević. 2020. "The Significance of Reactive
792  Oxygen Species and Antioxidant Defense System in Plants: A Concise Overview." *Front Plant*
793  *Sci* 11: 552969. https://doi.org/10.3389/fpls.2020.552969.
794  https://www.ncbi.nlm.nih.gov/pubmed/33488637.
795 Díaz-Tielas, C., E. Graña, M. J. Reigosa, and A. M. Sánchez-Moreiras. 2012. "The role of peroxidases
796  on the mode of action of chalcone in Arabidopsis roots." *Plant Signal Behav* 7 (10): 1274-6.
797  https://doi.org/10.4161/psb.21594. https://www.ncbi.nlm.nih.gov/pubmed/22895105.
798 Einset, J., and A. R. Collins. 2018. "Genome size and sensitivity to DNA damage by X-rays-plant
799  comets tell the story." *Mutagenesis* 33 (1): 49-51. https://doi.org/10.1093/mutage/gex029.
800  https://www.ncbi.nlm.nih.gov/pubmed/29045752.
801 Endo, M., Y. Ishikawa, K. Osakabe, S. Nakayama, H. Kaya, T. Araki, K. Shibahara, K. Abe, H. Ichikawa,
802  L. Valentine, B. Hohn, and S. Toki. 2006. "Increased frequency of homologous recombination
803  and T-DNA integration in Arabidopsis CAF-1 mutants." *EMBO J* 25 (23): 5579-90.
804  https://doi.org/10.1038/sj.emboj.7601434.
805  https://www.ncbi.nlm.nih.gov/pubmed/17110925.
806 Falk, M., E. Lukásová, and S. Kozubek. 2008. "Chromatin structure influences the sensitivity of DNA
807  to gamma-radiation." *Biochim Biophys Acta* 1783 (12): 2398-414.
808  https://doi.org/10.1016/j.bbamcr.2008.07.010.
809  https://www.ncbi.nlm.nih.gov/pubmed/18706456.
810 Garnier-Laplace, J., S. Geras'kin, C. Della-Vedova, K. Beaugelin-Seiller, T. G. Hinton, A. Real, and A.
811  Oudalova. 2013. "Are radiosensitivity data derived from natural field conditions consistent
812  with data from controlled exposures? A case study of Chernobyl wildlife chronically exposed
813  to low dose rates." *J Environ Radioact* 121: 12-21.
814  https://doi.org/10.1016/j.jenvrad.2012.01.013.
815  https://www.ncbi.nlm.nih.gov/pubmed/22336569.

Gichner, T. 2003. "DNA damage induced by indirect and direct acting mutagens in catalase-deficient transgenic tobacco. Cellular and acellular Comet assays." *Mutat Res* 535 (2): 187-93. https://doi.org/10.1016/s1383-5718(02)00320-0. https://www.ncbi.nlm.nih.gov/pubmed/12581537.

Hansen, E. L., O. C. Lind, D. H. Oughton, and B. Salbu. 2019. "A framework for exposure characterization and gamma dosimetry at the NMBU FIGARO irradiation facility." *Int J Radiat Biol* 95 (1): 82-89. https://doi.org/10.1080/09553002.2018.1539878. https://www.ncbi.nlm.nih.gov/pubmed/30407094.

International Atomic energy Agency, the annual report for 1992.

Jiang, H. W., M. J. Liu, I. C. Chen, C. H. Huang, L. Y. Chao, and H. L. Hsieh. 2010. "A glutathione S-transferase regulated by light and hormones participates in the modulation of Arabidopsis seedling development." *Plant Physiol* 154 (4): 1646-58. https://doi.org/10.1104/pp.110.159152. https://www.ncbi.nlm.nih.gov/pubmed/20935176.

Kevei, Z., M. Baloban, O. Da Ines, H. Tiricz, A. Kroll, K. Regulski, P. Mergaert, and E. Kondorosi. 2011. "Conserved CDC20 cell cycle functions are carried out by two of the five isoforms in Arabidopsis thaliana." *PLoS One* 6 (6): e20618. https://doi.org/10.1371/journal.pone.0020618. https://www.ncbi.nlm.nih.gov/pubmed/21687678.

Kim, D. S., J. B. Kim, E. J. Goh, W. J. Kim, S. H. Kim, Y. W. Seo, C. S. Jang, and S. Y. Kang. 2011. "Antioxidant response of Arabidopsis plants to gamma irradiation: Genome-wide expression profiling of the ROS scavenging and signal transduction pathways." *J Plant Physiol* 168 (16): 1960-71. https://doi.org/10.1016/j.jplph.2011.05.008. https://www.ncbi.nlm.nih.gov/pubmed/21665324.

Kim, J. H. 2019. "Chromatin Remodeling and Epigenetic Regulation in Plant DNA Damage Repair." *Int J Mol Sci* 20 (17). https://doi.org/10.3390/ijms20174093. https://www.ncbi.nlm.nih.gov/pubmed/31443358.

Kim, J. H., K. Hwangbo, E. Lee, S. K. Dubey, M. S. Chung, B. Y. Chung, and S. Lee. 2021. "Application of Gamma Ray-Responsive Genes for Transcriptome-Based Phytodosimetry in Rice." *Plants (Basel)* 10 (5). https://doi.org/10.3390/plants10050968. https://www.ncbi.nlm.nih.gov/pubmed/34067996.

Kim, J. H., T. H. Ryu, S. S. Lee, S. Lee, and B. Y. Chung. 2019. "Ionizing radiation manifesting DNA damage response in plants: An overview of DNA damage signaling and repair mechanisms in plants." *Plant Sci* 278: 44-53. https://doi.org/10.1016/j.plantsci.2018.10.013. https://www.ncbi.nlm.nih.gov/pubmed/30471728.

Kovalchuk, O., P. Burke, A. Arkhipov, N. Kuchma, S. J. James, I. Kovalchuk, and I. Pogribny. 2003. "Genome hypermethylation in Pinus silvestris of Chernobyl--a mechanism for radiation adaptation?" *Mutat Res* 529 (1-2): 13-20. https://doi.org/10.1016/s0027-5107(03)00103-9. https://www.ncbi.nlm.nih.gov/pubmed/12943916.

Kovács, E., and A. Keresztes. 2002. "Effect of gamma and UV-B/C radiation on plant cells." *Micron* 33 (2): 199-210. https://doi.org/10.1016/s0968-4328(01)00012-9. https://www.ncbi.nlm.nih.gov/pubmed/11567888.

Le Gall, H., F. Philippe, J. M. Domon, F. Gillet, J. Pelloux, and C. Rayon. 2015. "Cell Wall Metabolism in Response to Abiotic Stress." *Plants (Basel)* 4 (1): 112-66. https://doi.org/10.3390/plants4010112. https://www.ncbi.nlm.nih.gov/pubmed/27135320.

Lee, Y., C. Karunakaran, R. Lahlali, X. Liu, K. K. Tanino, and J. E. Olsen. 2017. "Photoperiodic Regulation of Growth-Dormancy Cycling through Induction of Multiple Bud-Shoot Barriers Preventing Water Transport into the Winter Buds of Norway Spruce." *Front Plant Sci* 8: 2109. https://doi.org/10.3389/fpls.2017.02109. https://www.ncbi.nlm.nih.gov/pubmed/29321789.

Lind, O. C., D. Helen Oughton, and B. Salbu. 2019. "The NMBU FIGARO low dose irradiation facility." *Int J Radiat Biol* 95 (1): 76-81. https://doi.org/10.1080/09553002.2018.1516906. https://www.ncbi.nlm.nih.gov/pubmed/30265167.

Livak, K. J., and T. D. Schmittgen. 2001. "Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method." *Methods* 25 (4): 402-8.
https://doi.org/10.1006/meth.2001.1262. http://www.ncbi.nlm.nih.gov/pubmed/11846609.
Mothersill, C., and C. B. Seymour. 1998. "Mechanisms and implications of genomic instability and
other delayed effects of ionizing radiation exposure." *Mutagenesis* 13 (5): 421-6.
https://doi.org/10.1093/mutage/13.5.421.
https://www.ncbi.nlm.nih.gov/pubmed/9800186.
Nisa, M. U., Y. Huang, M. Benhamed, and C. Raynaud. 2019. "The Plant DNA Damage Response:
Signaling Pathways Leading to Growth Inhibition and Putative Role in Response to Stress
Conditions." *Front Plant Sci* 10: 653. https://doi.org/10.3389/fpls.2019.00653.
https://www.ncbi.nlm.nih.gov/pubmed/31164899.
Nystedt, B., N. R. Street, A. Wetterbom, A. Zuccolo, Y. C. Lin, D. G. Scofield, F. Vezzi, N. Delhomme, S.
Giacomello, A. Alexeyenko, R. Vicedomini, K. Sahlin, E. Sherwood, M. Elfstrand, L. Gramzow,
K. Holmberg, J. Hällman, O. Keech, L. Klasson, M. Koriabine, M. Kucukoglu, M. Käller, J.
Luthman, F. Lysholm, T. Niittylä, A. Olson, N. Rilakovic, C. Ritland, J. A. Rosselló, J. Sena, T.
Svensson, C. Talavera-López, G. Theißen, H. Tuominen, K. Vanneste, Z. Q. Wu, B. Zhang, P.
Zerbe, L. Arvestad, R. Bhalerao, J. Bohlmann, J. Bousquet, R. Garcia Gil, T. R. Hvidsten, P. de
Jong, J. MacKay, M. Morgante, K. Ritland, B. Sundberg, S. L. Thompson, Y. Van de Peer, B.
Andersson, O. Nilsson, P. K. Ingvarsson, J. Lundeberg, and S. Jansson. 2013. "The Norway
spruce genome sequence and conifer genome evolution." *Nature* 497 (7451): 579-84.
https://doi.org/10.1038/nature12211. https://www.ncbi.nlm.nih.gov/pubmed/23698360.
Patro, R., G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford. 2017. "Salmon provides fast and bias-
aware quantification of transcript expression." *Nat Methods* 14 (4): 417-419.
https://doi.org/10.1038/nmeth.4197. https://www.ncbi.nlm.nih.gov/pubmed/28263959.
Pedroza-Garcia, J. A., Y. Xiang, and L. De Veylder. 2022. "Cell cycle checkpoint control in response to
DNA damage by environmental stresses." *Plant J* 109 (3): 490-507.
https://doi.org/10.1111/tpj.15567. https://www.ncbi.nlm.nih.gov/pubmed/34741364.
Qi, F., and F. Zhang. 2019. "Cell Cycle Regulation in the Plant Response to Stress." *Front Plant Sci* 10:
1765. https://doi.org/10.3389/fpls.2019.01765.
https://www.ncbi.nlm.nih.gov/pubmed/32082337.
Sheppard, B.D. Amiro and S.C. 1994. Effects of ionizing radiation on the boreal forest: Canada's FIG
experiment, with implications for radionuclides. The Science of the Total Environment.
Sidler, C., D. Li, O. Kovalchuk, and I. Kovalchuk. 2015. "Development-dependent expression of DNA
repair genes and epigenetic regulators in Arabidopsis plants exposed to ionizing radiation."
Radiat Res 183 (2): 219-32. https://doi.org/10.1667/RR13840.1.
https://www.ncbi.nlm.nih.gov/pubmed/25594540.
Sylvestre-Gonon, E., S. R. Law, M. Schwartz, K. Robe, O. Keech, C. Didierjean, C. Dubos, N. Rouhier,
and A. Hecker. 2019. "Functional, Structural and Biochemical Features of Plant Serinyl-
Glutathione Transferases." *Front Plant Sci* 10: 608. https://doi.org/10.3389/fpls.2019.00608.
https://www.ncbi.nlm.nih.gov/pubmed/31191562.
UNSCEAR. SOURCES AND EFFECTS

OF IONIZING RADIATION. United Nations Scientific Committee on the Effects of Atomic Radiation
UNSCEAR 1996 Report to the General Assembly, with Scientific Annexes.
van de Walle, J., N. Horemans, E. Saenen, M. Van Hees, J. Wannijn, R. Nauts, A. van Gompel, J.
Vangronsveld, H. Vandenhove, and A. Cuypers. 2016. "Arabidopsis plants exposed to gamma
radiation in two successive generations show a different oxidative stress response." *J
Environ Radioact* 165: 270-279. https://doi.org/10.1016/j.jenvrad.2016.10.014.
https://www.ncbi.nlm.nih.gov/pubmed/27814501.
Vanhoudt, N., N. Horemans, J. Wannijn, R. Nauts, M. Van Hees, and H. Vandenhove. 2014. "Primary
stress responses in Arabidopsis thaliana exposed to gamma radiation." *J Environ Radioact*

129: 1-6. https://doi.org/10.1016/j.jenvrad.2013.11.011.
https://www.ncbi.nlm.nih.gov/pubmed/24333636.

Verma, V., P. Ravindran, and P. P. Kumar. 2016. "Plant hormone-mediated regulation of stress responses." *BMC Plant Biol* 16: 86. https://doi.org/10.1186/s12870-016-0771-y. https://www.ncbi.nlm.nih.gov/pubmed/27079791.

Vishwakarma, K., N. Upadhyay, N. Kumar, G. Yadav, J. Singh, R. K. Mishra, V. Kumar, R. Verma, R. G. Upadhyay, M. Pandey, and S. Sharma. 2017. "Abscisic Acid Signaling and Abiotic Stress Tolerance in Plants: A Review on Current Knowledge and Future Prospects." *Front Plant Sci* 8: 161. https://doi.org/10.3389/fpls.2017.00161. https://www.ncbi.nlm.nih.gov/pubmed/28265276.

Weimer, A. K., S. Biedermann, H. Harashima, F. Roodbarkelari, N. Takahashi, J. Foreman, Y. Guan, G. Pochon, M. Heese, D. Van Damme, K. Sugimoto, C. Koncz, P. Doerner, M. Umeda, and A. Schnittger. 2016. "The plant-specific CDKB1-CYCB1 complex mediates homologous recombination repair in Arabidopsis." *EMBO J* 35 (19): 2068-2086. https://doi.org/10.15252/embj.201593083. https://www.ncbi.nlm.nih.gov/pubmed/27497297.

Xie, L., K. A. Solhaug, Y. Song, D. A. Brede, O. C. Lind, B. Salbu, and K. E. Tollefsen. 2019. "Modes of action and adverse effects of gamma radiation in an aquatic macrophyte Lemna minor." *Sci Total Environ* 680: 23-34. https://doi.org/10.1016/j.scitotenv.2019.05.016. https://www.ncbi.nlm.nih.gov/pubmed/31085442.

Yoshiyama, K. O., J. Kobayashi, N. Ogita, M. Ueda, S. Kimura, H. Maki, and M. Umeda. 2013. "ATM-mediated phosphorylation of SOG1 is essential for the DNA damage response in Arabidopsis." *EMBO Rep* 14 (9): 817-22. https://doi.org/10.1038/embor.2013.112. https://www.ncbi.nlm.nih.gov/pubmed/23907539.

Yu., Volkova Polina, Bondarenko Ekaterina V., and Kazakova Elizaveta A. 2022. Radiation hormesis in plants. Elesevier.

Zimin, A. V., K. A. Stevens, M. W. Crepeau, D. Puiu, J. L. Wegrzyn, J. A. Yorke, C. H. Langley, D. B. Neale, and S. L. Salzberg. 2017. "An improved assembly of the loblolly pine mega-genome using long-read single-molecule sequencing." *Gigascience* 6 (1): 1-4. https://doi.org/10.1093/gigascience/giw016. https://www.ncbi.nlm.nih.gov/pubmed/28369353.

Table 1. The gamma radiation dose rates and total doses applied in experiments with 12, 24 and 48 h exposure of seedlings of *Arabidopsis thaliana* and Norway spruce from day 7 after sowing, using a [60]Co source.

Dose rate (mGy h^{-1})	Dose rate interval (mGy h^{-1})		Total dose (Gy) after 48 h exposure	Total dose interval (Gy) after 48 h exposure		Total dose (Gy) after 24 h exposure	Total dose interval (Gy) after 24 h exposure		Total dose (Gy) after 12 h exposure	Total dose interval (Gy) after 12 h exposure	
	Min	Max		Min	Max		Min	Max		Min	Max
290	255	325	13.9	12.24	15.60	7.0	6.12	7.80	3.5	3.06	3.90
100	88	112	4.8	4.22	5.38	2.4	2.11	2.69	1.2	1.06	1.34
40	35	45	1.9	1.68	2.16	1.0	0.84	1.08	0.48	0.42	0.54
10	8,8	11,2	0.48	0.42	0.54	0.24	0.21	0.27	0.12	0.11	0.13
1	0,88	1,12	0.048	0.042	0.054	0.024	0.021	0.027	0.012	0.011	0.013
0	-	-	0	-	-		-	-		-	-

Fig. 1   *Bhattacharjee et al. 2025*

A. thaliana   *P. abies*   Dose rates
(mGy h^{-1})

Fig. 2   *Bhattacharjee et al. 2025*

Dose rate (mGy h⁻¹)

Fig. 3   *Bhattacharjee et al. 2025*

Fig. 4   *Bhattacharjee et al. 2025*

Fig. 5   *Bhattacharjee et al. 2025*

Fig. 6 Bhattacharjee et al., 2025

Fig. 7 *Bhattacharjee et al., 2025*

Fig. 1. Effect of 48 h of gamma irradiation from day 7 after sowing on growth in *A. thaliana* and Norway spruce (*P. abies*). (a) Hypocotyl and (b) cotyledon length in *A. thaliana* (n = 29-40) after the 48-h irradiation and (c) morphology 77 days post-irradiation (dpi). (d) Shoot and (e) root length of Norway spruce (n = 28-35) after the 48-h irradiation and (f) shoot diameter and (g) number of needles in time courses (n = 11-22) and (h) morphology 77 dpi. Scale bars: 3 cm. Means ± standard errors are shown. Different letters within each figure denotes significant difference (p ≤ 0.05) based on analysis of variance followed by Tukey`s test.

Fig. 2. Light micrographs of shoot apical meristems after 48 h of gamma irradiation from day 7 after sowing in *A. thaliana* and Norway spruce (*P. abies*). Five plants were investigated per treatment for each species. Scale bars: 25 µm.

Fig. 3. Transmission electron micrographs showing cellular organelles after 48 h of gamma irradiation from day 7 after sowing in (a) – (c) A. *thaliana* and (d) - (e) Norway spruce (*P. abies*). nu: nucleus, no: nucleoli, mt: mitochondrion, ch: chloroplast, sb: starch body. Arrows indicate nuclear membrane. Arrowheads indicate disrupted mitochondria. Five plants were investigated per treatment for each species. Scale bars: Scale nu: 1 µm, ch: 500 nm, mt: 500 nm.

Fig. 4. Effect of 48 h of gamma irradiation from day 7 after sowing on DNA damage detected by the COMET assay in (a) *A. thaliana* and (b) Norway spruce after the 48-h irradiation and 77 days post-irradiation (dpi). The line in each box represents the median values of 3 samples with 180-200 nuclei scored in each. Lower and upper box boundaries = 25 and 75 percentiles, error bars = 10 and 90 percentiles with data points outside these shown as dots. Different letters within each figure indicate significant difference (p ≤ 0.05) based on analysis of variance followed by Tukey's test.

Fig. 5. Number of differentially expressed genes (DEGs) and their $\log_2$ fold change after 48 h of gamma irradiation from day 7 after sowing relative to unexposed controls of (a)-(b). *A. thaliana* and (c)-(d) Norway spruce, with false discovery rate (FDR) < 0.05 (n = 4). Overlap of upregulated (red numbers) and downregulated (blue numbers) genes at different dose rates in (b)

A. thaliana and (d) Norway spruce. For each species, four repeated samples were analysed in duplicate per treatment.

Fig. 6. Relative transcript levels of DNA repair and cell cycle genes as well as the antioxidant-related *GLUTATION-S-TRANSFERASE* (*GSTU17*) gene after 12, 24 and 48 h of gamma irradiation from day 7 after sowing in shoots of *A. thaliana (At)* and Norway spruce *(Pa)*. Transcript levels were normalized against *ACTIN*, *ELONGATION FACTOR 1 α* and *UBIQUITIN 10* and shown relative to the unexposed controls at 12 h. The results are mean ± SE (n = 3-4 with 3-4 technical replicates). Different letters within each gene indicate significant difference (p ≤ 0.05) based on analysis of variance followed by Tukey's post-hoc test.

Fig. 7. Total antioxidant concentration or activity after 12, 24 and 48 h of gamma irradiation from day 7 after sowing in shoots of *A. thaliana* and Norway spruce (*Picea abies*). The antioxidant results were normalised based on absorbance of individual standards. The results are mean ± SE (n = 3 or 4 (flavonoids) with 2-3 technical replicates). Different letters within each concentration/activity indicate significant difference (p ≤ 0.05) based on analysis of variance followed by Tukey's post-hoc test.

Organisms

A. thaliana

γ- radiation

Norway spruce (Picea abies)

Effects / Outcomes

Mitochondria

Protection

Antioxidants

ROS

Signalling (MAPK)

DNA Damage 🔺

DDR 🔺

Regulation of transcription 🔺

• No effects on growth
• No/minor effects organelles
• Efficient DDR

ROS

Signalling (MAPK)

Antioxidants

Mitochondria damage

Less protection?

Respiration

Energy biosynthesis 🔺

DNA Damage 🔺

DDR 🔺

Regulation of transcription 🔺

Chloroplast damage

Photosynthesis 🔻

• Negative effects on growth
• Damaged organelles
• Partially efficient DDR

Gamma dose rate (mGy h⁻¹)

🔴 up regulated 🔵 down regulated

Arabidopsis DEGs: 1, 10, 40, 100, 290

Significance (–log₁₀ p-value adj.)

Direction (log₂ fold change)

Norway spruce DEGs

Probable causes

• Efficient AOX protection and DDR from low doses
• Small genome related flexibility

• AOX protection and DDR only in the higher doses
• Large genome related constraints

Significance

Prediction and preparation for adverse environmental effects in nuclear accidents/use of weapons

🔺 =Up and 🔻 = down-regulation in RNA-seq
Probable effects ┈┈➤   Direct effects ──➤

AOX= Antioxidants
DDR= DNA damage repair

Created with Biorender.com