High radiosensitivity in Norway spruce (*Picea abies*) is rendered by less comprehensive

mobilisation of protection, repair and stress responses compared to the radiotolerant

Arabidopsis thaliana

Payel Bhattacharjee[1,2], Dajana Blagojevic[1,2+], YeonKyeong Lee[1,2,3+], Gareth B Gillard[4], Lars

Grønvold[4], Torgeir Rhoden Hvidsten[5], Simen Rød Sandve[4], Ole Christian Lind[2,6], Brit Salbu

[2,6], Dag Anders Brede[2,6], Jorunn E. Olsen[1,2]

[1]Department of Plant Sciences, Faculty of Biosciences, Norwegian University of Life

Sciences, Ås, Norway

[2]Centre for Environmental Radioactivity (CERAD CoE), Norwegian University of Life

Sciences, Ås, Norway

[3]Department of Plant Biotechnology, Korea University Graduate School, 145 Anam-ro,

Seongbuk-ku, Seoul, Republic of Korea

[4]Department of Animal and Aquacultural Sciences, Faculty of Biosciences, Norwegian

University of Life Sciences, Ås, Norway

[5]Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life

Sciences, Ås, Norway

[6]Department of Environmental Sciences, Faculty of Environmental Sciences and Natural

Resource Management, Norwegian University of Life Sciences, Ås, Norway

[+] These authors contributed equally.

Authors for correspondence: Payel Bhattacharjee, Jorunn E. Olsen

Email: payel.bhattachrjee@nmbu.no

E-mail address: jorunn.olsen@nmbu.no

Word counts: Introduction: 939, Materials and Methods: 1728, Results: 2486, Discussion:

2329.

Number of figures: 7, number of tables: 1, number of supporting information files: 9 figures

and 11 tables.

ORCIDS:

Payel Bhattacharjee: orcid.org/0000-0003-1058-4233

Dajana Blagojevic

YeonKyeong Lee

Gareth B Gillard: orcid.org/0000-0001-9533-3227

Lars Grønvold: orcid.org/0000-0001-5404-4285

Torgeir Rhoden Hvidsten: orcid.org/0000-0001-6097-2539

Simen Rød Sandve: orcid.org/0000-0003-4989-5311

Ole Christian Lind: orcid.org/0000-0001-9074-8028

Brit Salbu

Dag Anders Brede: orcid.org/0000-0002-0906-3951

Jorunn E. Olsen: orcid.org/0000-0002-3380-3091

Summary

- Risk assessment and protection of plant communities in contaminated ecosystems require in-depth understanding of differential sensitivity to chronic ionising radiation in plants. However, the contributing molecular factors are poorly understood.

- We compared early molecular events associated with protection, repair, and stress responses, by analysing growth, organelle and DNA damage, and the dynamics of transcriptome and DNA damage repair (DDR) gene expression in gamma-irradiated seedlings of the radiosensitive conifer Norway spruce (*Picea abies*) and the radiotolerant *Arabidopsis thaliana*.

- Norway spruce showed reduced growth at 290 mGy h^{-1} and severe organelle damage, especially in mitochondria, at ≥ 1 mGy h^{-1} whereas *A. thaliana* showed unaffected development, minor organelle damage at ≥ 100 mGy h^{-1} and significantly less DNA damage, after 48-h of gamma exposure. Comparative transcriptomics revealed that *A. thaliana* showed massive activation of genes related to DDR, antioxidants, and other stress responses at ≥ 1 mGy h^{-1} while Norway spruce mobilized transcription of such pathways only at ≥ 40 mGy h^{-1}.

•   The transcriptional activation of repair and protection responses at higher gamma
dose-rates only and its absence in lower dose-rates, corelates with high
radiosensitivity of Norway spruce, compared to the massive transcriptional activation
from low dose-rates in the radiotolerant *A. thaliana*.

Key words: antioxidant protection, *Arabidopsis thaliana* (thale cress), comparative
transcriptomics, differential radiosensitivity, DNA damage repair, organelle damage, *Picea*
abies (Norway spruce), radiosensitivity in plants.

Introduction

Stress tolerance capacity in plants in general is highly variable and a defining factor for
species niche and habitat preference. Ionising radiation is a stressor that can induce harmful
effects to organisms due to damage to biological macromolecules such as DNA, proteins and
lipids (Caplin and Wiley 2018). Field experiments and surveys of radionuclide-contaminated
plant communities and ecosystems have shown that species that are sensitive to chronic
ionising radiation may succumb and be replaced by more tolerant species (Amiro and
Sheppard 1994; Caplin and Willey 2018). Laboratory studies have confirmed that
radiosensitivity of different plant species ranges over orders of magnitude (Caplin and Wiley
2018; Blagojevic et al., 2019). However, the underlying mechanisms conferring tolerance to
ionising radiation are not well understood. Such knowledge is needed to improve our ability
to predict environmental consequences of nuclear accidents and release of radionuclides
leading to exposure to high levels of ionising radiation.

The average natural background level of ionising radiation is estimated to 0.3 µGy h^{-1}, while
15 µGy h^{-1} has been measured in areas with naturally elevated radionuclide levels (Caplin
and Willey 2018). The nuclear power plant accident in Chernobyl in April 1986 as well as
other accidents and experiments have shown that acute high doses (10-1000 Gy) can be lethal
for plants and that radiosensitivity differs between species (UNSCEAR 1996). In response to

sub-lethal chronic radiation, adverse effects such as growth inhibition, reduced apical dominance (increased branching) and leaf chlorosis have been observed, particularly in conifers (Caplin and Wiley 2018; Kashparova et al., 2020). A dose rate of 10 mGy d^{-1} (417 µGy h^{-1}) has been suggested as a threshold value in the sense that lower dose rates have "no detrimental effects for populations of terrestrial plants in the field" (International Atomic Energy Agency [IAEA] 1992; (Caplin and Willey 2018; UNSCEAR 1996). However, this does not appear to hold true for boreal conifer forests. In a long-term field irradiation experiment in Canada persistent effects on forests including conifer trees were observed at chronic dose rates ≥100 µGy h^{-1} (Amiro and Sheppard 1994). In addition to the differential radiosensitivity between species (Blagojevic et al. 2019; Vanhoudt et al. 2014; Choi et al. 2021; IAEA1992), effects of ionising radiation depend on dose rate, duration (acute or chronic), total dose, life stages and the co-occurrence of other contaminants and environmental conditions (Garnier-Laplace et al. 2013).

The sensitivity of plant species to acute irradiation has been suggested to be attributed to the efficiency of DNA repair systems and genome organisation and size (De Micco et al., 2011). *A. thaliana* has a small genome of 135 Mb, whereas conifers like Norway spruce and Scots pine have giant genomes of 20 and 23 Gb (Nystedt et al. 2013; Zimin et al. 2017). On the other hand, the sensitivity to chronic irradiation appears to be regulated by delicate mechanisms on different levels of biological organisation (Caplin and Willey 2018; Kim et al. 2019). However, factors that define sensitivity to chronic exposure are poorly understood and apparently have functional rather than structural origin, depending on the molecular architecture of stress-responsive pathways of different plant species.

Ionising radiation can cause damage both directly and indirectly via the formation of free radicals and subsequent reactive oxygen species (ROS) (Caplin and Willey 2018). Plants possess antioxidant systems involving enzymes such as catalases (CAT), superoxide dismutase (SOD) and peroxidases (PER) as well as metabolites such as ascorbate, phenolic compounds and glutathione (GSH) that can neutralise ROS and thereby prevent oxidative damage (Dumanović et al. 2020). Studies of specific plant species including *A. thaliana*, have revealed induction of antioxidant-related genes, including *SOD* and *ASCORBATE PEROXIDASE* (*APX*) following ionising radiation exposure but there is limited information about antioxidant homeostasis in conifers (van de Walle et al. 2016; Kim et al. 2011).

DNA strand breaks induced by ROS or directly by the ionising radiation trigger the activation

of DNA damage repair (DDR) pathways (Kim, et al. 2019). A range of homologous-

recombination (HR) repair-mediated DDR genes known to be activated by the plant-specific

transcription factor *SUPRESSOR OF GAMMA RESPONSE 1 (SOG1)* were significantly

upregulated by gamma irradiation at 100 Gy in *A. thaliana* (Bourbousse et al. 2018). The

initiation of the DDR is closely interlinked with cell cycle stalling through the activation of

cell cycle checkpoints by regulating cyclin (CYC)-cyclin-dependent kinase (CDK)-

complexes (Qi and Zhang 2019), further leading to DNA damage-induced endoreduplication

(Nisa et al. 2019). In addition, transcriptional recruitment of DDR proteins in response to

gamma radiation requires chromatin remodelers involved in DNA methylation and histone

2A (H2A) modification (Choi et al. 2019; Kim 2019; Kim et al. 2021). However, there is a

lack of knowledge related to molecular events associated with DDR and other relevant

pathways in conifers exposed to ionising radiation.

Although transcriptional regulation in response to high doses of ionising radiation in *A.*

thaliana and a few other species has been explored (Choi, et al. 2021; Vanhoudt, et al. 2014),

information on the molecular mechanisms leading to the differential radiosensitivity among

plant species is yet to be clarified. Knowledge on the factors determining radiosensitivity of

plant species is crucial for risk assessment and protection of ecosystems from ionising

radiation hazard. Hence, by employing gamma irradiation under standardised laboratory

conditions, concomitant analyses of growth, organelle, and DNA damage, as well as the

dynamics of transcriptomics and DDR gene expression, this study aimed to provide an

overview of mechanisms involved in plant radiosensitivity. In this respect, early

physiological, cellular, and molecular effects were compared in the radiosensitive conifer

Norway spruce and the radio-tolerant *A. thaliana* to provide new insights regarding observed

effects in plant communities growing in areas contaminated with radionuclides.

Materials and methods

Plant materials and pre-growing conditions

Seeds of the Landsberg erecta (L*er*) accession of *A. thaliana thaliana* (L) Heyhn, and

Norway spruce (*Picea abies* L. H. Karst) from the provenance CØ1 from Halden, Norway

(59°N latitude, Skogfrøverket, Hamar, Norway) were surface-sterilised and sowed according

to the protocol in (Blagojevic, et al. 2019). The seeds were germinated for 6 days at 20 °C

under a 16 h photoperiod with a photosynthetic photon flux density (PPFD) of 30 μmol m^{-2} s^{-}

[1] at 400-700 nm (TL-D 58W/840 lamps, Philips, Eindhoven, The Netherlands), as measured

at the top of the Petri dishes with a Li-Cor Quantum/Radiometer/Photometer (LI-250, LI-

COR, USA).

Gamma irradiation of seedlings and growing conditions during the exposure

Six days old seedlings of *A. thaliana* and Norway spruce were exposed for 48 h using the

FIGARO gamma irradiation facility ([60]Co; 1173.2 and 1332.5 keV γ-rays) at the Norwegian

University of Life Sciences (Lind et al. 2019). Petri dishes (3.5 cm diameter) with seedlings

were placed at different distances from the gamma source to obtain the average dose rates to

water 1, 10, 20, 40, 100 and 290 mGy h^{-1} (Table 1). The irradiation experiments were

repeated three times. For one out three repeated experiments, the highest dose rate was 250

mGy h^{-1} as 290 mGy h^{-1} could not be obtained due to the decay of the [60]Co source (Table S1).

The dose rates to water and dose rate intervals (in front and back of the petri dishes) were

calculated from the air kerma rates according to (Hansen et al. 2019) as previously described

(Blagojevic et al. 2019). The total doses were calculated by multiplying the absorbed dose

rates to water and exposure time (48 h). During the exposure the room temperature was 20 ˚C

and the photoperiod 12 h with an irradiance of 55 μmol m^{-2} s^{-1} from high pressure metal

halide lamps (HPI-T Plus 250W lamps, Philips, The Netherlands). The red:far red (R:FR)

ratio was measured to 3.5 using a 660/730 nm Skye sensor (Skye Instruments, UK).

Post-irradiation growing conditions

178 The post-irradiation effects of the 48 h of gamma exposure were recorded after transfer of

179 seedlings to pots filled with S-soil (Hasselfors Garden AS, Örebro, Sweden), one plant per

180 pot. All plants were kept in growth chambers (manufactured by Norwegian University of Life

181 Sciences) with temperature, relative air humidity, R:FR ratio, and photoperiod maintained as

182 previously described (Blagojevic et al. 2019).

Growth parameter recordings

At the end of the gamma irradiation, seedlings from both species were placed between two
transparent plastic sheets with a millimetre paper on top and scanned. The shoot and root
lengths of Norway spruce and the lengths of the hypocotyl and cotyledons in *A. thaliana* were
measured using Image J (imagej.nih.gov), with 29-40 and 28-35 seedlings per gamma dose
rate for the respective species.

Post-irradiation, the number of rosette leaves as well as the number of plants with floral buds,
inflorescence and open flower buds were recorded in 23-24 *A. thaliana* plants per gamma
dose rate in time courses up to 75 days post-irradiation (dpi), and percentages of plants with
floral buds, inflorescence and open flowers were calculated. Leaf width, leaf length and
height of the inflorescence were measured at 34, 41 and 64 dpi, respectively. In Norway
spruce, shoot elongation, the number of needles and plant diameter were recorded in a time
course of 59 dpi in 11-22 plants per dose rate. The plant diameter was calculated by
averaging two perpendicular measurements from needle tip to needle tip across the plant at
the shoot apex. Also, plant height was measured from the pot edge to the shoot apical
meristem and the cumulative shoot elongation was calculated.

199 Histological and cytological studies of shoot tips

Histological studies of shoot apical meristems (SAMs) at the end of the gamma irradiation
were performed according to (Lee et al. 2017). Shoot tips of Norway spruce and *A. thaliana*
were then fixed in 4 % paraformaldehyde and 0.025 % glutaraldehyde in phosphate buffered
saline (PBS, pH 7.0), vacuum infiltrated at room temperature and stored at 4°C. The fixed
tissues were washed in PBS solution and dehydrated in a graded ethanol solution.
Approximately 3 mm of the shoot tips of each of 5 plants per treatment were infiltrated and
embedded in LR White (London Resin Company, UK). After polymerisation, 1 µm sections
were made using an Ultracut Leica EM UC6 microtome (Leica, Germany), stained with
toluidine blue O and inspected using a light microscope with bright field optics (DM6B,
Leica).

For cytological studies, LR white-embedded samples were sectioned into 70 nm ultrathin
sections, (Leica EMUC6), stained with uranyl acetate and lead citrate before examining by
transmission electron microscopy (TEM) (FEI Morgagni 268, United States).

213 Comet assay for analysis of DNA damage

Single and double-strand DNA breaks at the end of the gamma irradiation and 77 dpi were

quantified using the comet assay (Gichner 2003) with some modifications (Blagojevic et al.

2019). Briefly, this included isolation of cell nuclei and electrophoresis under alkaline pH,

resulting in a comet-like appearance of damaged DNA due to movement out of the nuclei.

This was followed by quantification of the DNA in the comet tail using fluorescence

microscopy. Three biological replicates per dose rate, each consisting of 3-4 mm of shoot tips

from 3 plants of Norway spruce or 5 plants of *A. thaliana*, were used to assess DNA damage.

For each sample, about 180-200 cell nuclei were scored, with 60-70 nuclei scored in each of

3 technical replicates (gels). and the median value was calculated.

RNA isolation for RNA sequencing

At the termination of the 48-h gamma irradiation, four biological replicates per gamma dose

rate (0, 1, 10, 40, 100 (both species) and 290 mGy h^{-1} (*A. thaliana* only due to lack of growth

inhibition, organelle and DNA damage at lower dose rates)) were harvested in liquid

nitrogen. For *A. thaliana* (20-30 shoots per dose rate), total RNA was extracted using the

RNeasy Plant Mini Kit (QIAGEN Inc., Valencia, CA) following the manufacturer's protocol.

For Norway spruce (four shoots per dose rate), total RNA was extracted using the Masterpure

Complete DNA and RNA Purification Kit (Epicenter, Madison, USA) following the

manufacturer's protocol, except that 3 µl beta-mercaptoethanol per sample replaced 1,4-

dithiothreitol (DTT) and 0.5% polyvinylpyrrolidone (PVP, mw 360 000, Sigma Aldrich,

Germany) was added to the extraction buffer. The purified RNA was quantified using a

NanoDrop ND-1000 Spectrophotometer (Thermo Scientific, USA) and the quality assessed

with an Agilent 2100 Bioanalyzer with an RNA 144000 Nano Kit (Agilent technologies,

USA).

RNA sequencing

The transcriptomes of the shoots were sequenced with two repeated lanes per sample, on an

Illumina HiSeq4000 platform at the Norwegian Sequencing Centre (Oslo, Norway) using the

Strand specific 20xTruseq RNA library preparation (paired end, read length 150 bp) as

recommended by the manufacturer (Illumina, San Diego, CA, USA). The raw sequence data

were submitted to the ArrayExpress under the accession number E-MTAB-8081 for Norway

spruce and E-MTAB-11120 for *A. thaliana.*

Differential gene expression analyses and functional term enrichment

The transcript abundances were estimated using the Salmon program (Patro et al. 2017),

comparing with the genomes of either Norway spruce (*Picea abies* 1.0 assembly,

plantgenie.org) (Nystedt, et al. 2013) or *A. thaliana* (TAIR10, arabidopsis.org) (Berardini et

al. 2015). Gene expression was compared between the samples at different gamma radiation

dose rates and the unexposed control samples using the DESeq2 (Love et al., 2014). Genes

were classified as differentially expressed genes (DEGs) if the False Discovery Rate

(Benjamini & Hochberg correction) adjusted p-values were < 0.05.

Gene Ontology (GO) annotations of Norway spruce genes, as well as spruce to *A. thaliana*

orthologs by best sequence matches, were downloaded from plantgenie.org. The *A. thaliana*

orthologs were used to associate the Norway spruce genes with KEGG pathways. Enrichment

of GO or KEGG terms in sets of DEGs was found using topGO (version 2.34.0) or kegga

from the limma package (version 3.38.2), respectively.

RT-qPCR analysis

Based on the significant regulation in one or both species in response to 48 h gamma

irradiation, 11 DNA repair and cell cycle control genes were selected for further RT-qPCR

validation and analysis of their regulation after 12, 24 and 48 h of gamma irradiation at 1, 10,

20, 40 and 100 mGy h^{-1} (Table S2, S3). RNA was isolated as described above from four

biological replicates per dose rate, each consisting of shoots from eight and 30 seedlings for

Norway spruce and *A. thaliana*, respectively. . cDNA was synthesised using SuperScript™

IV VILO™ cDNA Synthesis Kit (Invitrogen, Carlsbad, CA, US) following the

manufacturer's protocol, and no enzyme controls (-RT) were made for each sample.

Gene sequences were retrieved from NCBI (ncbi.nlm.nih.gov/gene) and TAIR

(arabidopsis.org) for *A. thaliana* and plantgenie.org for Norway spruce and exon-exon-

spanning primers were designed to avoid non-specific amplification from any genomic DNA

contamination in the RNA samples, using the online tools Oligoanalyzer

(eu.idtdna.com/calc/analyzer) and Primer Blast (ncbi.nlm.nih.gov/tools/primer-blast) with

default parameters. Amplifications were performed in a 7500 Fast Real-time PCR system

(Applied Biosystems, Foster city, USA) using Platinum Quantitative PCR Supermix-UDG

and SYBRGreen (Thermo Fisher, Carlsbad, CA, US), using standard protocol.

To compare the relative transcript levels of the selected genes in the samples exposed to

different gamma radiation dose rates at different time points with the unexposed controls (0

mGy h^{-1}), the comparative ΔΔCT method was used (Livak and Schmittgen 2001). The

expression of the target genes was normalised by the mean expression of the three reference

genes, *ACTIN* (*ACT*), *ELONGATION FACTOR 1α* (*EF-1α*) and *UBIQUITIN*.

Statistical analysis for growth recordings, DNA damage and qPCR results

DNA damage, hypocotyl, cotyledon and shoot length, as well as post-irradiation growth

parameters at the final sampling time (34 or 41 dpi for *A. thaliana* and 59 dpi for Norway

spruce) were tested for normality and homoscedasticity by the Ryan-Joiner and Levene tests,

respectively. One-way ANOVA tests (glm model in Minitab 21.1, Minitab Inc, PA, USA)

were then used to test for differences in these response variables between plants exposed to

48 h of gamma radiation versus control plants. Due to the lack of normality and

homoscedasticity of the non-transformed RT-qPCR gene expression data, log-transformed

data was used in two-way ANOVA for individual species (dose rate and gamma-irradiation

duration as factors) and three-way ANOVA for comparison of species (plant species, dose

rate and gamma-irradiation duration as factors). Tukey`s post hoc test was used to test for

differences between means ($p \leq 0.05$).

Results

Gamma irradiation for 48 h affects growth negatively in Norway spruce but not

A. thaliana

In *A. thaliana*, no significant effect of the gamma irradiation on hypocotyl and cotyledon

lengths was identified (Fig. 1a,b). At 41 dpi, no significant dose response-relationship was

observed for the length, width, or number of leaves (Fig. S1a,b,c). Although post-irradiation

flowering was slightly delayed at 250-290 mGy h^{-1}, eventually all plants flowered (63 dpi)

and showed very similar overall morphology, flowering, and silique development (77 dpi)

(Fig. 1c, S1d,e,f). In Norway spruce, the shoot length was significantly reduced (27%) after

48 h of gamma irradiation at 290 mGy h^{-1} only, but the root length was not affected (Fig.

1d,e). At 59 dpi, the shoot diameter and number of needles were significantly reduced by 5%

and 12%, in 100 mGy h^{-1} and 9% and 35% at 290 mGy h^{-1}, but no such effects were detected

at 77 dpi (Fig. 1f,g,h, S2). Hence, there was temporary growth impairment in Norway spruce

after exposure, while development in *A. thaliana* remained unaffected.

More severe gamma radiation-induced organelle damage in Norway spruce than *A. thaliana*

Normal dome-shaped shoot apical meristems with compact cells were observed in both species after 48 h of gamma irradiation (Fig. 2). In *A. thaliana*, slightly damaged mitochondria were observed at 100-290 mGy h^{-1}, but the chloroplasts and cell nuclei appeared normal at all dose rates compared to the unexposed control except that a few cells showed less dense nuclei morphology at $\geq$ 1 mGy h^{-1} (Fig. 3a,b,c). In comparison, in Norway spruce, disrupted nuclear and mitochondrial membranes were observed at $\geq$ 1 mGy h^{-1}, as well as swollen chloroplasts and disrupted thylakoid membranes at $\geq$ 40 mGy h^{-1} (Fig. 3d,e,f).

Higher levels of DNA damage in gamma-irradiated Norway spruce than *A. thaliana*

In *A. thaliana*, the Comet assay showed low but significant damage with $\leq$ 5% tail DNA at 1-100 mGy h^{-1} and 12% tail DNA at 290 mGy h^{-1} (Fig. 4a). In comparison, DNA damage was consistently 2-8-fold higher in Norway spruce with 10%, 12%, 15%, 17.5% and 40% DNA in the Comet tail after the 48-h of irradiation at 1, 10, 40, 100 and 290 mGy h^{-1}, respectively (Fig. 4b). At 77 dpi, the DNA damage was substantially reduced but still observed in *A. thaliana* at $\geq$ 10 mGy h^{-1} with $\leq$ 3% tail DNA and in Norway spruce seedlings at $\geq$ 1 mGy h^{-1}, with $\leq$ 5% tail DNA at all dose rates (Fig. 4a,b).

Profound transcriptome differences between gamma-irradiated Norway spruce and *A. thaliana*

Comparative RNA-seq analysis of gamma-irradiated shoot samples (1-100 mGy h^{-1} both species, 290 mGy h^{-1} *A. thaliana* only), revealed extensive differences between the species. *A. thaliana* expressed a total of 26,643 genes, of which 11,592 were DEGs between any dose rate and the unexposed control (43.5%) in one or multiple gamma dose rates (Fig. 5a). Norway spruce expressed a total of 66,069 genes, of which 5326 were DEGs (8.1%, Fig. 5c). In *A. thaliana*, a high number of DEGs was observed already from 1 mGy h^{-1} (3442 DEGs), with the higher dose rates having ~4400-5500 DEGs (Fig. 5a,b, S3a). At 100-290 mGy h^{-1}, there was a higher proportion of upregulated genes with more than 2-fold change as compared to the lower dose rates (Fig. 5a, S3b). By contrast, in Norway spruce, very few DEGs were observed at 1-10 mGy h^{-1} (38 and 43 DEGs), while 822 and 5047 DEGs were observed at 40-100 mGy h^{-1}, respectively (Fig. 5c,d). The increase in DEGs from 40 to 100

mGy h^{-1} was much more pronounced in Norway spruce than in *A. thaliana* (Fig. 5a,b). The

largest overlap of *A. thaliana* DEGs was between 100 and 290 mGy h^{-1}, and between 1, 10

and 40 mGy h^{-1} (Fig. 5b). A good proportion of Norway spruce DEGs at 40 and 100 mGy h^{-1}

overlapped, showing some consistency in the response even with a large difference in total

numbers of DEGs, including many unique ones (Fig. 5d).

Distinctive pathway enrichment in Norway spruce and *A. thaliana* at different

dose rates

GO and KEGG pathway analyses on sets of DEGs revealed a gamma dose rate-dependent

enrichment of different pathways. In *A. thaliana*, DNA replication, DNA repair, cell cycle,

endoreduplication, and endoplasmic reticulum (ER) stress were enriched at 1-40 mGy h^{-1}

(Fig. S4, S5). Hormone biosynthesis and signalling pathways showed considerable

enrichment along with stress response at 100-290 mGy h^{-1} (Fig. S4, S5). Oxidative stress,

defence response, growth and development, protein synthesis and catabolism as well as

mitogen-activated protein kinase (MAPK)-mediated signalling were significantly regulated at

all dose rates (Fig. S4, S5). In Norway spruce, pathways such as DNA replication, DNA

repair, cell cycle, endoreduplication, unfolded protein response (UPR) and ER stress,

response to gamma, energy metabolism and growth and development were all enriched at $\geq$

40 mGy h^{-1} (Fig. S6, S7). Photosynthesis and protein biosynthesis and metabolism were

enriched in downregulated DEGs at 100 mGy h^{-1} (Fig. S6, S7) in Norway spruce only.

In *A. thaliana* KEGG pathway analysis revealed that the DDR pathways such as HR,

mismatch repair (MMR), non-homologous end-joining (NHEJ) and base excision repair

(BER) were regulated at 10-40 mGy h^{-1}, and BER also at 290 mGy h^{-1} (Fig. S5). In Norway

spruce, NER, MMR, HR and NHEJ pathways were significantly regulated at 40-100 mGy h^{-1}

and BER at 40 mGy h^{-1} only (Fig. S7).

Distinct differences in stress response pathways between Norway spruce and *A.*

thaliana

The specific DEGs in DDR, antioxidant protection and other stress response pathways were

further analysed to compare *A. thaliana* and Norway spruce response after the 48- h of

gamma irradiation.

DNA damage repair

In *A. thaliana*, most DEGs in the DDR pathways showed a 30-70% change in expression (Table S3). Genes involved in DNA double strand break (DSB) repair via HR, including *RAD51*, *SMC6B*, *PCNA2* and *BRCA2B*, showed significant upregulation at all dose rates in the 10-290 mGy h^{-1} range. In Norway spruce, most DDR DEGs at 40-100 mGy h^{-1} had at least a 60% change in expression (Table S3), and genes involved in DSB repair via HR; *SRS2*, *RAD52*, *RPA1A*, *MCM8, XRCC2* and *XRCC3*, were all upregulated at 100 mGy h^{-1}. A few other genes such as *PARP1* and *PARP2* were significantly upregulated at the highest dose rates in both species. Some DEGs showed opposite changes in expression between species; *MRE11*, *RPA2A/ROR1* and *BRCA1* in the HR pathway, and *MSH7* in the MMR pathway. Unique DEGs to Norway spruce included upregulation of *KU80* at 40-100 mGy h^{-1} and *SOG1* at 100 mGy h^{-1}. Unique to *A. thaliana* was upregulation of *ATM* at 100 mGy h^{-1}.

Cell cycle and endoreduplication

In *A. thaliana*, most cell cycle-related DEGs including *CYCBs* and *CDKBs*, were observed at 1-40 mGy h^{-1} but not 100-290 mGy h^{-1} (Table S4). *CYCB1* was significantly upregulated at all dose rates ≤ 40 mGy h^{-1}, along with many other *CYCBs*. In Norway spruce, no such effects on B type cyclins were observed and most of the cell cycle genes showed changes in expression at ≥ 40 mGy h^{-1}. Anaphase-promoting factors *APC6* and *13*, chromatin assembly factor *FAS2*, and the cyclin-dependent kinases regulatory subunit *CKS1* showed upregulation at ≥ 40 mGy h^{-1}. Positive regulators of the cell cycle such as D type cyclins (e.g., *CYCD1*) were significantly upregulated in *A. thaliana* at ≤ 40 mGy h^{-1} but downregulated in Norway spruce at 100 mGy h^{-1}. *RBR1* was upregulated in both species. *E2Fb* was upregulated at 40 mGy h^{-1} in Norway spruce only. *WEE1,* an important G2/M checkpoint regulator, was upregulated at ≤ 40 mGy h^{-1} in *A. thaliana*, but at ≥ 40 mGy h^{-1} in Norway spruce. Most of the cell cycle-related genes, such as *CYCs* and *CDKs*, also involved in the endocycle and the endoreduplication-related biological pathway, were enriched in *A. thaliana* at ≤ 40 mGy h^{-1} but very few such DEGs were observed in Norway spruce (Table S5).

Chromatin remodelers and genes in epigenetic pathways

Significant activation of genes involved in the pathway associated with de-condensation of the nucleosome structure and activation of DDR pathways, was predominantly observed in Norway spruce at 40-100 mGy h^{-1} (Table S6). *In A. thaliana*, a few such genes were upregulated at ≤ 40 mGy h^{-1} only, and a few were downregulated at higher dose rates. The DNA helicase *DDM1* was upregulated in *A. thaliana* at ≤ 40 mGy h^{-1} and DNA

398 methyltransferase *CMT3,* histone 2A variant *HTA8* and a component of the core chromatin

399 remodelling complex *ARP6* were significantly downregulated at 100-290 mGy h^{-1} (Table S6).

400 In Norway spruce, histone deacetylase *HOS15* genes involved in the regulation of DNA

401 methylation, such as *DMT1*, *IDN2* and *SUVH4*, as well as the histone 2 A variant *H2AV*,

402 showed significant upregulation at 40-100 mGy h^{-1}.

Hormone biosynthesis and signalling

A. thaliana genes involved in biosynthesis and signalling of main growth-promoting

hormones such as auxin, cytokinin, auxin, cytokinin, gibberellin (GA) and brassinosteroid, as

well as important stress-response-related hormones such as abscisic acid (ABA) and ethylene,

were significantly regulated in response to the gamma irradiation (Table S7) especially at

higher dose rates. The auxin-responsive MYB transcription factors *AtMYB93* and *MYB61*, the

auxin-responsive genes *IAA13, IAA14, IAA27, SAUR10* and *GH3.6*, the auxin biosynthesis

gene *YUC9* as well as the auxin efflux carriers and transporters involved in polar auxin

transport, *PIN2*, *PIN3*, *AUX1* and several ABC transporters, were significantly upregulated in

this species (Table S6) but downregulated in Norway spruce. Although the auxin-

homeostasis-related gene *IAMT1*, and the auxin responsive genes *SAR1* and *SAUR76* showed

increased expression at 40-100 mGy h^{-1} in Norway spruce, several auxin signalling genes

including *ARF6, TIR1, AIR9* and *IAA26* were significantly downregulated (Table S7). In *A.*

thaliana, cytokinin biosynthesis, metabolism as well as cytokinin signalling genes such as

CKX5, LOG7, AHP1, SOT8 ARR2 and *ARR3*, were upregulated at 40-290 mGy h^{-1} (Table

S7). Contrastingly, none of these genes showed significant regulation in Norway spruce.

Similarly, the GA biosynthesis genes *GA2* (*ent-kaurene synthase 1*) and *GA 20-OXIDASE 2*

(*GA20OX2),* and the brassinosteroid biosynthesis and homeostasis genes *CYP90C1,*

CYP708A1, and *CYP85A1*, were upregulated in *A. thaliana* but not in Norway spruce.

In general, ABA biosynthesis and signalling genes did not show any gradual up- or

downregulation in *A. thaliana* (Table S7). However, the negative regulator of ABA

biosynthesis *PUB44*, and the ABA signalling genes *PYL4, CRK29* and *ATHB-6* were

downregulated at <40 mGy h^{-1}. Many stress- and ABA-responsive genes such as *ASPG1,*

PPCK1, and *RD22* showed consistent increasing expression from 10-290 mGy h^{-1}. Among

the genes regulated in both species, the ABA signalling gene *CIPK3* showed downregulation

in all dose rates in *A. thaliana*, and at 100 mGy h^{-1} in Norway spruce, and *ADH1* were

upregulated in both species at higher dose rates. In *A. thaliana*, ethylene biosynthesis and

signalling genes were mostly downregulated at lower dose rates with few exceptions (Table

S7). Ethylene-responsive genes such as *CTR1, RAP2-2, RAP2-12,* and *ERFs* as well as the

ethylene biosynthesis genes *EOL1* and *ACS5* showed significant regulation in *A. thaliana*, but

not in Norway spruce.

Genes in antioxidant metabolism pathways

Significant overall upregulation of antioxidant biosynthesis and signalling genes was

observed in *A. thaliana* even from 1 mGy h^{-1}. By contrast, in Norway spruce, most of the

antioxidant genes were not significantly affected below 100 mGy h^{-1} (Table S8). The key

flavonoid and anthocyanin biosynthesis genes *CHS* and *DFRA* were upregulated in *A.*

thaliana from 1 mGy h^{-1} but significantly downregulated in Norway spruce at 100 mGy h^{-1}.

Flavanol synthase (*FLS1*) was upregulated in *A. thaliana* at ≤ 10 mGy h^{-1} and Norway spruce

at 100 mGy h^{-1}. Many *PER* genes including *PER3* and *PER11* were regulated in both species

but from lower dose rates in *A. thaliana* than Norway spruce. *GLUTATHIONE-S-*

TRANFERASEs (*GSTU/GSTF*) were upregulated in both species.

Cell wall component genes

In *A. thaliana* genes involved in the biosynthesis of cell wall components, cell wall

maintenance and fortification, as well as membrane receptors were significantly upregulated

at 100-290 mGy h^{-1} with large fold-changes. In Norway spruce, only very few genes related

to cell wall biogenesis and maintenance showed regulation, i.e., at 40-100 mGy h^{-1} (Table

S9). In *A. thaliana*, *BOR2* involved in cellulose and pectin metabolism, *HHT1* involved in

pectin and suberin biosynthesis, the cell wall organisation and biosynthesis genes *FACT* and

CYP86B1, were upregulated at 100-290 mGy h^{-1}. The endoglucanases *GH9C2* and *GH9B13*

involved in cell wall organisation were upregulated in all dose rates. Although none of those

genes showed any significant changes at any dose rate in Norway spruce, a few other genes

involved in biogenesis and metabolism of expansins, cellulose and xyloglucan such as the

EXP4 and 8, CESA5 and *XTH 5* and *9* were significantly regulated in both species.

Photosynthesis and energy metabolism

Photosynthesis was massively downregulated in Norway spruce but not in *A. thaliana*.

Photosystem (PS) I and II-related genes such as *PS I reaction centre subunit*s *PSAG, PSAK*

and *PSAN*, PS 1 light harvesting complex *LHCA1*, PS II assembly factor *HCF136*, essential

enzymes such as NADPH-protochlorophyllide oxidoreductase *PORA*, and *Fructose-1,6-*

bisphosphatase CYFBP, chloroplast binding proteins and membrane efflux carriers, were

downregulated in Norway spruce at 100 mGy h^{-1} (Table S10). In *A. thaliana,* only a few

genes were regulated, such as upregulation of the PS I and II reaction centre components

PSAB and *PSBB*, and *NDHC* (Table S10).

Similarly, energy metabolism was far more affected in Norway spruce than *A. thaliana*

(Table S11). Norway spruce particularly showed upregulation at ≥ 40 mGy h^{-1} of a wide

range of genes involved in glycolysis, TCA cycle and gluconeogenesis, such as,

Glyceraldehyde-3-phosphate dehydrogenase *GAPC1*, *6-PHOSPHOFRUCTOKINASE 4*

(*PFK4*), *MALATE DEHYDROGENASE* (*MDH1*), *SUCCINATE DEHYDROGENASE (SDH2-*

1), and several homologs of Phosphoenolpyruvate carboxykinase *PCKA* and *PFK2*.

Additionally, the *CYTOCHROME C1* (*CYTC-1*) and *CYTC OXIDASE 6B1* (*COX6B-1*) genes

involved in mitochondrial electron transport were significantly upregulated at 100 mGy h^{-1}.

ALCOHOL DEHYDROGENASE 1 (*ADH1*) was significantly upregulated in both species

from 100 mGy h^{-1}.

Differential dynamic regulation of DNA repair genes in *A. thaliana* and Norway

spruce during gamma irradiation

To understand the dynamics of DDR and cell-cycle-related gene expression during radiation

stress, both species were exposed to 12, 24 and 48 h of gamma radiation and the transcript

levels of selected genes analysed by RT-qPCR. Consistent with the RNA-seq data,

significantly increased transcript levels of *WEE1, PARP2, RAD51* and *CYCB1.2* were

observed after 48 h in *A. thaliana*, by 5-10-fold as compared to the unexposed controls (Fig.

6). However, after 12 and 24 h exposure no significant effect of the gamma irradiation on

their expression was observed. By contrast, in Norway spruce the strongest response was

generally observed after 24 h of gamma irradiation (Fig. 7). The transcript levels of *WEE1*

increased significantly at 20-100 mGy h^{-1} after 24 h only and those of *SOG1* increased at 40-

100 mGy h^{-1} after 24 h, while *PARP2* expression increased significantly by 10-30-fold at 40-

100 mGy h^{-1} after 24 and 48 h. There were no significant effects on expression at 12 h.

Discussion

To improve the knowledge base for risk assessment and protection of plant communities in

radionuclide-contaminated ecosystems against ionising radiation, the current comparative

study has focussed on underlying mechanisms with respect to differential radiosensitivity in

the highly radio-sensitive conifer Norway spruce and the radio-tolerant *A. thaliana*. Through

exposure of seedlings to gamma radiation under standardised conditions and comparative assessment of phenotypic, cellular, and genotoxic effects, combined with RNA-seq and qPCR analysis, we present a novel in-depth understanding of such differences between the two species in the mechanisms involved in protection and repair.

Previously, we have shown dose rate-dependent impaired development in Norway spruce seedlings exposed to gamma radiation for 144 h, with significant growth inhibition at $\geq$ 40 mGy h^{-1} and progressively developing after-effects eventually causing mortality (Blagojevic et al. 2019). By contrast, *A. thaliana* showed no significant adverse effects at dose rates $\leq$ 540 mGy h^{-1} (Blagojevic et al. 2019a). Norway spruce recovers better from inhibited growth 77 dpi after 48 h exposure to 290 mGy h^{-1} (Fig. 1d,f,h) compared to the previous study (Blagojevic et al. 2019) despite the total dose after 48 h being higher (13,920 mGy) than after 144 h (5,760 mGy), emphasising the severity of chronic exposure. Consistent with previous findings, *A. thaliana* did not show any significant growth or developmental effects in this study (Fig. 1a,b,c). The RNA-seq analysis revealed, however, profound differences between the two species in response to gamma irradiation (Fig. 5, S4, S5). The fact that *A. thaliana* mounts a transcriptomic response at lower dose rates than Norway spruce signified that it was more poised towards rapid response to inflicted radiation stress.

The energy deposited at a specific gamma radiation dose rate in the two plant species was similar, with ionization and excitation of electrons occurring uniformly throughout the cells. Thus, induced damage should be randomly distributed in biomolecules such as DNA and relative to the DNA content (genome size) (Einset and Collins 2018). Consistent with previous report (Blagojevic et al. 2019) this study showed less DNA damage in *A. thaliana* than in Norway spruce both at the end of the irradiation and post-irradiation period (Fig. 4). We therefore hypothesised that *A. thaliana* has more effective DNA repair and/or protection systems. Yet, the DNA damage at 77 dpi in both species (Fig. 4), albeit at lower level than at the end of the 48-h irradiation, demonstrates that some DNA damage or genomic instability is not critical to survival or development. Although genomic instability may result into detrimental growth effects (Mothersill and Seymour 1998), no clear negative effects on the vegetative growth post-irradiation in *A. thaliana* even after exposure to 400-540 mGy h^{-1} in contrast to the mortality observed in Norway spruce and Scots pine at dose rates > 40 mGy h^{-1}, (Blagojevic et al. 2019) confirms better tolerance in *A. thaliana* compared to conifers. In our current study, no mortality after 48 h of gamma irradiation in either species again points to a certain tolerance to damage in both species.

The comparative RNA-seq analysis revealed that both *A. thaliana* and Norway spruce
initiated a plethora of DDR pathways to avoid harmful mutations and maintain genome
integrity. However, the response of the two species differed in two main aspects (Fig. S9).
Firstly, the upregulation of DDR-related genes at lower doses dose rates in *A. thaliana* than in
Norway spruce suggested that the former is far more responsive towards lower levels of DNA
damage. Secondly, the prominent DNA repair pathways in *A. thaliana* were mostly HR, and
to some extent MMR, whereas in Norway spruce, in addition to HR, several other pathways
such as NHEJ, MMR and DNA transleison synthesis were activated. Hence, it is tempting to
speculate that Norway spruce needed to engage a wider range of DDR because of the
significantly higher levels of DNA damage compared to *A. thaliana*, possibly because of less
antioxidant protection and a larger genome.

One of the key contributor genes in gamma radiation-induced DDR in plants*, SOG1*, which
activates several DDR pathways and cell cycle checkpoints (Kim et al. 2019), was not
affected at any time points or dose rates tested in the gene expression analysis (Table S3, Fig.
6). This might indicate that induction of *SOG1* expression is only modulated in higher total
doses or acute radiation that generate high level of DNA damage (Bourbousse et al. 2018).
However, no change in *SOG1* expression after 48 h of gamma irradiation does not necessarily
indicate the lack of SOG1 activity as that could rather depend on the post-translational
modification by ATM- or ATR-kinases (Bourbousse et al. 2018; Yoshiyama et al. 2013). The
involvement of *SOG1* in *A. thaliana* is indeed supported by the activation of many primary
target genes of SOG1 (i.e., *RAD51, BRCA1, PARP1* and *2, GMI1*) in response to the 48 h of
gamma irradiation. By contrast, significant upregulation of *SOG1* in Norway spruce after 24
h at $\geq$ 40 mGy h^{-1} and 48 h at 100 mGy h^{-1} (Fig. 7) but no significant regulation of ATM
(Table S3), might imply that elevated *SOG1* expression was triggered in Norway spruce
possibly due to higher levels of accumulated DNA damage. Moreover, upregulation of
WEE1, a direct target of SOG1 and key regulator of CDK-inhibition and stress-induced cell
cycle arrest (Kevei et al. 2011; De Schutter et al. 2007), coincided with *SOG1* expression in
Norway spruce, strongly suggesting SOG1 involvement in gamma-induced DDR and cell
cycle arrest (Pedroza-Garcia et al. 2022; Bourbousse et al. 2018).

Transcriptional activation of numerous B type *CYC-* and *CDKB*-genes in *A. thaliana* after 48
h at 1-40 mGy h^{-1} (Table S4) might indicate gamma radiation stress-induced cell cycle arrest
and subsequent DDR through HR by maintaining higher cyclin threshold (Endo et al. 2006).
This further supports the previous notion that enhanced CDKB1 activity is involved in

ionising radiation-induced cell cycle arrest leading to HR-repair in damaged mitotically

active cells (Weimer et al. 2016). On the contrary, there were few regulations of B-type *CYC*

or *CDK* genes in Norway spruce (Table S4), indicating no significant involvement of such

genes in cell cycle arrest. The regulation of the key contributors in cell cycle progression in

both species, albeit to a larger degree in *A. thaliana*, suggests that the cell cycle is one of the

most affected biological pathways in the applied gamma radiation. Most of the CYCs and

CDKs regulated in *A. thaliana* in our study are also involved in the regulation of the

endocycle (Table S5). However, stress-indued endoreduplication is less common in

gymnosperms than angiosperms (De Veylder et al. 2001) and very few genes related to

endocycle were upregulated in Norway spruce. Increased endoreduplication was previously

shown in *A. thaliana* at 1 Gy min^{-1} (150 Gy totally) and *Lemna minor* at 1500 mGy h^{-1} for 7

days, but not at lower dose rates (Adachi et al., 2009; Van Hoeck et al., 2015). Our

preliminary study did not reveal any effect of 48 h gamma radiation at 100 mGy h^{-1} on

endoreduplication in *A. thaliana* and Norway spruce (flow cytometry results not shown),

suggesting that the applied dose was not sufficient.to induce endocycle.

DDR is closely related to epigenetic regulation and chromatin structure must be decondensed

for the DDR proteins to carry out efficient repair (Choi, et al. 2019; Kim, et al. 2019).

Although shown previously in *A. thaliana* exposed to ionising radiation (Sidler et al. 2015),

very few chromatin remodelers were regulated in *A. thaliana* in our study. By contrast, the

upregulation of *H2A* variants, DNA methylation and deacetylation-related genes particularly

in Norway spruce (Table S6, Fig. S6), clearly indicated chromatin modification at $\geq$ 40 mGy

h^{-1} in this species. The strong enrichment of DNA methylation regulation is consistent with

the previously reported DNA hypermethylation in *P. sylvestris* trees exposed to high doses of

ionising radiation in the Chernobyl area, a mechanism suggested to stabilise the genome

(Kovalchuk et al. 2003). Despite the genome of Norway spruce being 150 times bigger than

that of *A. thaliana*, it only contains 2.5 times more genes (Nystedt, et al. 2013), and has large

heterochromatin regions which requires the activation of chromatin remodelers for

decondensation for efficient DDR (Fig. S9). Moreover, gamma radiation-induced damage to

heterochromatin regions does not initiate rapid activation of DDR pathways (Falk et al.

2008). It is thus plausible that Norway spruce requires a significantly more DNA damage for

the activation of DDR and chromosome remodelers.

The 48-h exposure did not affect the structural integrity of the SAMs (Fig. 2) in contrast to

the previous studies where 144 or 360-h gamma irradiation resulted in adverse effects in

SAMs of Norway spruce only (Blagojevic et al. 2019). We therefore proceeded to investigate

the effect of the radiation exposure on organelles. TEM image analysis revealed that integrity

of thylakoid and nuclear membranes was compromised in Norway spruce, but not in *A.*

thaliana (Fig. 3). Notably, in both species the mitochondria appeared to be the most

vulnerable organelle. Furthermore, the fact that disrupted mitochondria were evident from 1

mGy h^{-1} in Norway spruce, while *A. thaliana* showed no damage below 100 mGy h^{-1},

indicates a major difference in their capacity to maintain cell homeostasis.

Former studies of gamma-irradiated plant species such as *A. thaliana* and *L. minor* have

demonstrated activation of ROS-scavenging enzymes such as SOD and APX (Vanhoudt, et

al. 2014; Xie et al. 2019), but such information is limited in conifers. Consistent with the

notion that species-specific antioxidant response determines oxidative stress-induced damage,

less activation of such genes occurred in Norway spruce < 40 mGy h^{-1}, comprehensive

modulation of enzymatic and non-enzymatic antioxidant-related genes (*PERs, CHSs, GSTs*,

anthocyanin and flavanol synthesis genes) in *A. thaliana* (Table S8), could at least partly

explain the massive damage to mitochondria, chloroplasts, and nuclear membranes at low

dose rates in Norway spruce in contrast to only slight mitochondrial damage in *A. thaliana*

(Fig. 2 and 3). Moreover, antioxidants such as *PERs* can modulate cell wall composition

during stress (Díaz-Tielas et al. 2012). Cellulose and pectin were previously shown to be

damaged by very high doses of gamma radiation (1-2.5 kGy) (Kovács and Keresztes 2002).

Hence, in contrast to Norway spruce, the radical upregulation of cell wall component genes

even at lower gamma dose rates in *A. thaliana* (Table S9) might suggest efficient

replacement of damaged cell wall components and cell wall fortification in *A. thaliana* (Le

Gall et al. 2015).

Antioxidants such as *GST*s may also downregulate auxin-responsive genes (Sylvestre-Gonon

et al. 2019; Jiang et al. 2010). The upregulation of *GSTs* and growth retardation in Norway

spruce at higher dose rates as well as downregulation or no regulation of growth-promoting

hormone pathways are consistent with this notion (Fig.1, S1, Table S7). By contrast, the

gamma-irradiation-induced upregulation of auxin and cytokinin-related genes (biosynthesis,

metabolism, signalling, auxin transport) in *A. thaliana* complies with the maintenance of

meristem function to sustain normal growth. These results are in line with the previously

demonstrated activation of such genes in response to abiotic stress, indicating the importance

of growth-hormone homeostasis under stress (Verma *et al*., 2016). Similar to previous reports

on the induction of ABA biosynthesis and signalling genes triggered by ROS in gamma-

irradiated *A. thaliana* (Vishwakarma et al. 2017), upregulation of many such ABA-responsive

genes were observed in *A. thaliana* in our study, although the total doses applied were much

lower. Also, upregulation of the ethylene-responsive oxidative stress-pathway (*ERFs*, *ACSs*,

MPK6) and SA-responsive genes (*WRKY*, *MYB*) in *A. thaliana*, but not Norway spruce (Table

S7), indicate less induction of hormone response to gamma-stress in the latter and is

consistent with the greater tolerance of *A. thaliana* to gamma radiation. Hence, in *A. thaliana*

better antioxidant homeostasis, cell wall biosynthesis/fortification and hormone signalling

might contribute to the tolerance to the gamma irradiation. Conversely, less transcriptional

activation of the same protective pathways in Norway spruce possibly have resulted in

adverse effects on overall growth and higher susceptibility to radiation-induced oxidative

damage.

The RNA-seq analysis provided evidence that energy metabolism and photosynthesis in

Norway spruce were strongly affected by gamma irradiation in Norway spruce, but only

marginally in *A. thaliana*. In Norway spruce, the reduced expression of photosynthesis-

related genes coincided with the reduced growth and chloroplast impairment at higher dose

rates ($\geq$ 40 mGy h^{-1}). Concurrently, the strong upregulation of energy metabolism-related

genes (glycolysis, Krebs cycle, mitochondrial electron transport-related genes) (Table S11) in

this species may support the hypothesis that there is a high ATP requirement for damage

repair including maintenance of the large genome. Hence, it appears that radiation induces

energy deficiency, which in turn requires increased energy expenditure on damage repair at

the expense of growth and is reflected by the downregulation of growth-promoting hormones

in Norway spruce at higher dose rates. However, the upregulation of energy-metabolism-

genes might additionally indicate that the damage to mitochondria at high dose rates could

have led to inefficient energy production and thus a need for the activation of respiratory

genes to meet the energy demands.

In summary, this study demonstrates that compared to the radiosensitive Norway spruce, the

radiotolerant *A. thaliana* mobilises more comprehensive gamma radiation-induced defence,

repair, and homeostasis-maintaining stress responses, even at low dose rates. Despite

significant adverse effects in Norway spruce related to DNA and organelle damage and

growth at low dose rates, transcriptional activation of DDR was evident at 40-100 mGy h^{-1}

only Furthermore, in Norway spruce, mobilisation of antioxidants, as well as modulation of

growth-regulatory and stress-related hormones at lower dose were less efficient than in *A.*

thaliana. Thus, the short-lived *A. thaliana* with quick generation-time, and small genome

with less energy requirement for maintenance, shows better plasticity in response to radiation
stress compared to the long-lived Norway spruce with lengthy generation time and gigantic
genome. These observations expand our understanding of differential radiosensitivity in
plants, and thus improves the knowledge base for risk assessment and protection of
ecosystems from ionising radiation exposure. In the context of environmental impact, the
findings strongly support that under severe ecosystem exposure situations, radiosensitive
conifers require special attention with respect to remediating measures.

Acknowledgments

666   This study has been funded by the Research Council of Norway through its Centre of
667   Excellence (CoE) funding scheme (Project No. 223268/F50) and the Norwegian University
668   of Life Science (NMBU). Sincere thanks to Marit Siira and Svein Andre Kolltveit for skillful
669   assistance in plant growing and growth measurements and Tone Ingeborg Melby and Linda
670   Ripel for skillful work with qPCR analyses of gene expression.

Competing interests

672   None declared

Author contributions

Payel Bhattacharjee: Conceptualization, Data curation, Formal analysis, Investigation,
Methodology, Software, Validation, Visualization, Writing - original draft, Writing - review
& editing; Dajana Blagojevic: Conceptualization, Data curation, Formal analysis,
Investigation, Methodology, Writing – review and editing; YeonKyeong Lee: Methodology,
Visualization, Formal analysis, Writing – review and editing; Gareth B Gillard: Data
curation, Formal analysis, Visualization, Writing - review & editing; Lars Grønvold, Torgeir
R Hvidsten, Simen R Sandve: Data curation, Formal analysis, Writing – review and editing;
Ole C Lind: Conceptualization , Writing – review and editing; Brit Salbu : Resources;
Funding acquisition, Writing – review and editing; Dag A Brede: Investigation,
Methodology, Writing - review & editing, Jorunn E Olsen: Conceptualization, Funding
acquisition, Investigation, Methodology, Project administration, Resources, Supervision,
Validation, Visualization, Writing - original draft, Writing - review & editing.

Data availability

The raw RNA sequence data are available in ArrayExpress under the accession number E-
MTAB-8081 for Norway spruce and E-MTAB-11120 for *A. thaliana.* Other data that support
the findings of this study are available within the article, in the Supporting Information of this
article and in online repositories. The names of the online repository and the link of the
identifiers will be available in the Norwegian University of Life Sciences archive once the
submission is approved.

References

Berardini, T. Z., L. Reiser, D. Li, Y. Mezheritsky, R. Muller, E. Strait, and E. Huala. 2015. "The
Arabidopsis Information Resource: Making and Mining The "Gold Standard" Annotated
Reference Plant Genome." *Genesis* 53, no. 8 (Aug): 474-85.
https://dx.doi.org/10.1002/dvg.22877.

Blagojevic, D., Y. Lee, D. A. Brede, O. C. Lind, I. Yakovlev, K. A. Solhaug, C. G. Fossdal, B. Salbu, and J.
E. Olsen. 2019. "Comparative Sensitivity to Gamma Radiation at the Organismal, Cell and
Dna Level in Young Plants of Norway Spruce, Scots Pine and Arabidopsis Thaliana." *Planta*
(Aug). https://dx.doi.org/10.1007/s00425-019-03250-y.

705
706
707 Bourbousse, C., N. Vegesna, and J. A. Law. 2018. "Sog1 Activator and Myb3r Repressors Regulate a
708   Complex Dna Damage Network In." *Proc Natl Acad Sci U S A* 115, no. 52 (12 26): E12453-
709   E12462. https://dx.doi.org/10.1073/pnas.1810582115.

710
711 Caplin, N., and N. Willey. 2018. "Ionizing Radiation, Higher Plants, and Radioprotection: From Acute
712   High Doses to Chronic Low Doses." *Front Plant Sci* 9: 847.
713   https://dx.doi.org/10.3389/fpls.2018.00847.

714
715 Choi, H. I., S. M. Han, Y. D. Jo, M. J. Hong, S. H. Kim, and J. B. Kim. 2021. "Effects of Acute and Chronic
716   Gamma Irradiation on the Cell Biology and Physiology of Rice Plants." *Plants (Basel)* 10, no. 3
717   (Feb 25). https://dx.doi.org/10.3390/plants10030439.

718
719 Choi, S. H., T. H. Ryu, J. I. Kim, S. Lee, S. S. Lee, and J. H. Kim. 2019. "Mutation in Ddm1 Inhibits the
720   Homology Directed Repair of Double Strand Breaks." *PLoS One* 14, no. 2: e0211878.
721   https://dx.doi.org/10.1371/journal.pone.0211878.

722
723 De Schutter, K., J. Joubès, T. Cools, A. Verkest, F. Corellou, E. Babiychuk, E. Van Der Schueren, T.
724   Beeckman, S. Kushnir, D. Inzé, and L. De Veylder. 2007. "Arabidopsis Wee1 Kinase Controls
725   Cell Cycle Arrest in Response to Activation of the Dna Integrity Checkpoint." *Plant Cell* 19,
726   no. 1 (Jan): 211-25. https://dx.doi.org/10.1105/tpc.106.045047.

De Veylder, L., G. T. Beemster, T. Beeckman, and D. Inzé. 2001. "Cks1at Overexpression in Arabidopsis Thaliana Inhibits Growth by Reducing Meristem Size and Inhibiting Cell-Cycle Progression." *Plant J* 25, no. 6 (Mar): 617-26. https://dx.doi.org/10.1046/j.1365-313x.2001.00996.x.

Dumanović, J., E. Nepovimova, M. Natić, K. Kuča, and V. Jaćević. 2020. "The Significance of Reactive Oxygen Species and Antioxidant Defense System in Plants: A Concise Overview." *Front Plant Sci* 11: 552969. https://dx.doi.org/10.3389/fpls.2020.552969.

Díaz-Tielas, C., E. Graña, M. J. Reigosa, and A. M. Sánchez-Moreiras. 2012. "The Role of Peroxidases on the Mode of Action of Chalcone in Arabidopsis Roots." *Plant Signal Behav* 7, no. 10 (Oct 01): 1274-6. https://dx.doi.org/10.4161/psb.21594.

Einset, J., and A. R. Collins. 2018. "Genome Size and Sensitivity to Dna Damage by X-Rays-Plant Comets Tell the Story." *Mutagenesis* 33, no. 1 (Feb 24): 49-51. https://dx.doi.org/10.1093/mutage/gex029.

Endo, M., Y. Ishikawa, K. Osakabe, S. Nakayama, H. Kaya, T. Araki, K. Shibahara, K. Abe, H. Ichikawa, L. Valentine, B. Hohn, and S. Toki. 2006. "Increased Frequency of Homologous Recombination and T-Dna Integration in Arabidopsis Caf-1 Mutants." *EMBO J* 25, no. 23 (Nov 29): 5579-90. https://dx.doi.org/10.1038/sj.emboj.7601434.

Falk, M., E. Lukásová, and S. Kozubek. 2008. "Chromatin Structure Influences the Sensitivity of Dna to Gamma-Radiation." *Biochim Biophys Acta* 1783, no. 12 (Dec): 2398-414. https://dx.doi.org/10.1016/j.bbamcr.2008.07.010.

Garnier-Laplace, J., S. Geras'kin, C. Della-Vedova, K. Beaugelin-Seiller, T. G. Hinton, A. Real, and A. Oudalova. 2013. "Are Radiosensitivity Data Derived from Natural Field Conditions Consistent with Data from Controlled Exposures? A Case Study of Chernobyl Wildlife Chronically Exposed to Low Dose Rates." *J Environ Radioact* 121 (Jul): 12-21. https://dx.doi.org/10.1016/j.jenvrad.2012.01.013.

Gichner, T. 2003. "Dna Damage Induced by Indirect and Direct Acting Mutagens in Catalase-Deficient Transgenic Tobacco. Cellular and Acellular Comet Assays." *Mutat Res* 535, no. 2 (Mar 03): 187-93. https://dx.doi.org/10.1016/s1383-5718(02)00320-0.

Hansen, E. L., O. C. Lind, D. H. Oughton, and B. Salbu. 2019. "A Framework for Exposure Characterization and Gamma Dosimetry at the Nmbu Figaro Irradiation Facility." *Int J Radiat Biol* 95, no. 1 (01): 82-89. https://dx.doi.org/10.1080/09553002.2018.1539878.

International Atomic Energy Agency, the Annual Report for 1992.

Jiang, H. W., M. J. Liu, I. C. Chen, C. H. Huang, L. Y. Chao, and H. L. Hsieh. 2010. "A Glutathione S-Transferase Regulated by Light and Hormones Participates in the Modulation of Arabidopsis

Seedling Development." *Plant Physiol* 154, no. 4 (Dec): 1646-58.
https://dx.doi.org/10.1104/pp.110.159152.

Kevei, Z., M. Baloban, O. Da Ines, H. Tiricz, A. Kroll, K. Regulski, P. Mergaert, and E. Kondorosi. 2011.
"Conserved Cdc20 Cell Cycle Functions Are Carried out by Two of the Five Isoforms in
Arabidopsis Thaliana." *PLoS One* 6, no. 6: e20618.
https://dx.doi.org/10.1371/journal.pone.0020618.

Kim, D. S., J. B. Kim, E. J. Goh, W. J. Kim, S. H. Kim, Y. W. Seo, C. S. Jang, and S. Y. Kang. 2011.
"Antioxidant Response of Arabidopsis Plants to Gamma Irradiation: Genome-Wide
Expression Profiling of the Ros Scavenging and Signal Transduction Pathways." *J Plant Physiol*
168, no. 16 (Nov 01): 1960-71. https://dx.doi.org/10.1016/j.jplph.2011.05.008.

Kim, J. H. 2019. "Chromatin Remodeling and Epigenetic Regulation in Plant Dna Damage Repair." *Int*
J Mol Sci 20, no. 17 (Aug). https://dx.doi.org/10.3390/ijms20174093.

Kim, J. H., K. Hwangbo, E. Lee, S. K. Dubey, M. S. Chung, B. Y. Chung, and S. Lee. 2021. "Application of
Gamma Ray-Responsive Genes for Transcriptome-Based Phytodosimetry in Rice." *Plants*
(Basel) 10, no. 5 (May 13). https://dx.doi.org/10.3390/plants10050968.

Kim, J. H., T. H. Ryu, S. S. Lee, S. Lee, and B. Y. Chung. 2019. "Ionizing Radiation Manifesting Dna
Damage Response in Plants: An Overview of Dna Damage Signaling and Repair Mechanisms
in Plants." *Plant Sci* 278 (Jan): 44-53. https://dx.doi.org/10.1016/j.plantsci.2018.10.013.

Kovalchuk, O., P. Burke, A. Arkhipov, N. Kuchma, S. J. James, I. Kovalchuk, and I. Pogribny. 2003.
"Genome Hypermethylation in Pinus Silvestris of Chernobyl--a Mechanism for Radiation
Adaptation?" *Mutat Res* 529, no. 1-2 (Aug 28): 13-20. https://dx.doi.org/10.1016/s0027-
5107(03)00103-9.

Kovács, E., and A. Keresztes. 2002. "Effect of Gamma and Uv-B/C Radiation on Plant Cells." *Micron*
33, no. 2: 199-210. https://dx.doi.org/10.1016/s0968-4328(01)00012-9.

Le Gall, H., F. Philippe, J. M. Domon, F. Gillet, J. Pelloux, and C. Rayon. 2015. "Cell Wall Metabolism in
Response to Abiotic Stress." *Plants (Basel)* 4, no. 1 (Feb 16): 112-66.
https://dx.doi.org/10.3390/plants4010112.

Lee, Y., C. Karunakaran, R. Lahlali, X. Liu, K. K. Tanino, and J. E. Olsen. 2017. "Photoperiodic
Regulation of Growth-Dormancy Cycling through Induction of Multiple Bud-Shoot Barriers
Preventing Water Transport into the Winter Buds of Norway Spruce." *Front Plant Sci* 8: 2109.
https://dx.doi.org/10.3389/fpls.2017.02109.

Lind, O. C., D. Helen Oughton, and B. Salbu. 2019. "The Nmbu Figaro Low Dose Irradiation Facility."
Int J Radiat Biol 95, no. 1 (01): 76-81. https://dx.doi.org/10.1080/09553002.2018.1516906.

Livak, K. J., and T. D. Schmittgen. 2001. "Analysis of Relative Gene Expression Data Using Real-Time
Quantitative Pcr and the 2(-Delta Delta C(T)) Method." *Methods* 25, no. 4 (Dec): 402-8.
https://dx.doi.org/10.1006/meth.2001.1262.

Mothersill, C., and C. B. Seymour. 1998. "Mechanisms and Implications of Genomic Instability and
Other Delayed Effects of Ionizing Radiation Exposure." *Mutagenesis* 13, no. 5 (Sep): 421-6.
https://dx.doi.org/10.1093/mutage/13.5.421.

Nisa, M. U., Y. Huang, M. Benhamed, and C. Raynaud. 2019. "The Plant Dna Damage Response:
Signaling Pathways Leading to Growth Inhibition and Putative Role in Response to Stress
Conditions." *Front Plant Sci* 10: 653. https://dx.doi.org/10.3389/fpls.2019.00653.

Nystedt, B., N. R. Street, A. Wetterbom, A. Zuccolo, Y. C. Lin, D. G. Scofield, F. Vezzi, N. Delhomme, S.
Giacomello, A. Alexeyenko, R. Vicedomini, K. Sahlin, E. Sherwood, M. Elfstrand, L. Gramzow,
K. Holmberg, J. Hällman, O. Keech, L. Klasson, M. Koriabine, M. Kucukoglu, M. Käller, J.
Luthman, F. Lysholm, T. Niittylä, A. Olson, N. Rilakovic, C. Ritland, J. A. Rosselló, J. Sena, T.
Svensson, C. Talavera-López, G. Theißen, H. Tuominen, K. Vanneste, Z. Q. Wu, B. Zhang, P.
Zerbe, L. Arvestad, R. Bhalerao, J. Bohlmann, J. Bousquet, R. Garcia Gil, T. R. Hvidsten, P. de
Jong, J. MacKay, M. Morgante, K. Ritland, B. Sundberg, S. L. Thompson, Y. Van de Peer, B.
Andersson, O. Nilsson, P. K. Ingvarsson, J. Lundeberg, and S. Jansson. 2013. "The Norway
Spruce Genome Sequence and Conifer Genome Evolution." *Nature* 497, no. 7451 (May):
579-84. https://dx.doi.org/10.1038/nature12211.

Patro, R., G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford. 2017. "Salmon Provides Fast and Bias-
Aware Quantification of Transcript Expression." *Nat Methods* 14, no. 4 (Apr): 417-419.
https://dx.doi.org/10.1038/nmeth.4197.

Pedroza-Garcia, J. A., Y. Xiang, and L. De Veylder. 2022. "Cell Cycle Checkpoint Control in Response
to Dna Damage by Environmental Stresses." *Plant J* 109, no. 3 (02): 490-507.
https://dx.doi.org/10.1111/tpj.15567.

Qi, F., and F. Zhang. 2019. "Cell Cycle Regulation in the Plant Response to Stress." *Front Plant Sci* 10:
1765. https://dx.doi.org/10.3389/fpls.2019.01765.

Sheppard, B.D. Amiro and S.C. 1994. *Effects of Ionizing Radiation on the Boreal Forest: Canada's Fig
Experiment, with Implications for Radionuclides*. Vol. 157: The Science of the Total
Environment.

Sidler, C., D. Li, O. Kovalchuk, and I. Kovalchuk. 2015. "Development-Dependent Expression of Dna
Repair Genes and Epigenetic Regulators in Arabidopsis Plants Exposed to Ionizing Radiation."
Radiat Res 183, no. 2 (Feb): 219-32. https://dx.doi.org/10.1667/RR13840.1.

Sylvestre-Gonon, E., S. R. Law, M. Schwartz, K. Robe, O. Keech, C. Didierjean, C. Dubos, N. Rouhier,
and A. Hecker. 2019. "Functional, Structural and Biochemical Features of Plant Serinyl-
Glutathione Transferases." *Front Plant Sci* 10: 608.
https://dx.doi.org/10.3389/fpls.2019.00608.

862
863 UNSCEAR. *Sources and Effects*

864 *of Ionizing Radiation*: United Nations Scientific Committee on the Effects of Atomic Radiation
865 UNSCEAR 1996 Report to the General Assembly, with Scientific Annexes.

866
867 van de Walle, J., N. Horemans, E. Saenen, M. Van Hees, J. Wannijn, R. Nauts, A. van Gompel, J.
868 Vangronsveld, H. Vandenhove, and A. Cuypers. 2016. "Arabidopsis Plants Exposed to
869 Gamma Radiation in Two Successive Generations Show a Different Oxidative Stress
870 Response." *J Environ Radioact* 165 (Dec): 270-279.
871 https://dx.doi.org/10.1016/j.jenvrad.2016.10.014.

872
873 Vanhoudt, N., N. Horemans, J. Wannijn, R. Nauts, M. Van Hees, and H. Vandenhove. 2014. "Primary
874 Stress Responses in Arabidopsis Thaliana Exposed to Gamma Radiation." *J Environ Radioact*
875 129 (Mar): 1-6. https://dx.doi.org/10.1016/j.jenvrad.2013.11.011.

876
877 Verma, V., P. Ravindran, and P. P. Kumar. 2016. "Plant Hormone-Mediated Regulation of Stress
878 Responses." *BMC Plant Biol* 16 (Apr 14): 86. https://dx.doi.org/10.1186/s12870-016-0771-y.

879
880 Vishwakarma, K., N. Upadhyay, N. Kumar, G. Yadav, J. Singh, R. K. Mishra, V. Kumar, R. Verma, R. G.
881 Upadhyay, M. Pandey, and S. Sharma. 2017. "Abscisic Acid Signaling and Abiotic Stress
882 Tolerance in Plants: A Review on Current Knowledge and Future Prospects." *Front Plant Sci*
883 8: 161. https://dx.doi.org/10.3389/fpls.2017.00161.

884
885 Weimer, A. K., S. Biedermann, H. Harashima, F. Roodbarkelari, N. Takahashi, J. Foreman, Y. Guan, G.
886 Pochon, M. Heese, D. Van Damme, K. Sugimoto, C. Koncz, P. Doerner, M. Umeda, and A.
887 Schnittger. 2016. "The Plant-Specific Cdkb1-Cycb1 Complex Mediates Homologous
888 Recombination Repair in Arabidopsis." *EMBO J* 35, no. 19 (10): 2068-2086.
889 https://dx.doi.org/10.15252/embj.201593083.

890
891 Xie, L., K. A. Solhaug, Y. Song, D. A. Brede, O. C. Lind, B. Salbu, and K. E. Tollefsen. 2019. "Modes of
892 Action and Adverse Effects of Gamma Radiation in an Aquatic Macrophyte Lemna Minor."
893 *Sci Total Environ* 680 (Aug 25): 23-34. https://dx.doi.org/10.1016/j.scitotenv.2019.05.016.

894
895 Yoshiyama, K. O., J. Kobayashi, N. Ogita, M. Ueda, S. Kimura, H. Maki, and M. Umeda. 2013. "Atm-
896 Mediated Phosphorylation of Sog1 Is Essential for the Dna Damage Response in
897 Arabidopsis." *EMBO Rep* 14, no. 9 (Sep): 817-22.
898 https://dx.doi.org/10.1038/embor.2013.112.

899
900 Zimin, A. V., K. A. Stevens, M. W. Crepeau, D. Puiu, J. L. Wegrzyn, J. A. Yorke, C. H. Langley, D. B.
901 Neale, and S. L. Salzberg. 2017. "An Improved Assembly of the Loblolly Pine Mega-Genome
902 Using Long-Read Single-Molecule Sequencing." *Gigascience* 6, no. 1 (01 01): 1-4.
903 https://dx.doi.org/10.1093/gigascience/giw016.

904
905

906 **Table 1.** The gamma radiation dose rates and total doses applied in experiments with 48 h

907 exposure of seedlings of *Arabidopsis thaliana* and Norway spruce from day 7 after sowing,

908 using a ^{60}Co source.

Dose rate (mGy h^{-1})	Dose rate interval (mGy h^{-1})		Total dose (Gy) after 48 h exposure	Total dose interval (Gy) after 48 h exposure)	
	Minimum	Maximum		Minimum	Maximum
290	255	325	13.9	12.2	15.6
100	88	112	4.8	4.3	5.4
40	35	45	1.9	1.69	2.15
10	8,8	11,2	0.48	0.41	0.55
1	0,88	1,12	0.048	0.041	0.055
0	-	-	0	-	-

		Dose rates (mGy h⁻¹)
A. thaliana	P. abies	0
		1
		10
		40
		100
		290

Fig. 2   *Bhattacharjee et al. 2024*

Fig. 3   *Bhattacharjee et al. 2024*

Fig. 4   *Bhattacharjee et al. 2024*

Fig. 5   *Bhattacharjee et al. 2025*

Fig. 6   *Bhattacharjee et al. 2024*

Fig. 7 *Bhattacharjee et al. 2024*