1 **Distinct roles of dopamine and acetylcholine in delay- and effort-based**
2 **decision making in humans**
3
4 Short title: Dopaminergic and cholinergic modulation of cost-benefit decision making
5
6

7 Mani Erfanian Abdoust[a], Monja Isabel Froböse[a], Alfons Schnitzler[b], Elisabeth
8 Schreivogel[b], Gerhard Jocham[a]
9
10
11 [a] Biological Psychology of Decision Making, Institute of Experimental Psychology, Heinrich Heine
12 University Düsseldorf, Düsseldorf, Germany
13 [b] Institute of Clinical Neuroscience and Medical Psychology, Medical Faculty, Heinrich Heine University
14 Düsseldorf, Germany
15
16
17 [1] correspondence to
18 Mani Erfanian Abdoust
19 erfanian@hhu.de

Abstract

In everyday life, we encounter situations that require tradeoffs between potential rewards and associated costs, such as time and (physical) effort. The literature indicates a prominent role for dopamine in discounting of both delay and effort, with mixed findings for delay discounting in humans. Moreover, the reciprocal antagonistic interaction between dopaminergic and cholinergic transmission in the striatum suggests a potential opponent role of acetylcholine in these processes. We found opposing effects of dopamine D2 (haloperidol) and acetylcholine M1 receptor (biperiden) antagonism on specific components of effort-based decision making in healthy humans: haloperidol decreased, whereas biperiden increased the willingness to exert physical effort. In contrast, delay discounting was reduced under haloperidol, but not affected by biperiden. Together, our data suggest that dopamine, acting at D2 receptors, modulates both effort and delay discounting, while acetylcholine, acting at M1 receptors, appears to exert a more specific influence on effort discounting only.

Introduction

Consider the following scenarios: Would you prefer to order an average pizza that can be delivered within minutes or rather wait an extra hour for your most favourite pizza? Similarly, would you visit a highly-rated pizza place that requires climbing up a steep hill or opting for a conveniently accessible pizza place right in front of the next bus stop? In scenarios like this, decision making involves the balancing of potential rewards against distinct costs required to obtain them. The tendency to devalue rewards as a function of effort or delay costs are commonly described as effort and delay discounting, respectively (Bickel & Marsch, 2001; Hartmann et al., 2013).

Striatal dopamine has been suggested to play a central role in both effort and delay discounting. For effort-based decision making, studies in both humans and rodents indicate that diminished dopamine transmission makes individuals less willing to exert physical effort in exchange for larger rewards (Denk et al., 2005; Farrar et al., 2010; Le Heron et al., 2018; McGuigan et al., 2019; Salamone & Correa, 2012; Soder et al., 2020; Wardle et al., 2011). Pharmacological studies consistently demonstrate that agents that enhance dopaminergic transmission in humans increase reward sensitivity and decrease effort sensitivity, while studies directly investigating the effects of selective D2-receptor antagonism in healthy humans are rare (Bogdanov et al., 2022; Chong et al., 2023; Le Bouc et al., 2016). For delay discounting, the existing literature is somewhat more mixed. Some studies suggest that increased dopamine transmission decreases delay discounting (de Wit et al., 2002; Floresco et al., 2008; Kayser et al., 2012), whereas others have found either no effect (Acheson & de Wit, 2008; Hamidovic et al., 2008) or an increase in delay discounting (Pine et al., 2010). Notably, more recent evidence in humans indicates decreased delay discounting after blockade of D2 receptors (Soutschek & Tobler, 2023; Wagner et al., 2020). While

dopamine has received much attention in cost-benefit decision making, other

neuromodulators may also play an important role (Borderies et al., 2020; Meyniel et

al., 2016; Nunes et al., 2013). One modulator that has not received much attention in

this regard is acetylcholine, which is surprising given the literature on reciprocal

antagonistic interactions between acetylcholine and dopamine signalling in the

striatum (Chantranupong et al., 2023; Foster et al., 2016; Gerber et al., 2001;

Myslivecek, 2021; Threlfell et al., 2010). This suggests that pharmacological blockade

of M1 receptors will have effects on cost-benefit decision making that are opposite to

those of blocking dopamine D2 receptors. In line with this, animal studies indicate that

muscarinic agonists induce behavioural changes in effort-based choices that are

similar to those produced by dopamine antagonists (Hailwood et al., 2019; Nunes et

al., 2013), underscoring the potential interplay between dopamine and acetylcholine

in modulating cost-benefit decision making. Furthermore, research in animal models

points to a role for acetylcholine in modulating delay-based choices through both

muscarinic and nicotinic receptors, albeit with inconsistent findings (Dallery & Locey,

2005; Kolokotroni et al., 2011; Mendez et al., 2012; Ozga & Anderson, 2018).

Importantly, studies directly investigating these effects in human decision making are

lacking.

To our knowledge, there has been no study so far that tested the impact of

dopaminergic and cholinergic manipulations on both aspects of cost-benefit decision

making in one single experiment. To fill this gap, we investigated the effects of two

drugs, haloperidol and biperiden, that selectively block either dopamine D2-like or

muscarinic M1 acetylcholine receptors in human participants performing two decision-

making tasks involving effort- and delay-based decisions. The goal of our study was

threefold. First, we aimed to conceptually replicate the finding that dopamine D2

antagonists increase discounting of physical effort. Second, we aimed to assess the

contribution of acetylcholine to cost-benefit decision making and conceptually contrast

it with the effects of dopamine. In doing so, we investigated whether any effects of

muscarinic or dopaminergic receptor antagonism would modulate the computation of

both time and effort costs or have a more specific effect on one cost dimension. Third,

we sought to contribute new evidence to the thus far conflicting literature on the role

of dopamine in delay-based decision making using a large sample size and a within-

subjects design. Based on previous findings, we hypothesized that the administration

of haloperidol will (1) increase effort discounting and (2) reduce delay discounting. In

contrast, we expected opposite effects of biperiden, in particular (3) a decrease in

effort discounting and (4) an increase in delay discounting. In brief, we found opposing

effects of haloperidol and biperiden only on specific components of effort-based

choices. Specifically, haloperidol reduced the willingness to invest physical effort,

whereas biperiden increased it. Results for delay discounting were less consistent.

While haloperidol decreased delay discounting, there was no credible modulation by

biperiden.

105   –   insert Figure 1 here –

Results

62 healthy participants performed two cost-benefit decision-making tasks aimed to

quantify the extent to which the subjective value of a monetary reward is discounted

as a function of either effort or delay costs. Participants completed both tasks during

three sessions under the influence of either the D2 receptor antagonist haloperidol (2

mg), the M1 acetylcholine receptor antagonist biperiden (4 mg), or a placebo in a

within-subjects design (Fig. 1a). The effort-based decision-making task (effort discounting task, Fig. 1b) involved choices between options varying in reward magnitude and effort requirement (using handgrip force), while the delay-based decision-making task (delay discounting task, Fig. 1c) involved choices between one option varying in reward magnitude and delay versus a fixed one. Thus, both tasks required participants to choose between a high-reward/high-cost (high-cost option) and a low-reward/low-cost (low-cost option) alternative.

Drug Effects on Choice Behaviour

Participants consistently exhibited a preference for the high-cost option over the low-cost alternative in both tasks (effort discounting task: 78.50% ± 1.03; delay discounting task: 71.31% ± 1.12; Fig. 2a). For each of the two tasks, we used a Logistic Bayesian Generalized Linear Mixed Model (GLMM) to investigate the impact of drug manipulation, changes in task parameters, and the interaction between both, on choosing the high-cost option. Specifically, to regress participants' choices, we included the following fixed-effect predictors: drug condition, reward magnitude, cost level (i.e., effort or delay), and all possible interactions. In the effort discounting task, we used the difference terms (high-cost option minus low-cost option) for both reward and effort. For the delay discounting task, we used the absolute reward and delay levels of the varying high-cost option. To account for within-subject variability, we included random intercepts for each subject along with random slopes for all fixed-effect predictors (see Materials and Methods). We first confirmed that participants effectively discounted rewards based on costs and thereby adhered to the task requirements: As expected, in both discounting tasks higher reward magnitudes increased, while higher cost levels (effort or delay) decreased participants' likelihood

138 to select the high-cost option (reward effect on effort discounting: HDI_{Mean} = 3.44,

139 $HDI_{95\%}$ = [2.96; 3.95]; reward effect on delay discounting: HDI_{Mean} = 55.64, $HDI_{95\%}$ =

140 [41.76; 69.38], effort effect on effort discounting: HDI_{Mean} = -1.64, $HDI_{95\%}$ = [-1.87; -

141 1.40]; delay effect on delay discounting: HDI_{Mean} = -2.29, $HDI_{95\%}$ = [-3.30; -1.25], Fig.

142 2b-c).

144   –   insert Figure 2 here –

Notably, and in line with our predictions, we found that the two drugs had opposite

effects on the tendency to choose high-effort options in the effort discounting task

relative to placebo. Haloperidol reduced the willingness to invest higher effort for

greater reward, while biperiden increased this willingness (placebo: 79.17% ± 1.76;

haloperidol: 74.34% ± 1.85; biperiden: 81.99% ± 1.64; Fig. 2a (left panel)). These

effects were confirmed by credible main effects of both haloperidol (HDI_{Mean} = -0.532,

$HDI_{95\%}$ = [-0.943; -0.136]) and biperiden (HDI_{Mean} = 0.620, $HDI_{95\%}$ = [0.230; 1.048]; Fig.

3a) on choosing the high-cost option relative to placebo. Moreover, this analysis

revealed that biperiden increased reward sensitivity, as evidenced by a credible

interaction effect between biperiden and reward (HDI_{Mean} = 0.802, $HDI_{95\%}$ = [0.256;

1.409]; Fig. 3b). See Supplementary Table 1 for the full results.

In contrast, in the delay discounting task, both drugs had no effect on the

average rate of choosing the high-cost option (placebo: 71.59% ± 2.25; haloperidol:

71.55% ± 1.73; biperiden: 70.79% ± 1.82; Fig. 2a (right panel)), as indicated by the

absence of credible main effects for haloperidol (HDI_{Mean} = -0.046, $HDI_{95\%}$ = [-2.648;

1.884]) and biperiden (HDI_{Mean} = -1.315, $HDI_{95\%}$ = [-4.147; 0.655]; Fig. 3d). Importantly,

however, the analysis revealed a reduced sensitivity to delays under haloperidol,

evidenced by a credible interaction effect between haloperidol and delay (HDI_{Mean} =

1.332, $HDI_{95\%}$ = [0.328; 2.440]; Fig. 3f), while the other drug interactions did not reach

credibility (Fig. 3 and Supplementary Table 2). This indicates that diminished

dopaminergic activity attenuates the impact of time costs on decision making.

168   –   insert Figure 3 here –

Having established distinct task-specific drug effects on participants' choice behaviour,

we next investigated potential confounding order and fatigue effects using separate

GLMMs for effort- and delay-based decision making. These models mirrored the

previously described GLMMs but included additional predictors to test for fatigue and

session effects and their modulation by drugs.

To test for possible fatigue effects, we added trial number as well as the two-

way interaction with drug as additional predictors. This analysis revealed a credible

main effect of trial number for both the effort and delay discounting task (trial number

effect on effort discounting: HDI_{Mean} = -0.60, $HDI_{95\%}$ = [-0.75; -0.46]; trial number effect

on delay discounting: HDI_{Mean} = -0.20, $HDI_{95\%}$ = [-0.33; -0.08]), suggesting that the

tendency to choose high-cost options, irrespective of cost type, decreased over the

course of the experiment. Notably, we found a credible interaction effect between trial

number and haloperidol in the effort discounting task (HDI_{Mean} = -0.43, $HDI_{95\%}$ = [-0.63;

-0.23]), indicating that the fatigue effect was more pronounced under haloperidol

compared to placebo. Importantly, even after accounting for these fatigue-related

effects, the main effect of haloperidol on effort discounting remained credible (HDI_{Mean}

= -0.48, $HDI_{95\%}$ = [-0.90; -0.05]) (Supplementary Table 12 and 13).

Next, to investigate session and drug-order effects, we included session and

the two-way interactions between session and drug as additional predictors. These

analyses confirmed a main effect of session only for effort-based choices (HDI_{Mean} =

0.49, $HDI_{95\%}$ = [0.13; 0.85]), suggesting increased preference for high-effort options in

later sessions, likely reflecting task familiarity. Crucially, the interaction effects

between either drug and session were not credible in both tasks (Supplementary Table

14 and 15). This indicates that neither dopaminergic nor cholinergic manipulations

modulated learning effects across sessions. Moreover, this finding underscores that

the observed drug-induced effects cannot be explained by drug order effects.

To summarize, regression-based results show that haloperidol reduced the

overall propensity to choose the high-cost options in the effort domain. While

haloperidol had no effect on the average rate of selecting the high-cost option in delay-

based choices, it attenuated participants' sensitivity to delay costs. In contrast,

biperiden increased the overall propensity of selecting the high-cost option in the effort

discounting task, opposite to the effect of haloperidol. Consistent with this, biperiden

also increased sensitivity to reward magnitudes during effort-based choices.

Drug Effects on Computational Parameters

To obtain a more detailed understanding of the mechanism underlying the drug effects

on behaviour described above, we used hierarchical Bayesian modelling. To begin

with, we determined the best-fitting discounting models for both effort and delay,

comparing four commonly used models (linear, parabolic, hyperbolic, exponential). In

line with previous findings (Green & Myerson, 2004; Hartmann et al., 2013; Kirby &

Maraković, 1995; Klein-Flügge et al., 2015; Lockwood et al., 2017), model

comparisons revealed that effort discounting was best described by a parabolic model,

suggesting a greater impact of changes in high rather than low effort levels. In contrast,

delay discounting was best described by a hyperbolic model, indicating a greater

impact of changes in low rather than high delay levels (Supplementary Table 3). For

each model, we first calculated the subjective values of all choice options, based on

participant-specific weighing of reward magnitude and associated costs (i.e., effort and

delay levels). By introducing condition-specific shift parameters, we captured potential

drug effects on the effort and delay discounting parameter κ (denoted as $s\kappa_{HAL}$ for

haloperidol and $s\kappa_{BIP}$ for biperiden), with positive/negative shift parameter values

indicating increased/decreased effort and delay discounting, respectively. We then

used a softmax function to transform the option values into choice probabilities, with

choice stochasticity being modelled by an inverse temperature parameter β. Again,

condition-specific shift parameters for haloperidol and biperiden captured potential

drug effects on choice stochasticity ($s\beta_{HAL}$ for haloperidol and $s\beta_{BIP}$ for biperiden), with

positive/negative shift parameters indicating more deterministic/more stochastic

decision making (see Materials and Methods and Supplementary Methods). Model

validation and parameter recovery confirmed that both models accurately captured

key features of the choice data (see Supplementary Results).

230   –   insert Figure 4 here –

The results from the computational model of the effort discounting task align with and

extend the regression-based results presented above. Specifically, again biperiden

and haloperidol exerted opposing effects on both the discounting and inverse

temperature parameter. Haloperidol increased effort discounting, while biperiden

diminished it. Similarly, haloperidol induced more stochastic choices, while biperiden

led to more deterministic decisions (Table 1). For both drug-specific effects on the

effort discounting parameter κ, we acknowledge that, strictly speaking, the 95% HDI

of $s\kappa_{BIP}$ (HDI$_{Mean}$ = -0.012,; HDI$_{95\%}$ = [-0.000; 0.027]) and $s\kappa_{HAL}$ (HDI$_{Mean}$ = 0.013, HDI$_{95\%}$

= [-0.025; 0.000]) did overlap with zero, albeit to a very small extent (Fig. 4a). Notably,

the density that did not overlap with zero still accounted for more than 94% of the

posterior distribution (94.8% HDI > 0 for haloperidol and 94.4% HDI > 0 for biperiden).

This provides strong evidence for a credible modulation of effort discounting by

cholinergic M1 and dopaminergic D2 receptor manipulation, despite the slight overlap.

A similar pattern emerged in the analysis of the inverse temperature parameter,

further supporting the partly opposing effects of these neurotransmitters. The HDIs for

$s\beta_{HAL}$ (HDI$_{Mean}$ = -0.089, HDI$_{95\%}$ = [-0.164; -0.010]) and $s\beta_{BIP}$ (HDI$_{Mean}$ = 0.127, HDI$_{95\%}$ =

[0.048; 0.219]) did not overlap with zero, supporting the modulatory effects of these

drugs on choice stochasticity. Specifically, haloperidol increased choice stochasticity,

while biperiden exhibited the opposite effect (Fig. 4b).

252   --insert Table 1 here--

For the delay discounting task, computational modelling revealed credible evidence

only for a dopaminergic, but not cholinergic modulation of delay discounting (Fig. 4d

and Table 2). In line with the diminished delay sensitivity reported above, haloperidol

reduced delay discounting, making participants more willing to wait for greater financial

rewards with increasing levels of delay ($s\kappa_{HAL}$: HDI$_{Mean}$ = -0.630, HDI$_{95\%}$ = [-1.083; -

0.237]). For biperiden, we did not observe any shift of the discounting parameter ($s\kappa_{BIP}$:

HDI_{Mean} = -0.125, $HDI_{95\%}$ = [-0.410; 0.148]). Next, we examined the effects of

dopaminergic and muscarinic antagonism on choice stochasticity (Fig. 4e). Biperiden

credibly increased the choice stochasticity ($s\beta_{BIP}$: HDI_{Mean} = -0.043, $HDI_{95\%}$ = [-0.085; -

0.000]), while we did not find credible evidence for a modulatory effect of haloperidol,

as the 95% HDI overlapped with zero ($s\beta_{HAL}$: HDI_{Mean} = -0.055, $HDI_{95\%}$ = [-0.114;

0.006]).

267   --insert Table 2 here--

Lastly, to test whether putatively confounding side effects of the pharmacological

treatment account for our main effects of interest, we tested medication effects on

several control measures, including mood (alertness, calmness, and contentedness)

and basic physiological parameters (heart rate, systolic, and diastolic blood pressure),

using Bayesian Linear Mixed Models. In short, biperiden administration induced

reductions in systolic blood pressure and heart rate at T_1 and at T_2. Moreover, relative

to placebo, both biperiden and haloperidol caused decreases in subjective alertness

ratings at T_2 (all $HDI_{95\%}$ < 0). T_1 represents the measurement taken before participants

began the effort discounting task, and T_2 represents the measurement taken after they

completed the delay discounting task. Importantly, neither the changes in blood

pressure or heart rate, nor reductions in alertness showed credible correlations with

any shift parameter that was credibly modulated by haloperidol or biperiden in both

tasks. This implies that the drug-induced changes in behaviour were not linked to drug

effects on alertness, heart rate, or blood pressure. More detailed information can be

found in the Supplementary Results.

Taken together, the results derived from hierarchical Bayesian modelling are

consistent with findings obtained from the GLMMs. Diminishing dopaminergic D2

receptor activity decreases the willingness to invest effort for rewards, possibly by

amplifying effort discounting. Conversely, the administration of a cholinergic M1

receptor antagonist produces opposite effects. However, this pattern of opponent

dopaminergic and cholinergic effects on specific components of effort discounting was

not present in the delay discounting task. Instead, the findings suggest a reduced

impact of delay on decision making under haloperidol, while biperiden affected only

the choice stochasticity. This implies that contrasting effects of dopaminergic and

cholinergic manipulations may reflect computationally-specific, rather than universal

effects.

Across-Task Relationship of Discounting Behaviour

Lastly, as an exploratory analysis, we asked whether there was a relationship between

discounting of delays versus efforts across individuals. In other words, we tested

whether people showing steep discounting of physical efforts would also show steep

delay-based discounting. In line with previous findings (Klein-Flügge et al., 2015;

Seaman et al., 2018), we found no relationship between individuals' tendency to

discount rewards based on efforts compared to delays (no credible correlation

between the effort and delay discounting parameters κ under placebo; r = -0.17,

$HDI_{95\%}$ = [-0.06; 0.39]).

Discussion

In this study, we investigated the effects of pharmacologically manipulating

dopaminergic and cholinergic neurotransmission on cost-benefit decision making in

healthy young adults. Specifically, we administered haloperidol or biperiden, two drugs

that selectively block either dopamine D2-like or muscarinic M1 acetylcholine

receptors, respectively, and tested the effects on discounting a monetary reward as a

function of either effort or delay. In short, we found that reducing dopaminergic

transmission at D2-like receptors decreased participants' willingness to invest physical

effort for monetary rewards and attenuated the impact of delay on decision making. In

contrast, cholinergic M1 receptor manipulations promoted the preference for high-

effort options, without affecting delay-based choices.

The goal of our study was threefold. First, we sought to conceptually replicate

the boosting effects of dopamine on motivation. Second, we examined the contribution

of acetylcholine, acting at M1 receptors, on cost-benefit decision making, as the

influence of muscarinic acetylcholine receptors on human (value-based) decision

making has rarely been reported. Third, we aimed to address the conflicting literature

on dopamine's role in delay-based decision making by providing data from a large

sample using a within-subjects design.

Dopamine has been widely suggested to play a key role in various aspects of

reward processing and cost-benefit decision making (Jocham et al., 2011, 2014;

Webber et al., 2020). It modulates choices based on expected rewards and costs

associated with different options (Schultz, 2016a, 2016b; Tanaka et al., 2019).

However, the direction of dopaminergic manipulations on different aspects of costs

remains ambiguous. Traditionally, dopamine has been implicated in promoting

behaviour that maximizes reward outcomes, suggesting that increased dopaminergic

activity energizes behaviour to approach high-reward options despite associated costs

(Bailey et al., 2016; Berke, 2018; Phillips et al., 2007; Salamone & Correa, 2012).

According to this view, increasing dopaminergic transmission should bias decision

making toward high-reward options, even when they involve higher costs such as effort, delay, and risk. Conversely, decreasing dopaminergic transmission should have the opposite effect.

In line with this view, in our study, the administration of a D2-like receptor antagonist did in fact reduce the willingness to invest physical effort for reward. However, contrary to the notion that dopamine supports reward-maximizing behaviour, our findings in the delay discounting task did not confirm the expected pattern. We observed a decrease in delay discounting under haloperidol, rather than an increase. While there is indeed a substantial body of evidence suggesting enhancing effects of dopamine on motivation, as demonstrated by studies highlighting its involvement in promoting reward-seeking behaviour (Michely et al., 2020; Soder et al., 2020; Soutschek et al., 2020; Wardle et al., 2011), the impact of pharmacological manipulations targeting other cost-related factors, such as delay (de Wit et al., 2002; Hamidovic et al., 2008; Kayser et al., 2012; Petzold et al., 2019) and risk (Burke et al., 2018; Rutledge et al., 2015; Zack & Poulos, 2007), has produced rather inconsistent findings. These studies showed either contradicting results regarding dopaminergic manipulations or no effect at all. These inconsistencies challenge traditional views of dopamine and raise questions about the specific mechanisms through which dopamine modulates choices that incorporate time-dependent costs. A more recent theory proposes that dopamine, acting at D2-like receptors, biases action selection by increasing the preference for options with a proximity advantage over more distant alternatives (Soutschek et al., 2023; Westbrook & Frank, 2018). This theory is based on studies in rodents, where firing rates of neurons in the nucleus accumbens are increased by spatially proximal rewards, promoting a decision bias towards nearby low-reward options (Morrison & Nicola, 2014). This notion of proximity, originally

referring to spatial proximity, has been extended to the context of psychological

proximity (Westbrook & Frank, 2018). According to this view, dopamine not only

favours options that are physically closer in space, but also options that are

psychologically closer, as in our case, rewards that are available sooner in time.

Consequently, the administration of D2 receptor antagonists, which, according to this

theory, are believed to reduce the proximity bias, have been shown in some studies

to increase the preference for delayed and risky reward options, which lack the

proximity advantage (Burke et al., 2018; Ojala et al., 2018; Soutschek & Tobler, 2023;

Wagner et al., 2020). In other words, this theoretical framework proposes that

dopamine may play a dual role in (i) promoting choices towards options with a

psychological proximity advantage and (ii) weighing reward magnitudes against

associated costs. Our findings of decreased delay discounting under haloperidol align

with this theory, potentially suggesting a diminished preference for more proximal

options when D2-receptor activity is reduced, resulting in increased endurance for

higher (time) costs. On the other hand, the reduction of dopaminergic

neurotransmission had the opposite effect on effort discounting, leading to a

diminished willingness to bear higher (effort) costs. This apparently conflicting finding

between effort and time costs may be explained by the fact that neither the low-effort

nor the high-effort option had a pronounced psychological proximity advantage. Thus,

in this specific context of cost-benefit decision making, the proximity factor appears to

be less relevant, and dopamine instead may promote behaviour aimed at maximizing

rewards simply by weighing rewards against associated costs. Importantly, our

experimental paradigm did not directly manipulate psychological proximity, nor did it

distinguish between proximity effects and the impact of delay. Therefore, specific

effects of dopamine on psychological proximity in human delay discounting remain

highly speculative and warrant further investigation with experimental paradigms

specifically tailored to this question.

Low doses of dopamine receptor antagonists may facilitate dopaminergic

transmission by primarily blocking presynaptic autoreceptors, which may increase

rather than decrease dopamine release (Frank & O'Reilly, 2006; Schoemaker et al.,

1997). We would however argue that such a presynaptic mechanism of action is

unlikely to explain our findings. First, a PET study found that the same dose of

haloperidol as used here led to high levels of D2 receptor occupancy in the striatum

(Kapur et al., 1996). Second, our pattern of results aligns with studies in rodents

showing motivational deficits following ventral striatal dopamine depletion (Cousins et

al., 1996; Mai et al., 2012; Salamone et al., 1994). Third, high doses of haloperidol

have been shown to reduce alertness (Liem-Moolenaar et al., 2010), whereas drugs

like methylphenidate that increase synaptic dopamine levels, enhance subjective

ratings of alertness (Linssen et al., 2011; Swart et al., 2017). Consistent with this, we

observed a reduction in alertness ratings following haloperidol administration.

Together, this suggests that our results are best explained by a blockade of

postsynaptic D2 receptors by haloperidol.

Notably, prior research has highlighted the contribution of other

neurotransmitters such as serotonin (Meyniel et al., 2016; Miyazaki et al., 2020),

adenosine (Randall et al., 2011; Salamone et al., 2018), and acetylcholine (Betts et

al., 2021; Hailwood et al., 2019; Nunes et al., 2013), in the weighing of costs and

rewards. However, the exact role of these neurotransmitters in reward processing and

cost-benefit decision making in humans has been rarely investigated. Understanding

the involvement of acetylcholine, due to its reciprocal activity with dopamine in the

striatum, is of particular interest (Chantranupong et al., 2023; Gerber et al., 2001;

Moehle & Conn, 2019; Myslivecek, 2021). At the functional level, these mutual

interactions are evident from the fact that muscarinic receptor antagonist diminish the

extrapyramidal side effects of dopamine antagonists, and, vice versa, muscarinic

agonists display antipsychotic properties, resembling the action of D2 antagonists

(Brocks, 1999; Stanhope et al., 2001). Additionally, blocking M1 receptors has been

shown to reverse motivational impairments induced by dopaminergic antagonism,

further emphasizing its potential role in modulating dopamine-related processes

(Hailwood et al., 2019). At the cellular level, it has been shown that M1 receptor

activation inhibits D2-receptor-mediated effects in the striatum (Di Chiara et al., 1994).

These observations highlight an interplay between acetylcholine and dopamine

signalling. Therefore, in addition to testing dopaminergic D2-like receptor

manipulations, we also investigated the role of cholinergic M1 receptor

neurotransmission on decision making.

Indeed, we observed opposing effects of dopaminergic and cholinergic

manipulations within specific components of effort-, but not delay-based decision

making. Specifically, we found evidence for biperiden increasing the willingness to

invest effort for rewards and, in line with this, decreasing effort discounting. In contrast,

haloperidol reduced the general willingness to invest effort and increased effort

discounting. However, we also observed drug effects that were not in opposite

directions between biperiden and haloperidol. For example, biperiden increased

reward sensitivity, while haloperidol had no credible effect on the impact of rewards

on effort-based choices. Furthermore, haloperidol modulated decision times in both

experimental tasks, whereas biperiden did not affect them. These findings suggest

partially opposing effects between both neurotransmitters, mostly evident within

specific components of effort-based choices.

This partially opposing mechanistic relationship between dopaminergic and

cholinergic neurotransmission during the effort discounting task is further reflected in

changes in choice stochasticity. Previous studies have linked D2 receptor antagonism

with increased choice stochasticity (Eisenegger et al., 2014; Mikus et al., 2022).

Consistent with these findings, our results revealed that haloperidol administration

increased stochasticity and thereby reduced value dependency on choices, while

biperiden had the opposite effect. After lowering cholinergic M1 receptor activity,

participants were more likely to choose the option with the highest expected value

compared to placebo. However, it is important to note that biperiden had the opposite

effect in the delay discounting task, increasing, rather than reducing choice

stochasticity. This discrepancy indicates that the role of cholinergic neurotransmission

in balancing deterministic versus stochastic behaviour is more complex and needs

further investigation.

Some limitations should be noted. First, our focus was primarily on striatal D2

receptor activity, as haloperidol predominantly targets D2 receptors in the striatum

(Nordström et al., 1992). However, it is important to note that in the context of cost-

benefit decision making, dissociable roles of D1 versus D2 receptor activity have been

reported (Collins & Frank, 2014; Soutschek et al., 2023; Webber et al., 2020), and thus

the general role of dopamine beyond its selective activity on striatal D2 receptors

remains unclear. Conversely, biperiden primarily targets M1 receptors in the cortex

and striatum (Sudo et al., 1999), making it challenging to determine the precise

mechanisms underlying cortical and striatal cholinergic modulation and the reciprocal

effects on dopaminergic activity. Second, previous research has suggested a U-

shaped dose-response function for dopamine, indicating deleterious effects of both

extremely high and extremely low levels of dopamine (Cools & D'Esposito, 2011;

Goldman-Rakic et al., 2000). According to this idea, the same dopamine agent can produce opposing effects in different individuals. Therefore, it may be insightful to consider individual differences in baseline dopamine levels when studying the effects of dopaminergic manipulations. A recent study found evidence for the absence of a correlation between dopamine synthesis capacity and putative behavioural proxies of dopamine, such as working memory or trait impulsivity (van den Bosch et al., 2022). Consequently, investigating baseline dopamine levels require more costly and invasive techniques, such as positron emission tomography. Third, as participants were required to exert physical effort on each trial, fatigue effects could have developed in the effort discounting task. While our additional analysis confirmed a general fatigue effect and a dual role of haloperidol in both reducing the overall propensity to invest effort and in exacerbating the fatigue effect throughout the task, we acknowledge that recent studies revealed the existence of distinct states of fatigue (Matthews et al., 2023; Müller et al., 2021). Importantly, this was discovered by using task paradigms and computational models that were designed to distinguish between these different types of fatigue. Additionally, it is important to consider that potential motor effects, particularly in the context of dopaminergic manipulations, could also affect effort discounting behaviour. Future studies could extend our approach by incorporating measurements of force pulses to investigate potential motor-related effects and apply task designs and computational models specifically tailored to capture how drug manipulations might affect different forms of fatigue. Lastly, it is important to note that several factors preclude a direct quantitative comparison between the delay and effort discounting tasks. These include the distinct nature of rewards and costs (hypothetical vs. real), varying reward magnitudes in both tasks (large vs. small), and differences in task structure (fixed vs. variable alternative option).

Additionally, both tasks were consistently performed in a fixed order within and

between participants (effort followed by delay). While this ensured consistent drug

levels within each task, it is potentially introducing task order effects and possibly

leading to differential drug concentrations between tasks. These methodological

differences limit our ability to directly compare drug effects across the two cost

domains. Future research would benefit from applying experimental paradigms that

manipulate both delay and effort costs within the same task, allowing a more controlled

and direct comparison of how pharmacological manipulations differentially influence

sensitivity to these distinct cost types.

In conclusion, our findings support prior research indicating an invigorating

effect of dopamine on motivation in an effort-based decision-making task. Moreover,

our study contributed to understanding dopamine's involvement in temporal cost-

benefit tradeoffs by revealing decreased delay discounting following dopaminergic D2-

like receptor antagonism. Further, we demonstrate that the administration of biperiden,

a muscarinic M1 receptor antagonist, had contrasting effects to those of dopaminergic

D2 receptor antagonism in the general willingness to choose high-cost options and the

effort discounting parameter. This suggests that in the context of human cost-benefit

decision making, the previously reported reciprocal relationship between both

neurotransmitters may be limited only to specific components of behaviour. Our

findings indicate that, while D2 receptor activity plays a role in integrating both delay

and effort costs, acetylcholine, acting at M1 receptors, may have a more specific role

in effort processing.

Materials and Methods

Ethics

Ethical approval for the study (2021-1549) was obtained from the Ethics Committee of the Medical Faculty of the University of Düsseldorf, Germany. Prior to participating, all volunteers provided informed written consent.

Participants

Participants were recruited for the study via online advertisements and university postings. They were initially screened via email interviews to verify compliance with the inclusion and exclusion criteria. Inclusion criteria included an age range of 18 to 35 years, a body weight ranging from 60 to 90 kg, and a body mass index equal to or greater than 18 and less than 28. Exclusion criteria included a history of psychiatric or neurological illnesses, current intake of prescription medication (excluding oral contraceptives), current pregnancy or breastfeeding, and the presence of any medical conditions contraindicated for the drugs used in the study. Additionally, participants with a history of drug use were excluded, with exceptions for alcohol, nicotine, and cannabis, which were limited to consumption of less than 14 units of alcohol per week, less than 5 cigarettes per day, and no cannabis consumption within the past month.

From an initial group of 96 individuals attending the medical screening on site, 33 candidates decided not to proceed, or they were excluded due to not meeting the participation criteria (despite otherwise indicated in the prior interview). Eventually, a total of 63 volunteers were enrolled in the study. One participant had to be excluded due to experiencing side effects following the biperiden testing session, resulting in a final sample size of 62 participants (32 female; Mean Age (SD): 22.79 (3.20); Age

Range: 18 - 35). The sample did not include any non-binary participants. All

participants received a fixed reimbursement of 240 € for their participation in all testing

sessions and were provided with a flexible payment based on their performance in the

tasks.

Procedure

A double-blind, randomized, within-subject design was employed. Participation

involved three testing sessions, each separated by a minimum of one week to ensure

complete drug washout. Prior to participation, on a separate day, participants

underwent a medical screening session to determine their eligibility. This screening

session consisted of a clinical interview, medical assessment, practice trials of both

tasks, and completion of several questionnaires assessing personality traits, including

Apathy Evaluation Scale (AES) (Lueken et al., 2006; Marin et al., 1991), Beck's

Depression Inventory (BDI) (Beck et al., 1996; Kühner et al., 2007), and Barratt

Impulsiveness Scale (BIS-15) (Meule et al., 2011; Spinella, 2007). The final decision

regarding participant suitability for the study was made by a physician at the end of

the screening session.

Each testing day followed the same procedure, with the only difference being

the administration of a placebo, biperiden (4 mg), or haloperidol (2 mg) in separate

sessions. Testing sessions were scheduled between 9 a.m. and 12 p.m., with efforts

made to assign participants to the same time slot consistently. Participants were

instructed to fast overnight and consume only water before the testing session.

Additionally, female participants completed a pregnancy test before the start of each

testing day. Upon arrival, participants underwent a screening conducted by a

physician, who restated the potential effects of the drug and provided a recap of the

testing day's procedures. Subsequently, participants received the assigned drug treatment along with a standardized breakfast. Orally administered biperiden reaches its peak concentration between 1 and 1.5 hours after administration (Brocks, 1999; Grimaldi et al., 1986), while haloperidol reaches its peak concentration only between 2 and 6 hours after administration (Kudo & Ishizaki, 1999). In order to follow a double-blind procedure while also accounting for variations in the drugs' peak times, a dummy drug application was included. Thus, on each testing day, a second capsule was administered 120 minutes after the first drug application, resulting in the planned delay between drug and effort discounting task of approximately 180 minutes for haloperidol (179.57 ± 2.61) and, on a separate day, approximately 60 minutes (59.65 ± 1.30) for biperiden. The second task was performed 205.59 minutes (± 3.12) after haloperidol and 86.24 minutes (± 6.09) after biperiden administration. Shortly before starting the experimental task, participants further completed the trail-making test A (Tombaugh, 2004). Blood pressure and heart rate were measured three times throughout the testing day: at T_0 (before drug administration), T_1 (before starting the effort discounting task), and T_2 (after finishing the delay discounting task). These measurements coincided with mood assessments using Visual Analogue Scales (Bond & Lader, 1974). Please refer to Fig. 1a for a detailed explanation of the procedure. The tasks were presented in a fixed order during each session, starting with the effort discounting task (Fig. 1b) and concluding with the delay discounting task (Fig. 1c). These discounting paradigms were implemented using Visual Basic software and displayed on a 15.6" laptop screen (Dell Latitude E5550). Participants used an external keypad to indicate their choices, arranging the laptop and keypad in a manner that was most comfortable for them.

Experimental Tasks

Effort Discounting Task

Participants engaged in a modified version of the Apple Tree Task, which has been previously utilized with both healthy volunteers and patients with neurological diseases (Bonnelle et al., 2015; Chong et al., 2018; Le Heron et al., 2018). Unlike the original task where participants accepted or rejected offers, in this modified task, they were presented with two alternative options on each trial. Prior to each experimental session, the participants' maximum voluntary contraction (MVC) was assessed by having them grip a handheld dynamometer (Vernier, Orlando, USA) with their dominant hand as forcefully as possible. The MVC was determined immediately before starting the task by measuring the highest force exerted over three contractions.

During each trial, they had to choose between a high-reward/high-effort (high-cost) and a low-reward/low-effort (low-cost) offer. Participants had unlimited time for their decision. Notably, both the reward levels and effort levels of each option were varied among five possible levels. The reward levels ranged from 2 to 16 (2, 4, 8, 12, and 16), and the effort levels ranged from no effort to 80% (0, 20, 40, 60, and 80%) of the individually determined MVC. After making a choice, participants were given a 5-second window to squeeze the handheld dynamometer and reach the required effort level. They had to maintain the required force for at least 1 second. Throughout this effort production period, a bar visually represented the force exerted, providing real-time feedback. Following successful trials, participants received feedback on the reward earned during the trial. If participants failed to reach the designated effort level, no apples were gathered. In cases where participants chose an offer that required no effort, they had to wait for the same duration without engaging in squeezing. At the end of the experiment, we asked participants to rate the perceived level of demand for

each effort level using a Likert scale ranging from 0 ("not demanding at all") to 20

("extremely demanding"). This subjective rating allowed us to examine whether the

administration of the drugs had any effects on participants' perception of the demand

associated with each effort level.

Each participant completed a total of 125 trials, divided into 5 blocks. The trial

structure was full randomized for each participant and experimental session. To

prevent strategic behaviour and mitigate delay discounting effects, all blocks and trials

were the same length, regardless of previous choices made. Importantly, patients

were required to squeeze the handheld dynamometer after every trial (if effort

production was chosen), meaning that no hypothetical choices were made. Based on

their task performance, participants received a flexible payment consisting of 1 cent

for each apple they collected.

Delay Discounting Task

After completing the effort discounting task, participants proceeded to the delay-based

decision-making task. In this task, they were presented with a similar choice paradigm,

but this time between a high-reward/high-delay (high-cost) and a low-reward/low-delay

(low-cost) option on each trial, again without any time limit. The tasks shared a

common structure, with participants consistently facing decisions involving a tradeoff

between a more favourable outcome with higher associated costs and a less

favourable outcome with fewer costs. In contrast to the previous task, the low-cost

option was fixed at 20 € and available immediately across all trials. The high-cost

option varied between 20.20 € and 260 € (20.20, 20.40, 21, 22, 24, 30, 36, 40, 50, 60,

80, 100, 140, 200, 260 €), with associated delays ranging from 1 day to 60 days (1, 2,

3, 5, 8, 30, 60). This resulted in a total of 105 unique high-cost combinations.

Participants completed two blocks, with each block containing all unique

combinations, resulting in a total of 210 trials. We used a pseudorandomized trial

order, counterbalanced across experimental sessions but not across participants.

Consistent with previous studies, all choices in this task were hypothetical (Bickel et

al., 2009; Madden et al., 2003). However, participants were instructed to imagine that

one of their choices would be randomly selected and paid out.

Statistical Analysis

Regression-Based Analysis

To investigate the influence of drug administration (compared to placebo) and

experimental manipulations (i.e., reward magnitude, delay, and/or effort) on choice

behaviour, we employed Logistic Bayesian Generalized Linear Mixed Models, using

the brms package (Bürkner, 2017) in R (Version 4.1.3). In these models, we regressed

choices (choosing the high-cost option vs. choosing the low-cost option) on fixed-effect

predictors including drug, reward, cost (i.e., delay or effort), and all possible interaction

terms. Importantly, to test drug-specific effects on behaviour, we included the drug

condition (placebo, haloperidol, biperiden) as a 3-level categorical predictor with

placebo set as the reference. Thus, all drug-related main and interaction effects are

estimated in reference to the placebo. For the effort discounting task, as predictors we

used the difference term between both the reward magnitude and effort levels of the

two presented options (high-cost option – low-cost option). In the delay discounting

task, the reward and delay levels of the varying high-cost option were included as

regressors. To account for individual differences, all fixed-effect predictors were also

modelled as random slopes in addition to subject-specific random intercepts. To

ensure robust and informative analyses, we followed the approach suggested by

Gelman et al. (2008). Weakly informative priors were employed, with nonbinary

variables scaled to have a mean of 0 and a standard deviation of 0.5. Posterior

distributions of the parameter estimates were obtained by running four chains with

3000 samples (1000 samples for warmup). We present the 95% Highest Density

Intervals (HDI) of the estimates to capture the uncertainty in the parameter estimates.

The 95% HDI indicates that there is a 95% probability that the true parameter value

falls within this interval. If the 95% HDI does not overlap with zero, it provides credible

evidence that the respective model parameter is meaningful (Ahn et al., 2017;

Kruschke, 2014). For detailed information on the models employed in this analysis,

please refer to the Supplementary Methods.

Order and Fatigue Control Analysis

To further test for potential confounding effects on choice behaviour, we ran separate

control regression analyses. For these models, we used the structure of the original

GLMMs but included additional fixed-effect predictors to control for session and fatigue

effects. Specifically, we included trial number as a predictor to account for potential

systematic changes in choice behaviour throughout the task progress. In separate

models, session was included as a fixed effect to control for potential learning or drug

order effects across different experimental sessions. Importantly, to capture potential

interactions between these control variables and the drug manipulations, we included

the two-way interaction terms in the fixed-effects structure. This allows to investigate

whether the effects of the drug manipulation on choice behaviour differed as a function

of the task progress or experimental session. All other aspects of the model

specification, including random-effects structure, priors, and estimation procedures,

remained identical to the original regression models described above.

Hierarchical Bayesian Modelling

Next, to investigate the impact of dopaminergic and cholinergic manipulations on underlying cognitive processes, specifically how individuals integrate rewards and costs to discount subjective reward values, we employed hierarchical Bayesian modelling. This approach allowed us to estimate both group-level hyperparameters and individual subject-level estimates, leveraging the hierarchical structure of the experimental design. By incorporating information from each individual's estimates into the group estimates and vice versa, we obtained more robust and reliable parameter estimates compared to conventional methods like maximum likelihood estimation (Ahn et al., 2017).

To capture how changes in the subjective values are influenced by reward, effort, and delay, we employed a single-parameter discounting model to determine which discounting function best describes the observed behaviour. Initially, participants' responses from the placebo condition were fitted to four commonly used models for discounting: linear, parabolic, hyperbolic, and exponential (Białaszek et al., 2017; Chong et al., 2017; Hartmann et al., 2013; Klein-Flügge et al., 2015). Model fits were compared using the trial-based Leave-One-Out Information Criterion (LOOIC) from the loo package (Vehtari et al., 2017).

Consistent with previous findings, delay discounting was best described by a hyperbolic model (Green & Myerson, 2004; Kirby & Maraković, 1995), whereas effort discounting showed the best fit with a parabolic model (Hartmann et al., 2013; Klein-Flügge et al., 2015; Lockwood et al., 2017). See Supplementary Methods for more details. The discounting parameter in the delay discounting model was modelled in log space due to the skewed distribution of the values towards zero.

$$SV(HC_{(t)}) = \frac{R_{(t)}}{1+\exp(\kappa)* D_{(t)}} \tag{1}$$

$$SV(HC_{(t)}) = R_{HC_{(t)}} - \kappa * E_{HC_{(t)}}^{2} \tag{2}$$

$$SV(LC_{(t)}) = R_{LC_{(t)}} - \kappa * E_{LC_{(t)}}^{2} \tag{3}$$

In the delay discounting model (Eq. 1), R_t represents the reward magnitude and D_t

represents the delay in days of the high-cost option (HC) on trial t, while the subjective

value (SV) of the low-cost option (LC) was fixed at 20 for each trial. In contrast, the

effort discounting models (Eq. 2 and 3) involve varying levels of both options. In this

model, the SV of the HC option and the SV of the LC option are calculated separately.

Once again, R_t denotes the corresponding reward magnitude, and E_t represents the

associated effort in percentage on trial t.

$$\kappa = \kappa_{PLC} + I_{HAL_{(t)}} * s_{\kappa_{HAL}} + I_{BIP_{(t)}} * s_{\kappa_{BIP}} \tag{4}$$

Consistent with previous studies that examined drug effects on changes in discounting

(Mathar et al., 2022; Peters & D'Esposito, 2020; Wagner et al., 2020), we extended

the original single-parameter model by incorporating two additional free parameters

(Eq. 4). In both cases, κ is the discounting parameter, either reflecting delay

discounting or effort discounting. A higher κ value indicates a greater degree of

discounting, whereas a lower κ value suggests less discounting. To capture potential

drug effects (compared to placebo), two separate shift parameters ($s_{\kappa_{HAL}}$ for haloperidol

and $s\kappa_{BIP}$ for biperiden) were included to model changes in the discounting rate. A

positive shift parameter indicates that the corresponding drug increases discounting,

while a negative shift parameter suggests that the drug decreases discounting. The

condition for each trial is indicated by the dummy-coded variable I, which indicates the

drug condition of the current trial.

$$P\big(HC_{(t)}\big) = \frac{\exp{(SV(HC_{(t)})* \beta)}}{\exp\big(SV(HC_{(t)})* \beta\big)+\exp{(SV(LC_{(t)})* \beta)}}$$ (5)

$$\beta = \beta_{PLC} + I_{HAL_{(t)}} * s\beta_{HAL} + I_{BIP_{(t)}} * s\beta_{BIP}$$ (6)

We then used a softmax function to transform the option values to choice probabilities

(Eq. 5). The choice stochasticity was modelled using the inverse temperature

parameter β. A lower β value indicates more stochastic behaviour. Conversely, a

higher β value suggests more deterministic choices.

Similar to the previous analysis, two additional shift parameters ($s\beta_{HAL}$ for

haloperidol and $s\beta_{BIP}$ for biperiden) were introduced to capture potential drug effects on

choice stochasticity (Eq. 6). A positive $s\beta$ parameter indicates that the corresponding

drug decreases the level of stochasticity in choices, while a negative $s\beta$ parameter

suggests that the drug increases stochasticity.

Model estimation was performed using MCMC sampling as implemented in

STAN (Stan Development Team, 2020) via R and the rSTAN package (Version

2.21.0). We utilized separate group-level distributions for all parameters in the placebo

condition (i.e., κ and β), as well as for the shift parameters (i.e., $s\kappa_{HAL}$, $s\kappa_{BIP}$, $s\beta_{HAL}$, and s

β_{BIP}), which capture potential modulatory effects of the drugs. Prior distributions for the

parameter means and standard deviations were chosen within plausible ranges based

on previous findings (Knauth & Peters, 2022; Lockwood et al., 2021, 2022; Mathar et

al., 2022; Wagner et al., 2020). The sampling process involved running four chains

with 4000 iterations after a warmup period of 3000 iterations. Chain convergence was

assessed using the Gelman-Rubinstein convergence diagnostic r-hat, with values less

than 1.01 considered acceptable (Gelman & Rubin, 1992). We report the mean of the

posterior group distribution for all parameters, along with the associated 95% HDI.

Please refer to the Supplementary Methods for details of the prior distributions, model

comparison, and model estimation procedure.

Model Validation & Parameter Recovery

To assess the model's ability to capture and recover important characteristics of the

data, we conducted a model validation and parameter recovery analysis. This involved

generating 500 synthetic datasets per participant using the posterior distributions of

subject-level parameters obtained from the winning models. From these 500 datasets,

we randomly selected 10 datasets for each participant and conducted the same

analysis as described above, using the synthetic datasets instead of the actual data.

First, to validate that the simulated data accurately captures the key features of

the participants' behaviour, we simulated the relative choice rates and visualized how

behavioural patterns changed as a function of varying levels of costs and rewards.

Next, to evaluate the recovery of group-level parameters, we examined whether the

simulated group-level parameters fell within the 95% HDI of the actual group-level

parameter distribution. For the subject-level parameter estimates, we calculated the

correlation between the simulated (averaged across all 10 simulated datasets) and

true estimated subject-level parameters. The results of the model validation and

parameter recovery are presented in the Supplementary Results.

Drug Effects on Vital Signs, Mood, Trail-Making Performance, MVC, and Effort Rating

The effects of the drugs on mood (alertness, calmness, and contentedness) and

physiological measures (heart rate, systolic, and diastolic blood pressure) were

analysed using Bayesian Linear Mixed Models. If credible drug interactions were

found, we further explored the relationship between drug induced alterations in

behaviour and potential explanatory factors, including changes in subjective mood

ratings and/or physiological responses. Specifically, we calculated the absolute

differences in mood ratings and physiological parameters between T_0 (before drug

administration) and the time point where a credible drug effect was observed (either

T_1 or T_2). These difference values were then correlated with the shift parameters that

capture drug-induced changes in performance. The purpose of this Bayesian

correlation analysis was to exclude that drug-induced changes in mood ratings or

physiological measures could account for the observed drug-induced changes in

behaviour. We further investigated potential confounding drug effects on trail-making

response times, MVC, and subjective effort perception. These analyses did not reveal

significant effects of drug manipulation on any of these measures (see Supplementary

Results). More detailed descriptions of these analysis are presented in the

Supplementary Methods.

Association between task- and drug-specific Computational Parameter Estimates

We investigated the association between discounting tendencies across different cost

domains, specifically exploring whether participants who exhibit stronger effort

discounting also displayed stronger delay discounting. To examine this, we performed

a Bayesian correlation analysis using the mean estimates of the κ parameters from

the placebo conditions in both the effort and delay discounting task.

Computational parameter estimates, self-ratings scores, and demographics

In a supplementary analysis, we investigated potential associations between self-

reported questionnaire scores, demographic data (including age and sex of the

participants), and the estimated discounting parameters κ from the baseline (placebo)

condition of both tasks. To this end, we performed robust linear regressions with the

respective discounting parameter as the outcome variable (see Supplementary

Methods and Supplementary Results).

Data availability

Anonymized participant's datasets analysed in the study are available from: https://osf.io/gx76t/

Code availability

The regression analysis and computational modelling code is available from: https://osf.io/gx76t/

Acknowledgements

This work was supported by a grant from the Deutsche Forschungsgemeinschaft (DFG JO-787/5-1). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. We express our gratitude to the students who conducted and supported the data collection: Bibiana Ackers, Evelina Andronova, Alina Hansen, and Maria Angelina Morales Diaz. Computational infrastructure and support were provided by the Centre for Information and Media Technology at Heinrich Heine University Düsseldorf.

Competing interests

The authors declare no competing interests.

References

Acheson, A., & de Wit, H. (2008). Bupropion improves attention but does not affect impulsive behavior in healthy young adults. *Experimental and Clinical Psychopharmacology*, *16*(2), 113–123. https://doi.org/10.1037/1064-1297.16.2.113

Ahn, W.-Y., Haines, N., & Zhang, L. (2017). Revealing Neurocomputational Mechanisms of Reinforcement Learning and Decision-Making With the hBayesDM Package. *Computational Psychiatry (Cambridge, Mass.)*, *1*, 24–57. https://doi.org/10.1162/CPSY_a_00002

Bailey, M. R., Simpson, E. H., & Balsam, P. D. (2016). Neural substrates underlying effort, time, and risk-based decision making in motivated behavior. *Neurobiology of Learning and Memory*, *133*, 233–256. https://doi.org/10.1016/j.nlm.2016.07.015

Beck, A. T., Steer, R. A., & Brown, G. (1996). Beck Depression Inventory–II. *Psychol. Assess.* https://doi.org/10.1037/t00742-000

Berke, J. D. (2018). What does dopamine mean? *Nature Neuroscience*, *21*(6), 787–793. https://doi.org/10.1038/s41593-018-0152-y

Betts, G. D., Hynes, T. J., & Winstanley, C. A. (2021). Pharmacological evidence of a cholinergic contribution to elevated impulsivity and risky decision-making caused by adding win-paired cues to a rat gambling task. *Journal of Psychopharmacology* , 269881120972421. https://doi.org/10.1177/0269881120972421

Białaszek, W., Marcowski, P., & Ostaszewski, P. (2017). Physical and cognitive effort discounting across different reward magnitudes: Tests of discounting models. *PloS One*, *12*(7), e0182353. https://doi.org/10.1371/journal.pone.0182353

Bickel, W. K., & Marsch, L. A. (2001). Toward a behavioral economic understanding of drug dependence: delay discounting processes. *Addiction* , *96*(1), 73–86. https://doi.org/10.1046/j.1360-0443.2001.961736.x

Bickel, W. K., Pitcock, J. A., Yi, R., & Angtuaco, E. J. C. (2009). Congruence of BOLD response across intertemporal choice conditions: fictive and real money gains and losses. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *29*(27), 8839–8846. https://doi.org/10.1523/JNEUROSCI.5319-08.2009

Bogdanov, M., LoParco, S., Otto, A. R., & Sharp, M. (2022). Dopaminergic medication increases motivation to exert cognitive control by reducing subjective effort costs in Parkinson's patients. *Neurobiology of Learning and Memory*, *193*, 107652. https://doi.org/10.1016/j.nlm.2022.107652

Bond, A., & Lader, M. (1974). The use of analogue scales in rating subjective feelings. *The British Journal of Medical Psychology*, *47*(3), 211–218. https://doi.org/10.1111/j.2044-8341.1974.tb02285.x

Bonnelle, V., Veromann, K.-R., Burnett Heyes, S., Lo Sterzo, E., Manohar, S., & Husain, M. (2015). Characterization of reward and effort mechanisms in apathy. *Journal of Physiology, Paris*, *109*(1–3), 16–26. https://doi.org/10.1016/j.jphysparis.2014.04.002

Borderies, N., Bornert, P., Gilardeau, S., & Bouret, S. (2020). Pharmacological evidence for the implication of noradrenaline in effort. *PLoS Biology*, *18*(10), e3000793. https://doi.org/10.1371/journal.pbio.3000793

Brocks, D. R. (1999). Anticholinergic drugs used in Parkinson's disease: An overlooked class of drugs from a pharmacokinetic perspective. *Journal of Pharmacy & Pharmaceutical Sciences: A Publication of the Canadian Society for Pharmaceutical Sciences, Societe Canadienne Des Sciences Pharmaceutiques*, *2*(2), 39–46. https://www.ncbi.nlm.nih.gov/pubmed/10952768

Burke, C. J., Soutschek, A., Weber, S., Raja Beharelle, A., Fehr, E., Haker, H., & Tobler, P. N. (2018). Dopamine Receptor-Specific Contributions to the Computation of Value. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *43*(6), 1415–1424. https://doi.org/10.1038/npp.2017.302

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using
Stan. *Journal of Statistical Software*, *80*, 1–28.
https://doi.org/10.18637/jss.v080.i01

Chantranupong, L., Beron, C. C., Zimmer, J. A., Wen, M. J., Wang, W., & Sabatini, B.
L. (2023). Dopamine and glutamate regulate striatal acetylcholine in decision-
making. *Nature*. https://doi.org/10.1038/s41586-023-06492-9

Chong, T. T.-J., Apps, M. A. J., Giehl, K., Hall, S., Clifton, C. H., & Husain, M. (2018).
Computational modelling reveals distinct patterns of cognitive and physical
motivation in elite athletes. *Scientific Reports*, *8*(1), 11888.
https://doi.org/10.1038/s41598-018-30220-3

Chong, T. T.-J., Apps, M., Giehl, K., Sillence, A., Grima, L. L., & Husain, M. (2017).
Neurocomputational mechanisms underlying subjective valuation of effort
costs. *PLoS Biology*, *15*(2), e1002598.
https://doi.org/10.1371/journal.pbio.1002598

Chong, T. T.-J., Fortunato, E., & Bellgrove, M. A. (2023). Amphetamines improve the
motivation to invest effort in Attention-Deficit/Hyperactivity Disorder. *The*
Journal of Neuroscience: The Official Journal of the Society for Neuroscience.
https://doi.org/10.1523/JNEUROSCI.0982-23.2023

Collins, A. G. E., & Frank, M. J. (2014). Opponent actor learning (OpAL): modeling
interactive effects of striatal dopamine on reinforcement learning and choice
incentive. *Psychological Review*, *121*(3), 337–366.
https://doi.org/10.1037/a0037015

Cools, R., & D'Esposito, M. (2011). Inverted-U-shaped dopamine actions on human
working memory and cognitive control. *Biological Psychiatry*, *69*(12), e113-25.
https://doi.org/10.1016/j.biopsych.2011.03.028

Cousins, M. S., Atherton, A., Turner, L., & Salamone, J. D. (1996). Nucleus
accumbens dopamine depletions alter relative response allocation in a T-maze
cost/benefit task. *Behavioural Brain Research*, *74*(1–2), 189–197.
https://doi.org/10.1016/0166-4328(95)00151-4

Dallery, J., & Locey, M. L. (2005). Effects of acute and chronic nicotine on impulsive
choice in rats. *Behavioural Pharmacology*, *16*(1), 15–23.
https://doi.org/10.1097/00008877-200502000-00002

de Wit, H., Enggasser, J. L., & Richards, J. B. (2002). Acute administration of d-
amphetamine decreases impulsivity in healthy volunteers.
Neuropsychopharmacology: Official Publication of the American College of
Neuropsychopharmacology, *27*(5), 813–825. https://doi.org/10.1016/S0893-
133X(02)00343-3

Denk, F., Walton, M. E., Jennings, K. A., Sharp, T., Rushworth, M. F. S., & Bannerman,
D. M. (2005). Differential involvement of serotonin and dopamine systems in
cost-benefit decisions about delay or effort. *Psychopharmacology*, *179*(3), 587–
596. https://doi.org/10.1007/s00213-004-2059-4

Di Chiara, G., Morelli, M., & Consolo, S. (1994). Modulatory functions of
neurotransmitters in the striatum: ACh/dopamine/NMDA interactions. *Trends in*
Neurosciences, *17*(6), 228–233. https://doi.org/10.1016/0166-2236(94)90005-
1

Eisenegger, C., Naef, M., Linssen, A., Clark, L., Gandamaneni, P. K., Müller, U., &
Robbins, T. W. (2014). Role of dopamine D2 receptors in human reinforcement
learning. *Neuropsychopharmacology: Official Publication of the American*
College of Neuropsychopharmacology, *39*(10), 2366–2375.
https://doi.org/10.1038/npp.2014.84

Farrar, A. M., Segovia, K. N., Randall, P. A., Nunes, E. J., Collins, L. E., Stopper, C.
M., Port, R. G., Hockemeyer, J., Müller, C. E., Correa, M., & Salamone, J. D.
(2010). Nucleus accumbens and effort-related functions: behavioral and neural
markers of the interactions between adenosine A2A and dopamine D2
receptors. *Neuroscience*, *166*(4), 1056–1067.
https://doi.org/10.1016/j.neuroscience.2009.12.056

Floresco, S. B., Tse, M. T. L., & Ghods-Sharifi, S. (2008). Dopaminergic and
glutamatergic regulation of effort- and delay-based decision making.
Neuropsychopharmacology: Official Publication of the American College of

Neuropsychopharmacology, *33*(8), 1966–1979. https://doi.org/10.1038/sj.npp.1301565

Foster, D. J., Wilson, J. M., Remke, D. H., Mahmood, M. S., Uddin, M. J., Wess, J., Patel, S., Marnett, L. J., Niswender, C. M., Jones, C. K., Xiang, Z., Lindsley, C. W., Rook, J. M., & Conn, P. J. (2016). Antipsychotic-like Effects of M4 Positive Allosteric Modulators Are Mediated by CB2 Receptor-Dependent Inhibition of Dopamine Release. *Neuron*, *91*(6), 1244–1252. https://doi.org/10.1016/j.neuron.2016.08.017

Frank, M. J., & O'Reilly, R. C. (2006). A mechanistic account of striatal dopamine function in human cognition: psychopharmacological studies with cabergoline and haloperidol. *Behavioral Neuroscience*, *120*(3), 497–517. https://doi.org/10.1037/0735-7044.120.3.497

Gelman, A., Jakulin, A., Pittau, M. G., & Su, Y.-S. (2008). A weakly informative default prior distribution for logistic and other regression models. *The Annals of Applied Statistics*, *2*(4), 1360–1383. https://doi.org/10.1214/08-AOAS191

Gelman, A., & Rubin, D. B. (1992). Inference from Iterative Simulation Using Multiple Sequences. *Statistical Science*, *7*(4), 457–472. https://doi.org/10.1214/ss/1177011136

Gerber, D. J., Sotnikova, T. D., Gainetdinov, R. R., Huang, S. Y., Caron, M. G., & Tonegawa, S. (2001). Hyperactivity, elevated dopaminergic transmission, and response to amphetamine in M1 muscarinic acetylcholine receptor-deficient mice. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(26), 15312–15317. https://doi.org/10.1073/pnas.261583798

Goldman-Rakic, P. S., Muly, III, E.C, & Williams, G. V. (2000). D1 receptors in prefrontal cells and circuits. *Brain Research Reviews*, *31*(2), 295–301. https://doi.org/10.1016/S0165-0173(99)00045-4

Green, L., & Myerson, J. (2004). A discounting framework for choice with delayed and probabilistic rewards. *Psychological Bulletin*, *130*(5), 769–792. https://doi.org/10.1037/0033-2909.130.5.769

Grimaldi, R., Perucca, E., Ruberto, G., Gelmi, C., Trimarchi, F., Hollmann, M., &
Crema, A. (1986). Pharmacokinetic and pharmacodynamic studies following
the intravenous and oral administration of the antiparkinsonian drug biperiden
to normal subjects. *European Journal of Clinical Pharmacology*, *29*(6), 735–
737. https://doi.org/10.1007/BF00615970

Hailwood, J. M., Heath, C. J., Phillips, B. U., Robbins, T. W., Saksida, L. M., & Bussey,
T. J. (2019). Blockade of muscarinic acetylcholine receptors facilitates
motivated behaviour and rescues a model of antipsychotic-induced
amotivation. *Neuropsychopharmacology: Official Publication of the American*
College of Neuropsychopharmacology, *44*(6), 1068–1075.
https://doi.org/10.1038/s41386-018-0281-8

Hamidovic, A., Kang, U. J., & de Wit, H. (2008). Effects of low to moderate acute doses
of pramipexole on impulsivity and cognition in healthy volunteers. *Journal of*
Clinical Psychopharmacology, *28*(1), 45–51.
https://doi.org/10.1097/jcp.0b013e3181602fab

Hartmann, M. N., Hager, O. M., Tobler, P. N., & Kaiser, S. (2013). Parabolic
discounting of monetary rewards by physical effort. *Behavioural Processes*,
100, 192–196. https://doi.org/10.1016/j.beproc.2013.09.014

Jocham, G., Klein, T. A., & Ullsperger, M. (2011). Dopamine-mediated reinforcement
learning signals in the striatum and ventromedial prefrontal cortex underlie
value-based choices. *The Journal of Neuroscience: The Official Journal of the*
Society for Neuroscience, *31*(5), 1606–1613.
https://doi.org/10.1523/JNEUROSCI.3904-10.2011

Jocham, G., Klein, T. A., & Ullsperger, M. (2014). Differential modulation of
reinforcement learning by D2 dopamine and NMDA glutamate receptor
antagonism. *The Journal of Neuroscience: The Official Journal of the Society*
for Neuroscience, *34*(39), 13151–13162.
https://doi.org/10.1523/JNEUROSCI.0757-14.2014

Kapur, S., Remington, G., Jones, C., Wilson, A., DaSilva, J., Houle, S., & Zipursky, R.
(1996). High levels of dopamine D2 receptor occupancy with low-dose

haloperidol treatment: a PET study. *The American Journal of Psychiatry*,
153(7), 948–950. https://doi.org/10.1176/ajp.153.7.948

Kayser, A. S., Allen, D. C., Navarro-Cebrian, A., Mitchell, J. M., & Fields, H. L. (2012).
Dopamine, corticostriatal connectivity, and intertemporal choice. *The Journal of*
Neuroscience: The Official Journal of the Society for Neuroscience, *32*(27),
9402–9409. https://doi.org/10.1523/JNEUROSCI.1180-12.2012

Kirby, K. N., & Maraković, N. N. (1995). Modeling myopic decisions: Evidence for
hyperbolic delay-discounting within subjects and amounts. *Organizational*
Behavior and Human Decision Processes, *64*(1), 22–30.
https://doi.org/10.1006/obhd.1995.1086

Klein-Flügge, M. C., Kennerley, S. W., Saraiva, A. C., Penny, W. D., & Bestmann, S.
(2015). Behavioral modeling of human choices reveals dissociable effects of
physical effort and temporal delay on reward devaluation. *PLoS Computational*
Biology, *11*(3), e1004116. https://doi.org/10.1371/journal.pcbi.1004116

Knauth, K., & Peters, J. (2022). Trial-wise exposure to visual emotional cues increases
physiological arousal but not temporal discounting. *Psychophysiology*, *59*(4),
e13996. https://doi.org/10.1111/psyp.13996

Kolokotroni, K. Z., Rodgers, R. J., & Harrison, A. A. (2011). Acute nicotine increases
both impulsive choice and behavioural disinhibition in rats.
Psychopharmacology, *217*(4), 455–473. https://doi.org/10.1007/s00213-011-
2296-2

Kruschke, J. (2014). *Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and*
Stan. Academic Press.

Kudo, S., & Ishizaki, T. (1999). Pharmacokinetics of haloperidol: an update. *Clinical*
Pharmacokinetics, *37*(6), 435–456. https://doi.org/10.2165/00003088-
199937060-00001

Kühner, C., Bürger, C., Keller, F., & Hautzinger, M. (2007). [Reliability and validity of
the Revised Beck Depression Inventory (BDI-II). Results from German

samples]. *Der Nervenarzt, 78*(6), 651–656. https://doi.org/10.1007/s00115-
006-2098-7

Le Bouc, R., Rigoux, L., Schmidt, L., Degos, B., Welter, M.-L., Vidailhet, M.,
Daunizeau, J., & Pessiglione, M. (2016). Computational Dissection of
Dopamine Motor and Motivational Functions in Humans. The Journal of
Neuroscience: The Official Journal of the Society for Neuroscience, 36(25),
6623–6633. https://doi.org/10.1523/JNEUROSCI.3078-15.2016

Le Heron, C., Plant, O., Manohar, S., Ang, Y.-S., Jackson, M., Lennox, G., Hu, M. T.,
& Husain, M. (2018). Distinct effects of apathy and dopamine on effort-based
decision-making in Parkinson's disease. *Brain: A Journal of Neurology*, *141*(5),
1455–1469. https://doi.org/10.1093/brain/awy110

Liem-Moolenaar, M., Gray, F. A., de Visser, S. J., Franson, K. L., Schoemaker, R. C.,
Schmitt, J. A. J., Cohen, A. F., & van Gerven, J. M. A. (2010). Psychomotor and
cognitive effects of a single oral dose of talnetant (SB223412) in healthy
volunteers compared with placebo or haloperidol. *Journal of*
Psychopharmacology   , *24*(1), 73–82.
https://doi.org/10.1177/0269881108094524

Linssen, A. M. W., Vuurman, E. F. P. M., Sambeth, A., Nave, S., Spooren, W., Vargas,
G., Santarelli, L., & Riedel, W. J. (2011). Contingent negative variation as a
dopaminergic biomarker: evidence from dose-related effects of
methylphenidate. *Psychopharmacology*, *218*(3), 533–542.
https://doi.org/10.1007/s00213-011-2345-x

Lockwood, P. L., Abdurahman, A., Gabay, A. S., Drew, D., Tamm, M., Husain, M., &
Apps, M. A. J. (2021). Aging Increases Prosocial Motivation for Effort.
Psychological Science, 956797620975781.
https://doi.org/10.1177/0956797620975781

Lockwood, P. L., Hamonet, M., Zhang, S. H., Ratnavel, A., Salmony, F. U., Husain,
M., & Apps, M. A. J. (2017). Prosocial apathy for helping others when effort is
required. *Nature Human Behaviour*, *1*(7), 0131. https://doi.org/10.1038/s41562-
017-0131

Lockwood, P. L., Wittmann, M. K., Nili, H., Matsumoto-Ryan, M., Abdurahman, A.,
Cutler, J., Husain, M., & Apps, M. A. J. (2022). Distinct neural representations
for prosocial and self-benefiting effort. *Current Biology: CB*.
https://doi.org/10.1016/j.cub.2022.08.010

Lueken, U., Seidl, U., Schwarz, M., Völker, L., Naumann, D., Mattes, K., Schröder, J.,
& Schweiger, E. (2006). [Psychometric properties of a German version of the
Apathy Evaluation Scale]. *Fortschritte der Neurologie-Psychiatrie*, *74*(12), 714–
722. https://doi.org/10.1055/s-2006-932164

Madden, G. J., Begotka, A. M., Raiff, B. R., & Kastern, L. L. (2003). Delay discounting
of real and hypothetical rewards. *Experimental and Clinical*
Psychopharmacology, *11*(2), 139–145. https://doi.org/10.1037/1064-
1297.11.2.139

Mai, B., Sommer, S., & Hauber, W. (2012). Motivational states influence effort-based
decision making in rats: the role of dopamine in the nucleus accumbens.
Cognitive, Affective & Behavioral Neuroscience, *12*(1), 74–84.
https://doi.org/10.3758/s13415-011-0068-4

Marin, R. S., Biedrzycki, R. C., & Firinciogullari, S. (1991). Reliability and validity of the
Apathy Evaluation Scale. *Psychiatry Research*, *38*(2), 143–162.
https://doi.org/10.1016/0165-1781(91)90040-v

Mathar, D., Erfanian Abdoust, M., Marrenbach, T., Tuzsus, D., & Peters, J. (2022).
The catecholamine precursor Tyrosine reduces autonomic arousal and
decreases decision thresholds in reinforcement learning and temporal
discounting. *PLoS Computational Biology*, *18*(12), e1010785.
https://doi.org/10.1371/journal.pcbi.1010785

Matthews, J., Pisauro, M. A., Jurgelis, M., Müller, T., Vassena, E., Chong, T. T.-J., &
Apps, M. A. J. (2023). Computational mechanisms underlying the dynamics of
physical and cognitive fatigue. Cognition, 240, 105603.
https://doi.org/10.1016/j.cognition.2023.105603

McGuigan, S., Zhou, S.-H., Brosnan, M. B., Thyagarajan, D., Bellgrove, M. A., &
Chong, T. T.-J. (2019). Dopamine restores cognitive motivation in Parkinson's
disease. *Brain: A Journal of Neurology*, *142*(3), 719–732.
https://doi.org/10.1093/brain/awy341

Mendez, I. A., Gilbert, R. J., Bizon, J. L., & Setlow, B. (2012). Effects of acute
administration of nicotinic and muscarinic cholinergic agonists and antagonists
on performance in different cost-benefit decision making tasks in rats.
Psychopharmacology, 224(4), 489–499. https://doi.org/10.1007/s00213-012-
2777-y

Meule, A., Vögele, C., & Kübler, A. (2011). Psychometrische Evaluation der deutschen
Barratt Impulsiveness Scale – Kurzversion (BIS-15). *Diagnostica*, *57*(3), 126–
133. https://doi.org/10.1026/0012-1924/a000042

Meyniel, F., Goodwin, G. M., Deakin, J. W., Klinge, C., MacFadyen, C., Milligan, H.,
Mullings, E., Pessiglione, M., & Gaillard, R. (2016). A specific role for serotonin
in overcoming effort cost. *ELife*, *5*. https://doi.org/10.7554/eLife.17282

Michely, J., Viswanathan, S., Hauser, T. U., Delker, L., Dolan, R. J., & Grefkes, C.
(2020). The role of dopamine in dynamic effort-reward integration.
Neuropsychopharmacology: Official Publication of the American College of
Neuropsychopharmacology, *45*(9), 1448–1453.
https://doi.org/10.1038/s41386-020-0669-0

Mikus, N., Korb, S., Massaccesi, C., Gausterer, C., Graf, I., Willeit, M., Eisenegger,
C., Lamm, C., Silani, G., & Mathys, C. (2022). Effects of dopamine D2/3 and
opioid receptor antagonism on the trade-off between model-based and model-
free behaviour in healthy volunteers. *ELife*, *11*.
https://doi.org/10.7554/eLife.79661

Miyazaki, K., Miyazaki, K. W., Sivori, G., Yamanaka, A., Tanaka, K. F., & Doya, K.
(2020). Serotonergic projections to the orbitofrontal and medial prefrontal
cortices differentially modulate waiting for future rewards. *Science Advances*,
6(48). https://doi.org/10.1126/sciadv.abc7246

Moehle, M. S., & Conn, P. J. (2019). Roles of the M4 acetylcholine receptor in the basal ganglia and the treatment of movement disorders. *Movement Disorders: Official Journal of the Movement Disorder Society*, *34*(8), 1089–1099. https://doi.org/10.1002/mds.27740

Morrison, S. E., & Nicola, S. M. (2014). Neurons in the nucleus accumbens promote selection bias for nearer objects. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *34*(42), 14147–14162. https://doi.org/10.1523/JNEUROSCI.2197-14.2014

Müller, T., Klein-Flügge, M. C., Manohar, S., Husain, M., & Apps, M. A. J. (2021). Neural and computational mechanisms of momentary fatigue and persistence in effort-based choice. Nature Communications, 12(1), 4593. https://doi.org/10.1038/s41467-021-24927-7

Myslivecek, J. (2021). Two Players in the Field: Hierarchical Model of Interaction between the Dopamine and Acetylcholine Signaling Systems in the Striatum. *Biomedicines*, *9*(1). https://doi.org/10.3390/biomedicines9010025

Nordström, A. L., Farde, L., & Halldin, C. (1992). Time course of D2-dopamine receptor occupancy examined by PET after single oral doses of haloperidol. *Psychopharmacology*, *106*(4), 433–438. https://doi.org/10.1007/BF02244811

Nunes, E. J., Randall, P. A., Podurgiel, S., Correa, M., & Salamone, J. D. (2013). Nucleus accumbens neurotransmission and effort-related choice behavior in food motivation: effects of drugs acting on dopamine, adenosine, and muscarinic acetylcholine receptors. *Neuroscience and Biobehavioral Reviews*, *37*(9 Pt A), 2015–2025. https://doi.org/10.1016/j.neubiorev.2013.04.002

Ojala, K. E., Janssen, L. K., Hashemi, M. M., Timmer, M. H. M., Geurts, D. E. M., Ter Huurne, N. P., Cools, R., & Sescousse, G. (2018). Dopaminergic Drug Effects on Probability Weighting during Risky Decision Making. *ENeuro*, *5*(2). https://doi.org/10.1523/ENEURO.0330-18.2018

Ozga, J. E., & Anderson, K. G. (2018). Reduction in delay discounting due to nicotine and its attenuation by cholinergic antagonists in Lewis and Fischer 344 rats.

Psychopharmacology, 235(1), 155–168. https://doi.org/10.1007/s00213-017-4752-0

Peters, J., & D'Esposito, M. (2020). The drift diffusion model as the choice rule in inter-temporal and risky choice: A case study in medial orbitofrontal cortex lesion patients and controls. *PLoS Computational Biology*, *16*(4), e1007615. https://doi.org/10.1371/journal.pcbi.1007615

Petzold, J., Kienast, A., Lee, Y., Pooseh, S., London, E. D., Goschke, T., & Smolka, M. N. (2019). Baseline impulsivity may moderate L-DOPA effects on value-based decision-making. *Scientific Reports*, *9*(1), 5652. https://doi.org/10.1038/s41598-019-42124-x

Phillips, P. E. M., Walton, M. E., & Jhou, T. C. (2007). Calculating utility: preclinical evidence for cost-benefit analysis by mesolimbic dopamine. *Psychopharmacology*, *191*(3), 483–495. https://doi.org/10.1007/s00213-006-0626-6

Pine, A., Shiner, T., Seymour, B., & Dolan, R. J. (2010). Dopamine, time, and impulsivity in humans. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *30*(26), 8888–8896. https://doi.org/10.1523/JNEUROSCI.6028-09.2010

Randall, P. A., Nunes, E. J., Janniere, S. L., Stopper, C. M., Farrar, A. M., Sager, T. N., Baqi, Y., Hockemeyer, J., Müller, C. E., & Salamone, J. D. (2011). Stimulant effects of adenosine antagonists on operant behavior: differential actions of selective A2A and A1 antagonists. *Psychopharmacology*, *216*(2), 173–186. https://doi.org/10.1007/s00213-011-2198-3

Rutledge, R. B., Skandali, N., Dayan, P., & Dolan, R. J. (2015). Dopaminergic Modulation of Decision Making and Subjective Well-Being. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *35*(27), 9811–9822. https://doi.org/10.1523/JNEUROSCI.0702-15.2015

Salamone, J. D., Cousins, M. S., & Bucher, S. (1994). Anhedonia or anergia? Effects of haloperidol and nucleus accumbens dopamine depletion on instrumental

response selection in a T-maze cost/benefit procedure. *Behavioural Brain Research*, *65*(2), 221–229. https://doi.org/10.1016/0166-4328(94)90108-2

Salamone, J. D., & Correa, M. (2012). The mysterious motivational functions of mesolimbic dopamine. *Neuron*, *76*(3), 470–485. https://doi.org/10.1016/j.neuron.2012.10.021

Salamone, J. D., Correa, M., Ferrigno, S., Yang, J.-H., Rotolo, R. A., & Presby, R. E. (2018). The Psychopharmacology of Effort-Related Decision Making: Dopamine, Adenosine, and Insights into the Neurochemistry of Motivation. *Pharmacological Reviews*, *70*(4), 747–762. https://doi.org/10.1124/pr.117.015107

Schoemaker, H., Claustre, Y., Fage, D., Rouquier, L., Chergui, K., Curet, O., Oblin, A., Gonon, F., Carter, C., Benavides, J., & Scatton, B. (1997). Neurochemical characteristics of amisulpride, an atypical dopamine D2/D3 receptor antagonist with both presynaptic and limbic selectivity. *The Journal of Pharmacology and Experimental Therapeutics*, *280*(1), 83–97. https://www.ncbi.nlm.nih.gov/pubmed/8996185

Schultz, W. (2016a). Dopamine reward prediction error coding. *Dialogues in Clinical Neuroscience*, *18*(1), 23–32. https://doi.org/10.31887/DCNS.2016.18.1/wschultz

Schultz, W. (2016b). Dopamine reward prediction-error signalling: a two-component response. *Nature Reviews. Neuroscience*, *17*(3), 183–195. https://doi.org/10.1038/nrn.2015.26

Seaman, K. L., Brooks, N., Karrer, T. M., Castrellon, J. J., Perkins, S. F., Dang, L. C., Hsu, M., Zald, D. H., & Samanez-Larkin, G. R. (2018). Subjective value representations during effort, probability and time discounting across adulthood. *Social Cognitive and Affective Neuroscience*, *13*(5), 449–459. https://doi.org/10.1093/scan/nsy021

Soder, H. E., Cooper, J. A., Lopez-Gamundi, P., Hoots, J. K., Nunez, C., Lawlor, V. M., Lane, S. D., Treadway, M. T., & Wardle, M. C. (2020). Dose-response

effects of d-amphetamine on effort-based decision-making and reinforcement learning. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*. https://doi.org/10.1038/s41386-020-0779-8

Soutschek, A., Gvozdanovic, G., Kozak, R., Duvvuri, S., de Martinis, N., Harel, B., Gray, D. L., Fehr, E., Jetter, A., & Tobler, P. N. (2020). Dopaminergic D1 Receptor Stimulation Affects Effort and Risk Preferences. *Biological Psychiatry*, *87*(7), 678–685. https://doi.org/10.1016/j.biopsych.2019.09.002

Soutschek, A., Jetter, A., & Tobler, P. N. (2023). Toward a Unifying Account of Dopamine's Role in Cost-Benefit Decision Making. *Biological Psychiatry Global Open Science*, *3*(2), 179–186. https://doi.org/10.1016/j.bpsgos.2022.02.010

Soutschek, A., & Tobler, P. N. (2023). A process model account of the role of dopamine in intertemporal choice. *ELife*, *12*. https://doi.org/10.7554/eLife.83734

Spinella, M. (2007). Normative data and a short form of the Barratt Impulsiveness Scale. *The International Journal of Neuroscience*, *117*(3), 359–368. https://doi.org/10.1080/00207450600588881

Stanhope, K. J., Mirza, N. R., Bickerdike, M. J., Bright, J. L., Harrington, N. R., Hesselink, M. B., Kennett, G. A., Lightowler, S., Sheardown, M. J., Syed, R., Upton, R. L., Wadsworth, G., Weiss, S. M., & Wyatt, A. (2001). The muscarinic receptor agonist xanomeline has an antipsychotic-like profile in the rat. *The Journal of Pharmacology and Experimental Therapeutics*, *299*(2), 782–792. https://www.ncbi.nlm.nih.gov/pubmed/11602695

Sudo, Y., Suhara, T., Suzuki, K., Okubo, Y., Yoshikawa, K., Uchida, S., Sassa, T., Okauchi, T., Sasaki, Y., & Matsushita, M. (1999). Muscarinic receptor occupancy by biperiden in living human brain. *Life Sciences*, *64*(8), PL99-104. https://doi.org/10.1016/s0024-3205(98)00613-4

Swart, J. C., Froböse, M. I., Cook, J. L., Geurts, D. E., Frank, M. J., Cools, R., & den Ouden, H. E. (2017). Catecholaminergic challenge uncovers distinct Pavlovian

and instrumental mechanisms of motivated (in)action. *ELife*, *6*. https://doi.org/10.7554/eLife.22169

Tanaka, S., O'Doherty, J. P., & Sakagami, M. (2019). The cost of obtaining rewards enhances the reward prediction error signal of midbrain dopamine neurons. *Nature Communications*, *10*(1), 3674. https://doi.org/10.1038/s41467-019-11334-2

Threlfell, S., Clements, M. A., Khodai, T., Pienaar, I. S., Exley, R., Wess, J., & Cragg, S. J. (2010). Striatal muscarinic receptors promote activity dependence of dopamine transmission via distinct receptor subtypes on cholinergic interneurons in ventral versus dorsal striatum. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *30*(9), 3398–3408. https://doi.org/10.1523/JNEUROSCI.5620-09.2010

Tombaugh, T. N. (2004). Trail Making Test A and B: normative data stratified by age and education. *Archives of Clinical Neuropsychology: The Official Journal of the National Academy of Neuropsychologists*, *19*(2), 203–214. https://doi.org/10.1016/S0887-6177(03)00039-8

van den Bosch, R., Lambregts, B., Määttä, J., Hofmans, L., Papadopetraki, D., Westbrook, A., Verkes, R.-J., Booij, J., & Cools, R. (2022). Striatal dopamine dissociates methylphenidate effects on value-based versus surprise-based reversal learning. *Nature Communications*, *13*(1), 4962. https://doi.org/10.1038/s41467-022-32679-1

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432. https://doi.org/10.1007/s11222-016-9696-4

Wagner, B., Clos, M., Sommer, T., & Peters, J. (2020). Dopaminergic Modulation of Human Intertemporal Choice: A Diffusion Model Analysis Using the D2-Receptor Antagonist Haloperidol. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *40*(41), 7936–7948. https://doi.org/10.1523/JNEUROSCI.0592-20.2020

Wardle, M. C., Treadway, M. T., Mayo, L. M., Zald, D. H., & de Wit, H. (2011). Amping
up effort: effects of d-amphetamine on human effort-based decision-making.
The Journal of Neuroscience: The Official Journal of the Society for
Neuroscience, *31*(46), 16597–16602.
https://doi.org/10.1523/JNEUROSCI.4387-11.2011

Webber, H. E., Lopez-Gamundi, P., Stamatovich, S. N., de Wit, H., & Wardle, M. C.
(2020). Using pharmacological manipulations to study the role of dopamine in
human reward functioning: A review of studies in healthy adults. *Neuroscience*
and Biobehavioral Reviews. https://doi.org/10.1016/j.neubiorev.2020.11.004

Westbrook, A., & Frank, M. J. (2018). Dopamine and Proximity in Motivation and
Cognitive Control. *Current Opinion in Behavioral Sciences*, *22*, 28–34.
https://doi.org/10.1016/j.cobeha.2017.12.011

Zack, M., & Poulos, C. X. (2007). A D2 antagonist enhances the rewarding and priming
effects of a gambling episode in pathological gamblers.
Neuropsychopharmacology: Official Publication of the American College of
Neuropsychopharmacology, *32*(8), 1678–1686.
https://doi.org/10.1038/sj.npp.1301295

1290

1291

1292
1293

Fig. 1 Study procedure and experimental tasks. (**a**) The experimental sessions were standardized across all sessions except for the drug treatment, which was counterbalanced. To account for the different times of haloperidol and biperiden to reach peak plasma levels, a dummy drug application was introduced, ensuring that the maximum concentration of both drugs was aligned to task execution. Approximately 180 minutes after the first capsule and approximately 60 minutes after the second capsule, participants engaged in the effort discounting task (b), followed by the delay discounting task (c). Physiological measures (including heart rate and blood pressure) and mood ratings (using the Bond & Lader Visual Analogue Scales) were collected at three distinct time points. Additionally, participants completed the trail-making test part A before task execution (see Materials and Methods for more details). In both tasks, participants were presented with two alternative options, each providing information about a monetary reward in return for specific costs. (**b**) Effort Discounting Task. One option required less effort (indicated by the horizontal yellow line) and provided a smaller reward (indicated by the number of apples, low-cost option), while the other option required more effort and provided a higher reward (high-cost option). Participants then chose one option and exerted the required effort (adjusted to the maximum voluntary contraction, MVC) for at least 1 second. (**c**) Delay discounting task. Similarly, participants were presented with two offers: a smaller but immediately available reward (low-cost option) or a larger reward available after the delay indicated (high-cost option).

Fig. 2 Behavioural Performance in the Effort (left panel) and Delay (right panel) Discounting Task. (**a**) The overall proportions of high-cost choices were modulated by drug administration in the effort (left), but not in the delay discounting task (right). Biperiden increased and haloperidol decreased the willingness to invest physical effort in return for reward. (**b**) The probability of choosing the high-cost option increased as a function of increasing reward magnitude for both the effort and delay discounting tasks. In the effort discounting task (left), biperiden increased the impact of the reward level on choice. (**c**) Similarly, the tendency to choose the high-cost option decreased as a function of increasing levels of effort and delay. This effect is reduced by haloperidol in the delay discounting task (right). Values in **a** show group-level (single-subject) means represented by bold (light) dots. Values in **b** and **c** display averaged group-level means per reward and cost level, with error bars representing the standard error of the mean. In **b**, reward levels are presented as the difference in magnitude between the high- and low-cost option in the effort discontinuing task and as the absolute reward value of the high-cost option in the delay discounting task. Likewise in **c**, the effort level represents the difference between the proportions of the individually calibrated MVC of the high- versus low-cost option, while the delay levels indicate the delay of the high-cost option.

Fig. 3 Drug Effects on Choice Behaviour. Posterior distributions and 95% HDI of the Logistic Bayesian Generalized Linear Mixed Models depict the estimate of each effect on choosing the high-cost option. (**a**) Biperiden credibly increased the overall willingness to invest physical effort for a corresponding reward, while haloperidol had the opposite effect. (**b**) Biperiden increased reward sensitivity in the effort discounting task, as indicated by a biperiden-by-reward interaction, without a credible effect of haloperidol. (**c**) In contrast, neither drug affected effort sensitivity. (**d**) In the delay discounting task, the willingness to tolerate delays for rewards was not affected by either drug. (**e**) Likewise, neither drug modulated reward sensitivity in the delay discounting task. (**f**) However, haloperidol decreased delay sensitivity, with no credible effect of biperiden. Here, a positive estimate of the interaction effect between haloperidol and delay indicates a reduction of the negative parameter estimate associated with delay, suggesting an attenuation of the impact of delay on choice behaviour. Bold dots represent the mean group-level estimate of the posterior distribution. The horizontal bars represent the group-level 95% highest density interval.

Fig. 4 Drug Effect on Computational Parameters. Posterior distributions and changes in the subjective value from the hierarchical Bayesian models. (**a**) In the effort discounting task, the discounting parameter κ is modulated in opposite directions by the drugs, with haloperidol increasing and biperiden decreasing effort discounting. (**b**) Similarly, these opposite effects are also present in the modulation of the softmax inverse temperature β, reflecting choice stochasticity. Biperiden administration led to more deterministic choices, while haloperidol induced more stochastic choices. (**c**) Modelled discounting functions show steeper discounting under haloperidol and flatter discounting under biperiden. (**d**) In the delay discounting task, the discounting parameter κ is reduced by haloperidol, with no credible modulatory effect of biperiden. (**e**) In contrast, the softmax inverse temperature β is reduced by biperiden, indicating more stochastic choices. (**f**) Overall, participants showed flatter discounting of future rewards under haloperidol compared to placebo. In **a**, **b**, **d**, and **e**, bold (light) dots represent the group-level (participant-level) mean estimate. The horizontal bars represent the group-level 95% highest density interval. In **c** and **f**, the subjective values are displayed as a discount function of effort and delay. Parabolic (effort discounting task) and hyperbolic (delay discounting task) functions are fitted on group-level mean estimates for each drug (see Materials and Methods).

1355 **Table 1.** Summary of the Group-Level Parameter Estimates for the Effort Discounting Task,
1356 Including the Mean, Standard Deviation (SD), and the Lower and Upper Bounds of the 95% HDI
1357 Interval.
1358

Parameter	Mean	SD	2.5%	97.5%
κ	0.102	0.008	0.086	0.118
β	0.667	0.032	0.605	0.732
$S\kappa_{HAL}$	0.013	0.007	-0.000	0.027
$S\kappa_{BIP}$	-0.012	0.006	-0.025	0.000
$S\beta_{HAL}$	-0.089	0.038	-0.164	-0.010
$S\beta_{BIP}$	0.127	0.043	0.048	0.219

1359
1360

1361 **Table 2.** Summary of the Group-Level Parameter Estimates for the Delay Discounting Task,
1362 Including the Mean, Standard Deviation (SD), and the Lower and Upper Bounds of the 95% HDI
1363 Interval.
1364

Parameter	Mean	SD	2.5%	97.5%
κ	-4.621	0.271	-5.176	-4.107
β	0.356	0.045	0.272	0.449
$s\kappa_{HAL}$	-0.630	0.214	-1.083	-0.237
$s\kappa_{BIP}$	-0.125	0.142	-0.410	0.148
$s\beta_{HAL}$	-0.055	0.030	-0.114	0.006
$s\beta_{BIP}$	-0.043	0.021	-0.085	-0.000