1 **Disentangling abiotic and biotic effects of treated wastewater on stream biofilm**
2 **resistomes enables the discovery of a new planctomycete beta-lactamase**
3

4 Mustafa Attrah[1,2+], Milo R. Schärer[1+], Mauro Esposito[1], Giulia Gionchetta[3], Helmut Bürgmann[3],
5 Piet Lens[2,4], Kathrin Fenner[5,6], Jack van de Vossenberg[2], Serina L. Robinson[1]*

6 [+]co-first authors contributed equally to this work
7 *corresponding author: serina.robinson@eawag.ch
8

[1]Department of Environmental Microbiology, Eawag, Swiss Federal Institute of Aquatic Science
and Technology (Eawag), 8600 Dübendorf, Switzerland
[2]Water Supply, Sanitation and Environmental Engineering, IHE Delft Institute for Water
Education, Westvest 7, 2611 AX Delft, The Netherlands
[3]Department of Surface Waters - Research and Management, Eawag, Swiss Federal Institute of
Aquatic Science and Technology, 6047 Kastanienbaum, Switzerland
[4]National University of Ireland Galway, University Road, Galway H91 TK33, Ireland
[5]Department of Environmental Chemistry, Eawag, Swiss Federal Institute of Aquatic Science and
Technology (Eawag), 8600 Dübendorf, Switzerland
[6]Department of Chemistry, University of Zürich, 8057 Zürich, Switzerland

19 Abstract

Antibiotic resistance, mediated by environmental reservoirs, poses a threat to human and animal
health. Aquatic biofilms impacted by treated wastewater (WW) are known environmental
reservoirs for antibiotic resistance, but it remains unclear if primarily biotic or abiotic factors from
WW influence the abundance and dissemination of antibiotic resistance genes within biofilm
communities. To disentangle these factors, natural biofilms were previously grown in flume
systems with different proportions of stream water and either ultrafiltered or non-ultrafiltered
treated WW. In this study, we conducted deep shotgun metagenomic sequencing of 75 biofilm,
stream, and WW samples from these flume systems and compared microbiome and resistome
composition. Our analysis revealed a strong relationship between the resistome and prokaryotic
microbiome composition which was also associated with experimental treatment. Within biofilm
communities exposed to increasing percentages of WW, we observed a decrease in the relative
abundances of Cyanobacteriota and Verrucomicrobiota, while Bacteroidota increased. Notably,
several antibiotic resistance genes (ARGs) exhibited a significant increase in abundance in
biofilms grown with increasing WW%, while a small number of sulfonamide and beta-lactamase
ARGs showed higher abundance when biofilms were instead grown in ultrafiltered WW. Overall,
our results pointed toward the dominance of biotic over abiotic factors in determining ARG
abundances in WW-impacted stream biofilms. However, some ARGs exhibited divergent
abundance patterns through proposed gene family-specific mechanisms. To investigate these
gene families experimentally, we biochemically characterized a new, sequence-divergent beta-
lactamase from planctomycetes abundant in stream biofilms and in WW. This ARG discovery is
noteworthy, not only due to its unusual source organism belonging to the phylum
Planctomycetota, but also because it was not detected by standard computational ARG tools.

This underpins the need for continued development of ARG detection methods using workflows
with *in silico* sequence and structure analysis coupled to *in vitro* functional characterization. In
summary, this study assessed the relative importance of biotic and abiotic factors in the context
of WW discharge and its impact on antibiotic resistance genes in aquatic ecosystems.

Keywords

47   Antibiotic resistance genes, Stream resistomes, Biofilms, Flume systems, Metagenomics, Beta-
48   lactamases

Introduction

Antibiotic resistance is a major threat to public health. Pollution resulting from anthropogenic
activities alters the abundance and proliferation of antibiotic resistance genes (ARGs) in
environmental reservoirs (Singer et al., 2016). The rapidly increasing consumption of antibiotics
and release of micropollutants (MPs) into the environment (Rizzo et al., 2013) stimulates the
spread of antibiotic-resistance genes (ARGs), posing a threat to modern medical procedures and
treatments that rely on effective antibiotics (Shallcross and Davies, 2014).

Aquatic ecosystems that receive effluent from wastewater treatment plants (WWTPs) are
influenced by biotic factors, such as antibiotic-resistant bacteria released from WWTPs
(Alexander et al., 2020) and abiotic factors (Rodriguez-Mozaz et al., 2015; Rogowska et al.,
2020), such as nutrients and MPs that are not completely eliminated by conventional WWTPs
(Mukherjee et al., 2021). Several recent studies demonstrated that exposure to WWTP effluent
alters the taxonomic and ARG composition of microbial communities in downstream rivers (Carles
et al., 2021; Desiante et al., 2022; Lee et al., 2023; Matviichuk et al., 2022; Perveen et al., 2023).
However, disentangling the effects of abiotic and biotic factors on ARG abundance is challenging
in the field. Here, to systematically distinguish abiotic and biotic factors, we built on a previous
research study which used controlled flume systems to grow biofilms in ultrafiltered and non-
ultrafiltered WWTP effluent diluted with different proportions of stream water (Carles et al., 2021;
Desiante et al., 2022). Ultrafiltration (UF) treatment (pore size 0.4 µm) removed most
microorganisms and particles (biotic factors) while maintaining consistent nutrient and MP levels,
including antibiotics. Therefore, the UF treatment separating biotic from abiotic factors enabled
investigation of the factors shaping the 'resistome', i.e., the total composition of antibiotic
resistance genes (ARGs), of stream biofilms influenced by WWTP discharge.

Many studies on stream biofilms have relied on amplicon sequencing methods for taxonomic
and/or ARG profiling (Carles et al., 2021; Depetris et al., 2022; Haenelt et al., 2023; Romero et
al., 2019). As opposed to amplicon sequencing, short-read and long-read metagenomic
sequencing studies of streams and stream biofilms (Ju et al., 2022; Lee et al., 2023; Reddington
et al., 2020) do not introduce a polymerase chain reaction (PCR) amplification bias. As we
demonstrate in this study, total metagenomic sequencing also enables the identification of
sequence-divergent ARGs not detected by conventional primer pairs. Amplicon sequencing is

more sensitive in that it can aid in the detection of low abundance species, but it is also biased
towards specific taxa during the PCR process (Brumfield et al., 2020; Tremblay et al., 2015). The
PCR bias also limits the detection of some bacterial phyla such as the Planctomycetota for which
the commonly-used 16S rRNA PCR primers do not properly bind (Klindworth et al., 2013; Oyarzúa
et al., 2021). The Planctomycetota are a specific focus in this paper as an abundant phylum in
WWTPs (Y. Zhang et al., 2023), lacustrine (Andrei et al., 2019), and riverine biofilm (periphyton)
communities (Cai et al., 2021), which are undersampled by classical amplicon-based methods
(Oyarzúa et al., 2021). We therefore conducted deep shotgun metagenomic sequencing of stream
biofilms and water samples in this study, enabling us to sample less well-studied microorganisms
and their functional genes including ARGs.

Here we demonstrated a combined computational and biochemical workflow to investigate the
effects of treated wastewater (WW) on microbiome taxonomic composition and ARG composition
in flume biofilms grown in natural stream water mixed with different proportions of ultrafiltered and
non-ultrafiltered WW (Carles et al., 2021; Desiante et al., 2022). Specifically, we leveraged the
functional genes identified through shotgun metagenomic sequencing to heterologously express
an uncharacterized ARG from the phylum Planctomycetota. The enzyme displayed beta-
lactamase activity despite sharing low sequence identity with previously identified ARGs. Overall,
this study enabled dissection of the influences of biotic and abiotic factors on stream biofilm
resistomes in order to characterize both known and new ARGs.

Methods

Biofilm cultivation and sampling

The flume systems and biofilm cultivation procedures used in two independent experiments to
generate the samples sequenced in this study were described previously (Carles et al., 2021;
Desiante et al., 2022). Briefly, the system consisted of flow-through channels fed with different
mixtures of two different water sources: (1) a peri-urban stream (Chriesbach; Dübendorf,
Switzerland) and (2) treated wastewater from a pilot-size wastewater treatment plant (100-person
equivalents). In the first experiment (Exp. 1), the stream water was mixed with treated wastewater
(WW) at the following percentages: 0% (control), 10%, 30%, and 80% WW. In the second
experiment (Exp. 2), the 10% treatment was omitted and replaced by additional treatments with
30% and 80% ultrafiltered WW (UF), using a filter with a pore size of 0.4 µm to remove particles,
e.g., suspended solids and microorganisms. Glass slides were added to the flow-through
channels and biofilms naturally forming on the slides were allowed to develop for a period of four
weeks. Water parameters (pH, temperature, conductivity, dissolved oxygen) were monitored
weekly (Carles et al., 2021; Desiante et al., 2022).

DNA extraction and metagenomics assessment

Stream water, WW, UF WW samples and biofilms from glass slides were sampled as described
previously (Desiante et al., 2022; Carles et al., 2022). Total genomic DNA was extracted with the
DNeasy PowerBiofilm kit (QIAGEN) for a total of 75 samples to be submitted for metagenomic
sequencing (**Table S1**). The quality and quantity of the DNA extracts were measured using a

NanoDrop One spectrophotometer. Extracted DNA samples were stored at -20 °C and shotgun
metagenomic sequencing was performed using Illumina platform NovaSeq with a paired-end
(2×150 bp; average of 20 Gbp raw data per sample) strategy. Library construction and sequencing
was performed by Novogene (Cambridge, UK). The reads obtained from shotgun metagenomics
and containing adapters and low-quality reads (N > 10% and quality score ≤ 5) were removed by
Novogene, and the read qualities were double-checked by the authors using Fastp v.0.20.0
(https://github.com/OpenGene/fastp). After quality controls of metagenome reads, de-novo
assembly was performed using MEGAHIT v1.2.9 (Li et al., 2015). Open reading frames (ORFs)
were predicted from the assembled contigs using Prodigal v2.6.3 (Hyatt et al., 2010). Although
metagenomes from both Exp. 1 and Exp. 2 were sequenced and the data for both are re-reported
here, we focused on Exp. 2 for analysis since it included UF treatment, unlike Exp. 1.

Taxonomic profiling

The taxonomic composition of the metagenomes was profiled using a degenerate consensus
marker-gene profiling tool, mTAGs v1.0.4, with 1000 maximum accepts and 1000 maximum
rejects (Rognes et al., 2016; Salazar et al., 2021). mTAGs profiles for all samples were merged
using mTAGs' merge function. Genus- and phylum-level per-sample count tables were used for
subsequent analysis.

Reads-based ARG detection and ordination analyses

The ARG resistome was profiled using the DeepARG v1.0.2 short reads pipeline with default
settings (Arango-Argoty et al., 2018). ARG counts normalized by the 16S-rRNA counts within the
output files for each sample were merged into an abundance table. All data analysis and
visualization were done in R v4.3.0. Data manipulation and visualization were done using the
tidyverse package (Wickham et al., 2019). Principal coordinate analysis (PCoA) of microbiome
composition was done using the genus abundance table. Unaligned and unassigned reads were
removed. Genus counts were normalized using the trimmed mean of M-values (TMM) method
within the EdgeR package (Pereira et al., 2018; Robinson et al., 2010; Robinson and Oshlack,
2010). Normalized genus counts were square-root-transformed. PCoA analysis of ARG resistome
composition was done using the 16S-normalized abundance data. For both data sets, pairwise
Bray-Curtis distances between samples were computed using the 'vegdist' function from the
vegan package (Oksanen et al., 2007). PCoA was performed using the 'pcoa' function from the
ape package (Paradis et al., 2004). Procrustes analysis comparing the PCoA vectors of the two
data sets was done using the 'procrust' and 'protest' functions from the vegan package (Oksanen
et al., 2007). The Kruskal-Wallis test (McKnight and Najab, 2010) was applied to test for significant
correlations between the normalized average coverage of different ARGs and wastewater effluent
proportions (WW%), and p-values were corrected for multiple hypothesis testing using the
Benjamini-Hochberg method (Benjamini and Hochberg, 1995).

ARG identification and coverage analysis from metagenome assemblies

For a complementary and separate approach to the metagenomic short reads pipeline for ARG analysis, predicted open reading frames (ORFs) from assembled metagenomes were identified by protein BLAST v2.9.0 (McGinnis and Madden, 2004) search against the Structured Antibiotic Resistance Genes (SARG) protein database v2.2, with e-value and query coverage cut-offs of 0.1 and 20, respectively (Yin et al., 2018). Using methods established by (Lee et al., 2021) for detecting ARGs in riverine metagenomes, BLAST hits with an amino acid percent identity of over 85% and an alignment length of 100 amino acids were retained. BLAST hits were then clustered using CD-HIT v4.6.8 (Fu et al., 2012; Li and Godzik, 2006), with parameters –n 5 and –c 0.7. The 102 non-singleton cluster representatives of SARG hits were selected for further analysis. Metagenomic coverage of selected (clustered) BLAST hits within all metagenomes were calculated based on methods previously described (Lee et al., 2021). Briefly, raw reads were mapped back to BLAST hits using bowtie2 v2.4.4 (Langmead and Salzberg, 2012) using the parameters shown in **Table S2**. Coverage depth was calculated using samtools v1.12 and average coverage was calculated as described by (Albertsen et al., 2013). The average coverages of the 102 ORFs were normalized to reads per million using the method described by (Lee et al., 2021).

Bioinformatic analyses of beta-lactamases

Homologs of Planctomycetota ARGs were identified by BLAST v.2.9.0 (Camacho et al., 2009) querying assembled proteins from the 75 metagenomes. Trends across different WW and UF WW conditions were examined leading to prioritization of two metallo beta-lactamase homologs as described in the results (NCBI accession numbers MBX3358097.1 and MBX3359331.1) assigned to the *Phycisphaeraceae* (Planctomycetota). Structural modeling of MBX3358097.1 was conducted using ColabFold AlphaFold 2.0 (Mirdita et al., 2022) with default parameters. Models were assessed for quality based on predicted local distance difference test (pLDDT) scores and aligned pairwise with the reference structure in the Protein Data Bank (PDB ID: 3NKS) using the PyMOL 'super' command with default parameters.

Cloning and protein expression

For functional characterization, two full-length proteins (NCBI accession numbers MBX3358097.1 and MBX3359331.1) were codon-optimized for *E. coli* codon usage using the Integrated DNA Technologies Codon Optimization Tool (https://eu.idtdna.com/pages/tools/). Signal peptides were detected using SignalP v6.0 (Teufel et al., 2022) and removed where present. Genes (**Table S3**) were synthesized as gBlocks by Integrated DNA Technologies. The genes were cloned individually by Gibson Assembly into the multiple cloning site 1 of the pETDuet-1 vector (*ampR* resistance marker) with N-terminal hexahistidine tags (**Table S3**). Constructs were transformed into *E. coli* DH5α cells and verified by Sanger sequencing (Microsynth, Balgach, Switzerland). Sequence-verified constructs were transformed into *E. coli* BL21(DE3) cells (New England Biolabs Frankfurt, Germany). Starter cultures (5 mL) were grown in LB with 100 µg/ml ampicillin overnight at 37°C. From the starter cultures, 2.5 mL was added to 400 mL LB with 100 µg/mL ampicillin in a 1 L baffled Erlenmeyer flask and grown to an

205 optical density of 0.5-0.8 at 37°C. Protein expression was induced by adding 0.1 mM isopropyl
206 β-D-1-thiogalactopyranoside (IPTG) and cultures were incubated for 18h at 16°C.

Protein purification

Expression cultures were poured into 50 mL Falcon tubes and centrifuged at 4000 rcf for 15 min.
Supernatants were decanted, and cell pellets were resuspended in 4 mL buffer A (20 mM Tris,
500 mM NaCl, 20 mM imidazole, 10% glycerol, pH = 8) per gram of pellet weight. Cells were lysed
by 2 cycles of sonication using a 6 mm needle, 20% amplitude and 10 x 10 second pulses with a
rest time of 10 seconds in between (total sonication time per cycle = 3 minutes). Lysates were
centrifuged at 20,000 rcf at 4°C for 1h. Cleared lysate was filtered through a 0.2 µm syringe filter,
loaded into a ÄKTA Pure system (Cytiva Europe GmbH, Glattbrugg, Switzerland) and injected
onto a HisTrap FF 5 mL column (Cytiva, Europe GmbH, Glattbrugg, Switzerland). The column
was washed with 20 column volumes of buffer A. His-tagged proteins were eluted in 3 column
volumes of buffer B (same as buffer A but with an imidazole concentration of 500mM), and 8 x 2
mL fractions were collected. The fractions with the highest protein concentrations were
determined qualitatively by Bradford assay (Bradford, 1976). Imidazole was removed by running
the sample over a PD-10 desalting column (Cytiva Europe GmbH, Glattbrugg, Switzerland) and
eluting in buffer (5 mM Tris, 30 mM NaCl, 10% glycerol, 0.1 mM $ZnCl_2$, pH = 8). $ZnCl_2$ was added
because the enzyme of interest is suspected to be Zn^{2+}-dependent. The purified protein was flash
frozen using liquid nitrogen in 200µl aliquots and subsequently stored at -80°C.

β-Lactamase activity assay

Purified MBX3358097.1 was tested with standard β-lactamase activity detection assays using the
chromogenic cephalosporin analog, nitrocefin (O'Callaghan et al., 1972). Nitrocefin was prepared
in a 40 mM stock solution in DMSO and then diluted in Tris buffer to prepare a 1 mM working
solution. Briefly, standard nitrocefin assays were run in a 96-well plate by combining 180µl 20 mM
Tris buffer with 0.1 mM $ZnCl_2$ (pH = 7.0) with 10 µL purified enzyme and 10µL nitrocefin working
solution. The plate was incubated in a plate reader (BioTek Synergy H1, Agilent) at 25°C, and
absorbance at 482 nm was measured every minute for 2h. As negative controls, boiled enzyme
and buffer-only conditions were included in every assay. To obtain the boiled enzyme control,
aliquots of purified protein were incubated at 100°C for 15 minutes. Reaction rates were quantified
from continuous measurements as described previously by Ryu et al. using the molar extinction
coefficient of nitrocefin (12,124 M^{-1} cm^{-1}) in 20 mM Tris (Ryu et al., 2016).

Results and discussion

Biotic factors appear to be predominantly responsible for altered community composition in biofilms grown in WWTP effluent

We deeply sequenced metagenomes of 75 total biofilm and water samples to an average
sequencing depth of 138 million reads per sample (max. 204 million reads). After assembly, we
first used the metagenomic data to analyze the taxonomic composition of stream biofilm
communities to assess whether community composition was correlated with experimental

treatments: 30% WW, 80% WW, 30% UF WW, 80% UF WW, and stream water (0% WW).
Experimental treatment was correlated with taxonomic composition for both prokaryotes
(PERMANOVA, R^2 = 0.65, p < 0.001) (**Figure 1**) and eukaryotes (PERMANOVA, R^2 = 0.70, p <
0.001) (**Figures S1-S3**) at the genus level (Salazar et al., 2021). Growth in 80% WW relative to
stream water (0% WW) altered prokaryotic biofilm community composition, namely we observed
a decrease in the relative abundance of *Cyanobacteria* and *Verrucomicrobiota* and an increase
in the relative abundance of *Bacteroidota* (**Figure 1**), in agreement with Carles et al. (2022). These
shifts also mirror changes in microbial communities previously described downstream of WWTPs
relative to those upstream (Matviichuk et al., 2022). Fold changes between UF and non-UF WW
biofilm samples, and in other lower abundance taxa, e.g., the Planctomycetota, are further
discussed by Carles et al. (2022).

The taxonomic composition of biofilms grown in UF wastewater is more similar to biofilms grown
in 0% WW than biofilms grown in the same fraction of non-UF wastewater, suggesting that biotic
factors from WWTPs are responsible for community changes as opposed to abiotic factors
(**Figure 2, Figure S3**). This metagenomic analysis highlights that WW alters stream biofilm
composition and supports the findings of Carles et al. (2022) that some microorganisms from WW
were able to colonize and modify biofilm community composition.

Resistome composition correlates with microbiome composition
We next examined how the total set of ARGs detected in each sample - the 'resistome' - varied
with experimental treatment (e.g., WW% and non-UF vs. UF). Resistome composition was
significantly affected by experimental treatment (PERMANOVA, R^2 = 0.60, p < 0.001), much like
microbiome composition (**Figure S4**). Biofilms exposed to 30% or 80% UF WW clustered with 0%
WW samples as opposed to samples exposed to equivalent amounts of non-UF WW (**Figure 2**).

To test for a relationship between microbiome and resistome composition, we used Procrustes
superimposition of resistome PCoA vectors onto prokaryotic microbiome PCoA vectors (**Figure
2**). Procrustes analysis revealed that the composition of the resistome was structurally related to
the composition of the prokaryotic microbiome. The low Procrustes sum of squares was 0.1889
which indicated a strong alignment together with the calculated correlation value of 0.9 in a
symmetric Procrustes rotation (p-value < 0.001). The alignment between microbiome and
resistome has been previously observed in microbial communities downstream of WWTPs (Lee
et al., 2023; Matviichuk et al., 2022), but local environmental features of different sampling sites
are potential confounding factors that may have determined both microbiome and resistome
composition in the field. Here we demonstrated a microbiome-resistome relationship in a
controlled experiment with biofilms grown in a defined gradient of stream water to WW and
ultrafiltered WW matrices. Generally, the rotated PCoA vectors for the resistome are closer to the
prokaryotic microbiome PCoA vectors for biofilm samples than they are for water samples (**Figure
2**). These results further suggest that changes in resistome composition in WW-impacted stream
biofilms are likely due to changes in colonization of the biofilm communities by WWTP bacteria.

Figure 1. Mean relative abundance of prokaryotic phyla in metagenomes of stream biofilms grown with different percentages of treated wastewater (WW) with or without ultrafiltration (UF) treatment. Taxa which were in the upper decile of abundance in at least one experimental treatment were retained while the remaining phyla were binned as 'other'.

Figure 2. PCoA-based Procrustes analysis showing a structural alignment between microbiome and resistome composition. Points indicate position on PCoA axes 1 and 2 of prokaryotic composition (microbiome), while arrow tips indicate position on PCoA axes 1 and 2 of resistome composition following Procrustes rotation. The correlation coefficient in a symmetric Procrustes rotation was 0.901, the sum of squares (m12 squared) was 0.1889), and p-value was less than 0.001.

WW ultrafiltration treatment has ARG class-specific effects on biofilm resistomes

A total of 570 ARGs across 41 different higher-level ARG classes were identified in the biofilm metagenomes using a short reads pipeline for ARG detection (Arango-Argoty et al., 2018). In biofilms, the abundances of ARGs conferring multidrug resistance, in addition to ARGs conferring resistance to beta-lactams, fluoroquinolones, peptides, rifamycins, and tetracyclines, differed significantly between conditions **(Table S5)**. The normalized read counts of several, but not all, ARG classes increased with increasing WW%, as seen in **Figure 3**. In biofilms grown in comparable percentages of UF WW, the abundance of ARGs from the abovementioned classes did not increase (**Figure 3**). The UF treatment results thus further support that biotic factors in WW increased the abundance of many ARG classes in biofilms downstream.

As a notable exception, sulfonamide ARGs showed increasing abundance with higher WW% yet displayed the highest abundance in 80% WW UF conditions. The unusual abundance in UF-grown biofilms suggests that in addition to being impacted by biotic factors, abiotic factors such as micropollutants (MPs) and nutrients may play a role in sulfonamide ARG abundance. Additionally, ARG abundance in UF-grown biofilms may relate to gene-family specific mechanisms such as horizontal transfer which may occur through transient contact or free DNA passing through the ultrafiltration membrane. The sulfonamide resistance, ARGs *sul1-4,* are known to be "highly mobilized" in WWTP effluent microbiomes (Ju et al., 2019). Although many ARGs are mobile, *sul1* ARGs, in particular, are notable for their tight association with the class 1 integrons (Lee et al., 2021) which are among the most common multiresistance cassette systems found in biological systems (Gillings et al., 2008, 2015). In support of the mobility of the sulfonamide resistance genes, *sul1* genes detected in our metagenomes were co-localized on the same metagenomic contigs as mobile genetic elements including *int1* integron-integrases as well as recombinases and transposases (**Figure S5**). Notably, the presence of *intl1* genes in environmental samples has been suggested as a gene proxy to estimate levels of anthropogenic pollution including MPs (as proposed by (Gillings et al., 2015). Our results further support this relationship between *int1* genes, sulfonamide ARGs typically associated with *int1,* and abiotic factors including pollutants (Gillings et al., 2008).

328
Figure 3. Abundance (16S-normalized read counts) of the DeepARG hits assigned to ARG
classes which varied significantly with adjusted p-values ≤ 0.001 (Kruskal-Wallis test + Benjamini-
Hochberg multiple hypothesis correction) between different treatment conditions. MLS =
macrolide-lincosamide-streptogramin.

Association between assembled ARG abundances and WW percentage

To obtain more detailed information on the sequence identity of abundant ARGs in the biofilm
communities, we assembled the metagenomes into contigs and used stringent criteria (see

Methods) to query called ORFs against the Structural Antibiotic Resistance Genes (SARG)
database. Assembly-based approaches are likely to have lower sensitivity but higher specificity
than short read-based approaches (Tamames et al., 2019, Peng et al., 2021). The results of
querying 75 assembled metagenomes against the SARG database yielded 2,240 ARG hits. We
clustered ARGs at the 70% amino acid sequence similarity level to obtain 102 ARG 'cluster
representatives.' Annotating the cluster representatives by ARG class resulted in 15 beta-lactam,
14 multidrug, 9 aminoglycoside, 5 tetracycline, 3 macrolide-lincosamide-streptogramin (MLS), 2
sulfonamide, 1 spectinomycin, and 1 vancomycin ARG clusters, in addition to 51 other cluster
representatives that were classified by SARG into the 'other' ARG category.

Figure 4. Corrected normalized average coverage of ARG cluster representatives identified using
an assembly-based ARG detection method. ARG clusters with significant differences in
abundances between treatment conditions are displayed (adjusted p-values ≤ 0.05 using Kruskal-
Wallis test + Benjamini-Hochberg multiple hypothesis correction).

To quantify relative ARG abundances across treatment conditions, we mapped raw reads back
onto the 102 ARG cluster representatives. Ten ARG cluster representatives differed significantly
(Kruskal-Wallis test, $p < 0.05$) in abundance between at least one experimental treatment
condition (**Table S6**). One each of the multidrug, MLS, and aminoglycoside class ARG
representatives, in addition to two beta-lactam resistance ARGs, displayed the trend of increasing
abundance with increasing WW% previously found for most of the ARGs identified with the short-
reads pipeline. Two ARG classes encoding for beta-lactam resistance displayed the greatest
increase in abundance with %WW (Class-A beta-lactamase gene "beta-lactam 1", $p = 0.0054$ and

a Class-D beta-lactamase bla OXA-4 gene "beta-lactam 2", p = 0.0054, **Figure 4**). These ARGs
have significant increases in average coverage in biofilms with increasing WWTP effluent. In
contrast, a third beta-lactam ARG (extended-spectrum beta-lactamase BEL-2 "beta-lactam 3" p-
value = 0.0488) displayed the highest average coverage exclusively in the 80% UF WW condition.
These results suggest the abundance of this cluster of beta-lactamase ARGs may have been
affected mainly by abiotic factors such as MPs in the UF WW-grown biofilms, the effects of which
may have been outcompeted by biotic effects in the non-UF WW grown-biofilms.

Our combined results from short read-based and assembly-based ARG analysis generally aligned
with the exception of the sulfonamide ARGs which did not show a significant difference in the
assembly-based approach. This may be due to limitations or artifacts of the metagenomic
assembly. The significance of beta-lactamase ARGs in the short read-based approach was also
captured in the assembly-based approach. Overall, our results also align well with previous
studies indicating that exposure to WWTP effluent alters resistome composition and increases
ARG abundance in surface waters in Switzerland (Lee et al., 2023) and globally (Czekalski et al.,
2015; Matviichuk et al., 2022; Mukherjee et al., 2021; Thornton et al., 2020). Beta-lactamase
ARGs were also previously shown to increase in abundance in microbial communities exposed
to WWTP effluent (Fadare and Okoh, 2021; Mughini-Gras et al., 2019) and are consistently the
most abundant class of ARGs detected in surface waters (Laht et al., 2014; Osińska et al., 2020;
Rodriguez-Mozaz et al., 2015; Stoll et al., 2012). The prevalence is attributed to a variety of
factors, including the early discovery and continued widespread overuse of beta-lactam antibiotics
for humans and animals (Agarwal et al., 2023; Sta Ana et al., 2021). Moreover, one of the beta-
lactam clusters ("beta-lactam 3") in our dataset showed an unexpectedly high abundance only in
80% WW UF (**Figure 4)**, making beta-lactamases a target in this study for further investigation.

Micropollutants as an abiotic factor which may influence ARG transfer

We next examined MP concentrations as abiotic factors which may influence gene-family specific
ARG abundances, such as the *sul* and "beta-lactam 3" ARG clusters which showed unexpectedly
higher abundances in the UF WW over non-UF conditions (**Figure 3, Figure 4)**. Based on
previous chemical analysis by liquid chromatography-tandem mass spectrometry (Desiante et al.,
2022), MP concentrations measured in the flume biofilm samples increased with increasing WW%
(**Table S4**), but MP concentrations did not significantly differ between the UF and non-UF WW-
grown biofilms. Other antibiotics targeted, including sulfamethoxazole and sulfapyridine (**Table
S4**), were below or near the limit of quantification in WW-grown biofilms, preventing further
analysis. The macrolide antibiotic, clarithromycin, had the highest concentrations in 80% WW and
80% UF WW grown biofilms (**Table S4**). Clarithromycin concentrations increased with WW% and
did not vary significantly between UF and non-UF treatments. In contrast, the metagenomic
abundances of macrolide-lincosamide-streptogramin (MLS) genes - the ARGs typically conferring
resistance to clarithromycin - were significantly lower in UF compared to non-UF conditions
(**Figure 4**). This example represents a case of an antibiotic-ARG pair where ARG abundances
did not correlate with the expected abiotic factor (antibiotic concentration). However, this does not
preclude that other MPs which are not classified as antibiotics may have influenced MLS ARG
abundance.

A growing body of evidence supports that many anthropogenic pollutants which do not have
known antimicrobial bioactivities nonetheless influence ARG transfer and proliferation (Alderton
et al., 2021; Kimbell et al., 2023; Qiu et al., 2022; Wang et al., 2021; Yu and Guo, 2022; H. Zhang
et al., 2023). However, there is generally a lack of knowledge about how such 'non-antibiotic' MPs
may still be selective agents for ARG abundance. Previously, 75 different MPs were measured in
the flume biofilms, stream water, and WW (Desiante et al., 2022) including artificial sweeteners
(sugar replacements, e.g., acesulfame-K, saccharin and cyclamate), non-antibiotic
pharmaceuticals (e.g., antihypertensives), corrosion inhibitors, and pesticides (**Table S4)**.
Artificial sweeteners were noteworthy as the most abundant MPs detected in µg/L concentrations
in WW samples (Desiante et al., 2022) and exhibiting a pattern of increasing concentrations with
increasing WW% (**Table S4**). Notably, the artificial sweeteners measured here (acesulfame K,
cyclamate, and saccharin) all contain sulfonamide moieties, which have a chemical resemblance
to the moieties found in sulfonamide antibiotics (**Figure S6**). Sulfonamides were also one of the
few ARG classes with significantly higher abundances in UF vs. non-UF WW-grown biofilms
(**Figure 3**). These observations led us to speculate that abiotic factors, such as sulfonamide-
containing MPs, may play a role in promoting gene-family specific ARG abundances. In support
of this, several experimental studies have shown that artificial sweeteners promote high rates of
horizontal gene transfer and natural uptake of ARGs in bacteria (Yu et al., 2022; Yu and Guo,
2022). Further work has demonstrated that acesulfame-K and other artificial sweeteners directly
enhance the rate and quantity of horizontal plasmid-mediated transfers of *sul* genes (Li et al.,
2022). Interestingly, a plasmid-encoded microbial hydrolase capable of cleaving the sulfonamide
moiety of acesulfame-K was recently identified and shown to have a beta-lactamase-like
structural fold (Bonatelli et al., 2023). Taken together, these observations prompted us to conduct
functional investigation of beta-lactamase-fold genes that increased in abundance with WW% and
sulfonamide-containing MPs.

Planctomycetota beta-lactamases display differential abundances in response to sulfonamide-containing MPs and WWTP effluent

To investigate the relationship between ARG abundances and sulfonamide-containing MPs, we
consulted previous reports of enrichment studies of activated sludge communities grown on
sulfonamide-containing MPs including artificial sweeteners (Deng et al., 2019; Huang et al.,
2021). Multiple independent enrichment studies on either acesulfame-K (Huang et al., 2021),
saccharin, or cyclamate (Deng et al., 2019) all reported a significant increase in the relative
abundance of bacteria belonging to the phylum Planctomycetota and particularly the
planctomycete family *Phycisphaeraceae*. The *Phycisphaeraceae* are known to colonize the
surfaces of aquatic macroalgae (Rambo et al., 2020) and share a common evolutionary ancestor
with the anaerobic ammonia-oxidizing (anammox) planctomycetes promising in wastewater
treatment processes (Andrei et al., 2019; Wu et al., 2022). Previously, Desiante et al. (2022) also
identified significant differences in relative abundances of *Phycisphaeraceae* in biofilms grown in
80% WW and 80% WW UF in two independent flume biofilm experiments (log$_2$ fold-change > 4;
p < 0.05). Through differential abundance analysis conducted here, the Planctomycetota also
emerged as a phylum with statistically-signiTficant differences in relative abundances between

WW and UF WW treatments (**Figure 5A**). In terms of their ecological roles, taxa belonging to the
phylum Planctomycetota compose up to 2.1% relative abundance of the biofilm, WW, and stream
water metagenomes, confirming previous reports of the Planctomycetota as a numerically-
relevant member of both freshwater and wastewater microbial communities (Andrei et al., 2019;
Cai et al., 2021). Based on metagenomic analysis, algae-associated *Phycisphaeraceae* are likely
to impact the fitness of their algal hosts (Rambo et al., 2020) e.g., in stream biofilms, but the
current unculturability of most symbiotic microorganisms in the *Phycisphaeraceae* limits our
knowledge of their ARG phenotypes. We therefore targeted *Phycisphaeraceae* ARGs for
functional characterization.

Due to challenges in the laboratory cultivation of the *Phycisphaeraceae* (Rambo et al., 2020), we
opted for a heterologous expression approach to functionally validate *Phycisphaeraceae* ARGs.
We focused on beta-lactamases for four reasons: (1) the beta-lactamases were identified in our
ARG analysis as the most abundant and significantly-different class of ARGs affected by the
percentage of treated wastewater, (2) beta-lactamase fold enzymes were also previously
identified as plasmid-encoded enzymes active in the biotransformation the sulfonamide MP
acesulfame-K (Bonatelli et al., 2023), revealing the 'structural overlap' of a versatile beta-
lactamase fold capable of cleaving both sulfonamide- and beta-lactam moieties, and (3) previous
evidence of sulfonamide cleavage in the sweetener cyclamate was shown by enzymes which
were metal-dependent hydrolases but their sequences were not determined (Nimura et al., 1974).
Finally, (4), full-length homologs with >50% amino acid identity to the sulfonamide resistance
genes *sul1-4* encoding drug resistant variants of the sulfonamide target dihydropteroate synthase
were not detected in the *Phycisphaeraceae*. For these reasons, we hypothesized that metallo-
beta-lactamase-fold enzymes may play an overlooked role in sulfonamide resistance in these
planctomycetes.

Functional characterization of a new beta-lactamase ARG from Planctomycetota
To identify metallo-beta-lactamase fold enzymes assigned to the *Phycisphaeraceae*, we queried
the 75 metagenomes sequenced in this study as well as metagenome assembled genomes
(MAGs) from previous sulfonamide-containing MP enrichment studies (Deng et al., 2019; Huang
et al., 2021). We identified two full-length metallo beta-lactamase fold protein hits from the
Phycisphaeraceae which showed a trend of increasing abundance patterns with percentage of
wastewater relative to UF controls (**Figure 5B**). Both shared low similarity (less than 30% amino
acid identity and between 20-50% query cover) with known beta-lactamases. We heterologously
expressed each of the *Phycisphaeraceae* genes in *E. coli*. One protein did not express in *E. coli*
and despite optimization could not be produced or purified to homogeneity with nickel-affinity
chromatography. The second gene (MBX3358097.1) expressed sufficiently for purification with
an average yield of 500 μg/mL pure protein (**Figure S7**), therefore we proceeded with the
functional analysis of this protein.

Figure 5. A) Volcano plot displaying $\log_2$-fold change of differentially abundant taxa between WW30UF and WW30. Similar significant trends, although with a lower fold-change difference, were observed in the Planctomycetota in WW80UF vs. WW80 samples. B) Counts of hits showing significant homology to the putative beta-lactamase MBX3358097.1 in biofilm metagenomes. C) Purified enzyme assays demonstrated beta-lactamase activity of MBX3358097.1 with the colorimetric cephalosporin analog, nitrocefin, compared to boiled enzyme and enzyme buffer only controls. D) Two different views of a high-confidence AlphaFold2 model (mean pLDDT = 96.5) of MBX3358097.1 structurally-aligned with its closest structural homolog in the Protein Data Bank, the *Bacillus cereus* beta-lactamase (PDB ID: 3KNS), identified by FoldSeek (van Kempen et al., 2023) as of 20.11.2023.

Next, we tested whether the purified *Phycisphaeraceae* protein encoded a functional beta-lactamase despite sharing low sequence similarity to known beta-lactamases. The protein cleaved the cephalosporin analog, nitrocefin, at significant levels above the boiled enzyme and buffer control (**Figure 5C**), indicating beta-lactamase activity. We next tested its substrate specificity for sulfonamide containing compounds. Using HPLC analysis we determined that the *Phycisphaeraceae* enzyme did not degrade the sulfonamide-containing MPs acesulfame-K, saccharin, or cyclamate, nor did it degrade sulfamethoxazole (data not shown). Although this planctomycete-derived beta-lactamase was sequence divergent from known beta-lactamases (less than 22% amino acid identity) with known beta-lactamases in the PDB, AlphaFold2 models indicated the protein shared structural similarity (RMSD of 1.53 Å) with class B2 beta-lactamases (**Figure 5D**). This analysis therefore also exemplified the potential of AlphaFold2 to identify proteins with low sequence similarity yet higher structural similarity performing conserved

functions such as beta-lactam cleavage. More broadly, this result highlights how sequence-
structure-function analysis can be used to mine microbial functions of interest from metagenomes
(Robinson, 2023; Ruppé et al., 2019).

There are few previous reports of ARGs from the phylum Planctomycetota (Ivanova et al., 2021).
Planctomycetes were previously reported to have an unusual proteinaceous cell wall composition
devoid of peptidoglycan (König et al., 1984), however later findings demonstrated planctomycetes
do indeed produce peptidoglycan and are thus susceptible to cell-wall targeting antibiotics
including beta-lactams (Jeske et al., 2015). To our knowledge, this is the first biochemical
characterization of an active metallo-beta-lactamase from the *Phycisphaeraceae*. This ARG was
also not found by current computational ARG detection tools, including the SARG database used
in this study (Yin et al., 2018) due to its low sequence identity with known ARGs. This suggests
that functional antibiotic resistance determinants, particularly from understudied microbial taxa,
may be overlooked by state-of-the-art ARG detection methods. A knowledge gap remains
whether such sequence-diverse ARGs from the Planctomycetota and other difficult-to-culture
bacteria make significant contributions to environmental ARG reservoirs.

525   Conclusions

Our results from a controlled flume system showed that WWTP effluent altered both the
microbiome and resistome composition of stream biofilms. Through ultrafiltration treatment of
WWTP effluent, we could attribute changes in taxonomic and resistome composition primarily to
biotic rather than abiotic factors. Comparing the abundances of ARG classes in UF vs non-UF
conditions revealed that many ARGs had significantly higher normalized metagenomic
abundances in non-UF WW as opposed to UF WW conditions which can be attributed to the
introduction new microbial taxa from the WWTP effluents (biotic factors). Conversely, some
sulfonamide and beta-lactamase ARGs were more abundant in biofilms grown in UF WW. We
speculated this may be linked to the presence of sulfonamide-containing MPs that were not
removed by ultrafiltration, such as artificial sweeteners present at high concentrations in WW. We
therefore aimed to exemplify a metagenome mining approach to test hypotheses emerging from
relative abundance observations since previous work had shown that the planctomycete
Phycisphaeraceae were enriched with sulfonamide-containing sweeteners (Deng et al., 2019;
Desiante et al., 2022; Huang et al., 2021). Here, we demonstrated that a sequence-divergent
beta-lactam resistance gene, reconstructed from our metagenomes and originating from
Phycisphaeraceae, was not involved in sulfonamide artificial sweetener degradation but shows
beta-lactamase activity. This highlights the need to characterize new ARGs from understudied
taxa to create a more comprehensive ARG catalog for querying complex microbial communities
such as stream biofilms. Overall, this study investigated the interplay between MPs, ARGs, and
their microbial hosts, and exemplified new functional workflows to shed light on antibiotic
resistance determinants in aquatic environments.

Data availability

The metagenomes have been deposited in the NCBI Sequence Read Archive under BioProject: PRJNA733009. Scripts used for analysis and figures are available at: https://github.com/MSM-group/AMR-ecoimpact-paper. Data and supplemental figures are also available in the Supplemental Information associated with this manuscript.

CRediT authorship contribution statement

Conceptualization: Mustafa Attrah, Milo Schärer, Serina L. Robinson, Kathrin Fenner, Helmut Bürgmann. Methodology: Mustafa Attrah, Mauro Esposito, Milo Schärer, Giulia Gionchetta, Jack van de Vossenberg, Serina L. Robinson. Investigation: Mustafa Attrah, Milo Schärer, Mauro Esposito, Serina L. Robinson. Writing - Original draft: Mustafa Attrah, Milo Schärer, Serina L. Robinson, Jack van de Vossenberg. Writing - Review & editing: All authors. Visualization: Milo Schärer, Mustafa Attrah, Serina L. Robinson. Project administration: Serina L. Robinson, Piet Lens, Jack van de Vossenberg. Funding acquisition: Mustafa Attrah, Kathrin Fenner, Serina L. Robinson

Acknowledgements

We acknowledge Robert Niederdorfer for his support and expertise in the metagenomic assembly. We acknowledge Niklas Ferenc Trottman for his training and support for the cloning, expression, and protein purification. We thank the Eawag Partnership Program together with the Institute of Water Education (IHE Delft) under the auspices of UNESCO for supporting the exchange of Mustafa Attrah to Eawag. We thank Elia Ceppi for assistance in interpretation of chemical micropollutant data and Thierry Marti for support in biological interpretation. We thank Nik Zemp and Aria Minder from the Genetic Diversity Center at ETH Zürich for their computational guidance. We acknowledge Stuart Dennis for his computational infrastructure support. We acknowledge the EcoImpact teams 1.0 and 2.0 for fruitful discussions as well as conceptualizing and running the flume experiments which provided the samples necessary for this study.

Declaration of competing interests

None

Funding

S.L.R. acknowledges support from the Swiss National Science Foundation (grant no. PZPGP2_209124), the Vontobel Foundation, and the Pierre Mercier Foundation. K.F. acknowledges the Swiss National Science Funding grant no. 200021L_201006. H. Bürgmann and G. Gionchetta were supported by the ANTIVERSA project (BiodivERsA, European Union), Swiss National Science Foundation (Switzerland) grant no. 186531. Student funding for an advanced class at IHE Delft Institute of Water Education was provided to M.A. through IHE Delft's DUPC2 program on the project 'Supporting integrated and sustainable water management in Iraq through capacity development and research' grant no.109070.

References

Agarwal, V., Tiwari, A., Varadwaj, P., 2023. An extensive review on β-lactamase enzymes and
their inhibitors. Curr. Med. Chem. 30, 783–808.
Albertsen, M., Hugenholtz, P., Skarshewski, A., Nielsen, K.L., Tyson, G.W., Nielsen, P.H., 2013.
Genome sequences of rare, uncultured bacteria obtained by differential coverage binning of
multiple metagenomes. Nat. Biotechnol. 31, 533–538.
Alderton, I., Palmer, B.R., Heinemann, J.A., Pattis, I., Weaver, L., Gutiérrez-Ginés, M.J.,
Horswell, J., Tremblay, L.A., 2021. The role of emerging organic contaminants in the
development of antimicrobial resistance. Emerging Contaminants 7, 160–171.
Alexander, J., Hembach, N., Schwartz, T., 2020. Evaluation of antibiotic resistance
dissemination by wastewater treatment plant effluents with different catchment areas in
Germany. Sci. Rep. 10, 8952.
Andrei, A.-Ş., Salcher, M.M., Mehrshad, M., Rychtecký, P., Znachor, P., Ghai, R., 2019. Niche-
directed evolution modulates genome architecture in freshwater *Planctomycetes*. ISME J.
13, 1056–1071.
Arango-Argoty, G., Garner, E., Pruden, A., Heath, L.S., Vikesland, P., Zhang, L., 2018.
DeepARG: a deep learning approach for predicting antibiotic resistance genes from
metagenomic data. Microbiome 6, 23.
Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: A practical and powerful
approach to multiple testing. J. R. Stat. Soc. 57, 289–300.
Bonatelli, M.L., Rohwerder, T., Popp, D., Liu, Y., Akay, C., Schultz, C., Liao, K.-P., Ding, C.,
Reemtsma, T., Adrian, L., Kleinsteuber, S., 2023. Recently evolved combination of unique
sulfatase and amidase genes enables bacterial degradation of the wastewater
micropollutant acesulfame worldwide. Front. Microbiol. 14, 1223838.
Bradford, M.M., 1976. A Rapid and Sensitive Method for the Quantitation of Microgram
Quantities of Protein Utilizing the Principle of Protein-dye Binding.
Brumfield, K.D., Huq, A., Colwell, R.R., Olds, J.L., Leddy, M.B., 2020. Microbial resolution of
whole genome shotgun and 16S amplicon metagenomic sequencing using publicly
available NEON data. PLoS One 15, e0228899.
Cai, X., Yao, L., Sheng, Q., Jiang, L., Wang, T., Dahlgren, R.A., Deng, H., 2021. Influence of a
biofilm bioreactor on water quality and microbial communities in a hypereutrophic urban
river. Environ. Technol. 42, 1452–1460.
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L.,
2009. BLAST+: architecture and applications. BMC Bioinformatics 10, 421.
Carles, L., Wullschleger, S., Joss, A., Eggen, R.I.L., Schirmer, K., Schuwirth, N., Stamm, C.,
Tlili, A., 2022. Wastewater microorganisms impact microbial diversity and important
ecological functions of stream periphyton, Water Res. 225, 119119.
Carles, L., Wullschleger, S., Joss, A., Eggen, R.I.L., Schirmer, K., Schuwirth, N., Stamm, C.,
Tlili, A., 2021. Impact of wastewater on the microbial diversity of periphyton and its
tolerance to micropollutants in an engineered flow-through channel system. Water Res.
203, 117486.
Czekalski, N., Sigdel, R., Birtel, J., Matthews, B., Bürgmann, H., 2015. Does human activity
impact the natural antibiotic resistance background? Abundance of antibiotic resistance
genes in 21 Swiss lakes. Environ. Int. 81, 45–55.
Deng, Y., Wang, Y., Xia, Y., Zhang, A.N., Zhao, Y., Zhang, T., 2019. Genomic resolution of
bacterial populations in saccharin and cyclamate degradation. Sci. Total Environ. 658, 357–
366.
Depetris, A., Tagliavini, G., Peter, H., Kühl, M., Holzner, M., Battin, T.J., 2022. Biophysical
properties at patch scale shape the metabolism of biofilm landscapes. NPJ Biofilms
Microbiomes 8, 5.

Desiante, W.L., Carles, L., Wullschleger, S., Joss, A., Stamm, C., Fenner, K., 2022. Wastewater
microorganisms impact the micropollutant biotransformation potential of natural stream
biofilms. Water Res. 217, 118413.
Fadare, F.T., Okoh, A.I., 2021. The abundance of genes encoding ESBL, pAmpC and non-β-
lactam resistance in multidrug-resistant *Enterobacteriaceae* recovered from wastewater
effluents. Front. Environ. Sci. Eng. China 9. https://doi.org/10.3389/fenvs.2021.711950
Fu, L., Niu, B., Zhu, Z., Wu, S., Li, W., 2012. CD-HIT: accelerated for clustering the next-
generation sequencing data. Bioinformatics 28, 3150–3152.
Gillings, M., Boucher, Y., Labbate, M., Holmes, A., Krishnan, S., Holley, M., Stokes, H.W., 2008.
The evolution of class 1 integrons and the rise of antibiotic resistance. J. Bacteriol. 190,
5095–5100.
Gillings, M.R., Gaze, W.H., Pruden, A., Smalla, K., Tiedje, J.M., Zhu, Y.-G., 2015. Using the
class 1 integron-integrase gene as a proxy for anthropogenic pollution. ISME J. 9, 1269–
1279.
Haenelt, S., Richnow, H.-H., Müller, J.A., Musat, N., 2023. Antibiotic resistance indicator genes
in biofilm and planktonic microbial communities after wastewater discharge. Front.
Microbiol. 14, 1252870.
Huang, Y., Deng, Y., Law, J.C.-F., Yang, Y., Ding, J., Leung, K.S.-Y., Zhang, T., 2021.
Acesulfame aerobic biodegradation by enriched consortia and *Chelatococcus* spp.:
Kinetics, transformation products, and genomic characterization. Water Res. 202, 117454.
Hyatt, D., Chen, G.-L., Locascio, P.F., Land, M.L., Larimer, F.W., Hauser, L.J., 2010. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics
11, 119.
Ivanova, A.A., Miroshnikov, K.K., Oshkin, I.Y., 2021. Exploring antibiotic susceptibility,
resistome and mobilome structure of planctomycetes from *Gemmataceae* family. Sustain.
Sci. Pract. Policy 13, 5031.
Jeske, O., Schüler, M., Schumann, P., Schneider, A., Boedeker, C., Jogler, M., Bollschweiler,
D., Rohde, M., Mayer, C., Engelhardt, H., Spring, S., Jogler, C., 2015. *Planctomycetes* do
possess a peptidoglycan cell wall. Nat. Commun. 6, 7116.
Ju, F., Beck, K., Yin, X., Maccagnan, A., McArdell, C.S., Singer, H.P., Johnson, D.R., Zhang, T.,
Bürgmann, H., 2019. Wastewater treatment plant resistomes are shaped by bacterial
composition, genetic exchange, and upregulated expression in the effluent microbiomes.
ISME J. 13, 346–360.
Ju, F., Lee, J., Beck, K., Zhang, G., Gekenidis, M.-T., Hummerjohann, J., Bürgmann, H., 2022.
Phenotypic metagenomics tracks wastewater-associated clinically important beta-lactam
resistant bacteria invading river habitats. Research Square. https://doi.org/10.21203/rs.3.rs-
1589365/v1. PPR: PPR488559.
Kimbell, L.K., LaMartina, E.L., Kohls, S., Wang, Y., Newton, R.J., McNamara, P.J., 2023. Impact
of corrosion inhibitors on antibiotic resistance, metal resistance, and microbial communities
in drinking water. mSphere e0030723.
Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glöckner, F.O., 2013.
Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
generation sequencing-based diversity studies. Nucleic Acids Res. 41, e1.
König, E., Schlesner, H., Hirsch, P., 1984. Cell wall studies on budding bacteria of the
Planctomyces/Pasteuria group and on a *Prosthecomicrobium* sp. Arch. Microbiol. 138,
200–205.
Laht, M., Karkman, A., Voolaid, V., Ritz, C., Tenson, T., Virta, M., Kisand, V., 2014. Abundances
of tetracycline, sulphonamide and beta-lactam antibiotic resistance genes in conventional
wastewater treatment plants (WWTPs) with different waste load. PLoS One 9, e103705.
Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods
9, 357–359.

Lee, J., Ju, F., Beck, K., Bürgmann, H., 2023. Differential effects of wastewater treatment plant
effluents on the antibiotic resistomes of diverse river habitats. ISME J.
https://doi.org/10.1038/s41396-023-01506-w

Lee, J., Ju, F., Maile-Moskowitz, A., Beck, K., Maccagnan, A., McArdell, C.S., Dal Molin, M.,
Fenicia, F., Vikesland, P.J., Pruden, A., Stamm, C., Bürgmann, H., 2021. Unraveling the
riverine antibiotic resistome: The downstream fate of anthropogenic inputs. Water Res. 197,
117050.

Li, D., Liu, C.-M., Luo, R., Sadakane, K., Lam, T.-W., 2015. MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics 31, 1674–1676.

Li, W., Godzik, A., 2006. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. Bioinformatics 22, 1658–1659.

Li, Z., Gao, J., Guo, Y., Cui, Y., Wang, Y., Duan, W., Wu, Z., 2022. Enhancement of antibiotic
resistance dissemination by artificial sweetener acesulfame potassium: Insights from cell
membrane, enzyme, energy supply and transcriptomics. J. Hazard. Mater. 422, 126942.

Matviichuk, O., Mondamert, L., Geffroy, C., Gaschet, M., Dagot, C., Labanowski, J., 2022. River
Biofilms Microbiome and Resistome Responses to Wastewater Treatment Plant Effluents
Containing Antibiotics. Front. Microbiol. 13, 795206.

McGinnis, S., Madden, T.L., 2004. BLAST: at the core of a powerful and diverse set of
sequence analysis tools. Nucleic Acids Res. 32, W20–5.

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., Steinegger, M., 2022.
ColabFold: making protein folding accessible to all. Nat. Methods 19, 679–682.

Mughini-Gras, L., Dorado-García, A., van Duijkeren, E., van den Bunt, G., Dierikx, C.M.,
Bonten, M.J.M., Bootsma, M.C.J., Schmitt, H., Hald, T., Evers, E.G., de Koeijer, A., van
Pelt, W., Franz, E., Mevius, D.J., Heederik, D.J.J., ESBL Attribution Consortium, 2019.
Attributable sources of community-acquired carriage of Escherichia coli containing β-lactam
antibiotic resistance genes: a population-based modelling study. Lancet Planet Health 3,
e357–e369.

Mukherjee, M., Laird, E., Gentry, T.J., Brooks, J.P., Karthikeyan, R., 2021. Increased
Antimicrobial and Multidrug Resistance Downstream of Wastewater Treatment Plants in an
Urban Watershed. Front. Microbiol. 12, 657353.

Nimura, T., Tokieda, T., Yamaha, T., 1974. Partial purification and some properties of cyclamate
sulfamatase. J. Biochem. 75, 407–417.

O'Callaghan, C.H., Morris, A., Kirby, S.M., Shingler, A.H., 1972. Novel method for detection of
beta-lactamases by using a chromogenic cephalosporin substrate. Antimicrob. Agents
Chemother. 1, 283–288.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J., Suggests,
M., 2007. The vegan package. Community ecology package 10, 719.

Osińska, A., Korzeniewska, E., Harnisz, M., Felis, E., Bajkacz, S., Jachimowicz, P., Niestępski,
S., Konopka, I., 2020. Small-scale wastewater treatment plants as a source of the
dissemination of antibiotic resistance genes in the aquatic environment. J. Hazard. Mater.
381, 121221.

Oyarzúa, P., Bovio-Winkler, P., Etchebehere, C., Suárez-Ojeda, M.E., 2021. Microbial
communities in an anammox reactor treating municipal wastewater at mainstream
conditions: Practical implications of different molecular approaches. J. Environ. Chem. Eng.
9, 106622.

Paradis, E., Claude, J., Strimmer, K., 2004. APE: Analyses of Phylogenetics and Evolution in R
language. Bioinformatics 20, 289–290.

Peng, Z., Mao, Y., Zhang, N., Zhang, L., Wang, Z., Han, M., 2021. Utilizing metagenomic data
and bioinformatic tools for elucidating antibiotic resistance genes in environment. Front.
Environ. Sci. Eng. China 9. https://doi.org/10.3389/fenvs.2021.757365

Pereira, M.B., Wallroth, M., Jonsson, V., Kristiansson, E., 2018. Comparison of normalization
methods for the analysis of metagenomic gene abundance data. BMC Genomics 19, 274.
Perveen, S., Pablos, C., Reynolds, K., Stanley, S., Marugán, J., 2023. Growth and prevalence
of antibiotic-resistant bacteria in microplastic biofilm from wastewater treatment plant
effluents. Sci. Total Environ. 856, 159024.
Qiu, D., Ke, M., Zhang, Q., Zhang, F., Lu, T., Sun, L., Qian, H., 2022. Response of microbial
antibiotic resistance to pesticides: An emerging health threat. Sci. Total Environ. 850,
158057.
Rambo, I.M., Dombrowski, N., Constant, L., Erdner, D., Baker, B.J., 2020. Metabolic
relationships of uncultured bacteria associated with the microalgae *Gambierdiscus.*
Environ. Microbiol. 22, 1764–1783.
Reddington, K., Eccles, D., O'Grady, J., Drown, D.M., Hansen, L.H., Nielsen, T.K., Ducluzeau,
A.-L., Leggett, R.M., Heavens, D., Peel, N., Snutch, T.P., Bayega, A., Oikonomopoulos, S.,
Ragoussis, I., Barry, T., van der Helm, E., Jolic, D., Richardson, H., Jansen, H., Tyson,
J.R., Jain, M., Brown, B.L., 2020. Metagenomic analysis of planktonic riverine microbial
consortia using nanopore sequencing reveals insight into river microbe taxonomy and
function. Gigascience 9, giaa053. https://doi.org/10.1093/gigascience/giaa053
Rizzo, L., Manaia, C., Merlin, C., Schwartz, T., Dagot, C., Ploy, M.C., Michael, I., Fatta-
Kassinos, D., 2013. Urban wastewater treatment plants as hotspots for antibiotic resistant
bacteria and genes spread into the environment: a review. Sci. Total Environ. 447, 345–
360.
Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26, 139–140.
Robinson, M.D., Oshlack, A., 2010. A scaling normalization method for differential expression
analysis of RNA-seq data. Genome Biol. 11, R25.
Robinson, S.L., 2023. Structure-guided metagenome mining to tap microbial functional diversity.
Curr. Opin. Microbiol. 76, 102382.
Rodriguez-Mozaz, S., Chamorro, S., Marti, E., Huerta, B., Gros, M., Sànchez-Melsió, A.,
Borrego, C.M., Barceló, D., Balcázar, J.L., 2015. Occurrence of antibiotics and antibiotic
resistance genes in hospital and urban wastewaters and their impact on the receiving river.
Water Res. 69, 234–242.
Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open
source tool for metagenomics. PeerJ 4, e2584.
Rogowska, J., Cieszynska-Semenowicz, M., Ratajczyk, W., Wolska, L., 2020. Micropollutants in
treated wastewater. Ambio 49, 487–503.
Romero, F., Acuña, V., Font, C., Freixa, A., Sabater, S., 2019. Effects of multiple stressors on
river biofilms depend on the time scale. Sci. Rep. 9, 15810.
Ruppé, E., Ghozlane, A., Tap, J., Pons, N., Alvarez, A.-S., Maziers, N., Cuesta, T., Hernando-
Amado, S., Clares, I., Martínez, J.L., Coque, T.M., Baquero, F., Lanza, V.F., Máiz, L.,
Goulenok, T., de Lastours, V., Amor, N., Fantin, B., Wieder, I., Andremont, A., van Schaik,
W., Rogers, M., Zhang, X., Willems, R.J.L., de Brevern, A.G., Batto, J.-M., Blottière, H.M.,
Léonard, P., Léjard, V., Letur, A., Levenez, F., Weiszer, K., Haimet, F., Doré, J., Kennedy,
S.P., Ehrlich, S.D., 2019. Prediction of the intestinal resistome by a three-dimensional
structure-based method. Nat Microbiol 4, 112–123.
Ryu, B.H., Ngo, T.D., Yoo, W., Lee, S., Kim, B.-Y., Lee, E., Kim, K.K., Kim, T.D., 2016.
Biochemical and structural analysis of a novel esterase from *Caulobacter crescentus*
related to penicillin-binding protein (PBP). Sci. Rep. 6, 37978.
Salazar, G., Ruscheweyh, H.-J., Hildebrand, F., Acinas, S.G., Sunagawa, S., 2021. mTAGs:
Taxonomic profiling using degenerate consensus reference sequences of ribosomal RNA
genes. Bioinformatics 38, 270–272.
Shallcross, L.J., Davies, D.S.C., 2014. Antibiotic overuse: a key driver of antimicrobial

resistance. Br. J. Gen. Pract. 64, 604–605.

Singer, A.C., Shaw, H., Rhodes, V., Hart, A., 2016. Review of antimicrobial resistance in the
environment and its relevance to environmental regulators. Front. Microbiol. 7, 1728.

Sta Ana, K.M., Madriaga, J., Espino, M.P., 2021. β-Lactam antibiotics and antibiotic resistance
in Asian lakes and rivers: An overview of contamination, sources and detection methods.
Environ. Pollut. 275, 116624.

Stoll, C., Sidhu, J.P.S., Tiehm, A., Toze, S., 2012. Prevalence of clinically relevant antibiotic
resistance genes in surface water samples collected from Germany and Australia. Environ.
Sci. Technol. 46, 9716–9726.

Tamames, J., Cobo-Simón, M., Puente-Sánchez, F., 2019. Assessing the performance of
different approaches for functional and taxonomic annotation of metagenomes. BMC
Genomics 20, 960.

Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gíslason, M.H., Pihl, S.I., Tsirigos, K.D.,
Winther, O., Brunak, S., von Heijne, G., Nielsen, H., 2022. SignalP 6.0 predicts all five
types of signal peptides using protein language models. Nat. Biotechnol. 40, 1023–1025.

Thornton, C.N., Tanner, W.D., VanDerslice, J.A., Brazelton, W.J., 2020. Localized effect of
treated wastewater effluent on the resistome of an urban watershed. Gigascience 9.
https://doi.org/10.1093/gigascience/giaa125

Tremblay, J., Singh, K., Fern, A., Kirton, E.S., He, S., Woyke, T., Lee, J., Chen, F., Dangl, J.L.,
Tringe, S.G., 2015. Primer and platform effects on 16S rRNA tag sequencing. Front.
Microbiol. 6, 771.

van Kempen, M., Kim, S.S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C.L.M., Söding, J.,
Steinegger, M., 2023. Fast and accurate protein structure search with Foldseek. Nat.
Biotechnol. https://doi.org/10.1038/s41587-023-01773-0

Wang, Y., Lu, J., Zhang, S., Li, J., Mao, L., Yuan, Z., Bond, P.L., Guo, J., 2021. Non-antibiotic
pharmaceuticals promote the transmission of multidrug resistance plasmids through intra-
and intergenera conjugation. ISME J. 15, 2493–2508.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K.,
Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo,
K., Yutani, H., 2019. Welcome to the tidyverse. J. Open Source Softw. 4, 1686.

Wu, P., Chen, J., Garlapati, V.K., Zhang, X., Wani Victor Jenario, F., Li, X., Liu, W., Chen, C.,
Aminabhavi, T.M., Zhang, X., 2022. Novel insights into Anammox-based processes: A
critical review. Chem. Eng. J. 444, 136534.

Yin, X., Jiang, X.-T., Chai, B., Li, L., Yang, Y., Cole, J.R., Tiedje, J.M., Zhang, T., 2018. ARGs-
OAP v2.0 with an expanded SARG database and Hidden Markov Models for enhancement
characterization and quantification of antibiotic resistance genes in environmental
metagenomes. Bioinformatics 34, 2263–2270.

Yu, Z., Guo, J., 2022. Non-caloric artificial sweeteners exhibit antimicrobial activity against
bacteria and promote bacterial evolution of antibiotic tolerance. J. Hazard. Mater. 433,
128840.

Yu, Z., Wang, Y., Henderson, I.R., Guo, J., 2022. Artificial sweeteners stimulate horizontal
transfer of extracellular antibiotic resistance genes through natural transformation. ISME J.
16, 543–554.

Zhang, H., Song, J., Zheng, Z., Li, T., Shi, N., Han, Y., Zhang, L., Yu, Y., Fang, H., 2023.
Fungicide exposure accelerated horizontal transfer of antibiotic resistance genes via
plasmid-mediated conjugation. Water Res. 233, 119789.

Zhang, Y., Zhao, Z., Xu, H., Wang, L., Liu, R., Jia, X., 2023. Fate of antibiotic resistance genes
and bacteria in a coupled water-processing system with wastewater treatment plants and
constructed wetlands in coastal eco-industrial parks. Ecotoxicol. Environ. Saf. 252, 114606.