Transcriptional inhibition after irradiation occurs preferentially at highly expressed genes in a manner dependent on cell cycle progression

Zulong Chen[1,^], Xin Wang[2,3,^], Xinlei Gao[2,3], Nina Arslanovic[1], Kaifu Chen[2,3,*] and Jessica Tyler[1,*]

[1]Weill Cornell Medicine, Department of Pathology and Laboratory Medicine, New York, NY 10065, USA

[2]Basic and Translational Research Division, Department of Cardiology, Boston Children's Hospital, Boston, MA 02115, USA

[3]Department of Pediatrics, Harvard Medical School, Boston, MA 02115, USA

^Equal contribution

*Corresponding Authors: jet2021@med.cornell.edu, kaifu.chen@childrens.harvard.edu

Abstract

In response to DNA double strand damage, ongoing transcription is inhibited to facilitate accurate DNA repair while transcriptional recovery occurs after DNA repair is complete. However, the mechanisms at play and identity of the transcripts being regulated in this manner are unclear. In contrast to the situation following UV damage, we found that transcriptional recovery after ionizing radiation (IR) occurs in a manner independent of the HIRA histone chaperone. Sequencing of the nascent transcripts identified a programmed transcriptional response, where certain transcripts and pathways are rapidly downregulated after IR, while other transcripts and pathways are upregulated. Specifically, most of the loss of nascent transcripts occurring after IR is due to inhibition of transcriptional initiation of the highly transcribed histone genes and the rDNA. To identify factors responsible for transcriptional inhibition after IR in an unbiased manner, we performed a whole genome gRNA library CRISPR / Cas9 screen. Many of the top hits in our screen were factors required for protein neddylation. However, at short times after inhibition of neddylation, transcriptional inhibition still occurred after IR, even though neddylation was effectively inhibited. Persistent inhibition of neddylation blocked transcriptional inhibition after IR, and it also leads to cell cycle arrest. Indeed, we uncovered that many inhibitors and conditions that lead to cell cycle arrest in G_1 or G_2 phase also prevent transcriptional inhibition after IR. As such, it appears that transcriptional inhibition after IR occurs preferentially at highly expressed genes in cycling cells.

Introduction

DNA double strand breaks (DSB) are one of most deleterious types of DNA lesions. Failure to repair a single DSB can lead to loss of a chromosome arm or cell death and inaccurate repair can lead to changes such as insertions, deletions, and translocations. Accordingly, the cell has developed an intricate DNA damage response (Jackson and Bartek, 2009). In vertebrate cells, the DNA damage response is mediated through activation of three PI3-like kinases: ataxia telangiectasia mutated (ATM), Ataxia telangiectasia and Rad3-related (ATR) and DNA dependent protein kinase (DNA-PK) (Blackford and Jackson, 2017), which coordinates DNA repair and the DNA damage cell cycle checkpoint which arrests cells until DSBs are repaired.

Additionally, ATM and DNA-PK have been shown to inhibit transcription in response to DSBs in a manner that occurs so transiently that it can only be detected when examining the nascent, not bulk, transcripts (Pankotai and Soutoglou, 2013). The transcription resumes or "recovers" immediately after DSB repair (Pankotai and Soutoglou, 2013). This transient inhibition of transcription after DSBs was initially shown for RNA polymerase I (Pol I) transcripts, where ATM triggered the reduction of nascent ribosomal gene transcripts, shown by visualization of a labelled ribonucleotide analog within the nucleolus, after exposure to ionizing radiation (IR) (Kruhlak et al., 2007). Mechanistically, the DSBs triggered a reduction in Pol I initiation complex assembly and led to premature displacement of elongating Pol I from the rDNA genes (Kruhlak et al., 2007). Using a reporter that allowed for visualization of repair protein recruitment and local transcription within cells, it was subsequently shown that ATM also mediates the inhibition of RNA polymerase II (Pol II) transcriptional elongation of genes in the vicinity of I-SceI endonuclease induced DSBs (Shanbhag et al., 2010). This transcriptional

inhibition was partly dependent on the E3 ubiquitin ligases RNF8 and RNF168, whereas

transcriptional recovery depended on the USP16 enzyme that deubiquitylates histone H2A

(Shanbhag et al., 2010). Additional mechanistic analyses using this same system revealed that

ATM dependent phosphorylation of the ATP-dependent nucleosome remodeler PBAF is

required for local transcriptional inhibition of Pol II transcription flanking a DSB (Kakarougkas

et al., 2014), indicating that chromatin changes are also required for transcriptional inhibition in

response to DSBs. The purpose of local transcriptional inhibition is to allow efficient and

accurate DSB repair (Kakarougkas et al., 2014, Meisenberg et al., 2019). Polycomb group

proteins and cohesin have also been shown to be required for local transcriptional inhibition of

Pol II transcription flanking a DSB, although their role is unclear (Meisenberg et al., 2019,

Kakarougkas et al., 2014), further indicating that chromatin structure and potentially

chromosome architecture also regulate transcriptional inhibition in response to DSBs.

Somewhat surprisingly, a distinct mechanism has been reported for the inhibition of Pol II

transcription at genes containing a DSB induced by the endonuclease I-PpoI (Pankotai et al.,

2012). In this case, both Pol II initiation and elongation were reduced adjacent to the DSB, in a

manner dependent on DNA-PK and the proteasome (Pankotai et al., 2012). Mechanistically,

DNA-PK appeared to help recruit the E3 ubiquitin ligase WWP2 to DSBs, which then promoted

the proteosome-dependent eviction of Pol II (Caron et al., 2019). In the absence of WWP2, the

DNA repair machinery was not efficiently recruited, indicating again that the reason for

transcriptional inhibition *in cis* flanking a DSB is to promote DNA repair (Caron et al., 2019).

The papers examining transcriptional inhibition around DSBs induced by endonucleases

generally find the transient repression occurs locally or *in cis* to the DSB (Iannelli et al., 2017).

However, another study found induction of the same set of ~ 200 transcripts soon after

irradiation and endonuclease break induction that occurred in a manner dependent on ATM and

p53, while only 33 nascent transcripts were down regulated after DSB induction (Venkata

Narayanan et al., 2017). Yet another study found that more genes were repressed than induced

after inducing global DSBs with neocarzinostatin, and this occurred via p53 mediated down-

regulation of MYC (Porter et al., 2017). As such, there are contradictory findings in the field at

present. Furthermore, the mechanism of transcriptional recovery after DSB repair is far from

clear. In response to UV damage, global transcriptional inhibition and recovery occurs, and this

transcriptional recovery after UV repair is dependent on the histone variant H3.3 histone

chaperone HIRA (Bouvier et al., 2021, AdamPolo and Almouzni, 2013). Mechanistically, HIRA

functioned to repress the transcriptional repressor ATF3, in turn promoting transcriptional

recovery after UV repair (Bouvier et al., 2021).

In contrast to the studies to date that have only examined local transcription inhibition occurring

in cis after DSB damage, we sought to examine transient transcriptional inhibition after induction

of global DSBs by IR exposure, and the subsequent transcriptional recovery after DSB repair, via

fluorescent labelling of a ribonucleotide analog incorporated only into nascent transcripts. Unlike

the situation following UV repair, we do not find a role for HIRA in transcriptional recovery

after DSB repair. Our sequencing of the nascent transcripts after irradiation identified a

programmed transcriptional program where a larger number of protein-coding genes were

upregulated than downregulated. The genes that were immediately downregulated after IR

tended to be highly transcribed genes including the rRNAs and histones, while the upregulated

genes tended to be transcribed at a lower level. We developed a flow cytometry-based assay of

nascent transcripts and used it as the basis for a whole genome gRNA screen to identify factors

required for transcriptional inhibition after IR. In addition to finding ATM as being required for

inhibition of transcription after DSB induction, we found that depletion of factors leading to cell

cycle arrest also blocked transcriptional inhibition.

Results

HIRA independent transcriptional inhibition and recovery after ionizing radiation

To detect bulk changes in nascent transcripts after irradiation *in situ*, we added the uridine analog

ethynyl uridine (EU) to human U2OS cells for 30 minutes. The incorporated EU was detected by

click chemistry to a fluorescent azide followed by immunofluorescence microscopy (Jao and

Salic, 2008). A reduction in bulk nascent transcripts was apparent 30 minutes after exposure to

10 Gray ionizing radiation (IR), and the transcriptional recovery was already occurring two hours

after irradiation (Fig. 1A). Co-immunofluorescence with γH2AX showed that the DNA damage

signal was greatly reduced at the same time point after IR where transcriptional recovery

occurred (Figure 1-figure supplement 1), consistent with transcriptional recovery occurring after

DSB repair. Given that the histone variant H3.3 histone chaperone HIRA promotes

transcriptional recovery after UV repair (Bouvier et al., 2021, AdamPolo and Almouzni, 2013),

we tested whether that was also the case for transcriptional recovery after IR. We found that

shRNA depletion of HIRA (Figure 1-figure supplement 2) had no effect on transcriptional

inhibition nor recovery after IR (Fig. 1A). In agreement, depletion of transcripts encoded from

both H3.3 genes (Figure 1-figure supplement 2) had no effect on transcriptional inhibition or

transcriptional recovery after IR (Fig. 1B). As such, the requirement for HIRA for transcriptional

recovery differs following IR and UV exposure, suggesting differences in the mechanism of

these processes.

Establishment of a flow cytometry-based assay for transcriptional inhibition and recovery

after irradiation

Given that the read out of nascent EU labelled transcripts after irradiation is fluorescence, we

established a flow cytometry-based assay to allow us to screen for factors regulating

transcriptional inhibition and recovery after IR. We established this assay in a murine Abelson

virus transformed pre-B cell line (termed Abl pre-B cells) (Bredemeyer et al., 2006) which are

non-adherent and are stably transformed with doxycycline inducible Cas9 (Chen et al., 2021). By

flow cytometry analysis, effective incorporation of EU into nascent transcripts was apparent in a

manner dependent on ongoing transcription because it was inhibited by the global RNA

polymerase inhibitor Actinomycin D (Fig. 2A). The EU signal had 2 peaks (Fig. 2A) and we

asked whether this reflects cell cycle differences in the cells with higher and lower EU

incorporation into the nascent transcripts. We conducted cell cycle analysis by labeling newly

synthesized DNA with BrdU and staining DNA contents with FxCycle Violet at the same time as

using EU to label nascent transcripts. It was apparent that the peak with less EU was from the G_1

phase cells, while the peak with more EU was derived from S and G_2 phase cells (Fig. 2B). To

determine whether there was a detectable reduction in EU incorporation by flow cytometry after

irradiation, we irradiated cells, waited different lengths of times before EU labelling of nascent

RNA (Fig. 2C). The transcriptional inhibition after irradiation was clearly detectable by flow

cytometry as early as 15 minutes after IR, while transcriptional recovery was complete by 4

hours after IR (Fig. 2D). Also, the extent of transcriptional repression was similar regardless of

whether the IR dose was 2, 5 or 10 Gray (Fig. 2D, Figure 2-figure supplement 1). This is

consistent with the possibility that the reduction of nascent transcripts after IR is a programmed /

signaling response rather than due to proximity of the genes to the DSB, which would have led to

a dose response.

Bulk reduction of nascent transcripts after IR is mainly due to a decrease in rDNA and

histone gene transcription

To gain a better understanding of transcriptional inhibition after IR, we sought to identify the

genes whose transcription was being inhibited after IR. We isolated EU labelled nascent total

RNA transcripts 30 minutes after IR and prior to IR from two independent experiments (Figure

3-figure supplement 1) and sequenced the EU-RNA (Fig. 3A). Prior to isolation of the ER

labelled nascent RNA, we added equal amounts of the commercial ERCC spike-in to RNA from

the same number of cells. This enabled the subsequent normalization of the total read number

from the human genome to total reads from the ERCC control, to detect global changes between

samples (Chen et al., 2015). We observed that the total read count of nascent transcripts declined

after IR (Fig. 3B). Most of the read counts were due to rDNA transcripts, and the decline in bulk

transcripts after IR was mostly due to a significant decline in rDNA transcripts (Fig. 3B, Figure

3-figure supplement 2A). By contrast, the total read count from the protein coding transcripts

significantly increased after IR (Fig. 3B). Analysis of the protein coding transcripts showed that

the transcripts of 3,026 and 1,388 protein-coding genes increased and decreased after IR,

respectively (Fig. 3C and 3D, Supp. File 1). To validate our EU-RNA sequencing results, we

performed quantitative RT-PCR to measure the nascent transcript levels of genes that were up-

and down-regulated after IR. Consistently, we found rDNA transcripts of *28S* and *18S* were

significantly downregulated after IR; while the p53 regulated gene *Cyclin-dependent kinase*

inhibitor 1 (*Cdk1a*/p21) was highly induced (Figure 3-figure supplement 3). The gene ontology

terms describing the genes that were activated after IR included known DNA damage response

pathways and related genes (Fig. 3E, Supp. File 2). For examples, the intrinsic apoptotic

signaling pathway in response to DNA damage (GO:0008630), type 2 response (GO:0042092),

and cytokine-mediated signaling pathway (GO:0019221) were up regulated significantly. Pro-

inflammatory cytokines are the major components of immediate early gene programs, being

rapidly activated after irradiation in various cell types (SchaueKachikwu and McBride, 2012)

ultimately leading to radiation-induced fibrosis in cancer patients following radiation therapy

(KimJenrow and Brown, 2014) (Yu et al., 2023). Meanwhile many of the genes that were

downregulated after IR included gene products that are involved in chromatin organization and

nucleosome assembly (Fig. 3E, Supp. File 3). For examples, chromatin silencing (GO:0006342),

DNA packaging (GO:0006323), and nucleosome assembly (GO:0006334) genes were down

regulated significantly.

To gain more insights on how the transcription of protein coding genes was regulated after IR,

we defined differentially expressed genes (DEGs) between samples before and after IR. We

found read number from these DEGs became significantly greater after IR (Fig. 4A). We sorted

the protein-coding DEGs by average expression level of each gene in all 4 samples: two replicate

samples before IR and two replicates after IR. The number of reads derived from the mostly

highly expressed protein-coding DEGs became significantly smaller after IR (Fig. 4A, 4B). If the

gene repression after IR is due to their being *in cis* to the DNA lesion, it would be expected that

genes that were repressed after IR would tend to be longer, because they would be more likely to

be damaged. However, this was not the case because the length of the nascent transcripts was

equivalent regardless of whether their transcription was repressed, activated, or not changed after

IR (Fig. 4C). Intriguingly, we found that the repressed genes of the top 100 high-expression

DEGs tended to be shorter (Fig. 4C). Next, we inspected the expression level of individual

protein-coding genes and confirmed that most changes in gene expression after IR tended to

occur for the genes that were activated after IR, while many of the genes that had a high-

expression level were repressed after IR, for example, the histone encoding genes (Fig. 4D).

Strikingly, we found that vast majority of the histone genes showed reduced transcription after

IR (Fig. 4E, Figure 3-figure supplement 2B), and was validated by RT-PCR analysis (Figure 3-

figure supplement 3). Finally, to determine whether transcriptional repression was occurring at

the initiation, elongation or both stages of transcription, we examined the read counts throughout

the open reading frames of the repressed protein coding genes, before and after IR. We found

that the decrease of transcripts mainly occurred in the gene body of these genes with similar

intensity at both the 3' and 5' ends of the gene body, which indicates transcriptional repression

after IR occurred at the initiation stage of transcription (Fig. 4F). Therefore, these data indicate

that the bulk reduction in nascent transcripts after IR is mainly due to reduced transcriptional

initiation of the rDNA and histone genes.

A CRISPR-Cas9 screen identifies ATM, neddylation and CUL4B as promoting transcriptional

inhibition after IR

We performed a genome-wide gRNA library CRISPR-Cas9 screen in Abl pre-B cells, allowing 7

days for the gRNAs to inactivate their target genes (Fig. 5A). We then sorted the 10% of the cells

with the most nascent RNA (high EU) 30 minutes after IR, as these would include cells with

gRNAs corresponding to gene products that are required for transcriptional inhibition after IR.

We sequenced the gRNAs within the high EU cells and within the total cell population. We

calculated an enrichment score for each of the 5 gRNAs that were included in the library against

each gene (Supp. File 4, Figure 5-figure supplement 1). Fig. 5B shows the enrichment scores for

the 5 gRNAs for some of the top hits (high EU) from the screen. We found that gRNAs against

ATM were enriched in the high EU cells, consistent with the fact that transcriptional inhibition

of a gene after induction of a DSB requires ATM signaling (Shanbhag et al., 2010). Other top

hits included most of the machinery that mediates neddylation, the post translational covalent

addition of NEDD8, a small ubiquitin-like peptide, onto other proteins (Rabut and Peter, 2008).

These hits included the *Nedd8* gene encoding the NEDD8 ubiquitin like modifier, *Nae1* encoding

NEDD8 activating enzyme E1 subunit 1 NAE1, *Uba3* encoding the Ubiquitin like modifier

activating enzyme 3 UBA3, *Ube2m/Ubc12* encoding the NEDD8-conjugating enzyme UBC12,

Ube2f encoding a neddylation E2 enzyme UBE2F, and *Rbx1* and *Rbx2* which encode linkers that

facilitate NEDD8 transfer from the E2 enzyme to Cullins (Fig. 5B). We also uncovered gRNAs

against the gene encoding the neddylation substrate CUL4B enriched in the high EC cells (Fig.

5B). These data suggested that neddylation may have a novel role in transcriptional inhibition

after IR.

We validated the role of ATM in bulk transcriptional inhibition after IR with the ATM inhibitor

Ku55933 (Fig. 5C). To validate a role for neddylation in transcriptional shut off after IR, we

depleted NAE1 with a gRNA to *Nae1* (Figure 5-figure supplement 2) and showed that depletion

of NAE1 also greatly reduced transcriptional inhibition after IR (Fig. 5D). Similarly, a 16-hour

treatment with the neddylation inhibitor MLN4924 also greatly reduced transcriptional inhibition

after IR (Fig. 5E). To ensure that the role of neddylation in transcriptional inhibition after IR is

not unique to the murine Abl pre-B cells, we found that transcriptional inhibition, detected by

fluorescence microscopy, in U2OS cells was significantly reduced upon inhibition of neddylation

(Fig. 5F). As such, these data suggest that neddylation is required for efficient transcriptional

inhibition after IR.

While most neddylation substrates have only been identified upon overexpression of the

neddylation machinery, the Cullins have been shown to be *bone fide* neddylation substrates *in*

vivo (Pan et al., 2004). Given that we found gRNAs to Cul4b encoding CULLIN 4B enriched in

the EU high cells (Fig. 5B) we asked whether its neddylation is induced after IR. We found that

when compared to the level of unneddylated CUL4A there was not a significantly higher

proportion of neddylated CUL4A or CUL4B in U2OS cells after IR (Fig. 6A). We used the

neddylation inhibitor MLN4924 as a positive control to confirm which bands were neddylated

CUL4A and CUL4B (Fig. 6A). To determine whether CUL4B or CUL4A were required for IR

induced transcriptional inhibition, we made stable Abl pre-B cell lines lacking each protein by

gRNA-mediated disruption of the *Cul4a* or *Cul4b* genes. Loss of CUL4A had no effect on

transcriptional inhibition after IR, while loss of CUL4B partially reduced transcriptional

inhibition after IR (Fig. 6B, Figure 6-figure supplement 1A). Given that CUL4B and CUL4A

show some functional redundancy (Hannah and Zhou, 2015, Brown et al., 2015), we depleted

both proteins at the same time. Additional transient bulk gRNA transfection-mediated depletion

of CUL4A from cells lacking CUL4B (depletion of both CUL4A and CUL4B is lethal) did not

further increase the block of transcriptional inhibition after IR (Figure 6-figure supplement 1B).

As such, these data indicate that CUL4B but not CUL4A contributes to transcriptional inhibition

after IR. Given that neddylation of CUL4A/CUL4B regulates cell cycle progression (Hannah and

Zhou, 2015), and we saw changes in the distribution of the cell cycle phases upon

CUL4A/CUL4B depletion and upon blocking neddylation as indicated by the change in relative

heights of the EU low (G_1) and EU high (S/G_2) peaks (Fig. 5D, 5E, 6B and Figure 6-figure

supplement 1B,C), we wondered if cell cycle arrest may be influencing transcriptional inhibition

after IR. Accordingly, we inhibited neddylation for 1-3 hours in Abl pre-B cells, which

effectively inhibited neddylation (Fig. 6C) and tested the effect on transcriptional inhibition after

IR. Strikingly, we observed transcriptional inhibition after IR even upon neddylation inhibition

treatment for 1-3 hours (Fig. 6D), indicating that neddylation is not required for transcriptional

inhibition after IR. We also observed that the length of time required for treatment with the

neddylation inhibition to block transcriptional inhibition after IR (Fig. 5E) caused arrest of the

cell cycle in G_2 phase (Figure 6-figure supplement 2). Therefore, these data indicate that

neddylation *per se* is not required for transcriptional inhibition after IR. They also raise the

possibility that the cell cycle arrest caused by persistent loss of neddylation may prevent

transcriptional inhibition after IR.

Cell cycle arrest in G_1 or G_2 phase blocks inhibition of transcription after IR

To directly determine the relationship between cell cycle arrest caused by prolonged neddylation

inhibition and transcriptional inhibition after IR, we labelled the Abl pre-B cells with both EU

(nascent transcripts) and 7-Aminoactinomycin D (7-AAD) (DNA stain). Prolonged neddylation

inhibition led to accumulation of cells with a 4N DNA content and greatly reduced

transcriptional inhibition after IR (Fig. 7A, Figure 6-figure supplement 2). To investigate if this

correlation was specific to neddylation inhibition, we used an unrelated inhibitor that causes cell

cycle arrest with a 4N DNA content, RO-3306, a CDK1 inhibitor (Vassilev et al., 2006). We also

saw accumulation of cells with a 4N DNA content and no transcriptional inhibition after IR upon

CDK1 inhibition (Fig. 7B, Figure 6-figure supplement 2). To determine whether this effect was

unique to cells arrested with a 4N DNA content or was shared with conditions that cause arrest in

G_1 phase, we treated cells with 10% or 0.1% FBS, where the later lead to arrest in G_1 phase (due

to serum depletion) and prevented transcriptional inhibition after IR (Fig. 7C). It is relevant to

point out that the level of bulk nascent transcripts in cells arrested in G_1 by serum depletion is

still far higher than cells lacking EU (Fig. 7C), suggesting that the limited transcriptional

inhibition after IR in G_1 arrested cells is not just because there is only minimal transcription

occurring. In agreement, two CDK4/6 inhibitors, Ribociclib and Palbociclib, that lead to G_1

arrest also prevented transcriptional inhibition after IR (Fig. 7D, 7E). Intriguingly, these data

indicate that multiple different treatments that lead to cell cycle arrest in G_1 or G_2 *per se* prevent

transcriptional inhibition after IR.

To gain molecular insight into why ATM-dependent bulk transcriptional inhibition occurs after

IR in cycling cells but not in arrested cells, we investigated the possibility that ATM-dependent

pathways that inhibit rDNA transcription after IR in cycling cells (Kruhlak et al., 2007) fail in

arrested cells. In cycling cells, DSBs within the rDNA trigger formation of nucleolar caps due to

damaged rDNA and associated proteins relocalizing to the nucleolar periphery (van Sluis and

McStay, 2017). In cycling cells, DSBs within the rDNA trigger ATM-dependent phosphorylation

of Treacle which promotes recruitment of NBS1 and TOPBP1 to the nucleolar caps to inhibit

rDNA transcription (Larsen et al., 2014, Mooser et al., 2020). Accordingly, in cycling cells,

TOPBP1 accumulates in the nucleolar caps after induction of DSBs, concomitant with repression

of rDNA transcription (Sokka et al., 2015, Mooser et al., 2020). We asked whether TOPBP1-

eGFP recruitment to nucleolar caps was disrupted after DNA damage in cells arrested by

treatment with MLN4924 or RO-3306. The overall induction of TOPBP1 expression was not

affected by inhibitor treatment (Figure 7-figure supplement 1). However, the percentage of cells

with TOPBP1-eGFP localizing to nucleolar caps after IR was markedly reduced upon MLN4924

and RO-3306 treatment (Fig. 7F, 7G). These results are consistent with arrested cells failing to

repress rDNA expression after global DNA damage, as a consequence of compromised ATM-

dependent localization of TOPB1 to nucleolar caps.

Discussion

We find that while the bulk abundance of nascent transcripts is rapidly reduced after IR, more

protein coding genes are induced than inhibited after IR. Instead, the reduction in bulk nascent

transcript levels that occurs after IR is due to reduced transcriptional initiation of a subset of

genes that are the most highly expressed in the cell – the rDNA and histone encoding genes.

Notably, bulk transcriptional inhibition after IR did not occur in cells arrested in G_1 or G_2 phases

of the cell cycle indicating that cells need to be cycling for IR to rapidly inhibit bulk

transcription. The length-independent and dose-independent reduction in bulk abundance of

nascent transcript after IR (Figs. 5C, Supp. Fig 3) suggests that the reduced bulk abundance of

nascent transcripts after IR may occur *in trans* as a programmed event. This is in contrast to

studies that have found transcriptional inhibition *in cis* of a gene immediately adjacent to an

endonuclease-induced DSB. Our work indicates that the genome wide transcriptional response to

DSBs after IR cannot be extrapolated from single gene studies.

Analysis of nascent transcripts at early times after irradiation revealed a different transcriptional

response compared to changes in total mRNAs after irradiation (Lieberman et al., 2017). These

changes in mRNA levels typically occurred in an IR dose-dependent manner. By contrast, the

bulk changes in nascent transcripts occurred in a dose-independent manner (Fig. 2D). Analysis

of total mRNAs changes following IR showed altered expression of genes involved in signal

transduction, regulation of transcription, and metabolism (Su et al., 2004). Similarly, the

upregulated nascent transcript changes identified pathways including signal transduction, while

nascent transcripts from protein-coding genes that affected nucleosome assembly and chromatin

structure (histones) were downregulated after IR (Fig. 3B, 3E). Finally, most of the gene

expression changes detected by analysis of total mRNA continued to increase or decrease over

long periods of time, up to 48 hours (Su et al., 2004), whereas the reduction of bulk nascent

transcript levels occurred in a very transient manner and was already returning to normal by 4

hours after IR (Fig. 1, Fig. 2D).

How is bulk nascent transcription being inhibited after irradiation? Reminiscent of the ATM

dependence on the changes in mRNA levels after IR (Artuso et al., 1995), the reduction in bulk

nascent transcript levels after IR was also dependent on ATM (Fig. 5C). Given that most of the

reduction in bulk nascent transcript levels was due to the rRNA (Fig. 3B), this is consistent with

the previous report of ATM-dependent inhibition of RNA Polymerase I transcription in response

to DSBs (Kruhlak et al., 2007). In this case, the Pol I transcription appeared to be inhibited at

both the initiation and elongation stages (Kruhlak et al., 2007). Sequencing analysis of nascent

transcripts of protein-coding genes whose expression significantly decreased 30 minutes after IR

suggested that it is the initiation of Pol II transcription that is inhibited, given that the reduction

in sequencing reads at the 5' end and 3' ends of open reading frames was equivalent (Fig. 4F).

Studies that examined the mechanism of Pol II transcriptional inhibition of one gene adjacent to

an endonuclease induced DSB identified a reduction in transcriptional elongation as indicated by

reduced Pol II Ser2 phosphorylation (Shanbhag et al., 2010), while other studies found a defect

in both Pol II initiation and elongation of a different gene adjacent to an endonuclease induced

DSB, in a manner dependent on DNA-PK and the proteasome (Caron et al., 2019, Pankotai et

al., 2012). Our data is consistent with the possibility that the major mechanism for the repression

of the ~1,000 protein coding genes after IR is at the transcriptional initiation stage. However, our

data do not rule out that transcriptional elongation may be additionally repressed after IR, but

would not be observed in our analyses due to the repression of transcriptional initiation.

Rapid inhibition of transcription *in cis* has been observed following endonuclease-mediated DSB

induction, where a DSB-proximal transcriptional reporter was inhibited while a second

transcriptional reporter inserted elsewhere in the genome without a proximal FokI nuclease site

was not inhibited after induction of Fok1 (Shanbhag et al., 2010). If the transcriptional inhibition

after IR that we observe was occurring *in cis,* we would expect that longer genes would be more

inhibited than shorter genes after IR, as they are more likely to experience a DSB induced by IR.

However, this was not the case because the length of the nascent transcripts was equivalent

regardless of whether their levels were increased, didn't change, or were decreased after IR (Fig.

4C). We would also expect that bulk transcript levels would be more reduced with a higher dose

of IR if it occurred *in cis*, but that was not the case (Fig. 2D). In fact, transcriptional repression

around a nuclease-induced DSB can spread hundreds of kb away from the break, throughout a

whole topological associated domain marked by gamma H2AX (Purman et al., 2019).

Importantly, our data do not contradict that DSBs can induce transcriptional inhibition *in cis*,

rather it likely reflects the random nature of the DNA damage induced by IR is not sufficient to

detect inhibition *in cis*, as every cell will have DNA DSBs at different locations. We also are

examining the effect following global DNA damage induced IR versus the effect of induction of

a single or limited numbers of DSBs by endonuclease induction. It is noteworthy that the nascent

transcript levels of more genes rapidly increased after IR than were reduced (Figs. 3 and 4), such

that the situation, at least after IR, is more complex than transcriptional inhibition *in cis* to the

DSBs.

The change in transcript levels after irradiation tended to depend on the expression level of the

genes before irradiation. Those genes that were normally most highly transcribed were repressed

after IR, while genes that were normally expressed at intermediate or low levels tended to be

induced after IR (Fig. 4A). The mechanistic reason for this is unclear. Among the genes that

were most repressed after IR treatment were many of the histone encoding genes (Fig. 4D, 4E).

Histone gene expression has been shown to be reduced after IR previously in a manner

dependent on ATM and p53, as seen at the mRNA level 10 hours or more after irradiation (Su et

al., 2004). By contrast, reduction in nascent histone transcript levels occurred 30 mins after IR

(Fig. 4D, 4E). Interestingly, we observed a tremendous expression induction of *Cdkn1a/p21*

gene, which encodes a potent cyclin-dependent kinase inhibitor, after IR (Supp Fig. 6). The

histone transcriptional coactivator NPAT induces histone transcription when it is phosphorylated

by cyclin E/CDK2 (Zhao et al., 1998, Zhao et al., 2000, Ma et al., 2000). As such, highly

elevated levels of Cdkn1a/p21 after IR might inactivate cyclin E/CDK2, leading to hypo-

phosphorylation of NPAT and an immediate repression of histone gene expression after IR. It

also may be relevant that the loci found to be repressed by bulk IR are highly repetitive gene

arrays that tend to form nuclear sub-compartments (nucleoli, histone bodies). As such, their

likelihood of repeats being in the vicinity of a repeat with an IR-induced DNA lesion in three-

dimensional space is high, which may promote their transcriptional repression after IR in *trans*.

Moreover, silencing may spread through the relevant nuclear sub-compartments, consistent with

the formation of DNA damage compartments described recently (Arnould et al., 2023).

In response to UV exposure, bulk inhibition of transcription occurs, followed by transcriptional

recovery after repair of UV-induced damage, where transcriptional recovery after repair of UV-

induced damage is dependent on the histone variant H3.3 histone chaperone HIRA (Bouvier et

al., 2021, AdamPolo and Almouzni, 2013). Mechanistically, HIRA functioned to repress the

transcriptional repressor ATF3, in turn promoting transcriptional recovery after repair of UV-

induced damage (Bouvier et al., 2021). We find that neither HIRA nor H3.3 is required for

recovery of bulk nascent transcripts after DSB repair (Fig. 1). Additionally, we find that the

genes that have altered levels of nascent transcripts after UV damage (Bouvier et al., 2021) and

after IR are quite distinct (data not shown). However, the UV studies were performed on nascent

mRNA from a human cell line, while our studies were performed on total nascent RNA from a

mouse cell line.

Why didn't our CRISRP/Cas9 screen of factors responsible for bulk nascent transcript inhibition

after IR identify previously reported factors involved in Pol II transcriptional inhibition proximal

to an endonuclease-induced DSB? This is likely because most of the reduction in bulk nascent

transcript level that we were detecting after IR was due mainly to reduced transcript abundance

from the rDNA, rather than Pol II transcripts (Fig. 3B). We did find ATM in the screen (Fig. 5B,

5C), and this is consistent with the fact that ATM is required for rDNA transcriptional inhibition

after IR (Kruhlak et al., 2007). Many of the most significant hits from the screen encoded factors

that are involved in the neddylation pathway and the neddylation substrate CUL4B (Fig. 5D, 5E,

5F and Fig. 6B). However, short times of neddylation inhibition were sufficient to inhibit

neddylation but did not prevent transcriptional inhibition after IR (Fig. 6C, 6D). Longer times of

neddylation inhibition did prevent transcriptional inhibition after IR (Fig. 5E, 5F), but also

caused cell cycle arrest (Fig. 7A, Supp. Fig. 10). These results lead us to speculate that

neddylation is not directly involved in transcriptional inhibition after IR, but that neddylation

promotes cell cycle progression, and that it is the cell cycle arrest occurring upon neddylation

inhibition that prevents the bulk reduction of nascent transcript levels after IR. Consistent with

the idea that cell cycle arrest *per se* may be preventing transcriptional inhibition after IR, we also

found that CDK1 inhibition leading to G_2 arrest, and serum starvation and CDK4/6 inhibition

leading to G_1 arrest, also prevented bulk transcriptional inhibition after IR (Fig. 7).

Why would cell cycle arrest in G_1 or G_2 phases of the cell cycle prevent transcriptional

repression of rDNA and histone genes after IR? Transcription of rDNA is known to be reduced

in non-cycling cells when total transcript levels were measured (Moss and Stefanovsky, 1995,

O'Mahony et al., 1992) while transcription of histone genes requires ongoing DNA replication

(SittmanGraves and Marzluff, 1983). As such, one possibility for failure to see bulk reduction in

nascent transcript abundance after IR in arrested cells may be that rDNA and histone

transcription is already reduced in G_1 and G_2 arrested cells. However, we found that the total

level of bulk nascent transcripts during G_2 arrest before IR were equivalent to, or more than, the

level of bulk nascent transcripts before cell cycle arrest (compare grey (no arrest) to black lines

(cell cycle arrest) in Figs. 5C-E, 7A, 7B) which suggest that the levels of rDNA transcripts may

not be reduced during G_2 arrest in our experiments given that most of the nascent transcripts are

from the rDNA in the conditions of our experiments (Fig. 7B). It is also possible that a factor that

is required for transcriptional inhibition of the rDNA after IR is absent or inactive in arrested

cells.

In addition to ATM, NBS1 was previously shown to be required for transcriptional inhibition of

rDNA after IR (Kruhlak et al., 2007). The mechanism for this was unclear, but it is tempting to

speculate that the requirement of ATM for transcriptional inhibition after IR in cycling cells may

be mediated through ATM-dependent phosphorylation of Treacle. In this form, Treacle functions

with TOPBP1 to promote recruitment of NBS1 to nucleolar caps to repress rDNA transcription

(Larsen et al., 2014, Mooser et al., 2020). That TOPB1 fails to relocalize to nucleolar caps after

IR in arrested cells (Fig. 7F, 7G) is consistent with a potential loss of ATM-dependent treacle

phosphorylation in arrested cells, which would prevent reduction in rDNA transcription after

DNA damage. Future experiments will reveal further insight into the cell cycle dependent control

of transcriptional inhibition of highly transcribed genes after DNA damage.

Figure legends

Figure 1. Transcriptional inhibition after irradiation and transcriptional restart after DNA

repair in a HIRA independent manner. A. U2OS cells were transfected with either a

scrambled shRNA or shRNA against HIRA, and were either incubated with EU or not, as

indicated, and were irradiated (10 Gy) or not as indicated, followed by detection of EU by click

chemistry of a fluorophore and DNA was detected by DAPI staining. The right panel shows

quantitation of the mean intensity of EU in at least 80 cells for each condition. **** indicates

p<0.001, ** indicates p<0.01, by students T-test. **B.** U2OS cells were transfected with either a

control siRNA (siRNA-Ctrl) or two siRNAs against each gene encoding H3.3 (siRNA-H3). EU

and DAPI were detected as described in A and quantitated as described in A. **** indicates

p<0.001, *** indicates p<0.005, * indicates p<0.05 by students T-test.

Figure 2. **Development of a flow cytometry assay for nascent transcripts shows that**

transcriptional inhibition after IR is not dose dependent. A. The EU positive signal in murine

Abl pre-B cells detected by flow cytometry is due to transcripts, as indicated by the addition of

5μM of the general RNA polymerase inhibitor Actinomycin D for 1 hour. **B**. The two EU peaks

observed by flow cytometry correspond to G_1 (low peak) and G_2 (high peak) cells. Cycling Abl

pre-B cells (left panel) were gated for those with 2N DNA (G_1) content or 4N (G_2) content as

detected by FxCycle Violate or with BrdU incorporation (S) as indicated and were individually

analyzed for EU incorporation into nascent transcripts (right panel). **C.** Schematic of the assay to

detect transcriptional inhibition and transcriptional recovery after IR. **D.** Time course of

transcriptional inhibition and recovery in Abl pre-B cells after IR with the indicated times after

IR at the indicated doses of IR.

Figure 3. Reduction of nascent transcript levels after irradiation is mainly of the rDNA.

A. Schematic of nascent transcript sequencing. **B.** Read counts for the total nascent transcripts,

rDNA transcripts and protein coding transcripts before and 30 minutes after IR, normalized to

ERCC spike in controls. **C**. Significantly changed nascent transcripts from protein coding genes

are indicated upon irradiation, and the numbers indicate the number of upregulated and

downregulated genes 30 minutes after IR. Data shown are an average of the two independent

experimental repeats. **D.** Heat map of significantly increased and decreased nascent transcripts

30 minutes after IR, shown for two independent experimental repeats. Expression z-score was

calculated by subtracting the overall average gene abundance from the raw expression for each

gene and dividing that result by the standard deviation (SD) of all of the measured counts across

all samples. **E.** Gene Ontology analysis of the top significantly enriched GO terms most

upregulated after IR (pink) and most downregulated after IR (blue). Enriched gene number (red)

and fold enrichment (blue) were showed in each GO term.

Figure 4. The highly transcribed protein coding genes tend to be repressed after

irradiation, due to a decrease in the transcription of the histone genes. **A.** Plot of the

transcript abundance of differentially expressed genes (DEGs) showing that highly expressed

genes have reduced nascent transcript levels 30 minutes after IR, while moderately expressed

and low level expressed genes tend to have increased nascent transcript levels 30 minutes after

IR. Mean gene expression and standard deviation is shown in million reads mapped to genes

normalized by ERCC spike-in reads. Data are shown from two independent experimental repeats

(rep) of the experiment. **B.** Heat map showing nascent transcript levels of the top 100 highly

expressed DEGs, ranked by gene expression from top (highest) to bottom, 30 minutes after IR,

shown for two independent experimental repeats. Expression z-score was calculated by

subtracting the overall average gene abundance from the raw expression for each gene and

dividing that result by the standard deviation (SD) of all the measured counts across all four

samples. **C.** Among the top 100 of highly expressed protein-coding genes, repressed genes are

significantly shorter compared to activated genes. The activated, non-changed and repressed

genes show little difference in gene size (the data are averaged for each gene between two

independent experimental repeats). **D.** Plot of change in gene expression after IR against mean

gene expression (log_2), ranked by mean gene expression in samples before IR on the x axis, for

all nascent transcripts. Some of the highly expressed genes whose nascent transcript levels

decreased after IR are labelled in the rectangle, including histone genes. **E.** Heat map of nascent

transcripts of all histone genes 30 minutes after IR, shown for two independent experimental

repeats. **F**. The average read counts for repressed protein coding genes throughout their gene

length before and after IR for 2 independent repeats of the experiment.

Figure 5. Whole genome gRNA screen CRISPR-Cas9 screen for factors involved in

transcriptional inhibition after irradiation identifies the neddylation pathway. A. Schematic

of whole genome gRNA CRISPR-Cas9 screen for gene products that promote transcriptional

inhibition after IR**. B.** Fold enrichment of 5 guide RNAs against the indicated genes in the 10%

of cells with most EU incorporated into transcripts 30 minutes after IR. **C**. Inhibition of ATM

greatly reduces transcriptional inhibition 30 minutes after IR in Abl pre-B cells. The inhibitor

was used at 15 μM for 1 hours. **D.** gRNA mediated depletion of Nae1 greatly reduces

transcriptional inhibition 30 minutes after IR in Abl pre-B cells. **E.** Inhibition of neddylation

greatly reduces transcriptional inhibition 30 minutes after IR. The inhibitor was used at 1μM for

16 hours in Abl pre-B cells. **F.** Inhibition of neddylation reduces transcriptional inhibition after

IR in U2OS cells, as detected by fluorescence analysis of nascent transcripts as described in

legend to Fig. 1 and quantitated as in Fig. 1. Significant difference after IR compared to before

IR are indicated by asterisks, where **** indicates p<0.001, * indicates p<0.05 by students T-

test. All experiments in this figure are in murine Abl pre-B cells.

Figure 6. CUL4B but not CUL4A contributes to transcriptional inhibition after irradiation.

A. Analysis of CUL4A and CUL4B neddylation in U2OS cells after IR, in the absence of

presence of 10 μM treatment for 3 hours with the neddylation inhibitor. N8 indicates the

neddylated species. **B.** Analysis of nascent transcripts in Abl pre-B cell lines stably depleted of

CUL4A or CUL4B 30 minutes after IR, as indicated. **C.** Short treatment of Abl pre-B cells with

neddylation inhibitor MLN4924 is sufficient to block neddylation of CUL4B. **D.** The same cells

used in C were analyzed for EU incorporation into nascent transcripts 30 minutes after

irradiation, without or with the indicated time of MLN4924 treatment before irradiation.

Figure 7. Cell cycle arrest in G_1 or G_2 prevents transcriptional inhibition after DNA

damage.

A. Abl pre-B cells were treated with MLN4924 (1 μM) for 16 hours, followed by IR and staining

of DNA with 7-AAD and nascent transcripts with EU 30 minutes after IR. **B.** Abl pre-B cells

were treated with RO-3306 (10 μM) for 16 hours, followed by IR and staining of DNA with 7-

AAD and nascent transcripts with EU 30 minutes after IR. **C.** The left two panels show the cell

cycle distribution of Abl pre-B cells after growth for 72 hours in 1% FBS or 0.1% FBS. The

rectangles and numbers indicate the % of cells with a 2N DNA content. The right panel shows

the EU incorporated into nascent transcripts 30 minutes after IR for the same samples. **D.** Abl

pre-B cells were treated with Ribociclib (5 μM) for 24 hr., followed by IR and staining of DNA

with 7-AAD and nascent transcripts with EU 30 minutes after IR. **E.** Abl pre-B cells were treated

with Palbociclib (5 μM) for 24 hr, followed by IR and staining of DNA with 7-AAD and nascent

transcripts with EU 30 minutes after IR. **F.** The U2OS cells were treated with doxycycline (1

μg/mL) for 12 hours to express eGFP-TOPBP1. Then, DMSO, MLN4924 and RO3306 were

added to the cells for another 16 hours. TOPBP1 localization was shown in cells before IR or 2

hours after IR (10 Gy). Scale bar is 10 μm. **G.** Quantification of TOPBP1 expressing cells with

nucleolar caps before and after IR (10 Gy) in cells treated with MLN4924 and RO-3306. Data

shown are an average of the three independent experimental repeats. Significant differences are

indicated by asterisks, where ** indicates $p < 0.01$ by students T-test.

Figure 1-figure supplement 1. Transcriptional inhibition after irradiation and

transcriptional restart after DNA repair in U2OS cells. A. U2OS cells were either incubated

with EU or not, as indicated, and were irradiated (10 Gy) or not as indicated, followed by

detection of the EU by click chemistry of a fluorophore, and immunofluorescence staining of

gamma H2AX in the same cells and the DNA was detected by Hoechst staining.

Figure 1-figure supplement 2. Confirmation of knockdown of HIRA (A) and H3.3 (B). The

samples were from the same experiments shown in Figure 1.

Figure 2-figure supplement 1. The mean intensities of the EU peaks shown in Figure 2D are

indicated. The mean value of each sample was calculated by Flowjo software.

Figure 3-figure supplement 1. Nascent transcript levels before and after 30 minutes of IR.

The samples from two independent experiments were used for the EU-seq and analyses in Fig. 3

and 4.

Figure 3-figure supplement 2. Screen shot from the UCSC browser of nascent transcripts.

A. Nascent transcripts before and after IR over ribosomal DNA (rDNA). **B.** Nascent histone

transcripts before and after IR over histone cluster 1.

Figure 3-figure supplement 3. Validation of nascent transcript levels of DEGs from EU-

RNA seq by real-time quantitative RT-PCR. Samples from three independent experiments

were used for the analyses. Ct value is presented to show the absolute amounts of EU labeled

RNA transcripts from the same number of cells before and after 30 minutes of IR (10 gray).

Significant difference after IR compared to before IR are indicated by asterisks, where **

indicates $p < 0.01$, * indicates $p < 0.05$ and ns indicates non-significant by students T-test.

Figure 5-figure supplement 1. CRISPR-Cas9 screen identifies genes promoting

transcriptional inhibition after IR. A volcano plot of guide RNA changes between Eu high

cells and unsorted cells. Labeled genes are some of those that have P adjust $<= 10^{-6}$.

Figure 5-figure supplement 2. Confirmation of knockdown of Nae1. gEV is an empty vector.

The samples were from the same experiments shown in Figure 5D.

Figure 6-figure supplement 1. Analysis of CUL4A and CUL4B depletion. A. The western

blot shows the CUL4A and CUL4B levels from the experiment shown in Fig. 6B. Additionally

gRNAs were used to deplete CUL4A in cul4b cells and CUL4B in cul4a cells and their western

blot analyses are also shown. **B.** The EU analysis of these double depleted cells is shown**. C.** Cell

cycle analysis of the experiment shown in A and B and Fig. 6B.

Figure 6-figure supplement 2. Cell cycle analysis of cells treated with MLN4924 and

RO3306. MLN4924 or RO3306 treatment leads cell cycle arrest in G_2 phase.

Figure 7-figure supplement 1. Quantification of cells with TOPBP1 expression before and

after IR (10 Gy) in cells treated with MLN4924 and RO-3306. The samples were from the same

experiments shown in Fig. 7F, 7G.

Supplementary File 1. Nascent RNA profiles of each gene using EU RNA-seq.

Supplementary File 2. Significantly enriched GO terms for up-regulated gene after irradiation.

Supplementary File 3. Significantly enriched GO terms for down-regulated genes after

irradiation.

Supplementary File 4. Whole genome CRISPR-Cas9 screen detects the abundance of all

gRNAs and target genes for EU high cells and unsorted cells.

Figure 1-figure supplement 2 Source Data 1. Original file for the Western blot analysis in

Figure 1-Figure Supplement 1A (anti-HIRA and anti-GAPDH).

Figure 1-figure supplement 2 Source Data 2. PDF containing Figure 1-Figure Supplement 1A

and original scans of the relevant Western blot analysis (anti-CUL4A, anti-CUL4B and anti-

GAPDH) with highlighted bands and sample labels.

Figure 1-figure supplement 2 Source Data 3. Original file for the Western blot analysis in

Figure 1-Figure Supplement 1B (anti-H3.3 and anti-GAPDH).

Figure 1-figure supplement 2 Source Data 4. PDF containing Figure 1-Figure Supplement 1B

and original scans of the relevant Western blot analysis (anti-H3.3 and anti-GAPDH) with

highlighted bands and sample labels.

Figure 5-figure supplement 2 Source Data 1. Original file for the Western blot analysis in

Figure 5D (anti-Nae1 and anti-GAPDH).

Figure 5-figure supplement 2 Source Data 2. PDF of Western blot analysis in Figure 5D and

original scans of the relevant Western blot analysis (anti-Nae1 and anti-GAPDH) with

highlighted bands and sample labels.

Figure 6 Source Data 1. Original file for the Western blot analysis in Figure 6A (anti-CUL4A,

anti-CUL4B and anti-GAPDH).

Figure 6 Source Data 2. PDF containing Figure 6A and original scans of the relevant Western

blot analysis (anti-CUL4A, anti-CUL4B and anti-GAPDH) with highlighted bands and sample

labels.

Figure 6 Source Data 3. Original file for the Western blot analysis in Figure 6C (anti-CUL4B,

and anti-GAPDH).

Figure 6 Source Data 4. PDF containing Figure 6C and original scans of the relevant Western

blot analysis (anti-CUL4B and anti-GAPDH) with highlighted bands and sample labels.

Figure 6-figure supplement 1 Source Data 1. Original file for the Western blot analysis in

Figure 6-figure supplement 1A (anti-CUL4A, anti-CUL4B and anti-GAPDH).

Figure 6-figure supplement 1 Source Data 2. PDF containing Figure 6-figure supplement 1A

and original scans of the relevant Western blot analysis (anti-CUL4A, anti-CUL4B and anti-

GAPDH) with highlighted bands and sample labels.

Methods and materials

Cell culture and transfections

U2OS cells (ATCC, HTB-96) were cultured in McCoy's 5A (Corning, 10050CV) medium

supplemented with 10% fetal bovine serum (FBS) and 1% Penicillin-Streptomycin. Abelson

virus-transformed pre-B cells were maintained in DMEM (Thermo Fisher, 11960-077)

supplemented with 10% FBS, 1% Penicillin-Streptomycin, 1x nonessential amino acids, 1 mM

sodium pyruvate, 2 mM L-glutamine, and 0.4% beta-mercaptoethanol. HEK-293T cells were

maintained in DMEM (Corning, 10-013-CM) supplemented with 10% FBS and 1% Penicillin-

Streptomycin. All the cells were grown at 37 °C under a humidified atmosphere with 5% CO_2.

SiRNA oligos against human H3F3A and H3F3B (SMARTPool) for RNAi in U2OS cells were

purchased from Horizon Discovery (Dharmacon). 100 nM of H3F3A and H3F3B were mixed

with Lipofectamine RNAiMAX transfection reagent (Thermo Scientific, 13778150) according to

the manufacturer's protocol to knockdown H3.3 for 48 hours. The siRNA control (ON-

TARGETplus non-targeting) was also purchased from Horizon Discovery (Dharmacon) and used

as negative control. shRNA lentiviral plasmids against HIRA (5'-

TAGAGCATACCAAGATGCC-3') and the control were described in a previous study (Huang

et al., 2018). HEK-293T cells were transfected with a mixture of shRNA plasmids and the viral

packaging and envelope vectors, pCMV-dR8.2 and pCMV-VSVG. The media containing

shRNA virus particles were collected 48 to 72 hours after transfection and filtered through a 0.45

µm filter. Cells were incubated with the lentiviral supernatant containing 5 µg/ml polybrene

(Sigma-Aldrich, S2667) for 24 hours, followed by 1 µg/mL Puromycin selection for another 48

hours. To inactivate Nae1, Cul4a and Cul4b in bulk cell populations, guide RNAs (gRNAs)

against each gene were cloned into pKLV-U6gRNA-EF(BbsI)-PGKpuro2ABFP (Addgene,

#62348) modified to express human CD2 or Thy1.1 as cell surface markers. The pKLV-gRNAs

lentiviruses were prepared in 293T cells as described above. The Abl pre-B cells containing

pCW-Cas9 (addgene, #50661), which can express Cas9 with doxycycline induction, were mixed

with viral supernatant supplemented with 5 µg/ml polybrene and centrifuged at 1800 rpm for 1.5

hours at room temperature. After the spin-infection, the transduced cells were maintained in

DMEM with 3 µg/ml doxycycline (Sigma-Aldrich, D9891) for 3 days before flow cytometric

cell sorting based on hCD2 or Thy1.1 expression. To make stable cell lines depleted of Cul4a

and Cul4b, serial dilution of the sorted cells into 96-well plate was used to isolate single cells.

Western blot analysis was used to determine the knockdown efficiency of each target genes.

In EU flow cytometric and immunofluorescence analysis, final concentrations of 15 µM ATM

inhibitor Ku55933 (Selleck Chemicals, S1092) and 5 µM of Actinomycin D (Sigma-Aldrich,

A9415) were added to cell culture 1 hour prior to irradiation. 1 µM and 10 µM of neddylation

inhibitor MLN4924/Pevonedistat (Active Biochem, A-1139) were used for long (16 hours) and

short (1 to 3 hour) treatments, respectively. To arrest cells in G_1 cell cycle phase, cells were

incubated in media supplemented with 5 µM Palbociclib (Selleck Chemicals, S1116) or 5 µM

Ribociclib (Selleck Chemicals, S7440) for 24 hours. For serum starvation, Abl pre-B cells were

grown in complete medium containing 10% of FBS to desired density and collected and washed

in medium with reduced concentration (0.1%) of FBS. The cells were maintained in the medium

with reduced FBS for 72 hours to arrest in G_1 phase. Representative data are shown for

experiments repeated three of more times with consistent results.

U2OS cell lines were authenticated by STR profiling, and MCF10A and murine cell lines tested

negative for mycoplasma contamination.

Western blots

The following antibodies were used for western blot: CUL4A (Cell Signaling Technology,

2699S, 1:1000), CUL4B (Proteintech, 12916-1-AP, 1:1000), H3.3 (Millipore Sigma, 09-838,

1:1000), HIRA (Abcam, ab20655, 1:1000), NAE1 (Thermo Fisher, PA5-59836, 1:500), GAPDH

(Sigma-Aldrich, G8795, 1:5000). Representative data are shown for experiments repeated three

of more times with consistent results.

Fluorescence microscopy

For immunofluorescence, Click-iT RNA Alexa Fluor Imaging Kit (Thermo Fisher, C10330) was

used to label newly synthesized RNAs in the cells. Briefly, 50,000 U2OS cells grown on cover

slips in 24-well plate were irradiated with 10 Gray and allowed to recover for indicated times at

37°C with 5% CO2. 0.5 mM EU was added to the medium and incubated for 45 minutes for EU

incorporation. Cells were then washed with PBS, fixed in 4% paraformaldehyde PBS for 15

minutes at room temperature, and permeabilized in cold 0.5% Triton X-100 PBS for 10 minutes.

Cells were blocked in 3% BSA-PBS for 1 hour at room temperature and subsequently incubated

overnight at 4 °C in primary antibody (anti-γH2AX (S139), EMD Millipore, 05-636). Coverslips

were then washed 3x with PBST (0.05% Tween 20), incubated with secondary antibody diluted

in 3% BSA PBS (Alexa Fluor 488 Goat anti-mouse IgG, BioLegend, 405319) in the dark for 1

hour at room temperature and washed 3x with PBST. Click-iT reaction cocktail was prepared

according to the manufacturer's protocol and immediately added to the cells to perform click

reaction in the dark for 30 minutes at room temperature. After washes with Click-iT reaction

rinse buffer (Component F) and PBS, cells were stained with Hoechst (1:2000) or DAPI (Sigma-

Aldrich, D9542) in PBS for 10 minutes and mounted in Prolong Gold Antifade Mountant (Life

Technologies, P-36930). Images were taken on Biotek Lionheart Automatic Microscope and EU

intensity quantification was conducted using Biotek Gen5 software. For eGFP-TOPBP1

fluorescence microscopy, 50,000 cells grown on coverslips in 24-well plates were treated with

doxycycline (1 μg/mL) for 12 hours to express eGFP-TOPBP1. MLN4924 (1 μM) and RO3306

(10 μM) were added to the cell culture, and the cells were incubated for another 16 hours

followed by IR (10 Gy) and recovery for 2 hours. Cells were fixed in 4% Paraformaldehyde for

20 minutes followed by Hoechst (1:2000) staining for DNA and mounting in Prolong Gold

Antifade Mountant. Images were taken and quantified on Biotek Lionheart Automatic

Microscope. Representative data are shown for experiments repeated three of more times with

consistent results.

Flow cytometry and cell cycle analysis

Click-iT RNA Alexa Fluor Imaging Kit was adapted to label newly synthesized RNAs in the

cells for flow cytometry. Abl pre-B cells grown in 24-well plate were irradiated with 10 Gray

and allowed to recover for different times. 2 mM EU was added to the medium and incubated for

30 minutes for EU incorporation. Cells were then washed with PBS, fixed in 4%

paraformaldehyde PBS for 15 minutes at room temperature, and permeabilized in cold 0.5%

TritonX-100 PBS for 5 minutes. Click-iT reaction cocktail was prepared according to the

manufacturer's protocol and immediately added to the cells to perform click reaction in the dark

for 30 minutes at room temperature. Cells were then washed with Click-iT reaction rinse buffer

(Component F) and 3% BSA-PBS, respectively. For cell cycle analysis, BrdU (10 ug/mL) was

added to the cells and incubated for 30 minutes to label new DNAs. Cells were washed with

PBS, fixed in 4% paraformaldehyde PBS for 15 minutes at room temperature, and permeabilized

in cold 0.5% Triton X-100 PBS for 5 minutes. Cells were then digested with DNase (BD

Biosciences, 51-2358KC) for 1 hour at 37 °C. Subsequently, cells were incubated with Alexa

Fluor 488 Mouse anti-BrdU (BD Biosciences, 51-9004981, 1:500) in 3% BSA-FBS for 1 hour at

room temperature and washed 2x with PBS, followed by FxCycle Violet (Thermo Scientific,

R37166) or 7-AAD (BD Pharmingen, 559925) staining for 10 minutes. Cells were resuspended

in PBS and analyzed on BD LSRII Flow Cytometer or BD LSRFortessa Flow Cytometer. Flow

cytometry results were further analyzed using FlowJo software. Representative data are shown

for experiments repeated three of more times with consistent results.

CRISPR-Cas9 screen

More than 140 million wild type Abl pre-B cells carrying inducible Cas9 transgene were

transduced with a lentiviral gRNA library containing 90,230 gRNAs targeting over 18,000

mouse genes (Addgene, 67988) by spin-infection as described above. 3 days post infection, cells

transduced with gRNAs were sorted on a BD FACSAria II Cell Sorter based on BFP expression.

BFP positive cells were treated 3 µg/ml doxycycline for 7 days to induce gRNA expression and

gene inactivation. Cells were irradiated with 10 Gray, allowed to recover for 30 minutes,

processed as described above for EU labeling of newly synthesized RNAs and analyzed on BD

FACSAria II Cell Sorter. Cells with high (top 10%) EU staining and unsorted cells were

collected, and genomic DNA of the cells were isolated for library preparation using nested-PCR.

The library was sequenced on an Illumina HiSeq 2500 platform. Raw fastq files were

demultiplexed by the Genomics and Epigenomics Core Facility of the Weill Cornell Medicine

Core Laboratories Center. The gRNA sequence region was then retrieved from the sequencing

data using Seqkit (Shen et al., 2016) and mapped to the gRNA sequence library (Koike-Yusa et

al., 2014, Tzelepis et al., 2016). The number of reads of each library sequence was counted and

then normalized as follows (Shalem et al., 2014). Normalized reads per gRNA = reads pers

gRNA total reads for all sgRNAs in sample$\times 10^6$+1. Hereby, the generated normalized reads from

each guide RNA were used and compared between the EU high cell and unsorted cell. P values

were measured by Poisson test to compare guide RNAs between EU high cell and unsorted cell.

FDR was used for adjusting P value. CRISPR score = log2 (final sgRNA abundance/initial

sgRNA abundance). The EU high genes were defined as these genes that have at least one guide

RNA with P adjust value <=0.01 & FC >=1.5. Gene ontology (GO) analysis was performed by

the R package cluster Profiler v3.18.1.

Isolation and deep sequencing of EU labeled nascent transcripts

Click-iT Nascent RNA Capture Kit (Thermo Fisher, C10365) was used to label and capture the

nascent transcripts. In brief, the same number of Abl pre-B cells was plated in two T-25

Polystyrene flasks, one for irradiation (10 Gray) and the other for no IR control. After irradiation,

both flasks of cells were recovered for 30 minutes and incubated in medium with 2 mM EU for

30 minutes to allow for the incorporation of EU into the nascent transcripts. Total RNAs were

harvested using TRIzol reagent (Thermo Fischer, 15596018) following the manufacturer's

protocol. The click-iT reaction was performed as per manufacturer's protocol in 50 µL total

volume for 30 minutes in the dark. Subsequently, 1 µL ultrapure glycogen, 50 µL 7.5 M

ammonium acetate, and 700 µL of chilled 100% EtOH were added to the reaction. The mixture

was incubated at −80 °C for 16 hours. RNA pellet was spun down at 13000 ×g for 20 minutes at

4 °C, washed 2x with chilled 75% EtOH and resuspended in nuclease free water. EU-RNAs were

pulled down with Dynabeads MyOne Streptavidin T1 magnetic beads and extracted with TRIzol

reagents. The same amount of ERCC spike-ins (Thermo Fisher, 4456740) were added to the

purified EU-RNAs, followed by cDNA library generation using NEBNext Ultr II Directional

RNA Library Prep Kit for Illumina (NEB, E7760) according to the manufacturer's protocol and

deep sequencing on Illumina HiSeq 2500 platform.

Real-time quantitative RT-PCR of EU labeled nascent transcripts

EU labeled RNA was prepared from the same number of cells as described above. All the

isolated EU-RNAs were used for cDNA synthesis using Superscript III (Thermo Fischer,

18080044) reverse transcription with random hexamer as primers following the manufacturer's

protocol. The same proportion of cDNA products of each sample was used as template for the

quantitative RT-PCR reaction with Light Cycler 480 SYBR Green I Master Mix (Roche,

04707516001). The Ct value was used to represent the absolute amount of EU-RNAs in each

sample, in which a smaller Ct value indicates higher nascent transcript levels of an individual

gene, given the same initial number of cells and the same proportion of EU-RNAs were used for

the analyses. Primer sequences for the analyses are as follows:

ms28S-fwd, 5'-TGGGTTTTAAGCAGGAGGTG-; ms28S-rev, 5'-

GTGAATTCTGCTTCACAATG-3'(Watada et al., 2020); ms18S-fwd, 5'-

CTTAGAGGGACAAGTGGCG-3'; ms18S-rev, 5'-ACGCTGAGCCAGTCAGTGTA-3'

(StephensStephens and Morrison, 2011); msHist1h2ab-fwd, 5'-GCCTGCAGTTCCCCGTA-3';

msHisth2ab-rev, 5'- ATCTCGGCCGTCAGGTACTC-3'; msHist1h2ac-fwd, 5'-

GGCTGCTCCGCAAGGGT-3'; msHist1h2ac-rev, 5'-CTTGTTGAGCTCCTCGTCGTT-3';

msH2afz-fwd, 5'- ACTCCGGAAAGGCCAAGACA-3'; msH2afz-rev, 5'-

GTTGTCCTAGATTTCAGGTG-3' (Nishida et al., 2005); msCdkn1a-fwd, 5'-

GTGGCCTTGTCGCTGTCT-3'; msCdkn1a-rev, 5'-TTTTCTCTTGCAGAAGACCAATC-3'

(Béguelin et al., 2017).

Analysis of nascent transcripts

The mouse genome version GRCm38.p6 release M23 and the associated GENCODE version of

mouse reference gene set were downloaded from the GENCODE website

(https://www.gencodegenes.org/mouse/release_M23.html). We trimmed adapter sequences and

low-quality sequences in RNA-seq data using the Trim Galore v0.6.6 (Martin, 2011) with default

parameters. To avoid rRNA homologous sequences (i.e., in the intron regions of Zc3h7a or

Cdk8) prior to subsequent genomic and other RNA analysis, we first mapped the reads to mm10

rDNA sequences by TopHat v2.1.1 (Kim et al., 2013). The unmapped reads were then further

mapped to the mouse genome version mm10 and ERCC spike-in version ERCC92 using TopHat

v2.1.1 (Kim et al., 2013). Successfully mapping reads were sorted by SAMtools v1. 5.

Afterward, read counts in several types of genomic feature, i.e., protein-coding genes, rDNA and

ERCCs (ERCC92.gtf), were quantified by Htseq-count v0.11.2 (AndersPyl and Huber, 2015)

using the union gene region option. The read number per gene was normalized based on total

ERCC read numbers in each sample.

To visualize read coverage across the genome, DeepTools v3.5.0 (Ramírez et al., 2014) was used

to convert BAM files into bigwig files using scale factors calculated by total ERCC read number

in each sample. Next, DeepTools was used to plot average read depth per sample across

interested groups of genomic regions (i.e., repressed protein coding genes from 3 kb upstream to

3kb downstream of gene bodies). Screenshots of reads density at individual regions were

generated by IGV 2.8.13 (ThorvaldsdóttirRobinson and Mesirov, 2013).

We then used one tail Poisson test to evaluate difference in gene expression level based on the

read counts normalized by total ERCC read counts. We defined differentially expressed RNAs as

those with a fold change greater than 1.5 and an FDR value smaller than 0.05. To detect highly

expressed genes, we ranked genes by RPKM in the control cells, whereas RPKM was calculated

using ERCC-normalized read counts further normalized by gene length. Gene ontology (GO)

analysis was performed by the R package clusterProfiler v3.18.1 (Wu et al., 2021). Heatmap

were generated by pheatmap.

Data availability

All raw sequencing data from the nascent EU RNA-seq and CRISPR screen experiments have

been deposited in the NCBI project database under accession PRJNA895065. The genome-wide

gRNA library CRISPR-Case9 screen datasets comprise Abl pre-B cells of both unsorted

(SRX18076832) and sorted the 10% of the cells with the most nascent RNA (high EU) 30

minutes after IR (SRX18076831). The raw FASTQ files for nascent EU RNA-seq include pre-B

cells without irradiation (SRX18050529 and SRX18050531) and with 30 minutes after

irradiation (SRX18050530 and SRX18050532). Additionally, both FASTQ files and processed

data for nascent EU RNA-seq are accessible at GSE217123. The source codes employed in the

data analysis and figure generation have been uploaded to GitHub at the following

repository: https://github.com/gucascau/NascentDiff.git.

The following data sets were generated:

Chen et al (2024) NCBI BioProject ID PRJNA895065. Transcriptional inhibition after

irradiation occurs preferentially at highly expressed genes in a manner dependent on cell cycle

progression https://www.ncbi.nlm.nih.gov/bioproject/PRJNA895065

Chen et al (2024) NCBI Gene Expression Omnibus, GSE217123. Transcriptional inhibition after

irradiation occurs preferentially at highly expressed genes in a manner dependent on cell cycle

progression https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE217123

Acknowledgements

The authors thank Yinan Wang for performing the bioinformatics for the high throughput screen.

We thank the Weill Cornell Flow Cytometry Core for flow cytometry. We thank the Weill

Cornell Epigenomics Core for performing the sequencing for the high throughput screen and the

Transcriptional Regulation & Expression Facility at Cornell University, Ithaca for providing

advice and performing the sequencing of the nascent transcripts. The stable doxycycline-

inducible eGFP-TOPBP1 U2OS cell lines were a kind gift from Dr. Helmut Pospiech (Fritz

Lipmann Institute, Germany). We are grateful to Pengbo Zhou for advice on Cullin 4A and B.

We also thank Barry Sleckman, Bo-Ruei Chen and Faith Fowler for advice throughout the

project. JKT is supported by NIH R35 GM139816 and RO1 CA95641. K.C is supported by NIH

R01GM138407, R01GM125632, R01HL148338, and R01HL133254.

References

ADAM, S., POLO, S. E. & ALMOUZNI, G. 2013. Transcription recovery after DNA damage requires
chromatin priming by the H3.3 histone chaperone HIRA. *Cell,* 155**,** 94-106.
ANDERS, S., PYL, P. T. & HUBER, W. 2015. HTSeq—a Python framework to work with high-
throughput sequencing data. *bioinformatics,* 31**,** 166-169.
ARNOULD, C., ROCHER, V., SAUR, F., BADER, A. S., MUZZOPAPPA, F., COLLINS, S., LESAGE, E., LE
BOZEC, B., PUGET, N., CLOUAIRE, T., MANGEAT, T., MOURAD, R., AHITUV, N.,

NOORDERMEER, D., ERDEL, F., BUSHELL, M., MARNEF, A. & LEGUBE, G. 2023. Author
Correction: Chromatin compartmentalization regulates the response to DNA damage.
Nature, 624**,** E1.
ARTUSO, M., ESTEVE, A., BRESIL, H., VUILLAUME, M. & HALL, J. 1995. The role of the Ataxia
telangiectasia gene in the p53, WAF1/CIP1(p21)- and GADD45-mediated response to
DNA damage produced by ionising radiation. *Oncogene,* 11**,** 1427-35.
BÉGUELIN, W., RIVAS, M. A., CALVO FERNÁNDEZ, M. T., TEATER, M., PURWADA, A., REDMOND,
D., SHEN, H., CHALLMAN, M. F., ELEMENTO, O., SINGH, A. & MELNICK, A. M. 2017. EZH2
enables germinal centre formation through epigenetic silencing of CDKN1A and an Rb-
E2F1 feedback loop. *Nat Commun,* 8**,** 877.
BLACKFORD, A. N. & JACKSON, S. P. 2017. ATM, ATR, and DNA-PK: The Trinity at the Heart of
the DNA Damage Response. *Mol Cell,* 66**,** 801-817.
BOUVIER, D., FERRAND, J., CHEVALLIER, O., PAULSEN, M. T., LJUNGMAN, M. & POLO, S. E. 2021.
Dissecting regulatory pathways for transcription recovery following DNA damage reveals
a non-canonical function of the histone chaperone HIRA. *Nat Commun,* 12**,** 3835.
BREDEMEYER, A. L., SHARMA, G. G., HUANG, C. Y., HELMINK, B. A., WALKER, L. M., KHOR, K. C.,
NUSKEY, B., SULLIVAN, K. E., PANDITA, T. K., BASSING, C. H. & SLECKMAN, B. P. 2006.
ATM stabilizes DNA double-strand-break complexes during V(D)J recombination. *Nature,*
442**,** 466-70.
BROWN, J. S., LUKASHCHUK, N., SCZANIECKA-CLIFT, M., BRITTON, S., LE SAGE, C., CALSOU, P.,
BELI, P., GALANTY, Y. & JACKSON, S. P. 2015. Neddylation promotes ubiquitylation and
release of Ku from DNA-damage sites. *Cell Rep,* 11**,** 704-14.
CARON, P., PANKOTAI, T., WIEGANT, W. W., TOLLENAERE, M. A. X., FURST, A., BONHOMME, C.,
HELFRICHT, A., DE GROOT, A., PASTINK, A., VERTEGAAL, A. C. O., LUIJSTERBURG, M. S.,
SOUTOGLOU, E. & VAN ATTIKUM, H. 2019. WWP2 ubiquitylates RNA polymerase II for
DNA-PK-dependent transcription arrest and repair at DNA breaks. *Genes Dev,* 33**,** 684-
704.
CHEN, B. R., WANG, Y., TUBBS, A., ZONG, D., FOWLER, F. C., ZOLNEROWICH, N., WU, W.,
BENNETT, A., CHEN, C. C., FENG, W., NUSSENZWEIG, A., TYLER, J. K. & SLECKMAN, B. P.
2021. LIN37-DREAM prevents DNA end resection and homologous recombination at
DNA double-strand breaks in quiescent cells. *Elife,* 10.
CHEN, K., HU, Z., XIA, Z., ZHAO, D., LI, W. & TYLER, J. K. 2015. The Overlooked Fact:
Fundamental Need for Spike-In Control for Virtually All Genome-Wide Analyses. *Mol Cell*
Biol, 36**,** 662-7.
HANNAH, J. & ZHOU, P. 2015. Distinct and overlapping functions of the cullin E3 ligase
scaffolding proteins CUL4A and CUL4B. *Gene,* 573**,** 33-45.
HUANG, T. H., FOWLER, F., CHEN, C. C., SHEN, Z. J., SLECKMAN, B. & TYLER, J. K. 2018. The
Histone Chaperones ASF1 and CAF-1 Promote MMS22L-TONSL-Mediated Rad51 Loading
onto ssDNA during Homologous Recombination in Human Cells. *Mol Cell,* 69**,** 879-892
e5.
IANNELLI, F., GALBIATI, A., CAPOZZO, I., NGUYEN, Q., MAGNUSON, B., MICHELINI, F.,
D'ALESSANDRO, G., CABRINI, M., RONCADOR, M., FRANCIA, S., CROSETTO, N.,
LJUNGMAN, M., CARNINCI, P. & D'ADDA DI FAGAGNA, F. 2017. A damaged genome's

transcriptional landscape through multilayered expression profiling around in situ-
mapped DNA double-strand breaks. *Nat Commun,* 8**,** 15656.
JACKSON, S. P. & BARTEK, J. 2009. The DNA-damage response in human biology and disease.
Nature, 461**,** 1071-8.
JAO, C. Y. & SALIC, A. 2008. Exploring RNA transcription and turnover in vivo by using click
chemistry. *Proc Natl Acad Sci U S A,* 105**,** 15779-84.
KAKAROUGKAS, A., ISMAIL, A., CHAMBERS, A. L., RIBALLO, E., HERBERT, A. D., KUNZEL, J.,
LOBRICH, M., JEGGO, P. A. & DOWNS, J. A. 2014. Requirement for PBAF in
transcriptional repression and repair at DNA breaks in actively transcribed regions of
chromatin. *Mol Cell,* 55**,** 723-32.
KIM, D., PERTEA, G., TRAPNELL, C., PIMENTEL, H., KELLEY, R. & SALZBERG, S. L. 2013. TopHat2:
accurate alignment of transcriptomes in the presence of insertions, deletions and gene
fusions. *Genome biology,* 14**,** 1-13.
KIM, J. H., JENROW, K. A. & BROWN, S. L. 2014. Mechanisms of radiation-induced normal tissue
toxicity and implications for future clinical trials. *Radiat Oncol J,* 32**,** 103-15.
KOIKE-YUSA, H., LI, Y., TAN, E. P., VELASCO-HERRERA, M. E. C. & YUSA, K. 2014. Genome-wide
recessive genetic screening in mammalian cells with a lentiviral CRISPR-guide RNA
library. *Nat Biotechnol,* 32**,** 267-73.
KRUHLAK, M., CROUCH, E. E., ORLOV, M., MONTANO, C., GORSKI, S. A., NUSSENZWEIG, A.,
MISTELI, T., PHAIR, R. D. & CASELLAS, R. 2007. The ATM repair pathway inhibits RNA
polymerase I transcription in response to chromosome breaks. *Nature,* 447**,** 730-4.
LARSEN, D. H., HARI, F., CLAPPERTON, J. A., GWERDER, M., GUTSCHE, K., ALTMEYER, M.,
JUNGMICHEL, S., TOLEDO, L. I., FINK, D., RASK, M. B., GRØFTE, M., LUKAS, C., NIELSEN,
M. L., SMERDON, S. J., LUKAS, J. & STUCKI, M. 2014. The NBS1-Treacle complex controls
ribosomal RNA transcription in response to DNA damage. *Nat Cell Biol,* 16**,** 792-803.
LIEBERMAN, H. B., PANIGRAHI, S. K., HOPKINS, K. M., WANG, L. & BROUSTAS, C. G. 2017. p53
and RAD9, the DNA Damage Response, and Regulation of Transcription Networks.
Radiat Res, 187**,** 424-432.
MA, T., VAN TINE, B. A., WEI, Y., GARRETT, M. D., NELSON, D., ADAMS, P. D., WANG, J., QIN, J.,
CHOW, L. T. & HARPER, J. W. 2000. Cell cycle-regulated phosphorylation of p220(NPAT)
by cyclin E/Cdk2 in Cajal bodies promotes histone gene transcription. *Genes Dev,* 14**,**
2298-313.
MARTIN, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet. journal,* 17**,** 10-12.
MEISENBERG, C., PINDER, S. I., HOPKINS, S. R., WOOLLER, S. K., BENSTEAD-HUME, G., PEARL, F.
M. G., JEGGO, P. A. & DOWNS, J. A. 2019. Repression of Transcription at DNA Breaks
Requires Cohesin throughout Interphase and Prevents Genome Instability. *Mol Cell,* 73**,**
212-223 e7.
MOOSER, C., SYMEONIDOU, I. E., LEIMBACHER, P. A., RIBEIRO, A., SHORROCKS, A. K.,
JUNGMICHEL, S., LARSEN, S. C., KNECHTLE, K., JASROTIA, A., ZURBRIGGEN, D.,
JEANRENAUD, A., LEIKAUF, C., FINK, D., NIELSEN, M. L., BLACKFORD, A. N. & STUCKI, M.
2020. Treacle controls the nucleolar response to rDNA breaks via TOPBP1 recruitment
and ATR activation. *Nat Commun,* 11**,** 123.

MOSS, T. & STEFANOVSKY, V. Y. 1995. Promotion and regulation of ribosomal transcription in
eukaryotes by RNA polymerase I. *Prog Nucleic Acid Res Mol Biol,* 50**,** 25-66.
NISHIDA, H., SUZUKI, T., OOKAWA, H., TOMARU, Y. & HAYASHIZAKI, Y. 2005. Comparative
analysis of expression of histone H2a genes in mouse. *BMC Genomics,* 6**,** 108.
O'MAHONY, D. J., XIE, W. Q., SMITH, S. D., SINGER, H. A. & ROTHBLUM, L. I. 1992. Differential
phosphorylation and localization of the transcription factor UBF in vivo in response to
serum deprivation. In vitro dephosphorylation of UBF reduces its transactivation
properties. *J Biol Chem,* 267**,** 35-8.
PAN, Z. Q., KENTSIS, A., DIAS, D. C., YAMOAH, K. & WU, K. 2004. Nedd8 on cullin: building an
expressway to protein destruction. *Oncogene,* 23**,** 1985-97.
PANKOTAI, T., BONHOMME, C., CHEN, D. & SOUTOGLOU, E. 2012. DNAPKcs-dependent arrest
of RNA polymerase II transcription in the presence of DNA breaks. *Nat Struct Mol Biol,*
19**,** 276-82.
PANKOTAI, T. & SOUTOGLOU, E. 2013. Double strand breaks: hurdles for RNA polymerase II
transcription? *Transcription,* 4**,** 34-8.
PORTER, J. R., FISHER, B. E., BARANELLO, L., LIU, J. C., KAMBACH, D. M., NIE, Z., KOH, W. S., LUO,
J., STOMMEL, J. M., LEVENS, D. & BATCHELOR, E. 2017. Global Inhibition with Specific
Activation: How p53 and MYC Redistribute the Transcriptome in the DNA Double-Strand
Break Response. *Mol Cell,* 67**,** 1013-1025 e9.
PURMAN, C. E., COLLINS, P. L., PORTER, S. I., SAINI, A., GUPTA, H., SLECKMAN, B. P. & OLTZ, E.
M. 2019. Regional Gene Repression by DNA Double-Strand Breaks in G1 Phase Cells. *Mol*
Cell Biol, 39.
RABUT, G. & PETER, M. 2008. Function and regulation of protein neddylation. 'Protein
modifications: beyond the usual suspects' review series. *EMBO Rep,* 9**,** 969-76.
RAMÍREZ, F., DÜNDAR, F., DIEHL, S., GRÜNING, B. A. & MANKE, T. 2014. deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic acids research,* 42**,** W187-W191.
SCHAUE, D., KACHIKWU, E. L. & MCBRIDE, W. H. 2012. Cytokines in radiobiological responses: a
review. *Radiat Res,* 178**,** 505-23.
SHALEM, O., SANJANA, N. E., HARTENIAN, E., SHI, X., SCOTT, D. A., MIKKELSON, T., HECKL, D.,
EBERT, B. L., ROOT, D. E., DOENCH, J. G. & ZHANG, F. 2014. Genome-scale CRISPR-Cas9
knockout screening in human cells. *Science,* 343**,** 84-87.
SHANBHAG, N. M., RAFALSKA-METCALF, I. U., BALANE-BOLIVAR, C., JANICKI, S. M. &
GREENBERG, R. A. 2010. ATM-dependent chromatin changes silence transcription in cis
to DNA double-strand breaks. *Cell,* 141**,** 970-81.
SHEN, W., LE, S., LI, Y. & HU, F. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q
File Manipulation. *PLoS One,* 11**,** e0163962.
SITTMAN, D. B., GRAVES, R. A. & MARZLUFF, W. F. 1983. Histone mRNA concentrations are
regulated at the level of transcription and mRNA degradation. *Proc Natl Acad Sci U S A,*
80**,** 1849-53.
SOKKA, M., RILLA, K., MIINALAINEN, I., POSPIECH, H. & SYVÄOJA, J. E. 2015. High levels of
TopBP1 induce ATR-dependent shut-down of rRNA transcription and nucleolar
segregation. *Nucleic Acids Res,* 43**,** 4975-89.

STEPHENS, A. S., STEPHENS, S. R. & MORRISON, N. A. 2011. Internal control genes for
quantitative RT-PCR expression analysis in mouse osteoblasts, osteoclasts and
macrophages. *BMC Res Notes,* 4**,** 410.
SU, C., GAO, G., SCHNEIDER, S., HELT, C., WEISS, C., O'REILLY, M. A., BOHMANN, D. & ZHAO, J.
2004. DNA damage induces downregulation of histone gene expression through the G1
checkpoint pathway. *EMBO J,* 23**,** 1133-43.
THORVALDSDÓTTIR, H., ROBINSON, J. T. & MESIROV, J. P. 2013. Integrative Genomics Viewer
(IGV): high-performance genomics data visualization and exploration. *Briefings in*
bioinformatics, 14**,** 178-192.
TZELEPIS, K., KOIKE-YUSA, H., DE BRAEKELEER, E., LI, Y., METZAKOPIAN, E., DOVEY, O. M.,
MUPO, A., GRINKEVICH, V., LI, M., MAZAN, M., GOZDECKA, M., OHNISHI, S., COOPER, J.,
PATEL, M., MCKERRELL, T., CHEN, B., DOMINGUES, A. F., GALLIPOLI, P., TEICHMANN, S.,
PONSTINGL, H., MCDERMOTT, U., SAEZ-RODRIGUEZ, J., HUNTLY, B. J. P., IORIO, F., PINA,
C., VASSILIOU, G. S. & YUSA, K. 2016. A CRISPR Dropout Screen Identifies Genetic
Vulnerabilities and Therapeutic Targets in Acute Myeloid Leukemia. *Cell Rep,* 17**,** 1193-
1205.
VAN SLUIS, M. & MCSTAY, B. 2017. Nucleolar reorganization in response to rDNA damage. *Curr*
Opin Cell Biol, 46**,** 81-86.
VASSILEV, L. T., TOVAR, C., CHEN, S., KNEZEVIC, D., ZHAO, X., SUN, H., HEIMBROOK, D. C. &
CHEN, L. 2006. Selective small-molecule inhibitor reveals critical mitotic functions of
human CDK1. *Proc Natl Acad Sci U S A,* 103**,** 10660-5.
VENKATA NARAYANAN, I., PAULSEN, M. T., BEDI, K., BERG, N., LJUNGMAN, E. A., FRANCIA, S.,
VELOSO, A., MAGNUSON, B., DI FAGAGNA, F. D., WILSON, T. E. & LJUNGMAN, M. 2017.
Transcriptional and post-transcriptional regulation of the ionizing radiation response by
ATM and p53. *Sci Rep,* 7**,** 43598.
WATADA, E., LI, S., HORI, Y., FUJIKI, K., SHIRAHIGE, K., INADA, T. & KOBAYASHI, T. 2020. Age-
Dependent Ribosomal DNA Variations in Mice. *Mol Cell Biol,* 40.
WU, T., HU, E., XU, S., CHEN, M., GUO, P., DAI, Z., FENG, T., ZHOU, L., TANG, W. & ZHAN, L.
2021. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The*
Innovation, 2**,** 100141.
YU, Z., XU, C., SONG, B., ZHANG, S., CHEN, C., LI, C. & ZHANG, S. 2023. Tissue fibrosis induced by
radiotherapy: current understanding of the molecular mechanisms, diagnosis and
therapeutic advances. *J Transl Med,* 21**,** 708.
ZHAO, J., DYNLACHT, B., IMAI, T., HORI, T. & HARLOW, E. 1998. Expression of NPAT, a novel
substrate of cyclin E-CDK2, promotes S-phase entry. *Genes Dev,* 12**,** 456-61.
ZHAO, J., KENNEDY, B. K., LAWRENCE, B. D., BARBIE, D. A., MATERA, A. G., FLETCHER, J. A. &
HARLOW, E. 2000. NPAT links cyclin E-Cdk2 to the regulation of replication-dependent
histone gene transcription. *Genes Dev,* 14**,** 2283-97.
1073

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 1-figure supplement 1

Figure 1-figure supplement 2

A

B

Figure 2-figure supplement 1

Dose	No EU	No IR	15 m	30 m	1 h	2 h	4 h
2 Gy	314	2690	1440	1261	1282	1640	2349
5 Gy	314	2690	1379	1169	1101	1345	2004
10 Gy	314	2690	1224	1094	1053	1347	1803

Figure 3-figure supplement 1

Figure 3-figure supplement 2

A

rDNA: 1-14,000

B

chr13: 23,525,853-23,624,883

Figure 3-figure supplement 3

Figure 5-figure supplement 1

Figure 5-figure supplement 2

Figure 6-figure supplement 1

A

B

C

Figure 6-figure supplement 2

Figure 7-figure supplement 1