Whole-genome sequencing of major malaria vectors revealed an

evolution of new insecticide resistance variants in a longitudinal

study in Burkina Faso

Mahamadi Kientega[1,3,*], Chris S Clarkson[2], Nouhoun Traoré[1,3], Hui Tin-Yu J[4],

Samantha O'Loughlin[4], Abdoul-Azize Millogo[1,5], Patric S Epopa[1], Franck A Yao[1],

Adrien M G Belem[3], Jon Brenas[2], Alistair Miles[2], Austin Burt[4], Abdoulaye Diabaté[1,*]

[1]Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso 01 BP 545, Burkina Faso ;

[2]Vector Surveillance Programme, Genomic Surveillance Unit, Wellcome Sanger Institute, Hinxton,

Cambridge, UK ;

[3]Université Nazi Boni, Bobo-Dioulasso 01 BP 1091, Burkina Faso ;

[4]Department of Life Sciences, Imperial College, London SW7 2AZ, UK;

[5]Institut des Sciences des Sociétés, Ouagadougou 03 BP 7047, Burkina Faso.

[*]Corresponding authors: npiediab@gmail.com (A.D.), mkient54@gmail.com (M.K.).

Email addresses : *mkient54@gmail.com (Mahamadi Kientega); cc28@sanger.ac.uk (Chris S*

Clarkson); nouhoun89@gmail.com (Nouhoun Traoré); tin-yu.hui11@imperial.ac.uk (Hui Tin-Yu J);

s.oloughlin@imperial.ac.uk (Samantha O'Loughlin); belemamg@hotmail.fr (Adrien M G Belem);

azizemillogo@gmail.com (Abdoul-Azize Millogo); epopastef@yahoo.fr (Patric S Epopa);

afrayao@gmail.com (Franck A. Yao); jon.brenas@bdi.ox.ac.uk (Jon Brenas);

alistair.miles@sanger.ac.uk (Alistair Miles); a.burt@imperial.ac.uk (Burt Austin); npiediab@gmail.com

(Abdoulaye Diabaté).

Abstract

Intensive deployment of insecticide based malaria vector control tools results in the rapid evolution of phenotypes resistant to these chemicals. Understanding this process on the genomic level is essential for the deployment of successful interventions. Using whole genome sequencing data of 1409 individual *An. gambiae* s.l. collected from 2012 to 2017, we investigated the change in genetic structure and the evolution of the insecticide resistance variants in natural populations over time and space. The results showed similar and constant nucleotide diversity and negative Tajima's D between *An. gambiae* s.s. and *An. coluzzii*. PCA and F_{ST} showed a clear genetic structure in the *An. gambiae* s.l. species. Genome-wide F_{ST} and H12 scans identified genomic regions under divergent selection and also having an implication in the adaptations to ecological changes. Novel voltage-gated sodium channel pyrethroid resistance target-site alleles (*V402L, I1527T*) were identified at increasing frequencies alongside the established *kdr* alleles (*Vgsc-L995F, Vgsc-L995S* and *N1570Y*) within the *An. gambiae* s.l. populations. Organophosphate metabolic resistance markers were also identified, at increasing frequencies, within the *An. gambiae* s.s. populations from 2012 to 2017, including the SNP *Ace1-G280S* and its associated duplication. Variants simultaneously identified in the same vector populations raise concerns about the long-term efficacy of new-generation bednets and the recently introduced organophosphate pirimiphos-methyl indoor residual spraying. These findings highlighted the benefit of genomic malaria vector surveillance for the detection of new insecticide resistance variants, the monitoring of the existing resistance variants, and also to get insights into the evolutionary processes driving insecticide resistance.

Author Summary

Genomic surveillance of malaria vectors is crucial for understanding the genetic variation in natural vector populations and also guiding the implementation of novel and innovative vector control tools. Application of sequencing technologies in vector studies provide insights on the genetic and evolutionary phenomena of vectors that could have impact on vector control strategies. By analyzing the whole genome data of 1409 wild *An. gambiae* s.l. mosquito collected between 2012 and 2017, we showed an emergence of novel insecticide resistance markers alongside increasing frequencies of existing insecticide resistance variants over time in Burkina Faso. We showed the benefit of genomic surveillance of malaria vectors for the monitoring of the insecticide resistance variants and also providing insights into the evolutionary processes driving insecticide resistance.

Introduction

Malaria remains a major public health problem in Africa [1]. *Anopheles gambiae* s.l., the main malaria vector in Africa, is phenotypically and genotypically diverse, allowing rapid adaptation to environmental changes, such as introductions of new insecticidal control methods [2]. In Burkina Faso, major malaria vectors *An. gambiae* s.s., *An. coluzzii* and *An. arabiensis* can be found living in sympatry, with diverging resting and feeding behaviours [3,4]. These species, through their ability to colonize different ecological settings, their preference for human blood feeding, and their high susceptibility to a malaria parasite (*Plasmodium falciparum*) infection, are responsible for around 90% of the malaria burden in Burkina Faso and the whole sub-Saharan Africa [5,6].

The National Malaria Control Programme (NMCP) of Burkina Faso and their partners have invested much to increase the coverage of long-lasting insecticide treated bednets (LLINs) and other control tools such as indoor residual spraying (IRS) of insecticides to drop the malaria transmission line in the country. The coverage of LLINs in households increased significantly in the country over the years (5.6% in 2003 to about 80% in 2014) reaching 83% in 2021 [7,8]. The recent 2019 and 2022 bednet campaigns involved the free distribution of three types of LLINs, especially the piperonyl butoxide (PBO)-synergist, the dual-AI Interceptor G2 and the standard LLINs [9]. Despite these achievements, malaria remains a countrywide health problem responsible for about 12.2 million cases and 4355 deaths during the year 2021 [10].

In Burkina Faso and most sub-Saharan countries, the primary vector surveillance strategies relied on the routine tracking of the vector species density and distribution and the monitoring of insecticide resistance (IR) status, unable to properly explain the genetic and evolutionary processes driving the increasing spread of IR variants and

vector behavioral changes [11]. Recent advances in genomic sequencing

technologies, both price and throughput, and new cloud native analysis tools have

made genomic surveillance of malaria vectors possible. Genomic surveillance has

advantages to provide insights on the genetic and evolutionary phenomena of vectors

that could have any impact on vector control strategies. Towards this aim, the

Anopheles gambiae 1000 genomes (Ag1000G) project driven by the MalariaGEN

Vector Observatory sequenced more than 15000 *Anopheles* mosquitoes collected in

25 African countries [12,13]. A major goal of this project was to support malaria

elimination efforts by providing a high quality open access data resource on natural

genetic variation within *An. gambiae* s.s., *An. coluzzii* and *An. arabiensis* populations,

both for the public health and research communities [14].

Previous studies investigating insecticide resistance (IR) using these genomic data

have highlighted associated genetic variants distributed throughout the mosquito

genome [15–18]. The phenotypic effect of many of these variants has been

demonstrated through in-vivo and in-vitro experiments and their effects are evident in

low mortality rate observed during the IR bioassays, and in the reduction of the efficacy

of the insecticide-based control tools [19–21]. The strength of these resistance

phenotypes and the wide geographical distribution of these variants raise concerns

about the durability of the new insecticide-based vector control tools being used to

overcome resistance, including indoor residual spraying (IRS) with organophosphate

insecticides such as Actellic (pirimiphos-methyl) and pyrethroid bed-nets treated with

metabolic resistance blocking piperonyl butoxide (PBO) synergists.

There is an increasingly urgent need for sustainable strategies and effective tools to

support malaria elimination efforts in the face of an ever evolving vector. Research of

alternative and innovative tools such as genetic engineering, new insecticidal

compounds as well as natural symbionts has increased in recent years, giving a

glimmer of hope to people living in endemic areas [22,23]. However, given the genome

complexity of the malaria vectors, highlighted by many studies [12,13,24], any

implementation of vector control tools requires an understanding of the genomic

variation of vector species, particularly an understanding of the evolutionary

processes, emergence and the spread of the variants associated with IR in natural

populations over time. Thus, longitudinal sampling of vectors, followed by whole

genome population genomics becomes an essential tool upstream and downstream

of the implementation of the control tools to provide insights on the genetic and

evolutionary phenomena of vectors that could have any impact on vector control

strategies. Here we leverage the 1409 *Anopheles gambiae* s.l. samples, collected in

Burkina Faso between 2012-2017 and sequenced as part of the *Anopheles gambiae*

1000 Genomes Project (2017, 2020), to characterize the spread of known IR variants

over space and time, while analyzing these natural populations for evidence of the

evolution of novel IR mechanisms which could threaten current or future vector control

methods.

Results

Population Sampling

Following the *Anopheles gambiae* 1000 Genomes project*,* further mosquito collections were sequenced as part of the follow-up project, the MalariaGEN Vector Observatory*.* In total, this resulted in 1409, deep, whole-genome sequenced, mosquitoes collected from three villages surrounding Bobo-Dioulasso city in western Burkina Faso. Mosquitoes were collected longitudinally, across the years 2012, 2014, 2015, 2016 and 2017 (Fig 1), though species representation varies between collection sites and time-points (Table 1). *An. coluz*zii was the most predominant species found, accounting for 52.73% [743/1409] of samples, followed by *An. gambiae s.s.* with 38.96% (549/1409), while *An. arabiensis* only comprised 8.16% (115/1409) of samples collected. Two *An. gambiae* s.s. and *An. coluzzii* hybrids were also identified. As shown by previous work, *An. coluzzii* remains the most predominant species in Bana (83.87% [489/583]) and Souroukoudinga (60.90% [243/399]) while *An. gambiae* s.s. is prevalent in Pala (70.72% [302/427]) [6].

149

150 **<u>Fig 1</u>**. Map of the sampling sites around Bobo-Dioulasso city

153 **Table 1.** Species composition of *An. gambiae* complex in the three villages

154 surrounding Bobo-Dioulasso city over the five collection years

Species	Periods	Sampling sites		
		Bana	Souroukoudinga	Pala
An. gambiae s.s.	2012	22 [33.85%]	28 [49.12%]	48 [81.36%]
	2014	32 [24.81%]	49 [49.49%]	89 [89.%]
	2015	25 [20.66%]	40 [31.5%]	76 [46.34%]
	2016	13 [9.77%]	38 [44.19%]	53 [79.1%]
	2017	0	0	36 [97.3%]
An. coluzzii	2012	42 [64.62%]	29 [50.88%]	11 [18.64%]
	2014	96 [74.42%]	50 [50.51%]	0
	2015	96 [79.34%]	87 [68.5%]	0
	2016	120 [90.23%]	48 [55.81%]	0
	2017	135 [100.%]	29 [96.67%]	0
An. arabiensis	2014	1 [0.78%]	0	11 [11.%]
	2015	0	0	88 [53.66%]
	2016	0	0	14 [20.9%]
	2017	0	1 [3.33%]	0
intermediate_gambiae_coluzzii	2012	1 [1.54%]	0	0
	2017	0	0	1 [2.7%]

Genetic Diversity

Understanding genetic diversity allows us to compare demography of species over time and space. Analyses of these data showed around 27.94 million of segregating sites (~ 91% of biallelic SNPs) in the *An. gambiae* s.l. genome. Approximately 25.68 million of SNPs were identified in the autosomes and 2.262 million of SNPs in the X chromosome. Analysis to evaluate the genetic diversity of this variation (see Methods), demonstrated that nucleotide diversity (θ_π) and Watterson's theta (θ_w), were high and stable over all collection years for *An. gambiae* and *An. coluzzii*, similar to the majority of natural populations across the species range (Ag1000g, 2017) (Fig 2), potentially suggesting similar demographic histories between these two species. In contrast, *An. arabiensis* showed lower diversity, though with apparent increase over collection years. The diversity findings were reflected in a genome-wide negative Tajima's D, caused by an excess of rare variants due to either positive selection and/or rapid demographic changes through population expansion. In addition, the allelic frequency spectrum (Fig S1 & S2) also showed the same trend highlighting the excess of low frequency variants in the *An. gambiae* s.l. genome.

Fig 2. Average of the diversity statistic (A: nucleotide diversity (θ_π), B: Watterson theta (θ_w), C: Tajima's D) in the *An. gambiae* s.l. populations

Population Structure

Understanding population structure between, and within, vector species is not only important to understand due to its implications in gene flow of medically relevant alleles or role out of gene drive control methods, but hidden structure will confound downstream population genomic analyses. Consequently, the genetic structure and the patterns of gene flow were investigated using SNPs on the 3L chromosome (see methods). Pairwise F_{ST} was estimated between species collected in different villages and was found to be low between *An. gambiae* and *An. coluzzii* scaling from 0 to 0.05 (Fig 3A). These results suggest gene flow between *An. coluzzii* and *An. gambiae* s.s.. *An. arabiensis*, however, showed relatively high F_{ST} between both *An. coluzzii* and *An. gambiae*, ranging from 0.26 to 0.30 (Fig 3A), potentially suggesting weaker gene-flow between species. No differentiation ($F_{ST} \sim 0$) was observed between populations of the same species collected in different villages, suggesting no population structure on this scale.

Principal component analysis (PCA) showed a clear divergence between the *An. gambiae* s.l. species. Examining the first two principal components, samples fell into three clusters, where individuals of the same species were more genetically related than to others. No sub-clusters were observed within each species (Fig 3B). No change was observed in the population structure of each species throughout the years, and the sampling sites (Fig S3); however, some other genomic regions were shown to be under divergent selection (see Organophosphate and carbamate resistance: increasing threats to control plus a potential novel IR mechanism, Fig S4). The lack of genetic structure (low $F_{ST} \sim 0.05$) between *An. coluzzii* and *An. gambiae* s.s. showed that these species are capable of sharing advantageous variants in this region Burkina Faso, such those involved in insecticide resistance.

Fig 3. Genetic differentiation and population structure of *An. gambiae* s.l. using the
SNPs identified in the 3L chromosome. A: Pairwise F_{ST} between *An. gambiae* s.l.
populations collected in different villages. No differentiation between populations of
the same species collected in two different villages. B: PCA showing the genetic
structure of *An. gambiae* s.l. populations.

Insecticide Resistance

Pyrethroid Target-site Resistance: concerning increase of new mutations

We investigated the distribution of genetic variants in the voltage gated sodium
channel gene (*Vgsc)*, which codes for the target-site protein of the pyrethroid
insecticides used in bednets). Non-synonymous SNPs in insecticide target site genes,
can lead to insecticide resistance via physically altering the conformation of the target-
site protein, causing reduced insecticide binding [25].

Pyrethroid target-site resistance in *Anopheles gambiae* s.l. has, historically, been
primarily driven by two main "*kdr"* SNPs, causing the amino acid changes *Vgsc-L995F*
and *Vgsc-L995S* [17]. These mutations are widespread, and are at high frequencies
in many regions in Africa [12]. Improvements in sequencing technologies have allowed

the identification of many additional SNPs in the *Vgsc* gene that could enhance the

level of pyrethroid resistance over the time [26,27]. This study identified 634 non-

synonymous SNPs in all the *An. gambiae* s.l. populations (Table S1). The *kdr* allele

Vgsc-L995F was identified at high frequencies (> 30%) in all the populations, and was

found at fixation in *An. gambiae* s.s. collections from 2012. While *An. arabiensis*

populations also showed high frequencies of *Vgsc-L995S* (Fig 4, Fig S5). The *Vgsc-*

N1570Y allele, previously shown to increase the level of pyrethroid resistance when

in concert with *Vgsc-L995F* [26], was also present in both *An. gambiae s.s.* and *An.*

coluzzii populations at around 30% frequency. Some SNPs were found to be species

specific over the years. We identified the alleles *Vgsc-V402L, N1345I, I1527T,*

K1603T, L1667M, P1874S, A1934V and *I1940T* specifically in *An. coluzzii* populations

whereas *Vgsc-V1853I, I1868T, E1597G* were specific to *An. gambiae* s.s. populations

(Fig S5). Most of these species specific alleles were identified at low frequencies.

Fig 4. Heat map showing the frequencies of the non-synonymous SNPs frequencies

(max freq > 0.05) in the *Vgsc* gene in the *An. gambiae s.l.* populations over time. The

X axis shows the *An. gambiae s.l.* populations and the years of collection. The Y axis

shows the non-synonymous SNPs positions in the chromosome 2L. The gradient color

bar shows the distribution of the allelic frequencies.

A *Vgsc* haplotype carrying the apparently linked amino acid substitutions *Vgsc-*

V402L+I1527T was also identified in Burkina Faso *An. coluzzii* (Fig 4 and 5, Appendix

5). A recent *in-vivo* investigation has shown that the *Vgsc-V402L* substitution results

in a pyrethroid resistance phenotype with lower fitness costs than the classic *Vgsc-*

L995F kdr mutation [27]. Upon discovery of *Vgsc-V402L+I1527T,* it was suggested

that this haplotype may be a "relic" resistance haplotype, pre-dating and being

overtaken by a more effective *Vgsc-L995F* carrying haplotype [17]. However, our time

series data shows that the inverse appears to be the case. *Vgsc-V402L+I1527T*

increased from 12.19% in 2012 to 33.12% in 2017 (Fig 4 and 5, Fig S5). The increase

of these alleles coincides with the drop in the frequency of the *Vgsc-L995F* mutation

in *An. coluzzii* populations from 86.58% in 2012 to 66.46% in 2017 (Fig 4 and 5). Our

results and other works [17,27] suggest these two alleles are increasing in frequency

and replacing *Vgsc-L995F* in *An. coluzzii*, but in 2017 had not spread to *An. gambiae*

s.s. The increased emergence of new *kdr* alleles alongside the existing *kdr* mutations

(*Vgsc-L995F* and *Vgsc-L995S*) at high frequencies may increase the resistance level

of vector populations to pyrethroid compounds.

Fig 5. Dynamics of *Vgsc-L995F* and double mutant *Vgsc-V402L+I1527T* allelic

frequencies in the *An. coluzzii* populations.

Pyrethroid Metabolic Resistance: increasing CNV frequencies.

Gene duplications and deletions (copy number variants - CNVs) are known to

influence the expression level of genes by providing additional copies of the original

gene. Consequently, CNVs of genes involved in xenobiotic detoxification can increase

insecticide metabolism, producing a resistance phenotype [15]. To help overcome

pyrethroid resistance, now commonplace in many natural populations, PBO synergist

bednets have recently been introduced. These bednets enhance the pyrethroid

efficacy by inhibiting the cytochrome P450 enzymes that can be responsible for

metabolic pyrethroid resistance [28]. Often, mechanisms driving the resistance in a

given area remain unclear [20], however, whole genome sequencing allows CNV

calling, and the detection of resistance associated alleles [16].

CNVs were identified in 136 P450 genes distributed across the whole genome and

most of these (~ 63.97 %) were gene amplifications (Table S2). The majority of the

CNVs were species-specific, and some were found at increasing frequencies over time

in the three species. Some genes showing low CNVs frequencies in 2012 displayed

increasing CNVs frequencies over the time (Fig 6). This situation was observed in the

Cyp6z cluster of genes (*Cyp6z1, Cyp6z2 and Cyp6z3*), known to be associated with

pyrethroid resistance [29], whose CNVs frequencies increased from 17% in 2012 to

up to 60% in 2017 in *An. gambiae* s.s. populations. Additionally, the *Cyp9k1* gene,

also previously shown to be involved in pyrethroid resistance, showed CNVs at high

frequencies up to 80% from 2012 to 2017. In the *An. coluzzii* populations, CNVs were

identified at frequencies ranging from 50% in 2012 to up to 90% in 2017 within the

Cyp6aa1, Cyp6aa2, Cyp6p15p and *Cyp12f2* genes. Moreover, the *Cyp6af* (*Cyp6af1*

and *Cyp6af2*) and *Cyp9m* cluster genes (*Cyp9m1 and Cyp9m2*) exhibited high CNVs

frequencies (50 – 100%) in all the populations (Fig 6).

Fig 6. Heat map of the cytochrome P450 genes showing high CNVs frequencies in

the *An. gambiae* s.l. populations. The X axis shows the *An. gambiae* s.l. populations

and the years of collection. The Y axis shows the positions of the P450 genes in the

genome. The gradient color bar shows the distribution of the allelic frequencies.

Organophosphate and carbamate resistance: increasing threats to control plus

a potential novel IR mechanism

Indoor residual spraying (IRS) remains another major vector control strategy in Africa [30]. However, the coverage of this strategy dropped considerably over the years because of the emergence of resistance to the pyrethroid compounds and also the low residual efficacy of carbamates compounds [31]. Recently, IRS with Actellic 300 CS containing pirimiphos-methyl, an organophosphate as the active ingredient, showed interesting results including a prolonged residual efficacy up to seven months and a reduction of vector density in many sprayed areas in Africa [32–34]. This new formulation demonstrated better ability to target pyrethroid resistant and indoor resting mosquitoes and showed additional benefit when used in combination with LLINs [35,36].

Using the time series collection, we investigated the magnitude of resistance to organophosphates and carbamates resistance in Burkina Faso that could threaten the long term efficacy of pirimiphos-methyl-based IRS. A sliding window genome-wide F_{ST} scan showed a year to year (from 2012 to 2017) genetic difference in some regions of the genome especially on chromosomes 2 and X in *An. gambiae* s.s. and *An. coluzzii* samples. Most of these changes were observed in genomic regions previously shown to be associated with insecticide-resistance suggesting a signal of positive selection in these genes (Fig 8, 9 and 10).

Both *Ace1* and Diacylglycerol Kinase (DGK) loci showed no selection signal in 2012 in any vector populations we sampled, but by 2017 selection signals were detected at both loci in *An. gambiae* s.s. populations. The increasing gene amplification signal in the *Ace1* locus, from 32.65% in 2012 to up to 91.67% in 2017, was corroborated by the increase in the insecticide resistance associated *Ace1-G280S* variant previously

shown to be involved in organophosphate and carbamate resistance [37] (from

15.81% in 2012 to 65.27% in 2017), but only in *An. gambiae* s.s. populations (Fig 7).

We also identified 243 other non-synonymous SNPs in the *Ace1* gene (Table S3).

CNV amplifications were also found in *An. coluzzii* populations, but at low and constant

frequencies (~ 3%), but not in *An. arabiensis* populations.

Fig 7. Evolution of the *Ace1* and Diacylglycerol Kinase (AGAP000519) CNV

frequencies in the *An. gambiae* s.s. populations.

Genetic variants of the *Gste* genes have also been reported to be involved in

resistance to organophosphates [38]. In this study, we identified CNVs in 41 genes

(24 genes with amplifications and 17 with deletions) of the *Gste* genes. These CNVs

are distributed across the genome (Table S4). Most of these CNVs were found at low

frequencies (< 50%) except the *GstD5* which showed high CNV frequencies (60% to

100%) in all the populations (Fig S6, Table S4).

The carboxylesterases (COEs) enzymes have been reported to be involved in the

rapid metabolization of both pyrethroid and organophosphate compounds [39] and

consequently could have an impact on the benefit of the combined organophosphate-

based IRS and pyrethroid-based LLINs approach. In this study, CNVs were identified

in 39 genes across the genome, most of these CNVs were gene amplifications, only

25.64% were deletions (Table S5). The frequencies of these CNVs were relatively low,

except those of the *Coeae60* gene which were identified at frequencies above 60% in

An. coluzzii populations from 2012 to 2017 (Fig S7). Additionally, a positive selection

signal was also detected in the *Coeae60* gene (Fig 8). The *Coeae*G* cluster genes

(*Coeae2G, Coeae3G, Coeae5G, Coeae6G* and *Coeae7G*) and the *Coeae3H* gene

showed CNVs at increasing frequencies from 11% in 2012 to up to 25% in 2017 in the

An. gambiae s.s. populations. The *Coejhe4E* gene showed CNVs at 32.56%

frequencies in *An. arabiensis*, and also at frequencies higher than 30% in *An. gambiae*

s.s. populations (Fig S7).

The emergence of new forms of insecticide resistance may impact vector control

strategy in unexpected ways. Recent studies reported new forms of resistance,

especially behavioural changes (early and/or outdoor biting) in *An. gambiae*

mosquitoes, in response to the intense use of LLINs causing residual malaria

transmission [40,41]. Here, we detected a signal of positive selection in the DGK locus

and also an increased frequency of CNVs of this gene within the *An. gambiae s.s.*

populations (Fig 9 and 10). No selection signal was detected in vector populations in

2012, with no CNVs identified either. However, in 2017, a signal of positive selection

was detected in the DGK locus, with an associated increasing frequency (from 0.00 in

2012 to 16.67% in 2017) of CNVs in vector populations (Fig 7, 9 and 10). Considering

this evolution pattern of the DGK gene and also its clear role in the adaptation of *C.*

elegans and *Drosophila sp* to environmental change [42,43], this gene may play an

important role in the modulation of *An. gambiae* biology in response to the presence

of vector control tools.

Fig 8. Evolution of the Garud H12 in the chromosome arm 2R of the *An. gambiae s.l.* populations from 2012 to 2017. High values of H12 indicate a signal of positive selection in the corresponding genomic region within the populations: *Ace1: Acetylcholinesterase gene, Cyp6p: Cytochrome P450 gene, COEAE60: Carboxyl-esterase gene, Keap1: Kelch-like ECH-Associated Protein 1.*

Fig 9. Garud H12 statistic showing a signal of positive selection in the *DGK* and the *Cyp9k1* genes within the *An. gambiae* s.l. populations.

Fig 10. Year to year variation of the genetic differentiation between populations of the same species of *An. gambiae* complex collected in 2012 and in 2017. Signals of differentiation were observed in some regions involved in insecticides resistance including *Cyp9k1* and DGK gene.

Discussion

Significant advances have been made in the reduction of malaria burden using LLINs and IRS, but the efficacy of these tools is at risk due to the high adaptability of the vector populations to the anthropogenic ecological changes [30]. A better understanding of the molecular, ecological and evolutionary processes allowing vector populations to adapt to the environmental changes would help in the development of innovative tools to support the current control tools in malaria elimination. In this study, we used whole genome sequencing data to study the genetic diversity, the population structure and the evolution of insecticide resistance variants in the *An. gambiae* s.l. populations in the western part of Burkina Faso.

Bednets – evidence and impacts of target-site and metabolic pyrethroid resistance

The insecticides pressure on malaria vectors lead to an emergence of new insecticide resistance variants that confer fitness benefits to vector populations [44]. This seems to be consistent and evident for the SNPs *Vgsc-V402L+I1527T* which were identified at increasing frequencies in *An. coluzzii* populations. The rapid rise of their frequencies in Burkina Faso *An. coluzzii* populations may be expected given the results of a recent study which revealed the *Vgsc-V402L* produces a pyrethroid resistance phenotype with reduced fitness costs, when compared with *Vgsc-L995F* [27]. The simultaneous emergence of different types of genetic variants around genes and gene clusters known to be associated with pyrethroid metabolism and target site mutations within the *An. gambiae* s.l. populations raise major concerns in the long term efficacy of new pyrethroid insecticide-based control strategies such as LLINs.

Insecticides resistance has been widely studied in Africa, and recent works have reported multiple resistance variants spreading in vectors populations, jeopardizing

NMCP efforts in control and elimination [17,18,45]. The implementation of LLINs next-

generation bed nets incorporating piperonyl butoxide (PBO) aims to strengthen the

efficacy of LLINs by inhibiting the cytochromes P450 enzymes responsible for the

rapid detoxifying of pyrethroid insecticide [46]. The high expression of the P450

enzymes was shown to increase the pyrethroid resistance and pre-exposure to PBO

restores the susceptibility of malaria vector to pyrethroid [28]. In most cases, the main

genetic variants responsible for the insecticide resistance in a given area remain

unclear because of the high heterogeneity of the genetic markers mediating metabolic

resistance. Although PBO inhibits the cytochrome genes, the emergence of additional

new variants could enhance the level of pyrethroid resistance impacting the efficacy

of the new vector control tools.

IRS - evidence and impacts of organophosphates resistance

Another major concern related to this multi-modal resistance is its threat on the

effectiveness of the indoor residual spraying (IRS). This strategy is efficient when it is

used in combination with LLINs because of its ability to target indoor resting pyrethroid

resistant vectors [47]. Although the SNP *Ace1-G280S* has long been identified in *An.*

gambiae populations, its frequencies remained low in the populations probably due to

its fitness cost in association with the kdr mutation [48]. Indeed, the duplication of the

Ace1 gene (*ace1*^{D}) seems to have completely solved the problem of genetic cost

yielded by the SNP *Ace1-G280S* [49]. Recent reports showed a wide distribution of

this duplication in West African *An. coluzzii* and *An. gambiae* s.s. populations [18,49]

probably in response to the use of organophosphate and carbamates compounds in

vector control. Similarly, the duplication of *Ace1* gene follows the same trend as the

SNP *Ace1-G280S* raising up in frequencies in *An. gambiae* s.s. populations in our

study. This clearly showed the close related link between these two variants in

conferring high fitness to the mosquito populations. In addition, the CNVs identified at

high frequencies in some of the *Gste* and COE genes could increase the expression

level of these genes. The simultaneous presence of these variants (CNVs and SNPs)

in *Ace1*, *Gste* and esterases genes within the same vector populations strongly

contribute to increase the survival capacity of the vector against organophosphates

and carbamates. This situation threatens the long-term efficacy of vector control

interventions including the recently introduced pirimiphos-methyl-based IRS.

Whole genome surveillance enables detection of potential novel IR loci

Genome-wide scans for positive selection have proved their usefulness in the

identification of genomic regions under divergent selection. In malaria vectors, under

high insecticide mediated evolutionary pressure, these regions often have implications

in insecticide resistance [50]. We detected a signal of positive selection in a genomic

region of the X chromosome corresponding to the DGK gene. Though there is currently

no evidence of this gene's association with insecticide resistant phenotypes in malaria

vectors, studies in *Drosophila melanogaster* and *Caenorhabditis elegans* showed its

potential involvement in the adaptation to environmental changings [42,43]. In *D.*

melanogaster, the rdgA gene, an orthologue of the DGK gene, is eye-specific, strongly

involved in light signaling. Any genetic variant influencing the functioning of this gene

would probably affect its sensitivity to light, as reported previously [42,51].

Consequently, if light signaling is regulated the same way in mosquito species, the

DGK gene may also be involved in the regulation of *An. gambiae* vision. In *An.*

gambiae, a previous study showed a rhythmic expression of the genes involved in light

signaling under the control of the circadian clock [52]. The expression pattern of these

genes suggests their role in calibration of the mosquito visual system to start the

nocturnal activities such as flight and biting. In response to the high coverage of LLINs,

major changes were reported in vector biting behavior particularly early biting and later

biting activities in many areas of Africa [31,41]. Even though no evidence was

highlighted, these changes could be related to alterations in the expression of genes

involved in the mosquito visual system.

Another possible implication of the DGK gene in the adaptation of vector populations

is its indirect implication to insecticide resistance via the regulation of the acetylcholine

release at synaptic junctions. In *C. elegans*, the DGK gene regulates the amount of

acetylcholine in synaptic junctions and its inactivation causes a high susceptibility to

aldicarb, a carbamate insecticide [43]. This report clearly showed the potential

implication of the DGK gene in organophosphates and carbamates resistance *C.*

elegans. If synaptic junctions are regulated in the same way in mosquitoes, any

selective variant in the DGK gene would affect the acetylcholine release in synapses.

These two hypotheses of the potential implication of the DGK gene in adaptations,

either insecticide resistance or biting time change, were previously drawn up to explain

the selection signal observed in the DGK gene [50]. These hypotheses seem different

in both organisms, but the molecular mechanisms underlying the DGK gene function

remain similar. Given this intriguing role of DGK, further studies are needed to deeply

investigate the role of the DGK gene (AGAP000519) in modulating malaria vector

behavior and its potential involvement in behavioral and genetic resistance.

Conclusion

Investigation of genome variation in vector populations is crucial in understanding dynamics of populations in a given area and guiding strategies of malaria control programs. This study showed a weak differentiation and strong genetic connection between *An. gambiae* and *An. coluzzii* populations in the western area of Burkina Faso. The emergence of new variants together with the already existing pyrethroids resistance variants at high frequencies raise concerns about the long-term efficacy of new-generation nets impregnated with pyrethroids compounds. The long-term efficacy of pirimiphos-methyl-based IRS is also threatened because of the increasing frequency of the SNP *Ace1-G280S* and the duplication of the *Ace1* gene, which confer high fitness to vector populations carrying them. Genome-wide selection scans detected strong signals of positive selection in the genomic regions having an implication in insecticide resistance. This study highlighted the benefit of the molecular surveillance of malaria vectors in the detection of new insecticide resistance variants, the monitoring of the existing resistance variants and also to get insights in evolutionary processes underlying these variants.

Methods

Sample collection

Mosquito samples were collected by the Target Malaria project in three sites, Bana (11.233, -4.472), Souroukoudinga (11.235, -4.535) and Pala (11.150, -4.235), located in the western part of Burkina Faso (see Results - Figure 1). These villages are located in the humid savannah zone surrounding the Bobo-Dioulasso city and are characterized by two seasons: a rainy period from May to October and a dry season from November to April. The average annual rainfall in this area is above 800mm (average annual temperature ~ 27 °C and humidity ~ 60%) [6]. The main agricultural practices are cotton and cereal (maize, rice, sorghum etc.) cultivation utilizing a large amount of pesticides [53].

Malaria remains endemic in the area surrounding the Bobo-dioulasso city, causing 338 deaths in the three health districts of the city [10]. *An. gambiae* complex species remains the major malaria vector in the sampling area [6]. Malaria control relies on the antimalarial drugs and prevention of mosquito bites using long-lasting insecticidal nets (LLINs), the most widespread vector control tool, and to a lesser extent indoor residual spraying (IRS), sprays and mosquito-repellent coils.

Mosquitoes were collected during the rainy and dry seasons from 2012 to 2017 using three collection methods: Pyrethroid spray catches (PSC), human landing catches (HLC) and swarm collection [54]. PSC and HLC were used for female mosquito collection, whereas males were collected in swarms. After collection, mosquito samples were morphologically identified using the morphological keys [55] and *An. gambiae s.l.* specimens were stored in 80% ethanol and then shipped to Wellcome Sanger Institute for the whole genome sequencing.

Whole genome sequencing and genomic data management

An. gambiae s.l. mosquitoes were sequenced individually at high coverage (30x) using Illumina technology at the Wellcome Sanger Institute. Genomic data were analyzed at the MalariaGEN Resource Centre Team using a wide range of genome analysis software. Reads were aligned to the AgamP4 reference genome using BWA version 0.7.15. Indel realignment was performed using GATK version 3.7-0 RealignerTargetCreator and IndelRealigner. Single nucleotide polymorphisms were called using GATK version 3.7-0 UnifiedGenotyper. Genotypes were called for each sample independently, in genotyping mode, given all possible alleles at all genomic sites where the reference base was not "N". Coverage was capped at 250× by random down-sampling. Complete specifications of the alignment and genotyping pipelines are available from the malariagen/pipelines GitHub repository [56]. After variant calling, the raw sequences in FASTQ format and the aligned sequences in BAM format were stored in the European Nucleotide Archive (ENA, Study Accession n° PRJEB42254). The genetic variants in VCF and zarr formats, including the samples' metadata, were stored on Google Cloud and are accessible via the malariagen-data package or are directly downloadable. More details regarding the sequencing technology, the genomic data storage and management including the SNPs variants calling, haplotypes phasing, and copy number variants identification are available on the homepage of MalariaGEN (https://malariagen.github.io/vector-data/ag3/methods.html).

Data analyses

Genetic diversity and population structure were analyzed in the python programming language on the Google Colab platform. The genetic variants (SNPs, haplotypes and

CNVs) and the reference genome of *An. gambiae* s.l. were accessed using the

malariagen-data package for direct analyses of the genomic data in the cloud. Specific

python packages including malariagen-data [57], scikit-allel [58] and other python

standard data-management packages were used for the data analyses. The diversity

statistics (segregating sites, nucleotide diversity (θ_π), Tajima's D (D), Watterson theta

(θ_w), allele frequency spectrum) were computed in the whole chromosome in all the

mosquito populations. Tukey multiple comparison test was applied to compare the

diversity statistic between vector populations over the year. Genome wide selection

scan was performed using the Garud h12 [59] statistic to detect the genomic region

under positive selection. F_{ST} and PCA analyses were performed in each chromosome

to investigate the genetic structure and the differentiation of *An. gambiae* s.l.

populations. The extent and the spread of insecticide resistance variants were also

investigated in the *An. gambiae* s.l. populations using the malariagen_data python

package. Time series (from 2012 to 2017) SNPs frequencies were analyzed in known

target-site resistance associated genes, *Vgsc, Rdl* and *Ace1*. Also copy number

variation (CNV) frequencies were investigated in genomic regions associated with

metabolic resistance, cytochrome P450s, glutathione-s-transferases,

acetylcholinesterase, diacylglycerol kinase and carboxylesterases.

Ethics approval and consent to participate

All methods in this paper have been implemented in accordance with the relevant

guidelines/regulations/legislation in Burkina Faso. No ethics approval was required to

run all the activities related to this paper.

Consent for publication

Not applicable.

Availability of data and materials

Jupyter Notebooks and scripts to reproduce all the analyses, tables and figures are

available in the GitHub repository: https://github.com/mkient/BF_Ag1000G_analyses.

The SNPs and haplotypes data are available on the homepage of MalariaGEN and

can be accessed using the malariagen-data package.

Competing interests

The authors declare no competing interests.

Funding

The sample collection is supported by Target Malaria, which receives core funding

from the Bill and Melinda Gates Foundation and Open Philanthropy. Whole genome

sequencing and genomics data management were supported by MalariaGEN at the

Wellcome Sanger Institute.

Authors' contributions

MK, CSC and AM conceived the study. AB and AD provided the resources. AB, AMGB and

AD supervised the study. AAM, NT, PSA, SOL and FYA carried out samples collection in the

field. AM, CSC and JB produced the genomic data. MK, CSC, HTY and JB carried out data

analysis and visualization. MK and CSC drafted the manuscript. All authors have read and

approved this version of the manuscript.

Acknowledgements

The authors would like to acknowledge the Institut de Recherche en Sciences de la

Santé, its funders and partners for supporting the sample collection. We are grateful

to the Ag1000G project, the MalariaGEN, Sanger Core and GSU Vector Surveillance

Programme at the Wellcome Sanger Institute, and their partners in Africa for the

production of the *An. gambiae* genomic data. The authors also thank the populations

of the sampling sites for their sincere cooperation during the mosquito sampling.

References

1. WHO. World malaria report 2022. Geneva; 2022. Available: https://apps.who.int/iris/rest/bitstreams/1484818/retrieve

2. Hemming-Schroeder E, Zhong D, Machani M, Nguyen H, Thong S, Kahindi S, et al. Ecological drivers of genetic connectivity for African malaria vectors Anopheles gambiae and An. arabiensis. Sci Rep. 2020;10: 19946. doi:10.1038/s41598-020-76248-2

3. Costantini C, Ayala D, Guelbeogo WM, Pombi M, Some CY, Bassole IHN, et al. Living at the edge: Biogeographic patterns of habitat segregation conform to speciation by niche expansion in anopheles gambiae. BMC Ecol. 2009;9: 1–27. doi:10.1186/1472-6785-9-16

4. Fournet F, Cussac M, Ouari A, Meyer PE, Toé HK, Gouagna LC, et al. Diversity in anopheline larval habitats and adult composition during the dry and wet seasons in Ouagadougou (Burkina Faso). Malar J. 2010;9: 1–10. doi:10.1186/1475-2875-9-78

5. Soma DD, Zogo BM, Somé A, Tchiekoi BN, Hien DF de S, Pooda HS, et al. Anopheles bionomics, insecticide resistance and malaria transmission in southwest Burkina Faso: A pre-intervention study. Chitnis N, editor. PLoS One. 2020;15: e0236920. doi:10.1371/journal.pone.0236920

6. Epopa PS, Collins CM, North A, Millogo AA, Benedict MQ, Tripet F, et al. Seasonal malaria vector and transmission dynamics in western Burkina Faso. Malar J. 2019;18: 113. doi:10.1186/s12936-019-2747-5

7. Ministere de la Santé. Rapport De Revue De Performances Du Programme National De Lutte 2011-2015- Burkina Fasso. Ouagadougou; 2016.

8. INSD, ICF. Enquête Démographique et de Santé 2021 : Rapport de synthèse.

Rockville,   Maryland;   2023.   Available:

https://dhsprogram.com/pubs/pdf/SR279/SR279.pdf

9.   U.S. President's Malaria Initiative. Burkina Faso Malaria Operational Plan FY

2023. 2023. Available: www.pmi.gov

10.   Ministère de la Santé. Annuaire statistique 2021. Ouagadougou; 2022.

Available: http://cns.bf/IMG/pdf/annuaire_mshp_2021_signe.pdf

11.   Coulibaly ZI, Gowelo S, Traore I, Mbewe RB, Ngulube W, Olanga EA, et al.

Strengthening adult mosquito surveillance in Africa for disease control: learning

from   the   present.   Curr   Opin   Insect   Sci.   2023;60:   101110.

doi:10.1016/j.cois.2023.101110

12.   The Anopheles gambiae 1000 Genomes Consortium. Genetic diversity of the

African malaria vector Anopheles gambiae. Nature. 2017;552: 96–100.

doi:10.1038/nature24995

13.   The Anopheles gambiae 1000 Genomes Consortium. Genome variation and

population structure among 1142 mosquitoes of the African malaria vector

species Anopheles gambiae and Anopheles coluzzii. Genome Res. 2020;30:

1533–1546. doi:10.1101/gr.262790.120

14.   MalariaGEN. Ag1000G | MalariaGEN. MalariaGEN. 2014. Available:

https://www.malariagen.net/mosquito/ag1000g

15.   Weetman D, Djogbenou LS, Lucas E. Copy number variation (CNV) and

insecticide resistance in mosquitoes: evolving knowledge or an evolving

problem? Curr Opin Insect Sci. 2018;27: 82–88. doi:10.1016/j.cois.2018.04.005

16.   Lucas ER, Miles A, Harding NJ, Clarkson CS, Lawniczak MKN, Kwiatkowski DP,

et al. Whole-genome sequencing reveals high complexity of copy number

variation at insecticide resistance loci in malaria mosquitoes. Genome Res.

2019;29: 1250–1261. doi:10.1101/gr.245795.118

17.   Clarkson CS, Miles A, Harding NJ, O'Reilly AO, Weetman D, Kwiatkowski D, et

653   al. The genetic architecture of target-site resistance to pyrethroid insecticides in

654   the African malaria vectors Anopheles gambiae and Anopheles coluzzii. Mol

655   Ecol. 2021;30: 5303–5317. doi:10.1111/mec.15845

656   18.   Grau-Bové X, Lucas E, Pipini D, Rippon E, van 't Hof AE, Constant E, et al.

657   Resistance to pirimiphos-methyl in West African Anopheles is spreading via

658   duplication and introgression of the Ace1 locus. ffrench-Constant RH, editor.

659   PLOS Genet. 2021;17: e1009253. doi:10.1371/journal.pgen.1009253

660   19.   Amoudji AD, Ahadji-Dabla KM, Hien AS, Apétogbo YG, Yaméogo B, Soma DD,

661   et al. Insecticide resistance profiles of Anopheles gambiae s.l. in Togo and

662   genetic mechanisms involved, during 3-year survey: is there any need for

663   resistance management? Malar J. 2019;18: 177. doi:10.1186/s12936-019-

664   2813-z

665   20.   Moyes CL, Wiebe A, Gleave K, Trett A, Hancock PA, Padonou GG, et al.

666   Analysis-ready datasets for insecticide resistance phenotype and genotype

667   frequency in African malaria vectors. Sci Data. 2019;6. doi:10.1038/s41597-

668   019-0134-2

669   21.   Sovi A, Keita C, Sinaba Y, Dicko A, Traore I, Cisse MBM, et al. Anopheles

670   gambiae (s.l.) exhibit high intensity pyrethroid resistance throughout Southern

671   and Central Mali (2016–2018): PBO or next generation LLINs may provide

672   greater control. Parasit Vectors. 2020;13: 239. doi:10.1186/s13071-020-04100-

673   7

674   22.   Sougoufara S, Ottih EC, Tripet F. The need for new vector control approaches

675   targeting outdoor biting anopheline malaria vector communities. Parasit Vectors.

676   2020;13: 295. doi:10.1186/s13071-020-04170-7

677   23.   Wilson AL, Courtenay O, Kelly-Hope LA, Scott TW, Takken W, Torr SJ, et al.

678   The importance of vector control for the control and elimination of vector-borne

679   diseases. Barrera R, editor. PLoS Negl Trop Dis. 2020;14: e0007831.

680   doi:10.1371/journal.pntd.0007831

681   24.   Neafsey DE, Waterhouse RM, Abai MR, Aganezov SS, Alekseyev MA, Allen JE,

682   et al. Highly evolvable malaria vectors: The genomes of 16 Anopheles

683   mosquitoes. Science (80-). 2015;347. doi:10.1126/science.1258522

684   25.   Hemingway J, Ranson H. Insecticide resistance in insect vectors of human

685   disease.   Annu   Rev   Entomol.   2000;45:   371–91.

686   doi:10.1146/annurev.ento.45.1.371

687   26.   Jones CM, Liyanapathirana M, Agossa FR, Weetman D, Ranson H, Donnelly

688   MJ, et al. Footprints of positive selection associated with a mutation (N1575Y)

689   in the voltage-gated sodium channel of Anopheles gambiae. Proc Natl Acad Sci.

690   2012;109: 6614–6619. doi:10.1073/pnas.1201475109

691   27.   Williams J, Cowlishaw R, Sanou A, Ranson H, Grigoraki L. In vivo functional

692   validation of the V402L voltage gated sodium channel mutation in the malaria

693   vector   An.   gambiae.   Pest   Manag   Sci.   2022;78:   1155–1163.

694   doi:10.1002/ps.6731

695   28.   Protopopoff N, Mosha JF, Lukole E, Charlwood JD, Wright A, Mwalimu CD, et

696   al. Effectiveness of a long-lasting piperonyl butoxide-treated insecticidal net and

697   indoor residual spray interventions, separately and together, against malaria

698   transmitted by pyrethroid-resistant mosquitoes: a cluster, randomised

699   controlled, two-by-two fact. Lancet. 2018;391: 1577–1588. doi:10.1016/S0140-

700   6736(18)30427-6

701 29. Chandor-Proust A, Bibby J, Régent-Kloeckner M, Roux J, Guittard-Crilat E, Poupardin R, et al. The central role of mosquito cytochrome P450 CYP6Zs in insecticide detoxification revealed by functional expression and structural modelling. Biochem J. 2013;455: 75–85. doi:10.1042/BJ20130577

705 30. Ranson H. Current and Future Prospects for Preventing Malaria Transmission via the Use of Insecticides. Cold Spring Harb Perspect Med. 2017;7: a026823. doi:10.1101/cshperspect.a026823

708 31. Sherrard-Smith E, Skarp JE, Beale AD, Fornadel C, Norris LC, Moore SJ, et al. Mosquito feeding behavior and how it influences residual malaria transmission across Africa. Proc Natl Acad Sci. 2019;116: 15086–15095. doi:10.1073/pnas.1820646116

712 32. Kenea O, Balkew M, Tekie H, Deressa W, Loha E, Lindtjørn B, et al. Impact of combining indoor residual spraying and long-lasting insecticidal nets on Anopheles arabiensis in Ethiopia: results from a cluster randomized controlled trial. Malar J. 2019;18: 182. doi:10.1186/s12936-019-2811-1

716 33. Loha E, Deressa W, Gari T, Balkew M, Kenea O, Solomon T, et al. Long-lasting insecticidal nets and indoor residual spraying may not be sufficient to eliminate malaria in a low malaria incidence area: results from a cluster randomized controlled trial in Ethiopia. Malar J. 2019;18: 141. doi:10.1186/s12936-019-2775-1

721 34. Soma DD, Zogo B, Hien DF de S, Hien AS, Kaboré DA, Kientega M, et al. Insecticide resistance status of malaria vectors Anopheles gambiae (s.l.) of southwest Burkina Faso and residual efficacy of indoor residual spraying with microencapsulated pirimiphos-methyl insecticide. Parasit Vectors. 2021;14: 58. doi:10.1186/s13071-020-04563-8

726   35.   Rowland M, Boko P, Odjo A, Asidi A, Akogbeto M, N'Guessan R. A New Long-

727   Lasting Indoor Residual Formulation of the Organophosphate Insecticide

728   Pirimiphos Methyl for Prolonged Control of Pyrethroid-Resistant Mosquitoes: An

729   Experimental Hut Trial in Benin. Vontas J, editor. PLoS One. 2013;8: e69516.

730   doi:10.1371/journal.pone.0069516

731   36.   Hamainza B, Sikaala CH, Moonga HB, Chanda J, Chinula D, Mwenda M, et al.

732   Incremental impact upon malaria transmission of supplementing pyrethroid-

733   impregnated long-lasting insecticidal nets with indoor residual spraying using

734   pyrethroids or the organophosphate, pirimiphos methyl. Malar J. 2016;15: 100.

735   doi:10.1186/s12936-016-1143-7

736   37.   Weill M, Malcolm C, Chandre F, Mogensen K, Berthomieu A, Marquine M, et al.

737   The unique mutation in ace-1 giving high insecticide resistance is easily

738   detectable in mosquito vectors. Insect Mol Biol. 2004;13: 1–7.

739   doi:10.1111/j.1365-2583.2004.00452.x

740   38.   Riveron JM, Yunta C, Ibrahim SS, Djouaka R, Irving H, Menze BD, et al. A single

741   mutation in the GSTe2 gene allows tracking of metabolically based insecticide

742   resistance in a major malaria vector. Genome Biol. 2014;15: R27.

743   doi:10.1186/gb-2014-15-2-r27

744   39.   Cui F, Li M-X, Chang H-J, Mao Y, Zhang H-Y, Lu L-X, et al. Carboxylesterase-

745   mediated insecticide resistance: Quantitative increase induces broader

746   metabolic resistance than qualitative change. Pestic Biochem Physiol.

747   2015;121: 88–96. doi:10.1016/j.pestbp.2014.12.016

748   40.   Moiroux N, Damien GB, Egrot M, Djenontin A, Chandre F, Corbel V, et al.

749   Human Exposure to Early Morning Anopheles funestus Biting Behavior and

750   Personal Protection Provided by Long-Lasting Insecticidal Nets. Vontas J,

editor. PLoS One. 2014;9: e104967. doi:10.1371/journal.pone.0104967

41. Finda MF, Moshi IR, Monroe A, Limwagu AJ, Nyoni AP, Swai JK, et al. Linking

human behaviours and malaria vector biting risk in south-eastern Tanzania.

Carvalho   LH,   editor.   PLoS   One.   2019;14:   e0217414.

doi:10.1371/journal.pone.0217414

42. Masai I, Okazaki A, Hosoya T, Hotta Y. Drosophila retinal degeneration A gene

encodes an eye-specific diacylglycerol kinase with cysteine-rich zinc-finger

motifs and ankyrin repeats. Proc Natl Acad Sci. 1993;90: 11157–11161.

doi:10.1073/pnas.90.23.11157

43. Miller KG, Emerson MD, Rand JB. Goα and Diacylglycerol Kinase Negatively

Regulate the Gqα Pathway in C. elegans. Neuron. 1999;24: 323–333.

doi:10.1016/S0896-6273(00)80847-8

44. Benelli G, Beier JC. Current vector control challenges in the fight against

malaria.   Acta   Trop.   2017;174:   91–96.

doi:10.1016/J.ACTATROPICA.2017.06.028

45. Sougoufara S, Doucouré S, Sembéne PMB, Harry M, Sokhna C. Challenges for

malaria vector control in sub-Saharan Africa: Resistance and behavioral

adaptations in Anopheles populations. J Vector Borne Dis. 2017;54: 4–15.

Available: http://www.ncbi.nlm.nih.gov/pubmed/28352041

46. Feyereisen R. Insect P450 inhibitors and insecticides: challenges and

opportunities. Pest Manag Sci. 2015;71: 793–800. doi:10.1002/ps.3895

47. Okumu FO, Moore SJ. Combining indoor residual spraying and insecticide-

treated nets for malaria control in Africa: a review of possible outcomes and an

outline of suggestions for the future. Malar J. 2011;10: 208. doi:10.1186/1475-

2875-10-208

48.   Djogbénou L, Noel V, Agnew P. Costs of insensitive acetylcholinesterase

777   insecticide resistance for the malaria vector Anopheles gambiae homozygous

778   for the G119S mutation. Malar J. 2010;9: 12. doi:10.1186/1475-2875-9-12

779   49.   Assogba BS, Djogbénou LS, Milesi P, Berthomieu A, Perez J, Ayala D, et al. An

780   ace-1 gene duplication resorbs the fitness cost associated with resistance in

781   Anopheles gambiae, the main malaria mosquito. Sci Rep. 2015;5: 14529.

782   doi:10.1038/srep14529

783   50.   Miles A. Genomic epidemiology of malaria vectors in the Anopheles gambiae

784   species   complex.   University   of   Oxford.   2021.   Available:

785   https://ora.ox.ac.uk/objects/uuid:d0fc0f47-e24f-4804-9022-

786   26c4e3cf1428/download_file?safe_filename=thesis.pdf&type_of_work=Thesis

787   51.   Delgado R, Muñoz Y, Peña-Cortés H, Giavalisco P, Bacigalupo J. Diacylglycerol

788   Activates the Light-Dependent Channel TRP in the Photosensitive Microvilli of

789   Drosophila melanogaster Photoreceptors. J Neurosci. 2014;34: 6679–6686.

790   doi:10.1523/JNEUROSCI.0513-14.2014

791   52.   Rund SSC, Hou TY, Ward SM, Collins FH, Duffield GE. Genome-wide profiling

792   of diel and circadian gene expression in the malaria vector Anopheles gambiae.

793   Proc Natl Acad Sci. 2011;108. doi:10.1073/pnas.1100584108

794   53.   Naré RWA, Savadogo PW, Gnankambary Z, Nacro HB, Sedogo MP. Analyzing

795   Risks Related to the Use of Pesticides in Vegetable Gardens in Burkina Faso.

796   Agric For Fish. 2015;4: 165. doi:10.11648/j.aff.20150404.13

797   54.   Epopa PS, Millogo AA, Collins CM, North AR, Benedict MQ, Tripet F, et al.

798   Anopheles gambiae (s.l.) is found where few are looking: assessing mosquito

799   diversity and density outside inhabited areas using diverse sampling methods.

800   Parasit Vectors. 2020;13: 516. doi:10.1186/s13071-020-04403-9

55.   Gillies MT, Coetzee M. A SUPPLEMENT TO THE ANOPHELINAE OF AFRICA

SOUTH OF THE SAHARA (AFROTROPICAL REGION). The South African

Institute for Medical Research. Johannesburg; 1987.

56.   Malariagen. MalariaGEN pipelines - Pipelines for processing malaria parasite

and mosquito genome sequence data. In: MalariaGEN pipelines - releases 9

v0.0.9 [Internet]. 2022. Available: https://github.com/malariagen/pipelines

57.   Miles A. GitHub - malariagen/malariagen-data-python: A Python package

providing functions for accessing and analysing MalariaGEN data. In:

malariagen / malariagen-data-python [Internet]. 2022 [cited 6 May 2022].

Available: https://github.com/malariagen/malariagen-data-python

58.   Miles A, Bot P i., R. M, Ralph P, Harding N, Pisupati R, et al. Scikit-allel - Explore

and analyse genetic variation. Zenodo. 2021 [cited 6 May 2022].

doi:10.5281/ZENODO.4759368

59.   Garud NR, Messer PW, Buzbas EO, Petrov DA. Recent Selective Sweeps in

North American Drosophila melanogaster Show Signatures of Soft Sweeps.

Copenhaver GP, editor. PLOS Genet. 2015;11: e1005004.

doi:10.1371/journal.pgen.1005004

Supplementary information

<u>**Table S1**</u>. Non-synonymous SNPs frequencies of the *Vgsc* gene (2L: 2358158 -

2431617) within *An. gambiae* s.l. populations.

<u>**Table S2**</u>. Copy number variations of the cytochrome p450 genes and their

frequencies within *An. gambiae* s.l. populations.

<u>**Table S3**</u>. Non-synonymous SNPs frequencies of the *Ace1* gene (2R: 3484107 -

3495790) within An. gambiae s.l. populations.

<u>**Table S4**</u>. Copy number variations of the glutathione-s-transferase genes and their

frequencies within *An. gambiae* s.l. populations.

<u>**Table S5**</u>. Copy number variations of the carboxylesterase genes and their

frequencies within *An. gambiae* s.l. populations.

840 **Fig. S1.** Site frequency spectra in the 3L chromosome of the *An. gambiae* s.s. populations from 2012 to 2016 in the three villages.

843 **Fig.S2.** Site frequency spectra in the 3L chromosome of the *An. coluzzii* populations from 2012 to 2017 in the three villages.

849   **Fig. S3.** PCA showing the year-to-year (from 2012 to 2017) genetic structure of *An. gambiae* s.l. populations using the whole SNPs identified in

850   the 3L chromosome. B. Year-to-year genetic structure of *An. coluzzii* populations; C. Year-to-year genetic structure of *An. gambiae* s.s.

851   populations; D. Year-to-year genetic structure of *An. arabiensis* populations; These figures demonstrated the lack of geographic substructure

852   within each species of the *An. gambiae* s.l. populations collected in different villages over the years.

Fig. S4. Year to year variation of the genetic differentiation between populations of the same species of *An. gambiae* complex collected in 2012 and in 2017. Signals of differentiation were observed in some regions involved in insecticides resistance. *Ace1*: acetylcholinesterase gene, *Cyp6p*: Cytochrome P450 gene, *Keap1*: Kelch-like ECH-associated protein 1, *Vgsc*: voltage-gated sodium channel, *rdl*: resistance to dieldrin gene, *Gste1*: glutathione S-transferases epsilon 1, *Gste2*: glutathione-S-transferases epsilon 2, *Cyp9k1*: Cytochrome P450 gene.

Fig. S5. Heat map showing the evolution of the non-synonymous SNPs frequencies (max freq > 0.05) in the *Vgsc* gene within the *An. gambiae* s.l. populations and the sampling sites. The X axis shows the *An. gambiae* s.l. populations and the years of collection. The Y axis shows the non-synonymous SNPs positions in the chromosome 2L. The gradient color bar shows the distribution of the allelic frequencies. **BN:** Bana, **SK**: Souroukoudinga, **PL**: Pala, **MM**: Monomtenga.

870 **Fig. S6**. Heat map showing the CNVs frequencies (max freq > 0.05) of the glutathione-s-transferase genes in the *An. gambiae* s.l. populations.

871 The X axis shows the *An. gambiae* s.l. populations and the years of collection. The Y axis shows the positions of the glutathione-s-transferase

872 genes in the genome. The gradient color bar shows the distribution of the allelic frequencies.

Fig. S7. Heat map showing the CNVs frequencies (max freq > 0.05) of the carboxylesterase genes in the *An. gambiae* s.l. populations. The X

axis shows the *An. gambiae* s.l. populations and the years of collection. The Y axis shows the positions of the carboxylesterase genes in the

genomes. The gradient color bar shows the distribution of the allelic frequencies.