1 **Whole-genome sequencing of major malaria vectors reveals the evolution**

2 **of new insecticide resistance variants in a longitudinal study in Burkina**

3 **Faso**

4

5 Mahamadi Kientega[1,3], Chris S Clarkson [2], Nouhoun Traoré[1,3], Hui Tin-Yu J[4], Samantha

6 O'Loughlin[4], Azize Millogo[1], Epopa Patric Stephane[1], Franck A Yao[1], Adrien M G Belem[3],

7 Jon Brenas[2], Alistair Miles [2], Austin Burt[4], Abdoulaye Diabaté[1]

8

9 [1]Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso 01 BP 545, Burkina

10 Faso.

11 [2]Vector Surveillance Programme, Genomic Surveillance Unit, Wellcome Sanger Institute,

12 Hinxton, Cambridge, UK

13 [3]Université Nazi Boni, Bobo-Dioulasso 01 BP 1091, Burkina Faso.

14 [4]Department of Life Sciences, Imperial College, London SW7 2AZ, UK

15

16 **Abstract**

17 Intensive deployment of insecticide based malaria vector control tools results in the rapid

18 evolution of phenotypes resistant to these chemicals. Understanding this process on the

19 genomic level is essential for the deployment of successful interventions. Using whole genome

20 sequencing data of 1409 individual *An. gambiae* s.l. collected from 2012 to 2017, we

21 investigated the change in genetic structure and the evolution of the insecticide resistance

22 variants in natural populations over time and space. The results showed similar and constant

23 nucleotide diversity and negative Tajima's D between *An. gambiae* s.s. and *An. coluzzii*. PCA

24 and F_{ST} showed a clear genetic structure in the *An. gambiae* s.l. species. Genome-wide F_{ST} and

25 H12 scans identified genomic regions under divergent selection and also having an implication

in the adaptations to ecological changes. Novel voltage-gated sodium channel pyrethroid

resistance target-site alleles (*V402L, I1527T*) were identified at increasing frequencies

alongside the established *kdr* alleles (*Vgsc-L995F*, *Vgsc-L995S* and *N1570Y*) within the *An.*

gambiae s.l. populations. Organophosphate metabolic resistance markers were also identified,

at increasing frequencies, within the *An. gambiae* s.s. populations from 2012 to 2017, including

the SNP *Ace1-G280S* and its associated duplication. Variants simultaneously identified in the

same vector populations raise concerns about the long-term efficacy of new-generation bednets

and the recently introduced organophosphate pirimiphos-methyl indoor residual spraying.

These findings highlighted the benefit of genomic malaria vector surveillance for the detection

of new insecticide resistance variants, the monitoring of the existing resistance variants, and

also to get insights into the evolutionary processes driving insecticide resistance.

Author Summary

Genomic surveillance of malaria vectors is crucial for understanding the genetic variation in

natural vector populations and also guiding the implementation of novel and innovative vector

control tools. Application of sequencing technologies in vector studies provide insights on the

genetic and evolutionary phenomena of vectors that could have impact on vector control

strategies. By analyzing the whole genome data of 1409 wild *An. gambiae* s.l. mosquito

collected between 2012 and 2017, we showed an emergence of novel insecticide resistance

markers alongside increasing frequencies of existing insecticide resistance variants over time

in Burkina Faso. We showed the benefit of genomic surveillance of malaria vectors for the

monitoring of the insecticide resistance variants and also providing insights into the

evolutionary processes driving insecticide resistance.

Introduction

Malaria remains a major public health problem in Africa (WHO, 2022). *Anopheles gambiae*

s.l., the main malaria vector in Africa, is phenotypically and genotypically diverse, allowing

rapid adaptation to environmental changes, such as introductions of new insecticidal control

methods (Hemming-Schroeder *et al.*, 2020). In Burkina Faso, major malaria vectors *An.*

gambiae s.s., *An. coluzzii* and *An. arabiensis* can be found living in sympatry, with diverging

resting and feeding behaviours (Costantini *et al.*, 2009; Fournet *et al.*, 2010). These species,

through their ability to colonize different ecological settings, their preference for human blood

feeding, and their high susceptibility to a malaria parasite (*Plasmodium falciparum*) infection,

are responsible for around 90% of the malaria burden in Burkina Faso and the whole sub-

Saharan Africa (Epopa *et al.*, 2019; Soma *et al.*, 2020).

The National Malaria Control Programme (NMCP) of Burkina Faso and their partners have

invested much to increase the coverage of long-lasting insecticide treated bednets (LLINs) and

other control tools such as indoor residual spraying (IRS) of insecticides to drop the malaria

transmission line in the country. The coverage of LLINs in households increased significantly

in the country over the years (5.6% in 2003 to about 80% in 2014) reaching 83% in 2021

(Ministere de la Santé, 2016; INSD and ICF, 2023). The recent 2019 and 2022 bednet

campaigns involved the free distribution of three types of LLINs, especially the piperonyl

butoxide (PBO)-synergist, the dual-AI Interceptor G2 and the standard LLINs (U.S. President's

Malaria Initiative, 2023). Despite these achievements, malaria remains a countrywide health

problem responsible for about 12.2 million cases and 4355 deaths during the year 2021

(Ministère de la Santé, 2022).

In Burkina Faso and most sub-Saharan countries, the primary vector surveillance strategies

relied on the routine tracking of the vector species density and distribution and the monitoring

of insecticide resistance (IR) status, unable to properly explain the genetic and evolutionary

processes driving the increasing spread of IR variants and vector behavioral changes (Coulibaly

et al., 2023). Recent advances in genomic sequencing technologies, both price and throughput,

and new cloud native analysis tools have made genomic surveillance of malaria vectors

possible. Genomic surveillance has advantages to provide insights on the genetic and

evolutionary phenomena of vectors that could have any impact on vector control strategies.

Towards this aim, the *Anopheles gambiae* 1000 genomes (Ag1000G) project driven by the

MalariaGEN Vector Observatory sequenced more than 15000 *Anopheles* mosquitoes collected

in 25 African countries (The Anopheles gambiae 1000 Genomes Consortium, 2017, 2020). A

major goal of this project was to support malaria elimination efforts by providing a high quality

open access data resource on natural genetic variation within *An. gambiae* s.s., *An. coluzzii* and

An. arabiensis populations, both for the public health and research communities (MalariaGEN,

2014).

Previous studies investigating insecticide resistance (IR) using these genomic data have

highlighted associated genetic variants distributed throughout the mosquito genome (Weetman

et al., 2018; Lucas *et al.*, 2019; Clarkson *et al.*, 2021; Grau-Bové *et al.*, 2021). The phenotypic

effect of many of these variants has been demonstrated through in-vivo and in-vitro

experiments and their effects are evident in low mortality rate observed during the IR bioassays,

and in the reduction of the efficacy of the insecticide-based control tools (Amoudji *et al.*, 2019;

Moyes *et al.*, 2020; Sovi *et al.*, 2020). The strength of these resistance phenotypes and the wide

geographical distribution of these variants raise concerns about the durability of the new

insecticide-based vector control tools being used to overcome resistance, including indoor

residual spraying (IRS) with organophosphate insecticides such as Actellic (pirimiphos-

methyl) and pyrethroid bed-nets treated with metabolic resistance blocking piperonyl butoxide

(PBO) synergists.

There is an increasingly urgent need for sustainable strategies and effective tools to support

malaria elimination efforts in the face of an ever evolving vector. Research of alternative and

innovative tools such as genetic engineering, new insecticidal compounds as well as natural

symbionts has increased in recent years, giving a glimmer of hope to people living in endemic

areas (Sougoufara *et al.*, 2020; Wilson *et al.*, 2020). However, given the genome complexity

of the malaria vectors, highlighted by many studies (Neafsey *et al.*, 2015; The Anopheles

gambiae 1000 Genomes Consortium, 2017, 2020), any implementation of vector control tools

requires an understanding of the genomic variation of vector species, particularly an

understanding of the evolutionary processes, emergence and the spread of the variants

associated with IR in natural populations over time. Thus, longitudinal sampling of vectors,

followed by whole genome population genomics becomes an essential tool upstream and

downstream of the implementation of the control tools to provide insights on the genetic and

evolutionary phenomena of vectors that could have any impact on vector control strategies.

Here we leverage the 1409 *Anopheles gambiae* s.l. samples, collected in Burkina Faso between

2012-2017 and sequenced as part of the *Anopheles gambiae* 1000 Genomes Project (2017,

2020), to characterize the spread of known IR variants over space and time, while analyzing

these natural populations for evidence of the evolution of novel IR mechanisms which could

threaten current or future vector control methods.

Results

Population Sampling

Following the *Anopheles gambiae* 1000 Genomes project, further mosquito collections were

sequenced as part of the follow-up project, the MalariaGEN Vector Observatory. In total, this

resulted in 1409, deep, whole-genome sequenced, mosquitoes collected from three villages

surrounding Bobo-Dioulasso city in western Burkina Faso. Mosquitoes were collected

longitudinally, across the years 2012, 2014, 2015, 2016 and 2017 (Fig 1), though species

representation varies between collection sites and time-points (Table 1). *An. coluz*zii was the

most predominant species found, accounting for 52.73% [743/1409] of samples, followed by

An. gambiae s.s. with 38.96% (549/1409), while *An. arabiensis* only comprised 8.16%

(115/1409) of samples collected. Two *An. gambiae* s.s. and *An. coluzzii* hybrids were also

identified. As shown by previous work, *An. coluzzii* remains the most predominant species in

Bana (83.87% [489/583]) and Souroukoudinga (60.90% [243/399]) while *An. gambiae* s.s. is

prevalent in Pala (70.72% [302/427]) (Epopa *et al.*, 2019).

Fig 1. Map of the sampling sites around Bobo-Dioulasso city

Table 1. Species composition of *An. gambiae* complex in the three villages surrounding

Bobo-Dioulasso city over the five collection years

Species	Periods	Sampling sites		
		Bana	Souroukoudinga	Pala
An. gambiae s.s.	2012	22 [33.85%]	28 [49.12%]	48 [81.36%]
	2014	32 [24.81%]	49 [49.49%]	89 [89.%]
	2015	25 [20.66%]	40 [31.5%]	76 [46.34%]
	2016	13 [9.77%]	38 [44.19%]	53 [79.1%]
	2017	0	0	36 [97.3%]
An. coluzzii	2012	42 [64.62%]	29 [50.88%]	11 [18.64%]
	2014	96 [74.42%]	50 [50.51%]	0
	2015	96 [79.34%]	87 [68.5%]	0
	2016	120 [90.23%]	48 [55.81%]	0
	2017	135 [100.%]	29 [96.67%]	0
An. arabiensis	2014	1 [0.78%]	0	11 [11.%]
	2015	0	0	88 [53.66%]
	2016	0	0	14 [20.9%]
	2017	0	1 [3.33%]	0
intermediate_gambiae _coluzzii	2012	1 [1.54%]	0	0
	2017	0	0	1 [2.7%]

Genetic Diversity

Understanding genetic diversity allows us to compare demography of species over time and space. Analyses of these data showed around 27.94 million of segregating sites (~ 91% of biallelic SNPs) in the *An. gambiae* s.l. genome. Approximately 25.68 million of SNPs were identified in the autosomes and 2.262 million of SNPs in the X chromosome. Analysis to evaluate the genetic diversity of this variation (see Methods), demonstrated that nucleotide diversity (θ_π) and Watterson's theta (θ_w), were high and stable over all collection years for *An. gambiae* and *An. coluzzii*, similar to the majority of natural populations across the species range (Ag1000g, 2017) (Fig 2), potentially suggesting similar demographic histories between these two species. In contrast, *An. arabiensis* showed lower diversity, though with apparent increase over collection years. The diversity findings were reflected in a genome-wide negative Tajima's D, caused by an excess of rare variants due to either positive selection and/or rapid demographic changes through population expansion. In addition, the allelic frequency spectrum (Fig S1 & S2) also showed the same trend highlighting the excess of low frequency variants in the *An. gambiae* s.l. genome.

Fig 2. Average of the diversity statistic (A: nucleotide diversity (θ_π), B: Watterson theta (θ_w), C: Tajima's D) in the *An. gambiae* s.l. populations

Population Structure

Understanding population structure between, and within, vector species is not only important

to understand due to its implications in gene flow of medically relevant alleles or role out of

gene drive control methods, but hidden structure will confound downstream population

genomic analyses. Consequently, the genetic structure and the patterns of gene flow were

investigated using SNPs on the 3L chromosome (see methods). Pairwise F_{ST} was estimated

between species collected in different villages and was found to be low between *An. gambiae*

and *An. coluzzii* scaling from 0 to 0.05 (Fig 3A). These results suggest gene flow between *An.*

coluzzii and *An. gambiae* s.s.. *An. arabiensis*, however, showed relatively high F_{ST} between

both *An. coluzzii* and *An. gambiae*, ranging from 0.26 to 0.30 (Fig 3A), potentially suggesting

weaker gene-flow between species. No differentiation ($F_{ST} \sim 0$) was observed between

populations of the same species collected in different villages, suggesting no population

structure on this scale.

Principal component analysis (PCA) showed a clear divergence between the *An. gambiae* s.l.

species. Examining the first two principal components, samples fell into three clusters, where

individuals of the same species were more genetically related than to others. No sub-clusters

were observed within each species (Fig 3B). No change was observed in the population

structure of each species throughout the years, and the sampling sites (Fig S3); however, some

other genomic regions were shown to be under divergent selection (see Organophosphate and

carbamate resistance: increasing threats to control plus a potential novel IR mechanism, Fig

S4). The lack of genetic structure (low $F_{ST} \sim 0.05$) between *An. coluzzii* and *An. gambiae* s.s.

showed that these species are capable of sharing advantageous variants in this region Burkina

Faso, such those involved in insecticide resistance.

Fig 3. Genetic differentiation and population structure of *An. gambiae* s.l. using the SNPs

identified in the 3L chromosome. A: Pairwise F_{ST} between *An. gambiae* s.l. populations

collected in different villages. No differentiation between populations of the same species

collected in two different villages. B: PCA showing the genetic structure of *An. gambiae* s.l.

populations.

Insecticide Resistance

Pyrethroid Target-site Resistance: concerning increase of new mutations

We investigated the distribution of genetic variants in the voltage gated sodium channel gene

(*Vgsc)*, which codes for the target-site protein of the pyrethroid insecticides used in bednets).

Non-synonymous SNPs in insecticide target site genes, can lead to insecticide resistance via

physically altering the conformation of the target-site protein, causing reduced insecticide

binding (Hemingway and Ranson, 2000).

Pyrethroid target-site resistance in *Anopheles gambiae* s.l. has, historically, been primarily

driven by two main "*kdr*" SNPs, causing the amino acid changes *Vgsc-L995F* and *Vgsc-L995S*

(Clarkson *et al.*, 2021). These mutations are widespread, and are at high frequencies in many

regions in Africa (The Anopheles gambiae 1000 Genomes Consortium, 2017). Improvements

in sequencing technologies have allowed the identification of many additional SNPs in the

Vgsc gene that could enhance the level of pyrethroid resistance over the time (Jones *et al.*,

2012; Williams *et al.*, 2022). This study identified 634 non-synonymous SNPs in all the *An.*

gambiae s.l. populations (Table S1). The *kdr* allele *Vgsc-L995F* was identified at high

frequencies (> 30%) in all the populations, and was found at fixation in *An. gambiae* s.s.

collections from 2012. While *An. arabiensis* populations also showed high frequencies of *Vgsc-*

L995S (Fig 4, Fig S5). The N1570Y allele, previously shown to increase the level of pyrethroid

resistance when in concert with *Vgsc-L995F* (Jones *et al.*, 2012), was also present in both *An.*

gambiae s.s. and *An. coluzzii* populations at around 30% frequency. Some SNPs were found to

be species specific over the years. We identified the alleles *Vgsc-V402L, N1345I, I1527T,*

K1603T, L1667M, P1874S, A1934V and *I1940T* specifically in *An. coluzzii* populations

whereas *Vgsc-V1853I, I1868T, E1597G* were specific to *An. gambiae* s.s. populations (Fig S5).

Most of these species specific alleles were identified at low frequencies.

Fig 4. Heat map showing the frequencies of the non-synonymous SNPs frequencies (max freq

> 0.05) in the *Vgsc* gene in the *An. gambiae s.l.* populations over time. The X axis shows the

An. gambiae s.l. populations and the years of collection. The Y axis shows the non-synonymous

SNPs positions in the chromosome 2L. The gradient color bar shows the distribution of the

allelic frequencies.

A *Vgsc* haplotype carrying the apparently linked amino acid substitutions *Vgsc-V402L+I1527T*

was also identified in Burkina Faso *An. coluzzii* (Fig 4 and 5, Appendix 5). A recent *in-vivo*

investigation has shown that the *VGSC-V402L* substitution results in a pyrethroid resistance

phenotype with lower fitness costs than the classic *Vgsc-L995F kdr* mutation (Williams *et al.*,

2022). Upon discovery of *Vgsc-V402L+I1527T,* it was suggested that this haplotype may be a

"relic" resistance haplotype, pre-dating and being overtaken by a more effective *Vgsc-L995F*

carrying haplotype (Clarkson *et al.*, 2021). However, our time series data shows that the inverse

appears to be the case. *Vgsc-V402L+I1527T* increased from 12.19% in 2012 to 33.12% in 2017

(Fig 4 and 5, Fig S5). The increase of these alleles coincides with the drop in the frequency of

the *Vgsc-L995F* mutation in *An. coluzzii* populations from 86.58% in 2012 to 66.46% in 2017

(Fig 4 and 5). Our results and other works (Clarkson *et al.*, 2021; Williams *et al.*, 2022) suggest

these two alleles are increasing in frequency and replacing *Vgsc-L995F* in *An. coluzzii*, but in

2017 had not spread to *An. gambiae* s.s. The increased emergence of new *kdr* alleles alongside

the existing *kdr* mutations (*Vgsc-L995F* and *Vgsc-L995S*) at high frequencies may increase the

resistance level of vector populations to pyrethroid compounds.

Fig 5. Dynamics of *Vgsc-L995F* and double mutant *Vgsc-V402L+I1527T* allelic frequencies in

the *An. coluzzii* populations.

Pyrethroid Metabolic Resistance: increasing CNV frequencies.

Gene duplications and deletions (copy number variants - CNVs) are known to influence the

expression level of genes by providing additional copies of the original gene. Consequently,

CNVs of genes involved in xenobiotic detoxification can increase insecticide metabolism,

producing a resistance phenotype (Weetman *et al.*, 2018). To help overcome pyrethroid

resistance, now commonplace in many natural populations, PBO synergist bednets have

recently been introduced. These bednets enhance the pyrethroid efficacy by inhibiting the

cytochrome P450 enzymes that can be responsible for metabolic pyrethroid resistance

(Protopopoff *et al.*, 2018). Often, mechanisms driving the resistance in a given area remain

unclear (Moyes *et al.*, 2020), however, whole genome sequencing allows CNV calling, and the

detection of resistance associated alleles (Lucas *et al.*, 2019).

CNVs were identified in 136 P450 genes distributed across the whole genome and most of

these (~ 63.97 %) were gene amplifications (Table S2). The majority of the CNVs were species-

specific, and some were found at increasing frequencies over time in the three species. Some

genes showing low CNVs frequencies in 2012 displayed increasing CNVs frequencies over the time (Fig 6). This situation was observed in the *Cyp6z* cluster of genes (*Cyp6z1, Cyp6z2 and Cyp6z3*), known to be associated with pyrethroid resistance (refs), whose CNVs frequencies increased from 17% in 2012 to up to 60% in 2017 in *An. gambiae* s.s. populations. Additionally, the *Cyp9k1* gene, also previously shown to be involved in pyrethroid resistance, showed CNVs at high frequencies up to 80% from 2012 to 2017. In the *An. coluzzii* populations, CNVs were identified at frequencies ranging from 50% in 2012 to up to 90% in 2017 within the *Cyp6aa1, Cyp6aa2, Cyp6p15p* and *Cyp12f2* genes. Moreover, the *Cyp6af* (*Cyp6af1* and *Cyp6af2*) and *Cyp9m* cluster genes (*Cyp9m1 and Cyp9m2*) exhibited high CNVs frequencies (50 – 100%) in all the populations (Fig 6).

Fig 6. Heat map of the cytochrome P450 genes showing high CNVs frequencies in the *An. gambiae* s.l. populations. The X axis shows the *An. gambiae* s.l. populations and the years of collection. The Y axis shows the positions of the P450 genes in the genome. The gradient color bar shows the distribution of the allelic frequencies.

Organophosphate and carbamate resistance: increasing threats to control plus a potential

novel IR mechanism

Indoor residual spraying (IRS) remains another major vector control strategy in Africa (Ranson, 2017). However, the coverage of this strategy dropped considerably over the years because of the emergence of resistance to the pyrethroid compounds and also the low residual efficacy of carbamates compounds (Sherrard-Smith *et al.*, 2018). Recently, IRS with Actellic 300 CS containing pirimiphos-methyl, an organophosphate as the active ingredient, showed interesting results including a prolonged residual efficacy up to seven months and a reduction of vector density in many sprayed areas in Africa (Kenea *et al.*, 2019; Loha *et al.*, 2019; Soma *et al.*, 2021). This new formulation demonstrated better ability to target pyrethroid resistant and indoor resting mosquitoes and showed additional benefit when used in combination with LLINs (Rowland *et al.*, 2013; Hamainza *et al.*, 2016).

Using the time series collection, we investigated the magnitude of resistance to organophosphates and carbamates resistance in Burkina Faso that could threaten the long term efficacy of pirimiphos-methyl-based IRS. A sliding window genome-wide F_{ST} scan showed a year to year (from 2012 to 2017) genetic difference in some regions of the genome especially on chromosomes 2 and X in *An. gambiae* s.s. and *An. coluzzii* samples. Most of these changes were observed in genomic regions previously shown to be associated with insecticide-resistance suggesting a signal of positive selection in these genes (Fig 8, 9 and 10).

Both *Ace1* and Diacylglycerol Kinase (DGK) loci showed no selection signal in 2012 in any vector populations we sampled, but by 2017 selection signals were detected at both loci in *An. gambiae* s.s. populations. The increasing gene amplification signal in the *Ace1* locus, from 32.65% in 2012 to up to 91.67% in 2017, was corroborated by the increase in the insecticide resistance associated *Ace1-G280S* variant previously shown to be involved in organophosphate and carbamate resistance (Weill *et al.*, 2004) (from 15.81% in 2012 to 65.27% in 2017), but

only in *An. gambiae* s.s. populations (Fig 7). We also identified 243 other non-synonymous

SNPs in the *Ace1* gene (Table S3). CNV amplifications were also found in *An. coluzzii*

populations, but at low and constant frequencies (~ 3%), but not in *An. arabiensis* populations.

Fig7. Evolution of the *Ace1* and Diacylglycerol Kinase (AGAP000519) CNV frequencies in

the *An. gambiae* s.s. populations.

Genetic variants of the *Gste* genes have also been reported to be involved in resistance to

organophosphates (Riveron *et al.*, 2014). In this study, we identified CNVs in 41 genes (24

genes with amplifications and 17 with deletions) of the *Gste* genes. These CNVs are distributed

across the genome (Table S4). Most of these CNVs were found at low frequencies (< 50%)

except the *GstD5* which showed high CNV frequencies (60% to 100%) in all the populations

(Fig S6, Table S4).

The carboxylesterases (COEs) enzymes have been reported to be involved in the rapid

metabolization of both pyrethroid and organophosphate compounds (Cui *et al.*, 2015) and

consequently could have an impact on the benefit of the combined organophosphate-based IRS

and pyrethroid-based LLINs approach. In this study, CNVs were identified in 39 genes across

the genome, most of these CNVs were gene amplifications, only 25.64% were deletions (Table

S5). The frequencies of these CNVs were relatively low, except those of the *Coeae*60 gene

which were identified at frequencies above 60% in *An. coluzzii* populations from 2012 to 2017

(Fig S7). Additionally, a positive selection signal was also detected in the *Coeae*60 gene (Fig

8). The *Coeae*G* cluster genes (*Coeae2G, Coeae3G, Coeae5G, Coeae6G* and *Coeae7G*) and

the *Coeae3H* gene showed CNVs at increasing frequencies from 11% in 2012 to up to 25% in

2017 in the *An. gambiae* s.s. populations. The *Coejhe4E* gene showed CNVs at 32.56%

frequencies in *An. arabiensis*, and also at frequencies higher than 30% in *An. gambiae* s.s.

populations (Fig S7).

The emergence of new forms of insecticide resistance may impact vector control strategy in

unexpected ways. Recent studies reported new forms of resistance, especially behavioural

changes (early and/or outdoor biting) in *An. gambiae* mosquitoes, in response to the intense

use of LLINs causing residual malaria transmission (Moiroux *et al.*, 2014; Finda *et al.*, 2019).

Here, we detected a signal of positive selection in the DGK locus and also an increased

frequency of CNVs of this gene within the *An. gambiae s.s.* populations (Fig 9 and 10). No

selection signal was detected in vector populations in 2012, with no CNVs identified either.

However, in 2017, a signal of positive selection was detected in the DGK locus, with an

associated increasing frequency (from 0.00 in 2012 to 16.67% in 2017) of CNVs in vector

populations (Fig 7, 9 and 10). Considering this evolution pattern of the DGK gene and also its

clear role in the adaptation of *C. elegans* and *Drosophila sp* to environmental change (Masai

et al., 1993; Miller *et al.*, 1999), this gene may play an important role in the modulation of *An.*

gambiae biology in response to the presence of vector control tools.

Fig 8. Evolution of the Garud H12 in the chromosome arm 2R of the *An. gambiae s.l.* populations from 2012 to 2017. High values of H12 indicate a signal of positive selection in the corresponding genomic region within the populations: *Ace1: Acetylcholinesterase gene, Cyp6p: Cytochrome P450 gene, COEAE60: Carboxyl-esterase gene, Keap1: Kelch-like ECH-Associated Protein 1.*

Fig 9. Garud H12 statistic showing a signal of positive selection in the *DGK* and the *Cyp9k1* genes within the *An. gambiae* s.l. populations.

Fig 10. Year to year variation of the genetic differentiation between populations of the same species of *An. gambiae* complex collected in 2012 and in 2017. Signals of differentiation were observed in some regions involved in insecticides resistance including *Cyp9k1* and DGK gene.

Discussion

Significant advances have been made in the reduction of malaria burden using LLINs and IRS, but the efficacy of these tools is at risk due to the high adaptability of the vector populations to the anthropogenic ecological changes (Ranson, 2017). A better understanding of the molecular, ecological and evolutionary processes allowing vector populations to adapt to the environmental changes would help in the development of innovative tools to support the current control tools in malaria elimination. In this study, we used whole genome sequencing

data to study the genetic diversity, the population structure and the evolution of insecticide

resistance variants in the *An. gambiae* s.l. populations in the western part of Burkina Faso.

Bednets – evidence and impacts of target-site and metabolic pyrethroid resistance

The insecticides pressure on malaria vectors lead to an emergence of new insecticide resistance

variants that confer fitness benefits to vector populations (Benelli and Beier, 2017). This seems

to be consistent and evident for the SNPs *Vgsc-V402L+I1527T* which were identified at

increasing frequencies in *An. coluzzii* populations. The rapid rise of their frequencies in Burkina

Faso *An. coluzzii* populations may be expected given the results of a recent study which

revealed the *Vgsc-V402L* produces a pyrethroid resistance phenotype with reduced fitness

costs, when compared with *Vgsc-L995F* (Williams *et al.*, 2022). The simultaneous emergence

of different types of genetic variants around genes and gene clusters known to be associated

with pyrethroid metabolism and target site mutations within the *An. gambiae* s.l. populations

raise major concerns in the long term efficacy of new pyrethroid insecticide-based control

strategies such as LLINs.

Insecticides resistance has been widely studied in Africa, and recent works have reported

multiple resistance variants spreading in vectors populations, jeopardizing NMCP efforts in

control and elimination (Sougoufara *et al.*, 2017; Clarkson *et al.*, 2021; Grau-Bové *et al.*, 2021).

The implementation of LLINs next-generation bed nets incorporating piperonyl butoxide

(PBO) aims to strengthen the efficacy of LLINs by inhibiting the cytochromes P450 enzymes

responsible for the rapid detoxifying of pyrethroid insecticide (Feyereisen, 2015). The high

expression of the P450 enzymes was shown to increase the pyrethroid resistance and pre-

exposure to PBO restores the susceptibility of malaria vector to pyrethroid (Protopopoff *et al.*,

2018). In most cases, the main genetic variants responsible for the insecticide resistance in a

given area remain unclear because of the high heterogeneity of the genetic markers mediating

metabolic resistance. Although PBO inhibits the cytochrome genes, the emergence of

additional new variants could enhance the level of pyrethroid resistance impacting the efficacy

of the new vector control tools.

IRS - evidence and impacts of organophosphates resistance

Another major concern related to this multi-modal resistance is its threat on the effectiveness

of the indoor residual spraying (IRS). This strategy is efficient when it is used in combination

with LLINs because of its ability to target indoor resting pyrethroid resistant vectors (Okumu

and Moore, 2011). Although the SNP *Ace1-G280S* has long been identified in *An. gambiae*

populations, its frequencies remained low in the populations probably due to its fitness cost in

association with the kdr mutation (Djogbénou *et al.*, 2010). Indeed, the duplication of the *Ace1*

gene (*ace1^D*) seems to have completely solved the problem of genetic cost yielded by the SNP

Ace1-G280S (Assogba *et al.*, 2015). Recent reports showed a wide distribution of this

duplication in West African *An. coluzzii* and *An. gambiae* s.s. populations (Assogba *et al.*,

2015; Grau-Bové *et al.*, 2021) probably in response to the use of organophosphate and

carbamates compounds in vector control. Similarly, the duplication of *Ace1* gene follows the

same trend as the SNP *Ace1-G280S* raising up in frequencies in *An. gambiae* s.s. populations

in our study. This clearly showed the close related link between these two variants in conferring

high fitness to the mosquito populations. In addition, the CNVs identified at high frequencies

in some of the *Gste* and COE genes could increase the expression level of these genes. The

simultaneous presence of these variants (CNVs and SNPs) in *Ace1*, *Gste* and esterases genes

within the same vector populations strongly contribute to increase the survival capacity of the

vector against organophosphates and carbamates. This situation threatens the long-term

efficacy of vector control interventions including the recently introduced pirimiphos-methyl-

based IRS.

Whole genome surveillance enables detection of potential novel IR loci

Genome-wide scans for positive selection have proved their usefulness in the identification of

genomic regions under divergent selection. In malaria vectors, under high insecticide mediated

evolutionary pressure, these regions often have implications in insecticide resistance (Miles,

2021). We detected a signal of positive selection in a genomic region of the X chromosome

corresponding to the DGK gene. Though there is currently no evidence of this gene's

association with insecticide resistant phenotypes in malaria vectors, studies in *Drosophila*

melanogaster and *Caenorhabditis elegans* showed its potential involvement in the adaptation

to environmental changings (Masai *et al.*, 1993; Miller *et al.*, 1999). In *D. melanogaster,* the

rdgA gene, an orthologue of the DGK gene, is eye-specific, strongly involved in light signaling.

Any genetic variant influencing the functioning of this gene would probably affect its

sensitivity to light, as reported previously (Masai *et al.*, 1993; Delgado *et al.*, 2014).

Consequently, if light signaling is regulated the same way in mosquito species, the DGK gene

may also be involved in the regulation of *An. gambiae* vision. In *An. gambiae*, a previous study

showed a rhythmic expression of the genes involved in light signaling under the control of the

circadian clock (Rund *et al.*, 2011). The expression pattern of these genes suggests their role

in calibration of the mosquito visual system to start the nocturnal activities such as flight and

biting. In response to the high coverage of LLINs, major changes were reported in vector biting

behavior particularly early biting and later biting activities in many areas of Africa (Finda *et*

al., 2019; Sherrard-Smith *et al.*, 2019). Even though no evidence was highlighted, these

changes could be related to alterations in the expression of genes involved in the mosquito

visual system.

Another possible implication of the DGK gene in the adaptation of vector populations is its

indirect implication to insecticide resistance via the regulation of the acetylcholine release at

synaptic junctions. In *C. elegans*, the DGK gene regulates the amount of acetylcholine in

synaptic junctions and its inactivation causes a high susceptibility to aldicarb, a carbamate

insecticide (Miller *et al.*, 1999). This report clearly showed the potential implication of the

DGK gene in organophosphates and carbamates resistance *C. elegans*. If synaptic junctions are

regulated in the same way in mosquitoes, any selective variant in the DGK gene would affect

the acetylcholine release in synapses. These two hypotheses of the potential implication of the

DGK gene in adaptations, either insecticide resistance or biting time change, were previously

drawn up to explain the selection signal observed in the DGK gene (Miles, 2021). These

hypotheses seem different in both organisms, but the molecular mechanisms underlying the

DGK gene function remain similar. Given this intriguing role of DGK, further studies are

needed to deeply investigate the role of the DGK gene (AGAP000519) in modulating malaria

vector behavior and its potential involvement in behavioral and genetic resistance.

Conclusion

Investigation of genome variation in vector populations is crucial in understanding dynamics

of populations in a given area and guiding strategies of malaria control programs. This study

showed a weak differentiation and strong genetic connection between *An. gambiae* and *An.*

coluzzii populations in the western area of Burkina Faso. The emergence of new variants

together with the already existing pyrethroids resistance variants at high frequencies raise

concerns about the long-term efficacy of new-generation nets impregnated with pyrethroids

compounds. The long-term efficacy of pirimiphos-methyl-based IRS is also threatened because

of the increasing frequency of the SNP *Ace1-G280S* and the duplication of the *Ace1* gene,

which confer high fitness to vector populations carrying them. Genome-wide selection scans

detected strong signals of positive selection in the genomic regions having an implication in

insecticide resistance. This study highlighted the benefit of the molecular surveillance of

malaria vectors in the detection of new insecticide resistance variants, the monitoring of the

existing resistance variants and also to get insights in evolutionary processes underlying these

variants.

Methods

Sample collection

Mosquito samples were collected by the Target Malaria project in three sites, Bana (11.233, -

4.472), Souroukoudinga (11.235, -4.535) and Pala (11.150, -4.235), located in the western

part of Burkina Faso (see Results - Figure 1). These villages are located in the humid

savannah zone surrounding the Bobo-Dioulasso city and are characterized by two seasons: a

rainy period from May to October and a dry season from November to April. The average

annual rainfall in this area is above 800mm (average annual temperature ~ 27 °C and

humidity ~ 60%) (Epopa *et al.*, 2019). The main agricultural practices are cotton and cereal

(maize, rice, sorghum etc.) cultivation utilizing a large amount of pesticides (Naré *et al.*,

2015).

Malaria remains endemic in the area surrounding the Bobo-dioulasso city, causing 338 deaths

in the three health districts of the city (Ministère de la Santé, 2022). *An. gambiae* complex

species remains the major malaria vector in the sampling area (Epopa *et al.*, 2019). Malaria

control relies on the antimalarial drugs and prevention of mosquito bites using long-lasting

insecticidal nets (LLINs), the most widespread vector control tool, and to a lesser extent

indoor residual spraying (IRS), sprays and mosquito-repellent coils.

Mosquitoes were collected during the rainy and dry seasons from 2012 to 2017 using three

collection methods: Pyrethroid spray catches (PSC), human landing catches (HLC) and

swarm collection (Epopa *et al.*, 2020). PSC and HLC were used for female mosquito

collection, whereas males were collected in swarms. After collection, mosquito samples were

morphologically identified using the morphological keys (Gillies and Coetzee, 1987) and *An.*

gambiae s.l. specimens were stored in 80% ethanol and then shipped to Wellcome Sanger

Institute for the whole genome sequencing.

Whole genome sequencing and genomic data management

An. gambiae s.l. mosquitoes were sequenced individually at high coverage (30x) using Illumina

technology at the Wellcome Sanger Institute. Genomic data were analyzed at the MalariaGEN

Resource Centre Team using a wide range of genome analysis software. Reads were aligned to

the AgamP4 reference genome using BWA version 0.7.15. Indel realignment was performed

using GATK version 3.7-0 RealignerTargetCreator and IndelRealigner. Single nucleotide

polymorphisms were called using GATK version 3.7-0 UnifiedGenotyper. Genotypes were

called for each sample independently, in genotyping mode, given all possible alleles at all

genomic sites where the reference base was not "N". Coverage was capped at 250× by random

down-sampling. Complete specifications of the alignment and genotyping pipelines are

available from the malariagen/pipelines GitHub repository (Malariagen, 2022). After variant

calling, the raw sequences in FASTQ format and the aligned sequences in BAM format were

stored in the European Nucleotide Archive (ENA, Study Accession n° PRJEB42254). The

genetic variants in VCF and zarr formats, including the samples' metadata, were stored on

Google Cloud and are accessible via the malariagen-data package or are directly downloadable.

More details regarding the sequencing technology, the genomic data storage and management

including the SNPs variants calling, haplotypes phasing, and copy number variants

identification are available on the homepage of MalariaGEN

(https://malariagen.github.io/vector-data/ag3/methods.html).

Data analyses

Genetic diversity and population structure were analyzed in the python programming language

on the Google Colab platform. The genetic variants (SNPs, haplotypes and CNVs) and the

reference genome of *An. gambiae* s.l. were accessed using the malariagen-data package for

direct analyses of the genomic data in the cloud. Specific python packages including

malariagen-data (Miles, 2022), scikit-allel (Miles *et al.*, 2021) and other python standard data-

management packages were used for the data analyses. The diversity statistics (segregating

sites, nucleotide diversity (θ_π), Tajima's D (D), Watterson theta (θ_w), allele frequency

spectrum) were computed in the whole chromosome in all the mosquito populations. Tukey

multiple comparison test was applied to compare the diversity statistic between vector

populations over the year. Genome wide selection scan was performed using the Garud h12

(Garud *et al.*, 2015) statistic to detect the genomic region under positive selection. F_{ST} and PCA

analyses were performed in each chromosome to investigate the genetic structure and the

differentiation of *An. gambiae* s.l. populations. The extent and the spread of insecticide

resistance variants were also investigated in the *An. gambiae* s.l. populations using the

malariagen_data python package. Time series (from 2012 to 2017) SNPs frequencies were

analyzed in known target-site resistance associated genes, *Vgsc, Rdl* and *Ace1*. Also copy

number variation (CNV) frequencies were investigated in genomic regions associated with

metabolic resistance, cytochrome P450s, glutathione-s-transferases, acetylcholinesterase,

diacylglycerol kinase and carboxylesterases.

Ethics approval and consent to participate

All methods in this paper have been implemented in accordance with the relevant

guidelines/regulations/legislation in Burkina Faso. No ethics approval was required to run all

the activities related to this paper.

Consent for publication

Availability of data and materials

Jupyter Notebooks and scripts to reproduce all the analyses, tables and figures are available in

the GitHub repository: https://github.com/mkient/BF_Ag1000G_analyses. The SNPs and

haplotypes data are available on the homepage of MalariaGEN and can be accessed using the

malariagen-data package.

Competing interests

Funding

The sample collection is supported by Target Malaria, which receives core funding from the

Bill and Melinda Gates Foundation and Open Philanthropy. Whole genome sequencing and

genomics data management were supported by MalariaGEN at the Wellcome Sanger Institute.

Authors' contributions

Conceptualization:

Project administration:

Resources:

Investigation:

Supervision:

Methodology:

Data curation:

Formal analysis:

Software:

Validation:

Visualization:

Writing—original draft:

Writing—review & editing:

All authors have read and agreed to the published version of the manuscript.

Acknowledgements

The authors would like to acknowledge the Institut de Recherche en Sciences de la Santé, its

funders and partners for supporting the sample collection. We are grateful to the Ag1000G

project, the MalariaGEN, Sanger Core and GSU Vector Surveillance Programme at the

Wellcome Sanger Institute, and their partners in Africa for the production of the *An. gambiae*

genomic data. The authors also thank the populations of the sampling sites for their sincere

cooperation during the mosquito sampling.

References

Amoudji Adjovi D., Ahadji-Dabla Koffi M., Hien Aristide Sawdetuo, Apétogbo Yawo

Georges, Yaméogo Bienvenu, Soma Diloma Dieudonné, Bamogo Rabila, Atcha-Oubou

Rachid Tinah, Dabiré Roch Kounbobr and Ketoh Guillaume Koffivi (2019) 'Insecticide

resistance profiles of Anopheles gambiae s.l. in Togo and genetic mechanisms involved,

during 3-year survey: is there any need for resistance management?', *Malaria Journal*, 18(1),

p. 177. doi: 10.1186/s12936-019-2813-z.

Assogba Benoît S., Djogbénou Luc S., Milesi Pascal, Berthomieu Arnaud, Perez Julie, Ayala

Diego, Chandre Fabrice, Makoutodé Michel, Labbé Pierrick and Weill Mylène (2015) 'An

ace-1 gene duplication resorbs the fitness cost associated with resistance in Anopheles

gambiae, the main malaria mosquito', *Scientific Reports*, 5(1), p. 14529. doi:

10.1038/srep14529.

Benelli Giovanni and Beier John C. (2017) 'Current vector control challenges in the fight

against malaria', *Acta Tropica*. Elsevier, 174, pp. 91–96. doi:

10.1016/J.ACTATROPICA.2017.06.028.

Clarkson Chris S., Miles Alistair, Harding Nicholas J., O'Reilly Andrias O., Weetman David,

Kwiatkowski Dominic and Donnelly Martin J. (2021) 'The genetic architecture of target-site

resistance to pyrethroid insecticides in the African malaria vectors Anopheles gambiae and

Anopheles coluzzii', *Molecular Ecology*, 30(21), pp. 5303–5317. doi: 10.1111/mec.15845.

Costantini Carlo, Ayala Diego, Guelbeogo Wamdaogo M., Pombi Marco, Some Corentin Y.,

Bassole Imael H. N., Ose Kenji, Fotsing Jean Marie, Sagnon N'Falé, Fontenille Didier,

Besansky Nora J. and Simard Frédéric (2009) 'Living at the edge: Biogeographic patterns of

habitat segregation conform to speciation by niche expansion in anopheles gambiae', *BMC*

Ecology, 9, pp. 1–27. doi: 10.1186/1472-6785-9-16.

Coulibaly Zanakoungo I., Gowelo Steve, Traore Issouf, Mbewe Rex B., Ngulube Willy,

Olanga Evelyn A., DePina Adilson J., Sanou Antoine, Coleman Sylvester and Tangena Julie-

Anne A. (2023) 'Strengthening adult mosquito surveillance in Africa for disease control:

learning from the present', *Current Opinion in Insect Science*, 60, p. 101110. doi:

10.1016/j.cois.2023.101110.

Cui Feng, Li Mei-Xia, Chang Hai-Jing, Mao Yun, Zhang Han-Ying, Lu Li-Xia, Yan Shuai-

Guo, Lang Ming-Lin, Liu Li and Qiao Chuan-Ling (2015) 'Carboxylesterase-mediated

insecticide resistance: Quantitative increase induces broader metabolic resistance than

qualitative change', *Pesticide Biochemistry and Physiology*, 121, pp. 88–96. doi:

10.1016/j.pestbp.2014.12.016.

Delgado Ricardo, Muñoz Yorka, Peña-Cortés Hugo, Giavalisco Patrick and Bacigalupo Juan

(2014) 'Diacylglycerol Activates the Light-Dependent Channel TRP in the Photosensitive

Microvilli of *Drosophila melanogaster* Photoreceptors', *The Journal of Neuroscience*,

34(19), pp. 6679–6686. doi: 10.1523/JNEUROSCI.0513-14.2014.

Djogbénou Luc, Noel Valérie and Agnew Philip (2010) 'Costs of insensitive

acetylcholinesterase insecticide resistance for the malaria vector *Anopheles gambiae*

homozygous for the G119S mutation', *Malaria Journal*, 9(1), p. 12. doi: 10.1186/1475-2875-

9-12.

Epopa Patric Stephane, Collins Catherine Matilda, North Ace, Millogo Abdoul Azize,

Benedict Mark Quentin, Tripet Frederic and Diabate Abdoulaye (2019) 'Seasonal malaria

vector and transmission dynamics in western Burkina Faso', *Malaria Journal*, 18(1), p. 113.

doi: 10.1186/s12936-019-2747-5.

Epopa Patric Stephane, Millogo Abdoul Azize, Collins Catherine Matilda, North Ace R.,

Benedict Mark Quentin, Tripet Frederic, O'Loughlin Samantha, Dabiré Roch K., Ouédraogo

Georges Anicet and Diabaté Abdoulaye (2020) 'Anopheles gambiae (s.l.) is found where few

are looking: assessing mosquito diversity and density outside inhabited areas using diverse

sampling methods', *Parasites & Vectors*, 13(1), p. 516. doi: 10.1186/s13071-020-04403-9.

Feyereisen René (2015) 'Insect P450 inhibitors and insecticides: challenges and

opportunities', *Pest Management Science*, 71(6), pp. 793–800. doi: 10.1002/ps.3895.

Finda Marceline F., Moshi Irene R., Monroe April, Limwagu Alex J., Nyoni Anna P., Swai

Johnson K., Ngowo Halfan S., Minja Elihaika G., Toe Lea P., Kaindoa Emmanuel W.,

Coetzee Maureen, Manderson Lenore and Okumu Fredros O. (2019) 'Linking human

behaviours and malaria vector biting risk in south-eastern Tanzania', *PLOS ONE*. Edited by

Luzia Helena Carvalho, 14(6), p. e0217414. doi: 10.1371/journal.pone.0217414.

Fournet Florence, Cussac Maud, Ouari Ali, Meyer Pierre Erwann, Toé Hyacinthe K.,

Gouagna Louis Clément and Dabiré Roch K. (2010) 'Diversity in anopheline larval habitats

and adult composition during the dry and wet seasons in Ouagadougou (Burkina Faso)',

Malaria Journal, 9(1), pp. 1–10. doi: 10.1186/1475-2875-9-78.

Garud Nandita R., Messer Philipp W., Buzbas Erkan O. and Petrov Dmitri A. (2015) 'Recent

Selective Sweeps in North American Drosophila melanogaster Show Signatures of Soft

Sweeps', *PLOS Genetics*. Edited by Gregory P. Copenhaver, 11(2), p. e1005004. doi:

10.1371/journal.pgen.1005004.

Gillies M. T. and Coetzee M. (1987) *A SUPPLEMENT TO THE ANOPHELINAE OF*

AFRICA SOUTH OF THE SAHARA (AFROTROPICAL REGION), *The South African*

Institute for Medical Research. Johannesburg.

Grau-Bové Xavier, Lucas Eric, Pipini Dimitra, Rippon Emily, van 't Hof Arjèn E., Constant

Edi, Dadzie Samuel, Egyir-Yawson Alexander, Essandoh John, Chabi Joseph, Djogbénou

Luc, Harding Nicholas J., Miles Alistair, Kwiatkowski Dominic, Donnelly Martin J. and

Weetman David (2021) 'Resistance to pirimiphos-methyl in West African Anopheles is

spreading via duplication and introgression of the Ace1 locus', *PLOS Genetics*. Edited by

Richard H. ffrench-Constant, 17(1), p. e1009253. doi: 10.1371/journal.pgen.1009253.

Hamainza Busiku, Sikaala Chadwick H., Moonga Hawela B., Chanda Javan, Chinula

Dingani, Mwenda Mulenga, Kamuliwo Mulakwa, Bennett Adam, Seyoum Aklilu and Killeen

Gerry F. (2016) 'Incremental impact upon malaria transmission of supplementing pyrethroid-

impregnated long-lasting insecticidal nets with indoor residual spraying using pyrethroids or

the organophosphate, pirimiphos methyl', *Malaria Journal*, 15(1), p. 100. doi:

10.1186/s12936-016-1143-7.

Hemingway J. and Ranson H. (2000) 'Insecticide resistance in insect vectors of human

disease.', *Annual review of entomology*, 45, pp. 371–91. doi: 10.1146/annurev.ento.45.1.371.

Hemming-Schroeder Elizabeth, Zhong Daibin, Machani Maxwell, Nguyen Hoan, Thong

Sarah, Kahindi Samuel, Mbogo Charles, Atieli Harrysone, Githeko Andrew, Lehmann Tovi,

Kazura James W. and Yan Guiyun (2020) 'Ecological drivers of genetic connectivity for

African malaria vectors Anopheles gambiae and An. arabiensis', *Scientific Reports*, 10(1), p.

19946. doi: 10.1038/s41598-020-76248-2.

INSD and ICF (2023) *Enquête Démographique et de Santé 2021 : Rapport de synthèse*.

Rockville, Maryland. Available at: https://dhsprogram.com/pubs/pdf/SR279/SR279.pdf.

Jones C. M., Liyanapathirana M., Agossa F. R., Weetman D., Ranson H., Donnelly M. J. and Wilding C. S. (2012) 'Footprints of positive selection associated with a mutation (N1575Y) in the voltage-gated sodium channel of Anopheles gambiae', *Proceedings of the National Academy of Sciences*, 109(17), pp. 6614–6619. doi: 10.1073/pnas.1201475109.

Kenea Oljira, Balkew Meshesha, Tekie Habte, Deressa Wakgari, Loha Eskindir, Lindtjørn Bernt and Overgaard Hans J. (2019) 'Impact of combining indoor residual spraying and long-lasting insecticidal nets on Anopheles arabiensis in Ethiopia: results from a cluster randomized controlled trial', *Malaria Journal*, 18(1), p. 182. doi: 10.1186/s12936-019-2811-1.

Loha Eskindir, Deressa Wakgari, Gari Taye, Balkew Meshesha, Kenea Oljira, Solomon Tarekegn, Hailu Alemayehu, Robberstad Bjarne, Assegid Meselech, Overgaard Hans J. and Lindtjørn Bernt (2019) 'Long-lasting insecticidal nets and indoor residual spraying may not be sufficient to eliminate malaria in a low malaria incidence area: results from a cluster randomized controlled trial in Ethiopia', *Malaria Journal*, 18(1), p. 141. doi: 10.1186/s12936-019-2775-1.

Lucas Eric R., Miles Alistair, Harding Nicholas J., Clarkson Chris S., Lawniczak Mara K. N., Kwiatkowski Dominic P., Weetman David and Donnelly Martin J. (2019) 'Whole-genome sequencing reveals high complexity of copy number variation at insecticide resistance loci in malaria mosquitoes', *Genome Research*, 29(8), pp. 1250–1261. doi: 10.1101/gr.245795.118.

Malariagen (2022) *MalariaGEN pipelines - Pipelines for processing malaria parasite and mosquito genome sequence data*, *MalariaGEN pipelines - releases 9 v0.0.9*. Available at: https://github.com/malariagen/pipelines.

MalariaGEN (2014) *Ag1000G | MalariaGEN*, *MalariaGEN*. Available at: https://www.malariagen.net/mosquito/ag1000g (Accessed: 1 May 2023).

Masai I., Okazaki A., Hosoya T. and Hotta Y. (1993) 'Drosophila retinal degeneration A

gene encodes an eye-specific diacylglycerol kinase with cysteine-rich zinc-finger motifs and

ankyrin repeats.', *Proceedings of the National Academy of Sciences*, 90(23), pp. 11157–

11161. doi: 10.1073/pnas.90.23.11157.

Miles Alistair (2021) *Genomic epidemiology of malaria vectors in the Anopheles gambiae*

species complex. Available at: https://ora.ox.ac.uk/objects/uuid:d0fc0f47-e24f-4804-9022-

26c4e3cf1428/download_file?safe_filename=thesis.pdf&type_of_work=Thesis.

Miles Alistair (2022) *GitHub - malariagen/malariagen-data-python: A Python package*

providing functions for accessing and analysing MalariaGEN data., *malariagen /*

malariagen-data-python. Available at: https://github.com/malariagen/malariagen-data-python

(Accessed: 6 May 2022).

Miles Alistair, Bot Pyup. i., R. Murillo, Ralph Peter, Harding Nick, Pisupati Rahul, Rae

Summer and Millar Tim (2021) 'Scikit-allel - Explore and analyse genetic variation', *Zenodo*.

doi: 10.5281/ZENODO.4759368.

Miller Kenneth G., Emerson Melanie D. and Rand James B. (1999) 'Goα and Diacylglycerol

Kinase Negatively Regulate the Gqα Pathway in C. elegans', *Neuron*, 24(2), pp. 323–333.

doi: 10.1016/S0896-6273(00)80847-8.

Ministere de la Santé (2016) *Rapport De Revue De Performances Du Programme National*

De Lutte 2011-2015- Burkina Fasso. Ouagadougou.

Ministère de la Santé (2022) *Annuaire statistique 2021*. Ouagadougou. Available at:

http://cns.bf/IMG/pdf/annuaire_mshp_2021_signe.pdf.

Moiroux Nicolas, Damien Georgia B., Egrot Marc, Djenontin Armel, Chandre Fabrice,

Corbel Vincent, Killeen Gerry F. and Pennetier Cédric (2014) 'Human Exposure to Early

Morning Anopheles funestus Biting Behavior and Personal Protection Provided by Long-

Lasting Insecticidal Nets', *PLoS ONE*. Edited by John Vontas, 9(8), p. e104967. doi:

10.1371/journal.pone.0104967.

Moyes Catherine L., Athinya Duncan K., Seethaler Tara, Battle Katherine E., Sinka Marianne, Hadi Melinda P., Hemingway Janet, Coleman Michael and Hancock Penelope A. (2020) 'Evaluating insecticide resistance across African districts to aid malaria control decisions', *Proceedings of the National Academy of Sciences*, 117(36), pp. 22042–22050. doi: 10.1073/pnas.2006781117.

Naré Rayim Wendé Alice, Savadogo Paul Windinpsidi, Gnankambary Zacharia, Nacro Hassan Bismarck and Sedogo Michel Papaoba (2015) 'Analyzing Risks Related to the Use of Pesticides in Vegetable Gardens in Burkina Faso', *Agriculture, Forestry and Fisheries*, 4(4), p. 165. doi: 10.11648/j.aff.20150404.13.

Neafsey Daniel E., Waterhouse Robert M., Abai Mohammad R., Aganezov Sergey S., Alekseyev Max A., Allen James E., Amon James, Arcà Bruno, Arensburger Peter, Artemov Gleb, Assour Lauren A., Basseri Hamidreza, Berlin Aaron, Birren Bruce W., Blandin Stephanie A., Brockman Andrew I., Burkot Thomas R., Burt Austin, Chan Clara S., *et al.* (2015) 'Highly evolvable malaria vectors: The genomes of 16 Anopheles mosquitoes', *Science*, 347(6217). doi: 10.1126/science.1258522.

Okumu Fredros O. and Moore Sarah J. (2011) 'Combining indoor residual spraying and insecticide-treated nets for malaria control in Africa: a review of possible outcomes and an outline of suggestions for the future', *Malaria Journal*, 10(1), p. 208. doi: 10.1186/1475-2875-10-208.

Protopopoff Natacha, Mosha Jacklin F., Lukole Eliud, Charlwood Jacques D., Wright Alexandra, Mwalimu Charles D., Manjurano Alphaxard, Mosha Franklin W., Kisinza William, Kleinschmidt Immo and Rowland Mark (2018) 'Effectiveness of a long-lasting piperonyl butoxide-treated insecticidal net and indoor residual spray interventions, separately and together, against malaria transmitted by pyrethroid-resistant mosquitoes: a cluster, randomised controlled, two-by-two fact', *The Lancet*, 391(10130), pp. 1577–1588. doi:

10.1016/S0140-6736(18)30427-6.

Ranson Hilary (2017) 'Current and Future Prospects for Preventing Malaria Transmission via

the Use of Insecticides', *Cold Spring Harbor Perspectives in Medicine*, 7(11), p. a026823.

doi: 10.1101/cshperspect.a026823.

Riveron Jacob M., Yunta Cristina, Ibrahim Sulaiman S., Djouaka Rousseau, Irving Helen,

Menze Benjamin D., Ismail Hanafy M., Hemingway Janet, Ranson Hilary, Albert Armando

and Wondji Charles S. (2014) 'A single mutation in the GSTe2 gene allows tracking of

metabolically based insecticide resistance in a major malaria vector', *Genome Biology*, 15(2),

p. R27. doi: 10.1186/gb-2014-15-2-r27.

Rowland Mark, Boko Pelagie, Odjo Abibatou, Asidi Alex, Akogbeto Martin and N'Guessan

Raphael (2013) 'A New Long-Lasting Indoor Residual Formulation of the Organophosphate

Insecticide Pirimiphos Methyl for Prolonged Control of Pyrethroid-Resistant Mosquitoes: An

Experimental Hut Trial in Benin', *PLoS ONE*. Edited by John Vontas, 8(7), p. e69516. doi:

10.1371/journal.pone.0069516.

Rund Samuel S. C., Hou Tim Y., Ward Sarah M., Collins Frank H. and Duffield Giles E.

(2011) 'Genome-wide profiling of diel and circadian gene expression in the malaria vector

Anopheles gambiae', *Proceedings of the National Academy of Sciences*, 108(32). doi:

10.1073/pnas.1100584108.

Sherrard-Smith Ellie, Griffin Jamie T., Winskill Peter, Corbel Vincent, Pennetier Cédric,

Djénontin Armel, Moore Sarah, Richardson Jason H., Müller Pie, Edi Constant, Protopopoff

Natacha, Oxborough Richard, Agossa Fiacre, N'Guessan Raphael, Rowland Mark and

Churcher Thomas S. (2018) 'Systematic review of indoor residual spray efficacy and

effectiveness against Plasmodium falciparum in Africa', *Nature Communications*, 9(1), p.

4982. doi: 10.1038/s41467-018-07357-w.

Sherrard-Smith Ellie, Skarp Janetta E., Beale Andrew D., Fornadel Christen, Norris Laura C.,

Moore Sarah J., Mihreteab Selam, Charlwood Jacques Derek, Bhatt Samir, Winskill Peter,

Griffin Jamie T. and Churcher Thomas S. (2019) 'Mosquito feeding behavior and how it

influences residual malaria transmission across Africa', *Proceedings of the National*

Academy of Sciences, 116(30), pp. 15086–15095. doi: 10.1073/pnas.1820646116.

Soma Dieudonné Diloma, Poda Serge Bèwadéyir, Hien Aristide Sawdetuo, Namountougou

Moussa, Sangaré Ibrahim, Sawadogo John Marie Emmanuel, Fournet Florence, Ouédraogo

Georges Anicet, Diabaté Abdoulaye, Moiroux Nicolas and Dabiré Roch Kounbobr (2021)

'Malaria vectors diversity, insecticide resistance and transmission during the rainy season in

peri-urban villages of south-western Burkina Faso', *Malaria Journal*, 20(1), p. 63. doi:

10.1186/s12936-020-03554-5.

Soma Dieudonné Diloma, Zogo Barnabas Mahugnon, Somé Anthony, Tchiekoi Bertin

N'Cho, Hien Domonbabele François de Sales, Pooda Hermann Sié, Coulibaly Sanata,

Gnambani Jacques Edounou, Ouari Ali, Mouline Karine, Dahounto Amal, Ouédraogo

Georges Anicet, Fournet Florence, Koffi Alphonsine Amanan, Pennetier Cédric, Moiroux

Nicolas and Dabiré Roch Kounbobr (2020) 'Anopheles bionomics, insecticide resistance and

malaria transmission in southwest Burkina Faso: A pre-intervention study', *PLOS ONE*.

Edited by Nakul Chitnis, 15(8), p. e0236920. doi: 10.1371/journal.pone.0236920.

Sougoufara Seynabou, Doucouré Souleymane, Sembéne Pape M. Back., Harry Myriam and

Sokhna Cheikh (2017) 'Challenges for malaria vector control in sub-Saharan Africa:

Resistance and behavioral adaptations in Anopheles populations', *Journal of Vector Borne*

Diseases, 54(1), pp. 4–15. Available at: http://www.ncbi.nlm.nih.gov/pubmed/28352041.

Sougoufara Seynabou, Ottih Emmanuel Chinweuba and Tripet Frederic (2020) 'The need for

new vector control approaches targeting outdoor biting anopheline malaria vector

communities', *Parasites & Vectors*, 13(1), p. 295. doi: 10.1186/s13071-020-04170-7.

Sovi Arthur, Keita Chitan, Sinaba Youssouf, Dicko Abdourhamane, Traore Ibrahim, Cisse

Moussa B. M., Koita Ousmane, Dengela Dereje, Flatley Cecilia, Bankineza Elie, Mihigo

Jules, Belemvire Allison, Carlson Jenny, Fornadel Christen and Oxborough Richard M.

(2020) 'Anopheles gambiae (s.l.) exhibit high intensity pyrethroid resistance throughout

Southern and Central Mali (2016–2018): PBO or next generation LLINs may provide greater

control', *Parasites & Vectors*, 13(1), p. 239. doi: 10.1186/s13071-020-04100-7.

The Anopheles gambiae 1000 Genomes Consortium (2017) 'Genetic diversity of the African

malaria vector Anopheles gambiae', *Nature*, 552(7683), pp. 96–100. doi:

10.1038/nature24995.

The Anopheles gambiae 1000 Genomes Consortium (2020) 'Genome variation and

population structure among 1142 mosquitoes of the African malaria vector species Anopheles

gambiae and Anopheles coluzzii', *Genome Research*, 30(10), pp. 1533–1546. doi:

10.1101/gr.262790.120.

U.S. President's Malaria Initiative (2023) *Burkina Faso Malaria Operational Plan FY 2023*.

Available at: www.pmi.gov.

Weetman David, Djogbenou Luc S. and Lucas Eric (2018) 'Copy number variation (CNV)

and insecticide resistance in mosquitoes: evolving knowledge or an evolving problem?',

Current Opinion in Insect Science, 27, pp. 82–88. doi: 10.1016/j.cois.2018.04.005.

Weill M., Malcolm C., Chandre F., Mogensen K., Berthomieu A., Marquine M. and

Raymond M. (2004) 'The unique mutation in ace-1 giving high insecticide resistance is easily

detectable in mosquito vectors', *Insect Molecular Biology*, 13(1), pp. 1–7. doi:

10.1111/j.1365-2583.2004.00452.x.

WHO (2022) *World malaria report 2022*. Geneva. Available at:

https://apps.who.int/iris/rest/bitstreams/1484818/retrieve.

Williams Jessica, Cowlishaw Ruth, Sanou Antoine, Ranson Hilary and Grigoraki Linda

(2022) 'In vivo functional validation of the V402L voltage gated sodium channel mutation in

the malaria vector An. gambiae', *Pest Management Science*, 78(3), pp. 1155–1163. doi:

10.1002/ps.6731.

Wilson Anne L., Courtenay Orin, Kelly-Hope Louise A., Scott Thomas W., Takken Willem,

Torr Steve J. and Lindsay Steve W. (2020) 'The importance of vector control for the control

and elimination of vector-borne diseases', *PLOS Neglected Tropical Diseases*. Edited by

Roberto Barrera, 14(1), p. e0007831. doi: 10.1371/journal.pntd.0007831.

Supplementary information

Table S1. Non-synonymous SNPs frequencies of the *Vgsc* gene (2L: 2358158 - 2431617)

within *An. gambiae* s.l. populations.

Table S2. Copy number variations of the cytochrome p450 genes and their frequencies within

An. gambiae s.l. populations.

Table S3. Non-synonymous SNPs frequencies of the *Ace1* gene (2R: 3484107 - 3495790)

within An. gambiae s.l. populations.

Table S4. Copy number variations of the glutathione-s-transferase genes and their frequencies

within *An. gambiae* s.l. populations.

Table S5. Copy number variations of the carboxylesterase genes and their frequencies within

An. gambiae s.l. populations.

Fig. S1. Site frequency spectra in the 3L chromosome of the *An. gambiae* s.s. populations from 2012 to 2016 in the three villages.

Fig.S2. Site frequency spectra in the 3L chromosome of the *An. coluzzii* populations from 2012 to 2017 in the three villages.

844   **Fig. S3.** PCA showing the year-to-year (from 2012 to 2017) genetic structure of *An. gambiae* s.l. populations using the whole SNPs identified in the 3L

845   chromosome. B. Year-to-year genetic structure of *An. coluzzii* populations; C. Year-to-year genetic structure of *An. gambiae* s.s. populations; D. Year-to-year

846   genetic structure of *An. arabiensis* populations; These figures demonstrated the lack of geographic substructure within each species of the *An. gambiae* s.l.

847   populations collected in different villages over the years.

Fig. S4. Year to year variation of the genetic differentiation between populations of the same species of *An. gambiae* complex collected in 2012 and in 2017. Signals of differentiation were observed in some regions involved in insecticides resistance. *Ace1*: acetylcholinesterase gene, *Cyp6p*: Cytochrome P450 gene, *Keap1*: Kelch-like ECH-associated protein 1, *Vgsc*: voltage-gated sodium channel, *rdl*: resistance to dieldrin gene, *Gste1*: glutathione S-transferases epsilon 1, *Gste2*: glutathione-S-transferases epsilon 2, *Cyp9k1*: Cytochrome P450 gene.

Fig. S5. Heat map showing the evolution of the non-synonymous SNPs frequencies (max freq > 0.05) in the *Vgsc* gene within the *An. gambiae* s.l. populations

and the sampling sites. The X axis shows the *An. gambiae* s.l. populations and the years of collection. The Y axis shows the non-synonymous SNPs positions

in the chromosome 2L. The gradient color bar shows the distribution of the allelic frequencies. **BN:** Bana, **SK**: Souroukoudinga, **PL**: Pala, **MM**: Monomtenga.

Fig. S6. Heat map showing the CNVs frequencies (max freq > 0.05) of the glutathione-s-transferase genes in the *An. gambiae* s.l. populations. The X axis shows

the *An. gambiae* s.l. populations and the years of collection. The Y axis shows the positions of the glutathione-s-transferase genes in the genome. The gradient

color bar shows the distribution of the allelic frequencies.

Fig. S7. Heat map showing the CNVs frequencies (max freq > 0.05) of the carboxylesterase genes in the *An. gambiae* s.l. populations. The X axis shows the

An. gambiae s.l. populations and the years of collection. The Y axis shows the positions of the carboxylesterase genes in the genomes. The gradient color bar

shows the distribution of the allelic frequencies.

871