Hammerhead-type FXR agonists induce an eRNA *FincoR* that ameliorates nonalcoholic steatohepatitis in mice

Jinjing Chen[a, #], Ruoyu Wang[b, #], Feng Xiong[b], Hao Sun[a], Byron Kemper[a], Wenbo Li[b,*] and Jongsook Kim Kemper[a,*]

[a]Department of Molecular and Integrative Physiology, University of Illinois at Urbana-Champaign, Urbana, IL, USA and [b]Department of Biochemistry and Molecular Biology, McGovern Medical School, University of Texas Health Science Center, Houston, TX, USA

Running Title: Role of an FXR-induced eRNA, *FincoR*, in NASH

[#]JC and RW contributed equally to this study.

Present Address: Jinjing Chen, Department of Pharmacology and Toxicology, University of Arizona, Tucson, AZ, USA

* To whom correspondence may be addressed:
* Wenbo Li, Department of Biochemistry and Molecular Biology, McGovern Medical School, University of Texas Health Science Center, Houston, TX, USA
Tel: 713-500-6103, e-mail: Wenbo.Li@uth.tmc.edu
* J. Kim Kemper, Department of Molecular and Integrative Physiology,

24   University of Illinois at Urbana-Champaign, Urbana, IL

25   Tel: 217-722-7961, e-mail: jongsook@illinois.edu

Author Contributions:

31   JC, RW, WL and JK designed research; JC, RW, WL, FX, and HS performed

32   experiments; RW and WL performed bioinformatic analyses; JC, RW, WL, and BK

33   analyzed data; and JC, RW, BK, WL, and JK wrote the paper.

Abstract

The nuclear receptor, Farnesoid X Receptor (FXR/NR1H4), is increasingly recognized as a promising drug target for metabolic diseases, including nonalcoholic steatohepatitis (NASH). Protein coding genes regulated by FXR are well known, but whether FXR also acts through regulation of long non-coding RNAs (lncRNAs), which vastly outnumber protein-coding genes, remains unknown. Utilizing RNA-seq and GRO-seq analyses in mouse liver, we found that FXR activation affects the expression of many RNA transcripts from chromatin regions bearing enhancer features. Among these we discovered a previously unannotated liver-enriched enhancer-derived lncRNA (eRNA), termed *FincoR*. We show that *FincoR* is specifically induced by the hammerhead-type FXR agonists, including GW4064 and tropifexor. CRISPR/Cas9-mediated liver-specific knockdown of *FincoR* in dietary NASH mice reduced the beneficial effects of tropifexor, an FXR agonist currently in clinical trials for NASH and primary biliary cholangitis (PBC), indicating that that amelioration of liver fibrosis and inflammation in NASH treatment by tropifexor is mediated in part by *FincoR*. Overall, our findings highlight that pharmacological activation of FXR by hammerhead-type agonists induces a novel eRNA, *FincoR*, contributing to the amelioration of NASH in mice. *FincoR* may represent a new drug target for addressing metabolic disorders, including NASH.

Introduction

Nonalcoholic fatty liver disease (NAFLD) is the most common chronic liver disease and a leading cause of liver transplants and liver-related death *(Friedman et al., 2018)*. NAFLD begins with simple steatosis but may further progress to a severe form, non-alcoholic steatohepatitis (NASH), and later, fatal cirrhosis and liver cancer *(Friedman et al., 2018)*. Despite its striking global increase and clinical importance, there is no approved drug for NASH. The urgent need for development of therapeutic agents for NASH has greatly increased research interest in the nuclear receptor, Farnesoid X Receptor (FXR, NR1H4) *(Evans & Mangelsdorf, 2014)*.

FXR is activated by its physiological ligands, bile acids (BAs), and regulates expression of genes involved in BA, lipid, and glucose metabolism and hepatic autophagy, which maintain metabolite levels and metabolic homeostasis *(Calkin & Tontonoz, 2012; Kliewer & Mangelsdorf, 2015; Lee et al., 2006; Lee et al., 2014; Seok et al., 2014)*. Ligand-activated FXR also protects against hepatic inflammation and liver injury *(Jung et al., 2020; Wang et al., 2008)*. The action of FXR, similar to other nuclear receptors, is achieved primarily by its binding to chromatin loci to regulate the transcription of target genes *(Calkin & Tontonoz, 2012; Lee et al., 2006; Lee et al., 2012; Thomas et al., 2010)*. Consistent with its crucial physiological functions, FXR is increasingly recognized as a promising drug target, particularly for liver diseases, such as NASH and primary biliary cholangitis (PBC) *(Abenavoli et al., 2018; Ali et al., 2015; Downes et al., 2003; Kremoser, 2021)*. For example, semi-synthetic or non-steroidal synthetic agonists of FXR, including obeticholic acid (OCA) and hammerhead-type agonists, such as tropifexor and cilofexor, are currently in clinical trials for NASH and PBC patients *(Abenavoli et al., 2018; Kremoser,*

2021; Sanyal et al., 2023; Tully et al., 2017). However, how pharmacological activation of

FXR mediates such beneficial therapeutic effects is poorly understood.

Non-protein-coding RNAs (ncRNAs) are one of the fascinating discoveries of

modern biology *(Cech & Steitz, 2014)*. While a significant portion of the genome was

initially thought to be "junk DNA", it has been established that many non-coding regions

give rise to functional non-coding RNAs *(Cech & Steitz, 2014)*. Of these ncRNAs, long

non-coding RNAs (lncRNAs) are a group of transcripts longer than 200 nucleotides and

play important roles in diverse biological processes *(Lam et al., 2014; Li et al., 2016;*

Mattick et al., 2023; Sartorelli & Lauberth, 2020; Statello et al., 2021). A group of lncRNAs

are produced from genomic regions bearing epigenetic features of enhancers *(Hon et al.,*

2017; Mattick et al., 2023). This is consistent with the idea that many transcriptional

enhancers actively transcribe noncoding RNAs that are referred to as eRNAs (>85k in

humans and >57k in mice) *(Hirabayashi et al., 2019)*, some of which are functionally

important for enhancer functions *(Lai & Shiekhattar, 2014; Li et al., 2016; Sartorelli &*

Lauberth, 2020). A majority of eRNAs have not yet been annotated in current lncRNA

databases, such as GENCODE. Exploring eRNA landscapes and functions in diverse

biology and disease states will facilitate our understanding of both lncRNAs and

enhancers *(Li et al., 2016; Mattick et al., 2023)*. Indeed, the landscape of eRNAs in mouse

liver has been minimally explored *(Fang et al., 2014)*, and has not been well-studied in

response to specific nuclear receptor activation, such as by FXR. Moreover, functional

roles of eRNAs *in vivo* in intact organisms is understudied.

FXR was shown to regulate expression of small non-coding microRNA (miR)

genes, for example *miR-34a and miR-802 (Lee et al., 2010; Seok et al., 2021)*, but it has

not been reported whether FXR achieves its function through regulating lncRNAs, which

far outnumber miRs. Because FXR directly regulates expression of its target genes *(Lee*

et al., 2012; Thomas et al., 2010), we examined if FXR regulates enhancer-derived

lncRNAs (eRNAs), and if such transcripts participate in its physiological or

pharmacological functions. By utilizing RNA-seq and GRO-seq analyses of livers from

mice treated with FXR ligands, we identified a set of FXR-regulated eRNAs. Among these,

we focused on a highly induced and abundantly expressed eRNA that we referred to as

FXR-induced non-coding RNA (*FincoR*) for functional studies. *FincoR* is highly enriched

in mouse liver and is induced specifically by hammerhead-type FXR agonists, including

GW4064 and tropifexor. *In vivo* studies utilizing CRISPR/Cas9-mediated liver-specific

knockdown of *FincoR* in dietary NASH mice indicated that *FincoR* is critically involved in

mediating the beneficial pharmacological effects of tropifexor in reducing liver fibrosis and

inflammation.

Results

Activation of FXR by GW4064 induces a novel eRNA, *FincoR*, in mouse liver

To identify enhancer-derived lncRNAs (eRNAs) potentially regulated by FXR, we first

obtained a list of putative enhancers in mouse liver based on ENCODE H3K27ac ChIP-

Seq data (ENCFF001KMI, see **methods**). Then, we performed ribo-depleted total RNA-

seq in the livers from mice treated with a specific FXR agonist, GW4064, to identify

transcripts produced from these enhancer regions (see **Supplementary Table S1**). To

avoid confounding issues of RNA signals from genes, we specifically focused on

intergenic enhancer regions (+/- 3kb from H3K27ac peak center) that harbor discernible

RNA-Seq signals (RPKM > 1).

This genomic analysis resulted in identification of 190 high-confidence eRNAs in

mouse liver. Among these, 14 eRNAs were upregulated and 5 were downregulated by

GW4064 treatment (FDR < 0.05, Log2 FC > 1, **Figure 1A**) (see **Supplementary Table**

S2). FXR-regulated eRNAs were produced adjacent to many genes with important roles

in liver metabolism and disease, for example, *Hes1* (**Figure 1-figure supplement 1A, B**).

One of the most robustly induced eRNAs was <u>F</u>XR-<u>i</u>nduced <u>n</u>on-<u>co</u>ding <u>R</u>NA (*FincoR)*,

an unannotated novel transcript located on chromosome 19 (**Figure 1A, B, C**). Reverse

transcription qPCR (RT-qPCR) confirmed that treatment with GW4064 substantially

induced expression of *FincoR,* more than 10-fold in mouse liver, which is similar to

induction of *Shp*, a well-known FXR target gene (**Figure 1D**) *(Claudel et al., 2005; Evans*

& Mangelsdorf, 2014; Goodwin et al., 2000). Induction of *FincoR* by GW4064 was

transient, peaked within 1 h and then declined gradually, a pattern similar to that of *Shp*

(**Figure 1-figure supplement 1C**). Expression of genes adjacent to *FincoR*, including

Gcnt1, Rfk, Pcsk5 and Prune2, did not change after acute 1 h FXR activation as shown

by RNA-seq (**Figure 1C**), and confirmed for *Gcnt1* by time course qPCR (**Figure 1-figure**

supplement 1C).

We also found that the short-time GW4064 treatment resulted in 590 up-regulated

genes and 500 down-regulated genes (FDR < 0.05) (**Figure 1-figure supplement 2A**).

GO enrichment analysis of these differentially expressed genes (DEGs) revealed their

roles in the regulation of triglyceride, fatty acid, and cholesterol metabolism (**Figure 1-**

figure supplement 2B), which is consistent with known roles of FXR in these

physiological processes *(Claudel et al., 2005)*.

Ligand-activated FXR directly activates transcription of eRNAs, including *FincoR*

We sought to examine, 1) if these eRNAs were directly activated by FXR, and 2) if the

activation takes place transcriptionally using FXR liver-specific knockout (FXR-LKO) mice

that were treated with GW4064 to determine if FXR was required for induction of eRNAs

by GW4064. FXR-LKO mice were generated from FXR floxed mice (FXR-flox) **(Figure 2-**

figure supplement 1A) and transcription of eRNAs was detected by global run-on

sequencing (GRO-seq), a widely used method to detect nascent RNA transcription,

including eRNAs *(Lam et al., 2014; Li et al., 2016; Sartorelli & Lauberth, 2020)*. GRO-seq

showed that FXR-induced eRNAs were activated transcriptionally by GW4064 in FXR-

flox mice, but such induction was abolished in FXR-LKO mice (**Figure 2A**). In particular,

FincoR is robustly induced in the GRO-Seq analysis and its induction is dependent on

hepatic FXR (**Figure 2B**). RT-qPCR confirmed the FXR-dependent expression of *FincoR,*

similar to that of *Shp* (**Figure 2-figure supplement 1B**).

We next examined if FXR binds to the enhancers that produce the identified

eRNAs by analyzing published mouse liver ChIP-Seq data for FXR (**Supplementary**

Table S3) *(Lee et al., 2012; Thomas et al., 2010)*. FXR binding was strongly enriched at

the enhancers associated with FXR-induced eRNAs as were the enhancer marks

H3K27ac and H3K4me1 (**Figure 2-figure supplement 1C**). The binding of FXR and the

presence of histone marks at the *FincoR* enhancer region determined by ChIP-seq as

compared with nascent transcripts detected by Gro-seq is shown in **Figure 2B**. These

analyses support the conclusion that activation of FXR transcriptionally induces this

series of eRNAs via chromatin binding at these enhancers, including *FincoR*.

We validated FXR binding at the enhancer region that produces *FincoR* using

mouse liver ChIP (**Figure 2C**). We also examined binding of a well-known DNA binding

partner of FXR, Retinoid x Receptor alpha (RXRα/NR2B1) *(Evans & Mangelsdorf, 2014;*

Zheng et al., 2018), and bromodomain-containing protein 4 (BRD4), an acetylated histone

reader protein that often binds at active enhancers *(Chen et al., 2016; Li et al., 2016;*

Rahnamoun et al., 2018; Sartorelli & Lauberth, 2020) and is a transcriptional coactivator

of FXR *(Jung et al., 2020)*. GW4064 treatment resulted in substantial increases in

recruitment of both FXR and RXRα to the enhancer region close to the transcription start

site of *FincoR* (arrow shown below in Figure 2B), whereas binding was not detected at a

control region (**Figure 2C**). We also found that BRD4 occupancy was increased at this

enhancer region after GW4064 treatment (**Figure 2C**). These results indicate that

GW4064 activation of FXR leads to increased occupancy of the FXR/RXRα heterodimer

and BRD4 to the enhancer region to upregulate *FincoR* eRNA in the liver.

We identified an inverted repeat1 (IR1) motif that is known to bind FXR *(Calkin &*

Tontonoz, 2012; Lee et al., 2006) within the major FXR binding peak near the start site of

FincoR (arrow shown below in Figure 2B), which we also refer to as FXRE (**Figure 2B,**

C). We examined the functionality of this IR1 motif for mediating transcriptional activation

by GW4064 using reporter assays (**Figure 2D**). We cloned the region containing the IR1

motif into the pGL4.23 luciferase reporter and generated a mutated IR1 motif construct

as a comparison (**Figure 2D**). After transfection into human hepatic HepG2 cells,

GW4064 treatment significantly elevated the luciferase activity of the reporter with the

wild type IR1 motif, but not with the mutated IR1 motif (**Figure 2D**). Together, these results

suggest that GW4064-activated FXR directly upregulates *FincoR* expression.

***FincoR* is a liver-specific nucleus-enriched eRNA**

Because enhancers and eRNAs generally act in a tissue-specific manner *(Li et al., 2016;*

Sartorelli & Lauberth, 2020), we examined the tissue-specific expression of *FincoR* in

mice. Strikingly, *FincoR* is highly expressed in the liver and it is expressed at extremely

low levels in most other tissues, except for a detectable, but still fairly low, level in the

lung (**Figure 3A**). GW4064 treatment resulted in induction of *FincoR* specifically in the

liver (**Figure 3A**). The level of *FincoR* detected in primary mouse hepatocytes (PMHs)

isolated from GW4064-treated mouse liver was similar to that in the liver tissue from the

same mouse, suggesting that the majority of *FincoR* is present in hepatocytes (**Figure**

3B).

By using the 5' and 3' rapid amplification of cDNA ends (RACE), we identified one

transcript of *FincoR* that is of approximately 3.7 kb in length (**Figure 3C**). However, based

on RNA-seq, the length of *FincoR* is over 10 kb (**Figure 1C**), suggesting there are likely

additional multiple RNA isoforms that we were not able to identify by RACE. We next

analyzed the coding potential of *FincoR* utilizing a comparative genomic program,

PhyloCSF. While adjacent genes *Gcnt1* and *Prune2* (**Figure 3D, E**) were correctly

predicted to encode proteins, *FincoR* did not contain a potential protein-coding open

reading frame (**Figure 3E**). Consistent with this bioinformatic prediction, a vector

expressing *FincoR* failed to produce any proteins in an in vitro transcription/translation

assay (**Figure 3F**) confirming that the *FincoR* transcript is a noncoding RNA. *FincoR*

transcripts were enriched by binding to oligo-dT beads, suggesting that *FincoR* is 3'

polyadenylated (**Figure 3G**). Further, *FincoR* was detected in the nuclear compartment

and GW4064 treatment increased the nuclear abundance of *FincoR* (**Figure 3H**),

consistent with its potential transcriptional regulatory function. Together, these results

from molecular biochemical characterization studies reveal that *FincoR* is a liver-enriched

nuclear polyadenylated eRNA.

***FincoR* is induced specifically by the hammerhead-type synthetic FXR agonists**

To determine whether induction of the FXR-induced *FincoR* is ligand-specific, we

examined the effects of several hammerhead-type synthetic FXR agonists, including

GW4064, cilofexor, and tropifexor; a semi-synthetic agonist, OCA; and a non-

hammerhead-type gut-specific agonist, fexaramine (**Figure 4A**) *(Downes et al., 2003;*

Fang et al., 2015). Remarkably, treatment with each of the hammerhead type agonists

for 1 h resulted in a robust induction of *FincoR* (**Figure 4B**), whereas *FincoR* levels were

unchanged after treatment with OCA or fexaramine for 1 h (**Figure 4B**). Treatment with

OCA for 4 h or even one-week treatment with OCA failed to induce hepatic *FincoR* in

mice, while expression of *Shp* was significantly induced (**Figure 4C, D**). Acute feeding

with a diet supplemented with 0.5% cholic acid (CA), a primary BA, for 6 h also failed to

induce *FincoR* expression (**Figure 4E**). Collectively, these results demonstrate that

FincoR is induced specifically by hammerhead-type FXR agonists.

Generation of CRISPR/Cas9-mediated *FincoR* liver-specific knockdown mice

To explore the functional role of hepatic FincoR, we utilized the CRISPR/Cas9 technique

to generate *FincoR* liver-specific knockdown (FincoR-LKD) mice (**Figure 5-figure**

supplement 1A, B). Adenoviral-mediated sgRNA expression in Cas9 mice resulted in

downregulation of *FincoR* specifically in the liver by about 60%, but not in other tissues

(**Figure 5B**). As FXR is a key regulator of bile acid, cholesterol, lipid, and glucose

metabolism and *FincoR* is specifically regulated by FXR, *FincoR* may have a role in the

metabolic process in physiology and disease. We examined liver triglyceride, cholesterol,

bile acid, glycogen and serum non-esterified fatty acids (NEFA) but *FincoR*

downregulation did not result in any significant changes under physiological conditions

(**Figure 5C**).

To explore the molecular signatures and pathways affected by FincoR, we

examined global gene expression by RNA-seq analysis in mouse liver after *FincoR*

knock-down (**Figure 5D**). While *FincoR* was markedly downregulated, the neighboring

genes, such as *Gcnt1, Rfk, Pcsk5 and Prune2*, were largely unchanged (**Figure 5D;**

Figure 5-figure supplement 1C). The RNA-seq analysis revealed 18 up-regulated genes

and 53 downregulated genes in FincoR-LKD liver (**Supplementary Table S4; Figure 5E**).

GO analysis of those downregulated genes indicated that these genes were enriched in

pathways involved in fatty acid oxidation, organelle organization and metabolic process

(**Figure 5F**). Among the down-regulated genes, *Ppp1r3g,* which has a role in controlling

glycogen synthesis, and *Igfbp2,* which functions in insulin resistance, were markedly

reduced (**Supplementary Table S4; Figure 5-figure supplement 1D, E**). Among the up-

regulated genes, expression was substantially increased for *Eda2r*, which is a member

of the tumor necrosis factor receptor superfamily and is involved in inflammation, the

immune response, and development (**Figure 5-figure supplement 1F**). *Fndc1,* which is

involved in fibronectin matrix remodeling, was also suppressed by *FincoR* (and thus

upregulated upon its KO, **Figure 5-figure supplement 1G**). These studies suggest that

FincoR has a role in modulating metabolic homeostasis by regulating genes involved in

metabolism and inflammation.

Amelioration of hepatic steatosis mediated by tropifexor is independent of *FincoR*

in diet-induced NASH mice

Tropifexor, also known as LJN452, is a highly potent hammerhead type FXR agonist that

is currently under clinical trials for NASH and PBC patients *(Kremoser, 2021; Sanyal et*

al., 2023; Tully et al., 2017). Because *FincoR* is induced specifically by the hammerhead

class of FXR agonist (**Figure 4**) and has a potential role in the regulation of metabolism

and inflammation (**Figure 5**), we hypothesized that *FincoR* may play a role in tropifexor-

mediated beneficial effects on reducing NASH pathologies in mice.

We utilized a mouse model that had been fed the Amylin Liver NASH-promoting

(AMLN) diet *(Hernandez et al., 2019; Sun et al., 2022; Zhao et al., 2018)*, and examined

the potential impact of liver-specific downregulation of *FincoR* on tropifexor's effects on

NASH pathology. Cas9 mice fed the AMLN diet for 12 weeks were injected via tail veins

with adenovirus expressing control sgRNA or *FincoR* sgRNA, respectively, and then

treated daily with tropifexor (0.3 mg/kg) for 12 days (**Figure 6A**). In these mice, as a

technical validation of RNA induction and knockdown, *FincoR* levels were significantly

increased by FXR agonist tropifexor, and the increase was blocked by adenovirus

expressing sgRNA for *FincoR* (**Figure 6B**).

We then examined the effect of tropifexor treatment and *FincoR* downregulation

on hepatic steatosis in these mice. Tropifexor treatment markedly reduced neutral lipids

determined by Oil Red O staining of liver sections (**Figure 6C**) and liver TG levels (**Figure**

6D), and these beneficial effects on reducing fatty liver were not altered by *FincoR*

downregulation. Also, *FincoR* downregulation had little effect on liver cholesterol and

gallbladder BA levels, although gallbladder BA levels were reduced by tropifexor (**Figure**

6D).

Consistent with the phenotypes, hepatic expression of key genes involved in BA

synthesis was dramatically reduced by tropifexor treatment (i.e., *Cyp7a1 and Cyp8b1*),

which is consistent with decreased gallbladder BA levels mediated by this agonist (**Figure**

6E). Similarly, tropifexor also lowered lipid synthesis genes (*Srebp1c, Lpin1*, *Scd1*),

consistent with decreased liver TG levels. However, downregulation of *FincoR* did not

result in changes in mRNA levels of these genes (**Figure 6E**). These results indicate that

tropifexor-mediated beneficial effects on reducing hepatic steatosis are independent of

FincoR.

FincoR **facilitates alleviation of liver inflammation by tropifexor in diet-induced**

NASH

Tropifexor ameliorated fibrotic NASH pathologies in preclinical studies *(Hernandez et al.,*

2019; Tully et al., 2017) and has recently concluded phase 2 clinical trials for NASH

patients *(Sanyal et al., 2023)*. We, therefore, further examined the effects of *FincoR*

downregulation on altering other NASH pathologies, including hepatocellular apoptosis,

liver fibrosis and inflammation.

In the same AMLN diet-fed mice as described above (**Figure 6**), analyses of liver

sections revealed that tropifexor treatment reduced hepatocyte swelling/ballooning (H&E

staining), decreased numbers of apoptotic cells (TUNEL staining), alleviated fibrosis

(Sirius red staining), and lowered infiltration of macrophages (F4/80 staining) (**Figure 7A**).

Remarkably, these tropifexor-mediated beneficial effects on NASH pathologies were all

markedly diminished by *FincoR* downregulation (**Figure 7A**). This was consistently found

in various liver lobes and two representative pictures from two different lobes are shown

(**Figure 7A**). In control experiments, *FincoR* downregulation in vehicle-treated mice did

not result in marked changes in NASH pathologies (**Figure 7-figure supplement 1**).

These results demonstrate that *FincoR* is required for tropifexor-mediated beneficial

effects on reducing NASH pathologies, specifically reducing hepatic inflammation, fibrosis

and hepatocyte apoptosis. Consistent with these results, serum ALT and AST levels,

indicators of liver damage, were significantly elevated after *FincoR* downregulation

(**Figure 7B**). Protein levels of key inflammatory markers, IL1β and CCL2, in liver extracts

were also elevated after *FincoR* downregulation (**Figure 7C**).

Consistent with the phenotypes from histological analyses, hepatic expression of

fibrosis and inflammation was altered by *FincoR* knockdown. For example, tropifexor

treatment reduced mRNA levels of several genes that promote fibrosis (*Col1a1, Col1a2,*

Acta2) and hepatic inflammation (*Eda2r, Ifng, Ccl3*), whereas these reductions were

largely reversed by *FincoR* downregulation (**Figure 7D**). We also detected increased

expression of inflammatory genes (*Ccl2, Ccr2, Lcn2*) and an extracellular matrix

remodeling gene *(Fndc1)* in liver with *FincoR* knockdown (**Figure 7D**)*.* Tropifexor

treatment suppresses hepatic apoptosis by reducing pro-apoptotic genes (*Ctsb, Ctss*)

and upregulating anti-apoptotic genes such as *Bcl2 (Warren et al., 2019)*. Importantly,

these effects were significantly reversed by *FincoR* downregulation (**Figure 7D**).

Collectively, these results demonstrate that in diet-induced NASH mice,

pharmacological activation of FXR by tropifexor reduced fibrosis, apoptosis, and

inflammation, which was dependent, at least in part, on the induction of *FincoR*.

***FincoR* expression is increased in chronic liver disease with hepatic inflammation**

and liver injury

To determine whether expression of *FincoR* is altered in chronic liver disease, we utilized

mouse models of NAFLD/NASH and cholestatic liver injury. Hepatic *FincoR* levels were

significantly increased in mice fed with a high fat diet (HFD) for 12 weeks

(**Supplementary Figure S1A**) and in mice fed a HFD with high fructose (HFHF) in

drinking water for 12 weeks (**Supplementary Figure S1B**). Elevated hepatic *FincoR*

levels were also observed in mice treated with α-naphthylisothiocyanate (ANIT), a

chemical inducer of liver cholestasis *(Jung et al., 2020; Kim et al., 2016)* (**Supplementary**

Figure S1C), and in mice with bile duct ligation (BDL), a surgical method to induce

cholestatic liver injury (**Supplementary Figure S1D**).

The sequence of *FincoR* is moderately conserved between mice and humans as

displayed in the UCSC genome browser (**Supplementary Figure S1E**). Annotation in the

NCBI genome data viewer of the human sequence region with similarity to mouse *FincoR*

revealed an functionally uncharacterized human lncRNA, XR_007061585.1, in this region

(Supplementary Figure S1F). However, whether this sequence (or an as yet to be

identified lncRNA) is functional or not has not been determined. To explore the potential

changes or role of lncRNA XR_007061585.1 in human liver pathological conditions, we

measured hepatic levels of the transcripts in PBC and NAFLD patients. Compared to

normal individuals, hepatic lncRNA XR_007061585.1 levels were elevated in patients

with PBC or NAFLD, but not in severe NASH-fibrosis patients (**Supplementary Figure**

S1G, H). These results demonstrate that hepatic levels of a potential human analog of

FincoR are elevated in NAFLD and PBC patients, as *FincoR* is in mouse models of

chronic liver disease with hepatic inflammation and liver injury. However, whether human

lncRNA XR_007061585.1 is analogous to mouse *FincoR* in terms of functions and

mechanisms, and whether elevated XR_007061585.1 levels may have a role in the

disease progression or may be an adaptive response to liver injury remains to be

determined.

Discussion

FXR maintains metabolic homeostasis by transcriptional regulation of genes. Direct

regulation of protein-coding genes by FXR, including *Shp*, is well characterized. In this

study we show that FXR also mediates its functions by induction of lncRNA genes, which

vastly outnumber protein-coding genes. We further show that pharmacological activation

of FXR by hammerhead-type agonists induces a liver-specific enhancer-derived lncRNA,

which we named *FincoR*, that contributes to reduction of NASH pathologies in mice.

FincoR is specifically induced by the hammerhead class of FXR agonists, such as

GW4064, cilofexor, and tropifexor. GW4064 is the mother compound of these isoxazole-

type hammerhead ligands but is not an ideal therapeutic agent because of its poor water

solubility and pharmacokinetics *(Abel et al., 2010)*. In contrast, tropifexor and cilofexor

have better pharmacokinetics and are generally well-tolerated in clinical trials for NASH

and PBC patients *(Abenavoli et al., 2018; Kremoser, 2021; Sanyal et al., 2023; Tully et*

al., 2017), but the underlying mechanisms for their beneficial effects are poorly

understood. Intriguingly, a recent study showed that the gene signature regulated by

tropifexor-activated FXR appears to be broader than that of OCA, partly because the

tropifexor backbone allows a more favorable interaction of FXR with coactivators or

epigenomic modulators *(Hernandez et al., 2019)*. Further, tropifexor was shown to

regulate distinct sets of genes in experimental NASH as compared to other FXR agonists,

particularly genes involved in fibrosis, inflammation, and oxidative stress *(Hernandez et*

al., 2019). Utilizing CRISPR/Cas9-mediated liver-specific knockdown of *FincoR* in diet-

induced NASH mice, we demonstrate that beneficial effects on reducing liver fibrosis,

inflammation, and apoptosis mediated by tropifexor were largely dependent on *FincoR*

(Model, **Figure 7E**).

While this study focused on regulation of *FincoR* by pharmacological activation of

FXR, physiological and pathological regulations of *FincoR* appear to be complex.

Although *FincoR* can be induced by the hammerhead class of FXR agonists, it was not

induced by the endogenous FXR ligand, cholic acid. This implies that *FincoR* may not

contribute to the physiological functions of FXR. In an effort to investigate the potential

role of FincoR in the pathological conditions, we observed that *FincoR* levels were

elevated in mouse models of cholestasis and NASH as well as in human PBC and NAFLD

patients, where BA metabolism is dysregulated. Since different BAs can activate or

repress the gene-regulating function of FXR *(Wahlstrom et al., 2016)*, altered BA

composition in these pathological conditions may contribute to induction of *FincoR*.

Further, binding peaks for multiple nuclear receptors, FXR, LXR, PPARα, RXRα, and

HNF-4α, were detected in the *FincoR* locus so that regulation of *FincoR* likely involves

the combinatorial regulation by multiple nuclear receptors (**Supplementary Figure S2**).

Interestingly, occupancy of the nuclear receptor PPARα at the enhancer region was

increased in fasted mice (**Supplementary Figure S3**). It will be interesting to investigate

whether and how *FincoR* is differently regulated by these nuclear receptors in response

to physiological and pathological cues.

The roles of regulatory RNAs in liver function and diseases and their potentials as

therapeutic targets are increasingly being appreciated *(Brocker et al., 2020; Li et al., 2021;*

Sallam et al., 2016; Zhao et al., 2018). For example, a critical role for an oxysterol nuclear

receptor LXR-induced lncRNA, *LeXis*, in feedback modulation of cholesterol biosynthesis

has been shown *(Sallam et al., 2016)*. Recently, the role of a lncRNA, *Pair*, in liver

phenylalanine metabolism has been demonstrated *(Li et al., 2021)*. Enhancer RNAs are

a less characterized class of lncRNAs and are highly associated with enhancer functions

in gene regulation *(Li et al., 2016)*. Numerous studies have revealed transcriptional roles

for eRNAs in various cellular processes *(Lai & Shiekhattar, 2014; Lam et al., 2014; Li et*

al., 2016; Sartorelli & Lauberth, 2020) but most previous eRNA studies have used

cultured cells *(Hsieh et al., 2014; Li et al., 2013)*, with only a few *in vivo* studies in mouse

models *(Mirtschink et al., 2019; Tang et al., 2023)*. In our current study, through integrative

analysis of transcriptome and histone mark ChIP-Seq, we identified a group of FXR-

regulated eRNAs, including the highly induced *FincoR*. Our current work characterized

the role of *FincoR* in gene regulation and in mediating beneficial pharmacological effects

of tropifexor in NASH, representing important progress in understanding the roles of

eRNAs *in vivo*. Future work is warranted to elucidate the exact mechanisms by which

FincoR facilitates action of FXR agonists to alleviate inflammation, fibrosis and apoptosis.

RNA inside the cells usually associates with different RNA-binding proteins (RBPs)

(Gerstberger et al., 2014). We identified potential binding proteins of *FincoR* using the

ATtRACT database (*Giudice et al., 2016*). The top four candidates for *FincoR* binding are

KHDRBS1, RBM38, YBX2 and YBX3 (**Supplemental Table S5**). KHDRBS1 and RBM38

have been reported to have important roles in RNA processing (*Bielli et al., 2011; Zou et*

al., 2021). YBX2 and YBX3 belong to the Y-box (YBX) protein family, which have been

linked to diverse forms of RNA metabolism and many other processes, including cell

proliferation, DNA repair, stress responses, development, and inflammation (*Kleene,*

2018). Whether these predicted RBPs interact with *FincoR* and how they contribute to

phenotypes should be investigated in future experimentation to understand the

mechanisms involved in *FincoR*-regulated hepatocyte function.

There is limitation of some of our approaches that cannot fully dissect the

underlying mechanisms of *FincoR*. Currently, the function of eRNA loci can be attributed

to a functional eRNA transcript, or binding of transcription factors to this region, or the

transcription process itself, or some combination of these effects *(Li et al., 2016; Sartorelli*

& Lauberth, 2020). In our studies to decrease expression of *FincoR*, the region containing

the FXR binding site was deleted (**Figure 5-figure supplement 1A, right panel**). While

the expression of *FincoR* transcripts was significantly reduced, we cannot rule out

whether decreased binding of FXR or decreased transcription contribute to the observed

changes in phenotype. To directly investigate the function of *FincoR* transcripts,

downregulation of the transcript with antisense oligonucleotides together with

overexpression of *FincoR* in the liver will be required in future experiments.

FXR is increasingly recognized as an important therapeutic target for

enterohepatic diseases, but the development of clinically applicable and more targeted

FXR-based therapy is still challenging. In this study, we provided the first characterization

of an eRNA, *FincoR*, induced by pharmacological activation of FXR and show that *FincoR*

has a beneficial role in reducing liver fibrosis and inflammation in dietary NASH mice.

Complete understanding of the function and mechanisms of *FincoR* may provide novel

insights for the development of desirable therapy for NASH and other chronic liver

diseases.

Materials and Methods

Animal experiments

All animal studies were performed according to procedures approved by the Institutional

Animal Care & Use Committee at the University of Illinois at Urbana-Champaign and were

in accordance with National Institutes of Health guidelines. Mice were maintained in

12/12 h light/dark cycles and fed standard rodent chow. FXR-LKO mice were generated

by breeding FXR floxed mice with Albumin-Cre mice (The Jackson Lab). *FincoR*-LKD

mice were generated as previously reported *(Zhao et al., 2018)*. Briefly, Cas9 transgenic

mice (JAX #024858) were injected via the tail vein with adenoviruses (approximately

5×10^8 PFU) expressing two sgRNAs targeting *FincoR* (sgRNA1:

GGGTTAAGAGCTGTAGGCTG and sgRNA2: ACTTCTATGTCCAACAACCG). The

sequences of sgRNAs were designed using a CRISPR design tool (http://crispr.mit.edu/).

Mice were given a single dose of vehicle or 30 mg/kg GW4064 (in corn oil, Tocris

Bioscience, #2473) after overnight fasting. Mice were treated with 0.5 mg/kg tropifexor (in

corn oil, MedChem Express, HY-107418), 30 mg/kg cilofexor (in corn oil,

MedChemExpress, HY-109083), 100 mg/kg fexaramine (in 0.5% methylcellulose,

MedChem Express, HY-10912), and 20 mg/kg OCA (in 0.5% methylcellulose, MedChem

Express, HY-12222) as indicated. C57BL6 mice were fed with a chow diet containing

0.5% CA for 6 h *(Jung et al., 2020)*.

To induce cholestasis, mice were treated by gavage with 75 mg/kg α-naphthyl

isothiocyanate (ANIT) for 48 hours as previously reported *(Jung et al., 2020; Kim et al.,*

2016; Kim et al., 2020). Cholestasis was also induced by bile duct ligation or sham

operation in mice for 24 hours or 72 hours *(Li et al., 2018)*.

To induce dietary obesity, mice were fed a high-fat diet (TD88137; Harlan Teklad)

or a high-fat diet with 25% fructose in water (high fat/high fructose) for 12 weeks *(Seok et*

al., 2021).

To investigate the effect of liver-specific downregulation of *FincoR* on NASH, male

Cas9 mice were fed the AMLN diet (Research Diets, D09100310, 40 kcal% fat, 2%

cholesterol, 20 kcal% fructose) for 12 weeks and then, were injected with adenovirus

expressing control sgRNA or sgRNA targeting FincoR. Administration of tropifexor was

started 3 days later and given at 0.3 mg/kg dissolved in corn oil.

RNA-Seq

C57BL/6 mice were fasted overnight and *i.p.* injected with vehicle or GW4064 (30 mg/kg)

for 1 h, and livers were collected (n = 4 mice for either vehicle or treated group). Total

RNA from each liver was extracted by RNeasy kit (Qiagen), and two randomly selected

mice liver RNAs were pooled for RNA-seq (two RNA-seq reactions from 4 mice livers for

either vehicle or treated group). RNA-seq was performed as previously described *(Byun*

et al., 2018; Byun et al., 2020; Seok et al., 2018). Ribosomal RNA was removed with the

Ribozero HMR Gold kit (Illumina). The sequencing library was generated following

methods described below.

507   Construction of strand-specific RNAseq libraries

Construction of the RNAseq libraries and sequencing on the Illumina NovaSeq 6000 were

performed at the Roy J. Carver Biotechnology Center at the University of Illinois at

Urbana-Champaign. After DNase digestion, purified total RNAs were analyzed on a

Fragment Analyzer (Agilent) to evaluate RNA integrity. The total RNAs were converted

into individually barcoded polyadenylated mRNAseq libraries with the Kapa HyperPrep

mRNA kit (Roche). Libraries were barcoded with Unique Dual Indexes (UDI's) which have

been developed to prevent index switching. The adaptor-ligated double-stranded cDNAs

were amplified by PCR for 8 cycles with the Kapa HiFi polymerase (Roche). The final

libraries were quantitated with Qubit (Thermo Fisher) and the average cDNA fragment

sizes were determined on a Fragment Analyzer. The libraries were diluted to 10 nM and

further quantitated by qPCR on a CFX Connect Real-Time qPCR system (Bio-Rad) for

accurate pooling of barcoded libraries and maximization of number of clusters in the

flowcell.

522   Sequencing of libraries in the NovaSeq

The barcoded RNAseq libraries were loaded on one SP lane on a NovaSeq 6000 for

cluster formation and sequencing. The libraries were sequenced from one end of the

fragments for a total of 100 bp. The fastq read files were generated and demultiplexed

with the bcl2fastq v2.20 Conversion Software (Illumina, San Diego, CA). The quality of

the demultiplexed fastq files was evaluated with the FastQC software, which generates

reports with the quality scores, base composition, k-mer, GC and N contents, sequence

duplication levels and overrepresented sequences.

GRO-seq

To harvest the nuclei from mouse liver cells, the liver was harvested at indicated time and

washed with a cold swelling buffer (10 mM Tris-HCl, pH 7.5, 2 mM $MgCl_2$, 3 mM $CaCl_2$,

2 U/ml Superase-In). The nuclei were prepared by Dounce homogenization in cold

swelling buffer and filtered using a cell strainer (100 μm, BD Biosciences). Nuclei were

collected by centrifugation at 400 x g for 10 min, then resuspended in the lysis buffer

(swelling buffer with 10% glycerol and 1% IGEPAL) and incubated on ice for 5 min. Nuclei

were washed twice with the lysis buffer and resuspended at a concentration of 10^8

nuclei/ml in the freezing buffer (50 mM Tris-HCl, pH 8.3, 40% glycerol, 5 mM $MgCl_2$, 0.1

mM EDTA). We then followed our previous method *(Li et al., 2013; Oh et al., 2021)* to

conduct nuclear run-on and GRO-Seq library preparation. Briefly, the nuclei in freezing

buffer were subjected to the nuclear run-on reaction by mixing with an equal volume of

run-on buffer (10 mM Tris-HCl, pH 8.0, 5 mM $MgCl_2$, 1 mM dithiothreitol (DTT), 300 mM

KCl, 20 units of Superase-In, 1% sarkosyl, 500 μM ATP, GTP, Br-UTP and 2 μM CTP) for

5 min at 30 °C. The BrU-labeled run-on RNAs were extracted by TRIzol and purified by

anti-BrdU agarose beads (Santa Cruz Biotech, sc-32323 AC). The run-on RNAs were then subjected to end repair by T4 PNK, poly-adenylation, and then to cDNA first strand synthesis by a custom primer (oNTI223) that allows circularization of the cDNA. This cDNA then was re-linearized by Ape1 (NEB), size selected by TBE gel electrophoresis and the products of the desired size were excised (~320–350 bp) for final library prep and sequencing. GRO-Seq samples were run on a NextSeq 500 sequencer from Illumina with a single-end 80 nt model.

Histological analyses

For histology, tissues were dissected and immediately fixed in 10% formalin overnight and processed for paraffin embedding and H&E staining. Paraffin-embedded liver sections were incubated with F4/80 antibody, and antibody was detected using a peroxidase-based method (Abcam, ab64238). Liver collagen was detected by Sirius Red staining (Abcam, ab246832) and apoptosis was detected by TUNEL staining (Millipore, S7100). For Oil Red O staining, liver tissue was frozen in OCT compound (Sakura Finetek, 4583), sectioned, and stained. Liver sections were imaged with a NanoZoomer Scanner (Hamamatsu) and quantification was done using NIH ImageJ.

Metabolic analyses

Hepatic levels of TG (Sigma, MAK266), cholesterol (Sigma, MAK043), glycogen (Biovision, K646-100), total BA levels (Diazyme, DZ042A), serum NEFA (Sigma, MAK044), serum ALT (Sigma, MAK052) and serum AST (Sigma, MAK055) were determined according to the manufacturer's instructions. Mouse liver IL-1β (R&D

systems, MLB00C) and CCL2 (R&D systems, DY479-05) were detected by commercially

available ELISA kit.

Liver samples of PBC and NAFLD patients. Liver specimens from normal organ

donors, and patients with PBC or NAFLD were obtained from the Liver Tissue

Procurement and Distribution System. The samples were unidentifiable, and thus, ethical

approval was not required. Hepatic XR_007061585.1 levels were measured by qPCR.

Mouse liver ChIP

Liver ChIP assay was performed as described previously *(Jung et al., 2020; Seok et al.,*

2018). Briefly, chromatin extracts were prepared from FXR floxed and FXR-LKO mouse

livers after treatment of the mice with GW4064 which was followed by preclearing and

immunoprecipitation using control IgG or FXR antibody (Novus Biologicals, NBP2-16550;

Santa Cruz, sc-25309), RXRα antibody (Proteintech, catalog no. 21218-1-AP), or BRD4

antibody (Bethyl Laboratories, catalog #A301-985A50). Enrichment in chromatin

precipitates of gene sequences was measured by qRCR using primers listed

(**Supplementary Table S6**).

Primary mouse hepatocytes and HepG2 cells

Primary hepatocytes were isolated from C57BL/6 mice by collagenase (Worthington

Biochemical Corporation, LS004188) perfusion and maintained in William's E Medium

with primary hepatocyte maintenance supplements (Gibco #CM4000) in six-well plates

as described previously *(Jung et al., 2020; Kim et al., 2015)*. HepG2 cells were maintained

in DMEM supplemented with 10% fetal bovine serum at 37°C and 5% CO_2.

Luciferase reporter assay

The enhancer region containing the FXR binding element was amplified by PCR and

cloned into the pGL4.23 vector (Promega). HepG2 cells were transiently transfected with

FXR (100 ng/well) and RXRα (5 ng/well) in combination with reporters containing the wild

type FXRE or mutated FXRE (200 ng/well) (**Supplementary Table S6**). β-galactosidase

plasmid (200 ng/well) was also transfected as an internal control. Cells were treated with

GW4064 or DMSO after transient transfection. Six hours later, the cells were harvested.

All reporter assays were repeated at least three times in triplicates.

RACE

5' and 3' RACE assays were performed using a SMARTer RACE kit (Clontech) according

to the manufacturer's instructions. The resulting PCR products were separated by

electrophoresis in agarose gels and cloned into the pRACE vector provided by the kit.

The transcription start sites and end sites of *FincoR* were determined by sequencing. The

gene-specific primers used for 5'- and 3'-RACE are listed in **Supplementary Table S6**.

In vitro transcription and translation (TNT)

Expression plasmids for luciferase and *FincoR* were mixed with a Coupled Reticulocyte

Lysate System (Promega). After incubating at 30°C for 60 min, translated products were

separated on 4-20% gradient SDS polyacrylamide gels and transferred to PVDF

membranes. Chemiluminescent detection of in vitro translated protein was performed

following the manufacturer's protocol (Promega).

RT-qPCR

Total RNA was extracted using the RNeasy Mini Kit (Qiagen, 74104) and 2 μg of RNA

was reverse transcribed and mRNA expression was normalized relative to that of 36B4.

The qPCR primers are shown in **Supplementary Table S6**.

Subcellular fractionation

Using a Cytoplasmic and Nuclear RNA Purification Kit (Norgen, Thorold, ON, Canada),

the cytoplasm and nucleus fractions from primary hepatocytes isolated from livers of mice

treated with GW4064 or DMSO were separated, and both fractions were subjected to

RNA extraction and qRT-PCR.

Immunoblotting analysis

Total liver lysates were prepared as described before *(Sun et al., 2022)*. Antibodies for

immunoblotting for β-ACTIN (#4970) were purchased from Cell Signaling Technology.

Antibodies for immunoblotting for FXR (sc-25309) were purchased from Santa Cruz.

632
FincoR **polyadenylation study**

Total RNA was prepared using the RNeasy mini prep kit (Qiagen). Poly(A)+ and poly(A)-

RNA was separated using a Dynabeads™ mRNA Purification Kit (ThermoFisher

Scientific) according to the manufacturer's instructions. Briefly, total RNA was incubated

with the Dynabeads/binding buffer suspension at room temperature for 5 min and the

reaction tubes were placed on a magnet until solution was clear. The supernatant

containing poly(A)- RNA was saved. The beads with poly(A)+ RNA were washed three

times with washing buffer provided by the kit. RNA was extracted from the supernatant

and beads respectively using Trizol. qPCR was performed to analyze levels of *FincoR*

and 36b4 in the poly(A)+ and poly(A)- RNA fractions.

Genomics Analysis

RNA-Seq and GRO-Seq reads were mapped to the mouse reference genome mm10 with

STAR aligner *(Dobin et al., 2013)*. Transcript quantifications were done with the HOMER

tool set *(Heinz et al., 2010)* and enhancer RNAs were identified based on H3K27ac ChIP-

Seq in mouse liver. Briefly, intergenic H3K27ac ChIP-Seq peaks were selected as

putative enhancer regions in mouse liver.   Then +/- 3kb regions around putative

enhancers with RNA-Seq signal (>1 RPKM) were considered as putative eRNAs in

mouse livers. Overlapped eRNAs regions were merged, and redundant ones were

removed. In addition, any +/- 3kb extended eRNAs regions that overlapped with protein-

coding genes were further removed to avoid transcriptional readthrough from genes. We

used DESeq2 *(Love et al., 2014)* to identify significantly regulated eRNAs with a cutoff of

(FDR < 0.05, Log2FC > 1). Public ChIP-Seq datasets were obtained from ENCODE or

GEO (**Supplementary Table S3**).

Statistical analysis

Statistical analysis was performed using GraphPad Prism 9. Statistical differences were

evaluated using the two-tailed unpaired Student's t-test for comparisons between two

groups, or ANOVA and appropriate post hoc analyses for comparisons of more than two

groups. Statistical methods and corresponding *p* values for data shown in each panel are

indicated in figure legends.

Data availability

RNA-seq and Gro-seq data were deposited in GEO under the following accession

numbers GSE 221986.

Acknowledgments

We thank Deepa Prakashini Govindasamy Rajagopal for the help with tissue genotyping.

We thank Tiangang Li at University of Oklahoma for providing bile-duct ligated mouse

liver samples, and Kristina Schoonjans at Ecole Polytech, Switzerland, for providing FXR

floxed mice. We also thank the Liver Tissue Cell Distribution System, University of

Minnesota (NIH Contract # HHSN276201200017C), for providing liver specimens of

NAFLD and PBC patients and individuals without liver disease. This study was supported

by an American Diabetes Association Postdoctoral Fellowship to J.C. (1-19-PDF-117),

and by a John and Rebekah Harper Fellowship to R.W. W.L. is a Cancer Prevention and

Research Institute of Texas (CPRIT) Scholar (RR160083). This work is supported by

funding from NIH (K22CA204468 and R01GM136922), and Welch Foundation (AU-2000-

20220331) to W.L., and grants from the NIH (R01 DK062777 and R01 DK095842) to

J.K.K.

References

Abel, U., Schluter, T., Schulz, A., Hambruch, E., Steeneck, C., Hornberger, M., Hoffmann, T., Perovic-Ottstadt, S., Kinzel, O., Burnet, M., Deuschle, U., & Kremoser, C. (2010). Synthesis and pharmacological validation of a novel series of non-steroidal FXR agonists. *Bioorg Med Chem Lett*, *20*(16), 4911-4917. https://doi.org/10.1016/j.bmcl.2010.06.084

Abenavoli, L., Falalyeyeva, T., Boccuto, L., Tsyryuk, O., & Kobyliak, N. (2018). Obeticholic Acid: A New Era in the Treatment of Nonalcoholic Fatty Liver Disease. *Pharmaceuticals (Basel)*, *11*(4). https://doi.org/10.3390/ph11040104

Ali, A. H., Carey, E. J., & Lindor, K. D. (2015). Recent advances in the development of farnesoid X receptor agonists. *Ann Transl Med*, *3*(1), 5. https://doi.org/10.3978/j.issn.2305-5839.2014.12.06

Bielli, P., Busa, R., Paronetto, M. P., & Sette, C. (2011). The RNA-binding protein Sam68 is a multifunctional player in human cancer. *Endocr Relat Cancer*, *18*(4), R91-R102. https://doi.org/10.1530/ERC-11-0041

Brocker, C. N., Kim, D., Melia, T., Karri, K., Velenosi, T. J., Takahashi, S., Aibara, D., Bonzo, J. A., Levi, M., Waxman, D. J., & Gonzalez, F. J. (2020). Long non-coding RNA Gm15441 attenuates hepatic inflammasome activation in response to PPARA agonism and fasting. *Nat Commun*, *11*(1), 5847. https://doi.org/10.1038/s41467-020-19554-7

Byun, S., Kim, D. H., Ryerson, D., Kim, Y. C., Sun, H., Kong, B., Yau, P., Guo, G., Xu, H. E., Kemper, B., & Kemper, J. K. (2018). Postprandial FGF19-induced phosphorylation by Src is critical for FXR function in bile acid homeostasis. *Nat Commun*, *9*(1), 2590. https://doi.org/10.1038/s41467-018-04697-5

Byun, S., Seok, S., Kim, Y. C., Zhang, Y., Yau, P., Iwamori, N., Xu, H. E., Ma, J., Kemper, B., & Kemper, J. K. (2020). Fasting-induced FGF21 signaling activates hepatic autophagy and lipid degradation via JMJD3 histone demethylase. *Nat Commun*, *11*(1), 807. https://doi.org/10.1038/s41467-020-14384-z

Calkin, A. C., & Tontonoz, P. (2012). Transcriptional integration of metabolism by the nuclear sterol-activated receptors LXR and FXR. *Nat Rev Mol Cell Biol*, *13*(4), 213-224. https://doi.org/10.1038/nrm3312

Cech, T. R., & Steitz, J. A. (2014). The noncoding RNA revolution-trashing old rules to forge new ones. *Cell*, *157*(1), 77-94. https://doi.org/10.1016/j.cell.2014.03.008

Chen, J., Wang, Z., Hu, X., Chen, R., Romero-Gallo, J., Peek, R. M., Jr., & Chen, L. F. (2016). BET Inhibition Attenuates Helicobacter pylori-Induced Inflammatory Response by Suppressing Inflammatory Gene Transcription and Enhancer Activation. *J Immunol*, *196*(10), 4132-4142. https://doi.org/10.4049/jimmunol.1502261

Claudel, T., Staels, B., & Kuipers, F. (2005). The Farnesoid X receptor: a molecular link between bile acid and lipid and glucose metabolism. *Arterioscler Thromb Vasc Biol*, *25*(10), 2020-2030. https://doi.org/10.1161/01.ATV.0000178994.21828.a7

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15-21. https://doi.org/10.1093/bioinformatics/bts635

Downes, M., Verdecia, M. A., Roecker, A. J., Hughes, R., Hogenesch, J. B., Kast-Woelbern, H. R., Bowman, M. E., Ferrer, J. L., Anisfeld, A. M., Edwards, P. A., Rosenfeld, J. M., Alvarez, J. G., Noel, J. P., Nicolaou, K. C., & Evans, R. M. (2003). A chemical, genetic, and structural analysis of the nuclear bile acid receptor FXR. *Mol Cell*, *11*(4), 1079-1092. https://doi.org/10.1016/s1097-2765(03)00104-7

Evans, R. M., & Mangelsdorf, D. J. (2014). Nuclear Receptors, RXR, and the Big Bang. *Cell*, *157*(1), 255-266. https://doi.org/10.1016/j.cell.2014.03.012

Fang, B., Everett, L. J., Jager, J., Briggs, E., Armour, S. M., Feng, D., Roy, A., Gerhart-Hines, Z., Sun, Z., & Lazar, M. A. (2014). Circadian enhancers coordinate multiple phases of rhythmic gene transcription in vivo. *Cell*, *159*(5), 1140-1152. https://doi.org/10.1016/j.cell.2014.10.022

Fang, S., Suh, J. M., Reilly, S. M., Yu, E., Osborn, O., Lackey, D., Yoshihara, E., Perino, A., Jacinto, S., Lukasheva, Y., Atkins, A. R., Khvat, A., Schnabl, B., Yu, R. T., Brenner, D. A., Coulter, S., Liddle, C., Schoonjans, K., Olefsky, J. M., . . . Evans, R. M. (2015). Intestinal FXR agonism promotes adipose tissue browning and reduces obesity and insulin resistance. *Nat Med*, *21*(2), 159-165. https://doi.org/10.1038/nm.3760

Friedman, S. L., Neuschwander-Tetri, B. A., Rinella, M., & Sanyal, A. J. (2018). Mechanisms of NAFLD development and therapeutic strategies. *Nat Med*, *24*(7), 908-922. https://doi.org/10.1038/s41591-018-0104-9

Gerstberger, S., Hafner, M., & Tuschl, T. (2014). A census of human RNA-binding proteins. *Nat Rev Genet*, *15*(12), 829-845. https://doi.org/10.1038/nrg3813

Giudice, G., Sanchez-Cabo, F., Torroja, C., & Lara-Pezzi, E. (2016). ATtRACT-a database of RNA-binding proteins and associated motifs. *Database (Oxford)*, *2016*. https://doi.org/10.1093/database/baw035

Goodwin, B., Jones, S. A., Price, R. R., Watson, M. A., McKee, D. D., Moore, L. B., Galardi, C., Wilson, J. G., Lewis, M. C., Roth, M. E., Maloney, P. R., Willson, T. M., & Kliewer, S. A. (2000). A regulatory cascade of the nuclear receptors FXR, SHP-1, and LRH-1 represses bile acid biosynthesis. *Mol Cell*, *6*(3), 517-526. https://doi.org/10.1016/s1097-2765(00)00051-4

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H., & Glass, C. K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell*, *38*(4), 576-589. https://doi.org/10.1016/j.molcel.2010.05.004

Hernandez, E. D., Zheng, L., Kim, Y., Fang, B., Liu, B., Valdez, R. A., Dietrich, W. F., Rucker, P. V., Chianelli, D., Schmeits, J., Bao, D., Zoll, J., Dubois, C., Federe, G. C., Chen, L., Joseph, S. B., Klickstein, L. B., Walker, J., Molteni, V., . . . Laffitte, B. (2019). Tropifexor-Mediated Abrogation of Steatohepatitis and Fibrosis Is Associated With the Antioxidative Gene Expression Profile in Rodents. *Hepatol Commun*, *3*(8), 1085-1097. https://doi.org/10.1002/hep4.1368

Hirabayashi, S., Bhagat, S., Matsuki, Y., Takegami, Y., Uehata, T., Kanemaru, A., Itoh, M., Shirakawa, K., Takaori-Kondo, A., Takeuchi, O., Carninci, P., Katayama, S., Hayashizaki, Y., Kere, J., Kawaji, H., & Murakawa, Y. (2019). NET-CAGE characterizes the dynamics and topology of human transcribed cis-regulatory

elements. *Nat Genet*, *51*(9), 1369-1379. https://doi.org/10.1038/s41588-019-
0485-9
Hon, C. C., Ramilowski, J. A., Harshbarger, J., Bertin, N., Rackham, O. J., Gough, J.,
Denisenko, E., Schmeier, S., Poulsen, T. M., Severin, J., Lizio, M., Kawaji, H.,
Kasukawa, T., Itoh, M., Burroughs, A. M., Noma, S., Djebali, S., Alam, T.,
Medvedeva, Y. A., . . . Forrest, A. R. (2017). An atlas of human long non-coding
RNAs with accurate 5' ends. *Nature*, *543*(7644), 199-204.
https://doi.org/10.1038/nature21374
Hsieh, C. L., Fei, T., Chen, Y., Li, T., Gao, Y., Wang, X., Sun, T., Sweeney, C. J., Lee,
G. S., Chen, S., Balk, S. P., Liu, X. S., Brown, M., & Kantoff, P. W. (2014).
Enhancer RNAs participate in androgen receptor-driven looping that selectively
enhances gene activation. *Proc Natl Acad Sci U S A*, *111*(20), 7319-7324.
https://doi.org/10.1073/pnas.1324151111
Jung, H., Chen, J., Hu, X., Sun, H., Wu, S. Y., Chiang, C. M., Kemper, B., Chen, L. F., &
Kemper, J. K. (2020). BRD4 inhibition and FXR activation, individually beneficial
in cholestasis, are antagonistic in combination. *JCI Insight*, *6*(1).
https://doi.org/10.1172/jci.insight.141640
Kim, D. H., Kwon, S., Byun, S., Xiao, Z., Park, S., Wu, S. Y., Chiang, C. M., Kemper, B.,
& Kemper, J. K. (2016). Critical role of RanBP2-mediated SUMOylation of Small
Heterodimer Partner in maintaining bile acid homeostasis. *Nat Commun*, *7*,
12179. https://doi.org/10.1038/ncomms12179
Kim, Y. C., Fang, S., Byun, S., Seok, S., Kemper, B., & Kemper, J. K. (2015). Farnesoid
X receptor-induced lysine-specific histone demethylase reduces hepatic bile acid
levels and protects the liver against bile acid toxicity. *Hepatology*, *62*(1), 220-231.
https://doi.org/10.1002/hep.27677
Kim, Y. C., Jung, H., Seok, S., Zhang, Y., Ma, J., Li, T., Kemper, B., & Kemper, J. K.
(2020). MicroRNA-210 Promotes Bile Acid-Induced Cholestatic Liver Injury by
Targeting Mixed-Lineage Leukemia-4 Methyltransferase in Mice. *Hepatology*,
71(6), 2118-2134. https://doi.org/10.1002/hep.30966
Kleene, K. C. (2018). Y-box proteins combine versatile cold shock domains and
arginine-rich motifs (ARMs) for pleiotropic functions in RNA biology. *Biochem J*,
475(17), 2769-2784. https://doi.org/10.1042/BCJ20170956
Kliewer, S. A., & Mangelsdorf, D. J. (2015). Bile Acids as Hormones: The FXR-
FGF15/19 Pathway. *Dig Dis*, *33*(3), 327-331. https://doi.org/10.1159/000371670
Kremoser, C. (2021). FXR agonists for NASH: How are they different and what
difference do they make? *J Hepatol*, *75*(1), 12-15.
https://doi.org/10.1016/j.jhep.2021.03.020
Lai, F., & Shiekhattar, R. (2014). Enhancer RNAs: the new molecules of transcription.
Curr Opin Genet Dev, *25*, 38-42. https://doi.org/10.1016/j.gde.2013.11.017
Lam, M. T., Li, W., Rosenfeld, M. G., & Glass, C. K. (2014). Enhancer RNAs and
regulated transcriptional programs. *Trends Biochem Sci*, *39*(4), 170-182.
https://doi.org/10.1016/j.tibs.2014.02.007
Lee, F. Y., Lee, H., Hubbert, M. L., Edwards, P. A., & Zhang, Y. (2006). FXR, a
multipurpose nuclear receptor. *Trends Biochem Sci*, *31*(10), 572-580.
https://doi.org/10.1016/j.tibs.2006.08.002
Lee, J., Padhye, A., Sharma, A., Song, G., Miao, J., Mo, Y. Y., Wang, L., & Kemper, J.

K. (2010). A pathway involving farnesoid X receptor and small heterodimer partner positively regulates hepatic sirtuin 1 levels via microRNA-34a inhibition. *J Biol Chem*, *285*(17), 12604-12611. https://doi.org/10.1074/jbc.M109.094524

Lee, J., Seok, S., Yu, P., Kim, K., Smith, Z., Rivas-Astroza, M., Zhong, S., & Kemper, J. K. (2012). Genomic analysis of hepatic farnesoid X receptor binding sites reveals altered binding in obesity and direct gene repression by farnesoid X receptor in mice. *Hepatology*, *56*(1), 108-117. https://doi.org/10.1002/hep.25609

Lee, J. M., Wagner, M., Xiao, R., Kim, K. H., Feng, D., Lazar, M. A., & Moore, D. D. (2014). Nutrient-sensing nuclear receptors coordinate autophagy. *Nature*, *516*(7529), 112-115. https://doi.org/10.1038/nature13961

Li, J., Woolbright, B. L., Zhao, W., Wang, Y., Matye, D., Hagenbuch, B., Jaeschke, H., & Li, T. (2018). Sortilin 1 Loss-of-Function Protects Against Cholestatic Liver Injury by Attenuating Hepatic Bile Acid Accumulation in Bile Duct Ligated Mice. *Toxicol Sci*, *161*(1), 34-47. https://doi.org/10.1093/toxsci/kfx078

Li, W., Notani, D., Ma, Q., Tanasa, B., Nunez, E., Chen, A. Y., Merkurjev, D., Zhang, J., Ohgi, K., Song, X., Oh, S., Kim, H. S., Glass, C. K., & Rosenfeld, M. G. (2013). Functional roles of enhancer RNAs for oestrogen-dependent transcriptional activation. *Nature*, *498*(7455), 516-520. https://doi.org/10.1038/nature12210

Li, W., Notani, D., & Rosenfeld, M. G. (2016). Enhancers as non-coding RNA transcription units: recent insights and future perspectives. *Nat Rev Genet*, *17*(4), 207-223. https://doi.org/10.1038/nrg.2016.4

Li, Y., Tan, Z., Zhang, Y., Zhang, Z., Hu, Q., Liang, K., Jun, Y., Ye, Y., Li, Y. C., Li, C., Liao, L., Xu, J., Xing, Z., Pan, Y., Chatterjee, S. S., Nguyen, T. K., Hsiao, H., Egranov, S. D., Putluri, N., . . . Yang, L. (2021). A noncoding RNA modulator potentiates phenylalanine metabolism in mice. *Science*, *373*(6555), 662-673. https://doi.org/10.1126/science.aba4991

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Mattick, J. S., Amaral, P. P., Carninci, P., Carpenter, S., Chang, H. Y., Chen, L. L., Chen, R., Dean, C., Dinger, M. E., Fitzgerald, K. A., Gingeras, T. R., Guttman, M., Hirose, T., Huarte, M., Johnson, R., Kanduri, C., Kapranov, P., Lawrence, J. B., Lee, J. T., . . . Wu, M. (2023). Long non-coding RNAs: definitions, functions, challenges and recommendations. *Nat Rev Mol Cell Biol*, *24*(6), 430-447. https://doi.org/10.1038/s41580-022-00566-8

Mirtschink, P., Bischof, C., Pham, M. D., Sharma, R., Khadayate, S., Rossi, G., Fankhauser, N., Traub, S., Sossalla, S., Hagag, E., Berthonneche, C., Sarre, A., Stehr, S. N., Grote, P., Pedrazzini, T., Dimmeler, S., Krek, W., & Krishnan, J. (2019). Inhibition of the Hypoxia-Inducible Factor 1alpha-Induced Cardiospecific HERNA1 Enhance-Templated RNA Protects From Heart Disease. *Circulation*, *139*(24), 2778-2792. https://doi.org/10.1161/CIRCULATIONAHA.118.036769

Oh, S., Shao, J., Mitra, J., Xiong, F., D'Antonio, M., Wang, R., Garcia-Bassets, I., Ma, Q., Zhu, X., Lee, J. H., Nair, S. J., Yang, F., Ohgi, K., Frazer, K. A., Zhang, Z. D., Li, W., & Rosenfeld, M. G. (2021). Enhancer release and retargeting activates disease-susceptibility genes. *Nature*, *595*(7869), 735-740. https://doi.org/10.1038/s41586-021-03577-1

Rahnamoun, H., Lee, J., Sun, Z., Lu, H., Ramsey, K. M., Komives, E. A., & Lauberth, S. M. (2018). RNAs interact with BRD4 to promote enhanced chromatin engagement and transcription activation. *Nat Struct Mol Biol*, *25*(8), 687-697. https://doi.org/10.1038/s41594-018-0102-0

Sallam, T., Jones, M. C., Gilliland, T., Zhang, L., Wu, X., Eskin, A., Sandhu, J., Casero, D., Vallim, T. Q., Hong, C., Katz, M., Lee, R., Whitelegge, J., & Tontonoz, P. (2016). Feedback modulation of cholesterol metabolism by the lipid-responsive non-coding RNA LeXis. *Nature*, *534*(7605), 124-128. https://doi.org/10.1038/nature17674

Sanyal, A. J., Lopez, P., Lawitz, E. J., Lucas, K. J., Loeffler, J., Kim, W., Goh, G. B. B., Huang, J. F., Serra, C., Andreone, P., Chen, Y. C., Hsia, S. H., Ratziu, V., Aizenberg, D., Tobita, H., Sheikh, A. M., Vierling, J. M., Kim, Y. J., Hyogo, H., . . . Brass, C. A. (2023). Tropifexor for nonalcoholic steatohepatitis: an adaptive, randomized, placebo-controlled phase 2a/b trial. *Nat Med*, *29*(2), 392-400. https://doi.org/10.1038/s41591-022-02200-8

Sartorelli, V., & Lauberth, S. M. (2020). Enhancer RNAs are an important regulatory layer of the epigenome. *Nat Struct Mol Biol*, *27*(6), 521-528. https://doi.org/10.1038/s41594-020-0446-0

Seok, S., Fu, T., Choi, S. E., Li, Y., Zhu, R., Kumar, S., Sun, X., Yoon, G., Kang, Y., Zhong, W., Ma, J., Kemper, B., & Kemper, J. K. (2014). Transcriptional regulation of autophagy by an FXR-CREB axis. *Nature*, *516*(7529), 108-111. https://doi.org/10.1038/nature13949

Seok, S., Kim, Y. C., Byun, S., Choi, S., Xiao, Z., Iwamori, N., Zhang, Y., Wang, C., Ma, J., Ge, K., Kemper, B., & Kemper, J. K. (2018). Fasting-induced JMJD3 histone demethylase epigenetically activates mitochondrial fatty acid beta-oxidation. *J Clin Invest*, *128*(7), 3144-3159. https://doi.org/10.1172/JCI97736

Seok, S., Sun, H., Kim, Y. C., Kemper, B., & Kemper, J. K. (2021). Defective FXR-SHP Regulation in Obesity Aberrantly Increases miR-802 Expression, Promoting Insulin Resistance and Fatty Liver. *Diabetes*, *70*(3), 733-744. https://doi.org/10.2337/db20-0856

Statello, L., Guo, C. J., Chen, L. L., & Huarte, M. (2021). Gene regulation by long non-coding RNAs and its biological functions. *Nat Rev Mol Cell Biol*, *22*(2), 96-118. https://doi.org/10.1038/s41580-020-00315-9

Sun, H., Seok, S., Jung, H., Kemper, B., & Kemper, J. K. (2022). Obesity-induced miR-802 directly targets AMPK and promotes nonalcoholic steatohepatitis in mice. *Mol Metab*, *66*, 101603. https://doi.org/10.1016/j.molmet.2022.101603

Tang, X., Luo, Y., Yuan, D., Calandrelli, R., Malhi, N. K., Sriram, K., Miao, Y., Lou, C. H., Tsark, W., Tapia, A., Chen, A. T., Zhang, G., Roeth, D., Kalkum, M., Wang, Z. V., Chien, S., Natarajan, R., Cooke, J. P., Zhong, S., & Chen, Z. B. (2023). Long noncoding RNA LEENE promotes angiogenesis and ischemic recovery in diabetes models. *J Clin Invest*, *133*(3). https://doi.org/10.1172/JCI161759

Thomas, A. M., Hart, S. N., Kong, B., Fang, J., Zhong, X. B., & Guo, G. L. (2010). Genome-wide tissue-specific farnesoid X receptor binding in mouse liver and intestine. *Hepatology*, *51*(4), 1410-1419. https://doi.org/10.1002/hep.23450

Tully, D. C., Rucker, P. V., Chianelli, D., Williams, J., Vidal, A., Alper, P. B., Mutnick, D., Bursulaya, B., Schmeits, J., Wu, X., Bao, D., Zoll, J., Kim, Y., Groessl, T.,

McNamara, P., Seidel, H. M., Molteni, V., Liu, B., Phimister, A., . . . Laffitte, B. (2017). Discovery of Tropifexor (LJN452), a Highly Potent Non-bile Acid FXR Agonist for the Treatment of Cholestatic Liver Diseases and Nonalcoholic Steatohepatitis (NASH). *J Med Chem*, *60*(24), 9960-9973. https://doi.org/10.1021/acs.jmedchem.7b00907

Wahlstrom, A., Sayin, S. I., Marschall, H. U., & Backhed, F. (2016). Intestinal Crosstalk between Bile Acids and Microbiota and Its Impact on Host Metabolism. *Cell Metab*, *24*(1), 41-50. https://doi.org/10.1016/j.cmet.2016.05.005

Wang, Y. D., Chen, W. D., Wang, M., Yu, D., Forman, B. M., & Huang, W. (2008). Farnesoid X receptor antagonizes nuclear factor kappaB in hepatic inflammatory response. *Hepatology*, *48*(5), 1632-1643. https://doi.org/10.1002/hep.22519

Warren, C. F. A., Wong-Brown, M. W., & Bowden, N. A. (2019). BCL-2 family isoforms in apoptosis and cancer. *Cell Death Dis*, *10*(3), 177. https://doi.org/10.1038/s41419-019-1407-6

Zhao, X. Y., Xiong, X., Liu, T., Mi, L., Peng, X., Rui, C., Guo, L., Li, S., Li, X., & Lin, J. D. (2018). Long noncoding RNA licensing of obesity-linked hepatic lipogenesis and NAFLD pathogenesis. *Nat Commun*, *9*(1), 2986. https://doi.org/10.1038/s41467-018-05383-2

Zheng, W., Lu, Y., Tian, S., Ma, F., Wei, Y., Xu, S., & Li, Y. (2018). Structural insights into the heterodimeric complex of the nuclear receptors FXR and RXR. *J Biol Chem*, *293*(32), 12535-12541. https://doi.org/10.1074/jbc.RA118.004188

Zou, C., Wan, Y., He, L., Zheng, J. H., Mei, Y., Shi, J., Zhang, M., Dong, Z., & Zhang, D. (2021). RBM38 in cancer: role and mechanism. *Cell Mol Life Sci*, *78*(1), 117-128. https://doi.org/10.1007/s00018-020-03593-w

Figure legends

Figure 1. Activation of FXR by GW4064 induces *FincoR*, a novel eRNA, in mouse

liver. (A) Volcano plot from RNA-Seq showing significantly induced eRNAs (*FincoR* is

highlighed) in the livers of C57BL/6 male mice treated with GW4064 (*i.p*. injection, 30

mg/kg, 1 h) or vehicle. The *x* axis denotes $\log_2$ fold change (GW4064/Veh) of eRNAs and

the *y* axis denotes $-\log_{10}$ FDR of eRNAs. (B) A Bar plot showing the reads per million

(RPMs) of up regulated eRNAs by GW4064 treatment. (C) IGV genome browser track

showing RNA-seq signals from vehicle or GW4064 treated samples around the *FincoR*

locus and its neighboring regions. A zoom-in view of *FincoR* is shown below. Veh, vehicle;

GW, GW4064; pos, positive strand; neg, negative strand; Refseq, reference sequence.

(D) RT-qPCR data showing GW4064 induction of *FincoR* eRNA and *Shp* mRNA in the

liver (n=4/group). Data are presented as mean $\pm$ SEM. Statistical significance was

determined by the two-way ANOVA Sidak's multiple comparisons test with **p < 0.01 and

***p < 0.001.

Figure 2. Ligand-activated FXR directly activates transcription of eRNAs including

***FincoR* in the liver.** (A) Left: experimental outline. FXR-Flox and LKO male mice were

fasted overnight and treated with vehicle or GW4064, and livers were collected 1 h later

with nuclei isolated for Gro-seq (n=2/group). Right: a boxplot shows the Gro-Seq signals

for GW4064 up-regulated eRNAs in different conditions. RPKM: reads per kbp per million.

(B) IGV genome browser track showing RNA-seq, Gro-seq and ChIP-seq signals in the

FincoR locus. An arrow at the bottom points to the main enhancer FXR ChIP-seq peak

that overlaps the start site of *FincoR*. (C) ChIP assays were performed in the same liver

samples described in Fig. 1A to detect FXR, RXRα, and BRD4 occupancy at the FXR

binding peak region close to the transcription start site of *FincoR* (black arrow in Fig.2B).

(D) HepG2 cells were transfected with luciferase reporter expressing wild type FXRE or

mutant FXRE (see materials and methods) for 24 h before treatment with GW4064 for an

additional 6 h. Relative luciferase activities are shown. (C-D) Data are presented as

mean ± SEM. Statistical significance was determined by the Student's t test with *$p < 0.05$

and ***$p < 0.001$.

Figure 3. *FincoR* is a liver-specific nucleus-enriched eRNA. (A) Expression levels of

FincoR in various tissues after GW4064 treatment. Data from C57BL/6 male mice fasted

overnight and *i.p.* injected with vehicle or GW4064 (30 mg/kg) for 1 h (n=2/group). (B)

C57BL/6 male mice were fasted overnight and injected *i.p.* with vehicle or GW4064 (30

mg/kg) for 1 h. One small piece of liver was snap-frozen for later RNA isolation and the

remaining part was used for immediate primary hepatocyte isolation. Then, RNAs were

extracted from liver or primary hepatocytes and *FincoR* expression was measured

(n=3/group). (C) Agarose gel electrophoresis of PCR products generated in 5′ (left) and

3′ (right) RACE of *FincoR* in liver samples. Primer locations are shown. RACE, rapid

amplification of cDNA ends. GSP: gene specific primer. (D) A schematic diagram

showing location of *FincoR* relative to nearby genes in the mice genome. (E) PhyloCSF

analysis of the coding potential of *FincoR*. (F) In vitro translation of *FincoR* using the

Promega Transcend Non-Radioactive Translation Detection Systems. Luciferase is used

as a control for coding RNA. (G) qPCR analysis of *FincoR*, *36b4* in Poly(A)+ and Poly(A)-

RNA fractions from GW4064 treated mouse liver. (H) *FincoR* identified in the subcellular

fractions using cellular fractionation assays. Primary hepatocytes were isolated from

GW4064 or DMSO treated mice and the cytoplasm and nucleus fractions of these

hepatocytes were separated and both fractions were subjected to RNA extraction and

qPCR.

Figure 4. *FincoR* is induced by the hammerhead class of non-steroidal FXR

agonists, including GW4064 and tropifexor. (A) The chemical structures of the FXR

agonists including the hammerhead class of synthetic FXR agonists, non-hammerhead-

type synthetic agonists, semi-synthetic BA and natural BA. BA: bile acid. (B) qPCR data

showing *FincoR* expression levels in C57BL/6 mice liver respectively treated with

GW4064 (30 mg/kg), Cilofexor (30 mg/kg), Tropifexor (0.5 mg/kg), Fexaramine (100

mg/kg), or OCA (20 mg/kg) for 1 h (n=3~4/group). (C) *FincoR* expression levels in

C57BL/6 mice liver treated with OCA (20 mg/kg) or Fexaramine (100 mg/kg) for 4 hrs

(n=3/group). (D) Expression of *FincoR* in C57BL/6 mice liver after daily treatment with

OCA (20 mg/kg) for 7 days (n=5/group). *Shp* gene mRNA was measured as a positive

control. (E) Expression of *FincoR* in C57BL/6 mice fed with 0.5% cholic acid (CA) diet for

6 hrs (n=3~4/group). In panels B-E, all mice underwent over-night fasting. (B-E) Data are

presented as mean ± SEM. Statistical significance was determined by the Student's t test

with $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$.

Figure 5. Generation of CRISPR/Cas9-mediated *FincoR* liver-specific knockdown

mice. (A) Experimental scheme: male Cas9 mice were infected with adenovirus

expressing sgRNA for *FincoR* or a control for 1 week. Then the liver and serum were

collected from these mice after 4~5 hrs of fasting. (B) The expression of *FincoR* in the

liver was measured by qPCR (n=4/group). (C) Hepatic triglyceride, cholesterol, bile acid,

glycogen and serum NEFA were measured (n=5~7/group) (D) Male Cas9 mice were

infected with adenovirus expressing sgRNA for *FincoR* or control for 1 week. Then these

mice were fasted overnight and treated with GW4064 for 3 h before tissue collection.

RNA-seq profiles of expression of hepatic *FincoR* and the adjacent genes were shown

(n=2/group). (E) Genome-wide changes in mRNA expression shown in a volcano plot.

The numbers refer to the number of genes up- or down-regulated by 2-fold or more with

an adjusted *p*-value < 0.01. (F) Gene ontology analysis of biological pathways using

DAVID Tools for genes downregulated after *FincoR* knockdown. (B) Data are presented

as mean ± SEM. Statistical significance was determined by the Student's t test with ****p* <

0.001.

Figure 6. In diet-induced NASH mice, tropifexor-mediated beneficial effects on

reducing hepatic steatosis are largely independent of *FincoR*. (A-E) Male Cas9 mice

were fed a NASH diet for 12 weeks. The mice were randomly assigned to 3 groups and

infected with adenovirus expressing sgRNA for *FincoR* or control. Three days later, the

mice were treated with tropifexor (0.3 mg/kg) or vehicle for 12 days. The mice were

administered tropifexor or vehicle and fasted for 4 h before tissues were collected. (A)

Experimental scheme. (B) Hepatic *FincoR* expression was measured (n=6~7/group). (C)

Oil Red O staining of liver sections. Scale bar (50 μm). Image analyses were done using

Image J and the areas of stained field were quantified (n=5/group) (D) Hepatic TG,

hepatic cholesterol, gallbladder BA and hepatic BA levels were measured (n=6~7/group).

(E) mRNA levels in the liver of the indicated genes involved in bile acid regulation and

lipid regulation (n=6~7/group). (B, D and E) Data are presented as mean ± SEM.

Statistical significance was determined by the one-way ANOVA (Sidak's multiple

comparisons test) with *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$. Ad, adenovirus; H & E,

hematoxylin and eosin; TG, triglyceride; Veh, vehicle; ns, not significant.

Figure 7. In diet-induced NASH mice, tropifexor-mediated beneficial effects on

reducing liver fibrosis and inflammation are diminished by *FincoR* downregulation.

(A) Representative images from H & E, F4/80, Sirius Red and TUNEL staining of liver

sections from the same cohort of mice described in Figure 6. Scale bar (50 µm). Image

analyses were done using Image J and the area of collagen staining, TUNEL and F4/80

levels were quantified (n=5/group). (B) Serum ALT and AST levels were measured

(n=5/group). (C) IL-1β and CCL2 levels in the liver tissues were determined by ELISA

(n=5/group). (D) mRNA levels in the liver of the indicated genes involved in inflammation,

fibrosis and cell death (n=5/group). (E) Model: *FincoR* is a liver-enriched eRNA that is

induced specifically by hammerhead-type FXR agonists (top). In diet-induced NASH

mice, *FincoR* is required for tropifexor-mediated beneficial effects on reducing hepatic

inflammation, fibrosis, and cell death with the mechanisms to be determined (bottom). (A-

D) Data are presented as mean ± SEM. Statistical significance was determined by the

one-way ANOVA (Sidak's multiple comparisons test) with *$p < 0.05$, **$p < 0.01$ and ***$p <$

0.001.

1066

1067

 Supplementary Figure Legends

 Figure 1- figure supplement 1. (A, B) Examples of FXR-regulated eRNAs produced

 near the genes *Hes1* (A) and *Slc35g1* (B) were shown. (C) Time course expression of

 FincoR. C57BL/6 male mice were fasted overnight and injected *i.p.* with vehicle or

 GW4064 (30 mg/kg) for 1, 3, 6,12 h. Livers were collected at the indicated time points

 (n=3/group) and *FincoR* and mRNA levels of *Shp* and *Gcnt1* were measured. Data are

 presented as mean ± SEM. Statistical significance was determined by the two-way

 ANOVA Sidak's multiple comparisons test with *$p < 0.05$ and ***$p < 0.001$.

 Figure 1- figure supplement 2. (A) Volcano plot showing the differential expressed

 genes (DEGs) after GW4064 treatment (DESeq2 FDR<0.05). The numbers refer to the

 number of genes up- or down-regulated. (B) Bar plot showing the enriched Gene

 Ontology in terms of biological processes for up-regulated genes.

 Figure 2- figure supplement 1. (A) FXR protein levels in the livers isolated from FXR-

 Flox and FXR-LKO mice are shown. (B) Validation of hepatic FXR-dependent induction

 of *FincoR* by qPCR (n = 3 mice). The *Shp* gene was used as a control. Data are presented

 as mean ± SEM. Statistical significance was determined by the two-way ANOVA Tukey's

 multiple comparisons test with *$p < 0.05$ and **$p < 0.01$. (C) Metagene plots showing

 H3K27ac, H3K4me1, FXR, and RXRα ChIP-Seq profiles centered on up-regulated

 eRNAs.

 Figure 5- figure supplement 1. (A) Left: experimental scheme for the *FincoR* loss of

 function experiments. Right: illustration demonstrating the sequence targeted by the

 sgRNA in relation to the transcriptional and epigenetic profile. (B) The genomic DNAs

 from liver, spleen, intestine, brain, heart, muscle, kidney, lung and adipose tissue were

isolated and PCR was performed using the primers (**Supplementary Table S6**) to verify

tissue-specific knock-out. (C-G) RNA-seq profiles of expression of hepatic *Prune2* (C)*,*

PPP1r3g (D)*, Igfbp2* (E)*, Eda2r* (F) *and Fndc1* (G) are shown (n=2/group).

Figure 7- figure supplement 1. The effects of *FincoR* downregulation on NASH

pathologies. Male Cas9 mice were fed with a NASH diet for 12 weeks. Then these mice

were randomly assigned to 2 groups and infected with adenovirus expressing sgRNA for

FincoR or control. The tissues were collected 2 weeks later. Liver histology analysis was

performed and representative images were shown. Scale bar (50 µm).

1099

Supplementary Figure S1. Hepatic expression of *FincoR* is elevated in liver disease

associated with inflammation and fibrosis. (A) C57BL/6 mice were fed with a high fat

diet for 12 weeks. The liver RNAs were extracted and *FincoR* expression was measured

(n=5/group). (B) C57BL/6 mice were fed with a high fat diet with high fructose water for

12 weeks. The liver RNAs were extracted and *FincoR* expression was measured

(n=7/group). (C) C57BL/6 mice were treated with ANIT (75 mg/kg) for 48 h and then

sacrificed after 5 h of fasting. The liver RNAs were extracted, and *FincoR* expression was

measured (n=4~5/group). (D) C57BL/6 mice were bile duct ligated for 1 day or 3 days.

They were then sacrificed after 5 h of fasting. The liver RNAs were extracted and *FincoR*

expression was measured (n=4~5/group). (E) *FincoR* conservation between mice and

human as displayed in the UCSC Genome Browser. Red arrows indicate the conserved

region. (F) Human lncRNA XR_007061585.1 with sequence similarity to mouse *FincoR*

annotated in the NCBI genome data viewer. (G) Expression of lncRNA XR_007061585.1

in liver samples from normal individuals or patients with primary biliary cholangitis (PBC)

(n=14~15/group). (H) Expression of lncRNA XR_007061585.1 in liver samples from

normal individuals or patients with NAFLD-associated steatosis (n=12~15/group). (A-D,

G-H) Data are presented as mean ± SEM. Statistical significance was determined by the

Student's t test with *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$.

Supplementary Figure S2. Hepatic genome browser tracks of FXR, RXRα, LXR,

PPARα, and HNF4α binding peaks at the *FincoR* locus.

Supplementary Figure S3. PPARα occupancy in the *FincoR* enhancer region.

C57BL/6 male mice were fasted overnight with or without refeeding for 3 h and then

sacrificed. ChIP assays were performed in liver samples to detect PPARα occupancy at

the FXR binding peak region close to the transcription start site of *FincoR*.

1124

1125

1126

Supplemental Tables

Supplemental Table S1. Sequencing data generated in this study

Name	Total reads	Uniquely mapped reads
WT Liver Veh RNA-Seq rep1	50887391	38072659
WT Liver GW RNA-Seq rep1	54020150	40595856
WT Liver Veh RNA-Seq rep2	51268164	38898913
WT Liver GW RNA-Seq rep2	51899892	39487241
Flox Liver Veh GRO-Seq	62960996	30305910
Flox Liver GW GRO-Seq	59210073	26116795
FXR KO Liver Veh GRO-Seq	34153537	14379417
FXR KO Liver GW GRO-Seq	36430101	15008281
WT Liver RNA-Seq rep1	96262381	81801494
WT Liver RNA-Seq rep2	104979915	88365481
FincoR KD Liver RNA-Seq rep1	91654076	75346167
FincoR KD Liver RNA-Seq rep2	90211420	74745807

Supplemental Table S2 is attached as an Excel file.

Supplemental Table S3. Public ChIP-Seq dataset

target	accession ID
FXR	https://genome.ucsc.edu/cgi-bin/hgCustom?hgsid=1577671033_RPWVA7jbx3rDswKOeo2dazTsXxe0&hgct_table=ct_FxrLiver_2971
H3K27ac	ENCFF001KMI
H3K4me1	ENCFF001KNF
LXR	GSM864669
RXRα	GSM864674
PPARα	GSM864671
HNF4α	GSM2055887

Supplemental Table S4 is attached as an Excel file.

Supplemental Table S5: Predicted RNA binding proteins (RBPs) binding to *FincoR*

RBP	Length	Motif	Domain	number of binding sites
KHDRBS1	7	UAAAAAG	KH	1
RBM38	7	UUGUGUG, GUGUGUG	RRM	3
YBX2	7	CACACCA	CSD	11
YBX3	7	CACACCA	CSD	11
PCBP1	6	CUUUCC	KH	1
KHSRP	6	UGCAUG	KH	2
PTBP1	6	CUCUCU	RRM	1
PTBP2	6	CUCUCU	RRM	1
4KZD	5	GAAAC	N/A	1
4KZE	5	GAAAC	N/A	1
4Q9Q	5	GAAAC	N/A	1
DAZL	5	GUUCU	RRM	6

MSI1	5	GUAGU	RRM	1
TLR3	5	AAAGG	LRR;TIR	12

1143

1144

1145 Supplemental Table S6. Primer sequences used in this study

1146

1147 A. List of primer sequences for mouse qRT-PCR

Gene	Forward Primer (5'-3')	Reverse Primer (5'-3')
FincoR	GCAAAGCACCTTCTAGCACA	GTCAGGGAGCTAACGAATGC
Gcnt1	GCATCGCATCCTGCTTTGATA	GGTCTGCCTTAACCCGACTC
Shp	TCTGCAGGTCGTCCGACTAT	CAGGCAGTGGCTGTGAGAT
Cyp7a1	GGGATTGCTGTGGTAGTGAGC	GGTATGGAATCAACCCGTTGTC
Cyp8b1	CCTCTGGACAAGGGTTTTGTG	GCACCGTGAAGACATCCCC
Srebp1c	GCAGCCACCATCTAGCCTG	CAGCAGTGAGTCTGCCTTGAT
Lpin1	CATGCTTCGGAAAGTCCTTCA	GGTTATTCTTTGGCGTCAACCT
Scd1	TTCTTGCGATACACTCTGGTGC	CGGGATTGAATGTTCTTGTCGT
Ifng	GCG TCA TTG AAT CAC ACC TG	GAC CTG TGG GTT GTT GAC CT
Ccl3	TTCTCTGTACCATGACACTCTGC	CGTGGAATCTTCCGGCTGTAG
Ccl2	TTAAAAACCTGGATCGGAACCAA	GCATTAGCTTCAGATTTACGGGT
Ccr2	AAGAGGGCATTGGATTCACCACA	GCCGTGGATGAACTGAGGTAACA
Lcn2	GCAGGTGGTACGTTGTGGG	CTCTTGTAGCTCATAGATGGTGC
Col1a1	ATCGGTCATGCTCTCTCCAAACCA	ACTGCAACATGGAGACAGGTCAGA
Col1a2	CCTTTGTCAGAATACTGAGCAGC	GTAACTTCGTGCCTAGCAACA
Acta2	TCGGATACTTCAGCGTCAGGA	GTCCCAGACATCAGGGAGTAA
Eda2r	CACACTGCATAGTCTGCCCTC	GCCTTCTGGACCCGATTGA
Fndc1	GGGAGACATGGCAAACCTGT	TGGTAGGAGAGTATGTGGTGG
Ctsb	TCCTTGATCCTTCTTTCTTGCC	ACAGTGCCACACAGCTTCTTC
Ctss	GAAGTACGGCGTCTCATCTGG	CATGCCCACTTGGTAGGTATG
Bcl2	GTCGCTACCGTCGTGACTTC	CAGACATGCACCTACCCAGC
36b4	CCCTGAAGTGCTCGACATCA	TGCGGACACCCTCCAGAA
XR_007061585.1	TTGTCATCAAGCCCTGTTCA	TCTGCTTTGTCTGAGGACCA

1148

1149 B. List of primers used for mutagenesis

	Forward Primer (5'-3')	Reverse Primer (5'-3')
Mutation	GTGAATTCTCTCCTAGaTCcTTGACaTTCCAAGTCATGTAC	GTACATGACTTGGAAtGTCAAgGAtCTAGGAGAGAATTCAC

1150

1151

1152 ## C. List of primer sequences for mouse ChIP-qPCR

Gene	Forward Primer (5'-3')	Reverse Primer (5'-3')
FincoR	TCCTAGGTCTTTGACCTTCCAA	TGTGCCCACTGGTGAAATAG
Non-specific region	GTGTACACGCCCAAACTTGA	GTTCACCTCTCTAGCTCACCT

1153

1154 ## D. List of primer used for RACE

	Gene specific primer (5'-3')
5'RACE	GATTACGCCAAGCTT GGGAACTCACGTCTGGAAGCAGAAGCTG
3'RACE	GATTACGCCAAGCTT GGACACTCGGGAATTGACAGTGGGTCAGG
3'RACE nest primer	GATTACGCCAAGCTT GTGGGGGATAGCACAAGCGCCAGGATGC

1155

1156 ## E. List of primer used for genotyping

Forward Primer (5'-3')	Reverse Primer (5'-3')
GTGATTTCCTTTCCCACAGC	GTCAGGGAGCTAACGAATGC

1157

Figure 1

A RNA-seq (GW4064: 1 h)

B *Up regulated eRNAs*

C

chr19:17,131,719-17,526,635

chr19:17,300,000-17,340,000

D *FincoR*   *Shp*

Figure 2

Figure 3

Figure 4

A

GW4064　　　　Cilofexor　　　　Tropifexor (LJN452)

Hammerhead-type synthetic FXR agonists

Fexaramine　　　　Obeticholic acid (OCA)　　　　Cholic acid

non-hammerhead-type synthetic FXR agonist　　　semi-synthetic BA　　　natural BA

B *FincoR*

Rel RNA level

Veh
GW4064
Cilofexor
Tropifexor
OCA
Fex

C *FincoR*

Rel RNA level

Veh
OCA
Fex

D *Shp* *FincoR*

Rel RNA level

Veh
OCA

E *FincoR*

Rel RNA level

Normal chow
0.5% CA diet

Figure 5

A

Ad-sgRNA Ctrl or
Ad-sgRNA *FincoR* → FAST 4~5 hrs → Sac

Cas9   1 week

B

FincoR

Rel RNA level

☐ Ad-sgRNA Ctrl
▨ Ad-sgRNA *FincoR*

***

C

Triglyceride (mg/g liver)

Cholesterol (mg/g liver)

Liver BA (nmol/g liver)

Glycogen (µg/mg liver)

Serum NEFA (µM)

D

Ad-sgRNA Ctrl or
Ad-sgRNA *FincoR* → Fast O/N → GW4064 →(3 hrs)→ liver for RNA-seq

Cas9   1 week

MM10   chr19:17,158,000-17,483,300

325 Kb

17,200 kb   17,300 kb   17,400 kb

FincoR

Ctrl 1 (pos)
Ctrl 2 (pos)
LKD 1 (pos)
LKD 2 (pos)
Ctrl 1 (neg)
Ctrl 2 (neg)
LKD 1 (neg)
LKD 2 (neg)

Refseq

Prune2   *Gcnt1*   *Rfk*   *Pcsk5*

E

Down:53 Up:18

-log$_{10}$ FDR

log$_2$ Fold Change

F

GO Biological Process

positive regulation of fatty acid oxidation
positive regulation of organelle organization
tryptophan metabolic process
tryptophan catabolic process
indolalkylamine catabolic process
positive regulation of glucose metabolic process
regulation of fatty acid oxidation
positive regulation of fatty acid metabolic process
2-oxoglutarate metabolic process
regulation of insulin-like growth factor
receptor signaling pathway

-log$_{10}$ (Adjusted P-value)

Figure 6

Ad-sgRNA Ctrl + Veh   Ad-sgRNA Ctrl + Tropifexor   Ad-sgRNA *FincoR* + Tropifexor

A

Cas9 → NASH diet 12 weeks → 1: Ad-sgRNA Ctrl / 2: Ad sgRNA-Ctrl / 3: Ad sgRNA-*FincoR* → 3 days → D1 Tropifexor (0.3 mg/Kg)/Veh → D12 Tropifexor Fast 4 hrs → Sac

B

FincoR

Rel RNA level

C

Ad-sgRNA: Ctrl   Ctrl   *FincoR*

Tropifexor   -   +   +

Oil Red O

Oil Red O (% of the field)

D

Hepatic TG — Triglyceride (mg/g liver)

Hepatic cholesterol — Cholesterol (mg/g liver)

Hepatic BA — Liver bile acid (µmol/g liver)

Gallbladder BA — Gallbladder bile acid content (µmol)

E

Bile acid regulation

Shp   *Cyp7a1*   *Cyp8b1*

Rel RNA level

Lipid regulation

Srebp1c   *Lpin1*   *Scd1*

Ad-sgRNA Ctrl + Veh   Ad-sgRNA Ctrl + Tropifexor   Ad-sgRNA *FincoR* + Tropifexor

Figure 7

Figure 1-figure supplement 1

Figure 1–figure supplement 2

A

B

Figure 2-figure supplement 1

A

FXR-Flox FXR-LKO

FXR

actin

B

Shp

FincoR

Rel RNA level

GW4064 − + − + − + − +

C

H3K27ac   H3K4me1   FXR   RXRα

Chip-Seq reads

Distance from peak center (Kb)

Figure 5-figure supplement 1

Figure 7-figure supplement 1

Supplementary Figure S1

Supplementary Figure S2

Supplementary Figure S3

Fast O/N or

Fast O/N plus refed 3 hrs

→ Sac

C57BL/6