**Hammerhead-type FXR agonists induce an eRNA FincoR that ameliorates**

**nonalcoholic steatohepatitis in mice**

Jinjing Chen[a, #], Ruoyu Wang[b, #], Feng Xiong[b], Hao Sun[a], Byron Kemper[a],

Wenbo Li[b,*] and Jongsook Kim Kemper[a,*]

[a]Department of Molecular and Integrative Physiology, University of Illinois at Urbana-

Champaign, Urbana, IL, USA and [b]Department of Biochemistry and Molecular Biology,

McGovern Medical School, University of Texas Health Science Center, Houston, TX,

USA

Running Title: Role of an FXR-induced eRNA, FincoR, in NASH

[#]JC and RW contributed equally to this study.

Present Address: Jinjing Chen, Department of Pharmacology and Toxicology, University

of Arizona, Tucson, AZ, USA

* To whom correspondence may be addressed:

* Wenbo Li, Department of Biochemistry and Molecular Biology, McGovern Medical

School, University of Texas Health Science Center, Houston, TX, USA

Graduate School of Biomedical Sciences, University of Texas MD Anderson Cancer

Center and UTHealth, Houston, TX, USA

Tel: 713-500-6103, e-mail**:**  Wenbo.Li@uth.tmc.edu

* J. Kim Kemper, Department of Molecular and Integrative Physiology,

University of Illinois at Urbana-Champaign, Urbana, IL

Tel: 217-722-7961, e-mail: jongsook@illinois.edu

**Key words:** FXR, nuclear receptor, lncRNA, enhancer, enhancer RNA, tropifexor,

hepatic inflammation, NASH, NAFLD, RNA-seq, Gro-seq

**Author Contributions:**

JC, RW, WL and JK designed research; JC, RW, FX, and HS performed experiments;

RW and WL performed bioinformatic analyses; JC, RW, WL, and BK analyzed data;

and JC, RW, BK, WL, and JK wrote the paper.

## Abstract

The nuclear receptor, Farnesoid X Receptor (FXR/NR1H4), is increasingly recognized as a promising drug target for metabolic diseases, including nonalcoholic steatohepatitis (NASH). Protein coding genes regulated by FXR are well known, but whether FXR also acts through regulation of expression of long non-coding RNAs (lncRNAs), which vastly outnumber protein-coding genes, remains unknown. Utilizing RNA-seq and GRO-seq analyses in mouse liver, we found that FXR induces many RNA transcripts from chromatin regions bearing enhancer features. Among these we discovered a previously unannotated liver-enriched enhancer-derived lncRNA (eRNA), termed *FincoR*. We further show that *FincoR* is specifically induced by hammerhead-type FXR agonists, including GW4064 and tropifexor. CRISPR/Cas9-mediated liver-specific knockdown of *FincoR* in dietary NASH mice reduced the beneficial effects of tropifexor, an FXR agonist currently in clinical trials for NASH and primary biliary cholangitis (PBC), indicating that that amelioration of liver fibrosis and inflammation in NASH treatment by tropifexor is mediated in part by FincoR. Overall, our results indicate that pharmacological activation of FXR by hammerhead-type agonists induces a novel eRNA, *FincoR*, ameliorating NASH in mice. *FincoR* may represent a new drug target for combating metabolic disorders, including NASH.

**Introduction**

Nonalcoholic fatty liver disease (NAFLD) is the most common chronic liver disease and a leading cause of liver transplants and liver-related death *(Friedman and Sanyal 2018)*. NAFLD begins with simple steatosis but may further progress to a severe form, non-alcoholic steatohepatitis (NASH), and later, fatal cirrhosis and liver cancer *(Friedman and Sanyal, 2018)*. Despite its striking global increase and clinical importance, there is no approved drug for NASH. The urgent need for development of therapeutic agents for NASH has greatly increased research interest in the nuclear receptor, Farnesoid X Receptor (FXR, NR1H4) *(Evans and Mangelsdorf, 2014)*.

FXR is activated by its physiological ligands, bile acids (BAs), and regulates expression of genes involved in BA, lipid, and glucose metabolism and hepatic autophagy, which maintains metabolite levels and metabolic homeostasis *(Lee et al., 2006; Calkin and Tontonoz, 2012; Lee et al., 2014; Seok et al., 2014; Kliewer and Mangelsdorf, 2015)*. Ligand-activated FXR also protects against hepatic inflammation and liver injury *(Wang et al., 2008; Jung et al., 2020)*. The action of FXR, similar to other nuclear receptors, is achieved primarily by binding to chromatin at specific gene loci to regulate the transcription of target genes *(Lee et al., 2006; Thomas et al., 2010; Lee et al., 2012; Calkin and Tontonoz, 2012)*. Consistent with its crucial physiological functions, FXR is increasingly recognized as a promising drug target, particularly for liver diseases, such as NASH and primary biliary cholangitis (PBC) *(Downes et al., 2003; Ali et al., 2015; Abenavoli et al., 2018; Kremoser, 2021)*. For example, semi-synthetic or non-steroidal synthetic agonists of FXR, including obeticholic acid (OCA) and hammerhead-type agonists, such as tropifexor and cilofexor, are currently under clinical trials for NASH and

PBC patients *(Tully et al., 2017; Abenavoli et al., 2018; Kremoser, 2021; Sanyal et al.,*

*2023)*. However, how pharmacological activation of FXR mediates such beneficial

therapeutic effects is poorly understood.

Non-protein-coding RNAs (ncRNAs) are one of the fascinating discoveries of

modern biology *(Cech and Steitz, 2014)*. Indeed, only about 2% of the human genome

encodes proteins, while the vast majority are transcribed into ncRNAs *(The ENCODE*

*Project Consortium, 2012)*. Of these ncRNAs, long non-coding RNAs (lncRNAs) are a

group of transcripts longer than 200 nucleotides and play important roles in diverse

biological processes *(Lam et al., 2014; Li et al., 2016; Sartorelli and Lauberth, 2020;*

*Statello et al., 2021; Mattick et al., 2023)*. Many lncRNAs are produced from genomic

regions bearing epigenetic features of enhancers *(Hon et al., 2017; Mattick et al., 2023)*.

This is consistent with the idea that many transcriptional enhancers actively transcribe

noncoding RNAs that are referred to as eRNAs, some of which are functionally important

for enhancer functions *(Lai and Shiekhattar, 2014; Li et al., 2016)*. Due to the increasing

technical sensitivity of RNA detection, many enhancers have been found to produce

eRNAs (>85k in humans and >57k in mice) *(Hirabayashi et al., 2019)*. Many eRNAs have

not yet been annotated in current lncRNA databases, such as GENCODE. Exploring

eRNA landscapes and functions in diverse biology and disease states will facilitate our

understanding of both lncRNAs and enhancers *(Li et al., 2016; Mattick et al., 2023)*.

However, the landscape of eRNAs in mouse liver has been minimally explored *(Fang et*

*al., 2014)*, and has not been studied in response to specific nuclear receptor activation,

such as by FXR. Moreover, functional role of eRNAs in vivo in intact organisms is

understudied.

FXR was shown to regulate expression of small non-coding microRNA (miR)

genes, for example *miR-34a and miR-802 (Lee et al., 2010; Seok et al., 2021)*, but it has

not been reported whether FXR mediates its function through regulating lncRNAs, which

far outnumber miRs. Because FXR directly regulates expression of its target genes

*(Thomas et al., 2010; Lee et al., 2012)*, we examined if FXR regulates enhancer-derived

lncRNAs, and if such transcripts participate in its physiological or pharmacological

functions. By utilizing RNA-seq and GRO-seq analyses of mouse liver treated by FXR

ligands, we identified a series of FXR-regulated eRNAs, including many that were

previously unannotated. Among these, we focused on a highly induced and abundantly

expressed eRNA that we referred to as FXR-induced non-coding RNA (*FincoR*) for

functional studies. *FincoR* is highly enriched in mouse liver and is induced specifically by

hammerhead-type FXR agonists, including GW4064 and tropifexor. In vivo studies

utilizing CRISPR/Cas9-mediated liver-specific knockdown of *FincoR* in dietary NASH

mice indicated that *FincoR* is critically involved in mediating the beneficial

pharmacological effects of tropifexor in reducing liver fibrosis and inflammation.

**Results**

**Activation of FXR by GW4064 induces a novel eRNA, *FincoR*, in mouse liver**

To identify enhancer-derived lncRNAs (eRNAs) potentially regulated by FXR, we first

obtained a list of putative enhancers in mouse liver based on ENCODE H3K27ac ChIP-

Seq data (ENCFF001KMI, see **methods**). Then, we performed ribo-depleted total RNA-

seq in the livers from mice treated with a specific FXR agonist, GW4064, to identify

transcripts produced from these enhancer regions (see **Supplementary Table S1**). To

avoid confounding issues of RNA signals from genes, we specifically focused on

intergenic enhancer regions (+/- 3kb from H3K27ac peak center) that harbor discernible

RNA-Seq signals (RPKM > 1).

This genomic analysis resulted in identification of 190 high-confidence eRNAs in

mouse liver. Among these, 14 eRNAs were upregulated and 5 were downregulated by

GW4064 treatment (FDR < 0.05, Log2 FC > 1, **Figure 1A**) (see **Supplementary Table**

**S2**). FXR-regulated eRNAs were produced adjacent to many genes with important roles

in liver metabolism and disease, for example, *Hes1* (**Figure 1-Figure Supplement 1A,**

**B**). One of the most robustly induced eRNAs was <u>F</u>XR-<u>i</u>nduced <u>n</u>on-<u>co</u>ding <u>R</u>NA

(*FincoR)*, an unannotated novel transcript located on chromosome 19 (**Figure 1A, B, C**).

Reverse transcription qPCR (RT-qPCR) confirmed that treatment with GW4064

substantially induced expression of *FincoR,* more than 10-fold in mouse liver, which is

similar to induction of *Shp*, a well-known FXR target gene (**Figure 1D**) *(Goodwin et al.,*

*2000; Claudel et al., 2005; Evans and Mangelsdorf, 2014)*. Induction of *FincoR* by

GW4064 was transient, peaked within 1 h and then, declined gradually, similar pattern to

*Shp* (**Figure 1-Figure Supplement 1C**). Expression of genes adjacent to *FincoR*,

including *Gcnt1, Rfk, Pcsk5 and Prune2,* did not change after acute 1 h FXR activation

as shown by RNA-seq (**Figure 1C**), and confirmed for *Gcnt1* by time course qPCR

(**Figure 1-Figure Supplement 1C**).

We also found that the short GW4064 treatment resulted in 590 up-regulated

genes and 500 down-regulated genes (FDR < 0.05) (**Figure 1-Figure Supplement 2A**).

GO enrichment analysis of these differentially expressed genes (DEGs) revealed their

roles in the regulation of triglyceride, fatty acid, and cholesterol metabolism (**Figure 1-**

**Figure Supplement 2B**), which is consistent with known roles of FXR in these

physiological processes *(Claudel et al., 2005)*.

**Ligand-activated FXR directly activates transcription of eRNAs, including *FincoR*.**

We sought to examine, 1) if these eRNAs were directly activated by FXR, and 2) if the

activation takes place transcriptionally using FXR liver-specific knockout (FXR-LKO) mice

that were treated with GW4064 to determine if FXR was required for induction of eRNAs

by GW4064. FXR-LKO mice were generated from FXR floxed mice (FXR-flox) **(Figure 2-**

**Figure Supplement 1A**) and transcription of eRNAs was detected by global run-on

sequencing (GRO-seq), a widely used method to detect nascent RNA transcription,

including eRNAs *(Lam et al., 2014; Li et al., 2016; Sartorelli and Lauberth, 2020)*. GRO-

seq showed that FXR-induced eRNAs were activated transcriptionally by GW4064 in

FXR-flox mice, but such induction was abolished in FXR-LKO mice (**Figure 2A**). In

particular, *FincoR* is robustly induced in the GRO-Seq analysis and its induction is

dependent on hepatic FXR (**Figure 2B**). RT-qPCR confirmed the FXR-dependent

expression of *FincoR,* similar to that of *Shp* (**Figure 2-Figure Supplement 1B**).

We next examined if FXR binds to these enhancers that produce the identified

eRNAs by analyzing published mouse liver ChIP-Seq data for FXR (**Supplementary**

**Table S3**) *(Thomas et al., 2010; Lee et al., 2012)*. FXR binding was strongly enriched at

the enhancers associated with FXR-induced eRNAs as were the enhancer marks

H3K27ac and H3K4me1 (**Figure 2-Figure Supplement 1C**). In **Figure 2B**, the results

show the binding of FXR and the presence of histone marks at the *FincoR* enhancer

region determined by ChIP-seq as compared with nascent transcripts detected by Gro-

seq. These analyses support the conclusion that activation of FXR transcriptionally

induces this series of eRNAs via chromatin binding at these enhancers, including *FincoR*.

We validated FXR binding at the enhancer region that produces *FincoR* using

mouse liver ChIP (**Figure 2C**). We also examined binding of a well-known DNA binding

partner of FXR, Retinoid x Receptor alpha (RXRα/NR2B1) *(Evans and Mangelsdorf,*

*2014; Zheng et al., 2018)*, and bromodomain-containing protein 4 (BRD4), an acetylated

histone reader protein that often binds at active enhancers *(Chen et al., 2016; Li et al.,*

*2016; Rahnamoun et al., 2018; Sartorelli and Lauberth, 2020)* and a transcriptional

coactivator of FXR *(Jung et al., 2020)*. GW4064 treatment resulted in substantial

increases in recruitment of both FXR and RXRα to the enhancer region close to the

transcription start site of *FincoR* (arrow shown below in Figure 2B), whereas binding was

not detected at a control region (**Figure 2C**). We also found that BRD4 occupancy was

increased at this enhancer region after GW4064 treatment (**Figure 2C**). These results

indicate that GW4064 activation of FXR leads to increased occupancy of the FXR/RXRα

heterodimer and BRD4 to the enhancer region to upregulate *FincoR* eRNA in the liver.

We identified an inverted repeat1 (IR1) motif that was known to bind FXR *(Lee et*

*al., 2006; Calkin and Tontonoz, 2012)* within the major FXR binding peak near the start

site of *FincoR* (arrow shown below in Figure 2B), which we also refer to as FXRE (**Figure**

**2B, C**). We examined the functionality of this IR1 motif for mediating transcriptional

activation by GW4064 using reporter assays (**Figure 2D**). We cloned the region

containing the IR1 motif into the pGL4.23 luciferase reporter and generated a mutated

IR1 motif construct as a comparison (**Figure 2D**). After transfection into human hepatic

HepG2 cells, GW4064 treatment significantly elevated the luciferase activity of the

reporter with the wild type IR1 motif, but not with the mutated IR1 motif (**Figure 2D**).

Together, these results suggest that GW4064-activated FXR directly upregulates *FincoR*

expression.

***FincoR* is a liver-specific nucleus-enriched eRNA**

Because enhancers and eRNAs generally act in a tissue-specific manner *(Li et al., 2016;*

*Sartorelli and Lauberth, 2020)*, we examined the tissue-specific expression of *FincoR* in

mice. Strikingly, *FincoR* is highly expressed in the liver and it is expressed at extremely

low levels in most other tissues, except for a detectable, but still fairly low, level in the

lung. GW4064 treatment resulted in induction of *FincoR* specifically in the liver (**Figure**

**3A**). The level of *FincoR* detected in isolated primary mouse hepatocytes (PMHs) was

similar to that in mouse liver tissue treated with GW4064, suggesting that the majority of

*FincoR* is present in hepatocytes (**Figure 3B**).

219               By using the 5' and 3' rapid amplification of cDNA ends (RACE), we

identified one transcript of FincoR that is of approximately 3.7 kb in length (**Figure 3C**).

However, based on RNA-seq, the length of *FincoR* is over 10 kb (**Figure 1C**), suggesting

there are likely additional multiple RNA isoforms that we were not able to identify by

RACE. We next analyzed the coding potential of *FincoR* utilizing a comparative genomic

program, PhyloCSF. While adjacent genes *Gcnt1* and *Prune2* (**Figure 3D, E**) were

correctly predicted to encode proteins, *FincoR* did not contain a potential protein-coding

open reading frame (**Figure 3E**). Consistent with this bioinformatic prediction, a vector

expressing *FincoR* failed to produce any proteins in an in vitro transcription/translation

assay (**Figure 3F**) confirming that the *FincoR* transcript is a noncoding RNA. *FincoR*

transcripts were enriched by binding to oligo-dT beads, suggesting that *FincoR* is 3'

polyadenylated (**Figure 3G**). Further, *FincoR* was detected in the nuclear compartment

and GW4064 treatment increased the nuclear abundance of *FincoR* (**Figure 3H**),

consistent with its potential transcriptional regulatory function. Together, these results

from molecular biochemical characterization studies reveal that *FincoR* is a liver-enriched

nuclear polyadenylated eRNA.

***FincoR* is induced specifically by the hammerhead-type synthetic FXR agonists**

To determine whether induction of the FXR-induced FincoR is ligand-specific, we

examined the effects of hammerhead-type synthetic FXR agonists, GW4064, cilofexor,

and tropifexor; a semi-synthetic agonist, OCA; and a non-hammerhead-type gut-specific

agonist, fexaramine (**Figure 4A**) *(Downes et al., 2003; Fang et al., 2015)*. Remarkably,

treatment with each of the hammerhead type agonists for 1 h resulted in a robust induction

of *FincoR* (**Figure 4B**), whereas *FincoR* levels were unchanged after treatment with OCA

or fexaramine for 1 h (**Figure 4B**). Treatment with OCA for 4 h or even one-week

treatment with OCA failed to induce hepatic *FincoR* in mice, while expression of *Shp* was

significantly induced (**Figure 4C, D**). Acute feeding with a diet supplemented with 0.5%

cholic acid (CA), a primary BA, for 6 h also failed to induce *FincoR* expression (**Figure**

**4E**). Collectively, these results demonstrate that *FincoR* is induced specifically by

hammerhead type FXR agonists.

**Generation of CRISPR/Cas9-mediated *FincoR* liver-specific knockdown mice**

To explore the functional role of hepatic FincoR, we utilized the CRISPR/Cas9 technique

to generate FincoR liver-specific knockdown (FincoR-LKD) mice (**Figure 5-Figure**

**Supplement 1A**). Adenoviral-mediated sgRNA expression in Cas9 mice resulted in

downregulation of FincoR specifically in the liver by about 60%, but not in other tissues

(**Figure 5B**). As FXR is a key regulator of bile acid, cholesterol, lipid, and glucose

metabolism and FincoR is specifically regulated by FXR, FincoR may have a role in the

metabolic process in physiology and disease. We examined liver triglyceride, cholesterol,

bile acid, glycogen and serum non-esterified fatty acids (NEFA) but *FincoR*

downregulation did not result in any significant changes under physiological conditions

(**Figure 5C**).

To explore the molecular signatures and pathways affected by FincoR, we

examined the global gene expression by RNA-seq analysis in mouse liver after *FincoR*

knock-down (**Figure 5D**). While FincoR was markedly downregulated, the neighboring

genes, such as *Gcnt1, Rfk, Pcsk5*, were largely unchanged (**Figure 5D**). The RNA-seq

analysis revealed 18 up-regulated genes and 53 downregulated genes in FincoR-LKD

liver (**Supplementary Table S4; Figure 5E**). GO analysis indicated that these genes

were enriched in pathways involved in fatty acid oxidation, organelle organization and

metabolic process (**Figure 5F**). Among the down-regulated genes, *Ppp1r3g* which has a

role in controlling glycogen synthesis, and *Igfbp2* which functions in insulin resistance,

were markedly reduced (**Supplementary Table S4; Figure 5-Figure Supplement 1D,**

**E**). Among the up-regulated genes, expression of *Eda2r* was substantially increased,

which belongs to a member of tumor necrosis factor receptor superfamily and is involved

in inflammation, the immune response and development (**Figure 5-Figure Supplement**

**1F**). *Fndc1* which is involved in fibronectin matrix remodeling was also suppressed by

FincoR (and thus upregulated upon its KO, **Figure 5-Figure Supplement 1G**). These

studies suggested that FincoR may have a role in modulating metabolic homeostasis by

regulating genes involved in metabolism and inflammation.

**Amelioration of hepatic steatosis mediated by tropifexor is independent of FincoR**

**in diet-induced NASH mice**

Tropifexor, also known as LJN452, is a highly potent hammerhead type FXR agonist that

is currently under clinical trials for NASH and PBC patients *(Tully et al., 2017; Kremoser,*

*2021; Sanyal et al., 2023)*. Because FincoR is induced specifically by the hammerhead

class of FXR agonist (**Figure 4**) and has a potential role in the regulation of metabolism

and inflammation (**Figure 5**), we hypothesized that FincoR may play a role in tropifexor-

mediated beneficial effects on reducing NASH pathologies in mice.

We adopted a mouse model that had been fed the Amylin Liver NASH-promoting

(AMLN) diet *(Zhao et al., 2018; Hernandez et al., 2019; Sun et al., 2022)*, and examined

the effects of liver-specific downregulation of FincoR in potentially impacting tropifexor

effects on NASH pathology. Cas9 mice fed AMLN diet for 12 weeks were injected via tail

veins with adenovirus expressing control sgRNA or FincoR sgRNA, respectively, and then

treated daily with tropifexor (0.3 mg/kg) for 12 days (**Figure 6A**). In these mice, as a

technical validation of RNA induction and knockdown, FincoR levels were significantly

increased by FXR agonist tropifexor, and the increase was blocked by adenovirus

expressing sgRNA for FincoR (**Figure 6B**).

We then examined the effect of tropifexor treatment and FincoR downregulation

on hepatic steatosis in these mice. Tropifexor treatment markedly reduced neutral lipids

determined by Oil Red O staining of liver sections (**Figure 6C**) and liver TG levels (**Figure**

**6D**), and these beneficial effects on reducing fatty liver were not altered by FincoR

downregulation. Also, FincoR downregulation had little effects on liver cholesterol and

gallbladder bile acid levels, although these levels were reduced by Tropifexor (**Figure 6D**).

Consistent with the phenotypes, hepatic expression of key genes involved in BA

synthesis was dramatically reduced by tropifexor treatment (i.e., *Cyp7a1 and Cyp8b1*),

which is consistent with decreased gallbladder BA levels by this drug (**Figure 6E**).

Similarly, tropifexor also lowered lipid synthesis genes (*Srebp1, Lipin*, *Scd1*), consistent

with decreased liver TG levels.  However, downregulation of *FincoR* did not result in

changes in mRNA levels of these genes (**Figure 6E**). These results indicate that

tropifexor-mediated beneficial effects on reducing hepatic steatosis are independent of

*FincoR*.

**FincoR facilitates alleviation of liver inflammation by tropifexor in diet-induced**

**NASH**

Tropifexor ameliorated fibrotic NASH pathologies in preclinical studies *(Tully et al., 2017;*

*Hernandez et al., 2019)* and recently completed phase 2 clinical trials for NASH patients

*(Sanyal et al., 2023)*. We, therefore, further examined the effects of FincoR

downregulation on altering other NASH pathologies, including hepatocellular apoptosis,

liver fibrosis and inflammation.

In the same AMLN diet-fed mice above (**Figure 6**), analyses of liver sections

revealed that tropifexor treatment reduced hepatocyte swelling/ballooning (H&E staining),

decreased numbers of apoptotic cells (TUNEL staining), alleviated fibrosis (Sirius red

staining), and lowered infiltration of macrophages (F4/80 staining) (**Figure 7A**).

Remarkably, these tropifexor-mediated beneficial effects on NASH pathologies were all

markedly diminished by FincoR downregulation (**Figure 7A**). This was consistently found

in various liver lobes and two representative pictures from 2 different lobes are shown

(**Figure 7A**). In control experiments, FincoR downregulation in vehicle-treated mice did

not result in marked changes in NASH pathologies (**Figure 7-Figure Supplement 1**).

These results demonstrate that FincoR is required for tropifexor-mediated beneficial

effects on reducing NASH pathologies, specifically reducing hepatic inflammation, fibrosis

and hepatocyte apoptosis. Consistent with these results, serum ALT levels, an indicator

of liver damage, were significantly elevated after FincoR downregulation (**Figure 7B**).

Protein levels of key inflammatory markers, IL1β and CCL2, in liver extracts were also

elevated after FincoR downregulation (**Figure 7C**).

Consistent with the phenotypes from histological analyses, hepatic expression of

fibrosis and inflammation was altered by FincoR knockdown. For example, tropifexor

treatment reduced mRNA levels of several genes that promote fibrosis (*Col1a1, Col1a2,*

*Acta2*) and hepatic inflammation (*Eda2r, Ifng, Ccl3*), whereas these reductions were

largely reversed by FincoR downregulation (**Figure 7D**). We also detected increased

expression of inflammatory genes (*Ccl2, Ccr2, Lcn2) and* extracellular matrix remodeling

gene *(Fndc1)* in mice liver with FincoR knockdown (**Figure 7D**)*.* Tropifexor treatment

suppresses hepatic apoptosis by reducing pro-apoptotic genes (*Ctsb, Ctss*) and

upregulating anti-apoptotic genes such as *Bcl2 (Warren et al., 2019)*. Importantly, these

effects were significantly reversed by FincoR downregulation (**Figure 7D**).

Collectively, these results demonstrate that in diet-induced NASH mice,

pharmacological activation of FXR by tropifexor reduced fibrosis, apoptosis, and

inflammation in NASH mice, which was dependent, at least in part, on the induction of

*FincoR*.

**Discussion**

FXR maintains metabolic homeostasis by transcriptional regulation of genes. Direct

regulation of protein-coding genes by FXR, including *Shp*, is well characterized. In this

study we show that FXR also mediates its functions by induction of lncRNA genes, which

vastly outnumber protein-coding genes. We further show that pharmacological activation

of FXR by hammerhead-type agonists induces a liver-specific enhancer-derived lncRNA,

which we named FincoR, that contributes to reduction of NASH pathologies in mice.

FincoR is specifically induced by the hammerhead class of FXR agonists, such as

GW4064, cilofexor, and tropifexor. It was not induced by other FXR ligands, including its

endogenous ligand, cholic acid, suggesting that FincoR does not contribute to

physiological functions of FXR, but instead plays a role in pharmacological FXR

responses and possibly in pathological processes. GW4064 is the mother compound of

these isoxazole type ligands but is not an ideal therapeutic agent because of its poor

water solubility and pharmacokinetics *(Abel et al., 2010)*. In contrast, tropifexor and

cilofexor have better pharmacokinetics and were generally well-tolerated in clinical trials

for NASH and PBC patients *(Tully et al., 2017; Abenavoli et al., 2018; Kremoser, 2021;*

*Sanyal et al., 2023)*, but the underlying mechanisms for their beneficial effects are poorly

understood. Intriguingly, a recent study showed that the gene signature regulated by

tropifexor-activated FXR appears to be broader than that of OCA, partly because the

tropifexor backbone allows a more favorable interaction of FXR with coactivators or

epigenomic modulators *(Hernandez et al., 2019)*. Further, tropifexor was shown to

regulate distinct sets of genes, particularly those promoting fibrosis, inflammation, and

oxidative stress, in experimental NASH *(Hernandez et al., 2019)*. Utilizing CRISPR/Cas9-

mediated liver-specific knockdown of FincoR in diet-induced NASH mice, we demonstrate

that beneficial effects on reducing liver fibrosis, inflammation, and apoptosis mediated by

tropifexor were largely dependent on FincoR (Model, **Figure 7E**).

The roles of regulatory RNAs in liver function and diseases and as therapeutic

targets or drugs are increasingly being appreciated (*Sallam et al., 2016; Zhao et al., 2018;*

*Brocker et al., 2020; Li et al., 2021)*. For example, a critical role for an oxysterol nuclear

receptor LXR-induced lncRNA, LeXis, in feedback modulation of cholesterol biosynthesis

has been shown *(Sallam et al., 2016)*. Recently, the role of a lncRNA, Pair, in the liver

phenylalanine metabolism has been demosntrated *(Li et al., 2021)*. Enhancer RNAs are

a less characterized class of lncRNAs and are highly associated with enhancer functions

in gene regulation *(Li et al., 2016)*. Numerous studies have revealed transcriptional roles

for eRNAs in various cellular processes *(Lai and Shiekhattar, 2014; Lam et al., 2014; Li*

*et al., 2016)* but most previous eRNA studies have used culture cells *(Li et al., 2013;*

*Hsieh et al., 2014)*, with only a few in vivo studies in mouse models *(Mirtschink et al.,*

*2019; Tang et al., 2023)*. In our current study, through integrative analysis of

transcriptome and histone mark ChIP-Seq, we identified FXR-induced eRNAs, including

the highly induced *FincoR*. We further examined the expression of function of *FincoR*

utilizing mouse models. Our current work characterized the role of *FincoR* in gene

regulation and in mediating beneficial pharmacological effects of FXR agonists in NASH,

representing an important progress in understanding physiological/pathological roles of

eRNAs in vivo. Future work is warranted to understand the exact mechanisms by which

*FincoR* facilitates action of FXR agonists to alleviate inflammation, fibrosis and apoptosis.

While this study focused on regulation of *FincoR* by pharmacological activation of

FXR, physiological and pathological regulations of *FincoR* appear to be complex. *FincoR*

was not induced by acute cholic acid feeding, but *FincoR* levels were elevated in mouse

models of cholestatic liver (data not shown), where BA metabolism is dysregulated. Since

different BAs can activate or repress the gene-regulating function of FXR *(Wahlstrom et*

*al., 2016)*, altered BA composition in cholestatic liver may contribute to induction of

*FincoR*. Further, binding peaks for multiple nuclear receptors, FXR, LXR, PPARα, RXRα,

and HNF-4α, were detected in the *FincoR* gene locus so that regulation of *FincoR* likely

involves combinatorial regulation by multiple nuclear receptors (**Other Supplementary**

**Figure 1**). Interestingly, occupancy of the nuclear receptor PPARα at the enhancer region

was increased in fasted mice (**Other Supplementary Figure 2**). It will be interesting to

see whether expression of *FincoR* is differently regulated by these nuclear receptors in

response to physiological and pathological cues.

FXR is increasingly recognized as an important therapeutic target for

enterohepatic diseases, but the development of clinically applicable and more targeted

FXR-based therapy is still challenging. In this study, we provided the first characterization

of an eRNA, *FincoR*, induced by pharmacological activation of FXR and show that *FincoR*

has a beneficial role in reducing liver fibrosis and inflammation in dietary NASH mice.

Complete understanding of the function and mechanisms of *FincoR* may provide novel

insights for the development of desirable therapy for NASH and other chronic liver

diseases.

**Materials and Methods**

**Animal experiments**

All animal studies were performed according to procedures approved by the Institutional

Animal Care & Use Committee at the University of Illinois at Urbana-Champaign and were

in accordance with National Institutes of Health guidelines. Mice were maintained in

12/12 h light/dark cycles and fed standard rodent chow. FXR-LKO mice were generated

by breeding FXR floxed mice with Albumin-Cre mice (The Jackson Lab). FincoR-LKD

mice were generated as previously reported *(Zhao et al., 2018)*. Briefly, Cas9 transgenic

mice (JAX #024858) were via the tail vein injected with adenoviruses (approximately

$5x10^8$ PFU) expressing two sgRNAs targeting FincoR (sgRNA1:

*GGGTTAAGAGCTGTAGGCTG* and sgRNA2: ACTTCTATGTCCAACAACCG). The

sequences of sgRNAs were designed using a CRISPR design tool (http://crispr.mit.edu/).

Mice were given a single dose of vehicle or 30 mg/kg GW4064 (in corn oil, Tocris

Bioscience, #2473) after overnight fasting. Mice were treated with 0.5 mg/kg tropifexor (in

corn oil, MedChem Express, HY-107418), 30 mg/kg cilofexor (in corn oil,

MedChemExpress, HY-109083), 100 mg/kg fexaramine (in 0.5% methylcellulose,

MedChem Express, HY-10912), and 20 mg/kg OCA (in 0.5% methylcellulose, MedChem

Express, HY-12222) as indicated. C57BL6 mice were fed with a chow diet containing

0.5% CA for 6 hrs *(Jung et al., 2020)*.

For liver-specific downregulation of FincoR, mice were fed AMLN diet (Research

Diets, D09100310, 40 kcal% fat, 2% cholesterol, 20 kcal% fructose) for 12 weeks and

then, were injected with adenovirus expressing control sgRNA or sgRNA targeting FincoR

and administration of tropifexor was started 3 days later given at 0.3 mg/kg dissolved in

corn oil.

**RNA-Seq**

C57BL/6 mice were fasted overnight and treated with 30 mg/kg GW4064 or vehicle for

1 h. RNA-seq was performed as previously described *(Byun et al., 2018; Seok et al., 2018;*

*Byun et al., 2020)*. Briefly, the RNAs from livers (n = 2 mice/group, each group contains

RNAs pooled from 2 mice) were prepared (RNeasy, Qiagen). Ribosomal RNA was

removed with the Ribozero HMR Gold kit (Illumina).

**Construction of strand-specific RNAseq libraries**

Construction of the RNAseq libraries and sequencing on the Illumina NovaSeq 6000 were

performed at the Roy J. Carver Biotechnology Center at the University of Illinois at

Urbana-Champaign. After DNase digestion, purified total RNAs were analyzed on a

Fragment Analyzer (Agilent) to evaluate RNA integrity. The total RNAs were converted

into individually barcoded polyadenylated mRNAseq libraries with the Kapa HyperPrep

mRNA kit (Roche). Libraries were barcoded with Unique Dual Indexes (UDI's) which have

been developed to prevent index switching. The adaptor-ligated double-stranded cDNAs

were amplified by PCR for 8 cycles with the Kapa HiFi polymerase (Roche). The final

libraries were quantitated with Qubit (ThermoFisher) and the average cDNA fragment

sizes were determined on a Fragment Analyzer. The libraries were diluted to 10 nM and

further quantitated by qPCR on a CFX Connect Real-Time qPCR system (Bio-Rad) for

accurate pooling of barcoded libraries and maximization of number of clusters in the

flowcell.

**Sequencing of libraries in the NovaSeq**

The barcoded RNAseq libraries were loaded on one SP lane on a NovaSeq 6000 for

cluster formation and sequencing. The libraries were sequenced from one end of the

fragments for a total of 100 bp. The fastq read files were generated and demultiplexed

with the bcl2fastq v2.20 Conversion Software (Illumina, San Diego, CA). The quality of

the demultiplexed fastq files was evaluated with the FastQC software, which generates

reports with the quality scores, base composition, k-mer, GC and N contents, sequence

duplication levels and overrepresented sequences.

**GRO-seq**

To harvest the nuclei from mouse liver cells, the liver was harvested at indicated time and

washed with a cold swelling buffer (10 mM Tris pH 7.5, 2 mM $MgCl_2$, 3 mM $CaCl_2$, 2 U/ml

Superase-In). The nuclei were prepared by Dounce homogenization in cold swelling

buffer and filtered using the cell strainer (100 μm, BD Biosciences). Nuclei were collected

by centrifugation at 400 x g for 10 min, then resuspended in the lysis buffer (swelling

buffer with 10% glycerol and 1% IGEPAL) and incubated on ice for 5 min. Nuclei were

washed twice with the lysis buffer and resuspended at a concentration of $10^8$ nuclei/ml in

the freezing buffer (50 mM Tris pH 8.3, 40% glycerol, 5 mM $MgCl_2$, 0.1 mM EDTA). We

then followed our previous method *(Li et al., 2013; Oh et al., 2021)* to conduct nuclear

run-on and GRO-Seq library preparation. Briefly, the nuclei in freezing buffer were

subjected to the nuclear run-on reaction by mixing with an equal volume of run-on buffer

(10 mM Tris-Cl pH 8.0, 5 mM $MgCl_2$, 1 mM dithiothreitol (DTT), 300 mM KCl, 20 units of

Superase-In, 1% sarkosyl, 500 µM ATP, GTP, Br-UTP and 2 µM CTP) for 5 min at 30 °C.

The BrU-labeled run-on RNAs were extracted by TRIzol and purified by anti-BrdU

agarose beads (Santa Cruz Biotech, sc-32323 AC). The run-on RNAs were then

subjected to end repair by T4 PNK, poly-adenylation, and then to cDNA first strand

synthesis by a custom primer (oNTI223) that allows circularization of the cDNA. This

cDNA then was re-linearized by Ape1 (NEB), size selected by TBE gel and the products

of the desired size were excised (~320–350 bp) for final library prep and sequencing.

GRO-Seq samples were run on a NextSeq 500 sequencer from Illumina with a single-

end 80 nt model.

**Histological analyses**

For histology, tissues were dissected and immediately fixed in 10% formalin overnight

and processed for paraffin embedding and H&E staining. Paraffin-embedded liver

sections were incubated with F4/80 antibody, and antibody was detected using a

peroxidase-based method (Abcam, ab64238). Liver collagen was detected by Sirius Red

staining (Abcam, ab246832) and apoptosis was detected by TUNEL staining (Millipore,

S7100). For Oil Red O staining, liver tissue was frozen in OCT compound (Sakura

Finetek, 4583), sectioned, and stained. Liver sections were imaged with a NanoZoomer

Scanner (Hamamatsu) and quantification was done using NIH ImageJ.

**Metabolic analyses**

Hepatic levels of TG (Sigma, MAK266), cholesterol (Sigma, MAK043), glycogen

(Biovision, K646-100), total BA levels (Diazyme, DZ042A), serum NEFA (Sigma,

MAK044) and serum ALT (Sigma, MAK052) were determined according to the

manufacturer's instructions. Mouse liver IL-1β (R&D systems, MLB00C) and CCL2 (R&D

systems, DY479-05) were detected by commercially available ELISA kit.

**Mouse liver ChIP**

Liver ChIP assay was performed as described previously *(Seok et al., 2018; Byun et al.,*

*2020; Jung et al., 2020)*. Briefly, chromatin extracts were prepared from FXR floxed and

FXR-LKO mouse livers after treatment of the mice with GW4064 which was followed by

preclearing and immunoprecipitation using control IgG or FXR antibody (Novus

Biologicals, NBP2-16550; Santa Cruz, sc-25309), RXRα antibody (Proteintech, catalog

no. 21218-1-AP), or BRD4 antibody (Bethyl Laboratories, catalog #A301-985A50).

Enrichment in chromatin precipitates of gene sequences was measured by qRCR using

primers listed (**Supplementary Table S5**).

**Primary mouse hepatocytes and HepG2 cells**

Primary hepatocytes were isolated from C57BL/6 mice by collagenase (Worthington

Biochemical Corporation, LS004188) perfusion and maintained in William's E Medium

with primary hepatocyte maintenance supplements (Gibco #CM4000) in six-well plates

as described previously *(Kim et al., 2015; Jung et al., 2020)*. HepG2 cells were maintained

in DMEM supplemented with 10% fetal bovine serum at 37°C and 5% $CO_2$.

**Luciferase reporter assay**

The enhancer region containing the FXR binding element was amplified by PCR and

cloned into the pGL4.23 vector (Promega). HepG2 cells were transiently transfected with

FXR (100 ng/well) and RXRα (5 ng/well) in combination with reporters containing the wild

type FXRE or mutated FXRE (200 ng/well) (**Supplementary Table S5**). β-galactosidase

plasmid (200 ng/well) was also transfected as internal controls. Cells were treated with

GW4064 or DMSO after transient transfection. Six hours later, the cells were harvested.

All reporter assays were repeated at least three times in triplicates.

**RACE**

5' and 3' RACE assays were performed using a SMARTer RACE kit (Clontech) according

to the manufacturer's instructions. The resulting PCR products were separated by

electrophoresis in agarose gels and cloned into the pRACE vector provided by the kit.

The transcription start sites and end sites of FincoR were determined by sequencing. The

gene-specific primers used for 5'- and 3'-RACE are provided (**Supplementary Table S5**).

**In vitro transcription and translation (TNT)**

Expression plasmids for luciferase and FincoR were mixed with a Coupled Reticulocyte

Lysate System (Promega). After incubating at 30ºC for 60 min, translated products were

separated on 4-20% gradient SDS polyacrylamide gels and transferred to PVDF

membranes. Chemiluminescent detection of in vitro translated protein was performed

following the manufacturer's protocol (Promega).

**RT-qPCR**

Total RNA was extracted using RNeasy Mini Kit (Qiagen, 74104) and 2 µg of RNA was

reverse transcribed and relative mRNA expression was normalized to those of 36B4. The

qPCR primers are shown in **Supplementary Tables S5**.

**Subcellular fractionation**

Using a Cytoplasmic and Nuclear RNA Purification Kit (Norgen, Thorold, ON, Canada),

the cytoplasm and nucleus fractions from primary hepatocytes isolated from livers of mice

treated with GW4064 or DMSO were separated, and both fractions were subjected to

RNA extraction and qRT-PCR.

**Immunoblotting analysis**

Total liver lysates were prepared as described before *(Sun et al., 2022)*. Antibodies for

immunoblotting for $\beta$-ACTIN (#4970) were purchased from Cell Signaling Technology.

Antibodies for immunoblotting for FXR (sc-25309) were purchased from Santa Cruz.

568
**FincoR polyadenylation study**

Total RNA was prepared using the RNeasy mini prep kit (Qiagen). Poly(A)+ and poly(A)-

RNA was separated by Dynabeads™ mRNA Purification Kit (ThermoFisher Scientific)

according to the manufacturer's instructions. Briefly, total RNA was incubated with the

Dynabeads/binding buffer suspension at room temperature for 5 min and the reaction

tubes were placed on a magnet until solution was clear. The supernatant containing

poly(A)- RNA was saved. The beads with poly(A)+ RNA were washed three times with

washing buffer provided by the kit. RNA was extracted from the supernatant and beads

respectively using Trizol. qPCR was performed to analyze levels of FincoR and 36b4 in

the Poly(A)+ and poly(A)- RNA fractions.

**Genomics Analysis**

RNA-Seq and GRO-Seq reads were mapped to the mouse reference genome mm10 with

STAR aligner. Transcript quantifications were done with HOMER tool set and enhancer

RNAs were identified based on H3K27ac ChIP-Seq in mouse liver. Briefly, intergenic

H3K27ac ChIP-Seq peaks were selected as putative enhancer regions in mouse liver.

Then +/- 3kb regions around putative enhancers with RNA-Seq signal (>1 RPKM) were

considered as putative eRNAs in mouse livers.  Overlapped eRNAs regions were merged,

and redundant ones were removed. In addition, any +/- 3kb extended eRNAs regions that

overlapped with protein-coding genes were further removed to avoid transcriptional

readthrough from genes. We used DESeq2 to identify significantly regulated eRNAs with

a cutoff of (FDR < 0.05, Log2FC > 1).  Public ChIP-Seq datasets were obtained from

ENCODE or GEO (**Supplementary Table S3**).

**Statistical analysis**

Statistical analysis was performed using GraphPad Prism 9. Statistical differences were

evaluated using two-tailed unpaired Student's t-test for comparisons between two groups,

or ANOVA and appropriate post hoc analyses for comparisons of more than two groups.

Statistical methods and corresponding $p$ values for data shown in each panel were

indicated in figure legends.

**Data availability**

RNA-seq and Gro-seq data were deposited in GEO under the following accession numbers GSE 221986.

**Acknowledgments**

We thank Deepa Prakashini Govindasamy Rajagopal for the help with tissue genotyping. We also thank Kristina Schoonjans at Ecole Polytech, Switzerland, for providing FXR floxed mice. This study was supported by an American Diabetes Association Postdoctoral Fellowship to J.C. (1-19-PDF-117), and by a John and Rebekah Harper Fellowship to R.W. W.L. is a Cancer Prevention and Research Institute of Texas (CPRIT) Scholar. This work is supported by funding from NCI (K22CA204468), NIGMS (R21GM132778, R01GM136922), Welch Foundation (AU-2000-20220331), CPRIT (RR160083) to W.L., and grants from the NIH (R01 DK062777 and R01 DK095842) to J.K.K.

**References**

Abel U, Schluter T, Schulz A, et al. 2010. Synthesis and pharmacological validation of a

novel series of non-steroidal FXR agonists. *Bioorg Med Chem Lett* **20**: 4911-

4917.

Abenavoli L, Falalyeyeva T, Boccuto L, et al. 2018. Obeticholic Acid: A New Era in the

Treatment of Nonalcoholic Fatty Liver Disease. *Pharmaceuticals (Basel)* **11**.

Ali AH, Carey EJ, Lindor KD. 2015. Recent advances in the development of farnesoid X

receptor agonists. *Ann Transl Med* **3**: 5.

Brocker CN, Kim D, Melia T, et al. 2020. Long non-coding RNA Gm15441 attenuates

hepatic inflammasome activation in response to PPARA agonism and fasting.

*Nat Commun* **11**: 5847.

Byun S, Kim DH, Ryerson D, et al. 2018. Postprandial FGF19-induced phosphorylation

by Src is critical for FXR function in bile acid homeostasis. *Nat Commun* **9**: 2590.

Byun S, Seok S, Kim YC, et al. 2020. Fasting-induced FGF21 signaling activates

hepatic autophagy and lipid degradation via JMJD3 histone demethylase. *Nat*

*Commun* **11**: 807.

Calkin AC, Tontonoz P. 2012. Transcriptional integration of metabolism by the nuclear

sterol-activated receptors LXR and FXR. *Nat Rev Mol Cell Biol* **13**: 213-224.

Cech TR, Steitz JA. 2014. The noncoding RNA revolution-trashing old rules to forge

new ones. *Cell* **157**: 77-94.

Chen J, Wang Z, Hu X, et al. 2016. BET Inhibition Attenuates Helicobacter pylori-

Induced Inflammatory Response by Suppressing Inflammatory Gene

Transcription and Enhancer Activation. *J Immunol* **196**: 4132-4142.

Claudel T, Staels B, Kuipers F. 2005. The Farnesoid X receptor: a molecular link

between bile acid and lipid and glucose metabolism. *Arterioscler Thromb Vasc*

*Biol* **25**: 2020-2030.

Consortium EP. 2012. An integrated encyclopedia of DNA elements in the human

genome. *Nature* **489**: 57-74.

Downes M, Verdecia MA, Roecker AJ, et al. 2003. A chemical, genetic, and structural

analysis of the nuclear bile acid receptor FXR. *Mol Cell* **11**: 1079-1092.

Evans RM, Mangelsdorf DJ. 2014. Nuclear Receptors, RXR, and the Big Bang. *Cell*

**157**: 255-266.

Fang B, Everett LJ, Jager J, et al. 2014. Circadian enhancers coordinate multiple

phases of rhythmic gene transcription in vivo. *Cell* **159**: 1140-1152.

Fang S, Suh JM, Reilly SM, et al. 2015. Intestinal FXR agonism promotes adipose

tissue browning and reduces obesity and insulin resistance. *Nat Med* **21**: 159-

165.

Friedman SL, Neuschwander-Tetri BA, Rinella M, et al. 2018. Mechanisms of NAFLD

development and therapeutic strategies. *Nat Med* **24**: 908-922.

Goodwin B, Jones SA, Price RR, et al. 2000. A regulatory cascade of the nuclear

receptors FXR, SHP-1, and LRH-1 represses bile acid biosynthesis. *Mol Cell* **6**:

517-526.

Hernandez ED, Zheng L, Kim Y, et al. 2019. Tropifexor-Mediated Abrogation of

Steatohepatitis and Fibrosis Is Associated With the Antioxidative Gene

Expression Profile in Rodents. *Hepatol Commun* **3**: 1085-1097.

Hirabayashi S, Bhagat S, Matsuki Y, et al. 2019. NET-CAGE characterizes the

dynamics and topology of human transcribed cis-regulatory elements. *Nat Genet*

**51**: 1369-1379.

Hon CC, Ramilowski JA, Harshbarger J, et al. 2017. An atlas of human long non-coding

RNAs with accurate 5' ends. *Nature* **543**: 199-204.

Hsieh CL, Fei T, Chen Y, et al. 2014. Enhancer RNAs participate in androgen receptor-

driven looping that selectively enhances gene activation. *Proc Natl Acad Sci U S*

*A* **111**: 7319-7324.

Jung H, Chen J, Hu X, et al. 2020. BRD4 inhibition and FXR activation, individually

beneficial in cholestasis, are antagonistic in combination. *JCI Insight* **6**.

Kim YC, Fang S, Byun S, et al. 2015. Farnesoid X receptor-induced lysine-specific

histone demethylase reduces hepatic bile acid levels and protects the liver

against bile acid toxicity. *Hepatology* **62**: 220-231.

Kliewer SA, Mangelsdorf DJ. 2015. Bile Acids as Hormones: The FXR-FGF15/19

Pathway. *Dig Dis* **33**: 327-331.

Kremoser C. 2021. FXR agonists for NASH: How are they different and what difference

do they make? *J Hepatol* **75**: 12-15.

Lai F, Shiekhattar R. 2014. Enhancer RNAs: the new molecules of transcription. *Curr*

*Opin Genet Dev* **25**: 38-42.

Lam MT, Li W, Rosenfeld MG, et al. 2014. Enhancer RNAs and regulated

transcriptional programs. *Trends Biochem Sci* **39**: 170-182.

Lee FY, Lee H, Hubbert ML, et al. 2006. FXR, a multipurpose nuclear receptor. *Trends*

*Biochem Sci* **31**: 572-580.

Lee J, Padhye A, Sharma A, et al. 2010. A pathway involving farnesoid X receptor and

small heterodimer partner positively regulates hepatic sirtuin 1 levels via

microRNA-34a inhibition. *J Biol Chem* **285**: 12604-12611.

Lee J, Seok S, Yu P, et al. 2012. Genomic analysis of hepatic farnesoid X receptor

binding sites reveals altered binding in obesity and direct gene repression by

farnesoid X receptor in mice. *Hepatology* **56**: 108-117.

Lee JM, Wagner M, Xiao R, et al. 2014. Nutrient-sensing nuclear receptors coordinate

autophagy. *Nature* **516**: 112-115.

Li W, Notani D, Ma Q, et al. 2013. Functional roles of enhancer RNAs for oestrogen-

dependent transcriptional activation. *Nature* **498**: 516-520.

Li W, Notani D, Rosenfeld MG. 2016. Enhancers as non-coding RNA transcription units:

recent insights and future perspectives. *Nat Rev Genet* **17**: 207-223.

Li Y, Tan Z, Zhang Y, et al. 2021. A noncoding RNA modulator potentiates

phenylalanine metabolism in mice. *Science* **373**: 662-673.

Mattick JS, Amaral PP, Carninci P, et al. 2023. Long non-coding RNAs: definitions,

functions, challenges and recommendations. *Nat Rev Mol Cell Biol* **24**: 430-447.

Mirtschink P, Bischof C, Pham MD, et al. 2019. Inhibition of the Hypoxia-Inducible

Factor 1alpha-Induced Cardiospecific HERNA1 Enhance-Templated RNA

Protects From Heart Disease. *Circulation* **139**: 2778-2792.

Oh S, Shao J, Mitra J, et al. 2021. Enhancer release and retargeting activates disease-

susceptibility genes. *Nature* **595**: 735-740.

Rahnamoun H, Lee J, Sun Z, et al. 2018. RNAs interact with BRD4 to promote

enhanced chromatin engagement and transcription activation. *Nat Struct Mol Biol*

**25**: 687-697.

Sallam T, Jones MC, Gilliland T, et al. 2016. Feedback modulation of cholesterol

metabolism by the lipid-responsive non-coding RNA LeXis. *Nature* **534**: 124-128.

Sanyal AJ, Lopez P, Lawitz EJ, et al. 2023. Tropifexor for nonalcoholic steatohepatitis:

an adaptive, randomized, placebo-controlled phase 2a/b trial. *Nat Med* **29**: 392-

400.

Sartorelli V, Lauberth SM. 2020. Enhancer RNAs are an important regulatory layer of

the epigenome. *Nat Struct Mol Biol* **27**: 521-528.

Seok S, Fu T, Choi SE, et al. 2014. Transcriptional regulation of autophagy by an FXR-

CREB axis. *Nature* **516**: 108-111.

Seok S, Kim YC, Byun S, et al. 2018. Fasting-induced JMJD3 histone demethylase

epigenetically activates mitochondrial fatty acid beta-oxidation. *J Clin Invest* **128**:

3144-3159.

Seok S, Sun H, Kim YC, et al. 2021. Defective FXR-SHP Regulation in Obesity

Aberrantly Increases miR-802 Expression, Promoting Insulin Resistance and

Fatty Liver. *Diabetes* **70**: 733-744.

Statello L, Guo CJ, Chen LL, et al. 2021. Gene regulation by long non-coding RNAs and

its biological functions. *Nat Rev Mol Cell Biol* **22**: 96-118.

Sun H, Seok S, Jung H, et al. 2022. Obesity-induced miR-802 directly targets AMPK

and promotes nonalcoholic steatohepatitis in mice. *Mol Metab* **66**: 101603.

Tang X, Luo Y, Yuan D, et al. 2023. Long noncoding RNA LEENE promotes

angiogenesis and ischemic recovery in diabetes models. *J Clin Invest* **133**.

Thomas AM, Hart SN, Kong B, et al. 2010. Genome-wide tissue-specific farnesoid X

receptor binding in mouse liver and intestine. *Hepatology* **51**: 1410-1419.

Tully DC, Rucker PV, Chianelli D, et al. 2017. Discovery of Tropifexor (LJN452), a

Highly Potent Non-bile Acid FXR Agonist for the Treatment of Cholestatic Liver

Diseases and Nonalcoholic Steatohepatitis (NASH). *J Med Chem* **60**: 9960-9973.

Wahlstrom A, Sayin SI, Marschall HU, et al. 2016. Intestinal Crosstalk between Bile

Acids and Microbiota and Its Impact on Host Metabolism. *Cell Metab* **24**: 41-50.

Wang YD, Chen WD, Wang M, et al. 2008. Farnesoid X receptor antagonizes nuclear

factor kappaB in hepatic inflammatory response. *Hepatology* **48**: 1632-1643.

Warren CFA, Wong-Brown MW, Bowden NA. 2019. BCL-2 family isoforms in apoptosis

and cancer. *Cell Death Dis* **10**: 177.

Zhao XY, Xiong X, Liu T, et al. 2018. Long noncoding RNA licensing of obesity-linked

hepatic lipogenesis and NAFLD pathogenesis. *Nat Commun* **9**: 2986.

Zheng W, Lu Y, Tian S, et al. 2018. Structural insights into the heterodimeric complex of

the nuclear receptors FXR and RXR. *J Biol Chem* **293**: 12535-12541.

743
744

**Figure 1. Activation of FXR by GW4064 induces *FincoR*, a novel eRNA, in mouse**

**liver.** (A) C57BL/6 male mice were fasted overnight and *i.p.* injected with vehicle or

GW4064 (30 mg/kg) for 1 h, and livers were collected (n = 4 mice). RNAs from two mice

(per group) were pooled for RNA-seq. Volcano plot showing significantly induced eRNAs

(FincoR highlighed) by GW4064 treatment. The *x* axis denotes $\log_2$ fold change

(GW4064/Veh) of eRNAs and the *y* axis denotes $-\log_{10}$ FDR of eRNAs. (B) The bar plot

showing the RPMs of up-regulated eRNAs by GW4064 treatment. (C) FincoR induction

by GW4064 treatment was shown in the genome browser track. (D) GW4064 induction

of *FincoR* and *Shp* was measured in the liver by qPCR (n = 4 mice). Data are presented

as mean ± SEM. Statistical significance was determined by the two-way ANOVA Sidak's

multiple comparisons test with *$p < 0.05$ and ***$p < 0.001$. Veh, vehicle; GW, GW4064;

pos, positive strand; neg, negative strand; Refseq, reference sequence.

**Figure 2. Ligand-activated FXR directly activates transcription of eRNAs including**

***FincoR* in the liver.** (A) Experimental outline: FXR-Flox and LKO male mice were fasted

overnight and treated with vehicle or GW4064, livers were collected 1 h later (n = 2 each)

and nuclei was isolated for Gro-seq. Box plot shows the up-regulated eRNA GRO-Seq

RPKMs in different conditions. (B) Gro-seq profile of *FincoR* expression and genome

browser tracks of H3K4me1, H3K27ac, FXR and RXRα binding peaks at *FincoR* locus

were shown. (C) ChIP assays were performed in liver samples from Fig. 1A to detect

FXR, RXRα, and BRD4 occupancy to the FXR binding peak region close to the

transcription start site of *FincoR*. (D) HepG2 cells were transfected with luciferase reporter

expressing wild type FXRE or mutant FXRE (see materials and methods) for 24 hrs before

further subject to GW4064 for 6 hrs. The relative luciferase activity was shown.

**Figure 3. *FincoR* is a liver-specific nucleus-enriched eRNA.** (A) C57BL/6 male mice

were fasted overnight and *i.p.* injected with vehicle or GW4064 (30 mg/kg) for 1 h. Tissues

were collected and *FincoR* mRNA level was measured (n = 2 mice). (B) C57BL/6 male

mice were fasted overnight and *i.p.* injected with vehicle or GW4064 (30 mg/kg) for 1 h.

One small piece of liver was snap-frozen for later RNA isolation and the remaining part

was used for immediate primary hepatocyte isolation. Then, RNAs were extracted from

liver or primary hepatocytes and *FincoR* expression was measured. (C) Agarose gel

electrophoresis of PCR products generated in 5′ (left) and 3′ (right) RACE of *FincoR* in

liver samples. Primer locations were show. (D) Schematic diagrams of full-length *FincoR*.

(E) PhyloCSF analysis of the coding potential of *FincoR*. (F) In vitro translation of *FincoR*

using the Promega Transcend Non-Radioactive Translation Detection Systems.

Luciferase is used as controls for coding RNA. (G) qPCR analysis of *FincoR*, *36b4* in

Poly(A)+ and Poly(A)- RNA fractions from GW4064 treated mouse liver. (H) *FincoR*

identified in the subcellular fraction using cellular fractionation assays. The primary

hepatocytes were isolated from GW4064 or DMSO treated mouse liver and the cytoplasm

and nucleus fractions of these hepatocytes were separated and both fractions were

subjected to RNA extraction and qPCR. RACE, rapid amplification of cDNA ends.

**Figure 4. FincoR is induced by the hammerhead class of non-steroidal FXR**

**agonists, including GW4064 and tropifexor.** (A) The chemical structures of the FXR

agonists including the hammerhead class of non-synthetic FXR agonists, non-

hammerhead-type non-synthetic agonists, semi-synthetic BA and natural BA. (B)

C57BL/6 mice were treated with GW4064 (30 mg/kg) or Cilofexor (30 mg/kg) or Tropifexor

(0.5 mg/kg) or Fexaramine (100 mg/kg) or OCA (20 mg/kg) for 1 h after over-night fasting.

The liver RNAs were extracted and FincoR expression was measured (n = 3~4 mice). (C)

C57BL/6 mice were treated with OCA (20 mg/kg) or Fexaramine (100 mg/kg) for 4 hrs

after over-night fasting. The liver RNAs were extracted and FincoR expression was

measured (n = 3 mice). (D) C57BL/6 mice were daily treated with OCA (20 mg/kg) for 7

days after over-night fasting. The liver RNAs were extracted and FincoR expression was

measured (n = 5 mice). *Shp* gene was used as a positive control. (E) C57BL/6 mice were

fed with 0.5% cholic acid (CA) diet for 6 hrs after over-night fasting. The liver RNAs were

extracted and FincoR expression was measured (n = 3~4 mice). (B-E) Data are presented

as mean ± SEM. Statistical significance was determined by the Student's t test with *$p <$

$0.05$, ***$p < 0.01$ and ****$p < 0.001$.

**Figure 5. Generation of CRISPR/Cas9-mediated *FincoR* liver-specific knockdown**

**mice.** (A) Male Cas9 mice were infected with adenovirus expressing sgRNA for *FincoR*

or control for 1 week. Then the liver and serum were collected from these mice after 4~5

hrs of fasting. (B) The expression of FincoR in the liver was measured by qPCR (n = 4

mice). (C) Hepatic triglyceride, cholesterol, bile acid, glycogen and serum NEFA were

measured (n = 5~7 mice) (D) Male Cas9 mice were infected with adenovirus expressing

sgRNA for *FincoR* or control for 1 week. Then these mice were fasted overnight and

treated with GW4064 for 3 hrs before tissue collection. RNA-seq profiles of expression of

hepatic *FincoR* and the adjacent genes were shown (n = 2 mice). (E) Genome-wide

changes in mRNA expression shown in a volcano plot. The numbers refer to the number

812 of genes up- or down-regulated by 2-fold or more with a *p*-value < 0.01. (F) Gene ontology

813 analysis of biological pathways using DAVID Tools.

814 **Figure 6. In diet-induced NASH mice, tropifexor-mediated beneficial effects on**

815 **reducing hepatic steatosis are largely independent on FincoR.** (A-E) Male Cas9 mice

816 were fed with NASH diet for 12 weeks. Then these mice were randomly assigned to 3

817 groups and infected with adenovirus expressing sgRNA for *FincoR* or control. Three days

818 later, the mice were treated with Tropifexor (0.3 mg/kg) for 12 days. On the tissue

819 collection day, the mice were given the last shot of Tropifexor or vehicle and fasted for 4

820 hrs before tissues were collected (n = 6 ~ 7 mice). (A) Experimental scheme. (B) Hepatic

821 FincoR expression was measured (n = 6~7 mice). (C) Oil Red O staining of liver sections.

822 Scale bar (50 μm). (D) Hepatic TG, hepatic cholesterol, gallbladder BA and hepatic BA

823 levels were measured (n = 6 ~ 7 mice). (E) mRNA levels in the liver of the indicated genes

824 involved in bile acid regulation and lipid regulation (n = 6 ~ 7 mice). (B, D and E) Data are

825 presented as mean ± SEM. Statistical significance was determined by the one-way

826 ANOVA (Sidak's multiple comparisons test) with \**p* < 0.05, \*\**p* < 0.01 and \*\*\**p* < 0.001.

827 Ad, adenovirus; H & E, hematoxylin and eosin; TG, triglyceride; Veh, vehicle; ns, not

828 significant.

829 **Figure 7. In diet-induced NASH mice, tropifexor-mediated beneficial effects on**

830 **reducing liver fibrosis and inflammation are diminished by FincoR downregulation.**

831 (A) H & E, F4/80, Sirius Red and TUNEL staining of liver sections from Figure 6. Scale

832 bar (50 μm). Image analyses were done using Image J and the area of collagen staining,

833 TUNEL and F4/80 levels were quantified (n = 5/group). (B) Serum ALT level was

834 measured (n = 5/group). (C) IL-1β and CCL2 levels in the liver tissues were determined

by ELISA (n = 5/group). (D) mRNA levels in the liver of the indicated genes involved in

inflammation, fibrosis and cell death (n = 5/group). (E) Model: FincoR is a liver-specific

lncRNA that is induced specifically by hammerhead-type FXR agonists (top). In diet-

induced NASH mice, *FincoR* is required for tropifexor-mediated beneficial effects on

reducing hepatic inflammation, fibrosis, and cell death with the mechanisms to be

explored further (bottom). (A-D) Data are presented as mean ± SEM. Statistical

significance was determined by the one-way ANOVA (Sidak's multiple comparisons test)

with *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$.

**Supplementary Figure Legends**

**Figure 1- Figure Supplement 1.** (A, B) Examples of FXR-regulated eRNAs produced near the genes *Hes1* (A) and *Slc35g1* (B) were shown. (C) Time course expression of *FincoR*. C57BL/6 male mice were fasted overnight and *i.p.* injected with vehicle or GW4064 (30 mg/kg) for 1, 3, 6,12 h. Livers were collected at indicated time points (n = 3 mice) and *FincoR*, *Shp* and *Gcnt1* were measured. Data are presented as mean ± SEM. Statistical significance was determined by the two-way ANOVA Sidak's multiple comparisons test with *$p < 0.05$ and ***$p < 0.001$.

**Figure 1- Figure Supplement 2.** (A) Volcano plot showing the differential expressed genes (DEGs) by GW4064 treatment (DESeq2 FDR<0.05). The numbers refer to the number of genes up- or down-regulated. (B) Bar plot showing the enriched Gene ontology in terms of biological processes for up-regulated genes.

**Figure 2- Figure Supplement 1**. (A) FXR protein levels in the livers isolated from FXR-Flox and FXR-LKO mice were shown. (B) Validation of hepatic FXR-dependent induction of *FincoR* by qPCR (n = 3 mice). *Shp* gene was used as a control. Data are presented as mean ± SEM. Statistical significance was determined by the two-way ANOVA Tukey's multiple comparisons test with *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$. (C) Metagene plots showing H3K27ac, H3K4me1, FXR, and RXRα ChIP-Seq profiles centered on up-regulated eRNAs.

**Figure 5- Figure Supplement 1**. (A) The scheme for the *FincoR* loss of function experiments. (B) The genomic DNAs from liver, spleen, intestine, brain, heart, muscle, kidney, lung and adipose tissue were isolated and PCR was performed using the primers (**Supplementary Table S5E**) to verify tissue-specific knock-out. (C-G) RNA-seq profiles

of expression of hepatic *Prune2* (C)*, PPP1r3g* (D)*, Igfbp2* (E)*, Eda2r* (F) *and Fndc1* (G)

were shown (n = 2 mice).

**Figure 7- Figure Supplement 1. The effects of FincoR downregulation on NASH**

**pathologies.** Male Cas9 mice were fed with NASH diet for 12 weeks. Then these mice

were randomly assigned to 2 groups and infected with adenovirus expressing sgRNA for

*FincoR* or control. The tissues were collected 2 weeks later. Liver histology analysis was

performed. Scale bar (50 µm).

**Other Supplementary Figure 1. Hepatic genome browser tracks of FXR, RXRα, LXR,**

**PPARα, and HNF4α binding peaks at *FincoR* locus.**

**Other Supplementary Figure 2. PPARα occupnacy in the FincoR enhancer region.**

C57BL/6 male mice were fasted overnight with or without refed for 3 hrs and then

sacrificed. ChIP assays were performed in liver samples to detect PPARα occupancy to

the FXR binding peak region close to the transcription start site of *FincoR*.

## Supplemental Tables

Supplemental Table S1. Sequencing data generated in this study

| Name | Total reads | Uniquely mapped reads |
| --- | --- | --- |
| WT Liver Veh RNA-Seq rep1 | 50887391 | 38072659 |
| WT Liver GW RNA-Seq rep1 | 54020150 | 40595856 |
| WT Liver Veh RNA-Seq rep2 | 51268164 | 38898913 |
| WT Liver GW RNA-Seq rep2 | 51899892 | 39487241 |
| Flox Liver Veh GRO-Seq | 62960996 | 30305910 |
| Flox Liver GW GRO-Seq | 59210073 | 26116795 |
| FXR KO Liver Veh GRO-Seq | 34153537 | 14379417 |
| FXR KO Liver GW GRO-Seq | 36430101 | 15008281 |
| WT Liver RNA-Seq rep1 | 96262381 | 81801494 |
| WT Liver RNA-Seq rep2 | 104979915 | 88365481 |
| FincoR KD Liver RNA-Seq rep1 | 91654076 | 75346167 |
| FincoR KD Liver RNA-Seq rep2 | 90211420 | 74745807 |

Supplemental Table S2 is attached as an Excel file.

Supplemental Table S3. Public ChIP-Seq dataset

| target | accession ID |
| --- | --- |
| FXR | https://genome.ucsc.edu/cgi-bin/hgCustom?hgsid=1577671033_RPWVA7jbx3rDswKOeo2dazTsXxe0&hgct_table=ct_FxrLiver_2971 |
| H3K27ac | ENCFF001KMI |
| H3K4me1 | ENCFF001KNF |
| LXR | GSM864669 |
| RXR$\alpha$ | GSM864674 |
| PPAR$\alpha$ | GSM864671 |
| HNF4$\alpha$ | GSM2055887 |

Supplemental Table S4 is attached as an Excel file.

Supplemental Table S5. Primer sequences used in this study

A. List of primer sequences for mouse qRT-PCR

| Gene | Forward Primer (5'-3') | Reverse Primer (5'-3') |
| --- | --- | --- |
| FincoR | GCAAAGCACCTTCTAGCACA | GTCAGGGAGCTAACGAATGC |
| Gcnt1 | GCATCGCATCCTGCTTTGATA | GGTCTGCCTTAACCCGACTC |
| Shp | TCTGCAGGTCGTCCGACTAT | CAGGCAGTGGCTGTGAGAT |
| Cyp7a1 | GGGATTGCTGTGGTAGTGAGC | GGTATGGAATCAACCCGTTGTC |

| | | |
|---|---|---|
| *Cyp8b1* | CCTCTGGACAAGGGTTTTGTG | GCACCGTGAAGACATCCCC |
| *Srebp1c* | GCAGCCACCATCTAGCCTG | CAGCAGTGAGTCTGCCTTGAT |
| *Lpin* | CATGCTTCGGAAAGTCCTTCA | GGTTATTCTTTGGCGTCAACCT |
| *Scd1* | TTCTTGCGATACACTCTGGTGC | CGGGATTGAATGTTCTTGTCGT |
| *Ifng* | GCG TCA TTG AAT CAC ACC TG | GAC CTG TGG GTT GTT GAC CT |
| *Ccl3* | TTCTCTGTACCATGACACTCTGC | CGTGGAATCTTCCGGCTGTAG |
| *Ccl2* | TTAAAAACCTGGATCGGAACCAA | GCATTAGCTTCAGATTTACGGGT |
| *Ccr2* | AAGAGGGCATTGGATTCACCACA | GCCGTGGATGAACTGAGGTAACA |
| *Lcn2* | GCAGGTGGTACGTTGTGGG | CTCTTGTAGCTCATAGATGGTGC |
| *Col1a1* | ATCGGTCATGCTCTCTCCAAACCA | ACTGCAACATGGAGACAGGTCAGA |
| *Col1a2* | CCTTTGTCAGAATACTGAGCAGC | GTAACTTCGTGCCTAGCAACA |
| *Acta2* | TCGGATACTTCAGCGTCAGGA | GTCCCAGACATCAGGGAGTAA |
| *Eda2r* | CACACTGCATAGTCTGCCCTC | GCCTTCTGGACCCGATTGA |
| *Fndc1* | GGGAGACATGGCAAACCTGT | TGGTAGGAGAGTATGTGGTGG |
| *Ctsb* | TCCTTGATCCTTCTTTCTTGCC | ACAGTGCCACACAGCTTCTTC |
| *Ctss* | GAAGTACGGCGTCTCATCTGG | CATGCCCACTTGGTAGGTATG |
| *Bcl2* | GTCGCTACCGTCGTGACTTC | CAGACATGCACCTACCCAGC |
| *36b4* | CCCTGAAGTGCTCGACATCA | TGCGGACACCCTCCAGAA |

## B. List of primers used for mutagenesis

| | Forward Primer (5'-3') | Reverse Primer (5'-3') |
|---|---|---|
| Mutation | GTGAATTCTCTCCTAGaTCcTTGACaTTCCAAGTCATGTAC | GTACATGACTTGGAAtGTCAAgGAtCTAGGAGAGAATTCAC |

901
902

## C. List of primer sequences for mouse ChIP-qPCR

| Gene | Forward Primer (5'-3') | Reverse Primer (5'-3') |
|---|---|---|
| FincoR | TCCTAGGTCTTTGACCTTCCAA | TGTGCCCACTGGTGAAATAG |
| Non-specific region | GTGTACACGCCCAAACTTGA | GTTCACCTCTCTAGCTCACCT |

904
905

## D. List of primer used for RACE

| | Gene specific primer (5'-3') |
|---|---|
| 5'RACE | GATTACGCCAAGCTT GGGAACTCACGTCTGGAAGCAGAAGCTG |
| 3'RACE | GATTACGCCAAGCTT GGACACTCGGGAATTGACAGTGGGTCAGG |
| 3'RACE nest primer | GATTACGCCAAGCTT GTGGGGGATAGCACAAGCGCCAGGATGC |

907 <u>E. List of primer used for genotyping</u>

| Forward Primer (5'-3') | Reverse Primer (5'-3') |
|---|---|
| GTGATTTCCTTTCCCACAGC | GTCAGGGAGCTAACGAATGC |

# Figure 1

**A** RNA-seq (GW4064: 1 h)

**B** *Up regulated eRNAs*

**C** chr19:17,131,719-17,526,635

chr19:17,300,000-17,340,000

**D** *FincoR* *Shp*

# Figure 2

# Figure 3

# Figure 4

**A**

GW4064     Cilofexor     Tropifexor (LJN452)

Hammerhead-type synthetic FXR agonists

Fexaramine

non-hammerhead-type synthetic FXR agonist

Obeticholic acid (OCA)

semi-synthetic BA

Cholic acid

natural BA

**B** *FincoR*

Rel RNA level

- Veh
- GW4064
- Cilofexor
- Tropifexor
- OCA
- Fex

**C** *FincoR*

Rel RNA level

- Veh
- OCA
- Fex

**D** *Shp*     *FincoR*

Rel RNA level

- Veh
- OCA

**E** *FincoR*

Rel RNA level

- Normal chow
- 0.5% CA diet

# Figure 5

**A**

Cas9 → Ad-sgRNA Ctrl or Ad-sgRNA FincoR (1 week) → FAST 4~5 hrs → Sac

**B**

*FincoR*

□ Ad-sgRNA Ctrl
▨ Ad-sgRNA FincoR

Rel RNA level

***

**C**

Triglyceride (mg/g liver)

Cholesterol (mg/g liver)

Liver BA (nmol/g liver)

Glycogen (µg/mg liver)

Serum NEFA (µM)

**D**

Cas9 → Ad-sgRNA Ctrl or Ad-sgRNA FincoR (1 week) → Fast O/N → GW4064 → 3 hrs → liver for RNA-seq

MM10  chr19:17,158,000-17,483,300

325 Kb

17,200 kb    17,300 kb    17,400 kb

*FincoR*

Ctrl 1 (pos)
Ctrl 2 (pos)
LKD 1 (pos)
LKD 2 (pos)
Ctrl 1 (neg)
Ctrl 2 (neg)
LKD 1 (neg)
LKD 2 (neg)

Refseq

*Prune2*    *Gcnt1*    *Rfk*    *Pcsk5*

**E**

Down:53  Up:18

-log$_{10}$ FDR

log$_2$ Fold Change

**F**

## GO Biological Process

positive regulation of fatty acid oxidation
positive regulation of organelle organization
tryptophan metabolic process
tryptophan catabolic process
indolalkylamine catabolic process
positive regulation of glucose metabolic process
regulation of fatty acid oxidation
positive regulation of fatty acid metabolic process
2-oxoglutarate metabolic process
regulation of insulin-like growth factor receptor signaling pathway

-log$_{10}$ (Adjusted P-value)

# Figure 6

**A** Cas9 → NASH diet 12 weeks → 1: Ad-sgRNA Ctrl / 2: Ad sgRNA-Ctrl / 3: Ad sgRNA-FincoR → 3 days → D1 Tropifexor (0.3 mg/Kg)/Veh → D12 Tropifexor Fast 4 hrs → Sac

**B** *FincoR*

**C**
Ad-sgRNA: Ctrl / Ctrl / FincoR
Tropifexor: - / + / +
Oil Red O

**D** Hepatic TG / Hepatic cholesterol / Hepatic BA / Gallbladder BA

**E** Bile acid regulation — *Shp*, *Cyp7a1*, *Cyp8b1* ; Lipid regulation — *Srebp1c*, *Lpin*, *Scd1*

Legend: Ad-sgRNA Ctrl + Veh / Ad-sgRNA Ctrl + Tropifexor / Ad-sgRNA FincoR + Tropifexor

# Figure 7

# Figure 1-Figure Supplement 1

# Figure 1–Figure Supplement 2

# Figure 2-Figure Supplement 1

# Figure 5-Figure Supplement 1

# Figure 7-Figure Supplement 1

# Other Figure Supplement 1

# Other Figure Supplement 2

**C57BL/6**