The staphylococcal type VII secretion system impacts daptomycin sensitivity through

controlling bacterial cell envelope integrity

Victoria Smith[1], Giridhar Chandrasekharan[1], Kate E. Watkins[1], Ramon Garcia Marset[1$], Jeannifer

Yap[2], Sebastien Perrier[3], Arnaud Kengmo Tchoupa[1+], and Meera Unnikrishnan[1*].

1. Division of Biomedical Sciences, Warwick Medical School, University of Warwick,

Coventry, CV4 7AL, UK.

2. Department of Chemistry, University of Warwick, Coventry, CV4 7AL, UK

3. Department of Chemistry and Warwick Medical School, University of Warwick, Coventry,

CV4 7AL, UK.

$ current address: Centre for Urological Biology, Division of Medicine, University College

London, WC1E 6BT London, UK.

+ current address: Cluster of Excellence EXC 2124 Controlling Microbes to Fight Infections,

University of Tübingen, 72076 Tübingen, Germany

*Corresponding author m.unnikrishnan@warwick.ac.uk

Key words: Type VII secretion system, membrane integrity, membrane fluidity, daptomycin,

Summary

The *Staphylococcus aureus* type VII secretion system (T7SS) plays an important role in bacterial virulence during infection. However, T7SS functions during infection and in bacterial physiology are poorly understood. Here, we demonstrate that *S. aureus* strains lacking the T7SS transporter EssC or effectors, EsxC and EsxA, were highly sensitive to the important last resort drug, daptomycin. T7SS mutants displayed decreased membrane fluidity, altered cell membrane protein localisation, distinct cell surface morphologies, with effects on cell wall synthesis as well as surface charge under normal growth conditions, indicating that the T7SS is critical to cell envelope function. Daptomycin binding was enhanced in the mutants, whose membranes were more permeable when treated with the drug. Furthermore, the lack of EsxC enhanced daptomycin sensitivity during intracellular infection and during murine skin infection. Thus, our data show that the staphylococcal T7SS impacts sensitivity to membrane-acting drugs through modulating membrane integrity, indicating its potential as a drug target.

Keywords: Type VII secretion system, cell membrane integrity, membrane fluidity, daptomycin, intracellular infection

Introduction

S. aureus is a highly versatile pathogen and a commensal bacterium which causes infections in both humans and animals. Staphylococcal infections can be nosocomial or community-acquired, and can range from mild skin abscesses to severe life-threatening infections including septic shock and bacteraemia[1]. *S. aureus* infections place a huge cost burden on healthcare sectors across the world, especially due to the occurrence of antibiotic-resistant strains, such as methicillin-resistant *S. aureus* (MRSA) and strains with varying levels of vancomycin resistance[2,3]

S. aureus possesses an array of surface and secreted factors that help the bacterium colonise the host and establish an infection[4]. One such virulence-associated factor is the type VII secretion system (T7SS), which is widely found in Gram positive bacteria[5]. The T7SSb in *S. aureus* and other firmicutes is distinct from the systems in mycobacteria (T7SSa) although both are characterised by WXG containing small substrates such as EsxA and EsxB[6,7]. The T7SS in *S. aureus* is modular in nature, with heterogenous expression between strains[8,9]. Typically, in commonly studied *S. aureus* strains including USA300, Newman and RN6390, the T7SS locus comprises integral membrane proteins (EsaA, EssA, EssB and EssC), cytosolic proteins (EsaB and EsaG), several secreted substrates (EsxA, EsxB, EsxC, EsxD and EsaD) and chaperones (EsaE)[7,10]. EssC is the central transporter which is responsible for export of substrates and co-dependent export of secreted substrates has been demonstrated[6,11]. While specific interactions between the different T7SS components have been described, the overall structure of the secretion apparatus is poorly defined for the *S. aureus* T7SS[10,11]. Assembly of a functional system is dependent on the membrane microdomain protein, flotillin A (FloA)[12].

Recent studies have reported a role for T7SS in modulation of intraspecies competition. A T7SS substrate EsaD with nuclease activity mediates killing of *S. aureus* strains that lack the anti-toxin EsaG[13]. An LXG-domain containing protein, TspA, encoded distally to the T7SS cluster shows

T7SS-dependent toxic activity, which was neutralized by an antitoxin TsaI[14]. Some substrates

modulate host interactions during infection; EsxA modulates host cell death and EsaE elicits

cytokine production during infection[15,16]. However, substrates of this system are yet to be defined

in many *S. aureus* strains and specific functions of effectors remain elusive.

T7SS in other bacterial species has been described to support bacterial functions like DNA

transfer, membrane integrity and spore development, although for *S. aureus*, it is not clear if T7SS

is important in staphylococcal physiology[17-19]. Our recent study on the T7SS substrate EsxC and

the transporter EssC showed that mutants lacking these proteins were more sensitive to

unsaturated host-derived fatty acids[20]. T7SS defects compromised membrane integrity and

induced oxidative stress in presence of antimicrobial fatty acids, indicating a role for these proteins

in membrane homeostasis. Here we report that the absence of T7SS causes an increased

sensitivity to membrane targeting drugs like daptomycin, both during infection *in vitro* and *in vivo*.

We further demonstrate distinct effects on cell surface morphology, membrane fluidity, charge and

protein localization in T7SS transporter and substrate mutants indicating a role for T7SS in

controlling membrane integrity. Our data suggest that targeting the T7SS could augment activities

of membrane-acting drugs like daptomycin.

Results

T7SS mutants are more sensitive to membrane-acting drugs

The cyclic lipopeptide antibiotic, daptomycin, is a key drug used to treat resilient *S. aureus*

infections. Daptomycin binds to the cell membrane in a calcium-dependent manner[21]. As our

previous studies indicated a potential impact of T7SS on membrane homeostasis[20], we

investigated whether the *S. aureus* USA300 JE2 mutants lacking either the substrate EsxC

(Δ*esxC)* or the central T7SS transporter EssC, (Δ*essC)* respond differently to membrane-acting

drugs like daptomycin. Growth of T7SS mutants were compared to the WT USA300 JE2 strain in

the presence or absence of daptomycin. When cultured in presence of 5 μg/ml daptomycin, growth

of the WT USA300 JE2 strain is delayed, but growth of Δ*esxC* and Δ*essC* was significantly

impaired compared to the WT (Fig. 1 A, B). No differences were seen between the strains in growth

rates in tryptic soy broth (TSB) without antibiotic (Fig. S1 A). Further, a 10-fold decrease in colony

forming unit counts for the mutants was also observed in the presence of daptomycin (Fig. 1 C).

When Δ*esxC* was complemented with a plasmid containing the *esxC* gene, the phenotype of the

WT was restored (Fig. 1 D, E). To investigate if other effectors had similar effects, we studied a

strain lacking another T7SS substrate, EsxD[11]. However, the deletion of *esxD* in USA300 JE2

strain did not result in any growth defects in the presence of daptomycin (Fig. S2 A, B). The T7SS

mutants Δ*esxC* and Δ*essC* were also found to be more sensitive to daptomycin in comparison to

the WT in a killing assay where logarithmic phase bacteria were treated for 2 h with daptomycin

(Fig. 1 F, G).

We investigated the effect of T7SS on daptomycin sensitivity in other strain backgrounds. The

growth of RN6390 *S. aureus* was impacted more by the addition of daptomycin, with the bacterial

growth initiating much later (~14 h) compared to the USA300 JE2 strain (~8 h). However, a similar

trend was seen in the growth of mutants; Δ*esxC* and Δ*essC* mutants were more susceptible than

the WT to treatment with daptomycin (Fig. S2 C, D). A defect in growth was also seen in USA300

LAC Δ*esxA* mutant, while Δ*esxB* appeared to grow similar to the WT (Fig. S2 E, F) The Newman

strain was also more sensitive than USA300 JE2 to daptomycin with growth being delayed until

around 14 h. There was less difference between WT and the Δ*esxA* mutant however a similar

trend was observed (Fig. S2 G, H).

To determine if the differences observed between the WT and T7SS mutants were limited to

daptomycin, other membrane-targeting antimicrobials, such as gramicidin and bithionol, were

tested. Gramicidin is a polypeptide antimicrobial peptide that forms ion channels in the cell

membrane allowing diffusion of monovalent ions such as potassium (K^+) and sodium (Na^+),

disrupting cell ion concentrations[22]. Bithionol is an inhibitor of soluble adenylyl cyclase and its

membrane-disrupting activity was associated with its ability to kill *S. aureus* persister cells[23]. When

treated with 2 µg/ml gramicidin, the WT JE2 strain did not have any growth defects, however the

T7SS mutants displayed a slower growth (Fig. 1 H). A similar defect in growth was also seen in

the presence of biothionol, although the Δ*esxC* and Δ*essC* mutant showed only a slightly slower

growth in the presence of 1 µg/ml bithionol (Fig. 1 I). Finally, this effect appears to be specific to

membrane-acting drugs as other drugs which target the cell wall (vancomycin, oxacillin) or DNA

replication (ciprofloxacin, mitomycin) did not differentially affect the growth of T7SS mutants as

compared to the WT (Fig. S3).

Δ*esxC* displays decreased membrane fluidity

Changes in membrane fluidity can affect membrane function and antibiotic sensitivity. We probed

T7SS effects on the membrane by assessing membrane fluidity of the WT and Δ*esxC* mutant.

Pyrene decanoic acid, an excimer-forming lipid[24], was employed to measure the fluidity of WT and

Δ*esxC* membranes. Compared to the WT, the cytosolic membrane of the Δ*esxC* mutant showed

a mild but statistically significant increase in rigidity (Fig. 2 A). Furthermore, we stained WT and

with a fluorescent membrane dye DiI12C, which has been used detect regions of increased fluidity

in bacteria[25]. As reported previously for *S. aureus*, we observe a few fluid regions in the WT, and

an uneven staining of the membrane. In contrast, a very weak staining of the membranes was

observed for the Δ*esxC* mutant (Fig. 2 B), which may indicate a decrease in membrane fluidity.

Finally, membrane fluidity was measured using the fluorescent dye Laurdan (6-dodecanoyl-2-

dimethylaminonaphthalene) which is incorporated into the membrane bilayer[26]. Logarithmic phase

WT cells treated with Laurdan showed a lower average generalized polarisation value of 0.33

compared to 0.42 for Δ*esxC* suggesting that the membrane of the mutant is more rigid in

comparison to that of the WT (Fig. 2 C).

Overall, the data indicate that EsxC contributes to *S. aureus* cell surface structure through

modulating membrane fluidity.

T7SS mutants have altered protein localization and membrane protein profiles

Functional membrane microdomains (FMMs) act as scaffolds for protein interactions and

oligomerization[27]. Bacterial flotillins are key scaffolding proteins which have been shown to

regulate membrane fluidity and protein organisation[28]. FloA also interacts with structural T7SS

proteins EssA and EssB, however interactions with effector proteins were not investigated[12]. To

investigate possible protein localisation defects in the T7SS mutants, the distribution of the

staphylococcal flotillin FloA was examined.

When exponentially growing FloA-eYFP expressing WT and ∆*esxC* and ∆*essC* mutants were

studied by fluorescence microscopy, a different localisation of GFP-tagged flotillin was observed

in the ∆*esxC* and ∆*essC* mutants compared to the WT (Fig. 2 D). The WT appears to display a

more homogenous distribution of flotillin throughout the membrane while both T7SS mutants have

a more defined heterogenous punctate distribution. The number of FloA foci per cell were

quantified. The WT had a broader distribution of with generally higher numbers of foci, but smaller

and with lower levels of fluorescence, while mutants had fewer foci but with higher levels of

fluorescence. This suggests that the T7SS components may have a role in modulating flotillin

localisation.

Previously, we reported that the mutants ∆*esxC* and ∆*essC* had distinct total protein profiles with

and without treatment with linoleic acid[20]. Given the changes seen in bacterial membranes in the

absence of T7SS proteins, we examined global changes in the protein content in the membrane.

Membrane fractions extracted from logarithmic phase (OD$_{600}$ 3.0) cultures of WT, ∆*esxC* and

∆*essC* were analysed by LC/MS. 13 and 15 proteins were altered in abundance [log2 (fold

change) = 1.0, *P* < 0.05] in Δ*esxC* and Δ*essC* respectively in relation to the WT. While 6

differentially abundant proteins were common to both mutants, overall, both mutants displayed

distinct profiles (Table 1). Interestingly, EssD, a T7SS substrate which is a nuclease, is decreased

significantly in the membranes of Δ*esxC* but not in Δ*essC*. To study the impact of EssC and EsxC

on protein export, a secretome analysis from culture supernatants was performed. Secretome

analysis showed a lower abundance of EssD in Δ*esxC* compared to WT (Table S1). In agreement

with the requirement of transporter EssC for T7 secretion, EssD, and other T7SS proteins (EsxA

and EsxC) were depleted in the secretome of Δ*essC* (Table S1). Again, there was some overlap

in the proteins secreted by both T7SS mutants, although many proteins were distinct to either

mutant strain.

Thus, the T7SS mutants show distinct changes to the overall membrane protein profiles, along

with changes in localisation of specific proteins.

T7SS mutants display altered cell surface morphology

To examine the cell surface of T7SS mutants we visualised the WT, Δ*esxC* and Δ*essC* using

scanning electron microscopy. The cell surface of WT grown to early logarithmic phase have the

typical smooth spherical shape of *S. aureus*, however, the isogenic mutants Δ*esxC* and Δ*essC*

appear deflated with dents or hollows (Fig. 3 A). The WT phenotype was restored in Δ*esxC*

complemented with a plasmid expressing *esxC* (Fig. 3 B). Interestingly, this cell surface defect is

not seen in all T7SS mutants as strains lacking EsxD (Δ*esxD*) does not have a defective cell

surface phenotype (Fig. 3 A). *S. aureus* cell surface alterations were seen in strains USA300 (LAC)

and Newman lacking EsxA (Δ*esxA)*, but not in Δ*esxB* (Fig. S4). However, RN6390 strains lacking

esxC and *essC* showed very little surface defects (Fig. S4). Differences seen in surface

morphologies between WT and mutants were not due to growth rate differences, as growth rates

of all mutant strains in the different strain backgrounds (Δ*essC*, Δ*esxC*, Δ*esxA*, Δ*esxB,* and

Δ*esxD)* in TSB, were similar to that of their WT (Fig. S1). Our data suggest that the cell envelopes

of mutants lacking specific T7SS components were altered, potentially due to a defect in the cell

membrane and/or cell wall structure.

T7SS mutants show decreased rate of cell wall synthesis

To investigate if the surface structure changes were associated with a defect in cell wall structure

or synthesis, transmission electron microscopy was undertaken to investigate any differences

between the cell wall ultrastructure of the T7SS mutants. However, there were no clear differences

in cell wall diameters between strains (Fig. S5). To study whether peptidoglycan synthesis and

accumulation were affected in the T7SS mutants, HADA, a fluorescent D-amino acid was used[29].

The dye was either added to a fresh subculture and grown until the bacteria reached logarithmic

phase or added for 10 min once the culture reached an OD_{600} of 1. No differences were observed

in the cells which had been grown with TSB supplemented with HADA (Fig. 4 A, B), however

addition of HADA during exponential phase growth revealed a faster accumulation of HADA in the

WT compared to *ΔessC* and *ΔesxC* (Fig. 4 A, C). This suggests that T7SS mutants displayed a

decreased rate of new peptidoglycan synthesis.

***ΔesxC* and *ΔesxA* have a more negative cell surface charge**

As we observe cell surface defects and changes to peptidoglycan synthesis, we next tested if the

disruption of the T7SS could alter other cell surface properties like surface charge. A cytochrome

binding assay was employed to measure surface charge; the binding of cytochrome C, a highly

positively charged protein, is directly proportional to the net negative surface charge of *S. aureus*

cells[30]. A significantly higher level of binding of cytochrome to cell surface of Δ*esxC* and Δ*esxA*

was observed compared to the USA300 WT strains, JE2 and LAC respectively. However, there

was no difference in the levels of binding between the WTJE2 and Δ*essC* (Fig. 5 A, C). This

suggests that the cell surface of Δ*esxC* is overall more negative. Much lower levels of cytochrome

C bound to USA300 LAC WT and Δ*esxB* suggesting these cell surfaces are also less negative

(Fig. 5 C). Colony forming unit (CFU) counts from samples after cytochrome C addition showed

no differences in cell numbers and therefore it is unlikely that bacterial cell numbers influenced

cytochrome C binding (Fig.5 B, D). These data indicate that the T7SS effectors affect the surface

charge of *S. aureus*.

Δ*esxC* shows increased binding of daptomycin

S. aureus has been reported to resist daptomycin through possessing a more positive surface

charge, causing electrostatic repulsion of the positively charged daptomycin calcium complex, and

thus preventing binding[31,32]. Based on the increased negative surface charge observed for the

T7SS mutants in the cytochrome C assay, we hypothesised that daptomycin could bind more

easily in the absence of a functional T7SS. To test this, daptomycin was tagged with BODIPY and

used to quantify binding via fluorometry and to visualise bound daptomycin by fluorescence

microscopy. In the presence of daptomycin, a higher level of daptomycin seen to bind around the

septum for the WT (Fig. 5 E, F) as reported previously[33]. The Δ*esxC* mutant showed significantly

higher fluorescence in comparison to the WT. Thus, our data suggest that the increased

daptomycin binding may be due to reduced electrostatic repulsion of daptomycin by Δ*esxC*.

When we stained bacteria with propidium iodide (PI), a fluorescent intercalating agent which

cannot pass through the membranes of viable cells, cytosols of daptomycin-treated Δ*esxC* and

Δ*essC* showed a higher intensity of PI compared to the WT as quantified by fluorescence

microscopy (Fig. 6 A, B). We also stained cells with the lipid staining dye FM1-43. FM1-43 staining

was of similar intensity, although it appears less ordered in the daptomycin-treated Δ*esxC* and

Δ*essC* samples (Fig. 6 A, B). Thus, membranes of the Δ*esxC* and Δ*essC* mutants are more

permeable in presence of daptomycin.

Decreased survival of T7SS mutants during intracellular infection with daptomycin

S. aureus is a facultative intracellular pathogen, and daptomycin is considered to kill intracellular

bacteria effectively. Hence, we wanted to determine if deletion of *essC* and *esxC* from *S. aureus*

would enhance intracellular killing activity. A549 human lung epithelial cells[15] infected with *S.*

aureus were treated with a high concentration of daptomycin (20 µg/ml). While there were no

differences in the bacterial numbers between the WT and T7SS mutants in internalised bacteria

(1 h after infection), a significant decrease was observed in the numbers of Δ*esxC* and Δ*essC*

surviving within cells treated with daptomycin in comparison to WT after 4 h and 24 h after infection

(Fig. 7 A). Notably, there was a significant number of WT *S. aureus* bacteria surviving intracellularly

after treatment with daptomycin. Confocal microscopy imaging of infected cells also showed

similar number of bacterial cells entering the mammalian cells but more *S. aureus* WT remaining

compared to Δ*essC* and Δ*esxC* 24 h after infection in presence of daptomycin (Fig. 7 B, C).

USA300 JE2 WT, Δ*essC* and Δ*esxC* showed no differences in growth rate in the cell culture

medium, DMEM-10, indicating that the differences seen in the infection assay were not due to the

ability of the mutants to grow differently (Fig. S6A); as with the TSB, a growth defect was seen in

the T7SS mutants in presence of daptomycin (Fig. S6B). Therefore, our data suggest that

daptomycin is more effective in killing intracellular *S. aureus* in the absence of T7SS.

T7SS mutants are more sensitive to daptomycin in a skin infection model

We next investigated how effective daptomycin was in curbing *S. aureus* infection *in vivo* and if

the absence of T7SS had any impact on daptomycin killing *in vivo*. T7SS proteins have been

reported to be important in various murine infection models[6,8,34]. Here, we employed a localised

murine skin abscess model to investigate both the role of T7SS protein EsxC during skin infection

and if its presence could affect daptomycin activity.

7-8-week-old BALB/c mice were infected with *S. aureus* WT and Δ*esxC* intradermally, followed by

treatment (intravenous) 2 h post infection (h.p.i) and every 24 h after, with 10 mg/kg daptomycin.

Bacterial counts were obtained from skin tissue 24 or 72 h.p.i. Weight and skin lesion diameters

were recorded every day. In the groups not treated with daptomycin, bacterial counts from the site

of infection (lesions) showed no significant difference between WT and $\Delta esxC$ infected mice after

24 or 72 h of infection. A significant decrease was seen in the numbers of $\Delta esxC$ present in the

skin of daptomycin-treated mice compared to the untreated group 24 and 72 h.p.i, while this was

not the case for the WT infected group. The daptomycin treated $\Delta esxC$ infected group had lower

mean bacterial count (7.32×10^7 at 24 h and 7.39×10^6 at 72 h) than the WT-infected, daptomycin

treated group (1.12×10^8 at 24 h and 1.45×10^7 at 72 h) and (Fig. 8A and B). No significant

differences were observed between the areas of lesions measured for WT or $\Delta esxC$ after 24 h or

72 h infections and in daptomycin-treated mice (Fig S7). Although the dose of daptomycin used in

the time frame of treatment was not sufficient to treat WT infection, this data suggests that the

strain lacking EsxC was more susceptible to daptomycin in $vivo$. Interestingly, when cytokines

TNF-α and IL-β were measured, there was a significant difference between the treated and

untreated groups for both strains 72 h.p.i. (Fig 8 C and D). The mutant induced significantly less

IL1-β than the WT at 72 h.p.i., irrespective of daptomycin treatment (Fig 8 D).

Discussion

The $S.$ $aureus$ T7SS has been strongly associated with staphylococcal virulence and ability of this

pathogen to persist during infection[7]. Its importance for bacterial competition between $S.$ $aureus$

strains has been recently highlighted[13]. However, the role of this system in staphylococcal

physiology has been poorly defined. Here we report that the T7SS is critical in maintaining cell

membrane homeostasis through modulating membrane properties including fluidity, accompanied

by changes in membrane protein localisation, changes to the rate of cell wall synthesis and surface

charge. Consequently, T7SS impacts activity of membrane acting drugs including daptomycin, a

key drug used to treat resilient staphylococcal infections. Our data demonstrate that T7SS can

modulate the activity of daptomycin both intracellularly as well as during infection in $vivo$.

Several studies in mycobacterial species have linked the T7SS to cell surface integrity in different

ways[19,35-38]. The ESX-1 system has been previously reported to impact cell membrane

permeability in *M. tuberculosis*, although this effect appeared to be strain specific [35]. However,

ESX-5 which is considered essential for *in vitro* growth, has been linked to mycobacterial capsular,

cell wall and outer membrane permeability[19,36,39]. As ESX-5 was no longer essential in *M. marinum*

when the membrane was permeabilised, it was proposed that ESX-5 mediates transport of cell

envelope proteins to the outer membrane required for uptake of essential nutrients. While these

substrates have not been identified, a substrate PPE-10 was associated with capsular

integrity[19,36]. In *S. aureus,* we previously reported that the effectors, EsxC, and the ATPase, EssC,

were associated with maintenance of membrane integrity in response to antimicrobial host fatty

acids[20].

T7SS proteins have previously been linked with membrane fluidity and flotillins. Higher levels of

T7SS transcription were associated with a decrease in membrane fluidity, caused by the fatty acid

kinase complex, which incorporates host fatty acids into the membrane, inducing a change in

membrane dynamics[24]. *S. aureus* encodes one flotillin homolog, which shares 84% homology with

the FloA protein in *B. subtilis* [40]. FloA (also referred to as FloT) in *S. aureus,* has been linked to

T7SS assembly [12]. T7SS membrane proteins interact with FloA within functional membrane

microdomains (FMM) promoting more efficient interactions between T7SS membrane

components EssB and EssA, helping incorporate EssA into the EssC-EsaA-EssB pre-complex.

Here we see that the membrane of the *ΔesxC* mutant was more rigid than the WT, and floA

appears to localize differently in the membranes of this mutant. It is not clear whether the

localisation of the flotillin is due to the more rigid membrane in the mutants or if the flotillin

mislocalisation in the mutants causes the membrane to become more rigid, as flotillins have a role

in spatially organising membrane domains by acting as scaffolds to guide proteins to the correct

membrane areas [28]. Thus, EsxC may have a direct but transient interaction with FloA in the FMM

or an indirect impact upon other membrane-associated proteins which could impact the lipid

packing and fluidity of the membrane.

While we demonstrate that T7SS and specific substrates play a role in maintaining membrane

integrity, we do not know how the T7SS contributes to this. There are multiple possibilities 1.

substrates like EsxC are chaperones for membrane proteins and play a role in targeting them to

the membrane. A recent bacterial two-hybrid library screening we performed was not successful

in identifying any direct bacterial interactors, however it is possible that such interactions are

transient. 2. T7SS proteins may impact the state of the membrane by binding other molecules like

metal ions required for maintaining stability of the membrane. 3. destabilizing this complex

secretion apparatus has a non-specific effect on the physical structure of the membrane.

It is interesting that not all T7SS proteins/ substrates have a cell surface associated defect. This

indicates that only certain substrates of the system are mediating this effect. EsxD and EsxB did

not appear to be associated with surface morphology defects as observed for EsxC and EsxA.

Surface charges corelated well with daptomycin sensitivity indicating that these substrates did not

play a role in the membrane homeostasis. Surface charge measurements of WT and mutants

using a cytochrome C binding assay revealed that Δ*esxC* and Δ*esxA* had an increased negative

surface charge, whereas *ΔesxB* and *ΔessC* did not display any differences in charge to the WT.

This may be because esxC is still produced by *ΔessC*, although it is not secreted, thus

accumulating in the cytosol and membrane, further indicating that EsxC plays a key role in

maintaining surface charge.

While EsxA is conserved across all *S. aureus* strains, only the essC1 locus encodes EsxC [8,9]. It is

however likely that all *S. aureus* strains encode a protein orthologous to EsxC as proteins in

module 2, with variants essC2 – 4, have not yet been characterised. Therefore, it is possible that

one of the hypothetical proteins has functions like EsxC. Additionally, several of the phenotypes

reported for strains lacking EsxC were also seen in strains which lacked EsxA. Previous reports

have suggested that EsxC and EsxA interact[11]. Therefore, this could mean that interactions

between EsxC and EsxA may be important in modulating membrane integrity.

Daptomycin is thought to bind to the cell membrane before forming a complex with cell wall

biosynthesis machinery, preventing cell wall synthesis, before dispersing throughout the bacterial

membrane causing depolarisation[41]. Daptomycin, although not a cationic antimicrobial peptide,

forms a highly positively charged complex with calcium to become active. Significantly more

daptomycin was bound to the *ΔesxC* than the WT, which supports the hypothesis that the

decrease in surface charge measured in *ΔesxC* allows easier binding of daptomycin to the cell

membrane. However, the surface charge of *ΔessC* does not differ from the WT and therefore

increased susceptibility to the positively charged daptomycin is likely not caused solely by charge-

dependent binding. Also, in contrast to daptomycin, membrane targeting antimicrobials bithionol

and gramicidin are uncharged[23,42], therefore surface charge and thus decreased electrostatic

repulsion of the antibiotic in the T7SS mutants is not the only factor involved in increased

sensitivity. Daptomycin appears to have the highest impact on the T7SS mutants whereas

bithionol appears to have the lowest activity. As all three antimicrobials have different mechanisms

of action once bound to the membrane before causing membrane depolarization, the subsequent

steps after membrane binding appear to be important for efficacy. Therefore, perhaps daptomycin

treatment which impacts both the cell membrane, and the cell wall causes a larger effect in the

T7SS mutants.

While daptomycin increases killing of T7SS mutants *in vivo*, at the dose and strain of *S. aureus*

used, there was no impact of the T7SS in bacterial survival during acute skin infection. T7SS

proteins have been previously reported to be important for bacterial virulence using invasive

kidney abscess, nasal colonisation and skin infection models, although studies have used different

S. aureus strains[6,8,34]. The lack of decrease in the lesion size of the T7SS mutant-infected mice

treated with daptomycin, unlike the trend towards decreased lesion size observed for the WT (Fig

S7), could indicate that the increased killing of bacteria may cause an altered local immune

response. This is supported by the different levels of inflammatory cytokines seen in the

daptomycin treated WT and the mutant-infected mice. While our infection studies indicate a

potential role for EsxC (in a USA300 strain background) in modulation of IL1-β, the data indicate

that it may not play a role in bacterial survival during skin infection. A recent study reported that in

a *S. aureus* strain WU1, the T7SS was important only for invasive infection and not localized nasal

colonization [43].

Currently, levels of resistance to daptomycin are low, however, with increasing resistance in other

last line antibiotics, such as vancomycin, the use of daptomycin going forward will increase[44]. The

increased use of daptomycin will likely lead to the occurrence of more resistant isolates, therefore

research into new antibiotics, and targeting new proteins that can enhance existing treatments

could be a powerful approach. Our data indicate that inhibitors of T7SS proteins or of T7SS

secretion could have excellent potential in combination drug therapies.

Materials and Methods

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, m.unnikrishnan@warwick.ac.uk

Materials availability

This study did not generate new unique reagents.
416

Experimental model and study participant details

Animals

All animal procedures were carried out as per protocols in the Project Licence PCEC27E7D. All local ethical and home office approvals were obtained for animal experimentation. For the murine skin abscess model, female 7/8-week-old BALB/c mice (Charles River Laboratories, UK) flanks were shaved before injection, and inoculated through an intradermal route with 50 µl of bacterial suspension per flank (one injection on each flank), resulting in a final bacterial count of 1-2 x 10^6 per mouse. Overnight *S. aureus* cultures were subcultured, grown to OD_{600} of 1, centrifuged and washed twice with PBS, and diluted in PBS to achieve a final CFU/ml of 1-2 x 10^7 for infection. 2 h after infection mice were given an intravenous dose of daptomycin or PBS via tail vein injection. This was repeated once every 24 h. During the experiment, mice were weighed daily, and any abscess formation was measured and recorded. After 24 h or 72 h mice were euthanised by CO_2 inhalation and skin at the infection site was incised. The skin samples were weighed and homogenised in PBS using a FastPrep (MP Biomedicals), at 4 m/s for 60 seconds for a total of 6 cycles. Homogenates were diluted in PBS and CFU counts were determined and normalized to tissue weight.

Method details

Bacterial strains

S. aureus strains used in this study are included in Table 2. Strains were cultured aerobically in tryptic soy broth (TSB) (Sigma-Aldrich) at 37°C with aeration at 180 rpm overnight. Overnight cultures were subcultured into fresh TSB and grown to a density of optical density (OD_{600}) of 1, unless stated otherwise. For complemented strains TSB was supplemented with 10 µg/ml chloramphenicol.

Growth curves

Overnight bacterial cultures were diluted to an OD_{600} of 0.1 in TSB or TSB supplemented with antimicrobials where specified. Bacteria were grown in a 96-well plate with shaking and the OD_{600} was measured every 15 min with a FLUOstar OMEGA plate reader (BMG Labtech, UK). At appropriate time points 100 µl of the samples were taken for CFU determination.

Daptomycin killing assay

Bacterial cultures were grown overnight in TSB, diluted to an OD_{600} of 0.15 and grown to log phase. Bacterial cultures were then treated with 10 µg/ml of daptomycin (Acros Organics) in the presence of 1 mM $CaCl_2$ (Fisher Scientific) and incubated at 37ºC for a further 2 h. Aliquots were removed at indicated timepoints, serially diluted in PBS and plated on tryptic soy agar (TSA) for CFU determination.

Scanning Electron Microscopy

Overnight bacterial cultures were diluted to an OD_{600} of 0.15 and grown to an OD_{600} of 1 in TSB. The pellet was washed with PBS twice before dropping onto poly-lysine coated 12 mm circular coverslips. Samples were fixed with 2.5% glutaraldehyde in PBS for 1 h at 4°C. The bacterial cells were dehydrated by washing with a series of ethanol solutions (20%, 50%, 70%, 90% and 100%) and dried with hexamethyldisilazane. The samples were sputter coated with carbon and then observed using a Gemini SEM 500 (Zeiss).

Transmission Electron Microscopy

Overnight bacterial cultures were diluted to an OD_{600} of 0.15 and grown to an OD_{600} of 1 in TSB. The pellet was washed in PBS three times and fixed with paraformaldehyde and glutaraldehyde for 1 h at room temperature (RT). Pellets were rinsed a further three times in PBS before incubation with 1% osmium tetroxide for 1 h at RT. The bacterial cells were dehydrated by washing with a graded acetone series for 20 min each and transferred to graded acetone epoxy resin mixture for 45 min each. Cells were incubated in pure resin for 24 h before being hardened overnight at 60°C. The specimens were sectioned into 90 nm slices using an ultramicrotome

(RMC) before being placed onto grids and stained with heavy metals. The samples were

visualised using a TEM-2100Plus (Jeol).

Determination of membrane fluidity by Laurdan staining

Membrane fluidity was measured using the fluorescent dye Laurdan (Sigma). Logarithmic phase

S. aureus was grown, and 1 ml aliquots were washed in PBS. The samples were incubated with

a final concentration of 100 µM Laurdan for 5 min at RT in the dark. Samples were washed four

times in PBS and 200 µl was transferred to a black walled 96 well plate in triplicate. Membrane

fluidity was determined by using an excitation wavelength of 350 nm and measuring fluorescence

at 460 nm and 500 nm emission wavelengths on a BioTek Cytation 5 Cell Imaging Multimode

Reader (Agilent). The generalised polarisation (GP) value was calculated using the following

calculation, $GP = (I_{460} - I_{500}) / (I_{460} + I_{500})$.

DilC12 staining

O/N bacterial cultures were diluted to an OD600 of 0.15 in TSB with 1 µg/mL 1,1′-didodecyl-

3,3,3′,3′-tetramethylindocarbocyanine perchlorate (DilC12) (Invitrogen). Cultures were grown to

an OD600 of 1.0, centrifuged and washed twice with fresh TSB. Samples were spotted on to

agarose pads and imaged using a Leica DMi8 widefield microscope (Leica Microsystems, UK).

Acquired images were analysed with the ImageJ processing package, Fiji.

Cytochrome C binding assay

Overnight bacterial cultures were pelleted and washed with 20 mM MOPS buffer, pH 7. The cells

were resuspended in MOPS buffer at an OD_{600} of 1 and incubated with 50 µg/ml cytochrome C at

RT for 15 min. The cells were pelleted and the remaining cytochrome C in the supernatant was

measured spectrophotometrically at OD_{410} using a FLUOstar OMEGA plate reader (BMG Labtech,

UK).

HADA incorporation assay

Peptidoglycan synthesis was investigated by incorporation of HCC-amino-D-alanine (HADA)

incorporation. Overnight *S. aureus* cultures were diluted into TSB supplemented with 25 µM HADA

and incubated at 37ºC in the dark. When OD_{600} of 1.0 was reached cultures were washed three

times and fixed with 4% PFA. Alternatively, cultures were treated with daptomycin for 30 min before

washing and fixing. Microscopy was undertaken on these samples using the DAPI filter set on a

Leica DMi8 widefield microscope.

Localisation of FloA-eYFP

An overnight culture was diluted and grown to logarithmic phase. 1 ml of culture was centrifuged

and washed with PBS before fixing with 4% PFA at RT for 15 min. The bacteria were pelleted and

washed twice in PBS before resuspension in 100 µl of PBS. Bacteria were imaged using API

Deltavision 2 (Applied Precision Instruments) widefield microscope using YFP filter set at 100x

magnification.

Synthesis of Daptomycin BODIPY

A 1 mg/ml stock solution of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC),

N-Hydroxysuccinimide (NHS), daptomycin and BODIPY was prepared in DMSO. Th mixture was

stirred for 16-24 h at RT. 10 µl of the crude sample was removed and diluted with High

Performance Liquid Chromatography (HPLC) solvent, filtered and used for HPLC analysis. The

solution was purified by dialysis with ultrapure water to remove EDC, NHS and any other unwanted

by-products with a membrane cut-off of 100 – 500 Da. The water was replaced every 2 h for a

total of 2 times. The sample was freeze dried to remove the water.

Daptomycin BODIPY binding

S. aureus was grown as previously described and treated with daptomycin once logarithmic phase

had been reached. The samples were incubated at 37ºC for 30 min in the dark then washed with

PBS three times. 200 µl were aliquoted into black 96 well plates in triplicate and fluorescence was

measured using a BioTek Cytation 5 Cell Imaging Multimode Reader (Agilent). The excitation

wavelength used was 488 nm and emission wavelength used was 530 nm.

Propidium Iodide Staining

Bacterial cultures were grown in TSB to an OD_{600} of 1, centrifuged and resuspended in TSB

supplemented with 1 mM $CaCl_2$. Samples with and without daptomycin were incubated at 37°C

for a further hour before propidium iodide and FM1-43 were added to a final concentration of 3 μM

and 10 ng/μl, respectively. Samples were washed in PBS twice and spotted onto agarose pads.

Once dry, cover slips were mounted using DAPI anti-fade mounting fluid. Samples were imaged

on a Leica DMi8 widefield microscope (Leica Microsystems, UK) using 100x objective. Propidium

iodide, FM1-43 and DAPI were imaged using the Texas Red, FITC and DAPI filter sets

respectively.

Invasion and survival Assay

Infection assays were carried out in 24-well tissue culture treated plates (Falcon), or RS treated

glass chamber slides (154526, Nunc, Lab Tek II). Overnight *S. aureus* cultures were diluted to an

OD_{600} of 0.15 and grown to an OD_{600} of 1. Cultures were resuspended in DMEM media to a density

of 2.5 x10^6 colony forming units (CFU)/ml. A549 epithelial cells were seeded in 24 well plates at a

density of 2.5 x 10^5 a day before being infected with *S. aureus* at a multiplicity of infection (MOI)

of 10. Cells were infected for 1 h at 37°C with 5% CO_2. To kill any extracellular bacteria, cells were

washed with DMEM media containing 20 μg/ml lysostaphin (Sigma-Aldrich) and 50 μg/ml

gentamycin (Melford) and incubated for a further 30 min. Cells were washed twice with PBS, lysed

with 1 ml cold sterile water and vigorous pipetting before plating for CFU determination. To

investigate the course of infection after extracellular bacterial killing, the cells are washed twice

with PBS and refreshed with DMEM with 1 mM $CaCl_2$ only or DMEM supplemented with 1 mM

$CaCl_2$ and 10 μg/ml daptomycin and incubated until the desired timepoint. The above protocol to

lyse the cells is then followed.

Confocal microscopy

At required timepoints A549 cells infected with *S. aureus* were washed twice with PBS and fixed

with 4% paraformaldehyde (PFA) for 15 min. The chamber slide wells were washed in 1 ml PBS

and permeabilised with 1% saponin, 0.1% triton X-100 in PBS for 20 min at RT. Cells were blocked

with 3% BSA in PBS for 30 min at RT. Slides were then washed with 1% BSA in PBS and twice

with PBS only. Cells were stained with phalloidin 650 in 1% BSA for 40 min at RT and vancomycin-

BODIPY in 1% BSA for 20 min. DAPI anti-fade mounting fluid was added to the slide. Samples

were imaged using a spinning disk confocal UltraVIEW (PerkinElmer).

Preparation of samples for proteomics

Preparation of membrane fractions: *S. aureus* strains were grown to an OD_{600} of 3.0 in TSB.

After centrifugation of cultures, pellets were washed with PBS and resuspended PBS

supplemented with one cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail tablet, 250 mM

sucrose, 1 mM EDTA, and 50 µg/mL lysostaphin. Samples were incubated for 15 min at 37ºC.

Samples were aliquoted into lysing matrix B tubes (MP Biomedicals™), and bacteria were lysed

with a Fastprep machine (6.5 m/s, 2 x 45 s). Samples were centrifuged at 11 000 × g for 15 min,

and supernatants were collected and then ultracentrifuged (230 000 × g for 30 min at 4ºC) in clean,

thick wall polyallomer tubes. Pellets were resuspended in 400 µL PBS supplemented with 250 mM

sucrose and 10% glycerol.

Preparation of culture supernatant: *S. aureus* strains were grown to an OD600 of 3 in TSB.

After centrifugation of cultures, supernatants were sterile-filtered and incubated at 4°C overnight

with 10% trichloroacetic acid and 50 µM sodium deoxycholate in presence of one cOmplete™,

Mini, EDTA-free Protease Inhibitor Cocktail tablet (Sigma-Aldrich, UK). The precipitated proteins

were centrifuged at 10 000 g at 4°C for 15 min, gently washed with acetone twice, dried at RT for

10 min and resuspended in PBS. After denaturation, precipitated proteins were run on a gel until

all the proteins had moved from the stacking gel into the resolving gel. The gel was then stained

with InstantBlueTM (Sigma-Aldrich, UK) for 3 h, after which the protein bands were excised and

diced. Proteins were in-gel digested with trypsin as described elsewhere[45]. Briefly, proteins were

reduced and alkylated for 5 min at 70°C with 10 mM TCEP [tris(2-carboxyethyl) phosphine] and

40 mM CAA (2-chloroacetamide), respectively. Tryptic digestion was carried out at O/N at 37°C

in 50 mM ABC.

Label-free protein quantification.

Peptides prepared for proteome analyses were desalted and concentrated with a C18 cartridge in

40 µL MS buffer (2% acetonitrile plus 0.1% trifluoroacetic acid). For each sample, 20 µL were

analysed by nanoLC-ESI-MS/MS using the Ultimate 3000/Orbitrap Fusion instrumentation

(Thermo Scientific), and a 90-minute LC separation on a 50 cm column. The data were used to

interrogate the Uniprot *Staphylococcus aureus* USA300 database UP000001939, and the

common contaminant database from MaxQuant (Cox et al., 2014). MaxQuant software was used

for protein identification and quantification using 567 default settings. Intensities were log2-

transformed with the Perseus software, and proteins with one or no valid value for every sample

in triplicate were filtered.

The proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE

partner repository with the dataset identifier PXD047096 and 10.6019/PXD047096. Membrane

proteomic and secretome samples are labelled MS18-178 and MS19-126, respectively.

ELISA

The levels of mouse TNF-α and IL-1β were quantified from the supernatant of homogenised skin

samples using Mouse TNF-alpha DuoSet ELISA (DY410) or Mouse IL-1 beta/IL-1F2 DuoSet

ELISA (DY401) (R&D Systems) following the manufacturer's protocols. The optical density was

measured at 540 nm and subtracted from readings at 450 nm using the BioTek Cytation 5 Cell

Imaging Multimode Reader (Agilent). Concentrations were calculated from standard curves run in

parallel.

Image Analysis

All microscopy images were analysed using FIJI (ImageJ, Version 2.0.0-rc-69/1.52p). To calculate

fluorescence, a threshold value was set, and area or number of bacterial cells was used to

normalise fluorescent values against.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism 9.0 (MacOS version 9.5.1). Pairwise statistical analysis was carried out by the Man-Whitney U test for two groups, and one or two-way ANOVAs for multiple groups, followed by post-hoc Tukey's HSD tests where necessary. Significance was denoted by asterisks; * = P < 0.05, ** = P < 0.01, *** = P < 0.001, **** = P < 0.0001 and ns = not significant. The fold changes and *P* values of the proteomics data were calculated with the R package limma[46].

Acknowledgements

We would like to acknowledge funding from MIBTP (grant number) to Victoria Smith and the Medical Research Council grant (MR/N010140/1) for funding.

We would like to thank Novartis Vaccines, Siena (GSK biologicals) Italy for providing the *ΔesxA* and *ΔesxB* strains, and Prof Tracy Palmer for providing the RN6390 *ΔesxC* and *ΔessC* strains, used in this study. We acknowledge the Midlands Regional Cryo-EM Facility, hosted at the Warwick Advanced Bioimaging Research Technology Platform, for the use of the JEOL 2100Plus.

Author contributions

VS, AKT and MU conceptualised ideas, designed experiments, VS, GC, KEW, RGM, JY, AKT conducted experiments, MU and SP provided resources for the study, VS, AKT, GC, KEW, RGM and MU analysed data, AKT and MU supervised study, MU, VS and AKT drafted manuscript and all authors were involved in reviewing manuscript draft.

References

1.   Tong, S.Y., Davis, J.S., Eichenberger, E., Holland, T.L., and Fowler, V.G., Jr. (2015). Staphylococcus aureus infections: epidemiology, pathophysiology, clinical manifestations, and management. Clin Microbiol Rev *28*, 603-661. 10.1128/CMR.00134-14.

2.   Chambers, H.F., and Deleo, F.R. (2009). Waves of resistance: Staphylococcus aureus in

the antibiotic era. Nature reviews. Microbiology *7*, 629-641. 10.1038/nrmicro2200.

3.   Lee, B.Y., Singh, A., David, M.Z., Bartsch, S.M., Slayton, R.B., Huang, S.S., Zimmer, S.M.,

Potter, M.A., Macal, C.M., Lauderdale, D.S., et al. (2013). The economic burden of

community-associated methicillin-resistant Staphylococcus aureus (CA-MRSA). Clinical

microbiology and infection : the official publication of the European Society of Clinical

Microbiology and Infectious Diseases *19*, 528-536. 10.1111/j.1469-0691.2012.03914.x.

4.   Cheung, G.Y.C., Bae, J.S., and Otto, M. (2021). Pathogenicity and virulence of

Staphylococcus aureus. Virulence *12*, 547-569. 10.1080/21505594.2021.1878688.

5.   Bottai, D., Groschel, M.I., and Brosch, R. (2016). Type VII Secretion Systems in Gram-

Positive Bacteria. Curr Top Microbiol Immunol. 10.1007/82_2015_5015.

6.   Burts, M.L., Williams, W.A., DeBord, K., and Missiakas, D.M. (2005). EsxA and EsxB are

secreted by an ESAT-6-like system that is required for the pathogenesis of Staphylococcus

aureus infections. Proceedings of the National Academy of Sciences of the United States

of America *102*, 1169-1174.

7.   Unnikrishnan, M., Constantinidou, C., Palmer, T., and Pallen, M.J. (2017). The Enigmatic

Esx Proteins: Looking Beyond Mycobacteria. Trends in microbiology *25*, 192-204.

10.1016/j.tim.2016.11.004.

8.   Kneuper, H., Cao, Z.P., Twomey, K.B., Zoltner, M., Jager, F., Cargill, J.S., Chalmers, J.,

van der Kooi-Pol, M.M., van Dijl, J.M., Ryan, R.P., et al. (2014). Heterogeneity in ess

transcriptional organization and variable contribution of the Ess/Type VII protein secretion

system to virulence across closely related Staphyloccus aureus strains. Molecular

microbiology *93*, 928-943. 10.1111/mmi.12707.

9.   Warne, B., Harkins, C.P., Harris, S.R., Vatsiou, A., Stanley-Wall, N., Parkhill, J., Peacock,

S.J., Palmer, T., and Holden, M.T. (2016). The Ess/Type VII secretion system of

Staphylococcus aureus shows unexpected genetic diversity. BMC genomics *17*, 222.

10.1186/s12864-016-2426-7.

10. Bowman, L., and Palmer, T. (2021). The Type VII Secretion System of Staphylococcus.

Annu Rev Microbiol *75*, 471-494. 10.1146/annurev-micro-012721-123600.

11. Anderson, M., Aly, K.A., Chen, Y.H., and Missiakas, D. (2013). Secretion of atypical protein

substrates by the ESAT-6 secretion system of Staphylococcus aureus. Molecular

microbiology *90*, 734-743. 10.1111/mmi.12395.

12. Mielich-Suss, B., Wagner, R.M., Mietrach, N., Hertlein, T., Marincola, G., Ohlsen, K.,

Geibel, S., and Lopez, D. (2017). Flotillin scaffold activity contributes to type VII secretion

system assembly in Staphylococcus aureus. PLoS pathogens *13*, e1006728.

10.1371/journal.ppat.1006728.

13. Cao, Z., Casabona, M.G., Kneuper, H., Chalmers, J.D., and Palmer, T. (2016). The type

VII secretion system of Staphylococcus aureus secretes a nuclease toxin that targets

competitor bacteria. Nat Microbiol *2*, 16183. 10.1038/nmicrobiol.2016.183.

14. Ulhuq, F.R., Gomes, M.C., Duggan, G.M., Guo, M., Mendonca, C., Buchanan, G.,

Chalmers, J.D., Cao, Z., Kneuper, H., Murdoch, S., et al. (2020). A membrane-depolarizing

toxin substrate of the Staphylococcus aureus type VII secretion system mediates

intraspecies competition. Proceedings of the National Academy of Sciences of the United

States of America *117*, 20836-20847. 10.1073/pnas.2006110117.

15. Korea, C.G., Balsamo, G., Pezzicoli, A., Merakou, C., Tavarini, S., Bagnoli, F., Serruto, D.,

and Unnikrishnan, M. (2014). Staphylococcal Esx proteins modulate apoptosis and release

of intracellular Staphylococcus aureus during infection in epithelial cells. Infection and

immunity *82*, 4144-4153. 10.1128/IAI.01576-14.

16. Anderson, M., Ohr, R.J., Aly, K.A., Nocadello, S., Kim, H.K., Schneewind, C.E.,

Schneewind, O., and Missiakas, D. (2017). EssE Promotes Staphylococcus aureus ESS-

670   Dependent Protein Secretion To Modify Host Immune Responses during Infection. Journal

671   of bacteriology *199*. 10.1128/JB.00527-16.

672   17.   Fyans, J.K., Bignell, D., Loria, R., Toth, I., and Palmer, T. (2013). The ESX/type VII

673   secretion system modulates development, but not virulence, of the plant pathogen

674   Streptomyces scabies. Molecular plant pathology *14*, 119-130. 10.1111/j.1364-

675   3703.2012.00835.x.

676   18.   Flint, J.L., Kowalski, J.C., Karnati, P.K., and Derbyshire, K.M. (2004). The RD1 virulence

677   locus of Mycobacterium tuberculosis regulates DNA transfer in Mycobacterium smegmatis.

678   Proceedings of the National Academy of Sciences of the United States of America *101*,

679   12598-12603. 10.1073/pnas.0404892101.

680   19.   Ates, L.S., Ummels, R., Commandeur, S., van de Weerd, R., Sparrius, M., Weerdenburg,

681   E., Alber, M., Kalscheuer, R., Piersma, S.R., Abdallah, A.M., et al. (2015). Essential Role

682   of the ESX-5 Secretion System in Outer Membrane Permeability of Pathogenic

683   Mycobacteria. PLoS genetics *11*, e1005190. 10.1371/journal.pgen.1005190.

684   20.   Kengmo Tchoupa, A., Watkins, K.E., Jones, R.A., Kuroki, A., Alam, M.T., Perrier, S., Chen,

685   Y., and Unnikrishnan, M. (2020). The type VII secretion system protects Staphylococcus

686   aureus against antimicrobial host fatty acids. Sci Rep *10*, 14838. 10.1038/s41598-020-

687   71653-z.

688   21.   Huang, H.W. (2020). DAPTOMYCIN, its membrane-active mechanism vs. that of other

689   antimicrobial peptides. Biochim Biophys Acta Biomembr *1862*, 183395.

690   10.1016/j.bbamem.2020.183395.

691   22.   David, J.M., and Rajasekaran, A.K. (2015). Gramicidin A: A New Mission for an Old

692   Antibiotic. J Kidney Cancer VHL *2*, 15-24. 10.15586/jkcvhl.2015.21.

693   23.   Kim, W., Zou, G., Hari, T.P.A., Wilt, I.K., Zhu, W., Galle, N., Faizi, H.A., Hendricks, G.L.,

694   Tori, K., Pan, W., et al. (2019). A selective membrane-targeting repurposed antibiotic with

695   activity against persistent methicillin-resistant Staphylococcus aureus. Proceedings of the

National Academy of Sciences of the United States of America *116*, 16529-16534.

10.1073/pnas.1904700116.

24.   Lopez, M.S., Tan, I.S., Yan, D., Kang, J., McCreary, M., Modrusan, Z., Austin, C.D., Xu, M.,

and Brown, E.J. (2017). Host-derived fatty acids activate type VII secretion in

Staphylococcus aureus. Proceedings of the National Academy of Sciences of the United

States of America *114*, 11223-11228. 10.1073/pnas.1700627114.

25.   Strahl, H., Burmann, F., and Hamoen, L.W. (2014). The actin homologue MreB organizes

the bacterial cell membrane. Nat Commun *5*, 3442. 10.1038/ncomms4442.

26.   Scheinpflug, K., Krylova, O., and Strahl, H. (2017). Measurement of Cell Membrane

Fluidity by Laurdan GP: Fluorescence Spectroscopy and Microscopy. Methods Mol Biol

1520, 159-174. 10.1007/978-1-4939-6634-9_10.

27.   Lopez, D., and Kolter, R. (2010). Functional microdomains in bacterial membranes. Genes

Dev *24*, 1893-1902. 10.1101/gad.1945010.

28.   Bach, J.N., and Bramkamp, M. (2013). Flotillins functionally organize the bacterial

membrane. Molecular microbiology *88*, 1205-1217. 10.1111/mmi.12252.

29.   Kuru, E., Hughes, H.V., Brown, P.J., Hall, E., Tekkam, S., Cava, F., de Pedro, M.A., Brun,

Y.V., and VanNieuwenhze, M.S. (2012). In Situ probing of newly synthesized peptidoglycan

in live bacteria with fluorescent D-amino acids. Angew Chem Int Ed Engl *51*, 12519-12523.

10.1002/anie.201206749.

30.   Peschel, A., Otto, M., Jack, R.W., Kalbacher, H., Jung, G., and Gotz, F. (1999). Inactivation

of the dlt operon in Staphylococcus aureus confers sensitivity to defensins, protegrins, and

other antimicrobial peptides. The Journal of biological chemistry *274*, 8405-8410.

10.1074/jbc.274.13.8405.

31.   Bertsche, U., Yang, S.J., Kuehner, D., Wanner, S., Mishra, N.N., Roth, T., Nega, M.,

Schneider, A., Mayer, C., Grau, T., et al. (2013). Increased cell wall teichoic acid production

and D-alanylation are common phenotypes among daptomycin-resistant methicillin-

722   resistant Staphylococcus aureus (MRSA) clinical isolates. PloS one *8*, e67398.

723   10.1371/journal.pone.0067398.

724 32. Mishra, N.N., Bayer, A.S., Weidenmaier, C., Grau, T., Wanner, S., Stefani, S., Cafiso, V.,

725   Bertuccio, T., Yeaman, M.R., Nast, C.C., and Yang, S.J. (2014). Phenotypic and genotypic

726   characterization of daptomycin-resistant methicillin-resistant Staphylococcus aureus

727   strains: relative roles of mprF and dlt operons. PloS one *9*, e107426.

728   10.1371/journal.pone.0107426.

729 33. Monteiro, J.M., Fernandes, P.B., Vaz, F., Pereira, A.R., Tavares, A.C., Ferreira, M.T.,

730   Pereira, P.M., Veiga, H., Kuru, E., VanNieuwenhze, M.S., et al. (2015). Cell shape

731   dynamics during the staphylococcal cell cycle. Nat Commun *6*, 8055.

732   10.1038/ncomms9055.

733 34. Wang, Y., Hu, M., Liu, Q., Qin, J., Dai, Y., He, L., Li, T., Zheng, B., Zhou, F., Yu, K., et al.

734   (2016). Role of the ESAT-6 secretion system in virulence of the emerging community-

735   associated Staphylococcus aureus lineage ST398. Sci Rep *6*, 25163. 10.1038/srep25163.

736 35. Garces, A., Atmakuri, K., Chase, M.R., Woodworth, J.S., Krastins, B., Rothchild, A.C.,

737   Ramsdell, T.L., Lopez, M.F., Behar, S.M., Sarracino, D.A., and Fortune, S.M. (2010). EspA

738   acts as a critical mediator of ESX1-dependent virulence in Mycobacterium tuberculosis by

739   affecting bacterial cell wall integrity. PLoS pathogens *6*, e1000957.

740   10.1371/journal.ppat.1000957.

741 36. Ates, L.S., van der Woude, A.D., Bestebroer, J., van Stempvoort, G., Musters, R.J., Garcia-

742   Vallejo, J.J., Picavet, D.I., Weerd, R., Maletta, M., Kuijl, C.P., et al. (2016). The ESX-5

743   System of Pathogenic Mycobacteria Is Involved In Capsule Integrity and Virulence through

744   Its Substrate PPE10. PLoS pathogens *12*, e1005696. 10.1371/journal.ppat.1005696.

745 37. Nath, Y., Ray, S.K., and Buragohain, A.K. (2018). Essential role of the ESX-3 associated

746   eccD3 locus in maintaining the cell wall integrity of Mycobacterium smegmatis.

International journal of medical microbiology : IJMM *308*, 784-795.

10.1016/j.ijmm.2018.06.010.

38.   Bosserman, R.E., and Champion, P.A. (2017). Esx Systems and the Mycobacterial Cell

Envelope: What's the Connection? Journal of bacteriology *199*. 10.1128/JB.00131-17.

39.   Bottai, D., Di Luca, M., Majlessi, L., Frigui, W., Simeone, R., Sayes, F., Bitter, W., Brennan,

M.J., Leclerc, C., Batoni, G., et al. (2012). Disruption of the ESX-5 system of

Mycobacterium tuberculosis causes loss of PPE protein secretion, reduction of cell wall

integrity and strong attenuation. Molecular microbiology *83*, 1195-1209. 10.1111/j.1365-

2958.2012.08001.x.

40.   Bramkamp, M., and Lopez, D. (2015). Exploring the existence of lipid rafts in bacteria.

Microbiol Mol Biol Rev *79*, 81-100. 10.1128/MMBR.00036-14.

41.   Grein, F., Muller, A., Scherer, K.M., Liu, X., Ludwig, K.C., Klockner, A., Strach, M., Sahl,

H.G., Kubitscheck, U., and Schneider, T. (2020). Ca(2+)-Daptomycin targets cell wall

biosynthesis by forming a tripartite complex with undecaprenyl-coupled intermediates and

membrane lipids. Nat Commun *11*, 1455. 10.1038/s41467-020-15257-1.

42.   Rostovtseva, T.K., Aguilella, V.M., Vodyanoy, I., Bezrukov, S.M., and Parsegian, V.A.

(1998). Membrane surface-charge titration probed by gramicidin A channel conductance.

Biophys J *75*, 1783-1792. 10.1016/S0006-3495(98)77620-9.

43.   Bobrovskyy, M., Chen, X., and Missiakas, D. (2023). The Type 7b Secretion System of S.

aureus and Its Role in Colonization and Systemic Infection. Infection and immunity *91*,

e0001523. 10.1128/iai.00015-23.

44.   Shariati, A., Dadashi, M., Moghadam, M.T., van Belkum, A., Yaslianifard, S., and Darban-

Sarokhalil, D. (2020). Global prevalence and distribution of vancomycin resistant,

vancomycin intermediate and heterogeneously vancomycin intermediate Staphylococcus

aureus clinical isolates: a systematic review and meta-analysis. Sci Rep *10*, 12689.

10.1038/s41598-020-69058-z.

45.   Goodman, J.K., Zampronio, C.G., Jones, A.M.E., and Hernandez-Fernaud, J.R. (2018).

Updates of the In-Gel Digestion Method for Protein Analysis by Mass Spectrometry.

Proteomics *18*, e1800236. 10.1002/pmic.201800236.

46.   Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015).

limma powers differential expression analyses for RNA-sequencing and microarray

studies. Nucleic Acids Res *43*, e47. 10.1093/nar/gkv007.

 Figure Legends

Figure 1. T7SS mutants are more sensitive to membrane acting drugs.

(A) Growth curves of WT USA300 JE2, Δ*esxC* and Δ*essC* in TSB in the presence of 5 µg/ml

daptomycin and 1 mM $CaCl_2$. Mean ± standard error of the mean (SEM) is shown, N = 3. **(B)** The

area under the curve (AUC) of biological replicates in (A) was calculated, mean ± standard

deviation (SD) plotted, * P ≤ 0.05, ** P ≤ 0.01 **(C)** CFU/ml was calculated after 8 h daptomycin

treatment. N = 3, mean ± SD, *** P≤0.001 using a one-way ANOVA. **(D)** Growth curves of WT

pOS1, Δ*esxC* pOS1 and complemented strain Δ*esxC* pOS1-*esxC* in TSB supplemented with 5

µg/ml daptomycin and 1 mM $CaCl_2$. Mean ± SEM shown, N = 3.; AUC was calculated and the

mean ± SD plotted in **(E)** * P ≤ 0.05. **(F)** CFU/ml from a killing assay in the presence of 10 µg/ml

daptomycin and 1mM $CaCl_2$ at timepoints indicated. **(G)** CFU/ml from killing assay using WT

USA300 JE2, Δ*esxC* and Δ*esxC* complemented strains of in the presence of 10 µg/ml daptomycin

and 1mM $CaCl_2$ at timepoints indicated, mean ± SD, N = 3, * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001

using a two-way ANOVA with Tukey multiple comparison test. (H) Growth curves of USA300 JE2

WT, Δ*essC* and Δ*esxC* in the presence of 2 µg/ml gramicidin, Mean ± SEM, N = 3 (I) Growth

curves of USA300 JE2 WT, Δ*essC* and Δ*esxC* in the presence of 1 µg/ml bithionol. Mean ± SEM,

N = 3.

Figure 2. T7SS affects membrane fluidity and protein localisation.

A) The membrane fluidity of WT USA300 JE2 and Δ*esxC* as measured with a pyrene decanoic

acid staining-based assay. Mean ± SEM are shown, N = 6, * indicates P < 0.05 using a two-tailed

t-test. B) Widefield micrographs of *S. aureus* WT and Δ*esxC* after growth in TSB to OD_{600} of 1.0

in the presence of the lipophilic dye DiIC12. Images are representative of 4 independent

experiments. The DiIC12 fluorescence of 80 bacterial clusters from different fields per strain was

quantitated with ImageJ. Means ± SD are shown, N = 4; **** P < 0.0001, using a Kruskal-Wallis

rank test. C) One point measurement of fluidity in logarithmic phase *S. aureus* WT and Δ*esxC*,

grown in TSB at 37°C. Laurdan generalised polarisation (GP) was calculated using the equation

$(I_{460} – I_{500})/ (I_{460} – I_{500})$ where I refers to the fluorescent intensity at the emission wavelength. *** P

= 0.0003 using an unpaired t-test with Welch's correction, N=3, mean ± SD. D) Representative

widefield microscopy images of FloA-eYFP expressing WT and T7SS mutant strains.

Deconvolved images were acquired using a Deltavision microscope. N=3. Quantification of the

number of foci per cell (n = 100 per mutant).

Figure 3. T7SS mutants have distinct surface morphologies.

A) Representative SEM images of *S. aureus* WT USA300 JE2, *ΔessC*, *ΔesxC*, and *ΔesxD*

bacteria grown to early logarithmic phase. B) Representative SEM images of *S. aureus* USA300

JE2 WT pOS1, *ΔesxC* pOS1, *ΔesxC* pOS1 *esxC* grown to early logarithmic phase.

Figure 4. T7SS impacts cell wall synthesis.

(A) Representative widefield micrographs of *S. aureus* cultures either grown to logarithmic phase

in the presence of 25 µM HADA or after 10-minute treatment with HADA once OD_{600} of 1 was

reached. (B) Quantification of fluorescence in cells grown to logarithmic phase with HADA. (C)

Quantification of fluorescence in cells treated with HADA for 10 min Quantification performed on

five images each from 3 independent experiments. ** = P< 0.01, and **** =P < 0.0001 by One-

way ANOVA.

Figure 5. T7SS mutant membranes bind more to daptomycin.

Quantitative binding assay of cytochrome C to WT USA300 JE2, *ΔesxC* and *ΔessC* and WT

USA300 LAC, *ΔesxA*, *ΔesxB* mutants (A, C). The CFU of the starting inoculum was calculated for

each mutant (B, D). Mean ± SD shown (N = 3), ***P < 0.001 using a one-way ANOVA. E)

Fluorescence measurements of daptomycin-BODIPY binding to *S. aureus.* The excitation

wavelength was 488nm and emission wavelength was 530nm. N=3, mean ± SD. P = 0.0067,

using an unpaired t-test. F) Representative micrograph of daptomycin-BODIPY binding to *S.*

aureus.

Figure 6. T7SS mutant membranes are more permeable in the presence of daptomycin

A) Representative fluorescent micrographs of *S. aureus* T7SS mutants treated with daptomycin.

Cell membranes were stained by FM1-43 and imaged using FITC filter channel, PI was imaged

using Texas red filter channel and the DAPI was imaged using DAPI filter set. B) Images were

quantified using FIJI. Data presented is from 3 independent experiments. 10 images from each

experiment were quantified and normalised to the number of cells in the field of view. Statistical

analysis performed using an ANOVA * $P \leq 0.05$, ** $P \leq 0.01$.

Figure 7. Presence of T7SS affects intracellular bacterial killing by daptomycin.

A) CFU counts at various timepoints during the infection assay. 0 h represents the initial inoculum,

1 h represents the CFU counts that have infected the cell and 4 h and 24 h represent the survival

of *S. aureus* in the presence of daptomycin, N = 3, **$P<0.05$, two-way ANOVA. (B) Confocal image

quantification from five images each from 3 independent experiments. Mean ± SD of intensity, is

shown, ***$P<0.01$ using a one-way ANOVA. (C) Representative confocal micrographs of cells after

1 h infection or 24 h post-infection. DAPI (blue), F-actin (red), and *S. aureus* (green, white arrows).

Figure 8. EsxC mutants are more sensitive to daptomycin *in vivo*.

Skin-abscess model mice (n=8, 2 lesions/animal) were infected with WT USA300 and Δ*esxC*

mutant followed by treatment with 10 mg/kg daptomycin. (A) CFU/g skin tissue was calculated 24

h post infection. Graphs show mean ± SD, P = 0.0007 using one-way ANOVA (B) CFU//g was

calculated 72 h post infection (n=5, 2 lesions/animal), mean ± SD shown, P = 0.0020 using a one-

way ANOVA. (C) TNF-α levels: Two-way ANOVA revealed a significant effect on treatment groups

(P <0.01). Results of multiple comparisons shown on graph. (D) IL-1β levels, but significant effect

seen between treatment groups (P<0.01) using two-way ANOVA. Graphs show mean ± SD, n=5.

894 **Table 1.** Proteins significantly changed in Δ*esxC* and Δ*essC* mutant membrane preparations

895 relative to WT USA300 JE2. Proteins common to both mutants are in bold.

	Uniprot ID	log$_2$ fold change	Adjusted P value	Description
USA300 JE2 Δ*esxC* vs USA300 JE2 WT	A0A0H2XJR2	-5.46	6.21E-08	Pyridine nucleotide-disulfide oxidoreductase
	A0A0H2XJ15	**-5.34**	**3.80E-09**	**TPR domain protein**
	A0A0H2XIV9	-5.33	1.49E-07	T7SS protein, EssD
	A0A0H2XJ90	-4.82	6.92E-09	D-isomer specific 2-hydroxyacid dehydrogenase family protein
	A0A0H2XF36	**3.62**	**2.14E-08**	**Uncharacterised protein**
	A0A0H2XJV3	**3.67**	**2.26E-08**	**ABC transporter, ATP-binding protein**
	A0A0H2XGS2	3.70	1.71E-08	Putative sensor histidine kinase
	A0A0H2XG87	**4.01**	**1.59E-08**	**Bifunctional ligase/repressor BirA**
	A0A0H2XH85	4.14	1.04E-08	Cobalamin synthesis protein/P47K family protein
	A0A0H2XER9	4.38	9.36E-09	LysR family regulatory protein
	A0A0H2XEB7	**5.25**	**2.39E-07**	**Integral membrane protein**
	A0A0H2XJ96	**5.27**	**1.04E-08**	**NADH-dependent flavin oxidoreductase**
	A0A0H2XHU5	5.97	2.55E-09	Putative membrane protein
USA300 JE2 Δ*essC* vs USA300 JE2 WT	A0A0H2XEV3	-8.65	3.31E-08	T7SS protein, EssC
	A0A0H2XJ15	-5.34	5.71E-09	TPR domain protein
	A0A0H2XH73	-4.56	2.19E-08	NAD-dependent epimerase-domain-containing protein
	A0A0H2XFR1	3.33	2.52E-08	Energy-coupling factor transporter transmembrane protein EcfT
	A0A0H2XJF9	3.47	2.19E-08	Putative deoxyribodipyrimidine photolyase
	A0A0H2XF36	3.57	2.19E-08	Uncharacterised protein
	A0A0H2XJV3	3.59	2.42E-08	ABC transporter, ATP-binding protein
	A0A0H2XG87	3.93	1.82E-08	Bifunctional ligase/repressor BirA
	A0A0H2XGW9	4.26	9.38E-09	Deoxynucleoside kinase family protein
	A0A0H2XIK4	4.46	8.12E-09	ornithine aminotransferase
	A0A0H2XHF0	4.56	8.12E-09	Carboxylic ester hydrolase
	A0A0H2XHR8	4.80	8.12E-09	DJ-1/PfpI domain-containing protein

A0A0H2XEB7	4.82	4.09E-07	Integral membrane protein
A0A0H2XJ96	4.83	1.82E-08	NADH-dependent flavin oxidoreductase
A0A0H2XEH0	8.22	5.27E-10	Alpha, alpha-phosphotrehalase

Table 2. Strains and Plasmids used in this study

Strain or Plasmid	Description	Source or Reference
Staphylococcus aureus		
USA300 LAC JE2	Plasmid-cured USA300 LAC	BEI Resources (NARSA)
USA300 JE2 *ΔessC*	USA300 LAC JE2 defective for EssC	(Kengmo Tchoupa *et al.,* 2020)
USA300 JE2 *ΔesxC*	USA300 LAC JE2 defective for EsxC	(Kengmo Tchoupa *et al.,* 2020)
USA300 LAC	Community acquired MRSA	(Bae and Schneewind, 2006)
USA300 LAC *ΔesxA*	USA300 LAC defective for EsxA	(Korea *et al.,* 2014)
USA300 LAC Δ*esxB*	USA300 LAC defective for EsxB	(Korea *et al.,* 2014)
Newman	Methicillin-sensitive *S. aureus*	(Bae and Schneewind, 2006)
Newman *ΔesxA*	Newman defective for EsxA	(Korea *et al.,* 2014)
Newman Δ*esxB*	Newman defective for EsxB	(Korea *et al.,* 2014)
RN6390	NCTC8325 derivative, *ΔrbsU, ΔtcaR,* cured of φ11, φ12, and φ13	Tracy Palmer (Kneuper *et al.,* 2014)
RN6390 *ΔessC*	RN6390 defective for EssC	Tracy Palmer (Kneuper *et al.,* 2014)
RN6390 *ΔesxC*	RN6390 defective for EsxC	Tracy Palmer (Kneuper *et al.,* 2014)
Plasmids		
pOS1	Empty vector for complementation	(Kengmo Tchoupa *et al.,* 2020)
pOS1-*esxC*	*esxC* complementation vector	(Kengmo Tchoupa *et al.,* 2020)

Supplementary Figure legends and tables

Figure S1. Growth curves of (A) USA300 JE2 WT, *ΔessC, ΔesxC* and *ΔesxD* (B) USA300JE2 WT

pOS1, *ΔesxC* pOS1 and complemented strain *ΔesxC* pOS1-*esxC* C) RN6390 WT, *ΔessC* and

ΔesxC (D) Newman WT and *ΔesxA* and *ΔesxB* mutants (E) USA300 LAC WT and *ΔesxA* and

ΔesxB grown in TSB at 37ºC. OD_{600} measured every 15 min. Mean ± SEM plotted on graph, N=3.

Figure S2. Growth curves of (A) USA300 JE2 WT, and *ΔesxD*, (C) RN6390 WT, *ΔessC* and

ΔesxC, (E) USA300 *ΔesxA* and *ΔesxB* and (G) Newman *ΔesxA* and *ΔesxB* in TSB in the presence

of 5 µg/ml daptomycin and 1 mM $CaCl_2$. Mean ± SEM are shown, N = 3. The AUC was calculated

and the mean ± SD was plotted for the growth curve shown on the left (B, D, F, H). * P ≤ 0.05, **

P ≤ 0.01.

Figure S3. Growth curves of USA300 JE2 WT, *ΔessC* and *ΔesxC* in the presence of (A) 4 µg/ml

ciprofloxacin (B) 0.5 µg/ml mitomycin C. (C) 1 µg/ml oxacillin. (D) 8 µg/ml vancomycin. For all

growth curves mean and SEM are shown, N = 3.

Figure S4. Scanning electron micrographs of *S. aureus* RN6390, USA300, and Newman T7SS

mutants.

Figure S5 Transmission electron micrographs of *S. aureus* USA300 JE2 WT, *ΔessC* and *ΔesxC*

grown to early logarithmic phase.

Figure S6 (A) Growth curves of WT, ΔessC and *ΔesxC* in DMEM + FBS. **(B)** Growth curves of

WT, *ΔessC* and ΔesxC in DMEM + FBS in the presence of 1µg/ml daptomycin and 1 mM $CaCl_2$.

Graphs show mean and SEM of three independent experiments.

Figure S7 The area of lesions was measured and calculated daily from mice infected with *S.*

aureus WT or T7SS mutants. Area of lesions was measured and calculated daily. Graphs show

mean ± SD. Two-way ANOVA between groups revealed no significant differences between

treatments.

Table S1 Secreted proteins significantly changed in Δ*esxC* and Δ*essC* mutants relative to WT

USA300 JE2.

Figure 1. T7SS mutants are more sensitive to membrane acting drugs.
(A) Growth curves of WT USA300 JE2, Δ*esxC* and Δ*essC* in TSB in the presence of 5 µg/ml daptomycin and 1 mM CaCl$_2$. Mean ± standard error of the mean (SEM) is shown, N = 3. **(B)** The area under the curve (AUC) of biological replicates in (A) was calculated, mean ± standard deviation (SD) plotted, * P ≤ 0.05, ** P ≤ 0.01 **(C)** CFU/ml was calculated after 8 h daptomycin treatment. N = 3, mean ± SD, *** P≤0.001 using a one-way ANOVA. **(D)** Growth curves of WT pOS1, Δ*esxC* pOS1 and complemented strain Δ*esxC* pOS1-*esxC* in TSB supplemented with 5 µg/ml daptomycin and 1 mM CaCl$_2$. Mean ± SEM shown, N = 3.; AUC was calculated and the mean ± SD plotted in **(E)** * P ≤ 0.05. **(F)** CFU/ml from a killing assay in the presence of 10 µg/ml daptomycin and 1mM CaCl$_2$ at timepoints indicated. **(G)** CFU/ml from killing assay using WT USA300 JE2, Δ*esxC* and Δ*esxC* complemented strains of in the presence of 10 µg/ml daptomycin and 1mM CaCl$_2$ at timepoints indicated, Mean ± SD,N = 3, * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001 using a two-way ANOVA with Tukey multiple comparison test. (H) Growth curves of USA300 JE2 WT, Δ*essC* and Δ*esxC* in the presence of 2 µg/ml gramicidin, Mean ± SEM, N = 3 (I) Growth curves of USA300 JE2 WT, Δ*essC* and Δ*esxC* in the presence of 1 µg/ml bithionol. Mean ± SEM, N = 3.

Figure 2. T7SS affects membrane fluidity and protein localisation.

A) The membrane fluidity of WT USA300 JE2 and ΔesxC as measured with a pyrene decanoic acid staining-based assay. Mean ± SEM are shown, N = 6, * indicates P < 0.05 using a two-tailed t-test. B) Widefield micrographs of *S. aureus* WT and ΔesxC after growth in TSB to OD_{600} of 1.0 in the presence of the lipophilic dye DiIC12. Images are representative of 4 independent experiments. The DiIC12 fluorescence of 80 bacterial clusters from different fields per strain was quantitated with ImageJ. Means ± SD are shown, N = 4; **** P < 0.0001, using a Kruskal-Wallis rank test. C) One point measurement of fluidity in logarithmic phase *S. aureus* WT and *ΔesxC*, grown in TSB at 37ºC. Laurdan generalised polarisation (GP) was calculated using the equation $(I_{460} - I_{500})/ (I_{460} - I_{500})$ where I refers to the fluorescent intensity at the emission wavelength. *** P = 0.0003 using an unpaired t-test with Welch's correction, N=3, mean ± SD. D) Representative widefield microscopy images of FloA-eYFP expressing WT and T7SS mutant strains. Deconvolved images were acquired using a Deltavision microscope. N=3. Quantification of the number of foci per cell (n = 100 per mutant).

Figure 3. T7SS mutants have distinct surface morphologies.
A) Representative SEM images of *S. aureus* WT USA300 JE2, *ΔessC*, *ΔesxC*, and *ΔesxD* bacteria grown to early logarithmic phase. B) Representative SEM images of *S. aureus* USA300 JE2 WT pOS1, *ΔesxC* pOS1, *ΔesxC* pOS1 *esxC* grown to early logarithmic phase.

Figure 4. T7SS impacts cell wall synthesis.

(A) Representative widefield micrographs of *S. aureus* cultures either grown to logarithmic phase in the presence of 25 μM HADA or after 10-minute treatment with HADA once OD_{600} of 1 was reached. (B) Quantification of fluorescence in cells grown to logarithmic phase with HADA. (C) Quantification of fluorescence in cells treated with HADA for 10 minutes. Quantification performed on five images each from 3 independent experiments. ** = P< 0.01, and **** =P < 0.0001 by One-way ANOVA

Figure 5. T7SS mutant membranes bind more to daptomycin.
Quantitative binding assay of cytochrome C to WT USA300 JE2, Δ*esxC* and Δ*essC* and WT USA300 LAC, *ΔesxA*, Δ*esxB* mutants (A, C). The CFU of the starting inoculum was calculated for each mutant (B, D). Mean ± SD shown (N = 3), ***P < 0.001 using a one-way ANOVA. E) Fluorescence measurements of daptomycin-BODIPY binding to *S. aureus.* The excitation wavelength was 488nm and emission wavelength was 530nm. N=3, mean ± SD. P = 0.0067, using an unpaired t-test. F) Representative micrograph of daptomycin-BODIPY binding to *S. aureus.*

A

	WT Untreated	WT Daptomycin	ΔessC Untreated	ΔessC Daptomycin	ΔesxC Untreated	ΔesxC Daptomycin
BF						
FM1-43						
PI						
DAPI						

B

Figure 6. T7SS mutant membranes are more permeable in the presence of daptomycin
A) Representative fluorescent micrographs of *S. aureus* T7SS mutants treated with daptomycin. Cell membranes were stained by FM1-43 and imaged using FITC filter channel, PI was imaged using Texas red filter channel and the DAPI was imaged using DAPI filter set. B) Images were quantified using FIJI. Data presented is from 3 independent experiments. 10 images from each experiment were quantified and normalised to the number of cells in the field of view. Statistical analysis performed using an ANOVA * P ≤ 0.05, ** P ≤ 0.01.

Figure 7. Presence of T7SS affects intracellular bacterial killing by daptomycin.
A) CFU counts at various timepoints during the infection assay. 0 h represents the
initial inoculum, 1 h represents the CFU counts that have infected the cell and 4 h and
24 h represent the survival of *S. aureus* in the presence of daptomycin, N = 3,
**P<0.05, two-way ANOVA. (B) Confocal image quantification from five images each
from 3 independent experiments. Mean ± SD of intensity, is shown, ***P<0.01 using a
one-way ANOVA. (C) Representative confocal micrographs of cells after 1 h infection
or 24 h post-infection. DAPI (blue), F-actin (red), and *S. aureus* (green, white arrows).

Figure 8. EsxC mutants are more sensitive to daptomycin *in vivo*.
Skin-abscess model mice (n=8, 2 lesions/animal) were infected with WT USA300 and Δ*esxC* mutant followed by treatment with 10 mg/kg daptomycin. (A) CFU/g skin tissue was calculated 24 h post infection. Graphs show mean ± SD, P = 0.0007 using one-way ANOVA (B) CFU//g was calculated 72 h post infection (n=5, 2 lesions/animal), mean ± SD shown, P = 0.0020 using a one-way ANOVA. (C) TNF-α levels: Two-way ANOVA revealed a significant effect treatment groups (P <0.01). Results of multiple comparisons shown on graph. (D) IL-1β levels , but significant effect seen between treatment groups (P<0.01) using two-way ANOVA. Graphs show mean ± SD, n=5.

Fig S1

Figure S1. Growth curves of (A) USA300 JE2 WT, *ΔessC, ΔesxC* and *ΔesxD* (B) USA300JE2 WT pOS1, *ΔesxC* pOS1 and complemented strain *ΔesxC* pOS1-*esxC* C) RN6390 WT, *ΔessC* and *ΔesxC* (D) Newman WT and *ΔesxA* and *ΔesxB* mutants (E) USA300 LAC WT and *ΔesxA* and *ΔesxB* grown in TSB at 37ºC. OD_{600} measured every 15 minutes. Mean ± SEM plotted on graph, N=3.

Fig S2

Figure S2. Growth curves of (A) USA300 JE2 WT, and Δ*esxD*, (C) RN6390 WT, Δ*essC* and Δ*esxC*, (E) USA300 Δ*esxA* and Δ*esxB* and (G) Newman Δ*esxA* and Δ*esxB* in TSB in the presence of 5 μg/ml daptomycin and 1 mM CaCl$_2$. Mean ± SEM are shown, N = 3. The AUC was calculated and the mean ± SD was plotted for the growth curve shown on the left (B, D, F, H). * P ≤ 0.05, ** P ≤ 0.01.

Fig S3

Figure S3. Growth curves of USA300 JE2 WT, *ΔessC* and *ΔesxC* in the presence of (A) 4 μg/ml ciprofloxacin (B) 0.5 μg/ml mitomycin C. (C) 1 μg/ml oxacillin. (D) 8 μg/ml vancomycin. For all growth curves mean and SEM are shown, N = 3.

Fig S4

Figure S4. Scanning electron micrographs of *S. aureus* RN6390, USA300, and Newman T7SS mutants**.**

Fig S5

Figure S5. Transmission electron micrographs of *S. aureus* USA300 JE2 WT, Δ*essC* and Δ*esxC* grown to early logarithmic phase

Fig S6

Figure S6. (A) Growth curves of WT, ΔessC and Δ*esxC* in DMEM + FBS. **(B)** Growth curves of WT, Δ*essC* and ΔesxC in DMEM + FBS in the presence of 1μg/ml daptomycin and 1 mM CaCl$_2$. Graphs show mean and SEM of three independent experiments.

Fig S7

Figure S7 The area of lesions was measured and calculated daily from mice infected with *S. aureus* WT or T7SS mutants. Area of lesions was measured and calculated daily. Graphs show mean ± SD. Two-way ANOVA between groups revealed no significant differences between treatments.

Table S1. Secreted proteins significantly changed in Δ*esxC* and Δ*essC* mutants relative to WT USA300 JE2. Proteins common to both mutants are in bold.

		Uniprot ID	log$_2$ fold change	Adjusted P value	Description
JE2 WT	USA300 JE2 ΔesxC vs USA300	**A0A0H2XIV9**	**-4.27**	**7.11E-10**	**T7SS protein EssD**
		A0A0H2XHT7	**-2.78**	**7.96E-09**	**Phosphocarrier protein HPr**
		A0A0H2XH98	**-1.75**	**1.34E-07**	**Hydrolase, MutT/nudix family**
		A0A0H2XIK2	**-1.71**	**1.34E-07**	**T7SS protein EsxC**
		A0A0H2XI99	**-1.45**	**0.003045**	**T7SS protein EsxA**
		A0A0H2XFA5	-1.28	4.03E-05	Putative GTP-binding protein
		A0A0H2XEF1	**1.55**	**3.40E-06**	**2-oxoisovalerate dehydrogenase**
		Q2FEF1	**1.63**	**0.000836**	**Isopentenyl-diphosphate delta-isomerase**
		A0A0H2XKB6	**1.90**	**0.00012**	**Uncharacterized protein**
		A0A0H2XH53	2.58	0.042723	Uncharacterized protein
	USA300 JE2 ΔessC vs USA300 JE2 WT	A0A0H2XI99	-6.59779	1.63E-07	T7SS protein EsxA
		A0A0H2XIV9	-4.27062	7.11E-10	T7SS protein EssD
		A0A0H2XHT7	-2.77719	7.96E-09	Phosphocarrier protein HPr
		A0A0H2XKB6	-2.39392	2.34E-05	Uncharacterized protein
		A0A0H2XH98	-1.74886	1.34E-07	Hydrolase, MutT/nudix family
		A0A0H2XIK2	-1.70544	1.34E-07	T7SS protein EsxC
		A0A0H2XJ55	-1.24877	0.000168	Uncharacterized protein
		Q2FHI5	-1.04906	0.015225	ATP-dependent protease subunit HslV
		Q2FFT2	0.777989	0.026729	Serine protease
		A0A0H2XEF1	1.394088	5.08E-06	2-oxoisovalerate dehydrogenase
		A0A0H2XHA0	1.470262	0.041559	Uncharacterized protein
		A0A0H2XG38	1.580427	1.63E-07	Uncharacterized protein
		Q2FEF1	1.633554	0.000707	Isopentenyl-diphosphate delta-isomerase