The staphylococcal type VII secretion system effector EsxC impacts daptomycin sensitivity

through controlling bacterial cell envelope integrity

Victoria Smith[1], Giridhar Chandrasekharan[1], Robeena Farzand[1], Kate E. Watkins[1], Ramon Garcia

Marset[1$], Jeannifer Yap[2], Sebastien Perrier[3], Arnaud Kengmo Tchoupa[1+], and Meera

Unnikrishnan[1*].

1. Division of Biomedical Sciences, Warwick Medical School, University of Warwick,

Coventry, CV4 7AL, UK.

2. Department of Chemistry, University of Warwick, Coventry, CV4 7AL, UK

3. Department of Chemistry and Warwick Medical School, University of Warwick, Coventry,

CV4 7AL, UK.

$ current address: Centre for Urological Biology, Division of Medicine, University College

London, WC1E 6BT London, UK.

+ current address: Cluster of Excellence EXC 2124 Controlling Microbes to Fight Infections,

University of Tübingen, 72076 Tübingen, Germany

*Corresponding author m.unnikrishnan@warwick.ac.uk

Abstract

The human pathogen *Staphylococcus aureus* encodes a specialised type VII secretion system (T7SS), which plays an important role in bacterial virulence during infection. However, the functions the T7SS during infection and in bacterial physiology remain unclear. Here we demonstrate that *S. aureus* strains lacking the the T7SS effector EsxC (Δ*esxC*) was highly sensitive to the important last resort drug, daptomycin, as well as other membrane-targeting antibiotics, including gramicidin and bithionol. To understand how EsxC mediates increased antibiotic sensitivity, we investigated its functions in the staphylococcal cell envelope. Scanning electron microscopy analysis of an *esxC* mutant revealed a distinct cell surface morphology. Interestingly, Δ*esxC* displayed a decrease in membrane fluidity, altered membrane protein profiles and altered cell wall synthesis. The *esxC* mutant demonstrated enhanced daptomycin binding which correlated with the increased negative charge of mutant membranes. Calcium ions, which can bind membranes affecting charge, impacted growth of Δ*esxC* and sensitivity to daptomycin, suggesting that EsxC may modulate calcium binding to membranes. Furthermore, the *esxC* mutant displayed a heightened susceptibility to daptomycin during intracellular infection, and in a murine skin infection model. Thus, our data show that the T7SS effector EsxC impacts sensitivity of *S. aureus* to membrane-acting drugs such as daptomycin through modulation of cell membrane integrity, indicating its potential as a drug target.

Author Summary T7SS has a range of functions in bacteria including specific roles in bacterial physiology including DNA uptake, membrane integrity and bacterial development. In *S. aureus* T7SS has been shown to be critical for bacterial virulence, intra-species competition and in host cell interactions, although their functions in bacterial physiology are not clear. Here we report a role of the staphylococcal T7SS effector EsxC in the modulation of the cell membrane and surface integrity, which impacts the activity of membrane targeting drugs like daptomycin. Our data indicate that targeting this system could potentially enhance activity of existing therapeutic agents.

Introduction

S. aureus is a highly versatile pathogen and a commensal bacterium which causes infections in both humans and animals. Staphylococcal infections can be nosocomial or community-acquired, and can range from mild skin abscesses to severe life-threatening infections including septic shock and bacteraemia (1). *S. aureus* infections place a huge cost burden on healthcare sectors across the world, especially due to the occurrence of antibiotic-resistant strains, such as methicillin-resistant *S. aureus* (MRSA) and strains with varying levels of vancomycin resistance (2, 3).

S. aureus possesses an array of surface and secreted factors that help the bacterium colonise the host and establish an infection (4). One such virulence-associated factor is the type VII secretion system (T7SS), which is widely found in Gram positive bacteria (5). The T7SSb in *S. aureus* and other firmicutes is distinct from the systems in mycobacteria (T7SSa) although both are characterised by small substrates containing the WXG motif, such as EsxA and EsxB (6, 7). The T7SS in *S. aureus* is modular in nature, with heterogenous expression between strains (8, 9). Typically, in commonly studied *S. aureus* strains including USA300, Newman and RN6390, the T7SS locus comprises of integral membrane proteins (EsaA, EssA, EssB and EssC), cytosolic proteins (EsaB and EsaG), several secreted substrates (EsxA, EsxB, EsxC, EsxD and EsaD) and chaperones (EsaE) (7, 10). EssC is the central transporter which is responsible for export of substrates and co-dependent export of secreted substrates has been reported (6, 11). While specific interactions between the different T7SS components have been described, the overall structure of the secretion apparatus is poorly defined for the *S. aureus* T7SS (10, 11). Assembly of a functional system is dependent on the membrane microdomain protein, flotillin A (FloA)(12).

Recent studies have reported a role for T7SS in modulation of intraspecies competition. A T7SS

substrate EsaD with nuclease activity mediates killing of *S. aureus* strains that lack the anti-toxin

EsaG (13). An LXG-domain containing protein, TspA, encoded distally to the T7SS cluster shows

T7SS-dependent toxic activity, which can be neutralized by an antitoxin TsaI (14). Some

substrates modulate host interactions during infection; EsxA modulates host cell death and EsaE

elicits cytokine production during infection (15, 16). However, effectors of this system are yet to

be defined in many *S. aureus* strains and their specific functions remain elusive.

T7SS in other bacterial species has been described to support bacterial functions like DNA

transfer, membrane integrity and spore development, although for *S. aureus*, it is not clear if T7SS

is important in staphylococcal physiology (17-19). Our recent study on the T7SS substrate EsxC

showed that a mutant lacking this protein was more sensitive to unsaturated host-derived fatty

acids (20). T7SS defects compromised membrane integrity and induced oxidative stress in

presence of antimicrobial fatty acids, indicating a role for these proteins in membrane

homeostasis. Here we report that the absence of EsxC causes an increased sensitivity to

membrane targeting drugs like daptomycin, both during infection *in vitro* and *in vivo*. We

demonstrate distinct effects on cell surface morphology, membrane fluidity and charge, indicating

a role for T7SS in controlling membrane integrity potentially through modulating calcium binding.

Our data suggest that targeting the T7SS could augment activities of membrane-acting drugs like

daptomycin.

Results

A mutant lacking T7SS effector EsxC is more sensitive to membrane-acting drugs

The cyclic lipopeptide antibiotic, daptomycin, is a key drug used to treat resilient *S. aureus*

infections. Daptomycin binds to the cell membrane in a calcium-dependent manner (21). As our

previous studies indicated a potential impact of T7SS, in particular the effector EsxC, on

membrane homeostasis (20), we investigated whether the *S. aureus* USA300 JE2 mutants lacking

the T7SS substrate EsxC (Δ*esxC)* respond differently to membrane-acting drugs like daptomycin.

Growth of Δ*esxC* was compared to the WT USA300 JE2 strain in the presence or absence of

daptomycin. When cultured in presence of 5 µg/ml daptomycin, growth of the WT USA300 JE2

strain was delayed, but growth of USA300 JE2 Δ*esxC* was significantly impaired compared to the

WT (Fig. 1 A). No differences in growth rates were seen between the strains in tryptic soy broth

(TSB) without antibiotic. Further, a 10-fold decrease in colony forming unit counts for the mutant

was also observed in the presence of daptomycin (Fig. 1 B). Complementing Δ*esxC* with pOS1-

esxC restored WT phenotype (Fig. 1 C, D), although daptomycin did not inhibit the pOS1 empty

plasmid-bearing Δ*esxC* to the same extent as Δ*esxC* without plasmid. This could be attributed to

differences in growth profiles of pOS1 containing WT and mutant strains, possibly due to the

effects of chloramphenicol exposure prior to performing growth assays. Next, a daptomycin killing

assay was employed where logarithmic phase bacteria were treated for 2 h with daptomycin.

Δ*esxC* was more sensitive to daptomycin in comparison to the WT in this killing assay, with the

Δ*esxC* pOS1-*esxC* strain reversing daptomycin sensitivity (Fig. 1 E, F). When we stained bacteria

with propidium iodide (PI), a fluorescent intercalating agent which cannot pass through the

membranes of viable cells, cytosols of daptomycin-treated Δ*esxC* showed a higher intensity of PI

compared to the WT, as quantified by fluorescence microscopy (Fig. 2 A, B). Staining with the

lipid staining dye FM1-43 was of similar intensity for WT and Δ*esxC*, although it appeared less

ordered in the daptomycin-treated Δ*esxC* sample (Fig. 2 A, B). Thus, Δ*esxC* demonstrated

increased membrane permeability in the presence of daptomycin.

To investigate whether other T7SS effectors had similar effects, we studied strains lacking

substrates EsxA and EsxB. We used mutants in USA300 LAC and Newman backgrounds, as

USA300 JE2 mutants in these genes were not available. A defect in growth was seen in USA300

LAC Δ*esxA* mutant compared to the USA300 LAC WT strain, while Δ*esxB* did not appear to have

a growth defect (Fig. S1 A, B). The Newman strain was more sensitive than USA300 JE2 to

daptomycin, with growth being delayed until ~14 h. Again a growth defect was observed for the

ΔesxA mutant, but not the ΔesxB mutant in presence of daptomycin (Fig. S1 C, D). Furthermore,

a USA300 JE2 mutant lacking another T7SS substrate, EsxD did not display any growth defects

in the presence of daptomycin (Fig. S1 E,F). We also studied a strain lacking and the central

transporter EssC (ΔessC) responsible for transport of T7SS effectors. As expected, the deletion

of essC significantly impaired the growth phenotype compared to the WT (Fig S1 G,H). We further

studied the effect of EsxC on daptomycin sensitivity in other strain backgrounds. The growth of

RN6390 S. aureus was impacted more by the addition of daptomycin, with the bacterial growth

initiating much later (~14 h) compared to the USA300 JE2 strain (~8 h). However, similar to the

JE2 mutants, RN6390 ΔesxC and ΔessC mutants were more susceptible than the WT to

daptomycin (Fig. S1 I, J).

To determine if the differences observed between the WT and ΔesxC were limited to daptomycin,

other membrane-targeting antimicrobials, such as gramicidin and bithionol, were tested.

Gramicidin is a polypeptide antimicrobial peptide that forms ion channels in the cell membrane

allowing diffusion of monovalent ions such as potassium (K^+) and sodium (Na^+), disrupting cell ion

concentrations. Bithionol is an inhibitor of soluble adenylyl cyclase and its membrane-disrupting

activity was associated with its ability to kill S. aureus persister cells (22). When treated with 2

μg/ml gramicidin, the WT JE2 strain did not have any growth defects, however the ΔesxC mutant

displayed slower growth (Fig. 3 A, B). A similar defect in growth was also seen in the presence of

biothionol, although only a slightly slower growth was seen for the mutant in the presence of 1

μg/ml bithionol (Fig. 3 C,D). Finally, this effect appears to be specific to membrane-acting drugs

as other drugs which target the cell wall (vancomycin, oxacillin) or DNA replication (ciprofloxacin,

mitomycin) did not differentially affect the growth of the ΔesxC (or ΔessC) as compared to the WT

(Fig. S2).

Δ*esxC* displays decreased membrane fluidity and altered membrane protein profiles

Changes in membrane fluidity can affect membrane function and antibiotic sensitivity. To study

the effect of EsxC on membrane fluidity, we assessed the membrane fluidity of the WT and Δ*esxC*

mutant using multiple assays. WT and Δ*esxC* were stained with a fluorescent membrane dye

DiI12C, which has been used detect regions of increased fluidity in bacteria (23). As reported

previously for *S. aureus*, we observe a few fluid regions in the WT, and an uneven staining of the

membrane. In contrast, a very weak staining of the membranes was observed for the Δ*esxC*

mutant (Fig. 4 A), which may indicate a decrease in membrane fluidity. An alternate assay using

pyrene decanoic acid, an excimer-forming lipid (24), was employed to measure the fluidity of WT

and Δ*esxC* membranes. Compared to the WT, the cytosolic membrane of the Δ*esxC* mutant

showed a mild but statistically significant increase in rigidity (Fig. 4 B). Finally, membrane fluidity

was measured using the fluorescent dye Laurdan (6-dodecanoyl-2-dimethylaminonaphthalene)

which is incorporated into the membrane bilayer (25). Logarithmic phase WT cells treated with

laurdan showed a lower average generalized polarisation value of 0.33 compared to 0.42 for

ΔesxC, which was reversed in the complemented strain Δ*esxC* pOS1*esxC*, suggesting that the

membrane of the *ΔesxC* mutant is more rigid in comparison to that of the WT (Fig. 4 C).

Overall, the data indicate that EsxC contributes to modulation of *S. aureus* membrane fluidity.

Previously, we reported that the mutants Δ*esxC* and Δ*essC* had distinct total protein profiles with

and without treatment with linoleic acid (20). Given the changes seen in bacterial membranes in

the absence of EsxC, we examined global changes in the protein content in the membrane.

Membrane fractions extracted from logarithmic phase (OD_{600} = 3.0) cultures of WT and Δ*esxC*

were analysed by nanoLC-ESI-MS/MS. 13 proteins were altered in abundance [log2 (fold change)

= 1.0, P < 0.05] in Δ*esxC* in relation to the WT (Table 1). Membrane proteins were increased in

abundance and EssD, a T7SS substrate and nuclease, was decreased significantly in the

membranes of Δ*esxC.* A secretome analysis from culture supernatants was also performed, which

showed a lower abundance of substrates EsxA and EssD in Δ*esxC* compared to WT (Table S1).

Thus, the *esxC* mutant show distinct changes to the overall membrane protein profiles and

secretion of certain T7SS substrates appear to be co-dependent on substrate EsxC.

EsxC displays altered cell surface morphology and decreased rate of cell wall synthesis

As EsxC appeared to be involved in membrane function, we next examined the cell surface of

Δ*esxC* using scanning electron microscopy. The cell surface of WT grown to early logarithmic

phase have the typical smooth spherical shape of *S. aureus*. However, Δ*esxC* appears deflated

with dents or hollows (Fig. 5 A). The WT phenotype was restored in Δ*esxC* pOS1-*esxC*

complemented strain (Fig. 5 B). Interestingly, this cell surface defect is not seen in all T7SS

mutants; strains lacking EsxD (Δ*esxD*) did not have a defective cell surface phenotype (Fig. S3).

S. aureus cell surface alterations were seen in USA300 (LAC) and Newman strains lacking EsxA

(Δ*esxA)*, but not in Δ*esxB* (Fig. S3). However, RN6390 strains lacking *esxC* showed very little

surface defects (Fig. S3). Finally the JE2 transporter mutant Δ*essC* showed defects similar to

Δ*esxC* (Fig S3). Differences seen in surface morphologies between WT and mutants were not

due to growth rate differences, as growth rates of all mutant strains in the different strain

backgrounds (Δ*essC*, Δ*esxC*, Δ*esxA*, Δ*esxB,* and Δ*esxD)* in TSB were similar to that of their WT

(Fig. S1). Our data suggest that the cell envelopes of mutants lacking specific T7SS components

were altered, potentially due to a defect in the cell membrane and/or cell wall structure.

To investigate if the surface structure changes were associated with a defect in cell wall structure

or synthesis, transmission electron microscopy was undertaken. However, there were no clear

differences in cell wall thickness between strains (Fig. S4). To study whether peptidoglycan

synthesis and accumulation were affected in the T7SS mutants, HADA, a fluorescent D-amino

acid was used (26). The dye was either added to a fresh subculture and grown until the bacteria

reached logarithmic phase or added for 10 min once the culture reached an OD_{600} of 1. No

differences were observed in the cells which had been grown with TSB supplemented with HADA

(Fig. 6 A, B), however addition of HADA during exponential phase growth revealed a faster

accumulation of HADA in the WT compared to *ΔesxC* (Fig. 6 A, C). This suggests that T7SS

mutants displayed a decreased rate of new peptidoglycan synthesis.

ΔesxC displays an increased negative cell surface charge and increased binding of

daptomycin

As we observed cell surface defects and changes to peptidoglycan synthesis, we next tested if

the deletion of *esxC* could alter other cell surface properties like surface charge. A cytochrome

binding assay was employed to measure surface charge; the binding of cytochrome C, a highly

positively charged protein, is directly proportional to the net negative surface charge of *S. aureus*

cells (27). A significantly higher level of binding of cytochrome C to the cell surface of Δ*esxC* was

observed compared to the USA300 JE2 WT strain (Fig. 7 A). Similarly, Δ*esxA* also showed

increased levels of binding of cytochrome C (Fig. S5). This suggests that the cell surfaces of

Δ*esxC* and Δ*esxA* are more negatively charged. Much lower levels of cytochrome C bound to

USA300 LAC WT and Δ*esxB* suggesting these cell surfaces are less negative (Fig. S5). CFU

counts from samples after cytochrome C addition showed no differences in cell numbers and

therefore it is unlikely that bacterial cell numbers influenced cytochrome C binding (Fig. 7 B, Fig

S5). These data indicate that the T7SS effectors EsxC impacts the surface charge of *S. aureus*.

S. aureus has been reported to resist daptomycin through possessing a more positive surface

charge, causing electrostatic repulsion of the positively charged daptomycin calcium complex, and

thus preventing binding (28, 29). Based on the increased negative surface charge observed for

the T7SS mutants in the cytochrome C assay, we hypothesised that daptomycin could bind more

easily in the absence of a functional T7SS. To test this, daptomycin was tagged with BODIPY and

used to quantify binding via fluorometry (Fig. 7C) and to visualise bound daptomycin by fluorescence microscopy (Fig 7D). In the presence of daptomycin, a higher level of daptomycin was observed to bind around the septum for the WT as reported previously (30). *ΔesxC* showed significantly higher fluorescence in comparison to the WT. Thus, our data suggest that reduced electrostatic repulsion of daptomycin may contribute to the increased daptomycin binding in *ΔesxC*.

Calcium affects the growth of *ΔesxC* and sensitivity to daptomycin

Calcium is an important component in membranes, and affects both structure and charge of bacterial membranes (31). Calcium is also required for the activity of daptomycin and can enhance killing by gramicidin (32, 33). We hypothesied that EsxC may control membrane integrity through modulating the membrane binding of calcium. To test this, EsxC mutants were cultured in minimal medium with or without calcium chloride (50 mM). While both WT and mutant strains showed slow growth in minimal medium in the absence of calcium, in the presence of 50 mM calcium chloride the mutant grew much slower than the WT and the *esxC*-complemented strains (Fig 8 A,B,C).

We then examined if the daptomycin toxicity was dependent on the concentration of calcium available. WT, *ΔesxC*, and the complemented strains were cultured in the presence of 5 μg/ml daptomycin and increasing calcium chloride concentrations (0 to 20 mM). In the absence of calcium, the daptomycin did not have any effect as expected. A dose-dependent decrease in growth of the WT strain was observed with increasing concentrations of calcium chloride with the WT showing slower growth at 10 mM and 20 mM $CaCl_2$ compared to 1 mM (Figure 8 D, Fig S6). The mutant was more sensitive to daptomycin at all calcium concentrations compared to the WT and complemented strains, with increased growth inhibition at higher $CaCl_2$ concentrations .

Hence calcium binding to membranes and/or downstream signalling may be impacted in the *ΔesxC*, affecting its ability to grow in presence of calcium and its sensitivity to membrane targeting drugs like daptomycin.

Decreased survival of EsxC mutants during intracellular infection with daptomycin

S. aureus is a facultative intracellular pathogen, and daptomycin is considered to kill intracellular bacteria effectively. Hence, to determine if deletion of *esxC* from *S. aureus* would enhance intracellular killing activity, A549 human lung epithelial cells (15) were infected with *S. aureus* and treated with a high concentration of daptomycin (20 µg/ml). While there were no differences in the bacterial numbers between the WT and T7SS mutant in internalised bacteria (1 h after infection), a significant decrease was observed in the numbers of Δ*esxC* surviving within cells treated with daptomycin in comparison to WT after 4 h and 24 h after infection (Fig. 9 A). Notably, there was a significant number of WT *S. aureus* bacteria surviving intracellularly after treatment with daptomycin. Confocal microscopy imaging of infected cells also showed similar number of bacterial cells invading or internalized by mammalian cells but more *S. aureus* WT remaining 24 h after infection compared to Δ*esxC* in presence of daptomycin (Fig. 9 B, C). The differences seen in the infection assay were not due to the ability of the mutants to grow differently in DMEM-10, as strains had similar growth rates (Fig. S7A), with a growth defect in presence of daptomycin as expected (Fig. S7B). Therefore, our data suggest that daptomycin is more effective in killing intracellular *S. aureus* in the absence of T7SS.

***S. aureus* lacking EsxC is sensitive to daptomycin in a skin infection model**

As T7SS proteins have been reported to be important in various murine infection models (6, 8, 34), we next investigated effects of daptomycin on *S. aureus* infection and the impact of EsxC on daptomycin killing *in vivo* employing a localised murine skin abscess model. 7-8-week-old BALB/c mice were infected with *S. aureus* WT and Δ*esxC* intradermally, followed by treatment (intravenous) 2 h post infection (h.p.i) and every 24 h thereafter with 10 mg/kg daptomycin. Bacterial counts were obtained from skin tissue 24 or 72 h.p.i. Weight and skin lesion diameters were recorded every day. In the non-treated groups, bacterial counts from the site of infection

(lesions) showed no significant difference between WT and *ΔesxC* infected mice after 24 or 72 h

of infection. A significant decrease was seen in the numbers of *ΔesxC* present in the skin of

daptomycin-treated mice compared to the untreated group 24 and 72 h.p.i, while this was not the

case for the WT-infected group. (Fig. 8A and B). No significant differences were observed between

the areas of lesions measured for WT or *ΔesxC* after 24 h or 72 h infections and in daptomycin-

treated mice (Fig S8). Although the dose of daptomycin used in the time-frame of treatment was

not sufficient to treat WT *S. aureus* infection, this data suggests that the strain lacking EsxC was

more susceptible to daptomycin *in vivo* compared to the WT.

Discussion

The *S. aureus* T7SS has been strongly associated with staphylococcal virulence and the ability of

this pathogen to persist during infection (7). Its importance for bacterial competition between *S.*

aureus strains has been recently highlighted (13). However, the role of this system in

staphylococcal physiology has been poorly defined. Here we report that EsxC, a T7SS effector is

critical in maintaining cell membrane homeostasis through modulating membrane properties

including fluidity, accompanied by changes in membrane protein localisation, changes to the rate

of cell wall synthesis and surface charge. Our data indicate that EsxC may mediate these

membrane effects through modulating calcium binding to the membrane. Consequently, EsxC

impacts activity of membrane acting drugs including daptomycin, a key drug used to treat resilient

staphylococcal infections. Our data demonstrate that EsxC can modulate the activity of

daptomycin both intracellularly as well as during infection *in vivo*.

Several studies in mycobacterial species have linked the T7SS to cell surface integrity in different

ways (19, 35-38). The ESX-1 system has been previously reported to impact cell membrane

permeability in *M. tuberculosis*, although this effect appeared to be strain specific (35). However,

ESX-5 which is considered essential for *in vitro* growth, has been linked to mycobacterial capsular,

cell wall and outer membrane permeability (19, 36, 39). As ESX-5 was no longer essential in *M.*

marinum when the membrane was permeabilised, it was proposed that ESX-5 mediates transport

of cell envelope proteins to the outer membrane required for uptake of essential nutrients. While

these substrates have not been identified, a substrate PPE-10 was associated with capsular

integrity (19, 36). In *S. aureus,* we previously reported that the effectors, EsxC, and the ATPase,

EssC, were associated with maintenance of membrane integrity in response to antimicrobial host

fatty acids (20). T7SS proteins have been previously linked to membrane fluidity. Higher levels of

T7SS transcription were associated with a decrease in membrane fluidity, caused by the fatty acid

kinase complex, which incorporates host fatty acids into the membrane, inducing a change in

membrane dynamics (24).

To explain the membrane defects associated with EsxC we considered multiple possibilities; 1.

EsxC affects other membrane-associated proteins, impacting the lipid packing and fluidity of the

membrane; EsxC may be a chaperone for membrane proteins and play a role in targeting them

to the membrane. A recent bacterial two-hybrid library screening we performed was not successful

in identifying any direct bacterial interactors, however it is possible that such interactions are

transient. 2. EsxC may impact the state of the membrane by binding other molecules like metal

ions required for maintaining stability of the membrane (40). 3. T7SS substrates are secreted in a

co-dependent manner, so EsxC-mediated effects may be attributed to another T7SS substrate

like EsxA. Our data indicate that the T7SS substrates may be involved in modulating metal binding

to the membrane.

Calcium and other metal ions like magnesium are known to impact structure and function of

bacterial membranes (31, 41, 42). Calcium maintains bacterial outer membrane structure through

binding to bacterial lipopolysaccharide, and also influences bacterial membrane fluidity (43, 44).

Calcium can also have antibacterial effects; higher concentrations of calcium can disrupt

membrane integrity (42, 45, 46). Calcium affects the growth of *S. aureus* lacking EsxC indicating

that the protein may modulate calcium binding to the membrane. EsxC may directly or with other

T7SS effectors or membrane proteins, bind to calcium, and/ or be involved in transporting or

modulating calcium binding to the cell envelope. The molecular mechanisms underlying calcium

interactions of EsxC/ other T7SS substrates need further investigation.

It is also interesting that not all T7SS substrates have a cell surface associated defect which

indicates substrate specificity. EsxD and EsxB did not appear to be associated with surface

morphology defects by SEM as observed for EsxC and EsxA. Surface charges corelated well with

daptomycin sensitivity indicating that these substrates did not play a role in the membrane

homeostasis. Surface charge measurements of WT and mutants using a cytochrome C binding

assay revealed that ΔesxC and ΔesxA had an increased negative surface charge, whereas *ΔesxB*

did not display any differences in charge to the WT. The *esxC* and *essC* RN6390 strains however

did not show similar differences by SEM, which may be due to differences in global regulatory

circuits that may control cell wall synthesis (Agr, Sar) (47-49).

While *esxA* is conserved across all *S. aureus* strains, *esxC* is only found in the *essC1* locus (8, 9).

It is however likely that all *S. aureus* strains encode a protein orthologous to EsxC as proteins in

module 2, with variants essC2-4, have not yet been characterised and could function like EsxC.

Additionally, common phenotypes were seen for strains which lacked EsxC or EsxA. Previous

reports have suggested that EsxC and EsxA interact (11). Also, our previous study showed no

differences in esxA levels in ΔesxC total cell lysates, but we find decreased EsxA in the ΔesxC

secretome indicating co-dependent secretion of EsxA and EsxC (20). Therefore, interactions

between EsxC and EsxA may be important in modulating membrane integrity.

Daptomycin is thought to bind to the cell membrane before forming a complex with cell wall

biosynthesis machinery, preventing cell wall synthesis, before dispersing throughout the bacterial

membrane causing depolarisation (50). Daptomycin, although not a cationic antimicrobial peptide,

forms a highly positively charged complex with calcium to become active (50). Significantly more

daptomycin was bound to *ΔesxC* than the WT, which supports the hypothesis that the decrease

in surface charge measured in *ΔesxC* allows easier binding of daptomycin to the cell membrane.

In agreement with this, T7SS substrates may influence the free calcium available for activating

daptomycin. Daptomycin has also been reported to affect bacterial membrane fluidity through

perturbing membrane microdomains (51). Clinical daptomycin resistant isolates have generally

been associated with increased fluidity which is not associated with changes in fatty acid content

(52, 53). As we show that EsxC can impact membrane fluidity, it could further contribute to

daptomycin resistance by altering membrane fluidity and charge. Daptomycin resistance in *S.*

aureus has been mainly associated with mutations in genes encoding for multipeptide resistance

factor (MprF), a lipid biosynthetic enzyme, and the vancomycin-resistance associated

sensor/regulator (VraSR), two component systems (54-56). Genome sequencing did not reveal

any changes in the *ΔesxC* genome, and a proteomic analysis of mutant whole lysates did not

show any modulation of proteins implicated in daptomycin resistance (20).

In contrast to daptomycin, bithionol and gramicidin are uncharged (22, 57), therefore surface

charge and thus decreased electrostatic repulsion of the antibiotics is not the only factor involved

in increased sensitivity of the EsxC mutant, although calcium is known to increase gramicidin

activity (32). As all three antimicrobials have different mechanisms of action once bound to the

membrane, before causing membrane depolarization, the subsequent steps after membrane

binding appear to be important for efficacy. Therefore, daptomycin treatment which impacts both

the cell membrane, and the cell wall causes a larger effect in the the EsxC mutant.

While daptomycin kills the *esxC* mutant *in vivo*, at the dose and with the strains of *S. aureus* used,

there was no impact of the EsxC in bacterial survival during acute skin infection. T7SS proteins

have been previously reported to be important for bacterial virulence using invasive kidney

abscess, nasal colonisation and skin infection models, although studies have used different *S.*

aureus strains (6, 8, 34). The lack of decrease in the lesion size of the T7SS mutant-infected mice

treated with daptomycin, unlike the trend towards decreased lesion size observed for the WT (Fig

S7), could indicate that the increased killing of bacteria may cause an altered local immune

response. A recent study reported that in a *S. aureus* strain WU1, the T7SS was important only

for invasive infection and not localized nasal colonization (58).

Currently, levels of resistance to daptomycin are low, however, with increasing resistance in other

last line antibiotics, such as vancomycin, the use of daptomycin going forward will increase (59).

The increased use of daptomycin will likely lead to the occurrence of more resistant isolates,

therefore research into new antibiotics, and targeting new proteins that can enhance existing

treatments could be a powerful approach. Our data indicate that inhibitors of T7SS proteins or of

T7SS secretion could have excellent potential in combination drug therapies.

Materials and Methods

Bacterial strains: *S. aureus* strains used in this study are included in Table 2. Strains were

cultured aerobically in tryptic soy broth (TSB) (Sigma-Aldrich) at 37°C with aeration at 180 rpm

overnight. Overnight cultures were subcultured into fresh TSB and grown to a density of optical

density (OD_{600}) of 1, unless stated otherwise. For complemented strains TSB was supplemented

with 10 µg/ml chloramphenicol.

Growth curves: Overnight bacterial cultures were diluted to an OD_{600} of 0.1 in TSB or TSB

supplemented with antimicrobials where specified. Bacteria were grown in a 96-well plate with

shaking and the OD_{600} was measured every 15 min with a FLUOstar OMEGA plate reader (BMG

Labtech, UK). At appropriate time points 100 µl of the samples were taken for CFU determination.

For growth curves in defined medium, overnight cultures were subcultured into minimal defined

media, prepared as described by Machado and colleagues (60), at an initial OD_{600} of 0.15, and

supplemented with varying concentrations of $CaCl_2$ where stated. Bacterial growth was monitored

manually by measuring OD_{600} at different time points using a spectrophotometer (Biochrom), due

to issue with increased bacterial clumping in this medium on microtitre plates. 100 µl of the

samples were also taken for CFU determination at different time points.

Daptomycin killing assays: Bacterial cultures were grown overnight in TSB, diluted to an OD_{600}

of 0.15 and grown to log phase. Bacterial cultures were then treated with 10 µg/ml of daptomycin

(Acros Organics) in the presence of 1 mM $CaCl_2$ (Fisher Scientific) and incubated at 37ºC for a

further 2 h. Aliquots were removed at indicated timepoints, serially diluted in PBS and plated on

tryptic soy agar (TSA) for CFU determination.

Scanning Electron Microscopy: Overnight bacterial cultures were diluted to an OD_{600} of 0.15

and grown to an OD_{600} of 1 in TSB. The pellet was washed with PBS twice before dropping onto

poly-lysine coated 12 mm circular coverslips. Samples were fixed with 2.5% glutaraldehyde in

PBS for 1 h at 4°C. The bacterial cells were dehydrated by washing with a series of ethanol

solutions (20%, 50%, 70%, 90% and 100%) and dried with hexamethyldisilazane. The samples

were sputter coated with carbon and then observed using a Gemini SEM 500 (Zeiss).

Laurdan staining: Membrane fluidity was measured using the fluorescent dye Laurdan (Sigma)

and was based on protocols described by Wenzel and colleagues (61). Logarithmic phase *S.*

aureus was grown, and 1 ml aliquots were washed in PBS. The samples were incubated with a

final concentration of 100 µM Laurdan for 5 minutes at 37°C in the dark. Samples were washed

four times in pre-warmed PBS and 200 µl was transferred to a pre-warmed black walled 96 well

plate in triplicate. Membrane fluidity was determined by using an excitation wavelength of 350 nm

and measuring fluorescence at 460 nm and 500 nm emission wavelengths on a BioTek Cytation

Cell Imaging Multimode Reader (Agilent). The generalised polarisation (GP) value was calculated using the following calculation, GP = $(I_{460}-I_{500}) / (I_{460}+I_{500})$.

Pyrene decanoic acid staining: O/N bacterial cultures were diluted to an OD_{600} of 0.15 in TSB and were grown to an OD_{600} of 1.0. Bacteria were washed with PBS prior to treatment for 30 min at 37ºC with 37.5 µg/mL lysostaphin in PBS containing 20% sucrose. The spheroblasts were then centrifuged at 8000 × g for 10 min, and the pellet resuspended in the labelling solution (PBS, 20% sucrose, 0.01% F-127, 5 µM pyrene decanoic acid). The incubation in the dark was done for 1h at 25ºC under gentle rotation. PBS supplemented with 20% sucrose was used to wash the stained spheroblasts that were afterwards transferred to 96-well plates for fluorescence 450 nm measurements as previously described (24).

DiIC12 staining: O/N bacterial cultures were diluted to an OD_{600} of 0.15 in TSB with 1 µg/mL 1,1′-didodecyl-3,3,3′,3′-tetramethylindocarbocyanine perchlorate (DiIC12) (Invitrogen). Cultures were grown to an OD_{600} of 1.0, centrifuged and washed twice with fresh TSB. Samples were spotted on to agarose pads and imaged using a Leica DMi8 widefield microscope (Leica Microsystems, UK). Acquired images were analysed with the ImageJ processing package, Fiji.

Cytochrome C binding assay: Overnight bacterial cultures were pelleted and washed with 20 mM MOPS buffer, pH 7. The cells were resuspended in MOPS buffer at an OD_{600} of 1 and incubated with 50 µg/ml cytochrome C at room temperature for 15 min. The cells were pelleted and the remaining cytochrome C in the supernatant was measured spectrophotometrically at OD_{410} using a FLUOstar OMEGA plate reader (BMG Labtech, UK).

HADA incorporation assay

Peptidoglycan synthesis was investigated by measuring HCC-amino-D-alanine (HADA) incorporation. Overnight *S. aureus* cultures were diluted into TSB supplemented with 25 µM HADA and incubated at 37ºC in the dark. When OD_{600} of 1 was reached cultures were washed three times and fixed with 4% PFA. Alternatively, cultures were treated with daptomycin for 30 min.

before washing and fixing. Microscopy was undertaken on these samples using the DAPI filter set

on a Leica DMi8 widefield microscope.

Daptomycin BODIPY binding assay: *S. aureus* was grown as previously described and treated

with 1μg/ml daptomycin-BODIPY once logarithmic phase had been reached. The samples were

incubated at 37ºC for 30 min in the dark, then washed with PBS three times. 200 μl were aliquoted

into black 96 well plates in triplicate and fluorescence was measured using a BioTek Cytation 5

Cell Imaging Multimode Reader (Agilent). The excitation wavelength used was 488 nm and

emission wavelength used was 530 nm.

Invasion and survival assays: Infection assays were carried out in 24 well tissue culture treated

plates (Falcon), or RS treated glass chamber slides (154526, Nunc, Lab Tek II). Overnight *S.*

aureus cultures were diluted to an OD_{600} of 0.15 and grown to an OD_{600} of 1. Cultures were

resuspended in DMEM media to a density of 2.5 x 10^6 CFU/ml. A549 epithelial cells were seeded

in 24 well plates at a density of 2.5 x 10^5 one day before being infected with *S. aureus* at a

multiplicity of infection (MOI) of 10. Cells were infected for 1 h at 37°C with 5 % CO_2. To kill any

extracellular bacteria, cells were washed with DMEM media containing 20 μg/ml lysostaphin

(Sigma-Aldrich) and 50 μg/ml gentamycin (Melford) and incubated for a further 30 minutes. Cells

were washed twice with PBS, lysed with 1 ml cold sterile water and vigorous pipetting before

plating for CFU determination. To investigate the course of infection after extracellular bacterial

killing, the cells are washed twice with PBS and refreshed with DMEM with 1 mM $CaCl_2$ only, or

DMEM supplemented with 1 mM $CaCl_2$ and 10 μg/ml daptomycin and incubated until the desired

timepoint. The above protocol to lyse the cells is then followed.

Murine skin abscess model: All animal procedures were carried out as per protocols in the

Project Licence PCEC27E7D. All local ethical and home office approvals were obtained for animal

experimentation. For the murine skin abscess model, female 7/8-week-old BALB/c mice (Charles

River Laboratories, UK) flanks were shaved before injection, and inoculated through an

intradermal route with 50 μl of bacterial suspension per flank (one injection on each flank),

resulting in a final bacterial count of 1-2 x 10^6 per mouse. Overnight *S. aureus* cultures were

subcultured, grown to OD_{600} of 1, centrifuged and washed twice with PBS, and diluted in PBS to

achieve a final CFU/ml of 1-2 x 10^7 for infection. 2 h after infection mice were given an intravenous

dose of daptomycin or PBS via tail vein injection. This was repeated once every 24 h. During the

experiment, mice were weighed daily, and any abscess formation was measured and recorded.

After 24 h or 72 h mice were euthanised by CO_2 inhalation and skin at the infection site was

incised. The skin samples were weighed and homogenised in PBS using a FastPrep (MP

Biomedicals), at 4 m/s for 60 seconds for a total of 6 cycles. Homogenates were diluted in PBS

and CFU/ml was determined. CFU counts were normalized to lesion weight.

Statistical Analysis: All experiments were performed at least three independent times. Statistical

analysis was performed using GraphPad Prism 9.0 (MacOS version 9.5.1). Pairwise statistical

analysis was carried out by the Mann-Whitney U test or t test? for two groups, or One way

ANOVAs for multiple groups, followed by post-hoc Tukey's HSD tests where necessary.

Significance was denoted by asterisks; * = $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$, **** = $P <$

0.0001 and ns = not significant.

Acknowledgements

We would like to acknowledge funding from MIBTP (grant 2097375) to Victoria Smith and the

Medical Research Council grants (MR/N010140/1 and MR/X00161X/1) for funding.

We would like to thank Novartis Vaccines, Siena (GSK biologicals) Italy for providing the *ΔesxA*

and *ΔesxB* strains, and Prof Tracy Palmer for providing the RN6390 *ΔesxC* and *ΔessC* strains,

used in this study. We acknowledge the Midlands Regional Cryo-EM Facility, hosted at the

Warwick Advanced Bioimaging Research Technology Platform, for the use of the JEOL 2100Plus.

Author contributions

VS, AKT and MU conceptualised ideas, designed experiments, VS, GC, RF, KEW, RGM, JY, AKT

conducted experiments, MU and SP provided resources for the study, VS, AKT, GC, KEW, RGM,

RF and MU analysed data, AKT and MU supervised study, MU, VS, RF and AKT drafted

manuscript and all authors were involved in reviewing manuscript draft.

References

522   1.   Tong SY, Davis JS, Eichenberger E, Holland TL, Fowler VG, Jr. 2015. Staphylococcus aureus
523   infections: epidemiology, pathophysiology, clinical manifestations, and management. Clin
524   Microbiol Rev 28:603-61.

525   2.   Chambers HF, Deleo FR. 2009. Waves of resistance: Staphylococcus aureus in the antibiotic
526   era. Nat Rev Microbiol 7:629-41.

527   3.   Lee BY, Singh A, David MZ, Bartsch SM, Slayton RB, Huang SS, Zimmer SM, Potter MA,
528   Macal CM, Lauderdale DS, Miller LG, Daum RS. 2013. The economic burden of community-
529   associated methicillin-resistant Staphylococcus aureus (CA-MRSA). Clin Microbiol Infect
530   19:528-36.

531   4.   Cheung GYC, Bae JS, Otto M. 2021. Pathogenicity and virulence of Staphylococcus aureus.
532   Virulence 12:547-569.

533   5.   Bottai D, Groschel MI, Brosch R. 2016. Type VII Secretion Systems in Gram-Positive
534   Bacteria. Curr Top Microbiol Immunol doi:10.1007/82_2015_5015.

535   6.   Burts ML, Williams WA, DeBord K, Missiakas DM. 2005. EsxA and EsxB are secreted by an
536   ESAT-6-like system that is required for the pathogenesis of Staphylococcus aureus
537   infections. Proc Natl Acad Sci U S A 102:1169-74.

538   7.   Unnikrishnan M, Constantinidou C, Palmer T, Pallen MJ. 2017. The Enigmatic Esx Proteins:
539   Looking Beyond Mycobacteria. Trends Microbiol 25:192-204.

540   8.   Kneuper H, Cao ZP, Twomey KB, Zoltner M, Jager F, Cargill JS, Chalmers J, van der Kooi-Pol
541   MM, van Dijl JM, Ryan RP, Hunter WN, Palmer T. 2014. Heterogeneity in ess transcriptional
542   organization and variable contribution of the Ess/Type VII protein secretion system to
543   virulence across closely related Staphylocccus aureus strains. Mol Microbiol 93:928-43.

544   9.   Warne B, Harkins CP, Harris SR, Vatsiou A, Stanley-Wall N, Parkhill J, Peacock SJ, Palmer T,
545   Holden MT. 2016. The Ess/Type VII secretion system of Staphylococcus aureus shows
546   unexpected genetic diversity. BMC Genomics 17:222.

547   10.   Bowman L, Palmer T. 2021. The Type VII Secretion System of Staphylococcus. Annu Rev
548   Microbiol 75:471-494.

549   11.   Anderson M, Aly KA, Chen YH, Missiakas D. 2013. Secretion of atypical protein substrates
550   by the ESAT-6 secretion system of Staphylococcus aureus. Mol Microbiol 90:734-43.

551   12.   Mielich-Suss B, Wagner RM, Mietrach N, Hertlein T, Marincola G, Ohlsen K, Geibel S, Lopez
552   D. 2017. Flotillin scaffold activity contributes to type VII secretion system assembly in
553   Staphylococcus aureus. PLoS Pathog 13:e1006728.

554   13.   Cao Z, Casabona MG, Kneuper H, Chalmers JD, Palmer T. 2016. The type VII secretion
555   system of Staphylococcus aureus secretes a nuclease toxin that targets competitor
556   bacteria. Nat Microbiol 2:16183.

557   14.   Ulhuq FR, Gomes MC, Duggan GM, Guo M, Mendonca C, Buchanan G, Chalmers JD, Cao Z,
558   Kneuper H, Murdoch S, Thomson S, Strahl H, Trost M, Mostowy S, Palmer T. 2020. A
559   membrane-depolarizing toxin substrate of the Staphylococcus aureus type VII secretion
560   system mediates intraspecies competition. Proc Natl Acad Sci U S A 117:20836-20847.

561   15.   Korea CG, Balsamo G, Pezzicoli A, Merakou C, Tavarini S, Bagnoli F, Serruto D, Unnikrishnan
562   M. 2014. Staphylococcal Esx proteins modulate apoptosis and release of intracellular
563   Staphylococcus aureus during infection in epithelial cells. Infect Immun 82:4144-53.

564   16.   Anderson M, Ohr RJ, Aly KA, Nocadello S, Kim HK, Schneewind CE, Schneewind O,
565   Missiakas D. 2017. EssE Promotes Staphylococcus aureus ESS-Dependent Protein Secretion
566   To Modify Host Immune Responses during Infection. J Bacteriol 199.

567   17.   Fyans JK, Bignell D, Loria R, Toth I, Palmer T. 2013. The ESX/type VII secretion system
568   modulates development, but not virulence, of the plant pathogen Streptomyces scabies.
569   Mol Plant Pathol 14:119-30.

570   18.   Flint JL, Kowalski JC, Karnati PK, Derbyshire KM. 2004. The RD1 virulence locus of
571   Mycobacterium tuberculosis regulates DNA transfer in Mycobacterium smegmatis. Proc
572   Natl Acad Sci U S A 101:12598-603.

573   19.   Ates LS, Ummels R, Commandeur S, van de Weerd R, Sparrius M, Weerdenburg E, Alber
574   M, Kalscheuer R, Piersma SR, Abdallah AM, Abd El Ghany M, Abdel-Haleem AM, Pain A,
575   Jimenez CR, Bitter W, Houben EN. 2015. Essential Role of the ESX-5 Secretion System in
576   Outer Membrane Permeability of Pathogenic Mycobacteria. PLoS Genet 11:e1005190.

577   20.   Kengmo Tchoupa A, Watkins KE, Jones RA, Kuroki A, Alam MT, Perrier S, Chen Y,
578   Unnikrishnan M. 2020. The type VII secretion system protects Staphylococcus aureus
579   against antimicrobial host fatty acids. Sci Rep 10:14838.

580   21.   Huang HW. 2020. DAPTOMYCIN, its membrane-active mechanism vs. that of other
581   antimicrobial peptides. Biochim Biophys Acta Biomembr 1862:183395.

582   22.   Kim W, Zou G, Hari TPA, Wilt IK, Zhu W, Galle N, Faizi HA, Hendricks GL, Tori K, Pan W, Huang
583   X, Steele AD, Csatary EE, Dekarske MM, Rosen JL, Ribeiro NQ, Lee K, Port J, Fuchs BB,
584   Vlahovska PM, Wuest WM, Gao H, Ausubel FM, Mylonakis E. 2019. A selective membrane-
585   targeting repurposed antibiotic with activity against persistent methicillin-resistant
586   Staphylococcus aureus. Proc Natl Acad Sci U S A 116:16529-16534.

587   23.   Strahl H, Burmann F, Hamoen LW. 2014. The actin homologue MreB organizes the bacterial
588   cell membrane. Nat Commun 5:3442.

589   24.   Lopez MS, Tan IS, Yan D, Kang J, McCreary M, Modrusan Z, Austin CD, Xu M, Brown EJ.
590   2017. Host-derived fatty acids activate type VII secretion in Staphylococcus aureus. Proc
591   Natl Acad Sci U S A 114:11223-11228.

592   25.   Scheinpflug K, Krylova O, Strahl H. 2017. Measurement of Cell Membrane Fluidity by
593   Laurdan GP: Fluorescence Spectroscopy and Microscopy. Methods Mol Biol 1520:159-174.

594   26.   Kuru E, Hughes HV, Brown PJ, Hall E, Tekkam S, Cava F, de Pedro MA, Brun YV,
595   VanNieuwenhze MS. 2012. In Situ probing of newly synthesized peptidoglycan in live
596   bacteria with fluorescent D-amino acids. Angew Chem Int Ed Engl 51:12519-23.

27.   Peschel A, Otto M, Jack RW, Kalbacher H, Jung G, Gotz F. 1999. Inactivation of the dlt
operon in Staphylococcus aureus confers sensitivity to defensins, protegrins, and other
antimicrobial peptides. J Biol Chem 274:8405-10.

28.   Bertsche U, Yang SJ, Kuehner D, Wanner S, Mishra NN, Roth T, Nega M, Schneider A, Mayer
C, Grau T, Bayer AS, Weidenmaier C. 2013. Increased cell wall teichoic acid production and
D-alanylation are common phenotypes among daptomycin-resistant methicillin-resistant
Staphylococcus aureus (MRSA) clinical isolates. PLoS One 8:e67398.

29.   Mishra NN, Bayer AS, Weidenmaier C, Grau T, Wanner S, Stefani S, Cafiso V, Bertuccio T,
Yeaman MR, Nast CC, Yang SJ. 2014. Phenotypic and genotypic characterization of
daptomycin-resistant methicillin-resistant Staphylococcus aureus strains: relative roles of
mprF and dlt operons. PLoS One 9:e107426.

30.   Monteiro JM, Fernandes PB, Vaz F, Pereira AR, Tavares AC, Ferreira MT, Pereira PM, Veiga
H, Kuru E, VanNieuwenhze MS, Brun YV, Filipe SR, Pinho MG. 2015. Cell shape dynamics
during the staphylococcal cell cycle. Nat Commun 6:8055.

31.   Smith RJ. 1995. Calcium and bacteria. Adv Microb Physiol 37:83-133.

32.   Fang ST, Huang SH, Yang CH, Liou JW, Mani H, Chen YC. 2022. Effects of Calcium Ions on
the Antimicrobial Activity of Gramicidin A. Biomolecules 12.

33.   Jung D, Rozek A, Okon M, Hancock RE. 2004. Structural transitions as determinants of the
action of the calcium-dependent antibiotic daptomycin. Chem Biol 11:949-57.

34.   Wang Y, Hu M, Liu Q, Qin J, Dai Y, He L, Li T, Zheng B, Zhou F, Yu K, Fang J, Liu X, Otto M, Li
M. 2016. Role of the ESAT-6 secretion system in virulence of the emerging community-
associated Staphylococcus aureus lineage ST398. Sci Rep 6:25163.

35.   Garces A, Atmakuri K, Chase MR, Woodworth JS, Krastins B, Rothchild AC, Ramsdell TL,
Lopez MF, Behar SM, Sarracino DA, Fortune SM. 2010. EspA acts as a critical mediator of
ESX1-dependent virulence in Mycobacterium tuberculosis by affecting bacterial cell wall
integrity. PLoS Pathog 6:e1000957.

36.   Ates LS, van der Woude AD, Bestebroer J, van Stempvoort G, Musters RJ, Garcia-Vallejo JJ,
Picavet DI, Weerd R, Maletta M, Kuijl CP, van der Wel NN, Bitter W. 2016. The ESX-5 System
of Pathogenic Mycobacteria Is Involved In Capsule Integrity and Virulence through Its
Substrate PPE10. PLoS Pathog 12:e1005696.

37.   Nath Y, Ray SK, Buragohain AK. 2018. Essential role of the ESX-3 associated eccD3 locus in
maintaining the cell wall integrity of Mycobacterium smegmatis. Int J Med Microbiol
308:784-795.

38.   Bosserman RE, Champion PA. 2017. Esx Systems and the Mycobacterial Cell Envelope:
What's the Connection? J Bacteriol 199.

39.   Bottai D, Di Luca M, Majlessi L, Frigui W, Simeone R, Sayes F, Bitter W, Brennan MJ, Leclerc
C, Batoni G, Campa M, Brosch R, Esin S. 2012. Disruption of the ESX-5 system of
Mycobacterium tuberculosis causes loss of PPE protein secretion, reduction of cell wall
integrity and strong attenuation. Mol Microbiol 83:1195-209.

40.   Thomas KJ, 3rd, Rice CV. 2014. Revised model of calcium and magnesium binding to the
bacterial cell wall. Biometals 27:1361-70.

41.   Pedersen UR, Leidy C, Westh P, Peters GH. 2006. The effect of calcium on the properties of
charged phospholipid bilayers. Biochim Biophys Acta 1758:573-82.

42. Melcrova A, Pokorna S, Pullanchery S, Kohagen M, Jurkiewicz P, Hof M, Jungwirth P, Cremer
PS, Cwiklik L. 2016. The complex nature of calcium cation interactions with phospholipid
bilayers. Sci Rep 6:38035.

43. Ginez LD, Osorio A, Vazquez-Ramirez R, Arenas T, Mendoza L, Camarena L, Poggio S. 2022.
Changes in fluidity of the E. coli outer membrane in response to temperature, divalent
cations and polymyxin-B show two different mechanisms of membrane fluidity adaptation.
FEBS J 289:3550-3567.

44. Schindler M, Osborn MJ. 1979. Interaction of divalent cations and polymyxin B with
lipopolysaccharide. Biochemistry 18:4425-30.

45. Xie Y, Yang L. 2016. Calcium and Magnesium Ions Are Membrane-Active against Stationary-
Phase Staphylococcus aureus with High Specificity. Sci Rep 6:20628.

46. Asif A, Mohsin H, Tanvir R, Rehman Y. 2017. Revisiting the Mechanisms Involved in Calcium
Chloride Induced Bacterial Transformation. Front Microbiol 8:2169.

47. Fujimoto DF, Bayles KW. 1998. Opposing roles of the Staphylococcus aureus virulence
regulators, Agr and Sar, in Triton X-100- and penicillin-induced autolysis. J Bacteriol
180:3724-6.

48. Cheung GY, Wang R, Khan BA, Sturdevant DE, Otto M. 2011. Role of the accessory gene
regulator agr in community-associated methicillin-resistant Staphylococcus aureus
pathogenesis. Infect Immun 79:1927-35.

49. Zielinska AK, Beenken KE, Joo HS, Mrak LN, Griffin LM, Luong TT, Lee CY, Otto M, Shaw LN,
Smeltzer MS. 2011. Defining the strain-dependent impact of the Staphylococcal accessory
regulator (sarA) on the alpha-toxin phenotype of Staphylococcus aureus. J Bacteriol
193:2948-58.

50. Grein F, Muller A, Scherer KM, Liu X, Ludwig KC, Klockner A, Strach M, Sahl HG, Kubitscheck
U, Schneider T. 2020. Ca(2+)-Daptomycin targets cell wall biosynthesis by forming a
tripartite complex with undecaprenyl-coupled intermediates and membrane lipids. Nat
Commun 11:1455.

51. Muller A, Wenzel M, Strahl H, Grein F, Saaki TNV, Kohl B, Siersma T, Bandow JE, Sahl HG,
Schneider T, Hamoen LW. 2016. Daptomycin inhibits cell envelope synthesis by interfering
with fluid membrane microdomains. Proc Natl Acad Sci U S A 113:E7077-E7086.

52. Jones T, Yeaman MR, Sakoulas G, Yang SJ, Proctor RA, Sahl HG, Schrenzel J, Xiong YQ, Bayer
AS. 2008. Failures in clinical treatment of Staphylococcus aureus Infection with daptomycin
are associated with alterations in surface charge, membrane phospholipid asymmetry, and
drug binding. Antimicrob Agents Chemother 52:269-78.

53. Mishra NN, Bayer AS. 2013. Correlation of cell membrane lipid profiles with daptomycin
resistance in methicillin-resistant Staphylococcus aureus. Antimicrob Agents Chemother
57:1082-5.

54. Ernst CM, Peschel A. 2019. MprF-mediated daptomycin resistance. Int J Med Microbiol
309:359-363.

55. Mehta S, Cuirolo AX, Plata KB, Riosa S, Silverman JA, Rubio A, Rosato RR, Rosato AE. 2012.
VraSR two-component regulatory system contributes to mprF-mediated decreased
susceptibility to daptomycin in in vivo-selected clinical strains of methicillin-resistant
Staphylococcus aureus. Antimicrob Agents Chemother 56:92-102.

56.   Nguyen AH, Hood KS, Mileykovskaya E, Miller WR, Tran TT. 2022. Bacterial cell membranes
684   and their role in daptomycin resistance: A review. Front Mol Biosci 9:1035574.
685   57.   Rostovtseva TK, Aguilella VM, Vodyanoy I, Bezrukov SM, Parsegian VA. 1998. Membrane
686   surface-charge titration probed by gramicidin A channel conductance. Biophys J 75:1783-
687   92.
688   58.   Bobrovskyy M, Chen X, Missiakas D. 2023. The Type 7b Secretion System of S. aureus and
689   Its Role in Colonization and Systemic Infection. Infect Immun 91:e0001523.
690   59.   Shariati A, Dadashi M, Moghadam MT, van Belkum A, Yaslianifard S, Darban-Sarokhalil D.
691   2020. Global prevalence and distribution of vancomycin resistant, vancomycin
692   intermediate and heterogeneously vancomycin intermediate Staphylococcus aureus
693   clinical isolates: a systematic review and meta-analysis. Sci Rep 10:12689.
694   60.   Machado H, Weng LL, Dillon N, Seif Y, Holland M, Pekar JE, Monk JM, Nizet V, Palsson BO,
695   Feist AM. 2019. Strain-Specific Metabolic Requirements Revealed by a Defined Minimal
696   Medium for Systems Analyses of Staphylococcus aureus. Appl Environ Microbiol 85.
697   61.   Wenzel M, Vischer NOE, Strahl H, Hamoen LW. 2018. Assessing Membrane Fluidity and
698   Visualizing Fluid Membrane Domains in Bacteria Using Fluorescent Membrane Dyes. Bio
699   Protoc 8:e3063.

Figure Legends

Figure 1. An *esxC* mutant is highly sensitive to daptomycin.

A) Growth curves of *S. aureus* WT USA300 JE2 and Δ*esxC* in TSB in the absence or presence

of 5 µg/ml daptomycin and 1 mM $CaCl_2$. Mean ± standard error of the mean (SEM) is shown, N =

3 (biological replicates). **B)** CFU/ml was enumerated after 8 h daptomycin treatment. N = 3

(biological replicates), mean + standard deviation (SD), *** P ≤ 0.001 using an unpaired t-test **C)**

Growth curves of USA300 JE2 WTpOS1, Δ*esxC*pOS1 and complemented strain, Δ*esxC*pOS1-

esxC in TSB supplemented with 5 µg/ml daptomycin and 1 mM $CaCl_2$. Mean ± SEM shown, N =

3 (biological replicates); AUC was calculated for the strains grown in presence of daptomycin **(D)**

Mean + SD ** P ≤ 0.05 using a one-way ANOVA with Tukey's multiple comparison test. **E)** CFU/ml

of WT USA300 JE2 and Δ*esxC* were determined from a killing assay in the presence of 10 µg/ml

daptomycin and 1 mM $CaCl_2$ at different timepoints. **F)** CFU/ml from killing assay using WT

USA300 JE2, Δ*esxC* and Δ*esxC* complemented strains in the absence or presence of 10 µg/ml

daptomycin and 1 mM CaCl$_2$ at different timepoints, mean ± SD, N = 3 (biological replicates), * P

≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001 using a two-way ANOVA with Tukey multiple comparison test.

Figure 2. Membrane permeability of the *esxC* mutant in the presence of daptomycin

A) Representative fluorescent micrographs of *S. aureus* USA300 JE2 WT and Δ*esxC* treated with

daptomycin. Cell membranes were stained by FM1-43 and imaged using FITC filter channel, PI

was imaged using Texas red filter channel. **B)** Images of daptomycin-treated samples (A) were

quantified using FIJI. Mean fluorescence intensity was quantified from 10 images from each

experiment and normalised to the number of cells in the field of view. Data presented is from 3

independent experiments. ****P < 0.0001 using an unpaired t test.

Figure 3. The *esxC* mutant show higher sensitivity to other membrane acting drugs

A) Growth curves of USA300 JE2 WT and Δ*esxC* in the presence of 2 µg/ml gramicidin, Mean ±

SEM, N = 3 (biological replicates). **B)** Area under curves (AUC) were calculated and the mean

AUC + SD was plotted for growth curves in (A). **C)** Growth curves of USA300 JE2 WT and Δ*esxC*

in the presence of 1 µg/ml bithionol. Mean ± SEM, N = 3 (biological replicates). **D)** AUC were

calculated and the mean AUC + SD was plotted for growth curves in (C). *P ≤ 0.05 using an

unpaired t-test.

Figure 4. EsxC affects membrane fluidity.

A) Widefield micrographs of *S. aureus* WT USA300 JE2 and Δ*esxC* after growth in TSB to OD$_{600}$

of 1.0 in the presence of the lipophilic dye DiIC12. Images are representative of 4 independent

experiments. The DiIC12 fluorescence of 80 bacterial clusters from different fields per strain was

quantitated with ImageJ. Means + SD are shown, N = 4 (biological replicates); **** P < 0.0001,

using an unpaired t-test. **B)** The membrane fluidity of *S. aureus* WT USA300 JE2, Δ*esxC* and

Δ*esxC* pOS1-*esxC* as measured with a pyrene decanoic acid staining-based assay. Mean ± SEM

are shown, N = 6 (biological replicates), * indicates P < 0.05 using a one-way ANOVA. **C)** USA300

JE2 WT, ΔesxC and ΔesxC pOS1-esxC were grown to logarithmic phase in TSB at 37°C prior to

staining with Laurdan. To assess membrane fluidity, Laurdan generalised polarisation (GP) was

calculated using the equation $(I_{460} - I_{500})/(I_{460} + I_{500})$ where I refers to the fluorescence intensity at

the indicated emission wavelength. *** P <0.0001 using a one-way ANOVA, N = 3 (biological

replicates), mean + SD.

Figure 5. The *esxC* mutant displays distinct surface morphologies.

A) Representative SEM images of *S. aureus* WT USA300 JE2 and *ΔesxC* grown to early

logarithmic phase. **B)** Representative SEM images of *S. aureus* USA300 JE2 WT pOS1, *ΔesxC*

pOS1, *ΔesxC* pOS1-*esxC* grown to early logarithmic phase.

Figure 6. EsxC impacts cell wall synthesis.

A) Representative widefield micrographs of *S. aureus* WT USA300 JE2 and *ΔesxC* either grown

to logarithmic phase in the presence of 25 μM HADA or after 10-minute treatment with HADA once

OD_{600} of 1 was reached. **B)** Quantification of fluorescence in cells grown to logarithmic phase with

HADA. **C)** Quantification of fluorescence in cells treated with HADA for 10 minutes. Quantification

performed on five images each from 3 independent experiments. * P< 0.01, and using an unpaired

t test.

Figure 7. EsxC mutant binds more to daptomycin.

A-B) Quantitative binding assay of cytochrome C to USA300 JE2 WT, ΔesxC and ΔesxC pOS1-

esxC (A). The CFU of the starting inoculum was enumerated for each mutant (B). Mean ± SD

shown, N = 3 (biological replicates), ***P < 0.001 using a one-way ANOVA. **C)** Fluorescence

measurements of daptomycin-BODIPY upon binding to *S. aureus* WT USA300 JE2 and *ΔesxC.*

The excitation wavelength was 488 nm and emission wavelength was 530 nm. N = 3 (biological

replicates), mean + SD. P = 0.0067, using an unpaired t-test. **D)** Representative micrographs of

daptomycin-BODIPY binding to *S. aureus* WT and Δ*esxC* are shown.

Figure 8. Calcium affects growth of *esxC* mutant and daptomycin sensitivity

Growth curves of *S. aureus* WT USA300 JE2, Δ*esxC*, and Δ*esxC* pOS1-*esxC* in synthetic minimal

medium in the absence (A) or presence (B) of 50 mM $CaCl_2$. Mean ± standard error of the mean

(SEM) is shown **C)** Area under curves (AUC) were calculated from (A) and (B) and the mean AUC

± SD were plotted. **D)** *S. aureus* WT USA300 JE2, Δ*esxC*, and Δ*esxC* pOS1-*esxC* were grown in

TSB supplemented with increasing calcium chloride concentrations (0 to 20 mM) in the presence

of daptomycin (5 µg/ml). AUCs calculated from growth curves shown in Figure S6 are shown.

Mean + SD, ***P= 0.0001, ****P <0.0001, using two-way ANOVA.

Figure 9. Presence of EsxC affects intracellular bacterial killing by daptomycin.

A) CFU counts at various timepoints during epithelial cell infection assay with WT USA300 JE2,

and Δ*esxC*. 0 h represents the initial inoculum, 1 h represents the CFU counts that have infected

the A549 cells and 4 h and 24 h represent the survival of *S. aureus* in the presence of 20 µg/ml

daptomycin, N = 3 (biological replicates), **P<0.05, two-way ANOVA. **B)** Representative confocal

micrographs of cells after 1 h infection or 24 h post-infection. DAPI (blue), F-actin (red), and *S.*

aureus (green, white arrows). **C)** Confocal image quantification from five images each from 3

independent experiments after 24 h. Mean ± SD of intensity, is shown, ***P<0.001 using an

unpaired t-test.

Figure 10. Effect of daptomycin on *esxC* mutants *in vivo*.

Mice were infected in the skin with WT USA300 and Δ*esxC* mutant followed by treatment with 10

mg/kg daptomycin. **A)** CFU/g skin tissue was counted 24 h post infection (n = 8, 2 lesions/animal).

Graphs show mean ± SD, P = 0.0007 using one-way ANOVA. **B)** CFU/g was calculated 72 h post

infection (n = 5, 2 lesions/animal), mean ± SD shown, P = 0.0020 using a one-way ANOVA.

Table 1. Proteins significantly changed in Δ*esxC* mutant membrane preparations relative to WT

USA300 JE2.

Uniprot ID	log$_2$ fold change	Adjusted *P* value	Description
A0A0H2XJR2	-5.46	6.21E-08	Pyridine nucleotide-disulfide oxidoreductase
A0A0H2XJ15	-5.34	3.80E-09	TPR domain protein
A0A0H2XIV9	-5.33	1.49E-07	T7SS protein, EssD
A0A0H2XJ90	-4.82	6.92E-09	D-isomer specific 2-hydroxyacid dehydrogenase family protein
A0A0H2XF36	3.62	2.14E-08	Uncharacterised protein
A0A0H2XJV3	3.67	2.26E-08	ABC transporter, ATP-binding protein
A0A0H2XGS2	3.70	1.71E-08	Putative sensor histidine kinase
A0A0H2XG87	4.01	1.59E-08	Bifunctional ligase/repressor BirA
A0A0H2XH85	4.14	1.04E-08	Cobalamin synthesis protein/P47K family protein
A0A0H2XER9	4.38	9.36E-09	LysR family regulatory protein
A0A0H2XEB7	5.25	2.39E-07	Integral membrane protein
A0A0H2XJ96	5.27	1.04E-08	NADH-dependent flavin oxidoreductase
A0A0H2XHU5	5.97	2.55E-09	Putative membrane protein

800　**Table 2. Strains and Plasmids used in this study**

Strain or Plasmid	Description	Source or Reference
Staphylococcus aureus		
USA300 LAC JE2	Plasmid-cured USA300 LAC	BEI Resources (NARSA)
USA300 JE2 *ΔessC*	USA300 LAC JE2 defective for EssC	(Kengmo Tchoupa *et al.,* 2020)
USA300 JE2 *ΔesxC*	USA300 LAC JE2 defective for EsxC	(Kengmo Tchoupa *et al.,* 2020)
USA300 LAC	Community acquired MRSA	(Bae and Schneewind, 2006)
USA300 LAC *ΔesxA*	USA300 LAC defective for EsxA	(Korea *et al.,* 2014)
USA300 LAC *ΔesxB*	USA300 LAC defective for EsxB	(Korea *et al.,* 2014)
Newman	Methicillin-sensitive *S. aureus*	(Bae and Schneewind, 2006)
Newman *ΔesxA*	Newman defective for EsxA	(Korea *et al.,* 2014)
Newman *ΔesxB*	Newman defective for EsxB	(Korea *et al.,* 2014)
RN6390	NCTC8325 derivative, *ΔrbsU, ΔtcaR,* cured of φ11, φ12, and φ13	Tracy Palmer (Kneuper *et al.,* 2014)
RN6390 *ΔessC*	RN6390 defective for EssC	Tracy Palmer (Kneuper *et al.,* 2014)
RN6390 *ΔesxC*	RN6390 defective for EsxC	Tracy Palmer (Kneuper *et al.,* 2014)
Plasmids		
pOS1	Empty vector for complementation	Olaf Schneewind
pOS1-*esxC*	*esxC* complementation vector	(Kengmo Tchoupa *et al.,* 2020)

Supplementary Figure legends and tables

Figure S1. Growth curves in TSB for A) *S. aureus* USA300 LAC Δ*esxA* and Δ*esxB*, C) *S. aureus*

Newman Δ*esxA* and Δ*esxB*, E) *S. aureus* USA300 JE2 WT and Δ*esxD,* G) *S. aureus* USA300

JE2 WT and Δ*essC* I) *S. aureus* RN6390 WT, Δ*essC* and Δ*esxC*, in the absence or presence of

5 µg/ml daptomycin and 1 mM $CaCl_2$. Mean ± SEM are shown, N = 3. The AUC was calculated

for the growth curves of all strains cultured in presence of daptomycin in A, C, E and G and I are

presented as % relative to WT, mean + SD is shown (B, D, F, H, J). * $P \leq 0.05$, ** $P \leq 0.01$ using

a one-sample t test. J) shows AUC of I) ****$P \leq 0.0001$, unpaired t test

Figure S2. Growth curves of USA300 JE2 WT, Δ*essC* and Δ*esxC* in the presence of (A) 4 µg/ml

ciprofloxacin (B) 0.5 µg/ml mitomycin C. (C) 1 µg/ml oxacillin. (D) 8 µg/ml vancomycin. For all

growth curves mean and SEM are shown, N = 3 (biological replicates).

Figure S3. Scanning electron micrographs of *S. aureus* RN6390, USA300, Newman and their

T7SS mutants.

Figure S4. Transmission electron micrographs of *S. aureus* USA300 JE2 WT and Δ*esxC* grown

to early logarithmic phase.

Figure S5. Quantitative binding assay of cytochrome C to USA300 LAC WT, Δ*esxA* and Δ*esxB*

(A). The CFU of the starting inoculum was calculated for each mutant **(B)**. Mean ± SD shown, N

= 3 (biological replicates), ***$P < 0.001$ using a one-way ANOVA

Figure S6. Growth curves of WT, Δ*esxC*, Δ*esxC* pOS1-*esxC* in TSB supplemented with

increasing calcium chloride concentrations (0 to 20 mM) in the presence of 5 µg/ml daptomycin.

Mean +SE is shown

Figure S7. (A) Growth curves of WT, ∆*essC* and ∆*esxC* in DMEM + FBS. **(B)** Growth curves of WT, ∆*essC* and ∆*esxC* in DMEM + FBS in the presence of 1 µg/ml daptomycin and 1 mM CaCl$_2$. Graphs show mean and SEM of three independent experiments.

Figure S8 The area of lesions was measured and calculated daily from mice infected with *S. aureus* WT or T7SS mutant.. Graph shows mean ± SD. Two-way ANOVA between groups revealed no significant differences between treatments.

Table S1 Secreted proteins significantly changed in ∆*esxC* and ∆*essC* mutants relative to WT USA300 JE2.

Figure 1. An *esxC* mutant is highly sensitive to daptomycin.
A) Growth curves of *S. aureus* WT USA300 JE2 and Δ*esxC* in TSB in the absence or presence of 5 µg/ml daptomycin and 1 mM CaCl$_2$. Mean ± standard error of the mean (SEM) is shown, N = 3 (biological replicates). **B)** CFU/ml was enumerated after 8 h daptomycin treatment. N = 3 (biological replicates), mean + standard deviation (SD), *** P ≤ 0.001 using an unpaired t-test **C)** Growth curves of USA300 JE2 WTpOS1, Δ*esxC*pOS1 and complemented strain, Δ*esxC*pOS1-*esxC* in TSB supplemented with 5 µg/ml daptomycin and 1 mM CaCl$_2$. Mean ± SEM shown, N = 3 (biological replicates); AUC was calculated for the strains grown in presence of daptomycin **(D)** Mean + SD ** P ≤ 0.05 using a one-way ANOVA with Tukey's multiple comparison test. **E)** CFU/ml of WT USA300 JE2 and Δ*esxC* were determined from a killing assay in the presence of 10 µg/ml daptomycin and 1 mM CaCl$_2$ at different timepoints. **F)** CFU/ml from killing assay using WT USA300 JE2, Δ*esxC* and Δ*esxC* complemented strains in the absence or presence of 10 µg/ml daptomycin and 1 mM CaCl$_2$ at different timepoints, mean ± SD, N = 3 (biological replicates), * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001 using a two-way ANOVA with Tukey multiple comparison test.

Figure 2. Membrane permeability of the *esxC* mutant in the presence of daptomycin

A) Representative fluorescent micrographs of *S. aureus* USA300 JE2 WT and ΔesxC treated with daptomycin. Cell membranes were stained by FM1-43 and imaged using FITC filter channel, PI was imaged using Texas red filter channel. **B)** Images of daptomycin-treated samples (A) were quantified using FIJI. Mean fluorescence intensity was quantified from 10 images from each experiment and normalised to the number of cells in the field of view. Data presented is from 3 independent experiments. ****P < 0.0001 using an unpaired t test.

Figure 3. The *esxC* mutant show higher sensitivity to other membrane acting drugs
A) Growth curves of USA300 JE2 WT and Δ*esxC* in the presence of 2 µg/ml gramicidin, Mean ± SEM, N = 3 (biological replicates). **B)** Area under curves (AUC) were calculated and the mean AUC + SD was plotted for growth curves in (A). **C)** Growth curves of USA300 JE2 WT and Δ*esxC* in the presence of 1 µg/ml bithionol. Mean ± SEM, N = 3 (biological replicates). **D)** AUC were calculated and the mean AUC + SD was plotted for growth curves in (C). *P ≤ 0.05 using an unpaired t-test.

Figure 4. EsxC affects membrane fluidity
A) Widefield micrographs of *S. aureus* WT USA300 JE2 and Δ*esxC* after growth in TSB to OD_{600} of 1.0 in the presence of the lipophilic dye DilC12. Images are representative of 4 independent experiments. The DilC12 fluorescence of 80 bacterial clusters from different fields per strain was quantitated with ImageJ. Means + SD are shown, N = 4 (biological replicates); **** P < 0.0001, using an unpaired t-test. **B)** The membrane fluidity of *S. aureus* WT USA300 JE2, Δ*esxC* and Δ*esxC* pOS1-*esxC* as measured with a pyrene decanoic acid staining-based assay. Mean ± SEM are shown, N = 6 (biological replicates), * indicates P < 0.05 using a one-way ANOVA. **C)** USA300 JE2 WT, Δ*esxC* and Δ*esxC* pOS1-*esxC* were grown to logarithmic phase in TSB at 37ºC prior to staining with Laurdan. To assess membrane fluidity, Laurdan generalised polarisation (GP) was calculated using the equation $(I_{460} - I_{500})/ (I_{460} + I_{500})$ where I refers to the fluorescence intensity at the indicated emission wavelength. *** P <0.0001 using a one-way ANOVA, N = 3 (biological replicates), mean + SD.

Figure 5. The *esxC* mutant displays distinct surface morphology.

A) Representative SEM images of *S. aureus* WT USA300 JE2 and *ΔesxC* grown to early logarithmic phase. **B)** Representative SEM images of *S. aureus* USA300 JE2 WTpOS1, *ΔesxC*pOS1, *ΔesxC*pOS1-*esxC* grown to early logarithmic phase.

Figure 6. EsxC impacts cell wall synthesis.
A) Representative widefield micrographs of *S. aureus* WT USA300 JE2 and *ΔesxC* either grown to logarithmic phase in the presence of 25 µM HADA or after 10-minute treatment with HADA once OD_{600} of 1 was reached. **B)** Quantification of fluorescence in cells grown to logarithmic phase with HADA. **C)** Quantification of fluorescence in cells treated with HADA for 10 minutes. Quantification performed on five images each from 3 independent experiments. * P< 0.01, and by an unpaired t test.

Figure 7. EsxC mutant membranes bind more to daptomycin.
Quantitative binding assay of cytochrome C to USA300 JE2 WT, Δ*esxC* and
Δ*esxC*pOS1-*esxC* **(A)**. The CFU of the starting inoculum was calculated for each
mutant **(B)**. Mean ± SD shown, N = 3 (biological replicates), ***P < 0.001 using a
one-way ANOVA. **C)** Fluorescence measurements of daptomycin-BODIPY binding
to *S. aureus* WT USA300 JE2 and Δ*esxC.* The excitation wavelength was 488nm
and emission wavelength was 530nm. N = 3 (biological replicates), mean ± SD. P
= 0.0067, using an unpaired t-test. **D)** Representative micrographs of daptomycin-
BODIPY binding to *S. aureus* WT and Δ*esxC* are shown.

Figure 8. Calcium affects growth of *esxC* mutant and daptomycin sensitivity
Growth curves of *S. aureus* WT USA300 JE2, Δ*esxC*, and Δ*esxC* pOS1-*esxC* in synthetic minimal medium in the absence (A) or presence (B) of 50 mM CaCl$_2$. Mean ± standard error of the mean (SEM) is shown **C)** Area under curves (AUC) were calculated from (A) and (B) and the mean AUC ± SD were plotted. **D)** *S. aureus* WT USA300 JE2, Δ*esxC*, and Δ*esxC* pOS1-*esxC* were grown in TSB supplemented with increasing calcium chloride concentrations (0 to 20 mM) in the presence of daptomycin (5 µg/ml). AUCs calculated from growth curves shown in Figure S6 are shown. Mean + SD, ***P= 0.0001, ****P <0.0001, using two-way ANOVA.

Figure 9. Presence of EsxC affects intracellular bacterial killing by daptomycin.
A) CFU counts at various timepoints during epithelial cell infection assay with WT USA300 JE2, and Δ*esxC*. 0 h represents the initial inoculum, 1 h represents the CFU counts that have infected the A549 cells and 4 h and 24 h represent the survival of *S. aureus* in the presence of 20 μg/ml daptomycin, N = 3 (biological replicates), **P<0.05, two-way ANOVA. **B)** Representative confocal micrographs of cells after 1 h infection or 24 h post-infection. DAPI (blue), F-actin (red), and *S. aureus* (green, white arrows). **C)** Confocal image quantification from five images each from 3 independent experiments after 24 h. Mean ± SD of intensity, is shown, ***P<0.001 using an unpaired t-test.

Figure 10. Effect of daptomycin on *esxC* mutants *in vivo.*

Mice were infected in the skin with WT USA300 and Δ*esxC* mutant followed by treatment with 10 mg/kg daptomycin. **A)** CFU/g skin tissue was counted 24 h post infection (n = 8, 2 lesions/animal). Graphs show mean ± SD, P = 0.0007 using one-way ANOVA. **B)** CFU/g was calculated 72 h post infection (n = 5, 2 lesions/animal), mean ± SD shown, P = 0.0020 using a one-way ANOVA.