CEP44 is required for maintaining centriole duplication and spindle integrity

Donghui Zhang, [1,2,*] Wenlu Wei, [2] Xiaopeng Zou, [2] Hui Meng, [3,4] Fangyuan Li, [1] Minjun Yao, [1]

Junling Teng, [1,*] Ning Huang, [3,4,*] Jianguo Chen[1,5,*]

[1]Key Laboratory of Cell Proliferation and Differentiation of the Ministry of Education, College of

Life Sciences, Peking University, 100871, Beijing, China.

[2]Zhanjiang Institute of Clinical Medicine, Central People's Hospital of Zhanjiang, Guangdong

Medical University Zhanjiang Central Hospital, 524045, Zhanjiang, China.

[3]Institute of Neuroscience, Translational Medicine Institute, Health Science Center, Xi'an Jiaotong

University, Xi'an 710061, China.

[4]Department of Physiology and Pathophysiology, School of Basic Medical Sciences, Health

Science Center, Xi'an Jiaotong University, Xi'an 710061, China.

[5]Center for Quantitative Biology, Peking University, 100871, Beijing, China.

Authorship note: Donghui Zhang and Wenlu Wei are co–first authors.

Conflict of interest: The authors have declared that no conflict of interest exists.

Address correspondence to: Jianguo Chen, Key Laboratory of Cell Proliferation and

Differentiation of the Ministry of Education, College of Life Sciences, Peking University, 100871,

Beijing, China. Phone: 86.010.6275.5786; E-mail: chenjg@pku.edu.cn. Or to: Ning Huang,

Institute of Neuroscience, Translational Medicine Institute, Health Science Center, Xi'an Jiaotong

University, Xi'an 710061, China. Phone: 86.010.6275.5786; E-mail: ning.huang@xjtu.edu.cn. Or

to: Junling Teng, Key Laboratory of Cell Proliferation and Differentiation of the Ministry of

Education, College of Life Sciences, Peking University, 100871, Beijing, China. Phone:

86.010.6275.5786; E-mail: junlinteng@pku.edu.cn; Or to: Donghui Zhang, Key Laboratory of

Cell Proliferation and Differentiation of the Ministry of Education, College of Life Sciences,

Peking   University,   100871,   Beijing,   China.   Phone: 86.010.6275.5786;   E-mail:

Zdh2014@pku.edu.cn;

Abstract

In animal cells, the centrosome, consisting of two centrioles, duplicates only once per cell cycle

for bipolar spindle formation. Defective centriole duplication results in abnormal spindle

formation and chromosome missegregation, which is closely linked to tumor growth. However,

the molecular mechanisms licensing only one centriole duplication cycle within a cell cycle are less

well known. Here we found that CEP44 is negatively correlated with breast carcinoma. CEP44,

jointly with CEP57 and CEP57L1, maintains centriole engagement in the interphase to ensure

centriole duplication once per cell cycle. Depletion of CEP44 leads to centriole overduplication

because of premature centriole disengagement, and multipolar spindle formation. Additionally,

CEP44 is phosphorylated by Aurora A at the G2/M phase to facilitate spindle localization and

maintain spindle integrity. Collectively, our results show the function of CEP44 in spindle

formation by preventing centriole overduplication and maintaining spindle integrity, and CEP44

may serve as a potential marker for breast carcinoma prognosis.

Introduction

The centrosome, at the center of which is a pair of centrioles, is surrounded by pericentriolar material (PCM) and centriolar satellites, and acts as the major microtubule organizing center (MTOC) in mammalian cells (1). In mitosis, the two centrosomes move to the opposite side in a cell and function in the nucleation of microtubules important for forming the bipolar spindle to ensure even segregation of genetic material. Therefore, centriole duplication, occurring only once per cell cycle, is strictly initiated through assembly of the procentriole at the proximal parts of preexisting centrioles during the S phase (2-4). Anomalies in centriole duplication caused by improper centrosome number control lead to disrupted spindle formation and chromosomal segregation errors, and eventually culminate in cancer and other developmental diseases (5-8).

The molecular mechanisms controlling only one centriole duplication cycle within a cell cycle are not well understood, despite significant advancements in understanding copy number control of centriole duplication in the S phase (3). Maintenance of centriole engagement in the interphase has been reported to be fundamental for ensuring that centriole duplication occurs only once per cell cycle (9). Immediately after centriole duplication, the newly formed daughter centriole is perpendicularly engaged with preexisting centrioles until late mitosis (centrosome engagement). After mitotic exit, the close-connected mother and daughter centrioles are separated (centrosome disengagement), and this separation allows for centriole duplication in the next cell cycle. When premature centriole disengagement happens as early as the S phase, centrioles are reduplicated, producing numerous centrioles within the same cell cycle (10).

It has been reported that PCM plays a pivotal role not only in microtubule nucleation, but also in modulation of centriole engagement and disengagement both in mitosis and interphase (11).

Pericentrin (PCNT), a PCM scaffold protein, is critical for ensuring the maintenance of centriole

engagement in mitosis. For normal centriole disengagement during mitotic exit, PCNT is

specifically cleaved by separase to disassemble the expanded PCM (12). PLK1-dependent

phosphorylation of PCNT is a prerequisite for the cleavage of PCNT by separase (13).

CEP57, a PCM protein, is closely associated with tumor development and variegated

aneuploidy syndrome (14, 15). As a crucial binding partner of PCNT, CEP57 serves as a PCM

organization modulator (16); CEP57 depletion causes premature centriole disengagement in

mitosis (17). Notably, co-depletion of CEP57 with its paralog CEP57L results in premature

centriole disengagement in interphase, resulting in centriole reduplication and disturbed

chromosome segregation by amplified centrioles (9). Thus, centriole engagement in interphase is

of much greater importance than that in mitosis for centriole duplication and proper chromosome

segregation. However, the molecular mechanisms underlying centriole engagement in interphase

are largely unknown, although there has been some progress in understanding the basis of

centriole engagement in mitosis.

A centrosomal protein of 44 kDa (CEP44) was first identified as a centrosome protein, and

probable spindle protein, via complementary proteomics (18). It has been reported that CEP44 is a

centrosome linker protein that functions in centrosome cohesion (19). Recent studies also

demonstrate that CEP44, as a centriole luminal protein, plays a vital role in the centriole-to-

centrosome conversion program by promoting the formation of structurally-intact centrioles with a

9-fold symmetric triplet microtubule structure (20). In this study, we show that the expression of

CEP44 is negatively correlated with breast carcinogenesis. CEP44 ensures centriole duplication

once per cell cycle by maintaining centriole engagement with CEP57 and CEP57l1 in interphase.

In mitosis, CEP44 is required for spindle integrity and is activated by Aurora A-mediated

phosphorylation.

Results

CEP44 is negatively correlated with breast carcinogenesis

Several studies have conclusively demonstrated that certain centrosome proteins act as powerful

tumor suppressors or activators (21-24). The centrosomes in breast cancer cells show

characteristic structural and functional aberrations attributed to dysregulation of important

centrosome or non-centrosome proteins (8, 25-31). To reveal the mechanisms underlying

intracellular centrosome structure and function in breast cancer cells, we conducted a search for

new centrosome proteins that might function in intrinsic mechanisms of centrosome structure and

function in breast cancer cells. Intriguingly, MDA-MB-436 cells showed a nearly total loss of

CEP44 protein expression and centrosome localization (Figure 1A-C). Human CEP44 consists of

390 amino acids (aa) containing two short coiled-coil domains in the middle region and C-

terminus (Figure 1–figure supplement 1A). To confirm the specificity of the antibodies against

CEP44, CEP44 was depleted by CEP44 siRNAs and detected by immunoblots. Both our

homemade and commercial antibodies recognized a band at approximately 45 kDa by

immunoblotting in HeLa cells, and this band was specifically abolished by siRNAs targeting

CEP44 (Figure 1–figure supplement 1B and C).

Moreover, the expression data obtained from TCGA showed lower expression of CEP44 mRNA

in breast cancer patients in comparison with normal individuals (Figure 1D), as well as in breast

cancer tissue in comparison with normal tissue (Figure 1E). Consistently, immunohistology

analysis of CEP44 expression levels using human breast tissue arrays that included paired tumor

and tumor-adjacent tissues showed that CEP44 was expressed at a low level in breast carcinoma

samples (Figure 1F-G). Analysis of data from the cancer genome atlas program (TCGA) cohort

also revealed that lower CEP44 expression in breast cancer patients was associated with worse

overall survival (OS) and recurrence free survival (RFS) outcomes (Figure H-I). In conclusion,

these observations suggest a vital role for CEP44 in breast carcinogenesis.

Detailed localization of CEP44 by stimulated emission depletion (STED) microscopy showed

that CEP44 formed a small ring-like structure, enclosed by the circle marked by polyglutamylated

and acetylated tubulin, a centriole wall marker (from the top view), and localized tightly along the

inner side of the circle (from the side view) (Figure 1–figure supplement 1D-G), suggesting that

CEP44 is a luminal centriole protein as previously reported (20). 3D-structured illumination

microscopy (3D-SIM) showed that CEP44 colocalized with C-Nap1, a marker of the centrosome

linker at the proximal ends, more closely than it colocalized with CEP164 or GFP-Centrin-3 (both

markers of the distal ends of centrioles), suggesting that CEP44 is localized at the proximal halves

of centrioles. Co-staining of CEP44 with CEP152 (localized to the proximal halves of centrioles)

and CEP192 (localized along the entire walls of centrioles) (32) revealed that CEP44 was partly

colocalized with CEP192 and encircled by CEP152 (Figure 1–figure supplement 1H),

demonstrating that CEP44 is localized to the proximal halves of centrioles.

CEP44 is a negative regulator of centriole duplication

As reported previously (33, 34), centrioles are remarkably overduplicated in breast cancer cells

such as MDA-MB-436, MDA-MB-231, SKBR3 and HCC1806 cells, but not in the spontaneously

immortalized, non-malignant breast cells MCF10A (Figure 2A). In particular, centriole

overduplication was found to be particularly severe in the MDA-MB-436 cells (approximately

80%) (Figure 1B and Figure 2B), suggesting that it CEP44 may play critical role in regulating

centriole duplication in breast cancer cells.

To investigate the function of CEP44 in centriole duplication, we performed siRNA knockdown

experiments with two CEP44-targeting siRNAs in HeLa cells and observed remarkably increased

centriole amplification (more than four centrioles marked by CEP97 or Centrin-3), which was

rescued by overexpressing siRNA-resistant CEP44 (Figure 2C-F).

To investigate whether the increased number of centrioles in MDA-MB-436 cells is attributed a

loss of CEP44, centriole numbers in MDA-MB-436 cells were recalculated after CEP44 ectopic

overexpression. The fraction of cells showing centriole overduplication significantly decreased to

approximately 40% in CEP44 overexpressed MDA-MB-436 cells (Figure 2G-H), demonstrating

that CEP44 plays an important role in controlling centriole duplication in MDA-MB-436 cells.

CEP44 maintains centriole engagement

To dissect the mechanism of centriole overduplication after CEP44 depletion, we observed

centrosome phenotypes. Firstly, we generated CEP44 knockout HeLa cell lines using a CRISPR-

Cas9 system (Figure 3–figure supplement 1A-B). CEP44-depleted cells showed increased distance

(>0.75 μm) between newly formed daughter centriole and old centriole after centriole duplication

in interphase and mitosis (Figure 3A-C), indicating that precocious centriole disengagement

occurred after CEP44 depletion. To exclude the possibility that the four separated centrioles

observed in interphase after CEP44 depletion resulted from failed cytokinesis, we determined the

number of old mother centrioles; if cytokinesis failed, then two old mother centrioles were

expected. Immunostaining with two markers of old mother centrioles, CEP164 and ODF2,

revealed the presence of only one dot marked by CEP164 and ODF2 in ~70% of CEP44-depleted

cells with four separated or amplified centrioles (Figure 3–figure supplement 1C-D). These

findings suggest that the separated centrioles were likely caused by premature centriole

disengagement instead of cytokinesis failure.

Given that depletion of CEP44 resulted in both centriole disengagement and overduplication

(Figure 2C-F, 3A-C and Figure 3–figure supplement 1E-H), we performed experiments to

elucidate the mechanisms underlying these events. Firstly, we used an S phase marker (5-ethynyl-

29-deoxyuridine, EdU) and a G2 phase marker (centromere protein F, CENP-F) to observe the

occurrence of centriole disengagement and overduplication during particular stages of the cell

cycle. In the G1 phase (Edu and CENP-F negative), CEP44-depleted cells and control cells both

exhibited two centrioles (Figure 3D-F). However, CEP44-depleted cells exhibited a remarkably

increased percentage of cells with four or more separated centrioles in the S phase (Edu positive

and CENP-F negative) and G2 phase (Edu negative and CENP-F positive) (Figure 3D-F). These

results verify that CEP44 depletion results in centriole disengagement as early as the S phase, and

thus CEP44 knockout cells exhibit four or more separate centrioles. Notably, centriole

overduplication also occurred as early as the S phase after CEP44 depletion. However, these

results did not show the sequential order of centriole disengagement and overduplication caused

by CEP44 depletion, as both of these events occurred in the S phase. Therefore, to address the link

between these events, centriole dynamics were monitored in cells transfected with GFP-centrin-3

and H2B-RFP by tracing centrioles with time-lapse microscopy. In control cells, maintenance of

centriole engagement was observed after centriole duplication, whereas CEP44 knockout cells

showed premature centriole disengagement followed by centriole overduplication per cell cycle

(Figure 3G-H), suggesting that premature centriole disengagement in interphase contributes to

centriole overduplication. It is worth noting that centriole overduplication after premature

centriole disengagement resulting from CEP44 depletion generally led to the formation of only

one newly formed daughter centriole, instead of excessive centrioles, from each pre-existing

centriole (Figure 3H).

As the daughter centriole is formed, PCM is recruited to it to ensure initiation of the first stage

of centrosome duplication, referred to as centriole-to-centrosome conversion (CCC). It has been

reported that CEP44 is required for CCC; therefore, it is conceivable that the CCC of precociously

disengaged daughter centrioles was impaired in CEP44-depleted cells in interphase, which may

have impaired centriole duplication, rather than inducing centriole overduplication, as shown by

CEP44-depleted cells in our assay. To address this paradox, we co-stained γ-tubulin and centrin-3

in CEP44 knockout cells. CEP44-depleted cells with more than 2 γ-tubulin-positive centrioles

were quite rare before the G2 phase, but they gradually became more abundant during the G2

phase, during which cells exhibited two dots with strong γ-tubulin staining, and one or two dots

with weak γ-tubulin staining. The light dots stained by γ-tubulin most likely represent the

disengaged daughter centrioles. After centriole duplication in CEP44-depleted cells, we also

detected recruitment of γ-tubulin in the reduplicated centrioles (Figure 3–figure supplement 1I-J).

These findings demonstrate that PCM is initially absent in the precociously disengaged daughter

centrioles, and PCM components are gradually recruited to their surroundings, primarily during

the G2 phase.

CEP44 maintains centriole engagement via CEP57 and CEP57L1

To determine the molecular mechanism underlying centriole engagement regulation by CEP44, we searched for proteins capable of interacting with CEP44. Our immunoprecipitation assay showed that there was no interaction between CEP44 and several centriole engagement modulators, including PCNT, Sgo1 and SCC1, but the interaction was detected between CEP44 and CEP57/CEP57L (Figure 4A-B). Further immunoprecipitation assays showed that there was no interaction between CEP44 and several centriole luminal proteins known to be important for centriole duplication, including SAS6, STIL, PLK4 and CEP135. Given that CEP57/CEP57L1 is known to be involved in controlling centriole disengagement in interphase (9), these results suggest that the molecular mechanism by which CEP44 regulates centriole engagement is likely mediated by its interaction with CEP57/CEP57L1. Thus, we further examined the relationship among CEP44, CEP57, and CEP57L1. To determine whether CEP44 affects CEP57/CEP57L1 expression or localization, we first knocked down CEP44 expression in HeLa cells and observed CEP57/CEP57L1 expression and localization. Although depletion of CEP44 by two different siRNAs did not affect CEP57/CEP57L1 expression (Figure 4C), localization of CEP57/CEP57L1 at centrosomes was significantly reduced (Figure 4D-F). As previously reported (20), we also observed impaired γ-tubulin localization at centrosomes in CEP44-depleted cells (Figure 4D and Figure 4–figure supplement 1A-B). Conversely, depletion of CEP57 or CEP57L1 did not significantly affect CEP44 localization at centrosomes (Figure 4G-H and Figure 4–figure supplement 1A). Similar to the expression pattern of CEP44 in breast cancer patients vs normal individuals, the expression data obtained from TCGA showed lower expression of CEP57 mRNA

in patients with breast cancer compared with normal individuals (Figure 4–figure supplement 1C),

as well as in breast cancer tissue compared with normal tissue (Figure 4–figure supplement 1D).

Taken together, these results suggest that CEP44 maintains centriole engagement by regulating

CEP57/CEP57L1 localization at the centriole.

CEP44 depletion induced centriole overduplication leads to multipolar spindle formation

Next, we found that the percentage of multipolar and disorganized spindles increased

remarkably in mitotic CEP44 knockout cells. This phenotype was nearly fully rescued by re-

introduction of CEP44-Flag (Figure 5A-C). Similar phenotypes of increased percentages of

multipolar and disorganized spindles in CEP44 knockdown HeLa cells were rescued by

overexpressing siRNA-resistant CEP44 (Figure 5–figure supplement 1A-D).

Given that depletion of CEP44 led to centriole amplification, we speculated that multipolar

spindles in mitotic cells that formed after depletion of CEP44 were attributed to the presence of an

elevated number of centrioles. To test our hypothesis, we stained mitotic HeLa cells with centriole

marker CEP97 and calculated the centriole number. We observed an increased number of

centrioles in multipolar spindles following depletion of CEP44 (Figure 5D-E and Figure 5–figure

supplement 1E-F). These results indicate that depletion of CEP44 induced centrosome

overduplication and abnormal spindle formation.

As mentioned above, CEP44-depleted HeLa cells exhibited a remarkably increased percentage

of multipolar spindles (Figure 5A-B and Figure 5–figure supplement 1B-C). To determine

whether centriole disengagement or/and centriole overduplication in interphase led to the

increased percentage of multipolar spindles in CEP44 knockout cells, we transfected cells with

GFP-centrin-3 and H2B-RFP and observed centrioles and chromosomes by live cell imaging.

Based on the pattern of centriole disengagement and overduplication, we grouped cells into three

categories in interphase: normal, premature centriole disengagement in interphase without

centriole overduplication (pattern 1), or with centriole overduplication (pattern 2). In control cells,

the two paired centrioles were engaged during the entire interphase (normal); while in CEP44

knockout cells, paired centrioles disengaged in interphase without centriole overduplication

(pattern 1) or with centriole overduplication (pattern 2). Quantification of the percentage of

multipolar spindles in cells showing pattern 1 or pattern 2 revealed that CEP44 knockout cells

associated with centriole disengagement and centriole overduplication (pattern 2) exhibited the

highest frequency of multipolar spindles (Figure 5F and Figure 5–figure supplement 1G). These

results suggest that the formation of multipolar spindles after CEP44 depletion is mainly attributed

to centriole overduplication resulting from premature centriole disengagement.

Aurora A phosphorylates CEP44 in mitosis

Apart from its localization at centrosomes, CEP44 was also identified as a probable spindle

protein in a complementary proteomics study (18). Thus, we determined whether CEP44 localizes

at spindles, and found that it was readily detected at mitotic spindles throughout mitosis (Figure

6A and Figure 6–figure supplement 1A). To determine the pattern of CEP44 protein level changes

during the cell cycle, we synchronized HeLa cells with nocodazole treatment. CEP44 protein

levels peaked in the enriched M phase population (Figure 6B), similar to the well-known mitotic

regulator cyclin B, indicating that CEP44 plays a critical role in mitosis. Multiple kinases,

including Aurora A, PLK1, NEK2A, and CDK1, are localized to the spindle or centrosome and

serve to regulate spindle dynamics during mitosis (35-38). To test whether these kinases can

phosphorylate CEP44 to exert their function in mitosis, we firstly co-overexpressed CEP44 with

these mitosis-related kinases in HEK293T cells. The band representing CEP44 was shifted slightly

when CEP44 was co-overexpressed with wild-type Aurora A, but not kinase-dead Aurora A

(Figure 6C and Figure 6–figure supplement 1B). However, this shifted band was completely

abolished when the cell lysate was pretreated with lambda protein phosphatase (λ-PPase) (Figure

6C). Moreover, treatment with Aurora A inhibitor MLN8237 also abolished the shifted CEP44

band at the G2/M phase (Figure 6D and Figure 6–figure supplement 1C), indicating that CEP44 is

phosphorylated by Aurora A at the G2/M phase. Mass spectrometry analysis revealed three sites

that might be phosphorylated by Aurora A: Ser192, Ser324 and Ser342. To confirm the CEP44

phosphorylation site for Aurora A, we constructed unphosphorylatable mutants in which Ser192,

Ser324 or Ser342 was replaced by alanine (S192A, S324A or S342A) using site-directed

mutagenesis. The shifted band of CEP44-S324A was barely detected in cells co-overexpressing

CEP44 and Aurora A (Figure 6E), indicating that CEP44 Ser324 is the site of Aurora-A-mediated

phosphorylation. Additionally, Aurora A is known to recognize the consensus sequence R-X-

pS/T-L/V for phosphorylation (39-42), and Ser324 of CEP44 accords with the consensus sequence

(Figure 6F), confirming that Ser324 of CEP44 is phosphorylated by Aurora A. We then raised a

phospho-CEP44 antibody against CEP44 Ser324 and verified the specificity of the antibody

through peptide competition assays (Figure 6–figure supplement 1D). A band was detected when

nocodazole-treated HeLa cell protein samples were incubated with the phospho-CEP44 antibody,

but no band was detected from the control HeLa cells (Figure 6G). Moreover, depletion of Aurora

A resulted in the loss of the band detected with the phospho-CEP44 antibody in nocodazole-

synchronized HeLa cells (Figure 6H). Collectively, these results suggest that Aurora A

phosphorylates CEP44 at Ser324 in the G2/M phase.

The results showing that CEP44 is highly expressed and phosphorylated during mitosis (Figure

6B-D) suggest that phosphorylation of CEP44 by Aurora A might alter its protein level. However,

the CEP44 protein level did not change significantly after Aurora A knockdown in the G2/M

phase (Figure 6H and Figure 6–figure supplement 1E). Next, we further examined the effect of

Aurora-A-mediated phosphorylation on CEP44 localization. Immunostaining microscopy showed

that mitotic Aurora A-depleted cells exhibited remarkably increased fluorescence intensity of

CEP44 at the mitotic spindle pole, but less fluorescence intensity at the spindle, compared to the

control cells (Figure 6I-J). This abnormal distribution of CEP44 in mitosis after Aurora A

depletion was also detected in HeLa cells after treatment with MLN8237 (Figure 6–figure

supplement 1F-H), suggesting that Aurora A-mediated phosphorylation of CEP44 regulates its

localization between the spindle pole and spindle.

CEP44 phosphorylation by Aurora A maintains spindle integrity

To clarify the physiological function of CEP44 and its phosphorylation at the spindle, we

immunostained mitotic HeLa cells with α-tubulin to label spindle microtubules. Depletion of

CEP44 significantly decreased the intensity of spindle microtubule staining, which was rescued

after re-introduction of CEP44 (Figure 7A-B and Figure 7–figure supplement 1A-B), indicating

that CEP44 regulates spindle microtubule stability. This finding suggests that CEP44

phosphorylation may be responsible for spindle microtubule stability, because CEP44 is

phosphorylated in the G2/M phase. To determine whether CEP44 phosphorylation affects

microtubule stability, CEP44 phosphorylation was mimicked in interphase HeLa cells by

overexpressing phosphorylatable (CEP44-Flag S324D and S324E) or unphosphorylatable

(CEP44-Flag S324A) mutants. Microtubule stability was not dramatically altered by

overexpression of the phosphorylatable or unphosphorylatable mutants. However, after treatment

with nocodazole for 5 min, HeLa cells overexpressing CEP44-Flag S324A exhibited more un-

stabilized microtubules in comparison with HeLa cells overexpressing CEP44-Flag wild-type,

S324D or S324E (Figure 7–figure supplement 1C), suggesting that CEP44 phosphorylation plays

an important role in controlling microtubule stability. Similarly, in mitotic HeLa cells, ectopic

overexpression of CEP44-Flag S324A remarkably impaired spindle microtubule stability (Figure

7C-E). Collectively, these results show the critical role of CEP44 phosphorylation in maintenance

of microtubule stability and spindle structure. Notably, we found that CEP44-depleted cells

showed weakened localization of γ-tubulin at the spindle (Figure 7F-G), which negatively affected

spindle structure stability.

Next, we investigated whether chromosome segregation errors occurred in mitotic CEP44-

depleted cells due to impaired spindle structure (Figure 7A-B and 7F-G). CEP44-depleted cells

showed a dramatically increased proportion of chromosome segregation errors, including

misalignment and lagging chromosomes (Figure 7H-I), although bipolar spindle formation was

accomplished eventually in some CEP44-depleted cells (Figure 7–figure supplement 1D).

Therefore, CEP44 phosphorylation by Aurora A ensures normal spindle structure formation and

chromosome segregation.

Discussion

Our study indicates that centriole luminal protein CEP44 is a master regulator of centriole

disengagement in interphase via its interaction with CEP57/CEP57L1, and its depletion results in

premature centriole disengagement, centriole overduplication and multipolar spindle formation.

Additionally, CEP44 is a spindle-localized protein that ensures spindle integrity, and this

localization is modulated by Aurora A phosphorylation. Collectively, our results show that CEP44

maintains correct spindle formation through its localization at the centrosome and spindle (Figure

8).

In contrast to the effects of co-depletion of PCM proteins CEP57 and CEP57L1, depletion of

CEP44 alone was sufficient to result in premature centriole disengagement in interphase. A

previous study reported that depletion of CEP44 impaired centrosome localization of PCM

proteins, such as CEP152 and CEP192, which are required for centriole duplication (20). However,

we showed centriole overduplication caused by premature centriole disengagement in CEP44-

depleted cells, indicating that the effect of premature centriole disengagement in interphase, which

accelerates centriole overduplication, overrides the effect of defective PCM recruitment (impaired

CCC), which impedes centriole duplication, in the context of CEP44 depletion. These findings

suggest that the residual PCM around prematurely disengaged centrioles is sufficient to induce

centriole duplication after CEP44 deficiency. Collectively, our findings confirm that centriole

disengagement is a critical prerequisite for a subsequent round of centriole duplication, indicating

the fundamental role of centriole engagement/disengagement in modulating centriole duplication.

The centriole number is controlled based on two distinct rules to coordinate multiple cellular

processes (2, 4); the first rule ensures that centrosomes duplicate once and only once per cell cycle

(cell cycle control), and the second rule stipulates that only one progeny centriole may form

adjacent to each parental centriole in the S phase (copy number control). In contrast to

encouraging recent progress in understanding copy number control of centriole duplication in the

S phase, the mechanisms licensing only one centriole duplication cycle within a cell cycle are not

well understood. However, our study shows the great importance of CEP44 in controlling the

centriole number during cell cycle progression, providing important insight into the detailed

mechanisms underlying centriole duplication.

It also has been reported that CEP44 functions in centrosome cohesion by stabilizing rootletin

(19). Centrosome cohesion is mediated by a loose connection between the proximal ends of the

two parental centrioles, which is known as the G1-G2 tether (GGT). GGT assembly occurs in the

G1 phase, which occurs simultaneously with, or just after, dissolution of the centriole engagement

component termed the S-M linker (SML). Therefore, it has been speculated that GGT assembly

depends on SML dissociation (4), but few studies have assessed the relationship between the SML

and GGT. Apart from the function of CEP44 in centrosome cohesion, we found that CEP44 is

involved in centriole engagement, suggesting the existence of some relationship between the SML

and GGT, and additional studies will likely further illuminate the close relationship between the

SML and GGT.

The localization and function of CEP44 led us to speculate that CEP44 may be a part of the

SML. Previous studies found that SML functionality depends on the SAS-6–ANA2 complex (43)

and suggested that cohesin might constitute part of the SML (44). However, the composition of

the SML remains poorly understood. In this study, we found that CEP44 does not interact with the

SAS6-STIL complex or cohesin (data not shown). Based on our results, it is plausible that the

luminal protein CEP44 in the proximal part of the procentriole interacts with PCM protein

CEP57/CEP57L1 in pre-existing centrioles to maintain centriole engagement, indicating that

CEP44 is a potential SML component. Future studies aimed at determining whether CEP44 is a

component of the SML will shed new light on the molecular mechanisms regulating centriole

engagement.

We found that CEP44 is highly expressed in G2/M phase HeLa cells and nearly absent in

MDA-MB-436 breast cancer cells, indicating that the protein level of CEP44 is precisely

modulated during different stages of the cell cycle and in specific cell types. However, Aurora A-

mediated phosphorylation of CEP44 had no effect on the protein level of CEP44 in our assay

(Figure 6–figure supplement 1E). It is possible that epigenetic mechanisms or post-translational

modifications function to regulate the protein level of CEP44.

Centrosome proteins have been reported to represent potent tumor suppressors or activators,

because centrosome amplification is extremely common in a wide range of malignancies,

including breast, prostate, ovarian, and lung tumors, and it is closely linked to tumor growth and a

worse prognosis (45). Preclinical research has shown that extra centrosomes can directly cause

tumor initiation, at least partly by counteracting centrosome functionality-decreasing mutations or

by increasing chromosome instability through transiently formed abnormal spindles that result in

missegregation because of deficiency in mitotic kinetochore-spindle attachments (46, 47).

Therefore, due to insufficient CEP44, the presence of supernumerary centrosomes and abnormal

spindle structure may compensate for impaired centrosome functionality of CCC during breast

cancer initiation. Thus far, anti-mitotic drugs such as Aurora A and Plk1 inhibitors have largely

failed in clinical contexts due to their low efficacy and serious side effects (48, 49). Notably,

defective mitosis targeted by therapeutic strategies is always induced by centrosome amplification;

therefore, it should be noted whether patients show centrosome amplification, and whether these

inhibitors affect centrosome amplification, to ensure the achievement of a precise anti-tumor effect.

Materials and Methods
398

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*Homo-sapiens*)	KIAA1712	GenBank	NM_001040157.3	
cell line (*Homo-sapiens*)	HEK293T	ATCC	CRL-11268G-1	
cell line (*Homo-sapiens*)	HeLa	ATCC	CCL-2	
cell line (*Homo-sapiens*)	MDA-MB-436	ATCC	HTB-130	
cell line (*Homo-sapiens*)	MDA-MB-231	ATCC	CRM-HTB-26	

cell line (*Homo-sapiens*)	MCF10A	ATCC	CRL-10317	
transfected construct (*M. musculus*)	4E-BP1 shRNA #1	VectorBuilder	Cat#LVS (VB181217-1124dqm)-C	Lentiviral construct to transfect and express the shRNA.
antibody	anti-CEP44 (Rabbit polyclonal)	Genetex	Cat# GTX32692	IF (1:200), WB (1:500)
antibody	anti-CEP44 324 (mouse polyclonal)	Abclonal	This Paper	WB (1:300)
antibody	anti-CEP57L1 (Rabbit polyclonal)	Proteintech	Cat# 24957-1-AP	IF (1:200), WB (1:500)
antibody	anti-CEP57 (Rabbit polyclonal)	Genetex	Cat# GTX115931	IF (1:200), WB (1:500)
antibody	anti- Cyclin B1 (Rabbit polyclonal)	Cell signaling	Cat#4138	WB (1:500)
antibody	anti-CEP192 (Rabbit polyclonal)	Proteintech	Cat# 18832-1-AP	IF (1:200)
antibody	anti-CPAP (Rabbit polyclonal)	Proteintech	Cat# 11517-1-AP	WB (1:500)
antibody	anti-CEP192 (Rabbit polyclonal)	Bethyl	Cat# A302-324A	IF (1:200)

antibody	anti-CEP152 (Rabbit polyclonal)	Proteintech	Cat# 21815-1-AP	IF (1:200)
antibody	anti- CEP97 (Rabbit polyclonal)	Proteintech	Cat# 22050-1-AP	IF (1:200)
antibody	anti- C-Nap1 (Mouse monoclonal)	Santa Cruz	Cat# sc-390540	IF (1:200)
antibody	anti- Centrin-3 (Mouse monoclonal)	Abnova	Cat# H00001070-M01	IF (1:200)
antibody	anti-Polyglutamylation Modification (Mouse monoclonal)	Adipogen	Cat# AG-20B-0020-C100	IF (1:200)
antibody	anti- SAS6 (Mouse monoclonal)	Santa	Cat# SC-98506	IF (1:200)
antibody	anti- Aurora A (Rabbit polyclonal)	Cell signaling	Cat# 14475	IF (1:200), WB (1:500)
antibody	anti-GAPDH (Mouse monoclonal)	Proteintech	Cat# 60004-1-Ig	WB (1:2000)
antibody	anti-α-tubulin (Mouse monoclonal)	Sigma-Aldrich	Cat# T6199	IF (1:1000), WB (1:5000)
antibody	anti-γ-tubulin (Mouse monoclonal)	Sigma-Aldrich	Cat# T6557	IF (1:200)

antibody	anti-Flag (Mouse monoclonal)	Sigma-Aldrich	Cat# F9291	IF (1:500), WB (1:5000)
antibody	anti-γ-tubulin (Rabbit polyclonal)	Sigma-Aldrich	Cat# T3559	IF (1:200)
antibody	anti-CENP-F (Rabbit monoclonal)	Abcam	Cat# ab223847	IF (1:200)
antibody	anti-Acetylated-tubulin (Mouse monoclonal)	Sigma-Aldrich	Cat#: T7451	IF (1:200)
antibody	anti-GFP (Rabbit polyclonal)	Proteintech	Cat# 66002-1-Ig	WB (1:2000)
antibody	Goat anti-Rabbit IgG (H+L) Secondary Antibody, Alexa Fluor 488	Invitrogen	Cat# A-11034	IF (1:200)
antibody	Goat anti-Mouse IgG (H+L) Secondary Antibody, Alexa Fluor 488	Invitrogen	Cat# A-11029	IF (1:200)
antibody	Goat anti-Rabbit IgG (H+L) Secondary Antibody, Alexa Fluor 568	Invitrogen	Cat# A-11036	IF (1:200)

antibody	Goat anti-Mouse IgG (H+L) Secondary Antibody, Alexa Fluor 568	Invitrogen	Cat# A-11031	IF (1:200)
recombinant DNA reagent	pEGFP-N1 (plasmid)	Clontech Laboratories		
recombinant DNA reagent	p3×FLAG-CMV-14 (plasmid)	Sigma-Aldrich		
sequenced-based reagent	siRNA: negative control	tsingke		UUCUCCGA ACGUGUGU CACGU
sequenced-based reagent	siRNA: CEP44#1	tsingke		GGCUAUAG UACAGCAU CAUTT
sequenced-based reagent	siRNA: CEP44#2	tsingke		GGUGAUUC GUCACUUG UAUTT
sequenced-based reagent	siRNA: CEP57#1	tsingke		AAGCAUGC AGAAAUGG AGAGG
sequenced-based reagent	siRNA: CEP57#2	tsingke		AACCAUCA AGGUCUAA UGGAA
sequenced-based reagent	siRNA: CEP57L1#1	tsingke		CGUUGUAC UCUUCUAG AGATT
sequenced-based reagent	siRNA: CEP57L1#2	tsingke		GCUUCAAA CUGGACUU GAATT

sequenced-based reagent	siRNA: Aurora A#1	tsingke		AUGCCCUG UCUUACUG UCATT
sequenced-based reagent	siRNA: Aurora A#2	tsingke		UUCUUAGA CUGUAUGG UUATT
commercial assay or kit	In-Fusion HD Cloning	Clontech	Clontech:6396 47	
chemical compound, drug	Aurora A inhibitor-MLN8237	Selleck	S1133	
chemical compound, drug	PLK1 inhibitor-BI2536	Selleck	S1109	
chemical compound, drug	CDK5 inhibitor-Roscovitine	Selleck	S1153	
software, algorithm	GraphPad Prism 5	GraphPad Prism 5	version 5.01	
software, algorithm	Image J	NIH		

Ethics statement

Breast tissue samples were collected during surgical treatment at the Oncology Department of the

First Affiliated Hospital of Xinxiang Medical University (Henan, China). All samples were

derived from idiopathic breast cancer patients, and none of these patients had received any

treatment before surgery. These fresh resected biopsies of breast cancer were 4% formalin-fixed

for immunohistochemical and pathological grading studies. Written informed consent for the

samples obtained under protocols was approved by the Ethics Committee of the First Affiliated

Hospital of Xinxiang Medical University (XYLL-2021B001). Diagnosis of all samples was

confirmed by at least two independent pathologists.

Plasmid construction

Human CEP44 (NM_001040157.3 [https://www.ncbi.nlm.nih.gov/nuccore/NM_001040157.3]) ,

CEP44 fragments, and CEP44 mutants were amplified via PCR and cloned into pEGFP-N1

(Clontech Laboratories) and p3×FLAG-CMV-14 (Sigma-Aldrich). Other plasmids used in this

study, including Centrin-3, Aurora A, Plk1, Nek2A, and CDK1 were generated as described

previously (50).

Antibodies

All antibodies used in this study are listed in the Key Resources Table. The mouse polyclonal

antibody against CEP44 was generated using full-length human CEP44 fused with glutathione-S-

transferase (GST). Anti-phosphorylated S324-CEP44 antibodies were raised in rabbits using a

keyhole limpet hemocyanin (KLH)-conjugated DLLNPHRK(pS)EVERPAS peptide, which was

affinity-purified on nonphosphorylated- and/or phosphorylated-epitope-bound columns (ABclonal

Biotechnology).

Cell culture and transfection

HEK293T (CRL-11268G-1), MCF10A, MDA-MB-436, MDA-MB-231and HeLa (CCL-2) cells

were obtained from the American Type Culture Collection (ATCC). HEK293T (CRL-11268G-1),

HeLa (CCL-2), MDA-MB-436 and MDA-MB-231 cells were cultured in DMEM (GIBCO)

supplemented with 10% FBS (GIBCO or CellMax) at 37 ℃ under 5% CO2, with MDA-MB-436

cells adding 10 µg/ml insulin and 16 µg/ml glutathione. MCF10A was cultured in a base medium

from Lonza/Clonetics Corporation (MEGM, Kit Catalog No. CC-3150) supplemented with 5%

horse serum and 100 ng/ml cholera toxin. To avoid mycoplasma contamination, mycoplasma

detection was implemented in these cell lines using GMyc-PCR Mycoplasma Test Kit (YEASEN,

40601ES20). Plasmids were transfected into HEK293T and HeLa cells using jetPEI (Polyplus

transfection) or LipofectamineTM 3000 (Invitrogen), and siRNAs were transfected into HeLa

cells using LipofectamineTM 2000, according to the manufacturer's protocols.

Cell synchronization and drug treatment

Synchronized HeLa cells in mitosis were treated with 100 ng/ml Nocodazole for 16-18 h. For

double-thymidine blocking, HeLa cells were firstly subjected to 2.5 mM thymidine for 18-24 h,

then released for 12 h, and subsequently blocked again for 18-24 h. To identify the kinase of

CEP44, HeLa cells were mock-treated or treated with a kinase inhibitor: 100 nM MLN8237 (an

Aurora A inhibitor) for 24 h, 100 nM BI2536 (a PLK1 inhibitor) for 4 h or 25 µM Roscovitine (a

CDK5 inhibitor) for 18 h.

Gene silencing by siRNA

All siRNAs used in this study were purchased from GenePharma. CEP57- and CEP57L1-siRNAs

were designed according to a previous study (16, 51). The siRNA sequences are shown in Key

Resources Table. The siRNA transfection procedure was performed for 48-72 h based on the

manufacturer's protocols. For the rescue assay, six silent mutations were introduced into the

CEP44 sequence to construct siRNA-resistant CEP44 (resCEP44) (5'-

GCTATAGTACAGCATCAT-3' mutated to 5'-GATACAGCACCGCTTCATC-3'). HeLa cells

were transfected with resCEP44 for 48 h after 24 h siRNA transfection.

Western blotting

The samples were boiled at 100°C for 5 min in SDS loading buffer and then loaded in SDS-PAGE

gels. The resolved samples were transferred to PVDF membranes (Millipore). The membranes

were incubated with the primary antibodies listed in Key Resources Table, followed by incubation

with HRP-conjugated secondary antibodies.

λ-PPase treatment and Phos-tag assay

Cells cultured in a 35 mm dish were rinsed twice with tris-buffered saline (TBS) (50 mM

Tris-HCl, 150 mM NaCl, pH 7.5) for collection, followed by lysis with RIPA buffer (1% NP-

40, 150 mM NaCl, 0.5 mM EDTA, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris-HCl,

2 mM PMSF, protease inhibitor cocktail, pH 7.4) on ice for 30 min. After centrifugation at

$12,000 \times g$ for 15 min at 4 °C, the lysate supernatants were mixed with 10× NEBuffer Pack

for Protein MetalloPhosphatases (PMP), 10× $MnCl_2$, and 400 U λ-PPase to react at 30 °C for

30 min, and quenched by the addition of SDS loading buffer. To analyze phosphorylated protein

samples, Phos-tag gel electrophoresis (10% separating gel) was applied based on the

manufacturer's protocol (Wako Pure Chemical Industries). The gel was supplemented with 50

μM $MnCl_2$ and 25 μM Phos-tag Acrylamide AAL-107 and polymerized overnight. Samples

were run at constant current of 30 mA per gel until the bromophenol blue arrived at the

bottom of the separating gel. Before transfer of proteins, the gels were washed in 1 mM

EDTA-containing transfer buffer for 30 min and then soaked in transfer buffer without

EDTA for 20 min before the proteins were transferred to PVDF membranes for western

blotting.

Immunoprecipitation

HEK293T or HeLa cells were collected after washing twice in cold PBS and lysed in lysis buffer

(50 mM HEPES, 250 mM NaCl, 5 mM EDTA, 0.1% NP-40, 1 mM DTT, 10% glycerol, 2 mM

PMSF, protease inhibitor cocktail, pH 7.5) on ice for 30 min. Proteins were recovered by

centrifugation at $12,000 \times g$ for 15 min at 4 °C, and the supernatant was incubated with primary

antibodies, protein G-Sepharose or protein A-Sepharose beads (GE Healthcare) for 2 h at 4 °C.

The beads were washed 3-5 times with cold lysis buffer, boiled at 100 °C in SDS loading buffer

for 5 min, and subjected to western blotting.

Clinical samples and immunohistochemistry

Tissue microarray of primary HCC samples was obtained from Shanghai Outdo Biotech Co., Ltd.

and US Biomax Inc. (Rockville, MD, USA). Immunohistochemical (IHC) staining was performed

as the following steps40. Formalin-fixed, paraffin-embedded tissue slides were dewaxed with

xylene and rehydrated by a graded series of alcohols, followed by antigen retrieval and block with

5% BSA for 60 min. Incubation was carried out at 4 °C for overnight with the primary antibody.

Primary antibodies included: anti-TRIM25 polyclonal antibody (1:200; Abclonal), anti-KEAP1

polyclonal antibody (1:200; Proteintech), and anti-Nrf2 polyclonal antibody (1:200; Proteintech).

Signals were detected using Envision-plus detection system (Dako, Carpinteria, CA, USA) and

visualized following incubation with 3,3′-diaminobenzidine.

Immunofluorescence and live cell imaging

Cells on coverslips were fixed using cold methanol for 10 min at -20°C and washed 3 times with

cold PBS. Permeabilized cells were blocked with 4% Bovine Serum Albumin (BSA) and

subsequently incubated with primary antibodies, followed by Alexa Fluor 488/568/647-conjugated

secondary antibodies. Next, the cells were stained with 1 μg/ml DAPI. The samples were observed

and photographed using a fluorescence microscope (TH4-200, Olympus) equipped with an Apo

Oil Objective lens (60×, NA 1.42, Olympus), and images were taken using DP controller software

(Olympus). Three-dimensional super resolution images were acquired using a three-dimensional

structured illumination microscope (3D-SIM) equipped with an Apo Oil Objective lens (100×, NA

1.49, Nikon) (N-SIM System, Nikon), and images were taken and reconstructed to maximum

projections using NIS-Elements AR software (Nikon). A confocal microscope (LSM-710 NLO,

Zeiss) equipped with a 100× /1.4 NA objective lens and a super-resolution confocal microscope

(Leica TCS SP8 STED × 3) equipped with a 100× /1.4 NA objective lens were used for cell

observation. Images were taken using LAS X software (Leica), deconvolved using Huygens

software (Scientific Volume Imaging, Netherlands), and processed using Photoshop (CC; Adobe).

Cells transfected with H_2B-RFP for 18 h were observed using a microscope (Nikon) with

spinning-disk UltraView VoX system (PerkinElmer). Cells were cultured in a chamber with 5%

CO_2 at 37 □, and time-lapse images were taken and post-processed using Volocity software

(PerkinElmer).

Construction of CEP44 knockout cell lines

CEP44 knockout HeLa cell lines were constructed using the CRISPR/Cas9 approach (52). Target

oligos were generated and ligated into U6-sgRNA vectors, and then transfected with pSpCas9 and

pcDNA3.1-Puro vectors. Clonal isolation of puromycin-resistant HeLa cells was performed using

flow cytometry analysis. To identify CEP44 knockout clones, cells were subjected to genome

sequencing and western blotting analysis. The target oligos for gRNA are designed as

5'-AAAAGCAGTTTATCCAATGT-3' and   5'-GGTAAGTCAGAACCTCCTTT-3'.

Statistics

The RNA-sequencing (RNA-seq) data was obtained from TCGA. Kaplan-Meier analysis was

performed for overall survival analyses and recurrence free survival. Overall survival was defined

as time from the first treatment after primary diagnosis to death of any cause. Recurrence free

survival was defined as time from the first treatment after primary diagnosis to death or recurrence

of any cause. The target regions were drawn around the fluorescent dots and measured using

ImageJ software (NIH). Either a two-tailed unpaired Student's t-test or one-way ANOVA with

Dunnett's multiple comparisons test was applied using GraphPad Prism 5 (version 5.01; GraphPad,

US) to compare differences between groups. The results are presented as the means ± S.E.M, and

$p < 0.05$ was considered to indicate a statistically significant difference, unless otherwise stated.

Data availability

All data needed to evaluate the conclusions in the paper are present in the paper.

References

1. Bettencourt-Dias M, and Glover DM. Centrosome biogenesis and function:

centrosomics brings new understanding. *Nat Rev Mol Cell Biol.* 2007;8(6):451-63.

2. Nigg EA, and Holland AJ. Once and only once: mechanisms of centriole duplication

and their deregulation in disease. *Nat Rev Mol Cell Biol.* 2018;19(5):297-312.

3. Fu J, Hagan IM, and Glover DM. The centrosome and its duplication cycle. *Cold*

Spring Harb Perspect Biol. 2015;7(2):a015800.

4. Nigg EA, and Stearns T. The centrosome cycle: Centriole biogenesis, duplication and

inherent asymmetries. *Nat Cell Biol.* 2011;13(10):1154-60.

5. Zhang Y, Tian J, Qu C, Peng Y, Lei J, Sun L, et al. A look into the link between

centrosome amplification and breast cancer. *Biomed Pharmacother.*

2020;132:110924.

6. Zyss D, and Gergely F. Centrosome function in cancer: guilty or innocent? *Trends*

Cell Biol. 2009;19(7):334-46.

7. Zhao JZ, Ye Q, Wang L, and Lee SC. Centrosome amplification in cancer and

cancer-associated human diseases. *Biochim Biophys Acta Rev Cancer.*

2021;1876(1):188566.

8. Pujana MA, Han JD, Starita LM, Stevens KN, Tewari M, Ahn JS, et al. Network

modeling links breast cancer susceptibility and centrosome dysfunction. *Nat Genet.*

2007;39(11):1338-49.

9. Ito KK, Watanabe K, Ishida H, Matsuhashi K, Chinen T, Hata S, et al. Cep57 and

Cep57L1 maintain centriole engagement in interphase to ensure centriole duplication

cycle. *J Cell Biol.* 2021;220(3).

10.   Nigg EA. Centrosome duplication: of rules and licenses. *Trends Cell Biol.*

2007;17(5):215-21.

11.   Seo MY, Jang W, and Rhee K. Integrity of the Pericentriolar Material Is Essential for

Maintaining Centriole Association during M Phase. *PLoS One.* 2015;10(9):e0138905.

12.   Lee K, and Rhee K. Separase-dependent cleavage of pericentrin B is necessary and

sufficient for centriole disengagement during mitosis. *Cell Cycle.* 2012;11(13):2476-

85.

13.   Kim J, Lee K, and Rhee K. PLK1 regulation of PCNT cleavage ensures fidelity of

centriole separation during mitotic exit. *Nat Commun.* 2015;6:10076.

14.   Aziz K, Sieben CJ, Jeganathan KB, Hamada M, Davies BA, Velasco ROF, et al.

Mosaic-variegated aneuploidy syndrome mutation or haploinsufficiency in Cep57

impairs tumor suppression. *J Clin Invest.* 2018;128(8):3517-34.

15.   Snape K, Hanks S, Ruark E, Barros-Nunez P, Elliott A, Murray A, et al. Mutations in

CEP57 cause mosaic variegated aneuploidy syndrome. *Nat Genet.* 2011;43(6):527-9.

16.   Wu Q, He R, Zhou H, Yu AC, Zhang B, Teng J, et al. Cep57, a NEDD1-binding

pericentriolar material component, is essential for spindle pole integrity. *Cell Res.*

2012;22(9):1390-401.

17.   Watanabe K, Takao D, Ito KK, Takahashi M, and Kitagawa D. The Cep57-pericentrin

module organizes PCM expansion and centriole engagement. *Nat Commun.*

2019;10(1):931.

18.   Jakobsen L, Vanselow K, Skogs M, Toyoda Y, Lundberg E, Poser I, et al. Novel

asymmetrically localizing components of human centrosomes identified by

complementary proteomics methods. *EMBO J.* 2011;30(8):1520-35.

19.   Hossain D, Shih SY, Xiao X, White J, and Tsang WY. Cep44 functions in centrosome

cohesion by stabilizing rootletin. *J Cell Sci.* 2020.

20.   Atorino ES, Hata S, Funaya C, Neuner A, and Schiebel E. CEP44 ensures the

formation of bona fide centriole wall, a requirement for the centriole-to-centrosome

conversion. *Nat Commun.* 2020;11(1):903.

21.   Kazazian K, Go C, Wu H, Brashavitskaya O, Xu R, Dennis JW, et al. Plk4 Promotes

Cancer Invasion and Metastasis through Arp2/3 Complex Regulation of the Actin

Cytoskeleton. *Cancer Res.* 2017;77(2):434-47.

22.   Ko MA, Rosario CO, Hudson JW, Kulkarni S, Pollett A, Dennis JW, et al. Plk4

haploinsufficiency causes mitotic infidelity and carcinogenesis. *Nat Genet.*

2005;37(8):883-8.

23.   Mariappan A, Soni K, Schorpp K, Zhao F, Minakar A, Zheng X, et al. Inhibition of

CPAP-tubulin interaction prevents proliferation of centrosome-amplified cancer cells.

EMBO J. 2019;38(2).

24.   Ganapathi Sankaran D, Stemm-Wolf AJ, and Pearson CG. CEP135 isoform

dysregulation promotes centrosome amplification in breast cancer cells. *Mol Biol Cell.*

2019;30(10):1230-44.

25.   Li X, Song N, Liu L, Liu X, Ding X, Song X, et al. USP9X regulates centrosome

duplication and promotes breast carcinogenesis. *Nat Commun.* 2017;8:14866.

26.   Fan G, Sun L, Shan P, Zhang X, Huan J, Zhang X, et al. Loss of KLF14 triggers

centrosome amplification and tumorigenesis. *Nat Commun.* 2015;6:8450.

27.   Wang HF, Takenaka K, Nakanishi A, and Miki Y. BRCA2 and nucleophosmin

606   coregulate centrosome amplification and form a complex with the Rho effector kinase

607   ROCK2. *Cancer Res.* 2011;71(1):68-77.

608   28.   Lingle WL, Lutz WH, Ingle JN, Maihle NJ, and Salisbury JL. Centrosome hypertrophy

609   in human breast tumors: implications for genomic stability and cell polarity. *Proc Natl*

610   *Acad Sci U S A.* 1998;95(6):2950-5.

611   29.   Lingle WL, Barrett SL, Negron VC, D'Assoro AB, Boeneman K, Liu W, et al.

612   Centrosome amplification drives chromosomal instability in breast tumor development.

613   *Proc Natl Acad Sci U S A.* 2002;99(4):1978-83.

614   30.   Bergmann S, Royer-Pokora B, Fietze E, Jurchott K, Hildebrandt B, Trost D, et al. YB-

615   1 provokes breast cancer through the induction of chromosomal instability that

616   emerges from mitotic failure and centrosome amplification. *Cancer Res.*

617   2005;65(10):4078-87.

618   31.   Kalimutho M, Sinha D, Jeffery J, Nones K, Srihari S, Fernando WC, et al. CEP55 is a

619   determinant of cell fate during perturbed mitosis in breast cancer. *EMBO Mol Med.*

620   2018;10(9).

621   32.   Sonnen KF, Gabryjonczyk AM, Anselm E, Stierhof YD, and Nigg EA. Human Cep192

622   and Cep152 cooperate in Plk4 recruitment and centriole duplication. *J Cell Sci.*

623   2013;126(Pt 14):3223-33.

624   33.   Saatci O, Akbulut O, Cetin M, Sikirzhytski V, Uner M, Lengerli D, et al. Targeting

625   TACC3 represents a novel vulnerability in highly aggressive breast cancers with

centrosome amplification. *Cell Death Differ.* 2023;30(5):1305-19.

34.   Pannu V, Mittal K, Cantuaria G, Reid MD, Li X, Donthamsetty S, et al. Rampant

centrosome amplification underlies more aggressive disease course of triple negative

breast cancers. *Oncotarget.* 2015;6(12):10487-97.

35.   Guan Z, Wang XR, Zhu XF, Huang XF, Xu J, Wang LH, et al. Aurora-A, a negative

prognostic marker, increases migration and decreases radiosensitivity in cancer cells.

Cancer Res. 2007;67(21):10436-44.

36.   Zitouni S, Nabais C, Jana SC, Guerrero A, and Bettencourt-Dias M. Polo-like kinases:

structural variations lead to multiple functions. *Nat Rev Mol Cell Biol.* 2014;15(7):433-

52.

37.   Mardin BR, Lange C, Baxter JE, Hardy T, Scholz SR, Fry AM, et al. Components of

the Hippo pathway cooperate with Nek2 kinase to regulate centrosome disjunction.

Nat Cell Biol. 2010;12(12):1166-76.

38.   Lemmens B, Hegarat N, Akopyan K, Sala-Gaston J, Bartek J, Hochegger H, et al.

DNA Replication Determines Timing of Mitosis by Restricting CDK1 and PLK1

Activation. *Mol Cell.* 2018;71(1):117-28 e3.

39.   Chou EJ, Hung LY, Tang CJ, Hsu WB, Wu HY, Liao PC, et al. Phosphorylation of

CPAP by Aurora-A Maintains Spindle Pole Integrity during Mitosis. *Cell Rep.*

2016;14(12):2975-87.

40.   Fu J, Bian M, Xin G, Deng Z, Luo J, Guo X, et al. TPX2 phosphorylation maintains

metaphase spindle length by regulating microtubule flux. *J Cell Biol.* 2015;210(3):373-

83.

41.　Gallini S, Carminati M, De Mattia F, Pirovano L, Martini E, Oldani A, et al. NuMA

Phosphorylation by Aurora-A Orchestrates Spindle Orientation. *Curr Biol.*

2016;26(4):458-69.

42.　Burgess SG, Peset I, Joseph N, Cavazza T, Vernos I, Pfuhl M, et al. Aurora-A-

Dependent Control of TACC3 Influences the Rate of Mitotic Spindle Assembly. *PLoS*

Genet. 2015;11(7):e1005345.

43.　Stevens NR, Roque H, and Raff JW. DSas-6 and Ana2 coassemble into tubules to

promote centriole duplication and engagement. *Dev Cell.* 2010;19(6):913-9.

44.　Schockel L, Mockel M, Mayer B, Boos D, and Stemmann O. Cleavage of cohesin

rings coordinates the separation of centrioles and chromatids. *Nat Cell Biol.*

2011;13(8):966-72.

45.　Mittal K, Kaur J, Jaczko M, Wei G, Toss MS, Rakha EA, et al. Centrosome

amplification: a quantifiable cancer cell trait with prognostic value in solid

malignancies. *Cancer Metastasis Rev.* 2021;40(1):319-39.

46.　Ganem NJ, Godinho SA, and Pellman D. A mechanism linking extra centrosomes to

chromosomal instability. *Nature.* 2009;460(7252):278-82.

47.　Harrison LE, Bleiler M, and Giardina C. A look into centrosome abnormalities in colon

cancer cells, how they arise and how they might be targeted therapeutically. *Biochem*

Pharmacol. 2018;147:1-8.

48.　Navarro-Serer B, Childers EP, Hermance NM, Mercadante D, and Manning AL.

Aurora A inhibition limits centrosome clustering and promotes mitotic catastrophe in

cells with supernumerary centrosomes. *Oncotarget.* 2019;10(17):1649-59.

49.   Serrano-Del Valle A, Reina-Ortiz C, Benedi A, Anel A, Naval J, and Marzo I. Future

671   prospects for mitosis-targeted antitumor therapies. *Biochem Pharmacol.*

672   2021;190:114655.

673   50.   He R, Huang N, Bao Y, Zhou H, Teng J, and Chen J. LRRC45 is a centrosome linker

674   component required for centrosome cohesion. *Cell Rep.* 2013;4(6):1100-7.

675   51.   Zhao H, Yang S, Chen Q, Duan X, Li G, Huang Q, et al. Cep57 and Cep57l1 function

676   redundantly to recruit the Cep63-Cep152 complex for centriole biogenesis. *J Cell Sci.*

677   2020;133(13).

678   52.   Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, and Zhang F. Genome engineering

679   using the CRISPR-Cas9 system. *Nat Protoc.* 2013;8(11):2281-308.

Acknowledgments

The authors thank C. Zhang (School of Life Sciences, Peking University) for providing the H2B-

RFP plasmid; C. Shan (the Core Facilities of Life Sciences, Peking University) and J. Zhou

(Nikon) for 3D-SIM and confocal microscopy imaging; L. Liu (Mass Spectrometry Facility of

National Center for Protein Sciences at Peking University) for assistance with protein

modification analysis; C. Yue (Department of Laboratory Medicine, Central People's Hospital of

Zhanjiang) for providing breast cancer cell lines; and Y. Qi (Department of Pathology, Central

People's Hospital of Zhanjiang) for observation of pathological tissue slices. The authors thank all

members of the laboratory for helpful discussion. This work was supported by the National

Natural Science Foundation of China (31830110, 32130024 and 32100542) and Zhanjiang

Science and Technology Project (2022A01104).

Author contributions

DHZ, JLT, NH and JGC designed the experiments. DHZ, WLW, FYL and MJY performed

experiments. DHZ, WLW, FYL and MJY performed data analysis. FYL and MJY collected breast

cancer tissue samples. DHZ, WLW and NH wrote and edited the manuscript. DHZ and JGC

provided funding for this study. The order of the co-first authors was assigned on the basis of their

relative contributions to the study.

Figures

Figure 1. Downregulation of CEP44 in breast cancer correlates with poor prognosis. (A)

Immunoblots of CPAP and CEP44 in MCF10A, MDA-MB-436 and MDA-MB-231 cells. Tubulin

was used as a loading control. (B) Immunostaining of Centrin-3 (red) and CEP44 (green) in

MCF10A, MDA-MB-436 and MDA-MB-231 cells. (C) Quantification of the fluorescence

intensity of CEP44 in the cells from (B). (D) CEP44 transcript levels in normal human mammary

tissues and breast carcinoma samples based on data sets from TCGA. (E) CEP44 transcript levels

in paired tumor and tumor-adjacent tissue samples based on data sets from TCGA. (F-G)

Representative images (F) of IHC staining and the relative IHC scores (G) of CEP44 in breast

cancer tissues and adjacent normal tissues. (H-I) Overall survival (OS) (H) and recurrence free

survival (RFS) (I) were compared between breast cancer patients with high and low CEP44

expression using Kaplan-Meier Plotter. For C, error bars represent the means ± S.E.M for three

independent experiments. n.s., not significant, ***$p < 0.001$, as determined using one-way

ANOVA with Dunnett's multiple comparisons test. For D, E and G, error bars represent the means

± S.E.M for three independent experiments. ***$p < 0.001$, as determined using Student's t-test.

Bars: 2 μm (B), 100 μm (F, upper), 50 μm (F, lower).

Figure 1-soure data 1 Information of collected breast cancer patients.

Figure 1–figure supplement 1. CEP44 is a centriole luminal protein localized at the proximal end.

(A) Schematic of human CEP44. FL, Full length; Coiled-coil domains, blue. (B-C) Immunoblots

of CEP44 in control and CEP44-depleted HeLa cells with a homemade (B) or commercial (C)

antibody against CEP44. Tubulin was used as a loading control. Asterisk: non-specific band. (D-G)

Immunostaining of CEP44 (green) with Acet-tubulin (red) (D and F) or GT335 (red) (E and G) in

HeLa cells. Centrioles were duplicated in (F) and (G). (H) Immunostaining of CEP44 (blue) with

CEP164, C-Nap1, CEP192, CEP152, CPAP or SAS6 (red) in CEP44-overexpressed HeLa cells.

Bars: 500 nm (D-G), 2 μm (H).

Figure 2. Depletion of CEP44 results in centriole overduplication. (A) Immunostaining of CEP97

(red) and α-tubulin (green) in MCF10A, MDA-MB-436, MDA-MB-231, SKBR3 and HCC1806

cells. DNA was stained with DAPI (blue). (B) Quantification of the percentage of cells with >4

centrioles from (A). (C-F) Immunostaining of Centrin-3 (green) (C) or CEP97 (green) (E) with γ-

tubulin (red) in HeLa cells transfected with control- or CEP44-siRNA. DNA was stained with

DAPI (blue). Quantification of the percentage of cells with the indicated number of Centrin-3 dots

(D) and CEP97 dots (F) from (C and E). (G) Immunostaining of CEP97 (red) in CEP44-GFP-

overexpressed MDA-MB-436 cells. (H) Quantification of the percentage of cells with more than 4

centrioles from (G). For B, D, F and H, error bars represent the means ± S.E.M for three

independent experiments. n.s., not significant, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, as determined

using one-way ANOVA with Dunnett's multiple comparisons test (D-F) or Student's t-test (H).

Bars: 10 μm (A), 5 μm (C, E), 2 μm (G).

Figure 3. Depletion of CEP44 results in premature centriole disengagement. (A) Immunostaining of Centrin-3 (red) and CEP192 (green) in wild-type (WT) and CEP44 knockout (KO) HeLa cells. DNA was stained with DAPI (blue). (B) Quantification of the percentage of cells with centriole disengagement from (A). (C) Immunostaining of CEP192 (green) and Centrin-3 (red) in WT and CEP44 KO HeLa cells. DNA was stained with DAPI (blue). (D) Immunostaining of CENP-F (green) and Centrin-3 (red) in WT and CEP44 KO HeLa cells treated with EdU (S phase marker; purple) for 30 min before fixation. DNA was stained with DAPI (blue). (E-F) Quantification of the percentage of cells with the indicated phenotype (E) and number of centrioles (F) from (D). (G-H) Time-lapse images of WT (G) and CEP44 KO (H) HeLa cells stably expressing H2B-RFP and GFP-Centrin-3. Red arrowheads indicate centrioles after the first duplication. Green arrowheads indicate the overduplicated centrioles after the second overduplication. For B, E and F, error bars represent the means ± S.E.M obtained for three independent experiments. **$p < 0.01$, as determined using Student's t-test. Bars: 5 μm (A, C upper, D, G and H), 2 μm (C, lower).

Figure 3–video 1 Video of WT HeLa cells stably expressing H2B-RFP and GFP-Centrin-3.

Figure 3–video 2 Video of CEP44 KO HeLa cells stably expressing H2B-RFP and GFP-Centrin-3.

Figure 3–figure supplement 1. Depletion of CEP44 results in centriole disengagement and centriole overduplication. (A) Schematic of the edited sequence of CEP44 in CEP44 knockout HeLa cells. (B) Immunoblots of the indicated proteins in wild-type (WT), CEP44-knockout (KO) and CEP44-rescued HeLa cells. Tubulin was used as a loading control. (C) Immunostaining of Centrin-3 (red) with CEP164 (green) (upper) or ODF2 (green) (lower) in wild-type (WT) and CEP44-knockout (KO) HeLa cells. (D) Quantification of the indicated ODF2 number with the percentage of 4 separate centrioles in CEP44 KO HeLa cells. (E) Immunostaining of Centrin-3

(red) and CEP192 (green) in control and CEP44 knockdown HeLa cells. DNA was stained with

DAPI (blue). (F) Quantification of the percentage of centriole disengagement in control and

CEP44 knockdown HeLa cells in interphase. (G) Immunostaining of Centrin-3 (red) and CEP192

(green) in WT and CEP44 KO HeLa cells. DNA was stained with DAPI (blue). (H) Quantification

of the percentage of mitotic cells with the indicated number of centrioles in WT and CEP44 KO

cells. (I) Immunostaining of γ-tubulin (red) in CEP44 KO HeLa cells transfected with GFP-

Centrin-3 (green). DNA was stained with DAPI (blue). The white and black arrowheads indicate

the centriole dots co-labelled with or without γ-tubulin, respectively. (J) Quantification of the

percentage of cells with more than 2 γ-tubulin-positive-centrioles in cells with disengaged ≥4

centrioles in the indicated cell cycle phase. For D, F and H, error bars represent the means ±

S.E.M obtained for three independent experiments. **$p < 0.01$, ***$p < 0.001$, as determined using

one-way ANOVA with Dunnett's multiple comparisons test. Bars: 5 μm (C, E, G and I).

Figure 4. CEP44 recruits CEP57/CEP57L1 to the centriole. (A) Lysates of HEK293T cells co-

overexpressing V5-CEP57L1 and Flag or CEP44-Flag were subjected to immunoprecipitation (IP)

and immunoblotting with anti-Flag and anti-V5 antibodies. (B) Lysates of HEK293T cells co-

overexpressing V5-CEP57 and Flag or CEP44-Flag were subjected to immunoprecipitation (IP)

and immunoblotting with anti-Flag and anti-V5 antibodies. (C) Immunoblots of CEP44, CEP57L1

and CEP57 in control and CEP44-depleted HeLa cells. Tubulin was used as a loading control. (D)

Immunostaining of CEP57L1 (green) or CEP57 (green) and γ-tubulin (red) in control and CEP44-

depleted HeLa cells. DNA was stained with DAPI (blue). (E-F) Quantification of the fluorescence

intensity of CEP57L1 (E) and CEP57 (F) in HeLa cells from (D). (G-H) Quantification of the

fluorescence intensity of CEP44 in CEP57-depleted (G) and CEP57L1-depleted (H) HeLa cells.

For E, F, G and H, error bars represent the means ± S.E.M for three independent experiments. n.s.,

not significant, ***$p < 0.001$, as determined using one-way ANOVA with Dunnett's multiple

comparisons test. Bars: 5 μm (D).

Figure 4–figure supplement 1. CEP57/CEP57L1 do not affect CEP44 localization at the

centrosome. (A) Immunostaining of CEP44 (green) and γ-tubulin (red) in control and CEP44

knockdown HeLa cells. DNA was stained with DAPI (blue). (B) Quantification of the normalized

fluorescence intensity of γ-tubulin at centrosomes in the cells from (A). (C) CEP57 transcript levels in normal human mammary tissues and breast carcinoma samples based on data sets from TCGA. (D) CEP57 transcript levels in paired tumor and tumor-adjacent tissue samples based on data sets from TCGA. For B, C and D, error bars represent the means ± S.E.M obtained for three independent experiments. ***$p < 0.001$, as determined using one-way ANOVA with Dunnett's multiple comparisons test. Bar: 5 μm (A).

Figure 5. Depletion of CEP44 results in multipolar spindle formation. (A) Immunostaining of α-tubulin (green) and γ-tubulin (red) in wild-type (WT) or CEP44 knockout (KO) HeLa cells. DNA was stained with DAPI (blue). (B-C) Quantification of the percentage of cells with a multipolar spindle (B) or a disorganized spindle (C) from (A). (D) Immunostaining of CEP97 (green) and α-tubulin (red) in WT or CEP44 KO HeLa cells. DNA was stained with DAPI (blue). (E)

Quantification of the percentage of cells with a multipolar spindle with centriole overduplication

from (D). (F) Time-lapse images of mitotic CEP44 KO HeLa cells stably expressing H2B-RFP

and GFP-Centrin-3. For B, C and E, error bars represent the means ± S.E.M obtained for three

independent experiments. n.s., not significant, **$p < 0.01$, ***$p < 0.001$, as determined using one-

way ANOVA with Dunnett's multiple comparisons test. Bars: 5 μm (A, D and F).

Figure 5–video 1 Video of multipolar spindle formation in mitotic CEP44 KO HeLa cells (pattern

1) stably expressing H2B-RFP and GFP-Centrin-3.

Figure 5–video 2 Video of multipolar spindle formation in mitotic CEP44 KO HeLa cells (pattern

2) stably expressing H2B-RFP and GFP-Centrin-3.

Figure 5–figure supplement 1. Knockdown of CEP44-centriole overduplication results in

multipolar spindle. (A) Immunoblots of the indicated proteins in control, CEP44-depleted and

CEP44-rescued HeLa cells. GAPDH was used as a loading control. (B) Immunostaining of α-

tubulin (green) and γ-tubulin (red) in control, CEP44-depleted and CEP44-rescued HeLa cells.

DNA was stained with DAPI (blue). (C) Quantification of the percentage of multipolar spindles in

(B). (D) Quantification of the percentage of disorganized mitotic spindles in (B). (E)

Immunostaining of α-tubulin (red) and CEP97 (red) in control and CEP44-depleted HeLa cells.

DNA was stained with DAPI (blue). (F) Quantification of the percentage of multipolar spindles

with centriole overduplication in (E). (G) Quantification of the percentage of pattern 1 and pattern

2 cells in WT and CEP44 KO cells. Pattern 1: premature centriole disengagement in interphase

without centriole overduplication; Pattern 2: premature centriole disengagement in interphase with

centriole overduplication. For C, D, F and G, error bars represent the means $\pm$ S.E.M obtained for

three independent experiments. n.s., not significant, $**p < 0.01$, $***p < 0.001$, as determined using

one-way ANOVA with Dunnett's multiple comparisons test. Bars: 5 μm (B, E).

Figure 6. Aurora A phosphorylates CEP44 at S324 in the G2/M phase. (A) Immunostaining of α-tubulin (red) and CEP44 (green) in HeLa cells. DNA was stained with DAPI (blue). (B) Immunoblots of CEP44 and Cyclin B1 at the indicated time points after nocodazole release in HeLa cells. GAPDH was used as a loading control. (C) Immunoblots of lysates from HEK293T cells co-overexpressing the indicated proteins with or without λ-ppase treatment. Tubulin was used as a loading control. The red asterisks indicate the shifted band of CEP44. (D) Immunoblots of lysates from HEK293T cells with or without nocodazole and MLN8237 treatment. Tubulin was used as a loading control. The red asterisk indicates the shifted band of CEP44. (E) Immunoblots of lysates from HEK293T cells co-overexpressing the indicated proteins. Tubulin was used as a

loading control. (F) The consensus sequence (R-X-pS/T-L/V) of the Aurora A substrate including

NuMA, CPAP, TPX2, TACC3 and CEP44. (G) Immunoblots of lysates from HeLa cells with or

without nocodazole treatment. (H) Immunoblots of the indicated proteins in control and Aurora A-

depleted HeLa cells after nocodazole treatment. (I) Immunostaining of Aurora A (red) and CEP44

(green) in control and Aurora A-depleted mitotic HeLa cells. DNA was stained with DAPI (blue).

(J) Quantification of the fluorescence intensity of CEP44 at spindle microtubules in mitotic cells

from (I). For J, error bars represent the mean ± S.E.M. for three independent experiments. ***$p <$

0.001, as determined using one-way ANOVA with Dunnett's multiple comparisons test. Bars: 5

μm (A and I).

Figure 6–figure supplement 1. CEP44 phosphorylation affects localization and does not affect

abundance. (A) Immunostaining of CEP44 (green) and α-tubulin (red) in control and CEP44

knockdown HeLa cells. DNA was stained with DAPI (blue). (B) Immunoblots of the lysates of

CEP44-Flag-overexpressed HeLa cells with overexpression of the indicated kinases. Tubulin was

used as a loading control. Asterisk: the shifted band. (C) Immunoblots of CEP44 in nocodazole-

treated HeLa cells with the addition of the indicated inhibitors. Tubulin was used as a loading

control. (D) Peptide competition assays using a phospho-CEP44 antibody against serine 324. (E)

Immunoblots of Aurora A and CEP44 in control and Aurora A knockdown HeLa cells with or

without nocodazole treatment. Tubulin was used as a loading control. (F) Immunostaining of

Aurora A (red) and CEP44 (green) in control and MLN8237-treated HeLa cells. DNA was stained

with DAPI (blue). (G) Quantification of the normalized intensity of CEP44 at spindle

microtubules from (F). (H) Quantification of the normalized intensity of CEP44 at spindle poles

from (F). For G and H, error bars represent the means ± S.E.M obtained for three independent

experiments. ***$p < 0.001$, as determined using Student's t-test. Bars: 5 μm (A and F).

Figure 7. Phosphorylation of CEP44 by Aurora A maintains spindle integrity. (A) Immunostaining

of γ-tubulin (red) and α-tubulin (green) in wild-type (WT) and CEP44 knockout (KO) HeLa cells.

DNA was stained with DAPI (blue). (B) Quantification of fluorescence microtubule intensity in

mitotic cells from (A). (C) Immunostaining of α-tubulin (red) and Flag (green) in mitotic HeLa

cells expressing the indicated proteins. DNA was stained with DAPI (blue). (D-E) Quantification

of the microtubule intensity at spindles (D) and intensity of CEP44 at spindle microtubules (E) in

mitotic cells from (C). (F) Immunostaining of Centrin-3 (red) and γ-tubulin (green) in control and

CEP44 knockdown HeLa cells. DNA was stained with DAPI (blue). (G) Quantification of the

normalized intensity of γ-tubulin at spindle MTs in control and CEP44 knockdown HeLa cells. (H)

Time-lapse images of mitotic CEP44 KO HeLa cells stably expressing H2B-RFP and GFP-

Centrin-3. Chr, chromosome. (I) Quantification of the percentage of cells with the indicated

phenotypes from (H). For B, D, E, G and I, error bars represent the means ± S.E.M for three

independent experiments. n.s., not significant, ***$p < 0.001$, as determined using one-way

ANOVA with Dunnett's multiple comparisons test. Bars: 5 µm (A, C, F and H).

Figure 7–video 1 Video of chromosome misalignment in mitotic CEP44 KO HeLa cells (pattern 1)

stably expressing H2B-RFP and GFP-Centrin-3.

Figure 7–video 2 Video of chromosome misalignment in mitotic CEP44 KO HeLa cells (pattern 2)

stably expressing H2B-RFP and GFP-Centrin-3.

Figure 7–figure supplement 1. CEP44 is critical for spindle structure formation. (A)

Immunostaining of γ-tubulin (red) and α-tubulin (green) in control, CEP44-depleted and CEP44-

rescued HeLa cells. DNA was stained with DAPI (blue). (B) Quantification of the normalized

fluorescence intensity of MT in the groups from (A). (C) Immunostaining of α-tubulin (green) and

Flag (red) in HeLa cells expressing the indicated proteins with or without nocodazole treatment.

DNA was stained with DAPI (blue). (D) Time-lapse images of cells stably expressing H2B-RFP

and GFP-Centrin-3 in mitotic CEP44 knockout HeLa cells. For B, error bars represent the means

± S.E.M obtained for three independent experiments. n.s., not significant, ***$p < 0.001$, as

determined using one-way ANOVA with Dunnett's multiple comparisons test. Bars: 5 μm (A, C

and D).

Figure 8. Schematic of the role of CEP44 in the cell cycle. In wild-type HeLa cells, CEP44 is localized at the proximal halves of the centriole lumen. For centriole duplication during the S phase, CEP44 exists in the lumen of the procentrioles to maintain disengagement between the old centrioles and the procentrioles. During mitosis, CEP44 translocates to the spindle due to Aurora A phosphorylation at serine 324 to ensure spindle integrity. After depletion of CEP44, premature centriole disengagement occurs following centriole duplication in the S phase, which licenses centriole overduplication as early as the S phase. The supernumerary centrioles result in multipolar spindle formation during mitosis. In addition, a lack of CEP44 localization at the spindle impairs spindle integrity, leading to an increase in the percentage of disorganized spindles. Notably, after Aurora A depletion or inhibition, unphosphorylated CEP44 fails to translocate to spindle microtubules and accumulates at the spindle pole, leading to impaired spindle integrity.

908   Unedited blot image