Designer antisense circRNA$_{GFP}$ reduces GFP abundance in Arabidopsis protoplasts in a

sequence-specific manner, independent of RNAi pathways

[1]Hossain M, [2]Pfafenrot C, [1]Nasfi S, [3]Sede A, [1]Imani J, [1]Šečić E, [1]Galli M, [1]Schäfer P,

[2]Bindereif A, [3]Heinlein M, [1]Ladera-Carmona M, [1,3&]Kogel KH

[1]Institute of Phytopathology, Centre for BioSystems, Land Use and Nutrition, Justus Liebig

University, Heinrich-Buff-Ring 26, D-35392, Giessen, Germany

[2]Institute of Biochemistry, Heinrich-Buff-Ring 17, Justus Liebig University, D-35392

Giessen, Germany

[3]Institut de biologie moléculaire des plantes, CNRS, Université de Strasbourg, 12 rue du

Général Zimmer, 67084 Strasbourg, France

Moammar.Hossain@agrar.uni-giessen.de (ORCID no.: 0009-0006-1528-4167)

Christina.Pfafenrot@gmx.de (ORCID no.: 0009-0009-1846-3613)

Sabrine.Nasfi-1@agrar.uni-giessen.de (ORCID no.: 0000-0001-5916-0824)

jafargholi.imani@agrar.uni-giessen.de (ORCID no.: 0000-0001-8265-1656)

Ena.Secic@agrar.uni-giessen.de (ORCID no.: 0000-0002-6469-4895)

ana.sede@ibmp-cnrs.unistra.fr (ORCID no.: 0000-0002-7868-0638)

Matteo.Galli@agrar.uni-giessen.de (ORCID no.: 0000-0003-4514-9434)

Patrick.Schaefer@agrar.uni-giessen.de (ORCID no.: 0000-0002-0366-6858)

manfred.heinlein@ibmp-cnrs.unistra.fr (ORCID no.: 0000-0001-7322-1654)

albrecht.bindereif@chemie.bio.uni-giessen.de (ORCID no.: 0000-0002-7730-7380)

Maria.Ladera-Carmona@agrar.uni-giessen.de (ORCID no.: 0000-0003-2167-3135)

karl-heinz.kogel@agrar.uni-giessen.de (ORCID no.: 0000-0003-1226-003X)

[&]corresponding author

Date of submission:

Number of tables and figures:

Word count:

Keywords: circular RNA, double-stranded RNA, protoplast transfection, disease resistance,

RNA interference, small RNA

Running title: circRNA reduces target protein abundance

Highlights

We demonstrate the potential of non-immunogenic circRNA as a tool for targeted gene

regulation in plants, where circRNA acts in an isoform- and sequence-specific manner, paving

the way for future agronomic applications.

Abstract:

Circular RNAs (circRNAs) are single-stranded RNA molecules characterised by their covalently closed structure and are emerging as key regulators of cellular processes in mammals, including gene expression, protein function and immune responses. Recent evidence suggests that circRNAs also play significant roles in plants, influencing development, nutrition, biotic stress resistance, and abiotic stress tolerance. However, the potential of circRNAs to modulate target protein abundance in plants remains largely unexplored. In this study, we investigated the potential of designer circRNAs to modulate target protein abundance in plants using Arabidopsis as a model system. We demonstrate that treatment with a 50 nt circRNA$_{GFP}$, containing a 30 nt GFP antisense sequence stretch, results in reduced GFP reporter target protein abundance in a dose- and sequence-dependent manner. Notably, a single-stranded open isoform of circRNA$_{GFP}$ had little effect on protein abundance, indicating the importance of the closed circular structure. Additionally, circRNA$_{GFP}$ also reduced GFP abundance in Arabidopsis mutants defective in RNA interference (RNAi), suggesting that circRNA activity is independent of the RNAi pathway. We also show that circRNA, unlike dsRNA, does not induce pattern-triggered immunity (PTI) in plants. Findings of this proof-of-principle study together are crucial first steps in understanding the potential of circRNAs as versatile tools for modulating gene expression and offer exciting prospects for their application in agronomy, particularly for enhancing crop traits through metabolic pathway manipulation.

Introduction

Agricultural production is affected by a variety of biotic and abiotic stress factors, which will increase with higher temperatures and extreme weather conditions in the course of climate change (Pareek et al. 2020; IPPC Secretariat 2021). Further improvement or even maintenance of global yield levels will depend to a large extent on new scientific solutions and their rapid introduction into agronomic practice (Van Dijk et al. 2021). While there is a broad consensus in the scientific community and clear legal requirements in most countries that synthetic pesticides, including herbicides, should be used as little as possible (Deguine et al. 2021), the effectiveness of alternative crop protection measures in intensive production systems still needs to be developed, and their dependence on environmental factors is often poorly understood (Perez-Alvarez et al. 2019; Kremer et al. 2023; Galli et al. 2024).

RNA is key for the storage, transmission, and modification of genetic information. In higher

organisms, RNA exists predominantly in the linear form as protein-coding mRNA and non-

coding forms, such as ribosomal (r)RNAs, long non-coding RNA (lnc)RNAs, transfer

(t)RNAs, and different types of small (s)RNA duplexes mostly of 21 to 24 base pairs. For the

latter, their high significance for regulatory processes such as maintenance of genome stability

and regulation of gene activity had only been found in 1998, when their function in RNA

interference (RNAi) was discovered (Fire et al. 1998; Baulcombe 2004). As a way of

communication between interacting organisms, RNA is also exchanged between animals or

plants and their pathogens or parasites, a phenomenon known as cross-kingdom RNA

interference (ckRNAi; LaMonte et al. 2012; Weiberg et al. 2013; Buck et al. 2014; Zhang et

al. 2016; Shahid et al. 2018; for review see Cai et al. 2018; Hamby et al. 2025).

Consistent with the role of RNA in natural communication between plant hosts and microbial

pathogens and pests, designer RNAs, such as engineered sRNA duplexes or longer double-

stranded (ds)RNAs of up to several hundred nucleotides, can protect plants from biotic and

abiotic stresses (for review see Koch and Kogel 2014; Cai et al. 2018; Niehl et al. 2018; Liu et

al. 2020; Koch and Wassenegger 2021; Liu et al. 2024; Chen and Kim 2024). However, their

instability and rapid degradation still hamper the agronomic use of these RNAs, especially if

they are not protected by chemical formulations (Mitter et al. 2017; Demirer et al. 2019; Jain

et al. 2022; Kogel 2025; Yong et al. 2025, Moorlach et al. 2025). Moreover, the risk of

genetic cross-resistance to various sRNAs or dsRNAs all acting via RNAi in the target

microbe or pest is a realistic scenario in which the RNAi pathway components and dsRNA

uptake mechanisms are susceptible to counter-selection (Khajuria et al. 2018; OECD 2020;

Wytinck et al. 2020; Mishra et al. 2021; Šečić and Kogel 2021; Choudhary et al. 2021; Luo et

al. 2024; Mishra et al. 2024).

RNAs are also exchanged between plant hosts and their weed parasites (Westwood and Kim

2017). There is growing interest in exploring this potential use of RNA for weed control (Mai

et al. 2021; Zabala-Pardo et al. 2022; Panozzo et al. 2025). However, RNA uptake and

stability in plants have challenged the development of RNA herbicides (Dalakouras et al.

2016; Bennett et al. 2020; Liu et al. 2021; Yong et al. 2025), indicating the need for RNA

with novel modes of action and molecular properties for their application in weed control.

In the present work, we have taken a first step to test the suitability of circular (circ)RNA for

future agronomic applications. Unlike linear (lin)RNA molecules, circRNAs form a

covalently closed loop, which confers resistance to exonucleases, making them more resistant

to degradation (Nielsen et al. 2022; Liu et al. 2022). This circularization can be achieved through a process known as back-splicing, in which a downstream splice donor site joins with an upstream splice acceptor site, resulting in the formation of a closed loop. circRNAs can arise from exons (exonic circRNA), introns (intronic circRNA), and intergenic regions (Zhang et al. 2013; Jeck and Sharpless 2014). Knowledge about circRNAs has been generated mainly in animal systems, where they are involved in the regulation of gene expression at multiple levels, including their activity as microRNA (miRNA) sponges (binding to miRNAs and repressing their function), as protein scaffolds, or in sequestration and translocation of proteins, facilitation of interactions between proteins, or translation of proteins (Hansen et al. 2013; Memczak et al. 2013; Guo et al., 2014; Yang et al. 2022). As a result, circRNAs modulate various physiological processes such as cell differentiation, development, and cellular immune responses, and play a role in numerous diseases, including cancer and neurological disorders, with their therapeutic potential widely recognized (He et al. 2021; Pisignano et al. 2023; Liu et al. 2022; Guo et al. 2025). circRNAs also have been detected in plants, where they accumulate in response to biotic and abiotic stress (Zhang and Dai 2022; He et al. 2025). A comparison of 6,519 circRNAs from rice (*Oryza sativa*) with those from 46 other species revealed a high degree of conservation within the Oryza genus (46%), and as much as 8.5% were also found in dicotyledonous plants, indicating some conservation of circRNAs in plants (Chu et al. 2022). An endogenous antisense circRNA was reported to regulate the expression of the small subunit of RuBisCO in *Arabidopsis thaliana* (Zhang et al. 2021). Interestingly, Arabidopsis circRNAs have also been detected in leaf intercellular washing fluids (IWF), showing that they can be secreted to the plant apoplast where they potentially get in contact with plant attacking microbes (Zand Karimi et al. 2022). Notably, apoplastic circRNAs are highly enriched in the posttranscriptional modification N6-methyladenine (m6A), which is known to efficiently initiate circRNA translation (Yang et al. 2017).

Here we explore the potential of exogenously applied designer circRNAs to target an endogenous Green Fluorescence Protein (GFP) reporter protein in Arabidopsis. *GFP*-expressing cells treated with the GFP-specific circRNA$_{GFP}$, in contrast to its corresponding linear single-strand form linRNA$_{GFP}$ or a circRNA that does not contain GFP-specific target sequences (circRNA$_{CTR1}$), showed reduced GFP protein abundance in a sequence- and circRNA-isoform-specific manner. Moreover, using RNAi mutants compromised in DICER-LIKE (DCL) and ARGONAUTE (AGO) activities, we demonstrate that the circRNA-

mediated activity on reporter protein abundance is independent of the canonical RNAi
pathways.

Results

Design of *GFP*-antisense circRNA

In a previous study, Pfafenrot and co-workers (2021) showed in the mammalian system that
antisense-circRNAs can be designed to efficiently interfere with translation of a protein-
coding gene. To develop a new tool for targeting gene expression with exogenous RNA, we
synthesized circRNA targeting the ORF of a *GFP* reporter gene (Fig. S1A). The exact
position of the target sequence was selected based on the secondary structure model of the
ORF (Fig. S1B). The selection of this region was confirmed by measuring mRNA
accessibility using the RNAup webtool (Fig. S1C). Based on this information, we designed a
50 nucleotide (nt) long antisense circRNA (circRNA$_{GFP}$) that contained a central anti-*GFP*
sequence of 30 nt with perfect complementarity. In addition, two different non-specific
circRNAs were synthesized, which contained a randomized 25 nt or 46 nt sequence with a
common 20 nt backbone, forming 45 nt circRNA$_{CTR1}$ and 66 nt circRNA$_{CTR2}$, respectively.
Secondary structure models of all circRNAs are shown in Fig. S1D (for sequences, see Table
S1).

circRNA$_{GFP}$ reduces the GFP abundance in *GFP*-expressing protoplasts in a sequence-
specific manner

To evaluate the antisense activity of the designed circRNAs, mesophyll protoplasts isolated
from Arabidopsis leaves were cotransfected with 4 μg of circRNA$_{GFP}$ or the non-specific
circRNA$_{CTR1}$ and 20 μg of plasmid pGY1-35S::GFP:RFP (Fig. S2A). After 18 h of incubation
(hpt) in the dark, the transfected protoplasts were analysed by measuring the ratio of GFP
fluorescence to RFP fluorescence using ImageJ. Notably, we found that the GFP fluorescence
was significantly reduced only in the circRNA$_{GFP}$-treated sample as compared to the
circRNA$_{CTR1}$ or the untreated controls (Fig. 1A, B). To further substantiate this finding, we
used an alternative GFP-expressing plasmid to transfect protoplasts. Similarly and consistent
with our expectation, in protoplasts transfected with pGY1-35S::GFP (Fig. S2B), GFP
fluorescence was also reduced upon treatment with circRNA$_{GFP}$, but not in samples treated

with circRNA$_{CTR1}$ or untreated controls, when normalized to red chlorophyll autofluorescence

(Fig. S3A, B). This finding indicated that circRNA$_{GFP}$ exerted an inhibitory effect on GFP

abundance in a sequence-specific manner.

The impact of various doses of circRNA$_{GFP}$ on GFP abundance

To further confirm target specificity of the designed circRNA, Arabidopsis protoplasts were

cotransfected with 20 µg pGY1-35S::GFP and increasing amounts of circRNA$_{GFP}$ and

circRNA$_{CTR1}$. ImageJ analyses of the GFP fluorescence after 18 hpt indicated that the effect

of circRNA$_{GFP}$ was slightly concentration dependent and remained circRNA-sequence-

specific over a concentration range up to 8 µg (Fig. 2A; Fig. S4A,B). Next, we quantified the

effect of circRNA$_{GFP}$ on GFP abundance in protoplasts by immunoblot analyses. For this

purpose, protoplasts were isolated from stable, transgenic *GFP*-expressing Arabidopsis plants

and subsequently treated with increasing concentrations of circRNA$_{GFP}$ and circRNA$_{CTR1}$. At

18 hpt, total protoplast proteins were extracted and separated by gel electrophoresis. GFP

protein abundance was visualised after blotting with an anti-GFP antibody and an anti-actin

antibody was used for protein normalisation. Consistent with the fluorescence analyses, we

found that the amount of GFP was reduced in protoplasts treated with increasing

concentrations of circRNA$_{GFP}$, as compared to protoplasts treated with circRNA$_{CTR1}$ (Fig.

2B).

The impact of circRNA$_{GFP}$ on GFP abundance is isoform-specific

Next, we comparatively examined the effect circRNA and its single-stranded, non-

circularised linear antisense form (linRNA$_{GFP}$) on GFP abundance, where linRNA$_{GFP}$

consisted of the same nt sequence as circRNA$_{GFP}$. For this purpose, Arabidopsis wild-type

protoplasts were treated with the plasmid pGY1-35S::GFP and 4 µg of either circular

(circRNA$_{GFP}$) or linear (linRNA$_{GFP}$) configurations of the GFP antisense RNA, and incubated

for 10 h, 18 h, and 32 h. circRNA$_{GFP}$-mediated inhibition of GFP abundance was already

detectable at 10 h after protoplasts treatment, and this effect persisted until 32 h (Fig. 3). In

contrast, linRNA$_{GFP}$ showed a transient inhibitory effect on protein abundance after 10 h,

which disappeared over time. Overall, our analyses revealed an isoform-specific effect of

circRNA$_{GFP}$ on GFP abundance. Our data suggest that circRNA is more effective than its

corresponding single-stranded linear RNA in antisense targeting of plant gene expression.

circRNA$_{GFP}$ affects GFP abundance independently of functional DCLs and AGOs

To obtain further information on the mode of action of sequence-specific designer circRNA on target protein abundance, we investigated whether the observed effect of circRNA$_{GFP}$ was lost in Arabidopsis mutants impaired in RNAi. Accordingly, protoplasts isolated from Arabidopsis DCL and AGO mutants were cotransfected with 20 μg plasmid pGY1-35S::GFP and 4 μg of the respective circRNA. Like wild-type protoplasts, *dcl1-11* and *ago1-27* protoplasts showed reduced GFP fluorescence in response to circRNA$_{GFP}$ (Fig. 4A,B; Fig. S5A,B), suggesting that disruption of DCL1 and AGO1 activities had no effect on circRNA$_{GFP}$-mediated reduction in GFP protein abundance. Consistent with this finding, immunoblot analyses further confirmed that disruption of the RNAi pathway did not affect the circRNA$_{GFP}$ effect. We found reduced GFP abundance in circRNA$_{GFP}$-treated *dcl1-11* (57%), *ago1-27* (49%) and wild-type (78%) protoplasts as compared to untreated protoplasts, whereas circRNA$_{CTR1}$ did not affect GFP abundance in either wild-type or mutant protoplasts (Fig. 4C). Immunoblot analyses of GFP abundance in additional RNAi mutants, including the DCL triple mutant *dcl2,3,4* and the two AGO mutants *ago2-1* and *ago4-2*, further substantiated that circRNA$_{GFP}$ retained its effect on GFP abundance when the RNAi pathway was compromised (Fig. S6).

circRNA$_{GFP}$ has no impact on *GFP* transcript abundance

Next, we analyzed the effect of circRNA$_{GFP}$ on the level of the *GFP* transcript in transgenic Arabidopsis protoplasts. Based on our previous work (Pfafenrot et al. 2021) we hypothesized that target mRNA levels would not be reduced upon circRNA treatment. To this end, *GFP*-expressing protoplasts were transfected with 20 μg pGY1-35S::GFP:RFP alone or together with either 4 μg circRNA$_{GFP}$ or circRNA$_{CTR1}$/circRNA$_{CTR2}$ followed by measurements of *GFP* transcript levels at 18 hpt. RT-qPCR analyses showed that none of the circRNAs reduced *GFP* transcript levels significantly in the wild-type (Fig. 5), and in all the mutants comprising *dcl1-11*, *ago1-27*, *dcl2,3,4*, *ago2-1* and *ago4-1*. (Fig. S7A-E). These findings showed that circRNA$_{GFP}$ inhibited protein abundance in a sequence- and isoform-specific manner without affecting *GFP* transcript levels, through a process that was independent of canonical RNAi pathways.

circRNA does not induce typical PTI responses in leaves

dsRNA activates pattern-triggered immunity (PTI) in plants leading to various responses, including callose deposition at plasmodesmata and MAP kinase activation (Niehl et al. 2016; Huang et al. 2023). We wondered whether similar to dsRNA also circRNA triggers a PTI response. To this end, equal molar amounts of circRNAs (circRNA$_{CTR1}$, circRNA$_{CTR2}$), their corresponding linear forms linRNA$_{CTR1}$ and linRNA$_{CTR2}$, or the synthetic dsRNA analog poly(I:C) (as positive control) were vacuum-infiltrated into Arabidopsis leaf disks together with aniline blue to stain callose. Fluorescence microscopy revealed that poly(I:C) treatment induced a strong aniline blue fluorescence at the plasmodesmata, as compared to the other treatments (Fig. 6A, B). Quantification of fluorescence showed that poly(I:C) triggered the highest mean callose intensity. Equal molar amounts (corresponding to 50 ng μL^{-1}) of the linear form of RNA$_{CTR1}$ (linRNA$_{CTR1}$) also triggered callose deposition, although to a lesser extent. Surprisingly, neither of the circRNA molecules induced an increased callose intensity, even when the circRNA concentration was increased by 5 times to 250 ng μL^{-1} (Fig. S8). In line with this, immunoblot analyses to detect mitogen-activated protein kinase (MAPK) phosphorylation in *Arabidopsis thaliana* leaves using anti-phospho-p44/42 ERK antibodies consistently revealed a strong induction of MAPK phosphorylation in response to 1 µM flg22, but not upon treatment with ~3 µM linRNA$_{CTR1}$ or circRNA$_{CTR1}$ (Fig. 6C). Finally, to assess whether circRNAs trigger a reactive oxygen species (ROS) response, *Nicotiana benthamiana* leaf discs were treated with 1 µM flg22, ~1 µM poly(I:C), or ~1 µM linRNA$_{CTR2}$ or circRNA$_{CTR2}$. In line with expectations, flg22 elicited a robust ROS burst, whereas poly(I:C), linRNA$_{CTR2}$, and circRNA$_{CTR2}$ failed to induce ROS accumulation (Fig. 6D). These results support the possibility that, unlike dsRNA, circRNAs may be able to evade receptor recognition in plants.

Discussion

The use of exogenous RNA to directly influence gene expression in crops is understudied despite its potential applicability as, for example, selective herbicides or as antimicrobial agents targeting plant susceptibility genes (Zabala☐Pardo et al., 2022, Mai et al., 2021; van Schie and Takken, 2014). Current RNA strategies based on dsRNA and sRNA, are challenged by low stability upon leaf application under field conditions (Mitter et al. 2017; Bachman et al. 2020; Kogel 2025; Yong et al. 2025). This limitation prompted us to conduct a baseline study to investigate whether more stable circular RNAs have properties that could make them a potentially effective additional alternative in crop protection in the future. Our findings

demonstrate that exogenously applied circRNA containing an antisense sequence to a *GFP*
reporter gene (circRNA$_{GFP}$) significantly reduces GFP protein abundance in Arabidopsis cells.
This effect is *i.* topology-dependent as the circular configuration is crucial, *ii.* sequence-
specific as non-targeting control circRNA was inactive, *iii.* RNAi-independent, as responses
to circRNA were similar in wild-type and RNAi-deficient mutants, and *iv.* accordingly
different from the mode of action of dsRNA. Overall, these results suggest that circRNA
application for manipulation of plant gene and metabolism is feasible, justifying further
fundamental research for future practical application.

circRNA is generally more stable than dsRNA due to its intrinsic physical properties: *i.*
circRNA has a covalently closed-loop structure, which prevents degradation by exonucleases
– unlike dsRNA, which has free 5' and 3' ends that are susceptible to degradation; *ii.* circRNA
is resistant to most RNA-degrading enzymes that typically target linear RNA molecules,
while dsRNA can be degraded by endonucleases such as RNase III; *iii.* circRNA has a high
secondary structure stability and can maintain its integrity under harsher conditions, while the
stability of dsRNA is affected by environmental factors such as temperature and pH (Liu and
Chen 2022; Nielsen et al. 2022; Ren et al. 2022; Liu et al. 2022; Moorlach et al. 2025). Future
studies are needed to determine whether the physical stability of circRNA makes it more
suitable for use in crop protection, considering both efficacy and environmental safety
implications.

dsRNA can trigger immune responses in animals, plants and fungi (de Reuver and Maelfait
2024; Niehl et al. 2016; Zheng et al. 2025). circRNA, on the other hand, is known to be less
immunogenic in mammals (Wesselhoeft et al. 2019). In line with this, we show that circRNA
does not induce callose deposition, ROS accumulation or MAP kinase activity, three
hallmarks of PTI responses in plants, whereas the dsRNA analogue poly(I:C) induced robust
callose deposition at plasmodesmata and MAP kinase (see Figs. 6 and S8). These results also
confirm earlier reports that dsRNA does not trigger ROS in plants (Niehl et al. 2016; Huang et
al. 2023). These observations, in line with reports of circRNAs not triggering antiviral
responses in mammalian cells (Breuer et al. 2022), highlight the specificity of circRNAs and
suggests they are less likely to provoke off-target immune reactions. This biological property
is also an interesting aspect when considering the potential of circRNA as a crop protection
agent.

We used the Arabidopsis protoplast system as the first proxy for evaluating the effect of

circRNAs on target proteins in plants. To make the protoplast experiments robust, we used

two different reporter constructs for transfection in independent experiments. Both constructs

pGY1-35S::GFP:RFP and pGY1-35S::GFP were expressed in the protoplasts and targeting

the GFP reporter by $circRNA_{GFP}$ resulted in a reduction of GFP fluorescence in both cases.

Consistent with this observation, immunoblot-based assays confirmed that the exogenous

application of $circRNA_{GFP}$, but not $circRNA_{CTR1}$, reduced the abundance of GFP protein.

The observation that $linRNA_{GFP}$, in contrast to $circRNA_{GFP}$, only transiently downregulated

GFP abundance (see Fig. 3) is consistent with the observation by our earlier work (Pfafenrot

et al. 2021), where the inhibitory potency of circular forms of antisense sequences

consistently surpassed their linear version. Therefore, the lasting inhibitory effect of

circRNAs is likely due to the high metabolic stability over linear forms, being more resistant

to the attack of exonucleases (Pfafenrot et al. 2021).

We also show here that the sequence-specific activity of circRNA occurs at the translation

level, as the mRNA-GFP transcript level is not reduced, while the GFP protein level decreases

significantly. That circRNA acts on the translational level rather than affecting target

transcript abundance is also consistent with Pfafenrot et al. (2021), who showed that various

designer circRNAs targeted viral RNA in infected mammalian cells in a sequence-specific

manner. The possibility that endogenous circRNA act on RNA transcripts in mammalian

systems has been recently discussed (Wang et al. 2024). Based on these recent reports along

with our findings we suggest a mode of circRNA action in plant cells where the circRNA

binds in a sequence-specific manner on target mRNAs and therefore inhibits or delays

translation. The slightly (albeit insignificantly) higher transcript abundance of the target *GFP*

mRNA in $circRNA_{GFP}$-treated samples could indicate that the transcript may be stabilized by

the binding.

Protoplasts have no cell walls and PEG facilitates the penetration of plasmids and other

nucleic acids into eukaryotic cells (Wu et al. 2009). By using protoplasts, we were able to

circumvent the problem of topical application of RNA on plant leaves, which is still a major

technical challenge due to the numerous barriers that need to be overcome, including the plant

cuticle and cell wall (Bennett et al. 2020; Kogel 2025). The issue of uptake of circRNA by

spray application in the field is one that requires further significant scientific input. While

there is no robust data on circRNA uptake through plant leaves, it is possible that future

applications may require formulations or physical means, as has been shown for dsRNAs and

small RNAs (Dalakouras et al. 2016; Mitter et al. 2017; Demirer et al. 2019; Yong et al.

2025).

In summary, this study demonstrates that exogenously applied designer circRNAs can

regulate protein expression in plants through a sequence-specific activity and an RNAi-

independent mode of action. These results pave the way for future studies aimed at using

circRNAs to develop new RNA-based herbicides and antimicrobials.

Experimental Procedures

Plant material and isolation of Arabidopsis protoplasts

Arabidopsis thaliana and *Nicotiana benthamiana* plants were grown from seeds in soil (LAT-

Terra Standard Topferde Struktur 1b, Hawita, Vechta, Germany) complemented with 2,5 g/L

fertilizer (Osmocote 12-7-19 +TE) at 22°C/18°C under 12 h/12 h or 16 h/8 h, light:dark

cycles, respectively. *A. thaliana* (Col-0) wild-type (WT), the mutants *ago1-27, ago2-1, ago4-*

1, dcl1-11 and the triple mutant *dcl2,3,4* were obtained from NASC (https://arabidopsis.info/).

All mutants were verified by genotyping. *A. thaliana* plants constitutively expressing *GFP*

were published in Harvey *et al*. (2020). Mesophyll protoplasts were produced using the tape-

Arabidopsis-sandwich method (Wu *et al*. 2007) starting from leaves of 30-day-old plants

grown at 22°C/18°C (day/night cycle) with 60% relative humidity and a photoperiod of 8/16 h

(240 µmol m^{-2} s^{-1} photon flux density) in a combination of type-T soil (F.-E. Typ Nullerde,

Hawita) and sand with a ratio of 3:1. Protoplasts were enzymatically released from leaves in a

solution containing 0.4 M mannitol, 20 mM KCl, 20 mM MES, pH 5.7, 1% (w/v) cellulase

R10, and 0.25% (w/v) macerozyme R10 (Duchefa Biochemie B.V.). Before use, the enzyme

solution was heated to 55°C for 10 min to solubilize the enzymes. Subsequently, 10 mM

CaCl$_2$ and 0.2% BSA were added before the solution was filter sterilized with a 45 µm filter

(Merck SA). Ten to 15 leaves with the upper epidermis peeled off were shaken (50-60 rpm) in

the activated enzyme solution for 1 h in the dark. Subsequently, protoplasts were collected by

filtration through a nylon mesh and centrifuged at 100 x g at 4°C. The final concentration of

mesophyll protoplasts used in each experiment was 500,000 ml^{-1}.

Design of circRNAs

The structure models of all circRNAs and putative target mRNAs were predicted using mfold

(version 3.6, mfold_util 4.7 and RNA Folding Form V2.3; www.unafold.org; Waugh et al.

2002; Zuker 2003) and RNAfold (RNAfold web server, university of Vienna;
http://rna.tbi.univie.ac.at; (Gruber et al. 2008; Lorenz et al. 2011; Mathews et al. 2004). The
30 nt antisense target sequence for the *GFP* gene in the 50 nt circRNA$_{GFP}$ was retrieved from
the *GFP-ORF* region (Fig. S1A). *GFP* mRNA accessibility was assessed with the software
RNAup (Vienna RNA Package, http://rna.tbi.univie.ac.at). In addition, two non-specific
circRNAs (with no *GFP* sequences) were produced, which contain a randomized 25 nt or 46
nt sequence with a common 20 nt backbone, forming a 45 nt circRNA$_{CTR1}$ and 66 nt
circRNA$_{CTR2}$, respectively.

373 Production of circRNAs and linRNAs
The syntheses of circRNAs were performed as described (Nielsen et al. 2022; Pfafenrot et al.
2021). Briefly, the antisense and the unspecific control sequences were inserted between two
spacers consisting of three unrelated nucleotides between the constant backbone, and this
arrangement assured the stem-loop formation in both the antisense and control sequences. The
oligonucleotide sequences for the circRNA synthesis, including the T7 promoter sequence,
were commercially synthesized (Sigma-Aldrich). circRNAs were produced by *in-vitro*
transcription from annealed DNA oligonucleotide templates (Table S2) using HighScribe T7
high-yield RNA synthesis kit (NEB) along with ATP, CTP, UTP, GTP (7.5 mM each), GMP
(30 mM, Merck), and RNaseOut (Thermo Fisher Scientific) at 37°C for 2 h. Before
circularization, the template DNA was digested with RQ1DNase (Promega) at 37°C for 30
min. Transcripts were purified with a Monarch RNA purification kit (NEB) and quantified
with a Qubit™ RNA broad-range assay kit (Thermo Fisher Scientific). The RNA transcript
was circularized overnight at 16°C with 200 U of T4 RNA ligase (Thermo Fisher Scientific)
in 200 µL T4 RNA ligase buffer containing 0.1 mg mL^{-1} BSA and RNaseOut (Thermo Fisher
Scientific). Following this reaction, circRNA was cleaned by phenol/chloroform extraction
(Roth) and ethanol precipitation.
Single-stranded linear RNA (linRNA$_{GFP}$) with the same nt sequence as circRNA$_{GFP}$ was
produced in the same way, but without circularization. Both linRNA and circRNA were
further gel-purified from denaturing polyacrylamide gels as described (Breuer and Rossbach
2020). To confirm the circularity of circRNA, 250 ng of circRNA or linRNA were treated
with or without 2 U of RNase R enzyme for 25 min at 37°C (Biozym) and analyzed by
denaturing polyacrylamide gel electrophoresis followed by ethidium bromide staining.

397 pGY1-35S::GFP:RFP plasmid construction

The red fluorescent protein (RFP) cassette was digested from the pBeaconRFP_GR vector

(https://gatewayvectors.vib.be/index.php/ID:3_20, Bargmann and Birnbaum 2009) using the

restriction enzyme *Nde*I (NEB, R0111S) for 20 min at 37°C followed by a deactivation step at

65° C for 10 min. pGY1-35S::GFP vector containing an *Nde*I restriction site was similarly

digested, including a dephosphorylation with the enzyme Fast alkaline phosphatase (Thermo

Fisher Scientific, EF0651). Both digestion products were run in a 1% agarose gel.

Corresponding bands were excised from the gel, purified using Wizard® SV Gel and PCR

Clean-Up System (Promega), and ligated overnight at room temperature using T4 DNA ligase

(Thermo Fischer Scientific, EL0011), resulting in the pGY1-35S::GFP:RFP vector.

Transfection of protoplasts

Twenty µg of plasmid pGY1-35S::GFP:RFP (Fig. S2A) or pGY1-35S::GFP (Fig. S2B;

Schweizer et al. 1999) along with circRNA or the respective linRNA were carefully added in

a volume of 20 µl to 10^5 mesophyll protoplasts in 200 µl MMg buffer (0.4 M mannitol, 15

mM $MgCl_2$, 4 mM MES, pH 5.7). Then 220 µl of PEG-Ca^{2+} (40% PEG-4000, 0.2 M

mannitol, 0.1 M $CaCl_2$) were slowly added to the protoplast suspension and incubated at room

temperature (Yoo *et al*., 2007). After 15 min of incubation, protoplasts were washed by

centrifugation two times with W5 buffer (154 mM NaCl, 125 mM $CaCl_2$, 5 mM KCl, 2 mM

MES, pH 5.7). Subsequently, protoplasts ($5x\ 10^5\ ml^{-1}$) were resuspended in W1 buffer (0.5 M

mannitol, 20 mM KCl, 4 mM MES, pH 5.7) and incubated in the dark at room temperature.

After 18 h, protoplasts were inspected under the fluorescence microscope (MZ16F Leica,

Germany). At least three pictures were taken from the same treatment at different spots.

Fluorescence was measured from these pictures by using ImageJ 1.54p software

(https://imagej.net/ij/) and the ratio of fluorescence levels between the GFP fluorescent

protoplasts (λ_{exc} 470, λ_{em} 525 nm) and total protoplasts (red auto fluorescence from the

chlorophyll, λ_{exc} 480, λ_{em} 510 nm), or alternatively, RFP (λ_{exc} 550, λ_{em} 650 nm) was

calculated.

Plant material and growth conditions for callose deposition assay

A. thaliana Col-0 plants were grown from seeds in soil (LAT-Terra Standard Topferde

Struktur 1b, Hawita, Vechta, Germany) complemented with 2,5 g/L fertilizer (Osmocote 12-

7-19 +TE) and kept in a growth chamber equipped with LED lights under 12h/12h light/dark

periods at 22°C/18°C. Leaf disks were excised using a cork borer and incubated overnight in

1 ml of water in the same chamber where the plants were grown. Leaf disks were then washed

two times with water, placed on microscope slides, and covered with coverslips fixed with

tape. The leaf disks were treated with 200 μl of a 0.1% aniline blue solution (pH=9)

containing either water, poly(I:C) (Sigma-Aldrich) as a positive control, or different

concentrations of circRNA or linRNA by adding the respective solution to the space between

the slide and the coverslip and by evacuation for 2 min (0.08 MPa). After incubation in the

dark for 30 min the callose staining at epidermal plasmodesmata was imaged with a Zeiss

LSM 780 confocal laser scanning microscope equipped with ZEN 2.3 software (Carl Zeiss,

Jean, Germany) by applying a 405 nm diode laser for excitation and filtering the emission at

475-525 nm. Eight-bit images acquired with a 40× 1.3 N.A. Plan Neofluar objectives with oil

immersion were analyzed with ImageJ 1.53 software (https://imagej.net/ij/) using the plug-in

calloseQuant (Huang et al. 2022). The fluorescence intensity levels of the callose spots were

measured in 3-4 images taken from each leaf disk. Three leaf disks from three different plants

were analyzed per condition. Normal distribution of the data was estimated and differences in

p-values between treatments and the control (water) were determined by parametric one-way

ANOVA followed by Dunnett's multiple comparisons test using the Prism 8.4.0 software.

Protein isolation, immunoblotting, and imaging

Protein extraction from protoplasts was performed using 4x SDS buffer (1 M Tris HCl, pH

6.8, 80% glycerol (v:v), bromophenol blue 10 mg, 4% SDS (w:v), 1 M of dithiothreitol (DTT)

in 20 ml Milli-Q water). The sample was vortexed, heated to 95°C for 5 min and then

centrifuged at 12,500 rpm for 2 min at 4°C. Protein concentration in the supernatant was

determined according to the Bradford Ultra method (Bradford, 1976) on a Bio-

Spectrophotometer (Eppendorf) at 595 nm. Then, 4 μg of each protein sample was loaded

onto a 12.5 % SDS-PAGE gel. Afterwards, the proteins in the gel were transferred into the

PVDF membrane (Merck KGaA) and blocked for one hour. Subsequently, the membrane was

cut into two separate pieces based on the sizes of GFP (~26.9 kDa) and Actin (~45 kDa). The

membrane containing the GFP band was incubated with the living colors monoclonal

antibody JL-8, (Takara Bio Inc) and anti-mouse IgG-peroxidase conjugate (Sigma) as the

secondary antibody. The membrane carrying the Actin band was incubated with an Actin

polyclonal antibody (AS132640, Agrisera, Sweden) and goat anti-rabbit IgG HRP (AS09602,

Agrisera, Sweden) as secondary antibody. After antibody incubation, the built-in software

from the ChemiDoc MP imaging system (Bio-Rad) was used to evaluate protein band

intensity in all western blots. Band intensity of control plants was set to one, and protein

accumulation was calculated as a ratio between GFP and Actin. Protoplast counting for all the

samples was done by ImageJ analysis.

RNA isolation and gene expression analysis

Total RNA was extracted with the Direct-zol™ RNA Microprep kit (Zymo Research) and

treated with DNase I following the manufacturer's instructions. One µg or 500 ng of RNA

was used for cDNA synthesis using a cDNA kit (RevertAid RT Kit, Thermo Fischer

Scientific). GFP transcript levels were quantified by qPCR using SYBR Green JumpStart Taq

ReadyMix (Sigma Aldrich, 1003444642) with a QuantStudio5 Real-Time PCR System

(Applied Biosystems). The total volume of 10 µl and three technical replicates are considered

for each reaction and 2 µl of ROX (CRX reference dye, Promega, C5411) was added to 1 ml

of SybrGreen as a passive reference dye that allows fluorescent normalization for qPCR data.

PCR conditions were 95°C for 5 min, followed by 40 cycles of 95°C for 15 s, 60°C for 30 s,

and 72°C for 30 s, and then by a melting curve analysis. GFP expression levels were first

normalized to RFP expression to account for transformation efficiency. Subsequently, fold

changes of GFP expression were calculated using the ΔΔCt method (Livak and Schmittgen

2001) relative to the geometric mean of two endogenous housekeeping genes (Vandesompele

et al. 2002), *Ubiquitin* (*UBQ5*, *AT3G62250*) and *Elongation Factor-1 alpha* (*EF1α*,

AT5G60390), with protoplasts transformed with pGY1-35S::GFP:RFP vector serving as the

control condition. One transformation of 100.000 protoplasts was considered as one biological

replicate. The results of four biological replicates are included in the data analysis. The primer

pairs employed for expression analysis are listed in Table S2.

Analysis of ROS production

Leaf disks were collected from 4-weeks-old *N. benthamiana* plants (3 biological replicates)

and incubated overnight in autoclaved Milli-Q water at 22°C in the dark. The following day,

the water was replaced, and the incubation continued for additional 4 h. Leaf disks were

transferred to a 96-well plate with 150 µL of a solution containing 17 µg/mL of luminol

(Sigma-Aldrich) and 10 µg/mL horseradish peroxidase (HRP; Sigma-Aldrich) together with

either 1 µM flg22 (ProteoGenix), 500 ng/µL of poly(I:C) (~1 µM) (Sigma-Aldrich) or 17

ng/µL (~1 µM) of linRNA$_{CTR2}$ or circRNA$_{CTR2}$. For the negative control, the elicitor was

replaced by water. Luminescence detection was achieved using the microplate reader
Varioskan LUX (Thermo Fisher Scientific) at 2 min intervals during 35 min. Mean values
obtained from nine leaf disks per treatment were expressed as mean relative light units (RLU).
499

MAPK phosphorylation of leaf extracts

The experiment was performed as described in Huang et al. (2023) with minor modifications.
Arabidopsis leaf disks were collected from three plants and incubated overnight in autoclaved
Milli-Q water at 22°C in the dark. After acclimatation, leaf disks were washed two times, then
gently transferred to 96-well plate and incubated for an additional hour in autoclaved Milli-Q
water. The water was replaced by 250 µL of a solution containing either 1 µM flg22
(EZBiolabs), or 50 ng/µL (~3 µM) of $linRNA_{CTR1}$ or $circRNA_{CTR1}$. For the negative control,
the elicitor was replaced by water. Leaf disks were vacuum infiltrated (0.08 MPa) for 30 min
and immediately placed on liquid nitrogen. For the immunoblots, the frozen tissue was
disrupted and resuspended in 100 µL 2X Laemmli buffer following a 5 min incubation at
95°C. Samples were separated in a 12% polyacrylamide gel and immunoblots were probed
with antibodies against phosphor-p44/42 ERK (Cell Signaling Technology) and HRP-labeled
secondary antibody (Thermo Fisher Scientific) for luminescence detection. Ponceau red was
used for loading control.

Statistical analysis

For Statistical analysis we used GraphPad Prism 8. The statistical significance between sets of
parametric data was analyzed with either one sample Student's *t*-test, or one-way ANOVA
followed by Dunnett's post-hoc test, whereas the one sample Wilcoxon test and Kruskal-
Wallis tests were used for non-parametric sets. A description of the specific tests is given in
figure legends. Callose quantification experiments were repeated twice with similar outcomes.
All the other experiments were repeated at least three times.

Acknowledgment

We thank Deutsche Forschungsgemeinschaft (DFG) and Agence National de la Research
(ANR) for strong support. In later stages, this work was also supported by a grant from the

Legends to Figures

Figure 1 Microscopic imaging of GFP and RFP fluorescence in Arabidopsis protoplasts.

Protoplasts were transfected with 20 µg of plasmid pGY1-35S::GFP:RFP and 4 µg of *GFP*

antisense circRNA$_{GFP}$, or 4 µg of non-targeting circRNA$_{CTR1}$. (A). After 18 hpt, protoplasts

were examined under the microscope using two distinct filters to GFP fluorescence (λexc 470,

λem 525 nm) and RFP fluorescence (λexc 550, λem 650 nm). Fluorescence intensity was

quantified based on images by using ImageJ 1.54p software. (B). The ratio between green

pixels (GFP fluorescence) and red pixels (RFP fluorescence) as calculated with ImageJ is

represented in the graph. The bar represents the measurements of ≥6 individual pictures taken

at various positions. Statistical analysis was performed using one-way ANOVA, followed by

Dunnett's multiple comparison test, where '**' denotes $p \leq 0.01$ significance to the protoplast

transfected without circRNA (control). Bars show standard deviation (SD).

Figure 2 Dose dependence of GFP fluorescence in Arabidopsis protoplasts after treatment

with increasing concentrations of circRNAs. Protoplasts were cotransfected with 20 µg of

plasmid pGY1-35S::GFP and the indicated amounts of circRNA$_{GFP}$ or circRNA$_{CTR1}$. (A)

After 18 hpt, protoplasts were inspected under the fluorescence microscope using two

different filters to calculate the ratio in fluorescence levels between the GFP fluorescent

protoplasts (λ_{exc} 470, λ_{em} 525 nm) and total protoplasts (red autofluorescence from the

chlorophyll, λ_{exc} 480, λ_{em} 510 nm). Fluorescence was measured based on pictures by using

ImageJ 1.54p software. Bars represent the average of the measurements of at least 3 pictures

taken at different spots with standard error of the mean (SEM). Statistical analysis was

performed using one-way ANOVA, where stars denote significance to the protoplast

transfected without circRNA (control). (Dunnett's test). '*' $p \leq 0.05$, '**' $p \leq 0.01$, '***' $p \leq$

0.001. (B) Immunoblot analysis of proteins extracted from protoplasts treated with increasing

concentrations of circRNA$_{GFP}$ or circRNA$_{CTR1}$. The values below the GFP band indicate the

remaining amount of GFP protein in percent as detected using an anti-GFP antibody. An anti-

Actin antibody was used to visualize Actin as an internal loading control. The ratio of GFP

protein accumulation is indicated below the bands expressed as ratio between GFP vs. Actin

signals).

Figure 3 Immunoblot analysis of GFP abundance in Arabidopsis protoplasts after treatment

with the *GFP* antisense circRNA$_{GFP}$ or its corresponding single-stranded, linear form

linRNA$_{GFP}$. Protoplasts were transfected with pGY1-35S::GFP and 4 µg circRN$_{GFP}$ or

linRNA$_{GFP}$ and analysed for GFP abundance after 10 h, 18 h and 32 h after transfection using

an anti-GFP antibody. An anti-actin antibody was used to visualize actin as a loading control.

GFP abundance is indicated below the band as percentage of GFP in protoplasts not treated

with RNA (control). The ratio of GFP protein abundance is indicated below the bands

expressed as ratio between GFP vs. actin signals).

Figure 4 GFP abundance in Arabidopsis protoplasts of RNAi mutants *dcl1-11* and *ago1-27*

after circRNA treatment. (A, B) Imaging of GFP fluorescence in *dcl1-11* (A) and *ago1-27* (B)

at 18 hpt. Protoplasts were transfected with 20 µg of plasmid pGY1-35S::GFP and 4 µg of

GFP antisense circRNA$_{GFP}$ or non-targeting circRNA$_{CTR1}$. Depicted is the ratio between green

pixels (GFP fluorescence) and red pixels (chlorophyll fluorescence) as calculated with

ImageJ. Bars represent the average of the measurements of at least 3 pictures taken at

different spots with standard error of the mean (SEM). Statistical analysis was performed with

one-way ANOVA, where * denotes $p \leq 0.05$ significance vs. control protoplasts (Dunnett's

test). (C). Immunoblot analysis of the GFP abundance in protoplasts from wild-type and

RNAi mutants *dcl1-11* and *ago1-27* upon treatment with circRNA$_{GFP}$ or circRNA$_{CTR1}$.

Protoplasts were harvested at 18 hpt, and equal amounts of protein were analyzed by

immunoblotting, using an anti-GFP antibody and an anti-Actin antibody as a loading control.

The ratio of GFP abundance is indicated below the bands expressed as ratio between GFP vs.

Actin signals.

Figure 5 RT-qPCR analysis of the amount of *GFP* transcripts in Arabidopsis wild-type

protoplasts upon treatment with circRNA. Protoplasts were transfected with 20 µg of pGY1-

35S::GFP:RFP plasmid and 4 µg of circRNA$_{GFP}$ or non-targeting circRNA$_{CTR2,}$ respectively.

Relative *GFP* expression was measured after 18 hpt. Values were normalized to RFP and the

housekeeping genes *Ubiquitin* and *EF1-α*. Bars represent an average of three independent

biological experiments pooled together with standard deviation (SD). No statistically

significant differences between treatments and genotypes were detected (Kruskal-Wallis test).

Figure 6 Analysis of immune innate responses after treatment with various unspecific RNAs.

(A) Mean values of callose intensity levels at PD in individual images taken of epidermal

cells of leaf disks treated with a 0.1% aniline blue solution containing water or 500 ng/µL of

poly(I:C), or 50 ng/µL of either linRNA or circRNA. Error bars show the standard error of the

mean. Parametric mean value variances were tested by one-way ANOVA followed by

Dunnett's multiple comparisons test. ****, $p \leq 0.0001$; *, $p \leq 0.05$; ns, non-significant. **(B)**

Callose accumulation at plasmodesmata (PD) in Arabidopsis leaves upon staining with aniline

blue. Samples were treated with 0.1% aniline blue solution together either with water

(negative control) or 500 ng/µL of poly(I:C) (positive control), or with 50 ng/µL of either

linRNA$_{CTR1}$, circRNA$_{CTR1}$, linRNA$_{CTR2}$, or circRNA$_{CTR2}$. For a better visualization, color was

assigned to the 8-bit images by applying the "fire" look-up table using the Image J software.

Scale bar, 10 µm. **(C)** Immunoblots for the detection of mitogen-activated protein kinase

(MAPK) phosphorylation in Arabidopsis leaves probed with antibodies against phosphor-

p44/42 ERK. Leaf disks were vacuum infiltrated with either water (negative control), 1 µM

flg22 or ~3 µM of each linRNA$_{CTR1}$ or circRNA$_{CTR1}$. As control for equal loading, Ponceau

Red staining was used. **(D)** ROS production assay in *N. benthamiana* leaf discs after

incubation with either water (negative control), 1 µM flg22, ~1 µM poly(I:C) or ~1 µM of

linRNA$_{CTR2}$ or circRNA$_{CTR2}$. Mean values and the standard error of the mean (SEM) are

shown from nine independent replicates per treatment. RLU, relative luminescence units.

Supplementary data and figure legends

Figure S1 Topology of designer circRNAs and their target sequences in the *GFP* gene. (A)

GFP-ORF region nucleotide sequence with color code: the 30 nt target sequence of the 50 nt

GFP antisense circRNA$_{GFP}$ is highlighted in red. Highlighted in blue is the CaMV35S

promotor sequence; highlighted in yellow is the 5'UTR; red font represents the circRNA

target sequence (30 nt); start codon ATG underlined in green; stop codon TAA underlined in

red; 3'UTR highlighted in green; terminator sequence highlighted in grey. (B) The exact

position of the target sequence was selected based on the secondary structure model of the

ORF. The red box inside the *GFP* mRNA structure indicates the binding site for the

designated antisense circRNA$_{GFP}$. (C) The selection of this region was confirmed by

measuring mRNA accessibility using the RNAup software (Vienna RNA Package,

http://rna.tbi.univie.ac.at/). The *GFP* mRNA sequence (5'UTR and 3'UTR included) was used

with its own antisense RNA sequence to determine RNA accessibility. This analysis served as

a guide for selecting sequence sections for antisense circRNA design. Green box indicates

higher mRNA accessibility, chosen for the target sequence of the antisense circRNA$_{GFP}$. (D)

Sequences and structures of three circRNAs used in this study. The secondary structures of

designer circRNAs were determined with the program mfold (www.unafold.org; Zuker 2003).

The *GFP* target sequence is colored in green, while randomized control sequences are colored

in blue. Respective nt sequences of all circRNA are shown in Table S1.

Figure S2 Map of plasmids for Arabidopsis protoplast transfection. (A) pGY1-

35S::GFP:RFP: The plasmid was prepared by using the pGY1-35S::GFP (Schweizer et al.

1999; see B) plasmid backbone, where RFP was inserted by cloning. (B) pGY1-35S::GFP

backbone: The whole construct confers resistance to ampicillin and carbenicillin antibiotics.

GFP is inserted and its expression is under the control of *Cauliflower Mosaic Virus* 35S

(CaMV35S) promoter and terminator. The Maps were generated by SnapGene freeware

(https://en.freedownloadmanager.org/Windows-PC/SnapGene-Viewer-FREE.html.

Figure S3 Microscopic imaging of GFP fluorescence in Arabidopsis protoplasts. Protoplasts

were transfected with 20 µg of plasmid pGY1-35S::GFP and 4 µg of *GFP* antisense

circRNA$_{GFP}$, or 4 µg of non-targeting circRNA$_{CTR1}$. (A). After 18 hpt, protoplasts were

examined under the microscope using two distinct filters to analyze the ratio of fluorescence

levels between the GFP fluorescent protoplasts (λexc 470, λem 525 nm) and red chlorophyll

autofluorescence (λ_{exc} 480, λ_{em} 510 nm). Fluorescence intensity was quantified based on

images by using ImageJ 1.54p software. (B). The ratio between green pixels (GFP

fluorescence) and red pixels (chlorophyll fluorescence) as calculated with ImageJ represented

in the graph. The bar represents the measurements of ≥ 6 individual pictures taken at various

positions. Statistical analysis was performed using one-way ANOVA, where '*' denotes $p \leq$

0.05 significance to the protoplast transfected without circRNA (control).

Figure S4 Microscopic imaging of the dose dependency of GFP abundance in Arabidopsis
protoplasts in response to circRNA treatment. Protoplasts were transfected with 20 µg of
plasmid pGY1-35S::GFP and the indicated amount of circRNA$_{GFP}$ (A) or circRNA$_{CTR1}$ (B),
respectively. At 18 hpt, protoplasts were inspected under the fluorescence microscope using
two different filters to calculate the ratio in fluorescence levels between the GFP fluorescent
protoplasts (λ_{exc} 470, λ_{em} 525 nm) and total protoplasts (red chlorophyll autofluorescence, λ_{exc}
480, λ_{em} 510 nm). Fluorescence was measured based on pictures by using ImageJ 1.54p
software.

Figure S5 Microscopic imaging of the GFP fluorescence in Arabidopsis RNAi mutants.
Protoplats of mutants *dcl1-11* (A) and *ago1-27* (B) were transfected with 20 µg of plasmid
pGY1-35S::GFP and 4 µg of *GFP* antisense circRNA$_{GFP}$ or non-targeting circRNA$_{CTR1}$. At 18
hpt, protoplasts were inspected under the fluorescence microscope using two different filters
to calculate the ratio in fluorescence levels between the GFP fluorescent protoplasts (λ_{exc} 470,
λ_{em} 525 nm) and total protoplasts (red chlorophyll autofluorescence, λ_{exc} 480, λ_{em} 510 nm).
Fluorescence was measured based on pictures using ImageJ 1.54p software.

Figure S6 Immunoblot analysis of the GFP abundance in Arabidopsis protoplasts of RNAi
mutants *dcl2,3,4*, *ago2-1* and *ago4-1* upon treatment with circRNA$_{GFP}$ or non-targeting
circRNA$_{CTR1}$. Protoplasts were transfected with 20 µg of pGY1-35S::GFP plasmid and 4 µg
circRNA$_{GFP}$ or circRNA$_{CTR1}$, respectively. Equal amounts of protein were analyzed by
Ponceau-S staining.

Figure S7 RT-qPCR analysis of the amount of *GFP* transcripts in Arabidopsis protoplasts
from RNAi mutants upon treatment with circRNA. Protoplasts were transfected with 20 µg of
pGY1-35S::GFP plasmid and 4 µg of circRNA$_{GFP}$ or non-targeting circRNA$_{CTR1,}$ respectively.
Relative *GFP* expression was measured after 18 hpt. Values were normalized to the
housekeeping gene *Ubiquitin*. Bars represent an average of three independent biological
experiments pooled together with standard error of the mean (SEM). No statistically
significant differences between treatments and genotypes were detected (one sample *t*-test for
(A), (B), (C), and (E) and one sample Wilcoxon test for (D), p≥ 0.05).

Figure S8 Mean values of callose intensity levels at PD in individual images taken of
epidermal cells of leaf disks treated with a 0.1% aniline blue solution containing water
(control), 500 ng/µL of poly(I:C), or either 50 ng/µL or 250 ng/µL of circRNA$_{CTR2}$. Error bars
show the standard error of the mean. Parametric mean value variances were tested by one-way
ANOVA followed by Dunnett's multiple comparisons test. ***, p$\leq$ 0.001; ns = non-
significant.

Table S1 Nucleotide sequence of circRNAs used in this study.
Table S2 List of oligonucleotides used in this study.
Oligonucleotides are used for circRNA and dsRNA production, as well as for RT-qPCR.

References

Bachman P, Fischer J, Song Z, Urbanczyk-Wochniak E, Watson G (2020) Environmental
Fate and Dissipation of Applied dsRNA in Soil, Aquatic Systems, and Plants. Front Plant
Sci 11:21. https://doi: 10.3389/fpls.2020.00021

Banerjee S, Banerjee, A, Gill SS, Gupta OP, Dahuja A, Jain PK, Sirohi A (2017) RNA
Interference: A Novel Source of Resistance to Combat Plant Parasitic Nematodes. Front
Plant Sci 8:834.

Bargmann BOR, Birnbaum KD (2009) Positive Fluorescent Selection Permits Precise, Rapid,
and In-Depth Overexpression Analysis in Plant Protoplasts. Plant Physiol 149(3):1231–
1239. https://doi.org/10.1104/pp.108.133975

Baulcombe DC (2004) RNA silencing in plants. Nature 431:356–363.

Bennett M, Deikman J, Hendrix B, Iandolino A (2020) Barriers to efficient foliar uptake of
dsRNA and molecular barriers to dsRNA activity in plant cells. Front Plant Sci 11.

Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram
quantities of protein utilizing the principle of protein-dye binding. Analytical
Biochemistry 72(1-2): 248–254.

Breuer J, Barth P, Noe Y, Shalamova L, Goesmann A, Weber F, Rossbach O (2022) What
goes around comes around: Artificial circular RNAs bypass cellular antiviral responses.
Molecular Therapy Nucleic Acids 28:623–635.

Breuer J, Rossbach O (2020) Production and purification of artificial circular RNA sponges for application in molecular biology and medicine. Methods Protocols 3(2): 42.

Buck A, Coakley G, Simbari F et al. (2014) Exosomes secreted by nematode parasites transfer small RNAs to mammalian cells and modulate innate immunity. Nat Comm 5:5488.

Cai Q, He B, Kogel KH, Jin H (2018) Cross-kingdom RNA trafficking and environmental RNAi-nature's blueprint for modern crop protection strategies. Curr Opin Microbiol 46:58–64.

Chen, LL, Kim VN (2024) Small and long non-coding RNAs: Past, present, and future. Cell 187(23):6451–6485.

Choudhary C, Meghwanshi KK, Shukla N, Shukla JN (2021) Innate and adaptive resistance to RNAi: a major challenge and hurdle to the development of double stranded RNA-based pesticides. 3 Biotech 11(12):498. https://doi: 10.1007/s13205-021-03049-3

Chu Q, Ding Y, Xu X, Ye CY, Zhu QH, Guo L, Fan L (2022) Recent origination of circular RNAs in plants. New Phytol 233:515–525.

Dalakouras A, Wassenegger M, McMillan JN, Cardoza V, Maegele I, Dadami E, Runne M, Krczal G, Wassenegger M (2016) Induction of Silencing in Plants by High-Pressure Spraying of In vitro-Synthesized Small RNAs. Front Plant Sci 7.

de Reuver R, Maelfait J (2024) Novel insights into double-stranded RNA-mediated immunopathology. Nat Rev Immunol 24:235–249. https://doi.org/10.1038/s41577-023-00940-3

Deguine JP, Aubertot JN, Flor RJ et al. (2021) Integrated pest management: good intentions, hard realities. A review. Agronomy for Sustainable Development 41:38.

Demirer GS, Zhang H, Matos JL, Goh NS, Cunningham FJ, Sung Y, Chang R, Aditham AJ, Chio L, Cho MJ, Staskawicz B, Landry MP (2019) High aspect ratio nanomaterials enable delivery of functional genetic material without DNA integration in mature plants. Nat Nanotechnol 14(5):456-464. https://doi: 10.1038/s41565-019-0382-5

Fire A, Xu S, Montgomery MK, Kostas S A, Driver SE, Mello CC (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. Nature 391(6669):806–811.

Galli M, Feldmann F, Vogler UK et al. (2024) Can biocontrol be the game-changer in integrated pest management? A review of definitions, methods and strategies. J Plant Dis Prot 131:265–291. https://doi.org/10.1007/s41348-024-00878-1

Gruber AR, Lorenz R, Bernhart SH, Neuböck R, Hofacker IL (2008) The Vienna RNA
Websuite. Nucleic Acids Research 36:suppl_2, pp W70–W74.
https://doi.org/10.1093/nar/gkn18

Guo JU, Agarwal V, Guo H, Bartel DP (2014) Expanded identification and characterization
of mammalian circular RNAs. Genome Biology 15(7):1–14.

Guo SK, Liu CX, Xu YF *et al.* (2025) Therapeutic application of circular RNA aptamers in a
mouse model of psoriasis. Nat Biotechnol 43:236–246. https://doi.org/10.1038/s41587-
024-02204-4

Hamby R, Cai Q, Jin H (2025) RNA communication between organisms inspires innovative
eco-friendly strategies for disease control. Nat Rev Mol Cell Biol 26:81–82.
https://doi.org/10.1038/s41580-024-00807-y

Hansen TB, Jensen TI, Clausen BH, Bramsen JB, Finsen B, Damgaard CK, Kjems J (2013)
Natural RNA circles function as efficient microRNA sponges. Nature 495(7441):384–
388.

Harvey S, Kumari P, Lapin D, Griebel T, Hickman R. *et al.* (2020) Downy Mildew effector
HaRxL21 interacts with the transcriptional repressor TOPLESS to promote pathogen
susceptibility. PLOS Pathogens 16(8): e1008835.

He S, Bing J, Zhong Y, Zheng X, Zhou Z, Wang Y, Hu J, Sun X (2025) PlantCircRNA: a
comprehensive database for plant circular RNAs. Nucleic Acids Res 53(D1):D1595-
D1605. https://doi: 10.1093/nar/gkae709

He AT, Liu J, Li F, Yang BB (2021) Targeting circular RNAs as a therapeutic approach:
current strategies and challenges. Signal Transduction Targeted Therapy 6(1):185.

Hsu MT, Coca-Prados M (1979) Electron microscopic evidence for the circular form of RNA
in the cytoplasm of eukaryotic cells. Nature 280(5720):339–340.

Huang C, Mutterer J, Heinlein M (2022) In vivo aniline blue staining and semi-automated
quantification of callose deposition at plasmodesmata. Meth Mol Biol 2457:151–165.

Huang C, Sede AR, Elvira-González L, Yan Y, Rodriguez ME, Mutterer J, Emmanuel E,
Shan L, Heinlein M (2023) dsRNA-induced immunity targets plasmodesmata and is
suppressed by viral movement proteins. Plant Cell 35(10):3845–3869.

IPPC Secretariat (2021) Scientific review of the impact of climate change on plant pests – A
global challenge to prevent and mitigate plant pest risks in agriculture, forestry and

ecosystems. Rome. FAO on behalf of the IPPC Secretariat. https://doi.org/10.4060/cb4769en

Jain RG, Fletcher SJ, Manzie N *et al.* (2022) Foliar application of clay-delivered RNA interference for whitefly control. Nat Plants 8:535–548. https://doi.org/10.1038/s41477-022-01152-8

Jeck WR, Sharpless NE (2014) Detecting and characterizing circular RNAs. Nat Biotech 32(5):453–461.

Khajuria C, Ivashuta S, Wiggins E, Flagel L, Moar W, Pleau M, Miller K, Zhang Y, Ramaseshadri P, Jiang C, Hodge T (2018) Development and characterization of the first dsRNA-resistant insect population from western corn rootworm, *Diabrotica virgifera virgifera* LeConte. PloS one 13(5):e0197059.

Koch A, Kogel KH (2014) New wind in the sails: improving the agronomic value of crop plants through RNAi-mediated gene silencing. Plant Biotechnol J 12(7):821-31. doi: https://10.1111/pbi.12226

Koch A, Wassenegger M (2021) Host-induced gene silencing – mechanisms and applications. New Phytol 231:54–59.

Kogel KH (2025) Lysozyme-coated LDHs boost trait control. Nat Plants 11(1):9-10. doi: https://10.1038/s41477-024-01874-x

Kremer RJ (2023) Chapter 7 - Bioherbicide development and commercialization: challenges and benefits, Editor(s): Opender Koul, Development Commercialization Biopesticides, Academic Press, Pages 119-148, ISBN 9780323952903.

LaMonte G, Philip N, Reardon J, Lacsina JR, Majoros W, Chapman L, Thornburg CD, Telen MJ, Ohler U, Nicchitta CV, Haystead T, Chi JT (2012) Translocation of sickle cell erythrocyte microRNAs into *Plasmodium falciparum* inhibits parasite translation and contributes to malaria resistance. Cell Host & Microbe 12(2):187–99.

Liu C, Kogel KH, Ladera-Carmona M (2024) Harnessing RNA interference for the control of Fusarium species: A critical review. Mol Plant Pathol. Oct;25(10):e70011. https://doi: 10.1111/mpp.70011.

Liu R, Ma Y, Guo T, Li G (2022) Identification, biogenesis, function, and mechanism of action of circular RNAs in plants. Plant Commun 4(1):100430. https://doi: 10.1016/j.xplc.2022.100430

Liu S, Jaouannet M, Dempsey DA, Imani J, Coustau C, Kogel KH (2020) RNA-based
technologies for insect control in plant production. Biotechnology Advances 39:107463.

Liu S, Ladera-Carmona MJ, Poranen MM *et al*. (2021) Evaluation of dsRNA delivery
methods for targeting macrophage migration inhibitory factor MIF in RNAi-based aphid
control. J Plant Dis Prot 128:1201–1212.

Livak KJ, Schmittgen TD (2001) Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. Methods 25:402–408.

Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C, Stadler PF, Hofacker IL
(2011) "ViennaRNA Package 2.0", Algorithms for Molecular Biology, 6(1):26.

Luo X, Satyabrata N, Zhang Y, Zhou X, Yang C, Pan H (2024) Risk assessment of RNAi-
based biopesticides. New Crops 1,100019, ISSN 2949-9526.
https://doi.org/10.1016/j.ncrops.2024.100019

Mai J, Liao L, Ling R, Guo X, Lin J, Mo B, Chen W, Yu Y (2021) Study on RNAi-based
herbicide for *Mikania micrantha*. Synthetic and Systems Biotechnology 6(4): 437–445.

Mathews DH, Disney MD, Childs JL, Schroeder SJ, Zuker M, Turner DH (2004)
Incorporating chemical modification constraints into a dynamic programming algorithm
for prediction of RNA secondary structure. Proc Natl Acad Sci U S A 101(19):7287–
7292.

Memczak S, Jens M, Elefsinioti A, Torti F, Krueger J, Rybak A et al. (2013) Circular RNAs
are a large class of animal RNAs with regulatory potency. Nature 495(7441): 333–338.

Mishra S, Moar W, Jurat-Fuentes JL (2024) Larvae of Colorado potato beetle (Leptinotarsa
decemlineata Say) resistant to double-stranded RNA (dsRNA) remain susceptible to
small-molecule pesticides. Pest Manag Sci 80(2):905–909. https://doi: 10.1002/ps.7825

Mishra S, Dee J, Moar W *et al*. (2021) Selection for high levels of resistance to double-
stranded RNA (dsRNA) in Colorado potato beetle (*Leptinotarsa decemlineata* Say) using
non-transgenic foliar delivery. Scientific Reports 11: 6523.

Mitter N, Worrall EA, Robinson KE, Li P, Jain RG, Taochy C, Fletcher SJ, Carroll BJ, Lu
GQ, Xu ZP (2017) Clay nanosheets for topical delivery of RNAi for sustained protection
against plant viruses. Nature Plants 3:16207.

Moorlach BW, Sede AR, Hermann KM, Levanova AA, Poranen MM, Westphal M,
Wortmann M, Stepula E, Jakobs-Schönwandt D, Heinlein M, Keil W, Patel AV (2025)
Interpolyelectrolyte complexes of in vivo produced dsRNA with chitosan and alginate for

enhanced plant protection against tobacco mosaic virus, International Journal of
Biological Macromolecules 306(2):141579.
https://doi.org/10.1016/j.ijbiomac.2025.141579

Niehl A, Soininen M, Poranen MM, Heinlein M (2018) Synthetic biology approach for plant
protection using dsRNA. Plant Biotechnol J 16(9):1679–87. https://doi:
10.1111/pbi.12904

Niehl A, Wyrsch I, Boller T, Heinlein M (2016) Double-stranded RNAs induce a pattern-
triggered immune signaling pathway in plants. New Phytol 211:1008–1019.

Nielsen AF, Bindereif A, Bozzoni I, Hanan M, Hansen TB, Irimia M *et al.* (2022) Best
practice standards for circular RNA research. Nat Methods 19(10):1208–1220.

OECD (2020) Considerations for the Environmental Risk Assessment of the Application of
Sprayed or Externally Applied ds-RNA-Based Pesticides, Series on Pesticides and
Biocides, OECD Publishing, Paris, https://doi.org/10.1787/576d9ebb-en.

Panozzo S, Milani A, Bordignon S, Scarabel L, Varotto S (2025) RNAi technology
development for weed control: all smoke and no fire? Pest Manag Sci
https://doi.org/10.1002/ps.8729

Pareek A, Dhankher OP, Foyer CH (2020) Mitigating the impact of climate change on plant
productivity and ecosystem sustainability. J Exp Bot 71(2):451–456.

Perez-Alvarez R, Nault BA & Poveda K (2019) Effectiveness of augmentative biological
control depends on landscape context. Scientific Reports 9:8664.

Pfafenrot C, Schneider T, Müller C, Hung LH, Schreiner S, Ziebuhr J, Bindereif A (2021)
Inhibition of SARS-CoV-2 coronavirus proliferation by designer antisense-circRNAs.
Nucleic Acids Research 49(21):12502–12516.

Pisignano G, Michael DC, Visal TH *et al.* (2023) Going circular: history, present, and future
of circRNAs in cancer. Oncogene 42:2783–2800. https://doi.org/10.1038/s41388-023-
02780-w

Ren L, Jiang Q, Mo L *et al.* (2022) Mechanisms of circular RNA degradation. Commun Biol
5:1355. https://doi.org/10.1038/s42003-022-04262-3

Schweizer P, Pokorny J, Abderhalden O, Dudler R (1999) A transient assay system for the
functional assessment of defense-related genes in wheat. Molecular Plant Microbe
Interactions 12:647–654.

Šečić E, Kogel KH (2021) Requirements for fungal uptake of dsRNA and gene silencing in
RNAi-based crop protection strategies. Curr Opin Biotech 70:136–142, ISSN 0958-1669.

Shahid S, Kim G, Johnson NR, Wafula E, Wang F, Coruh C, Bernal-Galeano V, Phifer T,
dePamphilis CW, Westwood JH, Axtell MJ (2018) MicroRNAs from the parasitic plant
Cuscuta campestris target host messenger RNAs. Nature 553:82–85.

Van Dijk M, Morley T, Rau ML, Saghai Y (2021) A meta-analysis of projected global food
demand and population at risk of hunger for the period 2010–2050. Nat Food 2(7):494–
501.

van Schie CC, Takken FL (2014) Susceptibility genes 101: how to be a good host. Annu Rev
Phytopathol 52:551–581.

Vandesompele J, De Preter K, Pattyn F *et al.* (2002) Accurate normalization of real-time
quantitative RT-PCR data by geometric averaging of multiple internal control genes.
Genome Biol 3, research0034.1. https://doi.org/10.1186/gb-2002-3-7-research0034

Wang S, Li X, Liu G *et al.* (2024) Advances in the understanding of circRNAs that influence
viral replication in host cells. Med Microbiol Immunol 213:1.
https://doi.org/10.1007/s00430-023-00784-7

Waugh A, Gendron P, Altman R, Brown JW, Case D, Gautheret D, Harvey SC, Leontis N,
Westbrook J, Westhof E, Zuker M, Major F (2002) RNAML: a standard syntax for
exchanging RNA information. RNA 8(6):707–717.

Weiberg A, Wang M, Lin F M, Zhao H, Zhang Z, Kaloshian I, Jin H (2013) Fungal small
RNAs suppress plant immunity by hijacking host RNA interference pathways. Science
342(6154):118–123.

Wesselhoeft RA, Kowalski PS, Parker-Hale FC, Huang Y, Bisaria N, Anderson DG (2019)
RNA circularization diminishes immunogenicity and can extend translation duration in
vivo. Mol Cell 74(3):508–520.

Westwood JH, Kim G (2017) RNA mobility in parasitic plant–host interactions. RNA
Biology 14(4):450–455.

Wu FH, Shen SC, Lee LY, Lee SH, Chan MT and Lin CS (2009) Tape-Arabidopsis
Sandwich-a simpler Arabidopsis protoplast isolation method. Plant Methods 5:1–10.

Wytinck N, Manchur CL, Li VH, Whyard S, Belmonte MF (2020) dsRNA uptake in plant
pests and pathogens: Insights into RNAi-Based Insect and Fungal Control Technology.
Plants (Basel) 9(12): 1780.

Yang L, Wilusz JE, Chen LL (2022) Biogenesis and Regulatory Roles of Circular RNAs.
Annu Rev Cell Biol 38:263-289. https://doi.org/10.1146/annurev-cellbio-120420-125117

Yang Y, Fan X, Mao M, Song X, Wu P, Zhang Y, Jin Y, Yang Y, Chen LL, Wang Y, Wong
CC (2017) Extensive translation of circular RNAs driven by N 6-methyladenosine. Cell
Res 27:626–641.

Yong J, Xu W, Wu M, Zhang R, Mann CWG, Liu G, Brosnan CA, Mitter N, Carroll BJ, Xu
ZP (2025) Lysozyme-coated nanoparticles for active uptake and delivery of synthetic
RNA and plasmid-encoded genes in plants. Nat Plants 1–14.

Yoo SD, Cho YH, Sheen J (2007) Arabidopsis mesophyll protoplasts: A versatile cell system
for transient gene expression analysis. Nat Protocols 2(7):1565–1572.

Zabala-Pardo D, Gaines T, Lamego FP, Avila LA (2022) RNAi as a tool for weed
management: challenges and opportunities. Advances Weed Science 40(Spec1):
e020220096.

Zand Karimi H, Baldrich P, Rutter BD, Borniego L, Zajt KK, Meyers BC, Innes RW (2022)
Arabidopsis apoplastic fluid contains sRNA-and circular RNA–protein complexes that
are located outside extracellular vesicles. Plant Cell 34(5):1863–1881.

Zhang H, Liu S, Li X, Yao L, Wu H, Baluška F, Wan Y (2021) An antisense circular RNA
regulates expression of RuBisCO small subunit genes in Arabidopsis. Front Plant Sci
12:665014.

Zhang P, Dai M (2022) CircRNA: a rising star in plant biology. Journal of Genetics and
Genomics 49(12):1081–1092.

Zhang Y, Zhang XO, Chen T, Xiang JF, Yin QF, Xing YH et al. (2013) Circular intronic long
noncoding RNAs. Mol Cell 51(6):792–806.

Zheng Y, Moorlach B, Jakobs-Schönwandt D, Patel A, Pastacaldi C, Jacob S, Ladera
Carmona M (2025) Exogenous dsRNA triggers sequence-specific RNAi and fungal
stress responses to control *Magnaporthe oryzae* in *Brachypodium distachyon*. Commu
Biol 8(1):121.

Zuker M (2003) Mfold web server for nucleic acid folding and hybridization prediction.
Nucleic Acids Research 31(13):3406–3415.

Statements & Declarations

Funding

This work was funded by the Deutsche Forschungsgemeinschaft (DFG) in the research unit
5116 (exRNA) to KHK and AB/PS. It was also funded by ERA-NET SusCrop 2 program
(DFG grant 459501999 to KHK and Agence National de la Research (ANR) grant ANR-21-
SUSC-0003-01 to MH) as part of the project BioProtect coordinated by MH and carried out
under the second call of the ERA-NET Cofund SusCrop, being part of the Joint Programming
Initiative on Agriculture, Food Security and Climate Change (FACCE-JPI). SusCrop has
received funding from the European Union's Horizon 2020 research and innovation
programme under grant agreement No 771134. In later stages, this work was also supported
by a grant from the Cercle Gutenberg (Alsace, France) to KHK and MH. SN was partly
supported by the Ernst-Leopold Klipstein Foundation, Paderborn Gießen, Germany.

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

All authors contributed to the study conception and design. Material preparation, data
collection and analysis were performed by M. Hossain, C. Pfafenrot, S. Nasfi, A. Sede, J.
Imani, E. Šečić, A. Bindereif A, M. Heinlein, M. Ladera-Carmona and KH Kogel. The first
draft of the manuscript was written by KH Kogel and all authors commented on previous
versions of the manuscript. All authors read and approved the final manuscript.

Data Availability

The datasets generated during and/or analysed during the current study are available from the
corresponding author on reasonable request.

Fig. 2

A

B

| circRNA$_{CRT1}$ | | | | | circRNA$_{GFP}$ | | | | |

Fig. 3

	10 h			18 h			32 h		
pGY1-35S::GFP	+	+	+	+	+	+	+	+	+
circRNA$_{GFP}$	−	+	−	−	+	−	−	+	−
linRNA$_{GFP}$	−	−	+	−	−	+	−	−	+
GFP (~35 kDa / ~25 kDa)									
% remaining	100	40	116	100	34	113	100	15	116
Actin (~55 kDa / ~40 kDa)									
% remaining	100	232	221	100	143	99	100	107	69
GFP/ACTIN	**1**	**0.17**	**0.52**	**1**	**0.24**	**1.14**	**1**	**0.14**	**1.68**

Fig. 4

A

B

Fig. 4

C

	WT			dcl1-11			ago1-27		
pGY1-35S::GFP	+	+	+	+	+	+	+	+	+
circRNA	–	GFP	CTR1	–	GFP	CTR1	–	GFP	CTR1
GFP % remaining	100	32	81	100	37	100	100	37	63
Actin % remaining	100	147	85	100	87	85	100	73	59
GFP/ACTIN	**1**	**0.22**	**0.95**	**1**	**0.43**	**1.17**	**1**	**0.51**	**1.06**

Fig. 5

Fig. 6