Title: Short-term social isolation acts on hypothalamic neurons to promote social

behavior in a sex- and context-dependent manner.

Xin Zhao[1], Yurim Chae[1], Destiny Smith[1], Valerie Chen[1], Dylan DeFelipe[1], Joshua W Sokol[1],

Archana Sadangi[1], and Katherine Tschida[1]

[1]Department of Psychology, Cornell University, Ithaca NY 14853

*Corresponding Author: Katherine Tschida, kat227@cornell.edu

Abstract

Social animals, including both humans and mice, are highly motivated to engage in
social interactions. Short-term social isolation promotes social behavior, but the neural
circuits through which it does so remain incompletely understood. Here, we sought to
identify neurons that promote social behavior in single-housed female mice, which exhibit
increased rates of social investigation, social ultrasonic vocalizations (USVs), and mounting
during same-sex interactions that follow a period of short-term (3-day) isolation. We first
used immunostaining for the immediate early gene Fos to identify a population of neurons in
the preoptic hypothalamus (POA) that increase their activity in single-housed females
following same-sex interactions (POA_{social} neurons) but not in single-housed females that did
not engage in social interactions. TRAP2-mediated chemogenetic silencing of POA_{social}
neurons in single-housed females significantly attenuates the effects of short-term isolation
on social investigation, USV production, and mounting. In contrast, caspase-mediated
ablation of POA_{social} neurons in single-housed females robustly attenuates mounting but
does not decrease social investigation or USV production. Optogenetic activation of POA_{social}
neurons in group-housed females promotes social investigation and USV production but
does not recapitulate the effects of short-term isolation on mounting. To understand whether
a similar population of POA_{social} neurons promotes social behavior in single-housed males,
we performed Fos immunostaining in single-housed males following either same-sex or
opposite-sex social interactions. These experiments revealed a population of POA neurons
that increase Fos expression in single-housed males following opposite-sex, but not same-
sex, interactions. Chemogenetic silencing of POA_{social} neurons in single-housed males during
interactions with females reduces mounting but does not affect social investigation or USV
production. These experiments identify a population of hypothalamic neurons that promote
social behavior following short-term isolation in a sex- and social context-dependent manner.

Keywords: social isolation, ultrasonic vocalization, preoptic hypothalamus, mounting,
female

Introduction

Humans and other social mammals find social interactions rewarding and are highly motivated to seek out social connections. Consequently, the experience of social isolation is aversive and impacts both our brains and our behaviors. While long-term isolation can lead to the emergence of anti-social behaviors in both humans and rodents (An et al., 2017; Arrigo and Bullock, 2008; Check et al., 1985; Hossain et al., 2020; Killgore et al., 2021; Ma et al., 2011, 2022; Machimbarrena et al., 2019; Matsumoto et al., 2005; Mears and Bales, 2009; Reid et al., 2022; Toth et al., 2011; Valzelli, 1973; Weiss et al., 2004; Wiberg and Grice, 1963; Zelikowsky et al., 2018), short-term isolation typically increases levels of social motivation and promotes social-seeking behaviors (Baumeister and Leary, 1995; Cacioppo et al., 2006; Cacioppo and Cacioppo, 2018; House et al., 1988; Lee et al., 2021; Niesink and van Ree, 1982; Panksepp and Beatty, 1980; Zhao et al., 2021). Alterations in social behavior are characteristic of many neurodevelopmental disorders, including autism spectrum disorder (Chevallier et al., 2012; Clements et al., 2018). How short-term social isolation acts on the brain to promote social behavior remains incompletely understood.

The mesolimbic pathway (i.e., projections from dopaminergic ventral tegmental neurons to the ventral striatum) plays an important role in regulating social motivation and social reward, during courtship as well as during same-sex interactions (Bariselli et al., 2018; Dai et al., 2022; Dölen et al., 2013; Gunaydin et al., 2014; Hung et al., 2017; Love, 2014; Melis et al., 2022; Resendez et al., 2020; Robinson et al., 2011; Solié et al., 2022; Tang et al., 2014; Walum and Young, 2018; Xiao et al., 2017). In line with its role in regulating social behavior, changes in the function of the mesolimbic pathway have been reported following long-term (weeks-long) social isolation and/or early-life social isolation (McWain et al., 2022; Musardo et al., 2022; Tan et al., 2021; Yorgason et al., 2016). The neural circuit changes that mediate the effects of short-term isolation on social behavior in adult animals are comparatively less explored, but here as well, recent studies in both humans and rodents have implicated changes in various populations of midbrain dopamine neurons (Inagaki et al., 2016; Matthews et al., 2016; Tomova et al., 2020). Beyond its effects on mesolimbic circuits, whether social isolation acts on additional neuronal populations to promote social behavior is unknown.

In recent work, we found that short-term (3-day) social isolation exerts robust effects on the social behaviors of C57BL/6J female mice (Zhao et al., 2021). Relative to group-housed females, single-housed females that subsequently engaged in same-sex interactions exhibited increased rates of social investigation, increased rates of USVs, and were also

observed to mount female social partners, a behavior never observed in pairs of group-
housed females (Zhao et al., 2021). The robust effect of short-term isolation on these three
aspects of female social behavior provides a powerful paradigm to identify neurobiological
changes that mediate the effects of short-term isolation on social behavior. In the current
study, we combined this behavioral paradigm with Fos immunostaining and the TRAP2
activity-dependent labeling approach (Allen et al., 2017; DeNardo et al., 2019) to identify and
characterize a population of neurons in the preoptic hypothalamus that increase their activity
in single-housed females following same-sex social interactions (i.e., POA$_{social}$ neurons). We
next asked whether silencing or ablation of POA$_{social}$ neurons in single-housed females
attenuates female social behaviors, and whether artificial activation of POA$_{social}$ neurons in
group-housed females promotes social behaviors. Finally, we extended a subset of these
experiments to single-housed males engaged in opposite-sex and same-sex interactions, to
understand whether short-term isolation acts on the POA to promote social behavior in a
manner that depends on either sex or social context. This study identifies novel
neurobiological mechanisms through which short-term social isolation acts on the brain to
promote social interaction. Our findings also add to an emerging literature indicating that the
POA regulates not only sexual behavior but also female social behavior during same-sex
interactions.

Results

Neurons in the preoptic hypothalamus increase Fos expression in socially isolated
female mice following same-sex social interactions

To identify changes in neuronal activity that may underlie the effects of short-term
isolation on female social behavior, we performed immunostaining for the immediate early
gene Fos in brain sections collected from group-housed and single-housed (3-days) subject
females following 30-minute social encounters in their home cages with a novel, group-
housed visitor female (Fig. 1A). In line with our previous behavioral findings (Zhao et al.,
2021), we observed that single-housed female residents spent more time investigating
visitors (Fig. 1B; t-test, p = 0.001) and in many trials mounted visitors, a behavior that was
not observed in group-housed residents (Fig. 1C; 0 of 12 group-housed residents and 11 of
13 single-housed residents mounted visitors; z-test for independent proportions, p < 0.001;
see Table S1 for complete statistical details). Female pairs that contained a single-housed
resident also produced higher rates of ultrasonic vocalizations (USVs) than pairs with a
group-housed resident (Fig. 1D; Mann-Whitney U test, p < 0.001). Although either female in

a dyad can produce USVs (Warren et al., 2020), the robust effects of short-term isolation on

the non-vocal social behaviors of single-housed females suggest that at least some of the

elevation in total USVs is driven by increased USV production by the single-housed resident.

Given the robust effects of short-term isolation on these three aspects of female social

behavior, we focused our analyses on two hypothalamic regions implicated in regulating

these behaviors: the preoptic area (POA), which regulates social approach (McHenry et al.,

2017), social reward (Hu et al., 2021), mounting (Floody, 1989; Karigo et al., 2021; Wei et

al., 2018) and USV production (Chen et al., 2021; Gao et al., 2019; Green et al., 2018;

Karigo et al., 2021; Michael et al., 2020); and the ventromedial hypothalamus (VMH), which

regulates mounting (Hashikawa et al., 2017; Karigo et al., 2021; Lee et al., 2014; Liu et al.,

2022). We also examined Fos expression within the caudolateral periaqueductal gray

(PAG), based on the well-established role of this region in the control of vocalization in

vertebrates and USV production in mice (Chen et al., 2021; Jürgens, 1994; Michael et al.,

2020; Tschida et al., 2019). To test whether any observed differences in Fos expression in

these three regions were associated with isolation-induced changes in social behavior rather

than baseline differences between groups, we also measured Fos expression in the POA,

the VMH, and the PAG of group-housed and single-housed females that did not engage in

social interaction with novel female visitors (Fig. 1E-F; Fig. S1).

These analyses revealed that baseline levels of Fos expression within the POA and

the VMH did not differ between group-housed and single-housed females (Fig. 1F, left and

middle, open bars; two-way ANOVA to analyze Fos expression within each brain region,

factor 1 = housing status, factor 2 = social interaction, followed by post-hoc Tukey's HSD

tests). Following social interactions with novel female visitors, single-housed females

exhibited robust increases in Fos expression within the POA (Fig. 1F, left; $p < 0.001$) but not

within the VMH (Fig. 1F, middle; $p > 0.05$). In contrast, Fos expression within these two brain

areas did not increase significantly in group-housed females that interacted with novel

female visitors (Fig. 1F; $p > 0.05$ for both comparisons). Similar to the POA, baseline Fos

expression within the PAG did not differ between group-housed and single-housed females

($p > 0.05$), and only single-housed females displayed increased PAG Fos expression

following social interactions with novel female visitors (Fig. 1F, right; $p < 0.001$), a finding

that further supports the idea that single-housed females increase USV production during

same-sex interactions. POA Fos expression was significantly and positively correlated with

the total amount of time spent in resident-initiated investigation for both group-housed and

single-housed females (Fig. S1C, left; linear regression, $p < 0.05$), as well as with the total

time spent mounting by single-housed resident females (Fig. S1C, middle; $p = 0.01$). In both

group-housed and single-housed female residents, Fos expression tended to correlate

positively with total USVs, but these relationships were not significant (Fig. S1C, right; $p >$

0.05 for both comparisons; see Fig. S1D-E for relationships of VMH Fos and PAG Fos to

vocal and non-vocal social behaviors). In summary, POA Fos expression increases

selectively in single-housed females following same-sex social interactions, and levels of

POA Fos expression are also well related to the production of specific types of social

behaviors by both group-housed and single-housed females.

To ask whether the effects of short-term isolation on female social behavior and POA

Fos expression are long-lasting, we measured social behaviors of female residents at three

timepoints: (1) on day 0, when female subjects were still group-housed; (2) on day 3, after

female subjects had been single-housed for 3 days; and (3) on day 17, after half of the

subject females had been re-group-housed with their same-sex siblings for two weeks and

the other half of the subject females remained single-housed for an additional two weeks

(Fig. 1G-I). Brains of re-group-housed and 17-day single-housed subject females were

collected after the day 17 social interaction, and Fos expression within the POA was

examined (Fig. 1J). Consistent with our earlier findings, rates of social investigation and USV

production significantly increased following 3 days of social isolation (Fig. 1G, I; $p < 0.05$ for

day 0 vs. day 3 in both groups for both behaviors). Following re-group-housing, time spent in

social investigation and rates of USV production tended to decrease to pre-isolation levels

(Fig. 1G, I, top plots; $p = 0.08$ for day 0 vs. day 17 investigation time and $p = 0.06$ for day 0

vs. day 17 total USVs in re-group-housed females). In contrast, females that were single-

housed for 17 days continued to spend increased time in social investigation (Fig. 1G,

bottom plot; $p < 0.05$ for day 0 vs. day 3 investigation and for day 0 vs. day 17 investigation),

and pairs containing 17-day single-housed residents continued to produce elevated rates of

USVs (Fig. 1I, bottom plot; $p < 0.05$ for day 0 vs. day 3 USVs and for day 0 vs. day 17

USVs). Time spent mounting tended to follow the same trends as rates of social

investigation and USV production in re-group-housed and 14-day single-housed females

(Fig. 1H). Along with the attenuation of female social behaviors following re-group-housing,

we also found that POA Fos expression was significantly lower in re-group-housed females

relative to 17-day single-housed females (Fig. 1J; t-test, $p < 0.001$). These findings support

the idea that changes in female social behavior following short-term isolation are reversible

and are accompanied by decreased POA Fos expression. Hereinafter, we refer to the

population of POA neurons that increase Fos expression in single-housed females that have

engaged in same-sex interactions as POA$_{social}$ neurons, and we next conducted experiments

to test whether functional manipulations of POA$_{social}$ neuronal activity impact the effects of

short-term isolation on female social behavior.

Chemogenetic inhibition of POA$_{social}$ neurons attenuates social investigation,

mounting, and USV production in single-housed females

If increased activity of POA$_{social}$ neurons promotes social behavior in single-housed
females, one prediction is that reducing the activity of POA$_{social}$ neurons in single-housed
females will attenuate the effects of isolation on female social behavior. To test this idea, we
employed the TRAP2 activity-dependent labeling strategy to chemogenetically silence
POA$_{social}$ neurons in single-housed females during social interactions with novel, group-
housed female visitors (Fig. 2A). Briefly, the POA of TRAP2 female mice was injected
bilaterally with a virus driving the Cre-dependent expression of the inhibitory DREADDs
receptor hM4Di. Three weeks later, females were single-housed for 3 days and then given a
30-minute social encounter with a novel, group-housed female visitor in their home cage.
Following the social interaction, resident females were given an I.P. injection of 4-
hydroxytamoxifen (4-OHT), which drives the transient expression of Cre recombinase in
recently active neurons and thereby enables the expression of hM4Di in POA$_{social}$ neurons.
Subject females remained single-housed for an additional 24 hours and then were re-group-
housed with siblings for 2 weeks. Subject females were then single-housed a second time
for 3 days and subsequently given a 30-minute same-sex interaction following I.P. injection
of either saline (control) or clozapine-n-oxide (CNO) (saline and CNO tests were run 3 days
apart, and the order was counterbalanced across experiments).

Comparison of the social behaviors of single-housed females between CNO and
saline sessions revealed that chemogenetic silencing of POA$_{social}$ neurons significantly
reduced resident-initiated investigation (Fig. 2B; N = 12; red points; two-way ANOVA with
repeated measures on one factor; $p < 0.01$). Inhibition of POA$_{social}$ neurons also significantly
reduced the proportion of females that exhibited mounting (Fig. 2C; McNemar's test for
paired proportions; $p = 0.04$ for POA$_{social}$ CNO vs. saline). Finally, inhibition of POA$_{social}$
neurons significantly reduced USV production (Fig. 2D; two-way ANOVA with repeated
measures on one factor; $p < 0.01$). In contrast, CNO treatment did not affect the production
of any of these social behaviors in single-housed females with GFP expressed in POA$_{social}$
neurons (Fig. 2B-D; N = 14; black points; $p > 0.05$ for all CNO vs. saline comparisons in the
POA$_{social}$-GFP control group). To investigate the specificity of these effects to chemogenetic
silencing of POA$_{social}$ neurons, we also performed control experiments in which activity-
dependent chemogenetic silencing in single-housed females was performed caudal to the
POA within the anterior hypothalamus (AH) (Fig. 2B-D; N = 12; brown points) or within the
VMH (Fig. 2B-D; N = 5; gray points). No significant effects of CNO treatment on resident-
initiated investigation, mounting, or total USVs were observed in these control groups (Fig.
2C-D; $p > 0.05$ for all). The effect of chemogenetic inhibition of POA$_{social}$ neurons to decrease
female social behavior also cannot be attributed to an overall decrease in movement (Fig.

2E; paired t-test; p > 0.5 for difference in movement between saline and CNO sessions; see
Methods).

We next asked whether the magnitude of the effects of chemogenetic silencing of
POA$_{social}$ neurons on different female social behaviors was related to the rates at which these
behaviors were produced during TRAPing sessions that immediately preceded 4-OHT
treatment. To this end, we compared total resident-initiated investigation, total resident-
initiated mounting, and total USVs produced during TRAPing sessions to changes in these
behaviors for each female following chemogenetic silencing of POA$_{social}$ neurons (calculated
as (CNO behavior – saline behavior)). These comparisons revealed no significant
relationships (Fig. S2A-C; linear regression; p > 0.05 for all). Although we designed our
experiments to TRAP and subsequently manipulate the activity of POA neurons that
upregulated Fos expression following the production of social behaviors in single-housed
females, one possibility is that our strategy inadvertently led to the labeling of POA neurons
that upregulate Fos in a manner that reflects the experience of social isolation per se. To
test this idea, we performed chemogenetic silencing of POA neurons in females that were
single-housed but were not given a social interaction prior to 4-OHT treatment (N = 5 non-
social controls; Fig. S2D). In these control single-housed females, CNO treatment had no
effect on social investigation, mounting, or USV production during same-sex interactions
(Fig. S2E-G; p > 0.05). In summary, we demonstrate that chemogenetic silencing of
POA$_{social}$ neurons reduces social investigation, mounting, and USV production in single-
housed female mice, and that these effects in turn require the production of social behaviors
during the TRAPing sessions that precede 4-OHT treatment.

Ablation of POA$_{social}$ neurons attenuates mounting in single-housed females

In previous work investigating the role of the POA in regulating rodent social
behaviors, studies have reported different effects on behaviors according to whether they
employed reversible or irreversible neuronal silencing strategies. Studies that used
chemogenetic or optogenetic methods to reversibly silence genetically-defined subsets of
POA neurons report decreases in both USV production in males (Chen et al., 2021; Karigo
et al., 2021) and in mounting in males and females during interactions with female social
partners (Gao et al., 2019; Karigo et al., 2021). In contrast, studies employing caspase-
mediated ablation of genetically-defined subsets of POA neurons (Gao et al., 2019; Wei et
al., 2018) or electrolytic lesions of the POA (Bean et al., 1981) report decreased mounting
but no effects on rates of USV production. To test whether permanent ablation of POA$_{social}$
neurons attenuates social behaviors in single-housed females in a manner similar to the
effects of chemogenetic inhibition, we used the TRAP2 activity-dependent labeling strategy

to express caspase in and to thereby ablate POA$_{social}$ neurons (Fig. 3A; see Methods). Vocal

and non-vocal social behaviors of resident females were compared pre- and post-ablation,

and the same measurements were made in control females expressing GFP in POA$_{social}$

neurons.

Similar to the effects of chemogenetically inactivating POA$_{social}$ neurons, ablation of

POA$_{social}$ neurons significantly reduced mounting in single-housed females (Fig. 3C;

McNemar's test for paired proportions; p = 0.03 for proportion of trials with mounting on pre-

4-OHT vs. post-4-OHT tests in POA$_{social}$-caspase females; p > 0.05 for pre-4-OHT vs. post-

4-OHT in POA$_{social}$-GFP females). In contrast to the effects of chemogenetic inhibition of

POA$_{social}$ neurons, we found that caspase-mediated ablation of POA$_{social}$ neurons did not

affect rates of social investigation in single-housed females, although both experimental and

control females spent more time investigating visitors in the post-4-OHT session (Fig. 3B;

two-way ANOVA with repeated measures on one factor; p > 0.05 for main effect of group, p

< 0.01 for main effect of time, p > 0.05 for interaction effect). Ablation of POA$_{social}$ neurons

also failed to reduce USV production in pairs containing single-housed females (Fig. 3D;

two-way ANOVA with repeated measures on one factor; p > 0.05 for pre-4-OHT vs. post-4-

OHT total USVs in POA$_{social}$-caspase females). Together with our chemogenetic inhibition

data, these results show that both reversible inhibition or irreversible ablation of POA$_{social}$

neurons in single-housed female mice reduces mounting, whereas only chemogenetic

inhibition of POA$_{social}$ neurons attenuates the effects of short-term isolation on female social

investigation and USV production.

One possibility is that a difference in the efficacy of the strategies used to irreversibly

ablate vs. reversibly inhibit female POA$_{social}$ neurons accounts for the difference in their

effects on female social behaviors. Indeed, although we used TRAP2 homozygous females

in the POA$_{social}$-hM4Di dataset (Fig. 2), the majority of the females used in the POA$_{social}$-

caspase dataset were TRAP2 heterozygotes (11 of 15 females were generated by crossing

TRAP2 mice with Ai14 mice, and 4 of 15 were TRAP2 homozygotes; absence of tdTomato

labeling in TRAP2;Ai14 females was used to estimate viral spread; see Methods). To test

this possibility, we directly compared the effects on female social behavior of caspase-

mediated ablation of POA$_{social}$ neurons in TRAP2 heterozygous females (N = 11

TRAP2;Ai14, a subset of those plotted in Fig. 3B-D) and TRAP2 homozygous females (N =

9, including N = 4 from Fig. 3B-D). These experiments revealed that although ablation of

POA$_{social}$ neurons in TRAP2 heterozygous females tended to reduce mounting, this effect

was non-significant (Fig. S3B; McNemar's test for paired proportions; p = 0.11). As in the

original dataset, ablation of POA$_{social}$ neurons in TRAP2 homozygous females significantly

reduced mounting (Fig. S3B; p = 0.04). Ablation of POA$_{social}$ neurons failed to reduce social

investigation or USV production in either genotype, and both of these behaviors were slightly

and significantly elevated following 4-OHT treatment in TRAP2 heterozygous females (Fig.

S3A,C; p < 0.05 for pre-4-OHT vs. post-4-OHT social investigation and USVs in TRAP2;Ai14

females).

To ensure that ablation of POA$_{social}$ neurons in TRAP2 homozygous females was

effective in eliminating POA neurons that would normally upregulate Fos following same-sex

interactions in single-housed females, we compared counts of Fos-positive POA neurons in

a subset of TRAP2 homozygous POA$_{social}$-caspase females to those recorded in control

group-housed and single-housed females that either did or did not engage in same-sex

interactions (Fig. S3D; control female Fos data are the same as those plotted in Fig. 1D).

This analysis revealed that ablation of POA$_{social}$ neurons reduced Fos expression in the POA

below the levels normally seen in either group-housed or single-housed females following

same-sex interactions, down to levels observed in group-housed and single-housed females

that were not given social interactions (Fig. S3D; p < 0.05 for POA$_{social}$-caspase vs. group-

housed and single-housed social interaction control groups; p > 0.05 for POA$_{social}$-caspase

vs. group-housed and single-housed baseline control groups). Moreover, the remaining POA

Fos expression in TRAP2 homozygous POA$_{social}$-caspase females was not significantly

related to either rates of social investigation or to USV production (Fig. S3E-F; linear

regression; p > 0.05 for both; compare to relationships in Fig. S1C). Taken together, these

findings show that ablation of POA$_{social}$ neurons in TRAP2 homozygous females is more

effective at attenuating single-housed female social behaviors than ablation in TRAP2

heterozygotes. However, in spite of a robust reduction in social behavior-driven POA Fos

expression, the effects of irreversible ablation of POA$_{social}$ neurons on female social behavior

differ from those of chemogenetic inhibition.

Optogenetic activation of POA$_{social}$ neurons elicits USV production

To understand whether artificial activation of POA$_{social}$ neurons can recapitulate the

effects of short-term isolation on female social behavior, we assessed the effects of

optogenetic activation of POA$_{social}$ neurons on the social behaviors of group-housed females.

The TRAP2 strategy was used to express either channelrhodopsin (ChR2) or GFP in

POA$_{social}$ neurons (Fig. 4A; see Methods), and females were re-group-housed for two weeks

before beginning optogenetic activation experiments. The effects of optogenetically

activating POA$_{social}$ neurons were first assessed for each subject female in a 5-minute solo

session, in which the female was tested alone in a behavior chamber while pulses of blue

light were delivered unilaterally to the POA (473 nm, 10 mW, 20-50 Hz, 10-20 ms pulses, 5-

10s train durations). The effects of optogenetically activating POA$_{social}$ neurons were then

assessed for each subject female in a 20-minute social session, in which a novel, group-
housed female visitor was added to the behavior chamber. The pair was allowed to interact
in the absence of optogenetic stimulation for the first and last 5 minutes of the social
session, and pulses of blue light were delivered to the POA of the subject female throughout
the middle 10 minutes of the session (Fig. 4A).

321   When POA_{social}-ChR2 females were tested alone, we found that optogenetic
activation of POA_{social} neurons elicited weak-to-moderate USV production in 4 of 8 females,
but the comparison of USV rates from pre-laser baseline to the laser stimulation period was
not significant at the level of the entire group (Fig. 4B; Mann Whitney U test performed on
the difference in USV rates (laser - pre-laser), p = 0.09). In POA_{social}-GFP control females,
laser stimulation failed to elicit USV production (0 ± 0 USVs elicited in N = 6 POA_{social}-GFP
controls). Interestingly, we found that when laser stimulation was applied during social
sessions, optogenetic activation of POA_{social} neurons more readily elicited USV production
than in solo sessions (Fig. 4C; USVs elicited by blue laser stimulation in 7 of 8 POA_{social}-
ChR2 females; Mann Whitney U test performed on the difference in USV rates (laser - pre-
laser), p = 0.006). Moreover, optogenetic activation elicited higher rates of USVs when
applied at times when subject females were in close proximity to visitor females (within 2
mouse body lengths) as compared to times when the females were farther apart (mean
increase in USV rates from pre-laser to laser period was 2.96 ± 2.32 USVs/s for "near"
stimulations, 1.84 ± 1.75 USVs/s for "far" stimulations; paired t-test performed on the
difference in USV rates (laser - pre-laser) for "far" vs. "near" stimulations; p = 0.02). In
summary, optogenetic activation of POA_{social} neurons elicits USV production in group-housed
females, and the efficacy of this effect is modulated by social context and proximity to a
social partner.
340   We next considered whether optogenetic activation of POA_{social} neurons modulates
social investigation. To ask whether optogenetic activation could elicit social investigation,
we first excluded laser stimulations in which resident-initiated social investigation occurred at
any time in a 5-second window prior to laser onset. With these stimulations excluded, we
then calculated the proportion of stimulations in which resident-initiated social investigation
occurred following laser onset but prior to laser offset. This analysis revealed that POA_{social}-
ChR2 females were more likely to engage in social investigation of the visitor female
following laser stimulation than were POA_{social}-GFP females (Fig. 4E-F; p = 0.03 for
difference in proportion of laser stimulations followed by social investigation). To ask
whether optogenetic activation of POA_{social} neurons could prolong social investigation bouts,
we performed a second analysis in which social investigation bouts that occurred during the
middle 10 minutes of the social session were separated into those that temporally

overlapped with laser stimulation and those that did not. This analysis revealed that social

investigation bouts that overlapped with laser stimulation were significantly longer than those

without overlap in POA_{social}-ChR2 females but not in POA_{social}-GFP females (Fig. 4G; two-

way ANOVA with repeated measures on one factor; $p = 0.001$ for overlapping vs. non-

overlapping for POA_{social}-ChR2 females; $p > 0.05$ for overlapping vs. non-overlapping for

POA_{social}-GFP females). Finally, there was a non-significant trend toward increased overall

resident-initiated investigation during the middle 10 minutes of the social session for

POA_{social}-ChR2 females vs. POA_{social}-GFP females (mean total investigation time = 72.1 ±

54.3s for N = 9 POA_{social}-ChR2 females vs. 23.3 ± 16.3s for N = 6 POA_{social}-ChR2 females; t-

test, $p = 0.07$).

In contrast to the effects on USV production and social investigation, optogenetic

activation of POA_{social} neurons only infrequently elicited mounting (activation elicited n =1

bout of mounting in N = 1 POA_{social}-ChR2 female and n = 0 bouts of mounting in the

remaining N = 8 POA_{social}-ChR2 females). In summary, optogenetic activation of POA_{social}

neurons partially mimics the effects of short-term social isolation on female behavior by

promoting USV production and social investigation during same-sex interactions.

Experiments to test whether POA neurons regulate social behaviors during same-sex

interactions in group-housed females

Given our findings that POA_{social} neurons promote social behaviors in single-housed

females during same-sex interactions, we wondered whether a similar population of POA

neurons regulates the social behaviors of group-housed females during same-sex

interactions. This idea is supported by our earlier finding that Fos expression in the POA of

group-housed females is significantly related to rates of resident-initiated social investigation

(Fig. S1C, left; $p = 0.01$) and tends to be related to rates of USV production (Fig. S1C, right;

$p = 0.05$) during same-sex interactions.

To further test this idea, we performed two experiments to manipulate the activity of

POA neurons that were TRAPed following same-sex social interactions in group-housed

females (Fig. 5A,E). In the first set of experiments, we used the TRAP2 strategy to express

hM4Di in POA neurons that upregulated Fos expression following same-sex interactions in

group-housed females. Two weeks following 4-OHT treatment, these females were single-

housed, and the effect of CNO treatment on female social behavior during same-sex

interactions was assessed. We found that chemogenetic inhibition of group-housed-TRAPed

POA neurons failed to alter resident-initiated social investigation or total USVs when females

were subsequently single-housed and tested with female social partners (Fig. 5B,D; N = 5

group-housed-TRAPed hM4Di females; $p > 0.05$ for CNO vs. saline sessions for both

behaviors). Unfortunately, none of the females in this experimental cohort mounted in the saline or CNO sessions, so we were not able to evaluate the effect on mounting (Fig. 5C). In the second set of experiments, we used the TRAP2 strategy to express ChR2 in POA neurons that upregulate Fos expression following same-sex interactions in group-housed females, and female subjects remained group-housed for subsequent tests. Optogenetic activation of group-housed-TRAPed POA neurons failed to elicit statistically significant USV production (Fig. 5F), did not promote social investigation (Fig. 5G-H), and did not elicit mounting (mounting observed in 0 of 6 trials).

Because group-housed females spend a relatively small amount of time engaged in social investigation, do not exhibit mounting, and produce low rates of USVs relative to single-housed females (Fig. 1B-D), one important caveat to these experiments is that TRAP-based activity-dependent labeling may not work efficaciously in mice that produce low rates of social behaviors. That is to say that although POA neurons may regulate social behaviors in group-housed females, these POA neurons may not strongly upregulate Fos following group-housed social interactions (see Fig. 1F). To test whether the POA regulates the social behaviors of group-housed females using a non-activity-dependent viral strategy, we injected a non-Cre-dependent inhibitory DREAADs virus (AAV-hSyn-hM4Di) into the POA of group-housed B6 female mice (Fig. 5I). Three weeks later, the effects of CNO treatment on social behavior during same-sex interactions were tested. We found no effects of CNO treatment on resident-initiated social investigation in group-housed females (Fig. 5J; $p > 0.05$), and as expected, none of the group-housed females exhibited mounting in either saline or CNO sessions (Fig. 5K). Notably, we found that although rates of USV production were quite low, chemogenetic inhibition of POA neurons significantly reduced total USVs produced during same-sex interactions between group-housed females (Fig. 5L; paired t-test, $p = 0.048$). Following these tests, female subjects were single-housed for 3 days and then tested a second time. Consistent with our previous experiments, single-housing dramatically elevated rates of female social investigation, mounting, and USV production (compare saline day panels in Fig. 5J-L vs. Fig. 5M-O). We found that chemogenetic inactivation of POA neurons in single-housed females dramatically reduced all three behaviors (Fig. 5M-O; paired t-tests, $p < 0.05$ for all behaviors). These additional experiments reinforce our TRAP2-based findings that the POA regulates the social behaviors of single-housed female mice. These experiments also suggest that the POA may regulate the social behaviors of group-housed females, but that any effects may be more challenging to detect given the low rates of social behaviors produced by group-housed females during same-sex interactions.

Characterization of female POA$_{social}$ neuron molecular markers and axonal projections

Previous studies have found that USV production can be elicited in female and male

mice by artificial activation of VGAT$^+$ POA neurons (Gao et al., 2019), Esr1$^+$ POA neurons

(which are predominantly VGAT$^+$) (Chen et al., 2021; Michael et al., 2020), as well as POA

neurons that send axonal projections to the caudolateral PAG (which are predominantly

VGAT$^+$) (Chen et al., 2021; Michael et al., 2020). To ask to what extent POA$_{social}$ neurons

overlap with these previously described populations, we first evaluated the neurotransmitter

phenotype of POA$_{social}$ neurons by performing two-color in situ hybridization for c-fos mRNA

and vesicular GABA transporter (VGAT) mRNA and calculating the percentage of Fos$^+$ POA

neurons that co-expressed VGAT. This analysis revealed that a majority of POA$_{social}$ neurons

are GABAergic (Fig. S2A-B; N = 4, 76 ± 8.8%). We next used the TRAP2 activity-dependent

labeling strategy to express GFP in POA$_{social}$ neurons and found GFP-positive axons within

the caudolateral PAG, indicating that at least some POA$_{social}$ neurons send axonal

projections to the PAG (Fig. S2C-D; see Methods). Finally, we combined retrograde tracing

from the caudolateral PAG with Fos immunostaining to quantify the percentage of PAG-

projecting POA neurons that increase Fos expression in single-housed females following

same-sex interactions. This experiment revealed that around 20% of PAG-projecting POA

neurons express Fos in single-housed females following same-sex interactions (Fig. S2E-F;

N = 4 females, percentage of tdTomato neurons that are Fos-positive = 18.3 ± 2.9%). These

findings suggest that a subset of POA$_{social}$ neurons overlap with GABAergic, PAG-projecting

POA neurons that have been demonstrated in previous work to promote USVs via

disinhibition of excitatory PAG neurons important to USV production, although only ~20% of

PAG-projecting POA neurons upregulate Fos in association with female USV production

(Chen et al., 2021; Michael et al., 2020).

POA neurons increase their activity in single-housed male mice following opposite-
sex but not same-sex social interactions

Given our findings that POA$_{social}$ neurons contribute to isolation-induced changes in

the social behaviors of female mice, we next wondered whether a similar population of POA

neurons contributes to isolation-induced changes in social behavior in male mice. To

address this question, we measured the vocal and non-vocal social behaviors of sexually

naïve males, which were either group-housed with same-sex siblings or single-housed for

three days and then given a 30-minute social interaction with a novel, group-housed visitor.

To consider the effects of isolation on male social behavior in different social contexts, males

were given either a social encounter with a same-sex visitor (MM context) or with an

opposite-sex visitor (MF context). Following these social interaction tests, we collected the

brains of the subject males and performed immunostaining to measure Fos expression

within the POA. With respect to resident-initiated investigation, we found significant main

effects of both housing and social context: single-housed males spent more time

investigating visitors than group-housed males, and males spent more time investigating

female visitors than male visitors (Fig. 5A; two-way ANOVA, p = 0.02 for main effect of

housing; p < 0.001 for main effect of social context; p > 0.05 for interaction effect). With

respect to mounting, we found that single-housed males were more likely than group-housed

males to mount female visitors, but the proportion of trials with mounting did not differ for

single-housed males vs. group-housed males during male-male interactions (Fig. 5B; z-tests

for independent proportions; p = 0.046 for group-housed vs. single-housed with female

visitor, p > 0.05 for group-housed vs. single-housed with male visitor). Similarly, there was a

context-dependent effect of social isolation on male USV production, whereby only single-

housed males that interacted with female visitors exhibited increased USV production

relative to group-housed males (Fig. 5C; two-way ANOVA with post-hoc Tukey's HSD tests;

p < 0.001 for total USVs in single-housed MF vs. group-housed MF trials; p > 0.05 for total

USVs in single-housed MM vs. group-housed MM trials). The finding that short-term isolation

exerts larger effects on male social behavior during subsequent opposite-sex interactions

relative to same-sex interactions is consistent with prior work (Zhao et al., 2021). When we

examined POA Fos expression in these four groups of males, we found that POA Fos was

significantly elevated in single-housed males following interactions with females relative to

the other three groups (Fig. 5D; two-way ANOVA, Tukey's post-hoc HSD tests; p < 0.05 for

difference in POA Fos between single-housed MF and all other groups). In summary, the

effects of short-term isolation on male social behavior are context-dependent, and increased

Fos expression within the POA is seen in single-housed males following interactions with

females, a context marked by increased male social investigation, mounting, and USV

production.

To test whether neural activity in male POA_{social} neurons contributes to isolation-

induced changes in male social behavior, we used the TRAP2 strategy to chemogenetically

silence POA_{social} neurons in single-housed males during social interactions with novel, group-

housed females (see Methods). The vocal and non-vocal behaviors of single-housed subject

males were measured and compared during 30-minute social interactions following I.P.

injection of either saline or CNO. Control males were treated identically but were injected

with a virus to drive expression of GFP in POA_{social} neurons. As in females, chemogenetic

inhibition of male POA_{social} neurons significantly reduced the proportion of trials with

mounting (Fig. 5G; z-test for difference between proportions; p = 0.04 for proportion of trials

with mounting on CNO vs. saline days in POA_{social}-hM4Di males; p > 0.05 for proportion of

trials with mounting on CNO vs. saline days in POA_{social}-GFP males). In contrast to our

findings in females, chemogenetic inhibition of male POA_{social} neurons did not alter rates of

resident-initiated social investigation (Fig. 5F; two-way ANOVA with repeated measures on

one factor; $p < 0.001$ for main effect of group; $p > 0.05$ for main effect of drug and for

interaction effect) and also did not affect total USVs (Fig. 5H; two-way ANOVA with repeated

measures on one factor; $p > 0.05$ for main effects and interaction effect). Although we are

uncertain of the reason for the unexpected difference in social investigation times between

POA$_{social}$-hM4Di and POA$_{social}$-GFP males, we note that these cohorts of mice were tested at

different times and thus interacted with different groups of female visitors, which may have in

turn contributed to different rates of social investigation between experimental and control

males.

504   Given previous work implicating the POA in the regulation of male mounting (Floody,

1989; Karigo et al., 2021; Wei et al., 2018) as well as male USV production (Chen et al.,

2021; Gao et al., 2019; Green et al., 2018; Karigo et al., 2021; Michael et al., 2020), we were

surprised that chemogenetic silencing of POA$_{social}$ neurons in males attenuated mounting but

not USV production. To begin to understand this result, we considered rates of social

behaviors produced by POA$_{social}$-hM4Di males during the TRAPing sessions that preceded

4-OHT treatment and compared these to the TRAPing session behaviors produced by

POA$_{social}$-hM4Di females. Interestingly, we found that POA$_{social}$-hM4Di males spent

significantly more time mounting female visitors during TRAPing sessions than POA$_{social}$-

hM4Di females (Fig. S5B; Mann Whitney U test; $p = 0.004$). In comparison to females,

POA$_{social}$-hM4Di males spent less time engaged in social investigation of the female visitor

during TRAPing sessions (Fig. S5A; t-test; $p = 0.02$) and also tended to produce lower rates

of USVs (Fig. S5C; t-test, $p = 0.054$). Based on these sex differences in TRAPing session

social behavior, one possibility is that the bias of male TRAPing session behavior toward

mounting also resulted in a bias toward TRAP2-based labeling (and subsequent

chemogenetic inhibition) of POA mounting-related neurons. We note also that because all

animals in the study received the same total amount of 4-OHT (150 μL of 10 mg/mL 4-OHT)

and because males typically weigh more than females, the effectively smaller dosage of 4-

OHT delivered to males could have reduced TRAPing efficacy relative to females and could

have contributed to sex differences in the observed effects on behavior of chemogenetically

silencing POA$_{social}$ neurons. In summary, we find that chemogenetic silencing of male

POA$_{social}$ neurons reduces mounting during subsequent social interactions with females but

does not reduce social investigation or USV production, a pattern of results that differs from

the effects on single-housed female social behavior of chemogenetically silencing female

POA$_{social}$ neurons.

Discussion

In the current study, we identify and characterize a population of preoptic

hypothalamic neurons that contribute to the effects of short-term social isolation on the

social behaviors of mice. These POA$_{social}$ neurons exhibit increased Fos expression in

single-housed female mice following same-sex social interactions, and this increase in Fos

expression scales positively with the time females spend investigating and mounting female

visitors and tends also to scale with rates of USVs. Chemogenetic silencing of POA$_{social}$

neurons attenuates the effects of social isolation on female social behavior, significantly

reducing social investigation, mounting, and USV production. In contrast, irreversible

ablation of POA$_{social}$ neurons significantly reduces mounting in single-housed females but

does not reduce rates of social investigation or USVs. Optogenetic activation of POA$_{social}$

neurons partially recapitulates the effects of short-term isolation on female behavior and

promotes social investigation and USV production in female mice. Finally, we extended our

experiments to male mice to understand whether similar POA neurons may mediate

changes in male social behavior following short-term isolation. We find that short-term

isolation exerts more robust effects on male behavior during subsequent interactions with

females than during subsequent interactions with males, and increased POA Fos expression

is seen in single-housed males following social interactions with females but not following

interactions with males. Interestingly, chemogenetic silencing of these POA$_{social}$ neurons

reduces mounting but has no effect on social investigation and USV production, in contrast

to the effects of chemogenetically silencing POA$_{social}$ neurons in females. Together, these

experiments identify a population of preoptic hypothalamic neurons that promote social

behaviors in single-housed mice in a manner that depends on sex and social context.

An extensive body of past work has implicated the POA in the regulation of male

sexual behavior, including in the regulation of male courtship vocalizations in both rodents

and birds (Alger and Riters, 2006; Bean et al., 1981; Gao et al., 2019; Merari and Ginton,

1975; Riters, 2012; Riters et al., 2000; Riters and Ball, 1999; Wei et al., 2018). Past work

has also shown that activation of genetically-defined subsets of POA can elicit the

production of USVs in both male and female mice (Chen et al., 2021; Gao et al., 2019;

Karigo et al., 2021; Michael et al., 2020). However, whether the POA regulates natural USV

production in female mice remained unclear. In the current study, we demonstrate that

reversible silencing of POA neurons that increase their activity during same-sex interactions

decreases female USV production, indicating that the POA regulates the production of USVs

by single-housed females engaged in same-sex interactions. Using a non-activity-dependent

viral strategy, we also provide evidence that POA neurons regulate USV production in

group-housed females. Whether these same neurons regulate female USV production in

other behavioral contexts, including during interactions with male partners, remains an

important open question. In addition to attenuating USV production, we found that silencing

of POA$_{social}$ neurons in single-housed females significantly reduced social investigation and

mounting of same-sex partners. Our retrograde and anterograde tracing experiments

demonstrate that at least a subset of POA$_{social}$ neurons project to the caudolateral PAG

where neurons important for USV production reside, and an attractive possibility is that

POA$_{social}$ neurons promote USV production via disinhibition of PAG-USV neurons as

previously demonstrated for genetically-defined subsets of POA neurons (Chen et al., 2021;

Michael et al., 2020; Tschida et al., 2019). Future experiments will be required to determine

whether PAG-projecting POA$_{social}$ neurons also regulate social investigation and mounting, or

alternatively, whether distinct molecularly-defined or projection-defined subsets of POA$_{social}$

neurons differentially regulate these different aspects of female social behavior. A recent

study found that optogenetic activation of Tacr1[+] POA neurons elicits both mounting and

USV production in male mice and that optogenetic activation of the axon terminals of these

neurons within the PAG also promotes mounting, suggesting that some POA neurons may

regulate the production of multiple social behaviors. However, another study found that

distinct subsets of male Esr1[+] POA neurons exhibit selective tuning in their neuronal activity

to sniffing vs. mounting during interactions with females (Yang et al., 2023), although it

remains to be tested whether these neuronal subsets functionally regulate distinct social

behaviors during male-female (or female-female) interactions. The latter organization would

be reminiscent of how projection-defined subsets of galanin-expressing POA neurons

regulate different aspects of parental behavior (Kohl et al., 2018).

Although reversible inhibition of POA$_{social}$ neurons reduced both social investigation

and USV production in single-housed females, permanent caspase-mediated ablation of

these neurons significantly reduced mounting but had no effects on social investigation or

USV production. These differences are largely consistent with prior studies that reported

effects on both mounting and USV production following reversible manipulations of POA

activity and effects on mounting but not on USV production following irreversible

manipulations of POA activity (Bean et al., 1981; Chen et al., 2021; Gao et al., 2019; Karigo

et al., 2021; Wei et al., 2018). One possibility is that POA$_{social}$ neurons do not directly

regulate USV production and social investigation but rather that reversible silencing of these

neurons causes off-target disruptions of neural activity in interconnected brain regions that in

turn directly regulate USV production. Such a relationship was demonstrated for motor

cortex, whereby reversible silencing of motor cortex disrupted performance of a learned

forelimb reaching task in rats, while permanent lesions of motor cortex had no effect on task

performance after learning (Otchy et al., 2015). However, our finding that optogenetic

activation of POA$_{social}$ neurons elicits USV production, along with past work demonstrating

that POA activation elicits USV production (Chen et al., 2021; Gao et al., 2019; Karigo et al.,

2021; Michael et al., 2020) supports the idea that the POA directly regulates USV

production. Similarly, our finding that optogenetic activation of POA$_{social}$ neurons promotes

social investigation supports the interpretation that these neurons directly regulate this

behavior. An alternative explanation for the apparently contradictory effects of chemogenetic

silencing vs. ablation of POA$_{social}$ neurons is that while the POA plays an obligatory role in

mounting behavior (at least in certain contexts, see below), compensatory changes in

additional, non-POA circuits that promote USV production and social investigation are

sufficient to compensate for the permanent loss of POA$_{social}$ neurons, leaving these

behaviors unperturbed following POA$_{social}$ neuronal ablation. Our finding that ablation of

POA$_{social}$ neurons dramatically reduces POA Fos expression following female-female

interactions, and that the remaining Fos expression is uncorrelated with rates of USV

production and social investigation, support the latter idea. The identification of forebrain-to-

midbrain circuits that regulate USV production in both females and males remains an

important future goal.

Our finding that POA$_{social}$ neurons regulate female-female mounting extends recent

work examining the role of hypothalamic regions in regulating male mounting behavior in

different social contexts (Karigo et al., 2021). Although the POA plays a well-established role

in the regulation of mounting behavior by male rodents during interactions with females,

interactions which are typically marked by high rates of USV production and are considered

affiliative, the authors found that the VMH regulates male mounting behavior during

interactions with other males, which is typically not accompanied by USV production and

often precedes fighting. Taken together with the findings of the current study, we conclude

that the POA regulates mounting behavior in both male and female mice that is directed

toward female social partners and is accompanied by USV production. The question of

whether the social behaviors exhibited by single-housed female mice during same-sex

interactions are indeed affiliative and how these behaviors shape future interactions with

female social partners are important questions that remain to be addressed. Although

chemogenetic silencing and ablation of POA$_{social}$ neurons reduced female mounting, we were

not able to reliably elicit mounting behavior through optogenetic activation of POA$_{social}$

neurons. We note that previous studies that elicited mounting in male mice via optogenetic

activation of Esr1^{+} POA neurons, Tacr1^{+} POA neurons, or POA neurons that co-express

Esr1 and VGAT used longer laser stimulation trains (15-30s) than we employed in the

current study (Bayless et al., 2023; Karigo et al., 2021; Wei et al., 2018). It is possible that

longer periods of optogenetic activation of POA$_{social}$ neurons would be more effective in
eliciting mounting in group-housed females, and another possibility is that POA$_{social}$ neurons
overlap somewhat but not perfectly with the genetically-defined groups of POA neurons
manipulated in those prior studies. Regardless, our finding that optogenetic activation of
POA$_{social}$ neurons promotes USV production and social investigation efficaciously relative to
mounting indicates that effects on these behaviors are not secondary to effects on female
mounting.

To understand whether a similar population of POA neurons regulates changes in
social behavior following short-term isolation in males, we extended our experiments to
single-housed males that subsequently interacted with either a male social partner or with a
female social partner. Consistent with our prior work (Zhao et al., 2021), we found that short-
term isolation exerts more robust effects on male behavior during subsequent interactions
with females than during interactions with males. Although single-housed males exhibited
increased rates of social investigation while interacting with both males and females relative
to group-housed males, single-housed males that interacted with females exhibited
increases in all measured social behaviors (social investigation, USV production, and
mounting), a pattern of behavioral changes that is similar to what we observed in single-
housed females. We also observed that a population of POA neurons increased Fos
expression in single-housed males following interactions with female visitors but not
following interactions with male visitors. Unexpectedly, we found that the effects of
chemogenetically inhibiting POA$_{social}$ neurons in males differed from those in females.
Namely, while reversible inhibition of POA$_{social}$ neurons reduced mounting during interactions
between single-housed subject males and female visitors, there were no effects on rates of
social investigation and USV production. These findings differ also from recent work showing
that chemogenetic inhibition (Chen et al., 2021) or optogenetic inhibition (Karigo et al., 2021)
of Esr1$^+$ POA neurons reduces USV production in male mice. Although the factors that
account for this difference in results are unclear, one possibility is that our TRAP2-based
viral labeling in males is biased toward POA neurons that regulate mounting as compared to
POA neurons that regulate USV production, although as stated above, more work is
required to understand whether these different social behaviors are regulated by distinct or
overlapping subsets of POA neurons. Future experiments can also explore to what extent
female and male POA$_{social}$ neurons represent molecularly, anatomically, and functionally
similar or dissimilar neuronal populations.

The current study adds to an emerging body of literature implicating the POA in the
regulation of social behavior in both females and males, including in non-sexual social

contexts (Fukumitsu et al., 2022; Hu et al., 2021; Liu et al., 2023; McHenry et al., 2017; Wu et al., 2021). Our findings also complement recent work that has described a role for the POA in regulating behavioral responses to social isolation in female mice. A recent study using BALB/c mice found that reunion with same-sex cagemates following short-term "somatic isolation" (i.e., separation from cagemates via a partition that permits visual, auditory, and olfactory signaling but prevents physical contact) increases the activity of calcitonin-receptor (Calcr) expressing POA neurons (Fukumitsu et al., 2023). Knockdown of Calcr expression in these neurons reduced social-seeking behaviors directed at the cage partition exhibited by single-housed females, and chemogenetic activation of these neurons increased partition-biting behavior in single-housed females. Another recent study using female FVB/NJ mice applied a TRAP2-based intersectional approach to identify POA neurons that increase their activity following reunion with same-sex cagemates ($MPN_{reunion}$ neurons) (Liu et al., 2023). Optogenetic activation of these neurons did not alter social investigation in group-housed females, whereas activation of these neurons during reunion with cagemates following short-term social isolation decreased social investigation. However, optogenetic inhibition of $MPN_{reunion}$ during reunion did not alter rates of social investigation in single-housed females. Although we did not test the effects of activating POA_{social} neurons in single-housed females in the current study, the difference in the effects of inhibiting POA_{social} neurons on the behavior of single-housed females (reduced rates of social investigation, USV production, and mounting) relative to the effects of optogenetic inhibition of $MPN_{reunion}$ neurons in Liu et al. (no effect on social investigation during reunion) suggest that these represent distinct populations of POA neurons. We note that in the current study, single-housed female subjects remained single-housed for 24 hours following social interaction TRAPing sessions, a design intended to maximize viral labeling of POA neurons that promote increased female social behaviors and that would likely in turn minimize viral labeling of POA neurons that promote social satiety. These studies highlight a complex role for the POA in regulating multiple aspects of changes in social behavior following short-term social isolation. Although some studies of social isolation have avoided the use of C57BL/6J female mice, the robust triad of changes in social behavior exhibited by these females following short-term isolation affords a powerful opportunity to continue investigating neural circuit mechanisms through which short-term social isolation promotes social behaviors, as well as to investigate how hypothalamic circuits regulate the coordinated production of suites of social behaviors during female-female social interactions.

Acknowledgements: Thanks to Frank Drank and other CARE staff for their excellent
mouse husbandry. All experimental design schematics included in the figures were created
with Biorender.com.

Author contributions: X.Z. and K.A.T. designed the experiments. X.Z. and Y.C. conducted
the experiments. X.Z., Y.C., D.S., V.C., D.D., J.W.S., A.S., and K.A.T. analyzed data. X.Z.
and K.A.T. wrote the manuscript, and all authors approved the final version.

Declaration of Interests: The authors declare no competing interests.

 Figure legends

Figure 1. The POA contains neurons that increase Fos expression in single-housed
females following same-sex interactions. (A) Schematic of experiment to measure Fos
expression in group-housed and single-housed females following same-sex social
interactions. (B) Total time spent engaged in resident-initiated social investigation for group-
housed residents (teal) and single-housed residents (maroon). (C) Same as (B), for
proportion of trials with resident-initiated mounting. (D) Same as (B), for total USVs recorded
from pairs containing group-housed or single-housed residents. (E) Left-most image shows
the location of the POA in a coronal brain section. Representative confocal images show
Fos expression (green) in the POA of a group-housed female (left) and a single-housed
female (right) following same-sex social interactions. Blue, Neurotrace. (F) Quantification of
Fos-positive neurons is shown for the POA (left), the VMH (middle), and the caudolateral
PAG (right) for group-housed and single-housed females. Open bars show data from
females that did not engage in social interactions with novel females (baseline), and closed
bars show data from females following social interactions with novel females (interaction).
(G) Total time spent in resident-initiated interaction is plotted for 17-day single-housed
(maroon) and re-group-housed females (teal) during same-sex interactions that occurred
prior to isolation (day 0), following 3 days of isolation (day 3), and on the test day (day 17).
(H) Same as (G), for total resident-initiated mounting time. (I) Same as (G), for total USVs.
(J) Quantification of Fos-positive POA neurons is shown for 17-day single-housed females
(maroon) and re-group-housed females (teal).

Figure 2. Effects of chemogenetic inhibition of POA$_{social}$ neurons on the social
behaviors of single-housed female mice. (A) Experimental timeline and viral strategy to
chemogenetically inhibit the activity of POA$_{social}$ neurons in single-housed females. (B) Total
time spent in resident-initiated social investigation is shown on saline and CNO days for 4
experimental groups: (red symbols) experimental females in which hM4Di is expressed in
POA$_{social}$ neurons; (black symbols) control females in which GFP is expressed in POA$_{social}$
neurons; (brown symbols) control females in which hM4Di is expressed in 'TRAPed' AH
neurons; (gray symbols) control females in which hM4Di is expressed in 'TRAPed' VMH
neurons. (C) Same as (B), for proportion of trials with resident-initiated mounting. (D) Same
as (B), for total USVs. (E) Total movement is plotted for females with hM4Di expressed in
POA$_{social}$ neurons, on saline days vs. CNO days.

Figure 3. Effects of caspase-mediated ablation of POA$_{social}$ neurons on the social
behaviors of single-housed female mice. (A) Experimental timeline and viral strategy for

caspase-mediated ablation of POA$_{social}$ neurons in single-housed females. (B) Total time
spent in resident-initiated social investigation is shown pre- and post-4-OHT treatment for 2
experimental groups: (red symbols) experimental females in which caspase is expressed in
POA$_{social}$ neurons; (black symbols) control females in which GFP is expressed in POA$_{social}$
neurons. (C) Same as (B), for proportion of trials with resident-initiated mounting. (D) Same
as (B), for total USVs.

Figure 4. Effects of optogenetic activation of POA$_{social}$ neurons on the behaviors of
group-housed female mice. (A) Experimental timeline and viral strategy to optogenetically
activate POA$_{social}$ neurons in group-housed females. (B) Mean total USVs produced during
solo sessions shown for pre-laser periods and during laser stimulation periods for
experimental females with ChR2 expressed in POA$_{social}$ neurons (red symbols; N = 9) and for
control females with GFP expressed in POA$_{social}$ neurons (black symbols, N = 6). (C) Same
as (B), for social sessions. (D) Spectrograms are shown from a representative POA$_{social}$-
ChR2 female to illustrate USVs that were elicited through optogenetic activation of POA$_{social}$
neurons in a solo session (top) and a social session (bottom). Blue bars indicate timing of
laser stimulation. (E) Percentage of laser stimulations followed by a bout of social
investigation, plotted for POA$_{social}$-ChR2 and POA$_{social}$-GFP females. (F) Probability of social
investigation aligned with onset of laser stimulation is plotted over time for POA$_{social}$-ChR2
and POA$_{social}$-GFP females. (G) Mean duration of social investigation bouts that overlapped
with laser stimulation vs. bouts that did not overlap with laser stimulation is shown for
POA$_{social}$-ChR2 and POA$_{social}$-GFP females.

Figure 5. Experiments to test whether POA neurons regulate social behaviors of
group-housed females. (A) Experimental timeline and viral strategy to chemogenetically
inhibit POA neurons that were TRAPed following same-sex social interactions in group-
housed females. (B) Total time spent in resident-initiated social investigation is shown on
saline and CNO days. (C) Same as (B), for proportion of trials with resident-initiated
mounting. (D) Same as (B), for total USVs. (E) Experimental timeline and viral strategy to
optogenetically activate POA neurons that were TRAPed following same-sex social
interactions in group-housed females. (F) Mean total USVs produced during solo and social
sessions shown for pre-laser periods and during laser stimulation periods. (G) Percentage of
laser stimulations followed by a bout of social investigation, plotted for GH-TRAPed POA
ChR2 females (N = 6) and POA$_{social}$-GFP females (N = 6; same control group as in Figure 4).
(H) Probability of social investigation following onset of laser stimulation is plotted over time
for GH-TRAPed POA ChR2 and POA$_{social}$-GFP females. (I) Experimental timeline and viral

strategy to chemogenetically silence POA neurons in female mice, first while group-housed

and a second time following single-housing. (J) Total time spent in resident-initiated social

investigation is shown on saline and CNO days for group-housed tests. (K) Same as (J), for

proportion of trials with resident-initiated mounting. (L) Same as (J), for total USVs. (M) Total

time spent in resident-initiated social investigation is shown on saline and CNO days for

single-housed tests. (N) Same as (M), for proportion of trials with resident-initiated mounting.

(O) Same as (M), for total USVs.

Figure 6. Context-dependent differences in POA Fos expression and effects of

chemogenetic inhibition of POA$_{social}$ neurons on the social behaviors of single-housed

male mice. (A) Total time spent in resident-initiated social investigation is shown for group-

housed male residents (teal) and single-housed male residents (maroon) during interactions

with either female visitors (left) or male visitors (right). (B) Same as (A), for proportion of

trials with resident-initiated mounting. (C) Same as (A), for total USVs recorded from pairs

containing group-housed or single-housed male residents. (D) Total number of Fos-positive

POA neurons is shown for group-housed male residents (teal) and single-housed male

residents (maroon) following interactions with female visitors (left) or male visitors (right). (E)

Experimental timeline and viral strategy to chemogenetically inhibit the activity of POA$_{social}$

neurons in single-housed males. (F) Total time spent in resident-initiated social investigation

is shown on saline and CNO days for experimental males in which hM4Di is expressed in

POA$_{social}$ neurons (red symbols, N = 10) and control males in which GFP is expressed in

POA$_{social}$ neurons (black symbols, N = 10). (G) Same as (F), for proportion of trials with

resident-initiated mounting. (H) Same as (F), for total USVs.

Materials and methods

Key Resources

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
c-Fos (9F6) Rabbit mAb	Cell Signaling Technology	2250S
Alexa Fluor 488 goat-anti-rabbit	Invitrogen	A-11008
Neurotrace 435/455 Blue	Thermo Fischer Scientific	CAT#: N21479
Commercial assay or kit		
HCR v3.0	Molecular Instruments	
Bacterial and Virus Strains		
AAV2/1-hSyn-FLEX-hM4Di-mCherry	Addgene	RRID: Addgene_44262
AAV2/1-Ef1alpha-hChR2(h134R)-EYFP-WPRE-HGHpA	Addgene	RRID: Addgene_20298
AAVrg-pgk-Cre	Addgene	RRID: Addgene_24593
pAAV2/5-flex-taCasp3-TEVp	Addgene	RRID: Addgene_45580
pAAV2/5-hSyn-hM4Di-mCherry	Addgene	RRID: Addgene_50475
AAV2/1-pCAG-FLEX-EGFP-WPRE	Addgene	RRID: Addgene_51502
Chemicals, Peptides, and Recombinant Proteins		
Clozapine N-oxide dihydrochloride	Hello Bio	HB6149; E1458-2-1
4-Hydroxytamoxifen	Hello Bio	HB6040
Experimental Models: Organisms/Strains		
TRAP2 Fos$^{tm2.1(icre/ERT2)Luo}$/J	Jackson Labs	IMSR_JAX:030323
Ai14 B6.Cg$Gt(ROSA)26Sor^{tm14(CAG-tdTomato)Hze}$/J	Jackson Labs	IMSR_JAX:007914
C57BL/6J	Jackson Labs	IMSR_JAX:000664
Software and Algorithms		
MATLAB	Mathworks	http://www.mathworks.com RRID: SCR_001622
ZEN	Zeiss	https://www.zeiss.com RRID:SCR_013672
Spike2	CED	http://ced.co.uk RRID:SCR_000903
ImageJ	NIH	https://imagej.net/ij/ RRID:SCR_003070
Behavioral Observation Research Interactive Software	Open Behavior	https://github.com/olivierfriard/BORIS RRID:SCR_021509
R Project for Statistical Computing	R Core Team	http://www.r-project.org/ RRID:SCR_001905
R Studio	Posit	https://posit.co/ RRID:SCR_000432

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by

the lead contact, Katherine Tschida (kat227@cornell.edu).

Experimental models and subject details

Animal statement

All experiments and procedures were conducted according to protocols approved by the

Cornell University Institutional Animal Care and Use Committee (protocol #2020-001).

Animals

TRAP2 (Jackson Laboratories, 030323), Ai14 (Jackson Laboratories, 007914), TRAP2;Ai14,

and C57Bl/6J (Jackson Laboratories, 000664) mice were at least 8 weeks old at the time of

the experiments or surgeries. TRAP2;Ai14 mice were generated by crossing TRAP2 with

Ai14. All mice were kept on a 12:12 reversed light/dark cycle, were housed in ventilated

micro-isolator cages in a controlled environment with regulated temperature and humidity

and were provided with unrestricted access to food and water. A running wheel (Innovive)

was present in all home cages from the time of weaning and was subsequently removed

immediately before initiating the social interaction test. Mouse cages were cleaned weekly,

and experiments were never conducted on cage change days.

TRAP2 female and male subjects were used in all experiments, with the exception of N = 11

TRAP2;Ai14 females subjects used in the neuronal ablation experiments (Fig. 3, S3), N = 4

Ai14 females used in anatomical experiments to examine overlap between PAG-projecting

POA neurons and POA neurons that upregulate Fos following same-sex interactions in

single-housed females (Fig. S4), and N = 7 B6 females used for non-Cre-dependent

chemogenetic silencing of POA (Fig. 5I-O).

Methods details

Social isolation and social interaction tests

Female and male subject mice were either group-housed with same-sex siblings or

separated from their cage mates and individually housed in clean cages for three days prior

to behavioral tests. In the case of group-housed subject mice, siblings were temporarily

removed from the home cage for the duration of the test. The subject animal's home cage

was then placed in a sound-attenuating recording chamber (Med Associates) equipped with

an ultrasonic microphone (Avisoft), an infrared light source (Tendelux), and a webcam

(Logitech, with the infrared filter removed to enable video recording under infrared lighting

conditions). A novel, group-housed visitor mouse (female or male mouse on a C57BL/6

background) was placed in the home cage of the subject mouse, and vocal and non-vocal
behaviors were recorded for 30 minutes. Visitor mice were used across multiple experiments
(< 6 in total), including in interactions with both group-housed and single-housed subject
mice. Visitor females used in male-female interactions were never used for female-female
experiments, but a subset of visitors used in female-female interactions were subsequently
used in male-female experiments.

A separate cohort of female mice was used to investigate the effects on social behavior of
re-group-housing following a period of social isolation. For this cohort of mice, social
interaction tests with novel, group-housed female visitors were conducted at three
timepoints: (1) on day 0, when subject females were still group-housed; (2) On day 3, after
being single-housed for 3 days; (3) on day 17, after a randomly selected subset of subject
females were re-group-housed with their siblings for two weeks, and the remaining female
subjects remained single-housed for two weeks.

USV recording and detection

USVs were recorded with an ultrasonic microphone (Avisoft, CMPA/CM16), amplified
(Presonus TubePreV2), and digitized at 250 kHz (Avisoft UltrasoundGate 166H or CED
Power 1401). USVs were detected with custom MATLAB codes (Tschida et al., 2019) using
the following parameters (mean frequency > 45 kHz; spectral purity > 0.3; spectral
discontinuity < 1.00; minimum USV duration = 5 ms; minimum inter-syllable interval = 30
ms).

Analyses of non-vocal social behaviors

Trained observers used BORIS software (v.8.13; Friard and Gamba, 2016) to score the
following non-vocal behaviors: resident-initiated social investigation and resident-initiated
mounting. Social investigation included sniffing and following. Resident-initiated mounting of
the visitor typically occurred following a period of resident-initiated social investigation, with
the resident mouse positioning its forelimbs on top of the body of the visitor, sometimes with
pelvic thrusts and sometimes without. Neither visitor-initiated mounting nor fighting were
observed in our dataset.

In some trials, total movement was estimated using a custom MATLAB code that allows the
user to mark the position of a mouse in every 30^{th} frame (i.e., once per second). Total
movement was then calculated as the sum of changes in position across pairs of marked
frames.

Fos immunohistochemistry

Two hours following the start of the social interaction test, mice were deeply anesthetized
using isoflurane and then transcardially perfused with phosphate-buffered saline (PBS, pH
7.4), followed by 4% paraformaldehyde (PFA; Sigma-Aldrich, in 0.1 M PBS, pH 7.4). Brains
were subsequently dissected and post-fixed in 4% PFA for 24 hours at 4°C, followed by
immersion in 30% sucrose solution in PBS for 48 hours at 4°C. Afterward, brains were
embedded in frozen section embedding medium (Surgipath, VWR), flash frozen in a dry ice-
ethanol (100%) bath, and then stored at –80°C until sectioning. Sections were cut on a
cryostat (Leica CM1950) to a thickness of 80 μm, washed in PBS (3 x 5 mins at RT),
permeabilized for 2-3 hours in PBS containing 1% Triton X-100 (PBST), and then blocked in
0.3% PBST containing 10% Blocking One (Nacalai USA) for 1 hour at RT on a shaker.
Sections were then incubated for 24 hours at 4°C with primary antibody in blocking solution
(1:1000 rabbit-anti-Fos, Cell Signaling Technologies, 2250S), washed 3 x 30 minutes in
0.3% PBST, then incubated for 24 hours at 4°C with secondary antibody in blocking solution
(1:1000, Alexa Fluor 488 goat-anti-rabbit, Invitrogen, plus 1:500 NeuroTrace, Invitrogen)
Finally, sections were washed for 2 x 10 minutes in 0.3% PBST, followed by washing for 2 x
10 minutes in PBS. After mounting on slides, sections were dried and coverslipped with
Fluromount G (Southern Biotech). Slides were imaged with a 10x objective on a Zeiss 900
laser scanning confocal microscope, and Fos-positive neurons within regions of interest
were counted manually by trained observers.

Floating section two-color in situ hybridization

In situ hybridization was conducted using hybridization chain reaction (HCR v3.0, Molecular
Instruments). Ten minutes after the completion of the 30-minute social interaction tests, mice
underwent transcardial perfusion with RNase-free PBS (DEPC-treated), followed by 4%
PFA. Dissected brain samples were post-fixed overnight in 4% PFA at 4°C, cryoprotected in
a 30% sucrose solution in DEPC-PBS at 4°C for 48 hours, flash frozen in section embedding
medium, and stored at –80°C until sectioning. 40 μm-thick coronal floating sections were
collected into sterile 24-well plates in DEPC-PBS. These sections were briefly fixed once
again for 5 minutes in 4% PFA and subsequently immersed in 70% EtOH in DEPC-PBS
overnight. Sections were then rinsed in DEPC-PBS, incubated for 45 minutes in 5% SDS in
DEPC-PBS, followed by a series of rinses and incubations: 2x SSCT, pre-incubation in HCR
hybridization buffer at 37°C, and incubation in HCR hybridization buffer containing RNA
probes (VGAT and Fos) overnight at 37°C. Sections were then rinsed 4 x 15 minutes at
37°C in HCR probe wash buffer, rinsed in 2X SSCT, pre-incubated in HCR amplification
buffer, and then incubated in HCR amplification buffer containing HCR amplifiers at RT for
approximately 48 hours. On the final day, sections were rinsed in 2x SSCT, counterstained
with DAPI (Thermo Fisher, 1:5000), rinsed again with 2x SSCT, mounted on slides, and

coverslipped with Fluoromount-G (Southern Biotech). After drying, slides were imaged with a 10x or 20x objective on a Zeiss 900 laser scanning microscope. Neurons were scored from three sections of tissue from the POA from each mouse, and the absence of presence of staining for different probes was quantified manually by trained scorers.

Viruses

The following viruses and injection volumes were used: AAV2/1-hSyn-FLEX-hM4Di-mCherry (Addgene #44262, 200 nL), AAV2/1-CAG-FLEX-EGFP-WPRE (Addgene #51502, 200 nL), AAV2/5-Ef1alpha-FLEX-taCasp3-TEVp (Addgene #45580, 200 nL), AAV2/1-Ef1alpha-hChR2(h134R)-EYFP-WPRE-HGHpA (Addgene #20298, 200 nL), AAVrg-pgk-Cre (Addgene #24593, 200 nL), and AAV2/5-hSyn-hM4Di-mCherry (Addgene #50475, 200 nL). The final injection coordinates were as follows: POA, AP = -0.1 mm, ML = 0.6 mm, DV = 5.1 mm; AH, AP = -0.7 mm, ML = 0.6 mm, DV = 5.1 mm; VMH, AP = -1.5 mm, ML = 0.7 mm, DV = 5.4 mm; PAG, AP = -4.7 mm, ML = 0.6 mm, DV = 1.75 mm. Viruses were pressure-injected using a pulled glass pipettes mounted in a programmable nanoliter injector (Nanoject III, Drummond) at a rate of 15 nL every 60 s.

Stereotaxic Surgery

Mice were anesthetized using isoflurane (2.5% for induction, then 1.5 - 2.5% for maintenance) and then securely positioned in a stereotaxic apparatus (Angle Two, Leica). A midline incision in the scalp was made to expose the skull, and small craniotomies were created dorsal to each injection site using a surgical drill. Viral injection pipettes were left in place for a minimum of 10 minutes before and after viral injections to minimize backflow upon pipette withdrawal from the brain. Surgical sutures (LOOK 774B, Fisher Scientific) and tissue adhesive (3M) were used to close the incision.

For optogenetic activation experiments, an optogenetic ferrule (RWD Fiber Optic Cannula, Ø1.25 mm Ceramic Ferrule, 200 µm Core, 0.22 NA, L = 7 mm) was implanted approximately 250 µm above the viral injection site immediately following the viral injection and was secured in place with Metabond (Parkell).

TRAP activity-dependent labeling

Solutions of 4-hydroxytamoxifen (4-OHT, HelloBio, HB6040) were prepared by dissolving 4-OHT powder at 20 mg/mL in ethanol by shaking at 37°C, and aliquots (75 uL) were then stored at -20°C. Before use, 4-OHT was redissolved in ethanol by shaking at 37°C and filtered corn oil was added (150 uL). Ethanol was then evaporated by vacuum under

centrifugation to give a final concentration of 10 mg/mL, and the 4-OHT solution was used

on the same day it was prepared.

To express viral transgenes in recently active neurons, we used the Targeted

Recombination in Active Populations (TRAP2) strategy. Three weeks following viral

injection, TRAP2 and TRAP2;Ai14 mice were single-housed for 3 days and then given 30-

minute social encounters (as described above). Following the social encounter, subject mice

received I.P. injections of 4-OHT (150 uL of 10 mg/mL 4-OHT in filtered corn oil) to enable

expression of viral transgenes in recently active neurons. To minimize neural activity

triggered by stimuli outside of the social interaction test, all subject animals were individually

housed for an additional 24 hours following 4-OHT treatment before being re-group-housed

with their same-sex siblings. In some control experiments, mice remained group-housed

prior to the 30-minute social encounter and 4-OHT treatment (Fig. 5A-H) or were single-

housed but were not given a social encounter prior to 4-OHT treatment (Fig. S2D-G).

Chemogenetic inhibition

To reversibly reduce neuronal activity, TRAP2 female mice received bilateral injections of an

Cre-dependent inhibitory DREADDs virus into the hypothalamus (AAV2/1-hSyn-FLEX-

hM4Di-mCherry; injected into the POA, AH, or VMH) as described above. TRAP2 male mice

received the same viral injections into the POA only. Three weeks later, mice were single-

housed for 3 days and then were subsequently given a 30-minute social encounter in their

home cage with a novel, group-housed female visitor. Subject mice then received an I.P.

injection of 4-OHT to enable expression of hM4Di in activity-defined populations of

hypothalamic neurons. Following the TRAPing session, subjects remained single-housed for

an additional 24 hours. Subjects were then re-group-housed with same-sex siblings for two

weeks and were single-housed a second time prior to behavioral testing. On the first day of

testing, subject mice received an I.P. injection of either sterile saline (as a control) or

clozapine-n-oxide (CNO, 4 mg/kg, Hello Bio HB6149; to inhibit neurons expressing hM4Di)

30 minutes prior to a social interaction test. Three days later, mice that previously were

treated with saline received an I.P. injection of CNO, and mice that were previously treated

with CNO received an I.P. injection of saline, 30 minutes prior to another social interaction.

Rates of USV production and non-vocal social behaviors were compared between saline

and CNO days within animals to assess the effects of neuronal inhibition on social

behaviors. Control mice received unilateral injections into the POA of a Cre-dependent AAV

driving the expression of GFP (AAV2/1-CAG-FLEX-EGFP) and were otherwise treated

identically to experimental animals.

Neuronal ablation

To permanently ablate neurons, TRAP2;Ai14 female mice received bilateral injections of an

AAV2/5-ef1alpha-FLEX-taCasp3-TEVp virus into the POA. Following a three-week recovery

period, these animals were individually housed for three days and subsequently given a 30-

minute social encounter in their home cage with a novel, group-housed female visitor.

Subject mice then received an I.P. injection of 4-OHT to enable expression of caspase in

activity-defined POA neurons. Two weeks later, females were single-housed for three days

and then given a second 30-minute social interaction test. Social behaviors of subject

females were compared between the pre-ablation and post-ablation interaction tests to

assess the effects of neuronal ablation.

Optogenetic activation

Female TRAP2 mice received unilateral injections into the POA of AAV-ef1α-FLEX-ChR2

(experimental) or AAV-CAG-FLEX-GFP (control). In the same surgery, an optogenetic

ferrule was implanted approximately 250 μm above the viral injection site. Three weeks later,

females were single-housed for 3 days and then given a 30-minute social interaction with a

novel group-housed, female visitor. Subject females then received an I.P. injection of 4-

OHT. Two weeks later, females were first placed alone in a clean testing chamber for a 5-

minute habituation period after connecting the laser patch cable to the female's optogenetic

ferrule. Optogenetic activation sessions consisted of a 5-minute period in which optogenetic

activation was performed in solo females, followed by a 20-minute period in which activation

was performed as subject females interacted with a novel, group-housed female visitor. The

social session was further divided into three phases: 5 minutes without optogenetic

activation, 10 minutes with optogenetic activation, and 5 minutes without optogenetic

activation. During the middle 10 minutes of the social session, some laser stimuli were

delivered at times when the two females were near to one another (inter-animal distance <

~2 mouse body lengths), and other stimuli were delivered at times when the females were

not in close contact. POA$_{social}$ neurons were optogenetically activated with illumination from a

473 nM laser (10 mW) at 20-50 Hz (10-20 ms pulses, trains lasted 5-10s) Laser stimuli were

driven by computer-controlled voltage pulses (Spike 2 version 10.8, CED).

Anatomical tracing

Female TRAP2 mice used as GFP controls in the chemogenetic inhibition experiments were

subsequently used for anterograde mapping of the axonal projections of POA$_{social}$ neurons.

Three weeks following unilateral injection into the POA of a Cre-dependent AAV driving the

expression of GFP (AAV2/1-CAG-FLEX-EGFP), females were single-housed for 3 days and

subsequently given a 30-minute social interaction test. Subject mice then received an I.P.

injection of 4-OHT. Six weeks later, females were perfused, brains were collected and

sectioned, and a confocal microscope was used to image GFP-positive axon terminals
within coronal brain tissue sections.

To examine the overlap between PAG-projecting POA neurons and Fos expression, Ai14
females first received a unilateral injection into the PAG of an AAV driving the retrograde
expression of Cre-recombinase (AAVrg-pgk-Cre). Two weeks later, these females were
given a 30-minute social interaction test. Ninety minutes after the test, subject females were
perfused, brains were collected, and coronal brain sections containing the POA were
collected for Fos immunohistochemistry as described above. Brain tissue sections were
imaged with a 10x objective on a Zeiss 900 laser scanning confocal microscope, and POA
neurons that were Fos-positive and tdTomato-positive were counted manually by trained
observers.

Statistics

Two-sided statistical comparisons were used in all analyses (alpha = 0.05). The Shapiro-
Wilk test was performed to analyze the normality of each data distribution, and non-
parametric statistical tests were used for comparisons that included non-normally distributed
data. No statistical methods were used to pre-determine sample size. Mice were only
excluded from analysis in cases in which viral injections were not targeted accurately.
Details of the statistical analyses used in this study are included in Table S1.

References

Alger SJ, Riters LV. 2006. Lesions to the medial preoptic nucleus differentially affect singing and nest box-directed behaviors within and outside of the breeding season in European starlings (Sturnus vulgaris). *Behav Neurosci* **120**:1326–1336. doi:10.1037/0735-7044.120.6.1326

Allen WE, DeNardo LA, Chen MZ, Liu CD, Loh KM, Fenno LE, Ramakrishnan C, Deisseroth K, Luo L. 2017. Thirst-associated preoptic neurons encode an aversive motivational drive. *Science* **357**:1149–1155. doi:10.1126/science.aan6747

An D, Chen W, Yu D-Q, Wang S-W, Yu W-Z, Xu H, Wang D-M, Zhao D, Sun Y-P, Wu J-C, Tang Y-Y, Yin S-M. 2017. Effects of social isolation, re-socialization and age on cognitive and aggressive behaviors of Kunming mice and BALB/c mice. *Anim Sci J* **88**:798–806. doi:10.1111/asj.12688

Arrigo BA, Bullock JL. 2008. The psychological effects of solitary confinement on prisoners in supermax units: reviewing what we know and recommending what should change. *Int J Offender Ther Comp Criminol* **52**:622–640. doi:10.1177/0306624X07309720

Bariselli S, Hörnberg H, Prévost-Solié C, Musardo S, Hatstatt-Burklé L, Scheiffele P, Bellone C. 2018. Role of VTA dopamine neurons and neuroligin 3 in sociability traits related to nonfamiliar conspecific interaction. *Nat Commun* **9**:3173. doi:10.1038/s41467-018-05382-3

Baumeister RF, Leary MR. 1995. The need to belong: Desire for interpersonal attachments as a fundamental human motivation. *Psychol Bull* **117**:497–529. doi:10.1037/0033-2909.117.3.497

Bayless DW, Davis CO, Yang R, Wei Y, de Andrade Carvalho VM, Knoedler JR, Yang T, Livingston O, Lomvardas A, Martins GJ, Vicente AM, Ding JB, Luo L, Shah NM. 2023. A neural circuit for male sexual behavior and reward. *Cell* **186**: 3862-3881.e28. doi: 10.1016/j.cell.2023.07.021

Bean NY, Nunez AA, Conner R. 1981. Effects of medial preoptic lesions on male mouse ultrasonic vocalizations and copulatory behavior. *Brain Res Bull* **6**:109–112. doi:10.1016/s0361-9230(81)80033-0

Cacioppo JT, Cacioppo S. 2018. Chapter Three - Loneliness in the Modern Age: An Evolutionary Theory of Loneliness (ETL) In: Olson JM, editor. Advances in Experimental Social Psychology. Academic Press. pp. 127–197. doi:10.1016/bs.aesp.2018.03.003

Cacioppo JT, Hughes ME, Waite LJ, Hawkley LC, Thisted RA. 2006. Loneliness as a specific risk factor for depressive symptoms: cross-sectional and longitudinal analyses. *Psychol Aging* **21**:140–151. doi:10.1037/0882-7974.21.1.140

Check JVP, Perlman D, Malamuth NM. 1985. Loneuness and Aggressive Behaviour. *J Sol Pers Relat* **2**:243–252. doi:10.1177/0265407585023001

Chen J, Markowitz JE, Lilascharoen V, Taylor S, Sheurpukdi P, Keller JA, Jensen JR, Lim BK, Datta SR, Stowers L. 2021. Flexible scaling and persistence of social vocal communication. *Nature* **593**:108–113. doi:10.1038/s41586-021-03403-8

Chevallier C, Kohls G, Troiani V, Brodkin ES, Schultz RT. 2012. The Social Motivation Theory of Autism. *Trends Cogn Sci* **16**:231–239. doi:10.1016/j.tics.2012.02.007

1093
1094 Clements CC, Zoltowski AR, Yankowitz LD, Yerys BE, Schultz RT, Herrington JD. 2018.
1095 Evaluation of the Social Motivation Hypothesis of Autism. *JAMA Psychiatry* **75**:797–808.
1096 doi:10.1001/jamapsychiatry.2018.1100
1097
1098 Dai B, Sun F, Tong X, Ding Y, Kuang A, Osakada T, Li Y, Lin D. 2022. Responses and
1099 functions of dopamine in nucleus accumbens core during social behaviors. *Cell Rep*
1100 **40**:111246. doi:10.1016/j.celrep.2022.111246
1101
1102 DeNardo LA, Liu CD, Allen WE, Adams EL, Friedmann D, Fu L, Guenthner CJ, Tessier-
1103 Lavigne M, Luo L. 2019. Temporal Evolution of Cortical Ensembles Promoting Remote
1104 Memory Retrieval. *Nat Neurosci* **22**:460–469. doi:10.1038/s41593-018-0318-7
1105
1106 Dölen G, Darvishzadeh A, Huang KW, Malenka RC. 2013. Social reward requires
1107 coordinated activity of nucleus accumbens oxytocin and serotonin. *Nature* **501**:179–184.
1108 doi:10.1038/nature12518
1109
1110 Floody OR. 1989. Dissociation of hypothalamic effects on ultrasound production and
1111 copulation. *Physiol Behav* **46**:299–307. doi:10.1016/0031-9384(89)90271-0
1112
1113 Fukumitsu K, Huang AJ, McHugh TJ, Kuroda KO. 2023. Role of Calcr expressing neurons in
1114 the medial amygdala in social contact among females. *Mol Brain* **16**:10. doi:10.1186/s13041-
1115 023-00993-4
1116
1117 Fukumitsu K, Kaneko M, Maruyama T, Yoshihara C, Huang AJ, McHugh TJ, Itohara S,
1118 Tanaka M, Kuroda KO. 2022. Amylin-Calcitonin receptor signaling in the medial preoptic
1119 area mediates affiliative social behaviors in female mice. *Nat Commun* **13**:709.
1120 doi:10.1038/s41467-022-28131-z
1121
1122 Gao S-C, Wei Y-C, Wang S-R, Xu X-H. 2019. Medial Preoptic Area Modulates Courtship
1123 Ultrasonic Vocalization in Adult Male Mice. *Neurosci Bull* **35**:697–708. doi:10.1007/s12264-
1124 019-00365-w
1125
1126 Green DB, Shackleton TM, Grimsley JMS, Zobay O, Palmer AR, Wallace MN. 2018.
1127 Communication calls produced by electrical stimulation of four structures in the guinea pig
1128 brain. *PLoS One* **13**:e0194091. doi:10.1371/journal.pone.0194091
1129
1130 Gunaydin LA, Grosenick L, Finkelstein JC, Kauvar IV, Fenno LE, Adhikari A, Lammel S,
1131 Mirzabekov JJ, Airan RD, Zalocusky KA, Tye KM, Anikeeva P, Malenka RC, Deisseroth K.
1132 2014. Natural neural projection dynamics underlying social behavior. *Cell* **157**:1535–1551.
1133 doi:10.1016/j.cell.2014.05.017
1134
1135 Hashikawa K, Hashikawa Y, Tremblay R, Zhang J, Feng JE, Sabol A, Piper WT, Lee H,
1136 Rudy B, Lin D. 2017. Esr1+ cells in the ventromedial hypothalamus control female
1137 aggression. *Nat Neurosci* **20**:1580-1590. doi:10.1038/nn.4644
1138
1139 Hossain MM, Sultana A, Purohit N. 2020. Mental health outcomes of quarantine and
1140 isolation for infection prevention: a systematic umbrella review of the global evidence.
1141 *Epidemiol Health* **42**. doi:10.4178/epih.e2020038
1142
1143 House JS, Landis KR, Umberson D. 1988. Social relationships and health. *Science*
1144 **241**:540–545. doi:10.1126/science.3399889
1145

Hu RK, Zuo Y, Ly T, Wang J, Meera P, Wu YE, Hong W. 2021. An amygdala-to-hypothalamus circuit for social reward. *Nat Neurosci* **24**:831–842. doi:10.1038/s41593-021-00828-2

Hung LW, Neuner S, Polepalli JS, Beier KT, Wright M, Walsh JJ, Lewis EM, Luo L, Deisseroth K, Dölen G, Malenka RC. 2017. Gating of social reward by oxytocin in the ventral tegmental area. *Science* **357**:1406–1411. doi:10.1126/science.aan4994

Inagaki TK, Muscatell KA, Moieni M, Dutcher JM, Jevtic I, Irwin MR, Eisenberger NI. 2016. Yearning for connection? Loneliness is associated with increased ventral striatum activity to close others. *Soc Cogn Affect Neurosci* **11**:1096–1101. doi:10.1093/scan/nsv076

Jürgens U. 1994. The role of the periaqueductal grey in vocal behaviour. *Behav Brain Res* **62**:107–117. doi:10.1016/0166-4328(94)90017-5

Karigo T, Kennedy A, Yang B, Liu M, Tai D, Wahle IA, Anderson DJ. 2021. Distinct hypothalamic control of same- and opposite-sex mounting behaviour in mice. *Nature* **589**:258–263. doi:10.1038/s41586-020-2995-0

Killgore WDS, Cloonan SA, Taylor EC, Anlap I, Dailey NS. 2021. Increasing aggression during the COVID-19 lockdowns. *J Affect Disord Rep* **5**:100163. doi:10.1016/j.jadr.2021.100163

Kohl J, Babayan BM, Rubinstein ND, Autry AE, Marin-Rodriguez B, Kapoor V, Miyamishi K, Zweifel LS, Luo L, Uchida N, Dulac C. 2018. Functional circuit architecture underlying parental behaviour. *Nature* **556**:326–331. doi:10.1038/s41586-018-0027-0

Lee CR, Chen A, Tye KM. 2021. The neural circuitry of social homeostasis: Consequences of acute versus chronic social isolation. *Cell* **184**:1500–1516. doi:10.1016/j.cell.2021.02.028

Lee H, Kim D-W, Remedios R, Anthony TE, Chang A, Madisen L, Zeng H, Anderson DJ. 2014. Scalable control of mounting and attack by Esr1+ neurons in the ventromedial hypothalamus. *Nature* **509**:627–632. doi:10.1038/nature13169

Liu D, Rahman M, Johnson A, Tsutsui-Kimura I, Pena N, Talay M, Logeman BL, Finkbeiner S, Choi S, Capo-Battaglia A, Abdus-Saboor I, Ginty DD, Uchida N, Watabe-Uchida M, Dulac C. 2023. A Hypothalamic Circuit Underlying the Dynamic Control of Social Homeostasis. Preprint at bioRxiv. doi:10.1101/2023.05.19.540391

Liu M, Kim D-W, Zeng H, Anderson DJ. 2022. Make war not love: The neural substrate underlying a state-dependent switch in female social behavior. *Neuron* **110**:841-856.e6. doi:10.1016/j.neuron.2021.12.002

Love TM. 2014. Oxytocin, motivation and the role of dopamine. *Pharmacol Biochem Behav* **119**:49–60. doi:10.1016/j.pbb.2013.06.011

Ma X, Jiang D, Jiang W, Wang F, Jia M, Wu J, Hashimoto K, Dang Y, Gao C. 2011. Social isolation-induced aggression potentiates anxiety and depressive-like behavior in male mice subjected to unpredictable chronic mild stress. *PLoS One* **6**:e20955. doi:10.1371/journal.pone.0020955

Ma Y-K, Zeng P-Y, Chu Y-H, Lee C-L, Cheng C-C, Chen C-H, Su Y-S, Lin K-T, Kuo T-H. 2022. Lack of social touch alters anxiety-like and social behaviors in male mice. *Stress* **25**:134–144. doi:10.1080/10253890.2022.2047174

Machimbarrena JM, Álvarez-Bardón A, León-Mejía A, Gutiérrez-Ortega M, Casadiego-
Cabrales A, González-Cabrera J. 2019. Loneliness and Personality Profiles Involved in
Bullying Victimization and Aggressive Behavior. *School Ment Health* **11**:807–818.
doi:10.1007/s12310-019-09328-y

Matsumoto K, Pinna G, Puia G, Guidotti A, Costa E. 2005. Social isolation stress-induced
aggression in mice: a model to study the pharmacology of neurosteroidogenesis. *Stress*
8:85–93. doi:10.1080/10253890500159022

Matthews GA, Nieh EH, Vander Weele CM, Halbert SA, Pradhan RV, Yosafat AS, Glober
GF, Izadmehr EM, Thomas RE, Lacy GD, Wildes CP, Ungless MA, Tye KM. 2016. Dorsal
Raphe Dopamine Neurons Represent the Experience of Social Isolation. *Cell* **164**:617–631.
doi:10.1016/j.cell.2015.12.040

McHenry JA, Otis JM, Rossi MA, Robinson JE, Kosyk O, Miller NW, McElligott ZA, Budygin
EA, Rubinow DR, Stuber GD. 2017. Hormonal gain control of a medial preoptic area social
reward circuit. *Nat Neurosci* **20**:449–458. doi:10.1038/nn.4487

McWain MA, Pace RL, Nalan PA, Lester DB. 2022. Age-dependent effects of social isolation
on mesolimbic dopamine release. *Exp Brain Res* **240**:2803–2815. doi:10.1007/s00221-022-
06449-w

Mears DP, Bales WD. 2009. Supermax Incarceration and Recidivism. *Criminology* **47**:1131–
1166. doi:10.1111/j.1745-9125.2009.00171.x

Melis MR, Sanna F, Argiolas A. 2022. Dopamine, Erectile Function and Male Sexual
Behavior from the Past to the Present: A Review. *Brain Sci* **12**:826.
doi:10.3390/brainsci12070826

Merari A, Ginton A. 1975. Characteristics of exaggerated sexual behavior induced by
electrical stimulation of the medial preoptic area in male rats. *Brain Res* **86**:97–108.
doi:10.1016/0006-8993(75)90641-1

Michael V, Goffinet J, Pearson J, Wang F, Tschida K, Mooney R. 2020. Circuit and synaptic
organization of forebrain-to-midbrain pathways that promote and suppress vocalization. *Elife*
9:e63493. doi:10.7554/eLife.63493

Musardo S, Contestabile A, Knoop M, Baud O, Bellone C. 2022. Oxytocin neurons mediate
the effect of social isolation via the VTA circuits. *eLife* **11**:e73421. doi:10.7554/eLife.73421

Niesink RJ, van Ree JM. 1982. Short-term isolation increases social interactions of male
rats: a parametric analysis. *Physiol Behav* **29**:819–825. doi:10.1016/0031-9384(82)90331-6

Otchy TM, Wolff SBE, Rhee JY, Pehlevan C, Kawai R, Kempf A, Gobes SMH, Ölveczky BP.
2015. Acute off-target effects of neural circuit manipulations. *Nature* **528**:358–363.
doi:10.1038/nature16442

Panksepp J, Beatty WW. 1980. Social deprivation and play in rats. *Behav Neural Biol*
30:197–206. doi:10.1016/s0163-1047(80)91077-8

Reid JA, Chenneville T, Gardy SM, Baglivio MT. 2022. An Exploratory Study of COVID-19's
Impact on Psychological Distress and Antisocial Behavior Among Justice-Involved Youth.
Crime Delinquency **68**:1271–1291. doi:10.1177/00111287211054729

Resendez SL, Namboodiri VMK, Otis JM, Eckman LEH, Rodriguez-Romaguera J, Ung RL, Basiri ML, Kosyk O, Rossi MA, Dichter GS, Stuber GD. 2020. Social Stimuli Induce Activation of Oxytocin Neurons Within the Paraventricular Nucleus of the Hypothalamus to Promote Social Behavior in Male Mice. *J Neurosci* **40**:2282–2295. doi:10.1523/JNEUROSCI.1515-18.2020

Riters LV. 2012. The role of motivation and reward neural systems in vocal communication in songbirds. *Front Neuroendocrinol* **33**:194–209. doi:10.1016/j.yfrne.2012.04.002

Riters LV, Ball GF. 1999. Lesions to the medial preoptic area affect singing in the male European starling (Sturnus vulgaris). *Horm Behav* **36**:276–286. doi:10.1006/hbeh.1999.1549

Riters LV, Eens M, Pinxten R, Duffy DL, Balthazart J, Ball GF. 2000. Seasonal changes in courtship song and the medial preoptic area in male European starlings (Sturnus vulgaris). *Horm Behav* **38**:250–261. doi:10.1006/hbeh.2000.1623

Robinson DL, Zitzman DL, Smith KJ, Spear LP. 2011. Fast dopamine release events in the nucleus accumbens of early adolescent rats. *Neuroscience* **176**:296–307. doi:10.1016/j.neuroscience.2010.12.016

Solié C, Girard B, Righetti B, Tapparel M, Bellone C. 2022. VTA dopamine neuron activity encodes social interaction and promotes reinforcement learning through social prediction error. *Nat Neurosci* **25**:86–97. doi:10.1038/s41593-021-00972-9

Tan T, Wang W, Liu T, Zhong P, Conrow-Graham M, Tian X, Yan Z. 2021. Neural circuits and activity dynamics underlying sex-specific effects of chronic social isolation stress. *Cell Rep* **34**:108874. doi:10.1016/j.celrep.2021.108874

Tang Y, Chen Z, Tao H, Li C, Zhang X, Tang A, Liu Y. 2014. Oxytocin activation of neurons in ventral tegmental area and interfascicular nucleus of mouse midbrain. *Neuropharmacology* **77**:277–284. doi:10.1016/j.neuropharm.2013.10.004

Tomova L, Wang KL, Thompson T, Matthews GA, Takahashi A, Tye KM, Saxe R. 2020. Acute social isolation evokes midbrain craving responses similar to hunger. *Nat Neurosci* **23**:1597–1605. doi:10.1038/s41593-020-00742-z

Toth M, Mikics E, Tulogdi A, Aliczki M, Haller J. 2011. Post-weaning social isolation induces abnormal forms of aggression in conjunction with increased glucocorticoid and autonomic stress responses. *Horm Behav* **60**:28–36. doi:10.1016/j.yhbeh.2011.02.003

Tschida K, Michael V, Takatoh J, Han B-X, Zhao S, Sakurai K, Mooney R, Wang F. 2019. A Specialized Neural Circuit Gates Social Vocalizations in the Mouse. *Neuron* **103**:459-472.e4. doi:10.1016/j.neuron.2019.05.025

Valzelli L. 1973. The "isolation syndrome" in mice. *Psychopharmacologia* **31**:305–320. doi:10.1007/BF00421275

Walum H, Young LJ. 2018. The neural mechanisms and circuitry of the pair bond. *Nat Rev Neurosci* **19**:643–654. doi:10.1038/s41583-018-0072-6

Warren MR, Clein RS, Spurrier MS, Roth ED, Neunuebel JP. 2020. Ultrashort-range, high-frequency communication by female mice shapes social interactions. *Sci Rep* **10**:2637. doi:10.1038/s41598-020-59418-0

Wei Y-C, Wang S-R, Jiao Z-L, Zhang W, Lin J-K, Li X-Y, Li S-S, Zhang X, Xu X-H. 2018.
Medial preoptic area in mice is capable of mediating sexually dimorphic behaviors
regardless of gender. *Nat Commun* **9**:279. doi:10.1038/s41467-017-02648-0

Weiss IC, Pryce CR, Jongen-Rêlo AL, Nanz-Bahr NI, Feldon J. 2004. Effect of social
isolation on stress-related behavioural and neuroendocrine state in the rat. *Behav Brain Res*
152:279–295. doi:10.1016/j.bbr.2003.10.015

Wiberg GS, Grice HC. 1963. Long-term isolation stress in rats. *Science* **142**:507.
doi:10.1126/science.142.3591.507

Wu YE, Dang J, Kingsbury L, Zhang M, Sun F, Hu RK, Hong W. 2021. Neural control of
affiliative touch in prosocial interaction. *Nature* **599**:262–267. doi:10.1038/s41586-021-
03962-w

Xiao L, Priest MF, Nasenbeny J, Lu T, Kozorovitskiy Y. 2017. Biased Oxytocinergic
Modulation of Midbrain Dopamine Systems. *Neuron* **95**:368-384.e5.
doi:10.1016/j.neuron.2017.06.003

Yang B, Karigo T, Anderson DJ. 2023. Transformations of neural representations in a social
behavior network. *Nature* 608(7924):741-749. doi: 10.1038/s41586-022-05057-6

Yorgason JT, Calipari ES, Ferris MJ, Karkhanis AN, Fordahl SC, Weiner JL, Jones SR.
2016. Social isolation rearing increases dopamine uptake and psychostimulant potency in
the striatum. *Neuropharmacology* **101**:471–479. doi:10.1016/j.neuropharm.2015.10.025

Zelikowsky M, Hui M, Karigo T, Choe A, Yang B, Blanco MR, Beadle K, Gradinaru V,
Deverman BE, Anderson DJ. 2018. The Neuropeptide Tac2 Controls a Distributed Brain
State Induced by Chronic Social Isolation Stress. *Cell* **173**:1265-1279.e19.
doi:10.1016/j.cell.2018.03.037

Zhao X, Ziobro P, Pranic NM, Chu S, Rabinovich S, Chan W, Zhao J, Kornbrek C, He Z,
Tschida KA. 2021. Sex- and context-dependent effects of acute isolation on vocal and non-
vocal social behaviors in mice. *PLoS One* **16**:e0255640. doi:10.1371/journal.pone.0255640

Figure 1

Figure 2

Figure 3

A

B

Resident-initiated investigation (s)

Post > Pre, p < 0.01

POA$_{social}$ caspase   POA$_{social}$ GFP

C

Proportion of trials with resident-initiated mounting

| Pre | Post | Pre | Post |
| 13/15 | 4/15 | 8/13 | 6/13 |

*

n.s.

POA$_{social}$ caspase   POA$_{social}$ GFP

D

Total USVs

n.s.   n.s.

POA$_{social}$ caspase   POA$_{social}$ GFP

Figure 4

Figure 5.

Figure 6.