Parvalbumin interneuron mGlu$_5$ receptors govern sex differences in prefrontal cortex physiology and binge drinking

Carly B. Fabian[a,b,c], Nilah I. Jordan[c], Lily G. Carley[a,b,c], Shannon M. Thompson[a,b,c], Marianne L. Seney[a,b,c], and Max E. Joffe[a,b,c,*]

   a. Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, 15219, USA
   b. Translational Neuroscience Program, University of Pittsburgh, Pittsburgh, PA
   c. Center for Neuroscience University of Pittsburgh, Pittsburgh, PA

*Correspondence to:
Max E. Joffe, Ph.D.
Assistant Professor
University of Pittsburgh
219 Bridgeside Point II
450 Technology Dr
Pittsburgh, PA 15219
E-Mail: joffeme@upmc.edu
Phone: 412-383-6028
Twitter: @mejoffe

Short title: Group 1 mGlu receptors mediate sex differences in PFC PV-INs

Key words: prefrontal cortex, alcohol, G protein-coupled receptor (GPCR), cognition, GABA, addiction

Article Type: Research
Abstract: 232 words
Article: 3968 words (limit 4000)
Figures: 5
Tables: 0
Supplemental Information: 3 figures
References: 97

Abstract

Despite established sex differences in the prevalence and presentation of psychiatric disorders, little is known about the cellular and synaptic mechanisms that guide these differences under basal conditions. Proper function of the prefrontal cortex (PFC) is essential for the top-down regulation of motivated behaviors. Activity of the PFC is tightly controlled by parvalbumin-expressing interneurons (PV-INs), a key subpopulation of fast-spiking GABAergic cells that regulate cortical excitability through direct innervations onto the perisomatic regions of nearby pyramidal cells. Recent rodent studies have identified notable sex differences in PV-IN activity and adaptations to experiences such as binge drinking. Here, we investigated the cellular and molecular mechanisms that underlie sex-specific regulation of PFC PV-IN function. Using whole-cell patch clamp electrophysiology and selective pharmacology, we report that PV-INs from female mice are more excitable than those from males. Moreover, we find that $mGlu_1$ and $mGlu_5$ metabotropic glutamate receptors regulate cell excitability, excitatory drive, and endocannabinoid signaling at PFC PV-INs in a sex-dependent manner. Genetic deletion of $mGlu_5$ receptors from PV-expressing cells abrogates all sex differences observed in PV-IN membrane and synaptic physiology. Lastly, we report that female, but not male, PV-$mGlu_5^{-/-}$ mice exhibit decreased voluntary drinking on an intermittent access schedule, which could be related to changes in ethanol's stimulant properties. Importantly, these studies identify $mGlu_1$ and $mGlu_5$ receptors as candidate signaling molecules involved in sex differences in PV-IN activity and behaviors relevant for alcohol use.

Introduction

Sex differences contribute to the incidence, presentation, and severity of several psychiatric disorders [1,2]. For instance, women with major depressive disorder (MDD) experience greater symptom severity [3,4], men have an earlier age of onset of schizophrenia [5-7], and women display a faster progression through disease milestones of alcohol use disorder (AUD) [8,9]. Together, the broad sex differences in psychiatric diseases motivate continued preclinical research to identify the cells and plasticity mechanisms that mediate sex differences in brain function.

The prefrontal cortex (PFC) provides top-down regulation of motivated behaviors and displays sex differences in form and function [10-14]. Most PFC neurons are glutamatergic pyramidal cells that send, receive, and integrate information to and from various subcortical and cortical structures [15,16]. Pyramidal cell activity is tightly regulated by a small number of inhibitory neurons that transmit γ-aminobutyric acid (GABA). Parvalbumin-expressing inhibitory neurons (PV-INs) form a key subpopulation of these GABAergic cells which synapse onto perisomatic regions of pyramidal neurons and display fast-spiking firing properties [17-19]. Dysregulation of PFC PV-IN activity has been implicated in the etiology of many psychiatric disorders [20-23]. A growing body of literature in rodent models has also identified notable sex differences in PV-IN activity and adaptations to experiences. Indeed, male and female mice display sex differences in developmental trajectory of PFC PV expression [24], stress sex-specifically alters PFC PV-IN function [25-27], and PFC PV-INs undergo sex-dependent adaptations following four-weeks binge drinking [28]. Taken together, these studies suggest that sex differences in molecular mechanisms governing synaptic plasticity and intrinsic properties on PV-INs are critical in mediating experience-dependent adaptations to PFC function.

The group 1 metabotropic glutamate (mGlu) receptors, $mGlu_1$ and $mGlu_5$, are expressed on PFC PV-INs [29-32] and regulate cellular processes related to membrane physiology, synaptic strength, and endocannabinoid plasticity [33-35]. In addition, there are sex differences

in signaling and function of $mGlu_1$ and $mGlu_5$ receptors [36-38]. Here, we tested the hypothesis that $mGlu_1$ and $mGlu_5$ receptors mediate sex differences in PFC PV-IN function. Using genetically engineered mice, whole-cell patch clamp electrophysiology, and selective pharmacology, we examined sex differences in PV-IN function and regulation by $mGlu_1$ and $mGlu_5$ receptors. We then assessed a casual role for mGlu receptors in regulating PFC PV-IN function through mice with cell type-specific deletion of $mGlu_5$ receptors. The present results support the hypothesis that PV-IN $mGlu_5$ receptor signaling is critical for sex differences in PFC physiology and drinking.

Materials and Methods

Mice

All mice were maintained on a 12:12 light-dark cycle (lights on at 6:00 AM) with food and water *ad libitum*. Mice were bred at Charles River Laboratory (Wilmington, MA) and delivered to the University of Pittsburgh at 6-8 weeks of age. All mice were on a congenic C57BL/6J background and experiments began in adulthood (>10 weeks). Sex determination was limited to an assessment of external genitalia. Mice expressing tdTomato in PV-INs were bred by crossing female homozygous PV-Cre mice [39] (Jackson Laboratories, 017320) with male homozygous Rosa26-loxP-STOP-loxP-CAG-tdTomato "Ai9" mice [40] (Jackson Laboratories, Stock No: 007909). $PV-mGlu_5^{-/-}$ mice were generated by crossing PV-Cre and Ai9 mice with those harboring a floxed *Grm5* locus [31,41] (Jackson Laboratories, 028626). All experiments were approved by the Institutional Animal Care and Use Committee of the University of Pittsburgh and conducted in accordance with the NIH Guidelines.

Brain slice preparation

Mice (N=65) were decapitated under isoflurane anesthesia; brains were immediately dissected and placed in ice-cold N-methyl-D-glucamine (NMDG) cutting solution, containing (in mM): 93 NMDG, 20 HEPES, 2.5 KCl, 0.5 $CaCl_2$, 10 $MgCl_2$, 1.2 NaH_2PO_4, 25 glucose, 5 Na-ascorbate, and 3 Na-pyruvate. Acute coronal slices (300-µM) of the PL-PFC were prepared in room-temperature NMDG solution. Slices recovered for 10 minutes in warm (32-35°C) NMDG solution and were then transferred to artificial cerebral spinal fluid (ACSF), containing (in mM):119 NaCl, 2.5 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 1 NaH_2PO_4, 11 glucose, and 26 $NaHCO_3$ for at least 1 hour at room temperature before recordings. All solutions were oxygenated (95% O_2 / 5% CO_2).

Ex vivo whole-cell patch-clamp electrophysiology

Data were acquired using a Multiclamp 700B amplifier and pCLAMP 11 software (Axon Instruments, Molecular Devices, Sunnyvale, CA, USA). Slices were transferred to the recording chamber and perfused at 2mL/min with heated ACSF (30-32°C). For all whole-cell patch-clamp recordings, cells were patched with a borosilicate glass pipette pulled to 3-5 MΩ with a horizontal electrode puller (P-1000, Stutter Instruments). Pipettes were filled with a potassium-based internal solution, containing (in mM): 125 K-gluconate, 4 NaCl, 10 HEPES, 0.1 EGTA, 4 MgATP, 0.3 Na_2GTP, 10 Tris-phosphocreatine. PV-INs in layer 5 of the PL-PFC were identified by tdTomato fluorescence and cell identity was confirmed via characteristic fast-spiking phenotype. After gaining whole-cell access, PV-INs were dialyzed for 5 minutes under voltage-clamp at -80mV. Membrane properties were then assessed under current-clamp configuration as previously described [42]. A series of hyperpolarizing and depolarizing current injections (1s, starting at -150pA and ending at +500pA, in increments of 25pA) were applied to patched cells. Resting membrane potential (V_m) was determined as the average baseline membrane potential prior to current injection. Membrane resistance (R_m) was calculated as the slope of the hyperpolarizing membrane potential over three current injections [-75, -50, -25pA]. Rheobase

was determined as the minimal current injection needed to elicit firing (>1 spike). Number of spikes per current step were calculated using threshold-based event detection.

Cells were then returned to voltage-clamp configuration. Spontaneous excitatory postsynaptic currents (sEPSCs) were electrically isolated at -80mV, near the reversal potential for chloride [28]. Following a 5-minute baseline, 5µM and 50µM DHPG were sequentially added to the bath for a duration of 5-minutes each. For experiments using LY367385 (100µM) or MTEP (3µM), either drug was applied prior to baseline recordings and remained in the bath for the entire experiment. sEPSCs were identified using a template specific for PV-INs. Change in holding current and sEPSC frequency were calculated by comparing the average values over the last 2 minutes of drug application to the last 2 minutes of baseline.

In separate PV-INs, endocannabinoid-mediated short-term plasticity was assessed via depolarization-induced suppression of excitation (DSE) [43,44]. Bipolar nichrome wire stimulating electrodes were positioned in layer 2/3 medial to the recording electrode. EPSCs were evoked with a single pulse (10-50µA) at 0.2Hz using stimulation parameters to elicit EPSCs between 100-200pA. After establishing a stable 1-minute baseline, DSE was elicited by depolarizing patched PV-INs to +20mV for 10 seconds without stimulation, followed by an immediate return to -80mV and electrical stimulation. DSE was repeated 3-4 times per cell under basal conditions. DHPG (5µM) was then applied for 5 minutes and the DSE protocol was repeated. Changes in the peak amplitude of EPSCs were normalized to baseline. Data for all electrophysiology studies was processed using ClampFit (Molecular Devices).

Intermittent access (IA) ethanol

IA ethanol was performed as described [42,45]. For three 24-hour periods per week, ethanol was provided in home cages, with 24-48 hours between each exposure. Ethanol was diluted from undenatured ethyl alcohol (95%, Decon Labs, Incorporated) and was delivered by modified conical tubes with rubber stoppers open tip tubes Ethanol was generally provided and removed

3-4 hours prior to start of the dark cycle. After a ramp during the first three sessions [3, 6, 10%], the ethanol concentration was set at 20% ethanol for the remainder of the study.

Locomotor response to ethanol

We assessed the locomotor response to ethanol in a cohort of mice without previous exposure to ethanol. Mice received saline vehicle (10 µL/kg, *i.p.*) or ethanol (0.5, 1, or 2 g/kg) immediately before being placed in an open field chamber for 20 minutes. Over 4 consecutive days, all mice received all doses in a counterbalanced Latin-square design. Location and distance traveled were determined by video-based tracking (ANY-maze, Stoelting).

Drugs

Drugs were purchased from HelloBio (Princeton, NJ). Concentrations were based on previous studies [29,32,46,47]. Drugs were formulated, aliquoted, and frozen at 1000X, then thawed and diluted in ACSF on the day of recording. LY367385 [(*S*)-(+)-α-Amino-4-carboxy-2-methylbenzeneaceticacid] stock was made in 0.1M NaOH. MTEP [3-((2-Methyl-1,3-thiazol-4-yl)ethynyl)pyridine hydrochloride] and DHPG [(*RS*)-3,5-Dihydroxyphenylglycine] stocks were made in water.

Statistics

Data are presented as mean±SEM or box plots. Statistical analyses were performed using GraphPad Prism. For electrophysiological experiments, *n* represents the number of cells. For all analyses, $p<0.05$ was considered significant and $p<0.07$ was labeled trending. Data were analyzed using two-tailed Student's t-test. Two-way or three-way ANOVA, or mixed-effects analysis, with Bonferonni post hoc comparisons were used for analyses requiring multiple comparisons. Statistical analyses are reported in the figure legends.

Results

PL-PFC PV-INs from female are mice are more excitable than PV-INs from males

To assess intrinsic physiology of PFC PV-INs, we made whole-cell patch-clamp recordings from genetically engineered mice expressing tdTomato in PV-positive cells [32,42,48]. All (n=75/75) tdTomato-expressing cells from layer 5 of the PL-PFC displayed characteristic fast-spiking interneuron properties [49,50], including: high frequency firing, low input resistance, and minimal spike-frequency adaptation (Figure 1A). In general, PV-INs from male and female mice displayed comparable basal membrane properties, without significant differences in resting potential, membrane resistance, or medium afterhyperpolarization (Figure S1A-C). However, in response to positive current injections, cells from female mice exhibited greater spike-firing compared to cells from males (Figure 1B) and fired action potentials in response to a lower threshold current (Figure 1C). Together, these data indicate that while PV-INs exhibit similar basal membrane properties between sexes, PV-INs from female mice are more excitable than those from males.

mGlu$_1$ and mGlu$_5$ receptors depolarize PFC PV-IN in a sex-dependent manner

We and others previously demonstrated that mGlu$_1$ and mGlu$_5$ receptors modulate PV-INs in frontal cortex [29-31], but most of these studies were limited to males. We therefore aimed to determine whether mGlu$_1$ and mGlu$_5$ receptors modulate PV-IN function in a similar manner in males and females. We examined sensitivity to mGlu$_1$ and mGlu$_5$ receptor stimulation using a threshold concentration of the non-selective mGlu$_{1/5}$ agonist DHPG (5μM) alone, in the presence of the selective mGlu$_5$ NAM MTEP (3μM), or in the presence of the selective mGlu$_1$ antagonist LY367385 (100μM) (Figure 1D). Threshold DHPG induced modest depolarizing currents in PV-INs of both sexes; however, PV-INs from male mice displayed a trend towards greater depolarizing changes in holding current compared to females (Figure 1E), with no effect of either antagonist. We next assessed a high concentration of DHPG (50μM), which induced

greater depolarizing current relative to the threshold concentration in both sexes (Figure 1F). Depolarizing currents in response to 50µM DHPG were greater in PV-INs from males, and selective pharmacological antagonism revealed additional sex differences in the receptor subtypes involved. In PV-INs from all mice, MTEP attenuated DHPG-induced depolarizing currents. By contrast, LY367385 reduced DHPG currents in PV-INs from male mice but did reach significance in females. Together, these data suggest that PFC PV-IN membrane physiology is modulated by $mGlu_5$ receptors across sexes, but that PV-INs from male mice are more sensitive to $mGlu_1$ receptor activation.

$mGlu_1$ and $mGlu_5$ regulate synaptic strength on PFC PV-INs in a sex-dependent manner.

In addition to membrane physiology, Group 1 mGlu receptors can modulate synaptic strength on many cell types, including other types of interneurons in frontal cortex [32,51-54]. We measured AMPA receptor-mediated sEPSCs under voltage-clamp and found no evidence for basal sex differences in excitatory synaptic strength (Figure S1D-F). We next determined whether Group 1 mGlu receptors regulate excitatory transmission. Threshold DHPG (5µM) had no effect on sEPSC frequency (Figure 2A and 2B), but a higher concentration of DHPG (50µM) induced a three-fold increase in sEPSC frequency in PV-INs across sexes (Figure 2C). Further pharmacological studies revealed a sex-specific receptor mechanism underlying this effect: LY367385 blocked enhancement of excitatory drive in PV-INs from male mice, while the effects of DHPG on sEPSC frequency were attenuated by MTEP in PV-INs from females. These findings suggest a sex-specific role of the group 1 mGlu receptors in regulating synaptic strength onto PFC PV-INs, primarily driven by $mGlu_1$ receptor signaling in males and by $mGlu_5$ receptors in females.

$mGlu_5$ receptors facilitate short-term plasticity at PV-INs from male but not female mice

We next sought to examine mechanisms of mGlu receptor synaptic plasticity. In several brain regions, activation of postsynaptic Group 1 mGlu receptors can trigger endocannabinoid release and reductions in neurotransmitter release [55-58]. We elicited endocannabinoid plasticity by depolarization-induced suppression of excitation (DSE) [59,60], a form of short-term depression mediated by CB1 receptor activation and reduced glutamate release probability. Under control conditions, PV-IN EPSC amplitude was reduced by approximately 20% by cell depolarization across sexes (Figure 2D and 2E). We next assessed whether $mGlu_1$ and $mGlu_5$ receptor activation can facilitate DSE by applying the threshold concentration of DHPG (5μM). While no effect was observed in female mice, threshold DHPG enhanced DSE in PV-INs from male mice, which exhibited nearly a 40% reduction in EPSC amplitude (Figure 2E and 2F). Importantly, threshold DHPG did not affect EPSC amplitude in the absence of postsynaptic depolarization (Figure S3), suggesting the facilitation of DSE is a metaplastic change in endocannabinoid signaling, rather than an overall change in glutamate transmission. Based on our previous findings, we predicted that the enhancement of DSE in male mice would be lost in the presence of LY367385. To our surprise, DHPG retained the ability to enhance DSE in LY367385 (Figure 2G and 2I), while MTEP completely blocked DSE potentiation (Figure 2H and 2I). We did not perform experiments using the antagonists in female mice because there was no effect of DHPG alone. Taken together, these findings indicate that $mGlu_5$ receptors can potentiate endocannabinoid signaling in PV-INs from male mice.

Genetic deletion of $mGlu_5$ receptors from PV-INs abrogates sex-differences in membrane and synaptic physiology

We next used a genetic approach to examine $mGlu_5$ receptor function within PV-INs. We bred mice to selectively ablate $mGlu_5$ receptors from PV-expressing cells via Cre-mediated deletion of *Grm5*, as reported previously [31]. PV-$mGlu_5^{-/-}$ mice display postnatal reduction of $mGlu_5$ receptors to ~20% of control levels, with recombination beginning between around P28,

reflecting endogenous PV expression. We first assessed PV-IN membrane properties in PV-mGlu$_5^{-/-}$ mice. In contrast to controls (Figure 1A-1C), we observed no sex differences in PV-IN spike-firing (Figure 3A and 3B) or rheobase (Figure 3C) in PV-mGlu$_5^{-/-}$ mice. Furthermore, we observed no sex differences in any other membrane property or in basal synaptic strength (Figure S2). These data suggest that mGlu$_5$ receptors regulate the development and/or maintenance of sex differences in PV-IN excitability. We next assessed responses to mGlu$_{1/5}$ receptor activation, hypothesizing that DHPG would not depolarize PV-INs in female PV-mGlu$_5^{-/-}$ mice. To our surprise, DHPG induced comparable depolarizing currents in both sexes (Figure 3D-3F), and currents were attenuated by either MTEP or LY367385. These findings suggest that (1) there is residual expression of functional mGlu$_5$ receptors in PV-mGlu$_5^{-/-}$ mice, consistent with previous studies [31], and (2) female PV-mGlu$_5^{-/-}$ mice display unmasked sensitivity to mGlu$_1$ receptor antagonism. We next assessed DHPG potentiation of excitatory drive in PV-mGlu$_5^{-/-}$ mice (Figure 4A). As in controls, 5µM DHPG had little effect on sEPSC frequency (Figure 4B), while 50µM DHPG rapidly increased excitatory drive on PV-INs in both sexes (Figure 4C). In contrast to recordings from control mice, we observed a main effect of mGlu receptor antagonist without an effect of sex or interaction. We performed posthoc tests across sexes which revealed that LY367385, but not MTEP, reduced the ability of DHPG to potentiate sEPSC frequency, again indicating loss of mGlu$_5$ function in female mice and enhanced sensitivity to mGlu$_1$ receptor antagonism. Finally, we performed DSE experiments to assess eCB signaling (Figure 4D-4F). DHPG failed to potentiate DSE in PV-INs of PV-mGlu$_5^{-/-}$ mice. Taken together, all sex differences in PV-IN physiology and mGlu receptor regulation were absent in PV-mGlu$_5^{-/-}$ mice.

Deleting mGlu$_5$ receptors from PV-INs in female mice, but not males, decreases voluntary drinking

mGlu$_5$ receptor signaling regulates several behaviors related to alcohol use [33,61], and previous studies from our group have demonstrated that drinking and ethanol reward learning alter the intrinsic and synaptic properties of PV-INs [28,48]. Based on this literature and our current findings that mGlu$_5$ receptors regulate synaptic strength onto PV-INs in female mice but not males, we hypothesized that mGlu$_5$ receptors in PV-INs regulate binge drinking in a sex-dependent manner. To test this hypothesis, we provided control and PV-mGlu$_5$$^{-/-}$ mice with two-bottle choice IA ethanol over 7 weeks (Figure 5A) as in previous studies [42,45]. As observed by our lab and many others [42,62-64], female mice drank more than male mice overall. Moreover, a sex by genotype interaction emerged, such that female but not male PV-mGlu$_5$$^{-/-}$ mice drank less than their respective controls (Figure 5B). A similar effect was observed with respect to preference for ethanol relative to water (Figure 5C). We next performed follow-up studies in a new cohort of mice to assess sensitivity to ethanol's stimulant properties. Mice received controlled low-medium doses of ethanol immediately prior to assessing locomotor activity in an open field (Figure 5D). Ethanol dose-dependently increased locomotor activity. PV-mGlu$_5$$^{-/-}$ mice displayed enhanced hyperactivity relative to controls, but no effect of sex or sex by genotype interaction was observed. We assessed the time course of locomotor activity following the top dose (2 g/kg) and found that PV-mGlu$_5$$^{-/-}$ mice displayed increased locomotor activity during the first half of the test (Figure 5E). Finally, we found no differences in center time (Figure 5F), suggesting minimal effect of genotype on anxiety-like behavior. Together, these data implicate mGlu$_5$ receptors in PV cells as a key mediator of sex differences in drinking, potentially related in part to ethanol's stimulant properties.

Discussion

PV-INs are critical regulators of excitatory and inhibitory balance within the PFC [21]. Recent studies have identified sex differences in PV-IN activity and adaptations to experiences under basal conditions and in disease models [25-27]. Understanding the cellular elements and

molecular mechanisms mediating such sex differences in PV-IN function can help direct the development of novel therapeutic targets.

Here, we found that PV-INs from females exhibited lower rheobase and greater current-evoked spike-firing, while passive membrane properties, such as resting membrane potential and membrane resistance, did not differ between sex. These findings are consistent with prior studies showing modest sex differences in PFC PV-IN excitability, whereby PV-INs from female mice are more excitable, particularly during estrus [28,48,65]. The difference in spike-firing without accompanying changes in resting potential or membrane resistance suggests that sex differences in PV-IN intrinsic excitability may stem from voltage-gated and/or calcium-gated conductance. Furthermore, there were no sex differences in rheobase or spike-firing in PV-mGlu$_5^{-/-}$ mice, suggesting mGlu$_5$ receptors are critical in the sex-dependent development and/or maintenance of PV-IN excitability. An important caveat is that we did not conduct simultaneous, interleaved recordings from control mice and PV-mGlu$_5^{-/-}$ mice, so we cannot interpret whether *Grm5* deletion makes PV-INs from female mice less excitable (or those from males more excitable) relative to controls. Nonetheless, these data suggest that basal sex differences in PV-IN excitability stem from gated ion channels or effectors regulated by mGlu$_5$ receptors, potentially including calcium-activated potassium channels [46,66,67], TRP cation channels [68,69], and molecules related to Gα_q signaling.

The current findings reveal sex differences in mGlu$_{1/5}$ receptor signaling in PV-INs, as activation of group 1 mGlu receptors induced larger depolarizing currents in male PFC PV-INs compared to females. Furthermore, the MTEP blocked DHPG-induced inward currents in both sexes but LY367385 only reached significance in male mice. These findings suggest that mGlu$_1$ receptors have enhanced function in PV-INs from male mice. mGlu receptor signaling is regulated at several levels that could vary by sex. One possibility is that *Grm1* could display sex differences in splicing [70], leading to differences in subcellular protein expression and effector signaling [71,72]. Indeed, studies in hippocampal interneurons found that mGlu$_1$ and mGlu$_5$

receptors can activate distinct signaling pathways to regulate calcium mobilization [69], illustrating that $mGlu_1$ and $mGlu_5$ receptors could depolarize PV-INs through divergent mechanisms and these could also vary by sex. Finally, the propensity for $mGlu_1$ and $mGlu_5$ to form homodimers versus heterodimers [73,74] represents an intriguing potential source of sex differences in receptor function. Future studies should address these exciting possibilities.

In addition to membrane physiology, our findings reveal sex differences in $mGlu_1$ and $mGlu_5$ receptor regulation of excitatory synaptic strength on PV-INs. Previous studies found that $mGlu_5$ receptors potentiate spontaneous excitatory drive onto PFC pyramidal cells in male rats [54], whereas $mGlu_1$ receptors serve this role on somatostatin interneurons in both male and female mice [32,51]. Here, we found a striking sex difference in mGlu receptor subtype function on PV-INs, where $mGlu_1$ enhanced excitatory drive in male mice, but $mGlu_5$ receptors enhanced excitatory drive in females. To our surprise, genetically reducing $mGlu_5$ receptor expression in female mice unmasked $mGlu_1$ receptor-dependent responses. These findings suggest that $mGlu_5$ receptors may repress $mGlu_1$ receptor signaling in PV-INs of female mice, potentially at the level of gene expression [75], receptor heterodimerization [73,74], or homodimer signaling crosstalk [76]. Another possibility is that differential coupling with endocannabinoid systems could mediate sex differences in excitatory drive. Indeed, we found that $mGlu_5$ receptor activation potentiated DSE in PV-INs from male but not female mice. Thus, in males, it is possible that $mGlu_5$ receptor activation exerts simultaneous but nullifying actions to amplify excitatory drive and increase endocannabinoid-mediated inhibition in PV-INs. Another interesting possibility is that PV-INs from female mice may not have displayed DSE due to occlusion from enhanced tonic endocannabinoid signaling. Future studies should investigate whether sex differences in constitutive CB1 receptor activity, endocannabinoid production, and ligand affinity [37,77-80] contribute to differences we observed in endocannabinoid-mediated plasticity at PV-INs. Finally, we note these studies are limited by non-specific assessments of

excitatory transmission, so future work should identify the sources of glutamate underlying these effects.

Pharmacological antagonism and genetic deletion of mGlu$_5$ receptors reduce voluntary ethanol consumption, locomotor stimulation, and interoceptive effects [33,81-85]. Here, we found that genetic inhibition of mGlu$_5$ receptors on PV-INs decreased binge drinking in female but not male mice. We had previously observed that IA ethanol precipitates sex-specific changes in synaptic strength on PV-INs, such that drinking reduces and enhances excitatory drive onto PV-INs in female mice but enhances sEPSC frequency onto PV-INs of male mice [28]. The current findings suggest that mGlu$_5$ receptors mediate the reduction in excitatory drive onto PV-INs and this phenomenon facilitates escalated drinking in female mice. Results from our locomotor activity experiment may potentially implicate changes in ethanol's stimulant properties. An intriguing hypothesis for future studies is that female mice escalate drinking because they develop tolerance to ethanol's stimulant effects, and this process is regulated by mGlu$_5$ signaling on PV-INs. Interestingly, previous studies using systemic pharmacology or global genetic deletion of mGlu$_5$ receptors have found decreases in both drinking and stimulant and/or interoceptive effects; thus, the present findings indicate a curious cell type-specific disassociation of these attributes which merits further investigation. One important caveat to the present studies is that PV-expressing cells outside the PFC may have also contributed to the observed behavioral effects.

The current findings suggest that mGlu$_5$ receptors drive sex differences in PV-IN membrane properties and synaptic plasticity. However, mGlu$_5$ receptors are one of the most widely expressed GPCRs in the CNS but most cell types do not express overt physiological sex differences. Even in the closely related somatostatin-expressing interneurons, we and others found minimal evidence for sex differences in basal properties [42,86-88] or in mGlu receptor signaling [29,89]. These findings beg the question, what factors make PV-INs sensitive to sex-dependent mGlu receptor signaling? Sex steroid hormones represent the most likely candidates.

Clemens et al. [65] recently demonstrated that the excitability of PV-INs in somatosensory cortex is regulated by estradiol acting on membrane-associated estrogen receptor-beta (ERβ). Within neocortex, ERβ is nearly exclusively localized within PV-INs [65,90], providing a compelling molecular substrate to propagate sex-dependent signaling. Across many brain regions and cell types, cell-surface ERs can physically and functionally interact with group 1 mGlu receptors to regulate excitability and synaptic transmission [37,38,91-93]. While this literature has generally implicated the ERα subtype in coordinating mGlu receptor signaling in principal neurons, the extent to which these findings generalize to PV-INs is unclear. In addition, progesterone is another major sex hormone that can be metabolized into neurosteroids that alter GABA circuit function [94-96]. Previous studies have also linked testosterone with mGlu synaptic plasticity and eCBs [91,97]; thus, there are several hormone mechanisms that might mediate sex differences in PV-IN function.

Conclusion

Collectively, these findings illustrate that $mGlu_1$ and $mGlu_5$ receptors regulate multiple aspects of PFC PV-IN physiology in a sex-dependent manner. Gaining a better understanding of the cellular and molecular mechanisms that underlie sex differences in brain function should help guide the development and translation of novel therapeutic approaches for AUD and other psychiatric diseases.

Acknowledgments

We thank members of the University of Pittsburgh Department of Psychiatry and Translational Neuroscience Program for stimulating discussions.

Author Contributions

CBF – Conceptualization, Investigation, Writing – Original Draft, Visualization; NIJ – Investigation; LGC – Investigation; SMT – Investigation, Project administration; MLS – Conceptualization, Writing – Review & Editing; MEJ – Conceptualization, Investigation, Writing – Review & Editing, Supervision, Funding acquisition.

Funding

This work was supported by the National Institutes of Health [grant numbers R00AA027806 and MH120066], the Whitehall Foundation [grant number 2022-08-005], and the Brain and Behavior Research Foundation. CBF was supported by the Center for Neuroscience at the University of Pittsburgh and by an institutional predoctoral fellowship from the National Institutes of Health [grant number T32NS07433]. NIJ was supported by the Center for Neuroscience at the University of Pittsburgh Summer Undergraduate Research Program.

Competing Interests

The authors declare no competing interests.

Figure Legends

Figure 1. PV-INs from female mice are more excitable at baseline, and PV-INs from male mice are more responsive to mGlu$_{1/5}$ receptor stimulation. (a) Representative PV-IN current-clamp recordings from female (left, pink) and male (right, red) control mice. Traces depict responses to step current injections. The responses to -150 pA, 0 pA, +150 pA are shown on top and the responses to +300 pA are shown on the bottom. Scale bars 20mV, 0.5s. **(b)** PV-INs from female mice display greater action potential (AP) frequency relative to males in response to step current injections (Two-way mixed-effects analysis, main effect of sex: $F_{1,73} = 4.055$, *: p<0.05; sex x current interaction: $F_{15,1005} = 1.950$, p<0.05). n/N = 33-47/15-19 cells/mice per group. **(c)** PV-INs from female mice display lower rheobase than males ($t_{73} = 2.5$; *: p<0.05). **(d)** Representative traces showing effects of 5 µM and 50 µM DHPG on holding current in PV-INs from female (top) and male (bottom) mice. Scale bars indicate 20 pA, 3 min. **(e)** At a threshold concentration (5 µM), DHPG induced a trend toward greater depolarizing currents in male PV-INs than females (Two-way ANOVA, main effect of sex: $F_{1,56} = 3.4$, #: p<0.072; main effect of antagonist: $F_{2,56} = 2.9$, p<0.065). n/N = 9-11/8-10 cells/mice per group. **(f)** A saturating concentration (50 µM) of DHPG induced greater depolarizing inward currents in PV-INs from male mice. MTEP attenuated the effects of 50 µM DHPG in both sexes. In male mice only, LY367385 blocked DHPG-induced depolarizing currents in PV-INs. (Two-way ANOVA, main effect of sex: $F_{1,57} = 4.3$, p<0.05; main effect of antagonist: $F_{2,57} = 9.3$, p<0.001; *p<0.05, **: p<0.01 Bonferroni post-tests). n/N = 8-14/8-10 cells/mice per group.

Figure 2. mGlu$_1$ and mGlu$_5$ receptors sex-specifically regulate synaptic strength and endocannabinoid plasticity in PV-INs (a) Representative voltage-clamp recordings displaying excitatory postsynaptic current (sEPSC) traces from PV-INs from female (top) and male (bottom) mice. Recordings are from pre-drug baseline period (top) and following sequential 5-minute applications of 5 µM (middle) and 50 µM (bottom) DHPG. 20pA, 2s. **(b)** Minimal effects of 5 µM DHPG application on sEPSC frequency. LY367385 attenuated the effects of 5uM DHPG (Two-way ANOVA, main effect of antagonism: $F_{2, 56} = 4.0$, p<0.05; p<0.05 Control vs. LY367385 Bonferroni post-test). n/N = 8-12/8-10 cells/mice per group. **(c)** 50 µM DHPG induced a three-fold increase in sEPSC frequency in PV-INs from male and female mice. LY367385 inhibited DHPG-induced enhancement of sEPSC frequency in PV-INs from male mice, whereas MTEP attenuated this effect in PV-INs from female mice. (Two-way ANOVA, main effect of sex: $F_{1,56} = 4.2$, p<0.05; main effect of antagonism: $F_{2,56} = 3.4$, p<0.05; antagonism x sex interaction: $F_{2,56} = 3.4$, p<0.05; *: p<0.05; Bonferroni post-tests). n/N =7-15/8-10. Differences in sEPSC frequency were assessed on log-transformed data. **(d)** Representative experiment of depolarization-induced suppression of excitation (DSE) elicited by depolarizing patched PV-INs from -80 mV to +20 mV for 10 seconds. DSE was averaged across three technical replicates at baseline, then again following application of threshold DHPG (5 µM). DHPG did not affect DSE in PV-INs from female mice. Representative traces of baseline EPSCs (black) and DSE (pink) prior to (top) and during 5 µM DHPG application (bottom). Scale bars indicate 200 pA, 20ms. **(e)** Representative DSE experiment in PV-IN from male mouse. **(f)** DHPG potentiated DSE in PV-IN from male but not female mice (Two-way mixed-effects analysis, DHPG x sex interaction: $F_{1,12} = 5.5$, p<0.05; ** p<0.01; Bonferroni post-test). Representative experiments of DSE in male PV-INs in the presence of **(g)** LY367385 and **(h)** MTEP. **(i)** DHPG potentiated DSE in the presence of LY367385 but the effect was blocked by MTEP (Two-way mixed-effects analysis, DHPG x antagonist interaction: $F_{1,24} = 3.6$, p<0.069; *: p<0.05; Bonferroni post-test). n/N = 5-14/5-8 cells/mice per group.

Figure 3. Cell type-specific genetic deletion of *Grm5* abrogates sex differences in PV-IN membrane physiology. (a) Representative PV-IN current-clamp recordings from female (left, light purple) and male (right, dark purple) PV-mGlu$_5^{-/-}$ mice. Traces depict responses to step current injections at -150 pA, 0 pA, +150 pA. **(b)** No differences in current-evoked spiking were observed in PV-INs from male and female PV-mGlu$_5^{-/-}$ mice. **(c)** PV-mGlu$_5^{-/-}$ mice display no sex differences in rheobase (t_{60} = 0.37, ns,). n/N= 30-32/5-8 cells/mice per group. **(d)** Representative traces showing effects of 5 µM and 50 µM DHPG on holding current in female (top) and male (bottom) PV-INs. Scale bars indicate 10 pA, 2 min. **(e)** No effect of sex or antagonist on depolarizing currents following threshold DHPG (5 µM). **(f)** Saturating DHPG (50 µM) induced depolarizing currents to a comparable magnitude in PV-INs from male and female mice. Across sexes, depolarizing currents were blocked by application of either MTEP or LY367385. (Two-way ANOVA, main effect of sex: $F_{1,34}$ = 1.2, p=0.28; main effect of antagonist: $F_{2,34}$ = 5.3, p<0.05, *: p<0.05 Bonferroni post-test). n/N = 5-9/5-8 cells/mice per group.

Figure 4. Cell type-specific genetic deletion of *Grm5* abrogates sex differences in PV-IN synaptic physiology. (a) Representative voltage-clamp recordings displaying sEPSC traces from PV-INs from female (top) and male (bottom) PV-mGlu$_5^{-/-}$ mice. Recordings are from pre-drug baseline period (top) and following sequential 5-minute applications of 5 µM (middle) and 50 µM (bottom) DHPG. Scale bars 20 pA, 2 s. **(b)** Minimal effects of 5 µM DHPG application on sEPSC frequency. **(c)** No sex differences in the effects of 50 µM DHPG on sEPSC frequency in PV-INs from PV-mGlu$_5^{-/-}$ mice, but LY367385 blocked the enhancement of excitatory drive across sexes. (Two-way ANOVA, main effect of sex: $F_{1,34}$ = 2.8, p=0.099; main effect of antagonist: $F_{2,34}$ = 6.0, p<0.01, **: p<0.01 Bonferroni post-test). Differences in sEPSC frequency were assessed on log-transformed data. n/N = 5-9/5-8 cells/mice per group. Representative experiments of DSE from **(d)** female and **(e)** male PV-mGlu$_5^{-/-}$ mice. Representative traces of baseline EPSCs (black) and DSE (pink) prior to (top) and during 5 µM DHPG application (bottom). **(f)** DHPG did not affect DSE in PV-mGlu$_5^{-/-}$ mice. n/N= 7-8/3-4 cells/mice per group.

Figure 5. Female but not male PV-mGlu$_5^{-/-}$ mice display reduced binge drinking. (a) Schematic of intermittent access (IA) ethanol procedure. Mice had access to ethanol in a two-bottle choice procedure every other day for 7 weeks. **(b)** Female mice drank more than male mice overall (Three-way mixed-effects analysis, main effect of sex: $F_{1,34}$ = 22.9, p<0.0001). Follow-up tests indicated that female PV-mGlu$_5^{-/-}$ mice drank less than controls (Two-way mixed-effects analysis, main effect of genotype: $F_{1,16}$ = 7.5, p<0.05; genotype x sex interaction: $F_{6,93}$ = 2.6, p<0.05), but no effect of PV-mGlu$_5^{-/-}$ was detected in male mice. N = 6-12 mice per group. **(c)** Female mice had a greater preference for ethanol more than male mice overall (Three-way mixed-effects analysis, concentration x sex interaction: $F_{6,199}$ = 3.1, p<0.01). Follow-up tests indicated that female PV-mGlu$_5^{-/-}$ mice had lower preference than controls (Two-way mixed-effects analysis, main effect of genotype: $F_{1,16}$ = 8.7, p<0.01), but no effect of PV-mGlu$_5^{-/-}$ was detected in male mice. **(d)** Ethanol dose-dependently increased locomotor activity, and PV-mGlu$_5^{-/-}$ mice displayed lower hyperactivity than controls without any effect of sex (Three-way mixed-effects analysis, main effect of dose: $F_{3,75}$ = 3.6, p<0.05; main effect of genotype: $F_{1,99}$ = 19.3, ****:p<0.0001). N = 7-8 mice per group. **(d)** Ethanol dose-dependently increased locomotor activity, and PV-mGlu$_5^{-/-}$ mice displayed lower hyperactivity than controls without any effect of sex (Three-way mixed-effects analysis, main effect of dose: $F_{3,75}$ = 3.6, p<0.05; main effect of genotype: $F_{1,99}$ = 19.3, ****:p<0.0001). **(e)** PV-mGlu$_5^{-/-}$ mice displayed reduced hyperactivity immediately after the 2 g/kg ethanol injection (Three-way mixed-effects analysis, main effect of genotype: $F_{1,25}$ = 6.6, p<0.05; genotype x time interaction: $F_{3,75}$ = 5.0, p<0.01). **(f)**

Male mice spent more time in the center of the open field but no effect of sex or ethanol were detected (Three-way mixed-effects analysis, main effect of sex: $F_{1,25}$ = 6.8, p<0.05).

Figure S1. PV-INs display comparable resting membrane properties and basal synaptic strength across sexes. No sex differences in **(a)** resting membrane potential (V_m), **(b)** membrane resistance (R_m), or **(c)** medium afterhyperpolarization (mAHP) observed in PV-INs from male and female mice. n/N= 33-44/8 cells/mice per group. **(d)** Representative traces of baseline sEPSCs from female (top, pink) and male (bottom, red) PV-INs. **(e-f)** No sex differences in basal synaptic strength onto PV-INs. n/N= 14-15/8-10cells/mice per group.

Figure S2. PV-mGlu$_5$$^{-/-}$ mice display comparable basal membrane properties and synaptic strength across sexes. PV-mGlu$_5$$^{-/-}$ mice display no sex differences in **(a)** resting membrane potential (V_m), **(b)** membrane resistance (R_m), or **(c)** medium afterhyperpolarization (mAHP). **(d)** Representative traces of baseline EPSCs from PV-mGlu$_5$$^{-/-}$ female (top, light purple) and male (bottom, dark purple) mice. n/N= 22-24/8-10 cells/mice per group. **(e-f)** PV-mGlu$_5$$^{-/-}$ display no sex differences in basal synaptic strength onto PV-INs. n/N= 8-9-/3-4 cells/mice per group.

Figure S3. Threshold DHPG (5 µM) did not affect the amplitude of evoked EPSCs on PV-INs. Prior to depolarization-induced suppression of inhibition experiments, threshold DHPG (5 µM) was applied to cells for 5 minutes while electrically evoked EPSCs were monitored. Threshold DHPG had no effect on EPSC amplitude in control recordings **(a)**, in the presence of LY367385 or MTEP **(b)**, or in PV-mGlu$_5$$^{-/-}$ mice **(c)**. n/N = 5-8/4-5 mice per group.

References

Gobinath AR, Choleris E, Galea LA. Sex, hormones, and genotype interact to influence psychiatric disease, treatment, and behavioral research. J Neurosci Res. 2017;95(1-2):50-64.

Riecher-Rössler A. Sex and gender differences in mental disorders. Lancet Psychiatry. 2017;4(1):8-9.

Marcus SM, Kerber KB, Rush AJ, Wisniewski SR, Nierenberg A, Balasubramani GK, et al. Sex differences in depression symptoms in treatment-seeking adults: confirmatory analyses from the Sequenced Treatment Alternatives to Relieve Depression study. Compr Psychiatry. 2008;49(3):238-46.

Schuch JJ, Roest AM, Nolen WA, Penninx BW, de Jonge P. Gender differences in major depressive disorder: results from the Netherlands study of depression and anxiety. J Affect Disord. 2014;156:156-63.

Häfner H, Maurer K, Löffler W, Riecher-Rössler A. The influence of age and sex on the onset and early course of schizophrenia. Br J Psychiatry. 1993;162:80-6.

Ochoa S, Usall J, Cobo J, Labad X, Kulkarni J. Gender differences in schizophrenia and first-episode psychosis: a comprehensive literature review. Schizophr Res Treatment. 2012;2012:916198.

Goldstein JM, Link BG. Gender and the expression of schizophrenia. J Psychiatr Res. 1988;22(2):141-55.

Randall CL, Roberts JS, Del Boca FK, Carroll KM, Connors GJ, Mattson ME. Telescoping of landmark events associated with drinking: a gender comparison. J Stud Alcohol. 1999;60(2):252-60.

Diehl A, Croissant B, Batra A, Mundle G, Nakovics H, Mann K. Alcoholism in women: is it different in onset and outcome compared to men? Eur Arch Psychiatry Clin Neurosci. 2007;257(6):344-51.

Kouneiher F, Charron S, Koechlin E. Motivation and cognitive control in the human prefrontal cortex. Nat Neurosci. 2009;12(7):939-45.

Schlaepfer TE, Harris GJ, Tien AY, Peng L, Lee S, Pearlson GD. Structural differences in the cerebral cortex of healthy female and male subjects: a magnetic resonance imaging study. Psychiatry Res. 1995;61(3):129-35.

Gur RC, Turetsky BI, Matsui M, Yan M, Bilker W, Hughett P, et al. Sex differences in brain gray and white matter in healthy young adults: correlations with cognitive performance. J Neurosci. 1999;19(10):4065-72.

Goldstein JM, Seidman LJ, Horton NJ, Makris N, Kennedy DN, Caviness VS, Jr., et al. Normal sexual dimorphism of the adult human brain assessed by in vivo magnetic resonance imaging. Cereb Cortex. 2001;11(6):490-7.

Knouse MC, McGrath AG, Deutschmann AU, Rich MT, Zallar LJ, Rajadhyaksha AM, et al. Sex differences in the medial prefrontal cortical glutamate system. Biol Sex Differ. 2022;13(1):66.

Gabbott PL, Warner TA, Jays PR, Salway P, Busby SJ. Prefrontal cortex in the rat: projections to subcortical autonomic, motor, and limbic centers. J Comp Neurol. 2005;492(2):145-77.

Sesack SR, Deutch AY, Roth RH, Bunney BS. Topographical organization of the efferent projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study with Phaseolus vulgaris leucoagglutinin. J Comp Neurol. 1989;290(2):213-42.

Freund TF, Katona I. Perisomatic inhibition. Neuron. 2007;56(1):33-42.

Tremblay R, Lee S, Rudy B. GABAergic Interneurons in the Neocortex: From Cellular Properties to Circuits. Neuron. 2016;91(2):260-92.

Fish KN, Joffe ME. Targeting prefrontal cortex GABAergic microcircuits for the treatment of alcohol use disorder. Front Synaptic Neurosci. 2022;14:936911.

Marin O. Interneuron dysfunction in psychiatric disorders. Nat Rev Neurosci. 2012;13(2):107-20.

Ferguson BR, Gao WJ. PV Interneurons: Critical Regulators of E/I Balance for Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. Front Neural Circuits. 2018;12:37.

Chung DW, Fish KN, Lewis DA. Pathological Basis for Deficient Excitatory Drive to Cortical Parvalbumin Interneurons in Schizophrenia. Am J Psychiatry. 2016;173(11):1131-39.

Perlman G, Tanti A, Mechawar N. Parvalbumin interneuron alterations in stress-related mood disorders: A systematic review. Neurobiol Stress. 2021;15:100380.

Du X, Serena K, Hwang WJ, Grech AM, Wu YWC, Schroeder A, et al. Prefrontal cortical parvalbumin and somatostatin expression and cell density increase during adolescence and are modified by BDNF and sex. Mol Cell Neurosci. 2018;88:177-88.

Goodwill HL, Manzano-Nieves G, LaChance P, Teramoto S, Lin S, Lopez C, et al. Early Life Stress Drives Sex-Selective Impairment in Reversal Learning by Affecting Parvalbumin Interneurons in Orbitofrontal Cortex of Mice. Cell Rep. 2018;25(9):2299-307.e4.

Holland FH, Ganguly P, Potter DN, Chartoff EH, Brenhouse HC. Early life stress disrupts social behavior and prefrontal cortex parvalbumin interneurons at an earlier time-point in females than in males. Neurosci Lett. 2014;566:131-6.

Page CE, Shepard R, Heslin K, Coutellier L. Prefrontal parvalbumin cells are sensitive to stress and mediate anxiety-related behaviors in female mice. Sci Rep. 2019;9(1):19772.

Joffe ME, Winder DG, Conn PJ. Contrasting sex-dependent adaptations to synaptic physiology and membrane properties of prefrontal cortex interneuron subtypes in a mouse model of binge drinking. Neuropharmacology. 2020;178:108126.

Maksymetz J, Byun NE, Luessen DJ, Li B, Barry RL, Gore JC, et al. mGlu(1) potentiation enhances prelimbic somatostatin interneuron activity to rescue schizophrenia-like physiological and cognitive deficits. Cell Rep. 2021;37(5):109950.

Trantham-Davidson H, Burnett EJ, Gass JT, Lopez MF, Mulholland PJ, Centanni SW, et al. Chronic alcohol disrupts dopamine receptor activity and the cognitive function of the medial prefrontal cortex. J Neurosci. 2014;34(10):3706-18.

Barnes SA, Pinto-Duarte A, Kappe A, Zembrzycki A, Metzler A, Mukamel EA, et al. Disruption of mGluR5 in parvalbumin-positive interneurons induces core features of neurodevelopmental disorders. Mol Psychiatry. 2015;20(10):1161-72.

Joffe M, Maksymetz J, Luschinger J, Dogra S, Ferranti A, Luessen D, et al. Acute restraint stress redirects prefrontal cortex circuit function through mGlu5 receptor plasticity on somatostatin-expressing interneurons. Neuron. 2022.

Joffe ME, Centanni SW, Jaramillo AA, Winder DG, Conn PJ. Metabotropic Glutamate Receptors in Alcohol Use Disorder: Physiology, Plasticity, and Promising Pharmacotherapies. ACS Chem Neurosci. 2018;9(9):2188-204.

Niswender CM, Conn PJ. Metabotropic glutamate receptors: physiology, pharmacology, and disease. Annu Rev Pharmacol Toxicol. 2010;50:295-322.

Conn PJ, Pin JP. Pharmacology and functions of metabotropic glutamate receptors. Annu Rev Pharmacol Toxicol. 1997;37:205-37.

Fabian CB, Seney ML, Joffe ME. Sex differences and hormonal regulation of metabotropic glutamate receptor synaptic plasticity. Int Rev Neurobiol. 2023;168:311-47.

Tabatadze N, Huang G, May RM, Jain A, Woolley CS. Sex Differences in Molecular Signaling at Inhibitory Synapses in the Hippocampus. J Neurosci. 2015;35(32):11252-65.

Huang GZ, Woolley CS. Estradiol acutely suppresses inhibition in the hippocampus through a sex-specific endocannabinoid and mGluR-dependent mechanism. Neuron. 2012;74(5):801-8.

Hippenmeyer S, Vrieseling E, Sigrist M, Portmann T, Laengle C, Ladle DR, et al. A developmental switch in the response of DRG neurons to ETS transcription factor signaling. PLoS Biol. 2005;3(5):e159.

Kaiser T, Ting JT, Monteiro P, Feng G. Transgenic labeling of parvalbumin-expressing neurons with tdTomato. Neuroscience. 2016;321:236-45.

Xu J, Zhu Y, Contractor A, Heinemann SF. mGluR5 has a critical role in inhibitory learning. J Neurosci. 2009;29(12):3676-84.

Joffe ME, Winder DG, Conn PJ. Contrasting sex-dependent adaptations to synaptic physiology and membrane properties of prefrontal cortex interneuron subtypes in a mouse model of binge drinking. Neuropharmacology. 2020:108126.

Straiker A, Mackie K. Depolarization-induced suppression of excitation in murine autaptic hippocampal neurones. J Physiol. 2005;569(Pt 2):501-17.

Marcus DJ, Bedse G, Gaulden AD, Ryan JD, Kondev V, Winters ND, et al. Endocannabinoid Signaling Collapse Mediates Stress-Induced Amygdalo-Cortical Strengthening. Neuron. 2020;105(6):1062-76.e6.

Thompson SM, Fabian CB, Ferranti AS, Joffe ME. Acute alcohol and chronic drinking bidirectionally regulate the excitability of prefrontal cortex vasoactive intestinal peptide interneurons. Neuropharmacology. 2023;238:109638.

Mannaioni G, Marino MJ, Valenti O, Traynelis SF, Conn PJ. Metabotropic glutamate receptors 1 and 5 differentially regulate CA1 pyramidal cell function. J Neurosci. 2001;21(16):5925-34.

Di Menna L, Joffe ME, Iacovelli L, Orlando R, Lindsley CW, Mairesse J, et al. Functional partnership between mGlu3 and mGlu5 metabotropic glutamate receptors in the central nervous system. Neuropharmacology. 2018;128:301-13.

Ferranti AS, Johnson KA, Winder DG, Conn PJ, Joffe ME. Prefrontal cortex parvalbumin interneurons exhibit decreased excitability and potentiated synaptic strength after ethanol reward learning. Alcohol. 2022;101:17-26.

Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C. Interneurons of the neocortical inhibitory system. Nat Rev Neurosci. 2004;5(10):793-807.

Kawaguchi Y, Kubota Y. GABAergic cell subtypes and their synaptic connections in rat frontal cortex. Cereb Cortex. 1997;7(6):476-86.

Maksymetz J, Byun NE, Luessen DJ, Li B, Barry RL, Gore JC, et al. mGlu1 potentiation enhances prelimbic somatostatin interneuron activity to rescue schizophrenia-like physiological and cognitive deficits. Cell Rep. 2021;37(5):109950.

Chu Z, Hablitz JJ. Activation of group I mGluRs increases spontaneous IPSC frequency in rat frontal cortex. J Neurophysiol. 1998;80(2):621-7.

Whittington MA, Traub RD, Jefferys JG. Synchronized oscillations in interneuron networks driven by metabotropic glutamate receptor activation. Nature. 1995;373(6515):612-5.

Marek GJ, Zhang C. Activation of metabotropic glutamate 5 (mGlu5) receptors induces spontaneous excitatory synaptic currents in layer V pyramidal cells of the rat prefrontal cortex. Neurosci Lett. 2008;442(3):239-43.

Maejima T, Hashimoto K, Yoshida T, Aiba A, Kano M. Presynaptic inhibition caused by retrograde signal from metabotropic glutamate to cannabinoid receptors. Neuron. 2001;31(3):463-75.

Varma N, Carlson GC, Ledent C, Alger BE. Metabotropic glutamate receptors drive the endocannabinoid system in hippocampus. J Neurosci. 2001;21(24):Rc188.

Ohno-Shosaku T, Shosaku J, Tsubokawa H, Kano M. Cooperative endocannabinoid production by neuronal depolarization and group I metabotropic glutamate receptor activation. Eur J Neurosci. 2002;15(6):953-61.

Robbe D, Kopf M, Remaury A, Bockaert J, Manzoni OJ. Endogenous cannabinoids mediate long-term synaptic depression in the nucleus accumbens. Proc Natl Acad Sci U S A. 2002;99(12):8384-8.

Guo B, Wang J, Yao H, Ren K, Chen J, Yang J, et al. Corrigendum: Chronic Inflammatory Pain Impairs mGluR5-Mediated Depolarization-Induced Suppression of Excitation in the Anterior Cingulate Cortex. Cereb Cortex. 2019;29(4):1698.

Diana MA, Marty A. Endocannabinoid-mediated short-term synaptic plasticity: depolarization-induced suppression of inhibition (DSI) and depolarization-induced suppression of excitation (DSE). Br J Pharmacol. 2004;142(1):9-19.

Kasten CR, Holmgren EB, Wills TA. Metabotropic Glutamate Receptor Subtype 5 in Alcohol-Induced Negative Affect. Brain Sci. 2019;9(8).

Jury NJ, DiBerto JF, Kash TL, Holmes A. Sex differences in the behavioral sequelae of chronic ethanol exposure. Alcohol. 2017;58:53-60.

Finn DA, Helms ML, Nipper MA, Cohen A, Jensen JP, Devaud LL. Sex differences in the synergistic effect of prior binge drinking and traumatic stress on subsequent ethanol intake and neurochemical responses in adult C57BL/6J mice. Alcohol. 2018;71:33-45.

Becker HC, Lopez MF. Increased ethanol drinking after repeated chronic ethanol exposure and withdrawal experience in C57BL/6 mice. Alcohol Clin Exp Res. 2004;28(12):1829-38.

Clemens AM, Lenschow C, Beed P, Li L, Sammons R, Naumann RK, et al. Estrus-Cycle Regulation of Cortical Inhibition. Curr Biol. 2019;29(4):605-15 e6.

Saugstad JA, Marino MJ, Folk JA, Hepler JR, Conn PJ. RGS4 inhibits signaling by group I metabotropic glutamate receptors. J Neurosci. 1998;18(3):905-13.

Cannady R, McGonigal JT, Newsom RJ, Woodward JJ, Mulholland PJ, Gass JT. Prefrontal Cortex KCa2 Channels Regulate mGlu5-Dependent Plasticity and Extinction of Alcohol-Seeking Behavior. J Neurosci. 2017;37(16):4359-69.

Kim SJ, Kim YS, Yuan JP, Petralia RS, Worley PF, Linden DJ. Activation of the TRPC1 cation channel by metabotropic glutamate receptor mGluR1. Nature. 2003;426(6964):285-91.

Topolnik L, Azzi M, Morin F, Kougioumoutzakis A, Lacaille JC. mGluR1/5 subtype-specific calcium signalling and induction of long-term potentiation in rat hippocampal oriens/alveus interneurones. J Physiol. 2006;575(Pt 1):115-31.

Trabzuni D, Ramasamy A, Imran S, Walker R, Smith C, Weale ME, et al. Widespread sex differences in gene expression and splicing in the adult human brain. Nat Commun. 2013;4:2771.

Francesconi A, Duvoisin RM. Alternative splicing unmasks dendritic and axonal targeting signals in metabotropic glutamate receptor 1. J Neurosci. 2002;22(6):2196-205.

Naito R, Kassai H, Sakai Y, Schönherr S, Fukaya M, Schwarzer C, et al. New Features on the Expression and Trafficking of mGluR1 Splice Variants Exposed by Two Novel Mutant Mouse Lines. Front Mol Neurosci. 2018;11:439.

Lee J, Munguba H, Gutzeit VA, Singh DR, Kristt M, Dittman JS, et al. Defining the Homo- and Heterodimerization Propensities of Metabotropic Glutamate Receptors. Cell Rep. 2020;31(5):107605.

Doumazane E, Scholler P, Zwier JM, Trinquet E, Rondard P, Pin JP. A new approach to analyze cell surface protein complexes reveals specific heterodimeric metabotropic glutamate receptors. FASEB J. 2011;25(1):66-77.

Notartomaso S, Nakao H, Mascio G, Scarselli P, Cannella M, Zappulla C, et al. mGlu1 Receptors Monopolize the Synaptic Control of Cerebellar Purkinje Cells by Epigenetically Down-Regulating mGlu5 Receptors. Sci Rep. 2018;8(1):13361.

Sevastyanova TN, Kammermeier PJ. Cooperative signaling between homodimers of metabotropic glutamate receptors 1 and 5. Mol Pharmacol. 2014;86(5):492-504.

Ferraro A, Wig P, Boscarino J, Reich CG. Sex differences in endocannabinoid modulation of rat CA1 dendritic neurotransmission. Neurobiol Stress. 2020;13:100283.

Rodríguez de Fonseca F, Cebeira M, Ramos JA, Martín M, Fernández-Ruiz JJ. Cannabinoid receptors in rat brain areas: sexual differences, fluctuations during estrous cycle and changes after gonadectomy and sex steroid replacement. Life Sci. 1994;54(3):159-70.

Bara A, Manduca A, Bernabeu A, Borsoi M, Serviado M, Lassalle O, et al. Sex-dependent effects of in utero cannabinoid exposure on cortical function. Elife. 2018;7.

Bernabeu A, Bara A, Murphy Green MN, Manduca A, Wager-Miller J, Borsoi M, et al. Sexually Dimorphic Adolescent Trajectories of Prefrontal Endocannabinoid Synaptic Plasticity Equalize in Adulthood, Reflected by Endocannabinoid System Gene Expression. Cannabis Cannabinoid Res. 2023.

Backstrom P, Bachteler D, Koch S, Hyytia P, Spanagel R. mGluR5 antagonist MPEP reduces ethanol-seeking and relapse behavior. Neuropsychopharmacology. 2004;29(5):921-8.

Bird MK, Kirchhoff J, Djouma E, Lawrence AJ. Metabotropic glutamate 5 receptors regulate sensitivity to ethanol in mice. Int J Neuropsychopharmacol. 2008;11(6):765-74.

Stevenson RA, Hoffman JL, Maldonado-Devincci AM, Faccidomo S, Hodge CW. MGluR5 activity is required for the induction of ethanol behavioral sensitization and associated changes in ERK MAP kinase phosphorylation in the nucleus accumbens shell and lateral habenula. Behav Brain Res. 2019;367:19-27.

Besheer J, Grondin JJ, Cannady R, Sharko AC, Faccidomo S, Hodge CW. Metabotropic glutamate receptor 5 activity in the nucleus accumbens is required for the maintenance of ethanol self-administration in a rat genetic model of high alcohol intake. Biol Psychiatry. 2010;67(9):812-22.

Besheer J, Faccidomo S, Grondin JJ, Hodge CW. Regulation of motivation to self-administer ethanol by mGluR5 in alcohol-preferring (P) rats. Alcohol Clin Exp Res. 2008;32(2):209-21.

Dao NC, Brockway DF, Suresh Nair M, Sicher AR, Crowley NA. Somatostatin neurons control an alcohol binge drinking prelimbic microcircuit in mice. Neuropsychopharmacology. 2021.

Dao NC, Suresh Nair M, Magee SN, Moyer JB, Sendao V, Brockway DF, et al. Forced Abstinence From Alcohol Induces Sex-Specific Depression-Like Behavioral and Neural Adaptations in Somatostatin Neurons in Cortical and Amygdalar Regions. Front Behav Neurosci. 2020;14:86.

Sicher AR, Starnes WD, Griffith KR, Dao NC, Smith GC, Brockway DF, et al. Adolescent binge drinking leads to long-lasting changes in cortical microcircuits in mice. Neuropharmacology. 2023;234:109561.

Joffe ME, Maksymetz J, Luschinger JR, Dogra S, Ferranti AS, Luessen DJ, et al. Acute restraint stress redirects prefrontal cortex circuit function through mGlu(5) receptor plasticity on somatostatin-expressing interneurons. Neuron. 2022;110(6):1068-83.e5.

Blurton-Jones M, Tuszynski MH. Estrogen receptor-beta colocalizes extensively with parvalbumin-labeled inhibitory neurons in the cortex, amygdala, basal forebrain, and hippocampal formation of intact and ovariectomized adult rats. J Comp Neurol. 2002;452(3):276-87.

Gross KS, Moore KM, Meisel RL, Mermelstein PG. mGluR5 Mediates Dihydrotestosterone-Induced Nucleus Accumbens Structural Plasticity, but Not Conditioned Reward. Front Neurosci. 2018;12:855.

Fagan MP, Ameroso D, Meng A, Rock A, Maguire J, Rios M. Essential and sex-specific effects of mGluR5 in ventromedial hypothalamus regulating estrogen signaling and glucose balance. Proc Natl Acad Sci U S A. 2020;117(32):19566-77.

Vandegrift BJ, Hilderbrand ER, Satta R, Tai R, He D, You C, et al. Estrogen Receptor alpha Regulates Ethanol Excitation of Ventral Tegmental Area Neurons and Binge Drinking in Female Mice. J Neurosci. 2020;40(27):5196-207.

Maguire JL, Stell BM, Rafizadeh M, Mody I. Ovarian cycle-linked changes in GABA(A) receptors mediating tonic inhibition alter seizure susceptibility and anxiety. Nat Neurosci. 2005;8(6):797-804.

Maguire J, Mody I. Neurosteroid synthesis-mediated regulation of GABA(A) receptors: relevance to the ovarian cycle and stress. J Neurosci. 2007;27(9):2155-62.

Abramian AM, Comenencia-Ortiz E, Modgil A, Vien TN, Nakamura Y, Moore YE, et al. Neurosteroids promote phosphorylation and membrane insertion of extrasynaptic GABAA receptors. Proc Natl Acad Sci U S A. 2014;111(19):7132-7.

Conde K, Fabelo C, Krause WC, Propst R, Goethel J, Fischer D, et al. Testosterone Rapidly Augments Retrograde Endocannabinoid Signaling in Proopiomelanocortin Neurons to Suppress Glutamatergic Input from Steroidogenic Factor 1 Neurons via Upregulation of Diacylglycerol Lipase-alpha. Neuroendocrinology. 2017;105(4):341-56.

Figure 1: PV-INs from female mice are more excitable at baseline, and PV-INs from male mice are more responsive to mGlu$_{1/5}$ receptor stimulation

Figure 2. mGlu$_1$ and mGlu$_5$ receptors sex-specifically regulate synaptic strength and endocannabinoid plasticity in PV-INs

Figure 3. Cell type-specific genetic deletion of *Grm5* abrogates sex differences in PV-IN membrane physiology

Figure 4. Cell type-specific genetic deletion of *Grm5* abrogates sex differences in PV-IN synaptic physiology

Figure 5: Female but not male PV-mGlu$_5$$^{-/-}$ mice display reduced binge drinking

Figure S1: PV-INs display comparable resting membrane properties and basal synaptic strength across sexes

Figure S2: PV-mGlu$_5$$^{-/-}$ mice display comparable basal membrane properties and synaptic strength across sexes

Figure S3: Threshold DHPG (5µM) did not affect the amplitude of evoked EPSCs on PV-INs