# Parvalbumin interneuron mGlu$_5$ receptors govern sex differences in prefrontal cortex physiology and binge drinking

Carly B. Fabian[a,b,c], Nilah D. Jordan[c], Rebecca H. Cole[a,b,c], Lily G. Carley[a,b,c], Shannon M. Thompson[a,b,c], Marianne L. Seney[a,b,c], and Max E. Joffe[a,b,c,*]

  a. Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, 15219, USA
  b. Translational Neuroscience Program, University of Pittsburgh, Pittsburgh, PA
  c. Center for Neuroscience University of Pittsburgh, Pittsburgh, PA

*Correspondence to:
Max E. Joffe, Ph.D.
Assistant Professor
University of Pittsburgh
219 Bridgeside Point II
450 Technology Dr
Pittsburgh, PA 15219
E-Mail: joffeme@upmc.edu
Phone: 412-383-6028
Twitter: @mejoffe

Short title: Group 1 mGlu receptors mediate sex differences in PFC PV-INs

Key words: prefrontal cortex, alcohol, G protein-coupled receptor (GPCR), cognition, GABA, addiction

Article Type: Research
Abstract: 232 words
Article: 3996 words (limit 4000)
Figures: 5
Tables: 0
Supplemental Information: 6 figures, 2 tables
References: 100

**Abstract**

Despite established sex differences in the prevalence and presentation of psychiatric disorders,

little is known about the cellular and synaptic mechanisms that guide these differences under

basal conditions. Proper function of the prefrontal cortex (PFC) is essential for the top-down

regulation of motivated behaviors. Activity of the PFC is tightly controlled by parvalbumin-

expressing interneurons (PV-INs), a key subpopulation of fast-spiking GABAergic cells that

regulate cortical excitability through direct innervations onto the perisomatic regions of nearby

pyramidal cells. Recent rodent studies have identified notable sex differences in PV-IN activity

and adaptations to experiences such as binge drinking. Here, we investigated the cellular and

molecular mechanisms that underlie sex-specific regulation of PFC PV-IN function. Using

whole-cell patch clamp electrophysiology and selective pharmacology, we report that PV-INs

from female mice are more excitable than those from males. Moreover, we find that $mGlu_1$ and

$mGlu_5$ metabotropic glutamate receptors regulate cell excitability, excitatory drive, and

endocannabinoid signaling at PFC PV-INs in a sex-dependent manner. Genetic deletion of

$mGlu_5$ receptors from PV-expressing cells abrogates all sex differences observed in PV-IN

membrane and synaptic physiology. Lastly, we report that female, but not male, PV-$mGlu_5^{-/-}$

mice exhibit decreased voluntary drinking on an intermittent access schedule, which could be

related to changes in ethanol's stimulant properties. Importantly, these studies identify $mGlu_1$

and $mGlu_5$ receptors as candidate signaling molecules involved in sex differences in PV-IN

activity and behaviors relevant for alcohol use.

**Introduction**

Sex differences contribute to the incidence, presentation, and severity of several

psychiatric disorders [1,2]. For instance, women with major depressive disorder (MDD)

experience greater symptom severity [3,4], men have an earlier age of onset of schizophrenia

[5-7], and women display a faster progression through disease milestones of alcohol use

disorder (AUD) [8,9]. Together, the broad sex differences in psychiatric diseases motivate

continued preclinical research to identify the cells and plasticity mechanisms that mediate sex

differences in brain function.

The prefrontal cortex (PFC) provides top-down regulation of motivated behaviors and

displays sex differences in form and function [10-14]. Most PFC neurons are glutamatergic

pyramidal cells that send, receive, and integrate information to and from various subcortical and

cortical structures [15,16]. Pyramidal cell activity is tightly regulated by a small number of

inhibitory neurons that transmit γ-aminobutyric acid (GABA). Parvalbumin-expressing inhibitory

neurons (PV-INs) form a key subpopulation of these GABAergic cells which synapse onto

perisomatic regions of pyramidal neurons and display fast-spiking firing properties [17-19].

Dysregulation of PFC PV-IN activity has been implicated in the etiology of many psychiatric

disorders [20-23]. A growing body of literature in rodent models has also identified notable sex

differences in PV-IN activity and adaptations to experiences. Indeed, male and female mice

display sex differences in developmental trajectory of PFC PV expression [24], stress sex-

specifically alters PFC PV-IN function [25-27], and PFC PV-INs undergo sex-dependent

adaptations following four-weeks binge drinking [28]. Taken together, these studies suggest that

sex differences in molecular mechanisms governing synaptic plasticity and intrinsic properties

on PV-INs are critical in mediating experience-dependent adaptations to PFC function.

The group 1 metabotropic glutamate (mGlu) receptors, $mGlu_1$ and $mGlu_5$, are expressed

on PFC PV-INs [29-32] and regulate cellular processes related to membrane physiology,

synaptic strength, and endocannabinoid plasticity [33-35]. In addition, there are sex differences

in signaling and function of $mGlu_1$ and $mGlu_5$ receptors [36-38]. Here, we tested the hypothesis

that $mGlu_1$ and $mGlu_5$ receptors mediate sex differences in PFC PV-IN function. Using

genetically engineered mice, whole-cell patch clamp electrophysiology, and selective

pharmacology, we examined sex differences in PV-IN function and regulation by $mGlu_1$ and

$mGlu_5$ receptors. We then assessed a causal role for mGlu receptors in regulating PFC PV-IN

function through mice with cell type-specific deletion of $mGlu_5$ receptors. The present results

support the hypothesis that PV-IN $mGlu_5$ receptor signaling is critical for sex differences in PFC

physiology and drinking.

**Materials and Methods**

**Mice**

Mice were maintained on a 12:12 light-dark cycle (lights on at 6:00 AM) with food and water *ad*

*libitum*. Mice were bred at Charles River Laboratory (Wilmington, MA) and delivered to the

University of Pittsburgh at 6-8 weeks of age. All mice were on a congenic C57BL/6J background

and experiments began in adulthood (>10 weeks). Sex determination was limited to an

assessment of external genitalia. Mice expressing tdTomato in PV-INs were bred by crossing

female homozygous PV-Cre mice [39] (Jackson Laboratories, 017320) with male homozygous

Rosa26-loxP-STOP-loxP-CAG-tdTomato "Ai9" mice [40] (Jackson Laboratories, 007909). PV-

$mGlu_5^{-/-}$ mice were generated by crossing PV-Cre and Ai9 mice with those harboring a floxed

*Grm5* locus [31,41] (Jackson Laboratories, 028626). All experiments were approved by the

University of Pittsburgh IACUC and conducted in accordance with the NIH Guidelines.

**Brain slice preparation**

Mice (N=68) were decapitated under isoflurane anesthesia; brains were immediately dissected

and placed in *N*-methyl-D-glucamine (NMDG) cutting solution, (in mM: 93 NMDG, 20 HEPES,

2.5 KCl, 0.5 CaCl$_2$, 10 MgCl$_2$, 1.2 NaH$_2$PO$_4$, 25 glucose, 5 Na-ascorbate, and 3 Na-pyruvate).

Coronal slices (300-µM) of the PL-PFC were prepared in room-temperature NMDG solution.

Slices recovered for 10 minutes in warm (32-35°C) NMDG solution and were then transferred to

room temperature artificial cerebral spinal fluid (ACSF) (in mM:119 NaCl, 2.5 KCl, 2 CaCl$_2$, 1

MgCl$_2$, 1 NaH$_2$PO$_4$, 11 glucose, and 26 NaHCO$_3$) for at least 1 hour before recordings. All

solutions were oxygenated (95% O$_2$ / 5% CO$_2$).

**Ex vivo whole-cell patch-clamp electrophysiology**

Data were acquired using a Multiclamp 700B amplifier and pCLAMP 11 software (Molecular

Devices). Slices were transferred to the recording chamber and perfused at 2mL/min with

heated ACSF (30-32°C). For all whole-cell patch-clamp recordings, cells were patched with a

borosilicate glass pipette pulled to 3-5 MΩ with a horizontal electrode puller (P-1000, Stutter

Instruments). Pipettes were filled with a potassium-based internal solution (in mM: 125 K-

gluconate, 4 NaCl, 10 HEPES, 0.1 EGTA, 4 MgATP, 0.3 Na$_2$GTP, 10 Tris-phosphocreatine).

PV-INs in layer 5 of the PL-PFC were identified by tdTomato fluorescence and cell identity was

confirmed via characteristic fast-spiking phenotype. After gaining whole-cell access, PV-INs

were dialyzed for 5 minutes under voltage-clamp at -80mV. Membrane properties were then

assessed under current-clamp as described [42]. A series of current injections (1s, starting at -

150pA, ending at +500pA, increments of 25pA) were applied to patched cells. Resting

membrane potential (V$_m$) was determined as the average baseline membrane potential prior to

current injections. Membrane resistance (R$_m$) was calculated as the slope of the hyperpolarizing

membrane potential over three injections [-75, -50, -25pA]. Rheobase was determined as the

minimal current injection needed to elicit firing.

Cells were then returned to voltage-clamp configuration. Spontaneous excitatory

postsynaptic currents (sEPSCs) were electrically isolated at -80mV, near the reversal potential

for chloride [28]. DHPG-induced changes in depolarizing currents and excitatory transmission

onto PFC PV-INs were assessed sequentially in the same cell. Following a 5-minute baseline,

PV-INs underwent sequential 5-min application of 5µM and then 50µM DHPG. In some

experiments, LY367385 (100µM) or MTEP (3µM) was applied prior to baseline recordings and

was present for the entire experiment. The holding current did not drift in 15-minute vehicle

control experiments (0.3 ± 3.5pA, n=5 cells). AMPAR-mediated sEPSCs on PV-INs were

identified using pattern-matching event detection software (ClampFit). The template was

generated from PV-IN recordings (match threshold 3.5) and did not detect events in NBQX.

Change in holding current and sEPSC frequency were calculated by comparing the average

values over the last 2 minutes of drug application to the last 2 minutes of baseline. At the

conclusion of the 5-minute 50µM DHPG application, cells were returned to current-clamp and

membrane properties were re-assessed.

In separate PV-INs, endocannabinoid-mediated short-term plasticity was assessed via

depolarization-induced suppression of excitation (DSE) [43,44]. Bipolar nichrome wire

stimulating electrodes were positioned in layer 2/3 medial to the recording electrode. EPSCs

were evoked with a single pulse (10-50µA) at 0.2Hz to elicit EPSCs between 100-200pA. After

establishing a stable 1-minute baseline, DSE was elicited by depolarizing patched PV-INs to

+20mV for 10 seconds without stimulation, followed by an immediate return to -80mV and

electrical stimulation. DSE was repeated 3-4 times per cell under basal conditions. DHPG (5µM)

was then applied for 5 minutes and the DSE protocol was repeated. Changes in the EPSC

amplitude were normalized to baseline. To assess inhibitory transmission, some PV-INs were

patched with a high-chloride cesium-based internal solution (in mM: 125 CsCl, 5 NaCl, 10

HEPES, 4 MgATP, 0.3 $Na_2$GTP, 10 Tris-phosphocreatine, 5 Qx-314-Br) and held at -70mV in

NBQX (10 µM). Data for electrophysiology studies was processed using ClampFit (Molecular

Devices).

**Intermittent access (IA) ethanol**

IA ethanol was performed as described [42,46]. For three 24-hour periods per week, ethanol

was provided in home cages, with 24-48 hours between each exposure. Ethanol was diluted

from undenatured 95% ethyl alcohol and was delivered by modified conical tubes with rubber

stoppers and open tip tubes. Ethanol was generally provided and removed 3-4 hours prior to

start of the dark cycle. After a ramp during the first three sessions [3, 6, 10%], the ethanol

concentration was set at 20% ethanol for the remainder of the study.

**Locomotor response to ethanol**

We assessed the locomotor response to ethanol in mice without previous exposure to ethanol.

Mice received saline vehicle (10 µL/kg, *i.p.*) or ethanol (0.5, 1, or 2 g/kg) immediately before

being placed in an open field chamber for 20 minutes. Over 4 consecutive days, all mice

received all doses in a counterbalanced Latin-square design (see schematic in Figure 5).

Location and distance traveled were determined by video-based tracking (ANY-maze,

Stoelting).

**Drugs**

Drugs were purchased from HelloBio. Concentrations were based on previous studies

[29,32,47,48]. DHPG has no effect on Group 2 or Group 3 mGlu receptors ($mGlu_{2,3,4,7,8}$) at

concentrations up to 1mM [49,50]. Drugs were formulated, aliquoted, and frozen at 1000X, then

thawed and diluted in ACSF on the day of recording. LY367385 [(*S*)-(+)-α-Amino-4-carboxy-2-

methylbenzeneaceticacid] stock was made in 0.1M NaOH. MTEP [3-((2-Methyl-1,3-thiazol-4-

yl)ethynyl)pyridine hydrochloride], DHPG [(*RS*)-3,5-Dihydroxyphenylglycine], and NBQX stocks

were made in water.

**Statistics**

Data are presented as mean±SEM or box plots (min, Q1, median, Q3, max). Mean±SEM for key

experiments is reported in Table S1. Statistical analyses were performed using GraphPad

Prism. For electrophysiology, *n* represents the number of cells. For all analyses, p<0.05 was

considered significant and p<0.07 was labeled trending. Data were analyzed using two-tailed

Student's t-test. Two-way or three-way ANOVA, or mixed-effects analysis, with Bonferonni post-

hoc comparisons were used for analyses requiring multiple comparisons. Statistical analyses

are reported in the legends and in Table S2.

**Results**

**PL-PFC PV-INs from female are mice are more excitable than PV-INs from males**

To assess intrinsic physiology of PFC PV-INs, we made whole-cell patch-clamp

recordings from genetically engineered mice expressing tdTomato in PV-positive cells

[32,42,51]. All (n=75/75) tdTomato-expressing cells from layer 5 of the PL-PFC displayed

characteristic fast-spiking interneuron properties [52,53], including: high frequency firing, low

input resistance, and minimal spike-frequency adaptation (Figure 1A). In general, PV-INs from

male and female mice displayed comparable basal membrane properties, without significant

differences in resting potential, membrane resistance, or medium afterhyperpolarization (Figure

S1A-C). However, in response to positive current injections, cells from female mice exhibited

greater spike-firing compared to cells from males (Figure 1B) and fired action potentials in

response to a lower threshold current (Figure 1C). Together, these data indicate that while PV-

INs exhibit similar basal membrane properties between sexes, PV-INs from female mice are

more excitable than those from males.

**mGlu$_1$ and mGlu$_5$ receptors depolarize PFC PV-IN in a sex-dependent manner**

We and others previously demonstrated that mGlu$_1$ and mGlu$_5$ receptors modulate PV-

INs in frontal cortex [29-31], but most of these studies were limited to males. We therefore

aimed to determine whether $mGlu_1$ and $mGlu_5$ receptors modulate PV-IN function in a similar

manner in males and females. We examined sensitivity to $mGlu_1$ and $mGlu_5$ receptor stimulation

using a threshold concentration of the non-selective $mGlu_{1/5}$ agonist DHPG (5µM) alone, in the

presence of the selective $mGlu_5$ NAM MTEP (3µM), or in the presence of the selective $mGlu_1$

antagonist LY367385 (100µM) (Figure 1D). Threshold DHPG induced modest depolarizing

currents in PV-INs of both sexes; however, PV-INs from male mice displayed a trend towards

greater depolarizing changes in holding current compared to females (Figure 1E), with no effect

of either antagonist. We next assessed a high concentration of DHPG (50µM), which induced

greater depolarizing current relative to the threshold concentration in both sexes (Figure 1F).

Depolarizing currents in response to 50µM DHPG were greater in PV-INs from males, and

selective pharmacological antagonism revealed additional sex differences in the receptor

subtypes involved. In PV-INs from all mice, MTEP attenuated DHPG-induced depolarizing

currents. By contrast, LY367385 reduced DHPG currents in PV-INs from male mice but did

reach significance in females. Dual application of both MTEP and LY367385 attenuated but did

not completely block DHPG currents in both male and female PFC PV-INs (Figure S2A),

potentially due to competition between DHPG and LY367385 at the orthosteric site or

partial/biased NAM activity of MTEP in this assay. Together, these data suggest that PFC PV-IN

membrane physiology is modulated by $mGlu_5$ receptors across sexes, but that PV-INs from

male mice are more sensitive to $mGlu_1$ receptor activation. We also returned cells to current-

clamp at the conclusion of the DHPG wash-on and found PV-INs from males were depolarized

to a greater extent than females (Figure S3A).

**$mGlu_1$ and $mGlu_5$ regulate synaptic strength on PFC PV-INs in a sex-dependent manner.**

In addition to membrane physiology, Group 1 mGlu receptors can modulate synaptic

strength on many cell types, including other types of interneurons in frontal cortex [32,54-57].

We measured AMPA receptor-mediated sEPSCs under voltage-clamp and found no evidence

for basal sex differences in excitatory or inhibitory synaptic strength (Figure S1D-I). We next

determined whether Group 1 mGlu receptors regulate excitatory transmission. Threshold DHPG

(5µM) had no effect on sEPSC frequency (Figure 2A and 2B), but a higher concentration of

DHPG (50µM) induced a three-fold increase in sEPSC frequency in PV-INs across sexes

(Figure 2C). Further pharmacological studies revealed a sex-specific receptor mechanism

underlying this effect: LY367385 blocked enhancement of excitatory drive in PV-INs from male

mice, while the effects of DHPG on sEPSC frequency were attenuated by MTEP in PV-INs from

females. Dual application of MTEP/LY367385 blocked the DHPG-induced increase in excitatory

drive (Figure S2B). These findings suggest a sex-specific role of the group 1 mGlu receptors in

regulating synaptic strength onto PFC PV-INs, primarily driven by $mGlu_1$ receptor signaling in

males and by $mGlu_5$ receptors in females.

**$mGlu_5$ receptors facilitate short-term plasticity at PV-INs from male but not female mice**

We next sought to examine mechanisms of mGlu receptor synaptic plasticity. In several

brain regions, activation of postsynaptic Group 1 mGlu receptors can trigger endocannabinoid

release and reductions in neurotransmitter release [58-61]. We elicited endocannabinoid

plasticity by depolarization-induced suppression of excitation (DSE) [62,63], a form of short-term

depression mediated by CB1 receptor activation and reduced glutamate release probability.

Under control conditions, PV-IN EPSC amplitude was reduced by approximately 20% by cell

depolarization across sexes (Figure 2D and 2E). We next assessed whether $mGlu_1$ and $mGlu_5$

receptor activation can facilitate DSE by applying the threshold concentration of DHPG (5µM).

While no effect was observed in female mice, threshold DHPG enhanced DSE in PV-INs from

male mice (Figure 2E and 2F). Importantly, threshold DHPG did not affect EPSC amplitude in

the absence of postsynaptic depolarization (Figure S4), suggesting the facilitation of DSE is a

metaplastic change in endocannabinoid signaling, rather than an overall change in glutamate

transmission. Based on our previous findings, we predicted that the LY367385 would block the

enhancement of DSE in male mice. To our surprise, DHPG retained the ability to enhance DSE

in LY367385 (Figure 2G and 2I), while MTEP completely blocked DSE potentiation (Figure 2H

and 2I). We did not perform experiments using the antagonists in female mice because there

was no effect of DHPG alone. Taken together, these findings indicate that $mGlu_5$ receptors can

potentiate endocannabinoid signaling in PV-INs from male mice.

**Genetic deletion of $mGlu_5$ receptors from PV-INs abrogates sex-differences in membrane**

**and synaptic physiology**

235             We next used a genetic approach to examine $mGlu_5$ receptor function within PV-INs. We

bred mice to selectively ablate $mGlu_5$ receptors from PV-expressing cells via Cre-mediated

deletion of *Grm5* [31]. PV-$mGlu_5^{-/-}$ mice display postnatal reduction of $mGlu_5$ receptors to ~20%

of control levels, with recombination beginning between around P28, reflecting endogenous PV

expression. We first assessed PV-IN membrane properties in PV-$mGlu_5^{-/-}$ mice. In contrast to

controls (Figure 1A-1C), we observed no sex differences in PV-IN spike-firing (Figure 3A and

3B) or rheobase (Figure 3C) in PV-$mGlu_5^{-/-}$ mice. Furthermore, we observed no sex differences

in any other membrane property or in basal synaptic strength (Figure S5 and Table S2). These

data suggest that $mGlu_5$ receptors regulate the development and/or maintenance of sex

differences in PV-IN excitability. We next assessed responses to $mGlu_{1/5}$ receptor activation,

hypothesizing that DHPG would not depolarize PV-INs in female PV-$mGlu_5^{-/-}$ mice. To our

surprise, DHPG induced comparable depolarizing currents in both sexes (Figure 3D-3F), and

currents were attenuated by either MTEP or LY367385. These findings suggest that (1) there is

residual expression of $mGlu_5$ receptors in PV-$mGlu_5^{-/-}$ mice, consistent with previous studies

[31], and (2) female PV-$mGlu_5^{-/-}$ mice display unmasked sensitivity to $mGlu_1$ receptor

antagonism. We next assessed DHPG potentiation of excitatory drive in PV-$mGlu_5^{-/-}$ mice

(Figure 4A). 5µM DHPG had little effect on sEPSC frequency (Figure 4B), while 50µM DHPG

rapidly increased excitatory drive on PV-INs in both sexes (Figure 4C). In contrast to control

mice, we observed a main effect of mGlu receptor antagonist without an effect of sex or

interaction. Posthoc tests across sexes revealed that LY367385, but not MTEP, reduced the

ability of DHPG to potentiate sEPSC frequency, again indicating loss of $mGlu_5$ function in

female mice and enhanced sensitivity to $mGlu_1$ receptor antagonism. Finally, we performed

DSE experiments to assess eCB signaling (Figure 4D-4F). DHPG failed to potentiate DSE in

PV-INs of PV-$mGlu_5^{-/-}$ mice. Taken together, all sex differences in PV-IN physiology and mGlu

receptor regulation were absent in PV-$mGlu_5^{-/-}$ mice.

**Deleting $mGlu_5$ receptors from PV-INs in female mice, but not males, decreases voluntary**

**drinking**

$mGlu_5$ receptor signaling regulates behaviors related to alcohol use [33,64], and

previous studies from our group have demonstrated that drinking and ethanol reward learning

alter the PV-IN physiology [28,51]. Based on this literature, and our current findings that $mGlu_5$

receptors regulate synaptic strength onto PV-INs in female mice but not males, we

hypothesized that $mGlu_5$ receptors in PV-INs regulate binge drinking in a sex-dependent

manner. To test this, we provided control and PV-$mGlu_5^{-/-}$ mice with two-bottle choice IA ethanol

over 7 weeks (Figure 5A) [42,46]. As observed by our lab and many others [42,65-67], female

mice drank more than male mice overall. Moreover, a sex by genotype interaction emerged,

such that female but not male PV-$mGlu_5^{-/-}$ mice drank less than their respective controls (Figure

5B). Female mice had greater ethanol preference than males but there was no effect of

genotype on this measurement (Figure 5C). We next performed follow-up studies in a new

cohort of mice to assess sensitivity to ethanol's stimulant properties (Figure 5D). Low-medium

doses of ethanol increased locomotor activity. PV-$mGlu_5^{-/-}$ mice displayed enhanced

hyperactivity relative to controls, but no effect of sex or interaction was observed. We assessed

the time course of locomotor activity following the top dose (2 g/kg) and found that PV-$mGlu_5^{-/-}$

mice displayed increased locomotor activity during the first half of the test (Figure 5E). Finally,

we found no differences in center time (Figure 5F), suggesting minimal effect of genotype on

anxiety-like behavior. Together, these data implicate mGlu$_5$ receptors in PV cells as a key

mediator of sex differences in drinking, potentially related in part to ethanol's stimulant

properties.

**Discussion**

Recent studies have identified sex differences in PFC PV-IN activity and adaptations to

experiences under basal conditions and in disease models [25-27]. Understanding the

molecular mechanisms mediating such sex differences in PV-IN function can help direct the

development of novel therapeutic targets.

Here, we found that PV-INs from females exhibited lower rheobase and greater current-

evoked spike-firing, while passive membrane properties, such as resting membrane potential

and membrane resistance, did not differ between sex. These findings are consistent with prior

studies showing modest sex differences in PFC PV-IN excitability, whereby PV-INs from female

mice are more excitable, particularly during estrus [28,51,68]. The difference in spike-firing

without accompanying changes in resting potential or membrane resistance suggests that sex

differences in PV-IN intrinsic excitability may stem from voltage-gated and/or calcium-gated

conductance. Furthermore, there were no sex differences in rheobase or spike-firing in PV-

mGlu$_5^{-/-}$ mice, suggesting mGlu$_5$ receptors are critical in the sex-dependent development and/or

maintenance of PV-IN excitability. An important caveat is that we did not conduct simultaneous,

interleaved recordings from control mice and PV-mGlu$_5^{-/-}$ mice, so we cannot interpret whether

*Grm5* deletion makes PV-INs from female mice less excitable (or those from males more

excitable) relative to controls. Nonetheless, these data suggest that basal sex differences in PV-

IN excitability stem from gated ion channels or effectors regulated by mGlu$_5$ receptors,

potentially including calcium-activated potassium channels [47,69,70], TRP cation channels

[71,72], and molecules related to G$\alpha_q$ signaling.

The current findings reveal sex differences in $mGlu_{1/5}$ receptor signaling in PV-INs, as

activation of group 1 mGlu receptors induced larger depolarizing currents in male PFC PV-INs

compared to females (Supplemental Figure S6). Furthermore, the MTEP blocked DHPG-

induced inward currents in both sexes but LY367385 only reached significance in male mice.

These findings suggest that $mGlu_1$ receptors have enhanced function in PV-INs from male mice.

mGlu receptor signaling is regulated at several levels that could vary by sex. One possibility is

that *Grm1* could display sex differences in splicing [73], leading to differences in subcellular

protein expression and effector signaling [74,75]. Indeed, studies in hippocampal interneurons

found that $mGlu_1$ and $mGlu_5$ receptors can activate distinct signaling pathways to regulate

calcium mobilization [72], illustrating that $mGlu_1$ and $mGlu_5$ receptors could depolarize PV-INs

through divergent mechanisms and these could also vary by sex. Finally, the propensity for

$mGlu_1$ and $mGlu_5$ to form homodimers versus heterodimers [76,77] represents an intriguing

potential source of sex differences in receptor function. Future studies should address these

exciting possibilities.

In addition to membrane physiology, our findings reveal sex differences in $mGlu_1$ and

$mGlu_5$ receptor regulation of excitatory synaptic strength on PV-INs. Previous studies found that

$mGlu_5$ receptors potentiate spontaneous excitatory drive onto PFC pyramidal cells in male rats

[57], whereas $mGlu_1$ receptors serve this role on somatostatin interneurons in both male and

female mice [32,54]. Here, we found a striking sex difference in mGlu receptor subtype function

on PV-INs, where $mGlu_1$ enhanced excitatory drive in male mice, but $mGlu_5$ receptors enhanced

excitatory drive in females. To our surprise, genetically reducing $mGlu_5$ receptor expression in

female mice unmasked $mGlu_1$ receptor-dependent responses. These findings suggest that

$mGlu_5$ receptors may repress $mGlu_1$ receptor signaling in PV-INs of female mice, potentially at

the level of gene expression [78], receptor heterodimerization [76,77], or homodimer signaling

crosstalk [79]. Another possibility is that differential coupling with endocannabinoid systems

could mediate sex differences in excitatory drive. Indeed, we found that $mGlu_5$ receptor

activation potentiated DSE in PV-INs from male but not female mice. Thus, in males, it is

possible that mGlu$_5$ receptor activation exerts simultaneous but nullifying actions to amplify

excitatory drive and increase endocannabinoid-mediated inhibition in PV-INs. Importantly, we

note these studies are limited by non-specific assessments of excitatory transmission, so future

work should identify the sources of glutamate underlying these effects.

Pharmacological antagonism and genetic deletion of mGlu$_5$ receptors reduce voluntary

ethanol consumption, locomotor stimulation, and interoceptive effects [33,84-88]. Here, we

found that genetic inhibition of mGlu$_5$ receptors on PV-INs decreased binge drinking in female

but not male mice. We had previously observed that IA ethanol precipitates sex-specific

changes in synaptic strength on PV-INs, such that drinking reduces and enhances excitatory

drive onto PV-INs in female mice but enhances sEPSC frequency onto PV-INs of male mice

[28]. The current findings suggest that mGlu$_5$ receptors mediate the reduction in excitatory drive

onto PV-INs and this phenomenon facilitates escalated drinking in female mice. Results from

our locomotor activity experiment may potentially implicate changes in ethanol's stimulant

properties. An intriguing hypothesis for future studies is that female mice escalate drinking to

overcome tolerance to ethanol's stimulant effects, and this process is regulated by mGlu$_5$

signaling on PV-INs. Interestingly, previous studies using systemic pharmacology or global

genetic deletion of mGlu$_5$ receptors have found decreases in both drinking and stimulant and/or

interoceptive effects; thus, the present findings indicate a curious cell type-specific

disassociation of these attributes which merits further investigation. One important caveat to the

present studies is that PV-expressing cells outside the PFC may have also contributed to the

observed behavioral effects.

The current findings suggest that mGlu$_5$ receptors drive sex differences in PV-IN

membrane properties and synaptic plasticity. However, mGlu$_5$ receptors are one of the most

widely expressed GPCRs but most cell types do not express overt physiological sex differences.

Even in the closely related somatostatin-expressing interneurons, we and others found minimal

evidence for sex differences in basal properties [42,89-91] or in mGlu receptor signaling [29,92].

These findings beg the question, what factors make PV-INs sensitive to sex-dependent mGlu

receptor signaling? Sex steroid hormones represent the most likely candidates. Clemens et al.

[68] recently demonstrated that the excitability of PV-INs in somatosensory cortex is regulated

by estradiol acting on membrane-associated estrogen receptor-beta (ERβ). Within neocortex,

ERβ is highly localized within PV-INs [68,93], providing a compelling molecular substrate to

propagate sex-dependent signaling. Across many brain regions and cell types, cell-surface ERs

interact with group 1 mGlu receptors to regulate excitability and synaptic transmission [37,38,94-

96]. In addition to estrogens, progesterone is another major sex hormone that can be

metabolized into neurosteroids that alter GABA circuit function [97-99]. Previous studies have

also linked testosterone with mGlu synaptic plasticity and eCBs [94,100]; thus, there are several

hormone mechanisms that might mediate sex differences in PV-IN function.

**Conclusion**

Collectively, these findings illustrate that $mGlu_1$ and $mGlu_5$ receptors regulate multiple

aspects of PFC PV-IN physiology in a sex-dependent manner. Gaining a better

understanding of the cellular and molecular mechanisms that underlie sex differences in

brain function should help guide the development and translation of novel therapeutic

approaches for AUD and other psychiatric diseases.

**Acknowledgments**

We thank members of the University of Pittsburgh Department of Psychiatry and Translational

Neuroscience Program for stimulating discussions.

**Author Contributions**

CBF – Conceptualization, Investigation, Writing – Original Draft, Visualization; NDJ –

Investigation; RHC – Investigation; LGC – Investigation; SMT – Investigation, Project

administration; MLS – Conceptualization, Writing – Review & Editing; MEJ – Conceptualization,

Investigation, Writing – Review & Editing, Supervision, Funding Acquisition.

**Funding**

This work was supported by the National Institutes of Health [grant numbers R00AA027806 and

MH120066], the Whitehall Foundation [grant number 2022-08-005], and the Brain and Behavior

Research Foundation. CBF and RHC were supported by the Center for Neuroscience at the

University of Pittsburgh. CBF was supported by an institutional predoctoral fellowship from the

National Institutes of Health [grant number T32NS07433]. NDJ was supported by the Center for

Neuroscience at the University of Pittsburgh Summer Undergraduate Research Program.

**Competing Interests**

The authors declare no competing interests.

    **Figure Legends**

**Figure 1. PV-INs from female mice are more excitable at baseline, and PV-INs from male**
**mice are more responsive to mGlu$_{1/5}$ receptor stimulation. (a)** Representative PV-IN current-
clamp recordings from female (left, pink) and male (right, red) control mice. Traces depict
responses to step current injections. The responses to -150 pA, 0 pA, +150 pA are shown on
top and the responses to +300 pA are shown on the bottom. Scale bars 20mV, 0.5s. **(b)** PV-INs
from female mice display greater action potential (AP) frequency relative to males in response
to step current injections (Two-way mixed-effects analysis, main effect of sex: $F_{1,73} = 4.1$, *:
$p<0.05$; sex x current interaction: $F_{15,1005} = 2.0$, $p<0.05$). n/N = 33-42/15-19 cells/mice per group.
**(c)** PV-INs from female mice display lower rheobase than males ($t_{73} = 2.5$; *: $p<0.05$). **(d)**
Representative traces showing effects of 5 µM and 50 µM DHPG on holding current in PV-INs
from female (top) and male (bottom) mice. Scale bars indicate 20 pA, 3 min. **(e)** At a threshold
concentration (5 µM), DHPG induced a trend toward greater depolarizing currents in male PV-
INs than females (Two-way ANOVA, main effect of sex: $F_{1,56} = 3.4$, #: $p<0.072$; main effect of
antagonist: $F_{2,56} = 2.9$, $p<0.065$). n/N = 9-11/8-10 cells/mice per group. **(f)** A saturating
concentration (50 µM) of DHPG induced greater depolarizing inward currents in PV-INs from
male mice. MTEP attenuated the effects of 50 µM DHPG in both sexes. In male mice only,
LY367385 blocked DHPG-induced depolarizing currents in PV-INs. (Two-way ANOVA, main
effect of sex: $F_{1,57} = 4.3$, $p<0.05$; main effect of antagonist: $F_{2,57} = 9.3$, $p<0.001$; *$p<0.05$, **:
$p<0.01$ Bonferroni post-tests). n/N = 8-14/8-10 cells/mice per group.
416
**Figure 2. mGlu$_1$ and mGlu$_5$ receptors sex-specifically regulate synaptic strength and**
**endocannabinoid plasticity in PV-INs (a)** Representative voltage-clamp recordings displaying
excitatory postsynaptic current (sEPSC) traces from PV-INs from female (top) and male
(bottom) mice. Recordings are from pre-drug baseline period (top) and following sequential 5-
minute applications of 5 µM (middle) and 50 µM (bottom) DHPG. 20pA, 2s. **(b)** Minimal effects
of 5 µM DHPG application on sEPSC frequency. LY367385 attenuated the effects of 5uM
DHPG (Two-way ANOVA, main effect of antagonism: $F_{2,56} = 4.0$, $p<0.05$). n/N = 8-12/8-10
cells/mice per group. **(c)** 50 µM DHPG induced a three-fold increase in sEPSC frequency in PV-
INs from male and female mice. LY367385 inhibited DHPG-induced enhancement of sEPSC
frequency in PV-INs from male mice, whereas MTEP attenuated this effect in PV-INs from
female mice. (Two-way ANOVA, main effect of sex: $F_{1,56} = 4.2$, $p<0.05$; main effect of
antagonism: $F_{2,56} = 3.4$, $p<0.05$; antagonism x sex interaction: $F_{2,56} = 3.4$, $p<0.05$; *: $p<0.05$
Bonferroni post-test). n/N =7-15/8-10. Differences in sEPSC frequency were assessed on log-
transformed data. **(d)** Representative experiment of depolarization-induced suppression of
excitation (DSE) elicited by depolarizing patched PV-INs from -80 mV to +20 mV for 10
seconds. DSE was averaged across three technical replicates at baseline, then again following
application of threshold DHPG (5 µM). DHPG did not affect DSE in PV-INs from female mice.
Representative traces of baseline EPSCs (black) and DSE (pink) prior to (top) and during 5 µM
DHPG application (bottom). Scale bars indicate 200 pA, 20ms. **(e)** Representative DSE
experiment in PV-IN from male mouse. **(f)** DHPG potentiated DSE in PV-IN from male but not
female mice (Two-way mixed-effects analysis, DHPG x sex interaction: $F_{1,12} = 5.5$, $p<0.05$; **
$p<0.01$ Bonferroni post-test). Representative experiments of DSE in male PV-INs in the
presence of **(g)** LY367385 and **(h)** MTEP. **(i)** DHPG potentiated DSE in the presence of
LY367385 but the effect was blocked by MTEP (Two-way mixed-effects analysis, DHPG x
antagonist interaction: $F_{1,24} = 3.6$, $p<0.069$; *: $p<0.05$ Bonferroni post-test). n/N = 5-14/5-8
cells/mice per group.
443
444

**Figure 3. Cell type-specific genetic deletion of *Grm5* abrogates sex differences in PV-IN membrane physiology. (a)** Representative PV-IN current-clamp recordings from female (left, light purple) and male (right, dark purple) PV-mGlu$_5^{-/-}$ mice. Traces depict responses to step current injections at -150 pA, 0 pA, +150 pA. **(b)** No differences in current-evoked spiking were observed in PV-INs from male and female PV-mGlu$_5^{-/-}$ mice. **(c)** PV-mGlu$_5^{-/-}$ mice display no sex differences in rheobase ($t_{60}$ = 0.37, ns,). n/N= 30-32/5-8 cells/mice per group. **(d)** Representative traces showing effects of 5 µM and 50 µM DHPG on holding current in female (top) and male (bottom) PV-INs. Scale bars indicate 10 pA, 2 min. **(e)** No effect of sex or antagonist on depolarizing currents following threshold DHPG (5 µM). **(f)** Saturating DHPG (50 µM) induced depolarizing currents to a comparable magnitude in PV-INs from male and female mice. Across sexes, depolarizing currents were blocked by application of either MTEP or LY367385. (Two-way ANOVA, main effect of sex: $F_{1,34}$ = 1.2, p=0.28; main effect of antagonist: $F_{2,34}$ = 5.3, p<0.05, *: p<0.05 Bonferroni post-test). n/N = 5-9/5-8 cells/mice per group.

**Figure 4. Cell-type specific genetic deletion of *Grm5* abrogates sex differences in PV-IN synaptic physiology. (a)** Representative voltage-clamp recordings displaying sEPSC traces from PV-INs from female (top) and male (bottom) PV-mGlu$_5^{-/-}$ mice. Recordings are from pre-drug baseline period (top) and following sequential 5-minute applications of 5 µM (middle) and 50 µM (bottom) DHPG. Scale bars 20 pA, 2 s. **(b)** Minimal effects of 5 µM DHPG application on sEPSC frequency. **(c)** No sex differences in the effects of 50 µM DHPG on sEPSC frequency in PV-INs from PV-mGlu$_5^{-/-}$ mice, but LY367385 blocked the enhancement of excitatory drive across sexes. (Two-way ANOVA, main effect of sex: $F_{1,34}$ = 2.8, p=0.099; main effect of antagonist: $F_{2,34}$ = 6.0, p<0.01, *: p<0.05, **: p<0.01 Bonferroni post-test). Differences in sEPSC frequency were assessed on log-transformed data. n/N = 5-9/5-8 cells/mice per group. Representative experiments of DSE from **(d)** female and **(e)** male PV-mGlu$_5^{-/-}$ mice. Representative traces of baseline EPSCs (black) and DSE (pink) prior to (top) and during 5 µM DHPG application (bottom). **(f)** DHPG did not affect DSE in PV-mGlu$_5^{-/-}$ mice. n/N= 7-8/3-4 cells/mice per group.

**Figure 5. Female but not male PV-mGlu$_5^{-/-}$ mice display reduced binge drinking. (a)** Schematic of intermittent access (IA) ethanol procedure. Mice had access to ethanol in a two-bottle choice procedure every other day for 7 weeks. **(b)** Female mice drank more than male mice overall (Three-way mixed-effects analysis, main effect of sex: $F_{1,34}$ = 22.9, p<0.0001). Follow-up tests indicated that female PV-mGlu$_5^{-/-}$ mice drank less than controls (Two-way mixed-effects analysis, main effect of genotype: $F_{1,16}$ = 7.5, p<0.05; genotype x sex interaction: $F_{6,93}$ = 2.6, p<0.05), but no effect of PV-mGlu$_5^{-/-}$ was detected in male mice. N = 6-12 mice per group. **(c)** Female mice had a greater preference for ethanol more than male mice overall (Three-way mixed-effects analysis, concentration x sex interaction: $F_{6,199}$ = 3.1, p<0.01), but there was no effect of genotype on ethanol preference. **(d)** PV-mGlu$_5^{-/-}$ mice displayed enhanced hyperactivity following ethanol injections, without any effect of sex (Three-way mixed-effects analysis, main effect of dose: $F_{3,75}$ = 3.6, p<0.05; main effect of genotype: $F_{1,99}$ = 19.3, ****:p<0.0001). N = 7-8 mice per group. main effect of genotype: $F_{1,99}$ = 19.3, ****:p<0.0001). **(e)** PV-mGlu$_5^{-/-}$ mice displayed enhanced hyperactivity immediately after the 2 g/kg ethanol injection (Three-way mixed-effects analysis, main effect of genotype: $F_{1,25}$ = 6.6, p<0.05; genotype x time interaction: $F_{3,75}$ = 5.0, p<0.01). **(f)** Male mice spent more time in the center of the open field but no effect of sex or ethanol were detected (Three-way mixed-effects analysis, main effect of sex: $F_{1,25}$ = 6.8, p<0.05).

**Supplemental Figure Legends**

**Figure S1. PV-INs display comparable resting membrane properties and basal synaptic strength across sexes**. No sex differences in **(a)** resting membrane potential ($V_m$), **(b)** membrane resistance ($R_m$), or **(c)** medium afterhyperpolarization (mAHP) observed in PV-INs from male and female mice. n/N= 33-44/8 cells/mice per group. **(d)** Representative traces of baseline spontaneous excitatory postsynaptic currents (sEPSCs) from female (top, pink) and male (bottom, red) PV-INs. **(e-f)** No sex differences in basal synaptic strength onto PV-INs. n/N= 14-15/8-10 cells/mice per group. **(g)** Representative traces of baseline spontaneous inhibitory postsynaptic currents (sIPSCs) from PV-INs from female (top, pink) and male (bottom, red) mice. No sex differences in average **(h)** sIPSC frequency or **(i)** sIPSC amplitude onto PFC PV-INs. n/N= 8/3 cells/mice per group.

**Figure S2. Dual application of MTEP and LY367385 attenuated DHPG currents and blocked increase in synaptic transmission in PFC PV-INs. (a)** DHPG currents in the presence of a combination of MTEP (3 µM) and mGlu$_1$ antagonist LY367385 (100 µM). Currents are reduced compared to control or single NAM experiments (Figure 1); however, DHPG still exerted a concentration-dependent and sex-dependent depolarizing current (Two-way ANOVA, main effect of concentration: $F_{1,11} = 20.1$, $p<0.001$, main effect of sex: $F_{1,11} = 4.5$, $p<0.06$, concentration x sex interaction: $F_{1,11} = 3.7$, $p<0.08$, *:$p<0.05$, **:$p<0.01$ Bonferonni post-test). **(b)** MTEP/LY367385 application blocked the ability of DHPG to enhance excitatory drive onto PFC PV-INs (Two-way ANOVA, main effect of concentration: $F_{1,11} = 0.3$, n.s.; one-sample t-test $t_{25} = 1.2$, n.s.). n/N = 6-7/3.

**Figure S3. Male PV-INs undergo greater depolarization following mGlu$_1$ and mGlu$_5$ receptor stimulation.** Membrane properties of PFC PV-INs assessed under current clamp configuration during end of 50 µM DHPG wash-on. Change calculated as the difference relative to the baseline before DHPG wash-on. **(a)** PV-INs from males exhibit a greater DHPG-induced depolarization in resting membrane potential (Vm) than cells from females (Two-way ANOVA, main effect of sex: $F_{2,37} = 1.6$, *:$p<0.05$). No main effect of sex or antagonist on the change in **(b)** rheobase or **(c)** membrane resistance ($R_m$) in PV-INs. n/N = 6-9/8-10 cells/mice per group.

**Figure S4. Threshold DHPG (5 µM) did not affect the amplitude of evoked EPSCs on PV-INs.** Prior to depolarization-induced suppression of inhibition experiments, threshold DHPG (5 µM) was applied to cells for 5 minutes while electrically evoked EPSCs were monitored. Threshold DHPG had no effect on EPSC amplitude in control recordings **(a)**, in the presence of LY367385 or MTEP **(b)**, or in PV-mGlu$_5$$^{-/-}$ mice **(c)**. n/N = 5-8/4-5 mice per group.

**Figure S5. PV-mGlu$_5$$^{-/-}$ mice display comparable basal membrane properties and synaptic strength across sexes.** PV-mGlu$_5$$^{-/-}$ mice display no sex differences in **(a)** resting membrane potential ($V_m$), **(b)** membrane resistance ($R_m$), or **(c)** medium afterhyperpolarization (mAHP). **(d)** Representative traces of baseline EPSCs from PV-mGlu$_5$$^{-/-}$ female (top, light purple) and male (bottom, dark purple) mice. n/N= 22-24/8-10 cells/mice per group. **(e-f)** PV-mGlu$_5$$^{-/-}$ display no sex differences in basal synaptic strength onto PV-INs. n/N= 8-9/3-4 cells/mice per group.

**Figure S6. Schematic depicting proposed sex differences in mGlu$_1$ and mGlu$_5$ receptor signaling in PFC PV-INs.** In female mice, mGlu$_5$ receptor signaling depolarizes PV-INs, likely through inhibition of potassium channels. In males, cell significant depolarization of PV-INs is mediated by both mGlu$_1$ and mGlu$_5$ receptor signaling. Synaptic strength onto PV-INs is enhanced by mGlu$_5$ receptor activation in female mice; by contrast, in males synaptic strength

543 onto PV-INs is potentiated by $mGlu_1$ receptor activation. Finally, $mGlu_5$ receptor activation
544 potentiates endocannabinoid short-term plasticity onto PV-INs from male mice but not females.

**References**

1    Gobinath AR, Choleris E, Galea LA. Sex, hormones, and genotype interact to influence
psychiatric disease, treatment, and behavioral research. J Neurosci Res. 2017;95(1-2):50-
64.

2    Riecher-Rössler A. Sex and gender differences in mental disorders. Lancet Psychiatry.
2017;4(1):8-9.

3    Marcus SM, Kerber KB, Rush AJ, Wisniewski SR, Nierenberg A, Balasubramani GK, et al.
Sex differences in depression symptoms in treatment-seeking adults: confirmatory
analyses from the Sequenced Treatment Alternatives to Relieve Depression study.
Compr Psychiatry. 2008;49(3):238-46.

4    Schuch JJ, Roest AM, Nolen WA, Penninx BW, de Jonge P. Gender differences in major
depressive disorder: results from the Netherlands study of depression and anxiety. J
Affect Disord. 2014;156:156-63.

5    Häfner H, Maurer K, Löffler W, Riecher-Rössler A. The influence of age and sex on the
onset and early course of schizophrenia. Br J Psychiatry. 1993;162:80-6.

6    Ochoa S, Usall J, Cobo J, Labad X, Kulkarni J. Gender differences in schizophrenia and
first-episode psychosis: a comprehensive literature review. Schizophr Res Treatment.
2012;2012:916198.

7    Goldstein JM, Link BG. Gender and the expression of schizophrenia. J Psychiatr Res.
1988;22(2):141-55.

8    Randall CL, Roberts JS, Del Boca FK, Carroll KM, Connors GJ, Mattson ME. Telescoping of
landmark events associated with drinking: a gender comparison. J Stud Alcohol.
1999;60(2):252-60.

9    Diehl A, Croissant B, Batra A, Mundle G, Nakovics H, Mann K. Alcoholism in women: is it
different in onset and outcome compared to men? Eur Arch Psychiatry Clin Neurosci.
2007;257(6):344-51.

10   Kouneiher F, Charron S, Koechlin E. Motivation and cognitive control in the human
prefrontal cortex. Nat Neurosci. 2009;12(7):939-45.

11   Schlaepfer TE, Harris GJ, Tien AY, Peng L, Lee S, Pearlson GD. Structural differences in the
cerebral cortex of healthy female and male subjects: a magnetic resonance imaging
study. Psychiatry Res. 1995;61(3):129-35.

12   Gur RC, Turetsky BI, Matsui M, Yan M, Bilker W, Hughett P, et al. Sex differences in brain
gray and white matter in healthy young adults: correlations with cognitive performance.
J Neurosci. 1999;19(10):4065-72.

13   Goldstein JM, Seidman LJ, Horton NJ, Makris N, Kennedy DN, Caviness VS, Jr., et al.
Normal sexual dimorphism of the adult human brain assessed by in vivo magnetic
resonance imaging. Cereb Cortex. 2001;11(6):490-7.

14 Knouse MC, McGrath AG, Deutschmann AU, Rich MT, Zallar LJ, Rajadhyaksha AM, et al.
Sex differences in the medial prefrontal cortical glutamate system. Biol Sex Differ.
2022;13(1):66.

15 Gabbott PL, Warner TA, Jays PR, Salway P, Busby SJ. Prefrontal cortex in the rat:
projections to subcortical autonomic, motor, and limbic centers. J Comp Neurol.
2005;492(2):145-77.

16 Sesack SR, Deutch AY, Roth RH, Bunney BS. Topographical organization of the efferent
projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study
with Phaseolus vulgaris leucoagglutinin. J Comp Neurol. 1989;290(2):213-42.

17 Freund TF, Katona I. Perisomatic inhibition. Neuron. 2007;56(1):33-42.

18 Tremblay R, Lee S, Rudy B. GABAergic Interneurons in the Neocortex: From Cellular
Properties to Circuits. Neuron. 2016;91(2):260-92.

19 Fish KN, Joffe ME. Targeting prefrontal cortex GABAergic microcircuits for the treatment
of alcohol use disorder. Front Synaptic Neurosci. 2022;14:936911.

20 Marin O. Interneuron dysfunction in psychiatric disorders. Nat Rev Neurosci.
2012;13(2):107-20.

21 Ferguson BR, Gao WJ. PV Interneurons: Critical Regulators of E/I Balance for Prefrontal
Cortex-Dependent Behavior and Psychiatric Disorders. Front Neural Circuits. 2018;12:37.

22 Chung DW, Fish KN, Lewis DA. Pathological Basis for Deficient Excitatory Drive to Cortical
Parvalbumin Interneurons in Schizophrenia. Am J Psychiatry. 2016;173(11):1131-39.

23 Perlman G, Tanti A, Mechawar N. Parvalbumin interneuron alterations in stress-related
mood disorders: A systematic review. Neurobiol Stress. 2021;15:100380.

24 Du X, Serena K, Hwang WJ, Grech AM, Wu YWC, Schroeder A, et al. Prefrontal cortical
parvalbumin and somatostatin expression and cell density increase during adolescence
and are modified by BDNF and sex. Mol Cell Neurosci. 2018;88:177-88.

25 Goodwill HL, Manzano-Nieves G, LaChance P, Teramoto S, Lin S, Lopez C, et al. Early Life
Stress Drives Sex-Selective Impairment in Reversal Learning by Affecting Parvalbumin
Interneurons in Orbitofrontal Cortex of Mice. Cell Rep. 2018;25(9):2299-307.e4.

26 Holland FH, Ganguly P, Potter DN, Chartoff EH, Brenhouse HC. Early life stress disrupts
social behavior and prefrontal cortex parvalbumin interneurons at an earlier time-point
in females than in males. Neurosci Lett. 2014;566:131-6.

27 Page CE, Shepard R, Heslin K, Coutellier L. Prefrontal parvalbumin cells are sensitive to
stress and mediate anxiety-related behaviors in female mice. Sci Rep. 2019;9(1):19772.

28 Joffe ME, Winder DG, Conn PJ. Contrasting sex-dependent adaptations to synaptic
physiology and membrane properties of prefrontal cortex interneuron subtypes in a
mouse model of binge drinking. Neuropharmacology. 2020;178:108126.

29  Maksymetz J, Byun NE, Luessen DJ, Li B, Barry RL, Gore JC, et al. mGlu(1) potentiation
enhances prelimbic somatostatin interneuron activity to rescue schizophrenia-like
physiological and cognitive deficits. Cell Rep. 2021;37(5):109950.

30  Trantham-Davidson H, Burnett EJ, Gass JT, Lopez MF, Mulholland PJ, Centanni SW, et al.
Chronic alcohol disrupts dopamine receptor activity and the cognitive function of the
medial prefrontal cortex. J Neurosci. 2014;34(10):3706-18.

31  Barnes SA, Pinto-Duarte A, Kappe A, Zembrzycki A, Metzler A, Mukamel EA, et al.
Disruption of mGluR5 in parvalbumin-positive interneurons induces core features of
neurodevelopmental disorders. Mol Psychiatry. 2015;20(10):1161-72.

32  Joffe M, Maksymetz J, Luschinger J, Dogra S, Ferranti A, Luessen D, et al. Acute restraint
stress redirects prefrontal cortex circuit function through mGlu5 receptor plasticity on
somatostatin-expressing interneurons. Neuron. 2022.

33  Joffe ME, Centanni SW, Jaramillo AA, Winder DG, Conn PJ. Metabotropic Glutamate
Receptors in Alcohol Use Disorder: Physiology, Plasticity, and Promising
Pharmacotherapies. ACS Chem Neurosci. 2018;9(9):2188-204.

34  Niswender CM, Conn PJ. Metabotropic glutamate receptors: physiology, pharmacology,
and disease. Annu Rev Pharmacol Toxicol. 2010;50:295-322.

35  Conn PJ, Pin JP. Pharmacology and functions of metabotropic glutamate receptors. Annu
Rev Pharmacol Toxicol. 1997;37:205-37.

36  Fabian CB, Seney ML, Joffe ME. Sex differences and hormonal regulation of
metabotropic glutamate receptor synaptic plasticity. Int Rev Neurobiol. 2023;168:311-
47.

37  Tabatadze N, Huang G, May RM, Jain A, Woolley CS. Sex Differences in Molecular
Signaling at Inhibitory Synapses in the Hippocampus. J Neurosci. 2015;35(32):11252-65.

38  Huang GZ, Woolley CS. Estradiol acutely suppresses inhibition in the hippocampus
through a sex-specific endocannabinoid and mGluR-dependent mechanism. Neuron.
2012;74(5):801-8.

39  Hippenmeyer S, Vrieseling E, Sigrist M, Portmann T, Laengle C, Ladle DR, et al. A
developmental switch in the response of DRG neurons to ETS transcription factor
signaling. PLoS Biol. 2005;3(5):e159.

40  Kaiser T, Ting JT, Monteiro P, Feng G. Transgenic labeling of parvalbumin-expressing
neurons with tdTomato. Neuroscience. 2016;321:236-45.

41  Xu J, Zhu Y, Contractor A, Heinemann SF. mGluR5 has a critical role in inhibitory learning.
J Neurosci. 2009;29(12):3676-84.

42  Joffe ME, Winder DG, Conn PJ. Contrasting sex-dependent adaptations to synaptic
physiology and membrane properties of prefrontal cortex interneuron subtypes in a
mouse model of binge drinking. Neuropharmacology. 2020:108126.

43    Straiker A, Mackie K. Depolarization-induced suppression of excitation in murine autaptic hippocampal neurones. J Physiol. 2005;569(Pt 2):501-17.

44    Marcus DJ, Bedse G, Gaulden AD, Ryan JD, Kondev V, Winters ND, et al. Endocannabinoid Signaling Collapse Mediates Stress-Induced Amygdalo-Cortical Strengthening. Neuron. 2020;105(6):1062-76.e6.

45    Maksymetz J, Byun N, Luessen D, Li B, Barry WG, JC, Niswender C, et al. mGlu1 potentiation enhances cortical somatostatin interneuron activity to rescue schizophrenia-like

physiological and cognitive deficits. Cell Reports. 2021;37(5):109950.

Thompson SM, Fabian CB, Ferranti AS, Joffe ME. Acute alcohol and chronic drinking bidirectionally regulate the excitability of prefrontal cortex vasoactive intestinal peptide interneurons. Neuropharmacology. 2023;238:109638.

Mannaioni G, Marino MJ, Valenti O, Traynelis SF, Conn PJ. Metabotropic glutamate receptors 1 and 5 differentially regulate CA1 pyramidal cell function. J Neurosci. 2001;21(16):5925-34.

Di Menna L, Joffe ME, Iacovelli L, Orlando R, Lindsley CW, Mairesse J, et al. Functional partnership between mGlu3 and mGlu5 metabotropic glutamate receptors in the central nervous system. Neuropharmacology. 2018;128:301-13.

Gereau RWt, Conn PJ. Roles of specific metabotropic glutamate receptor subtypes in regulation of hippocampal CA1 pyramidal cell excitability. J Neurophysiol. 1995;74(1):122-9.

Wu S, Wright RA, Rockey PK, Burgett SG, Arnold JS, Rosteck PR, Jr., et al. Group III human metabotropic glutamate receptors 4, 7 and 8: molecular cloning, functional expression, and comparison of pharmacological properties in RGT cells. Brain Res Mol Brain Res. 1998;53(1-2):88-97.

Ferranti AS, Johnson KA, Winder DG, Conn PJ, Joffe ME. Prefrontal cortex parvalbumin interneurons exhibit decreased excitability and potentiated synaptic strength after ethanol reward learning. Alcohol. 2022;101:17-26.

Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C. Interneurons of the neocortical inhibitory system. Nat Rev Neurosci. 2004;5(10):793-807.

Kawaguchi Y, Kubota Y. GABAergic cell subtypes and their synaptic connections in rat frontal cortex. Cereb Cortex. 1997;7(6):476-86.

Maksymetz J, Byun NE, Luessen DJ, Li B, Barry RL, Gore JC, et al. mGlu1 potentiation enhances prelimbic somatostatin interneuron activity to rescue schizophrenia-like physiological and cognitive deficits. Cell Rep. 2021;37(5):109950.

Chu Z, Hablitz JJ. Activation of group I mGluRs increases spontaneous IPSC frequency in rat frontal cortex. J Neurophysiol. 1998;80(2):621-7.

56    Whittington MA, Traub RD, Jefferys JG. Synchronized oscillations in interneuron
networks driven by metabotropic glutamate receptor activation. Nature.
1995;373(6515):612-5.

57    Marek GJ, Zhang C. Activation of metabotropic glutamate 5 (mGlu5) receptors induces
spontaneous excitatory synaptic currents in layer V pyramidal cells of the rat prefrontal
cortex. Neurosci Lett. 2008;442(3):239-43.

58    Maejima T, Hashimoto K, Yoshida T, Aiba A, Kano M. Presynaptic inhibition caused by
retrograde signal from metabotropic glutamate to cannabinoid receptors. Neuron.
2001;31(3):463-75.

59    Varma N, Carlson GC, Ledent C, Alger BE. Metabotropic glutamate receptors drive the
endocannabinoid system in hippocampus. J Neurosci. 2001;21(24):Rc188.

60    Ohno-Shosaku T, Shosaku J, Tsubokawa H, Kano M. Cooperative endocannabinoid
production by neuronal depolarization and group I metabotropic glutamate receptor
activation. Eur J Neurosci. 2002;15(6):953-61.

61    Robbe D, Kopf M, Remaury A, Bockaert J, Manzoni OJ. Endogenous cannabinoids
mediate long-term synaptic depression in the nucleus accumbens. Proc Natl Acad Sci U S
A. 2002;99(12):8384-8.

62    Guo B, Wang J, Yao H, Ren K, Chen J, Yang J, et al. Corrigendum: Chronic Inflammatory
Pain Impairs mGluR5-Mediated Depolarization-Induced Suppression of Excitation in the
Anterior Cingulate Cortex. Cereb Cortex. 2019;29(4):1698.

63    Diana MA, Marty A. Endocannabinoid-mediated short-term synaptic plasticity:
depolarization-induced suppression of inhibition (DSI) and depolarization-induced
suppression of excitation (DSE). Br J Pharmacol. 2004;142(1):9-19.

64    Kasten CR, Holmgren EB, Wills TA. Metabotropic Glutamate Receptor Subtype 5 in
Alcohol-Induced Negative Affect. Brain Sci. 2019;9(8).

65    Jury NJ, DiBerto JF, Kash TL, Holmes A. Sex differences in the behavioral sequelae of
chronic ethanol exposure. Alcohol. 2017;58:53-60.

66    Finn DA, Helms ML, Nipper MA, Cohen A, Jensen JP, Devaud LL. Sex differences in the
synergistic effect of prior binge drinking and traumatic stress on subsequent ethanol
intake and neurochemical responses in adult C57BL/6J mice. Alcohol. 2018;71:33-45.

67    Becker HC, Lopez MF. Increased ethanol drinking after repeated chronic ethanol
exposure and withdrawal experience in C57BL/6 mice. Alcohol Clin Exp Res.
2004;28(12):1829-38.

68    Clemens AM, Lenschow C, Beed P, Li L, Sammons R, Naumann RK, et al. Estrus-Cycle
Regulation of Cortical Inhibition. Curr Biol. 2019;29(4):605-15 e6.

69    Saugstad JA, Marino MJ, Folk JA, Hepler JR, Conn PJ. RGS4 inhibits signaling by group I
metabotropic glutamate receptors. J Neurosci. 1998;18(3):905-13.

70 Cannady R, McGonigal JT, Newsom RJ, Woodward JJ, Mulholland PJ, Gass JT. Prefrontal
Cortex KCa2 Channels Regulate mGlu5-Dependent Plasticity and Extinction of Alcohol-
Seeking Behavior. J Neurosci. 2017;37(16):4359-69.

71 Kim SJ, Kim YS, Yuan JP, Petralia RS, Worley PF, Linden DJ. Activation of the TRPC1 cation
channel by metabotropic glutamate receptor mGluR1. Nature. 2003;426(6964):285-91.

72 Topolnik L, Azzi M, Morin F, Kougioumoutzakis A, Lacaille JC. mGluR1/5 subtype-specific
calcium signalling and induction of long-term potentiation in rat hippocampal
oriens/alveus interneurones. J Physiol. 2006;575(Pt 1):115-31.

73 Trabzuni D, Ramasamy A, Imran S, Walker R, Smith C, Weale ME, et al. Widespread sex
differences in gene expression and splicing in the adult human brain. Nat Commun.
2013;4:2771.

74 Francesconi A, Duvoisin RM. Alternative splicing unmasks dendritic and axonal targeting
signals in metabotropic glutamate receptor 1. J Neurosci. 2002;22(6):2196-205.

75 Naito R, Kassai H, Sakai Y, Schönherr S, Fukaya M, Schwarzer C, et al. New Features on
the Expression and Trafficking of mGluR1 Splice Variants Exposed by Two Novel Mutant
Mouse Lines. Front Mol Neurosci. 2018;11:439.

76 Lee J, Munguba H, Gutzeit VA, Singh DR, Kristt M, Dittman JS, et al. Defining the Homo-
and Heterodimerization Propensities of Metabotropic Glutamate Receptors. Cell Rep.
2020;31(5):107605.

77 Doumazane E, Scholler P, Zwier JM, Trinquet E, Rondard P, Pin JP. A new approach to
analyze cell surface protein complexes reveals specific heterodimeric metabotropic
glutamate receptors. FASEB J. 2011;25(1):66-77.

78 Notartomaso S, Nakao H, Mascio G, Scarselli P, Cannella M, Zappulla C, et al. mGlu1
Receptors Monopolize the Synaptic Control of Cerebellar Purkinje Cells by Epigenetically
Down-Regulating mGlu5 Receptors. Sci Rep. 2018;8(1):13361.

79 Sevastyanova TN, Kammermeier PJ. Cooperative signaling between homodimers of
metabotropic glutamate receptors 1 and 5. Mol Pharmacol. 2014;86(5):492-504.

80 Ferraro A, Wig P, Boscarino J, Reich CG. Sex differences in endocannabinoid modulation
of rat CA1 dendritic neurotransmission. Neurobiol Stress. 2020;13:100283.

81 Rodríguez de Fonseca F, Cebeira M, Ramos JA, Martín M, Fernández-Ruiz JJ. Cannabinoid
receptors in rat brain areas: sexual differences, fluctuations during estrous cycle and
changes after gonadectomy and sex steroid replacement. Life Sci. 1994;54(3):159-70.

82 Bara A, Manduca A, Bernabeu A, Borsoi M, Serviado M, Lassalle O, et al. Sex-dependent
effects of in utero cannabinoid exposure on cortical function. Elife. 2018;7.

83 Bernabeu A, Bara A, Murphy Green MN, Manduca A, Wager-Miller J, Borsoi M, et al.
Sexually Dimorphic Adolescent Trajectories of Prefrontal Endocannabinoid Synaptic
Plasticity Equalize in Adulthood, Reflected by Endocannabinoid System Gene Expression.
Cannabis Cannabinoid Res. 2023.

84 Backstrom P, Bachteler D, Koch S, Hyytia P, Spanagel R. mGluR5 antagonist MPEP reduces
ethanol-seeking and relapse behavior. Neuropsychopharmacology. 2004;29(5):921-8.

85 Bird MK, Kirchhoff J, Djouma E, Lawrence AJ. Metabotropic glutamate 5 receptors
regulate sensitivity to ethanol in mice. Int J Neuropsychopharmacol. 2008;11(6):765-74.

86 Stevenson RA, Hoffman JL, Maldonado-Devincci AM, Faccidomo S, Hodge CW. MGluR5
activity is required for the induction of ethanol behavioral sensitization and associated
changes in ERK MAP kinase phosphorylation in the nucleus accumbens shell and lateral
habenula. Behav Brain Res. 2019;367:19-27.

87 Besheer J, Grondin JJ, Cannady R, Sharko AC, Faccidomo S, Hodge CW. Metabotropic
glutamate receptor 5 activity in the nucleus accumbens is required for the maintenance
of ethanol self-administration in a rat genetic model of high alcohol intake. Biol
Psychiatry. 2010;67(9):812-22.

88 Besheer J, Faccidomo S, Grondin JJ, Hodge CW. Regulation of motivation to self-
administer ethanol by mGluR5 in alcohol-preferring (P) rats. Alcohol Clin Exp Res.
2008;32(2):209-21.

89 Dao NC, Brockway DF, Suresh Nair M, Sicher AR, Crowley NA. Somatostatin neurons
control an alcohol binge drinking prelimbic microcircuit in mice.
Neuropsychopharmacology. 2021.

90 Dao NC, Suresh Nair M, Magee SN, Moyer JB, Sendao V, Brockway DF, et al. Forced
Abstinence From Alcohol Induces Sex-Specific Depression-Like Behavioral and Neural
Adaptations in Somatostatin Neurons in Cortical and Amygdalar Regions. Front Behav
Neurosci. 2020;14:86.

91 Sicher AR, Starnes WD, Griffith KR, Dao NC, Smith GC, Brockway DF, et al. Adolescent
binge drinking leads to long-lasting changes in cortical microcircuits in mice.
Neuropharmacology. 2023;234:109561.

92 Joffe ME, Maksymetz J, Luschinger JR, Dogra S, Ferranti AS, Luessen DJ, et al. Acute
restraint stress redirects prefrontal cortex circuit function through mGlu(5) receptor
plasticity on somatostatin-expressing interneurons. Neuron. 2022;110(6):1068-83.e5.

93 Blurton-Jones M, Tuszynski MH. Estrogen receptor-beta colocalizes extensively with
parvalbumin-labeled inhibitory neurons in the cortex, amygdala, basal forebrain, and
hippocampal formation of intact and ovariectomized adult rats. J Comp Neurol.
2002;452(3):276-87.

94 Gross KS, Moore KM, Meisel RL, Mermelstein PG. mGluR5 Mediates
Dihydrotestosterone-Induced Nucleus Accumbens Structural Plasticity, but Not
Conditioned Reward. Front Neurosci. 2018;12:855.

95 Fagan MP, Ameroso D, Meng A, Rock A, Maguire J, Rios M. Essential and sex-specific
effects of mGluR5 in ventromedial hypothalamus regulating estrogen signaling and
glucose balance. Proc Natl Acad Sci U S A. 2020;117(32):19566-77.

96   Vandegrift BJ, Hilderbrand ER, Satta R, Tai R, He D, You C, et al. Estrogen Receptor alpha
Regulates Ethanol Excitation of Ventral Tegmental Area Neurons and Binge Drinking in
Female Mice. J Neurosci. 2020;40(27):5196-207.

97   Maguire JL, Stell BM, Rafizadeh M, Mody I. Ovarian cycle-linked changes in GABA(A)
receptors mediating tonic inhibition alter seizure susceptibility and anxiety. Nat
Neurosci. 2005;8(6):797-804.

98   Maguire J, Mody I. Neurosteroid synthesis-mediated regulation of GABA(A) receptors:
relevance to the ovarian cycle and stress. J Neurosci. 2007;27(9):2155-62.

99   Abramian AM, Comenencia-Ortiz E, Modgil A, Vien TN, Nakamura Y, Moore YE, et al.
Neurosteroids promote phosphorylation and membrane insertion of extrasynaptic
GABAA receptors. Proc Natl Acad Sci U S A. 2014;111(19):7132-7.

100  Conde K, Fabelo C, Krause WC, Propst R, Goethel J, Fischer D, et al. Testosterone Rapidly
Augments Retrograde Endocannabinoid Signaling in Proopiomelanocortin Neurons to
Suppress Glutamatergic Input from Steroidogenic Factor 1 Neurons via Upregulation of
Diacylglycerol Lipase-alpha. Neuroendocrinology. 2017;105(4):341-56.

**Figure 1. PV-INs from female mice are more excitable at baseline, and PV-INs from male mice are more responsive to mGlu$_{1/5}$ receptor stimulation**

**Figure 2. mGlu$_1$ and mGlu$_5$ receptors sex-specifically regulate synaptic strength and endocannabinoid plasticity in PV-INs**

**Figure 3. Cell type-specific genetic deletion of *Grm5* abrogates sex differences in PV-IN membrane physiology**

**Figure 4. Cell-type specific genetic deletion of *Grm5* abrogates sex differences in PV-IN synaptic physiology**

**Figure 5. Female but not male PV-mGlu$_5^{-/-}$ mice display reduced binge drinking**

**Figure S1. PV-INs display comparable resting membrane properties and basal synaptic strength across sexes**

**Figure S2. Dual application of MTEP and LY367385 attenuated DHPG currents and blocked increase in synaptic transmission in PFC PV-INs**

**Figure S3. Male PV-INs undergo greater depolarization following mGlu$_1$ and mGlu$_5$ receptor stimulation**

**Figure S4. Threshold DHPG (5 μM) did not affect the amplitude of evoked EPSCs on PV-INs**

**Figure S5. PV-mGlu$_5$$^{-/-}$ mice display comparable basal membrane properties and synaptic strength across sexes**

**Figure S6. Schematic depicting proposed roles of mGlu$_1$ and mGlu$_5$ receptor signaling in male and female PFC PV-INs**