Natural variation in yeast reveals multiple paths for acquiring higher stress resistance

Amanda N. Scholes[1,2], Tara N. Stuecker[1], Stephanie E. Hood[1], Cader J. Locke[1], and Jeffrey A. Lewis[1*]

[1] Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas, United States of America

[2] Interdisciplinary Graduate Program in Cell and Molecular Biology, University of Arkansas, Fayetteville, Arkansas, United States of America

 * Corresponding author:
Jeffrey A. Lewis
Department of Biological Sciences
University of Arkansas
850 W. Dickson St., SCEN 601
Fayetteville, AR 72701
E-mail: lewisja@uark.edu
Tel: +1.479.575.7740

Abstract

Organisms frequently experience environmental stresses that occur in predictable patterns and combinations. For wild *Saccharomyces cerevisiae* yeast growing in natural environments, cells may experience high osmotic stress when they first enter broken fruit, followed by high ethanol levels during fermentation, and then finally high levels of oxidative stress resulting from respiration of ethanol. Yeast have adapted to these patterns by evolving sophisticated "cross protection" mechanisms, where mild 'primary' doses of one stress can enhance tolerance to severe doses of a different 'secondary' stress. For example, in many yeast strains, mild osmotic or mild ethanol stresses cross protect against severe oxidative stress, which likely reflects an anticipatory response important for high fitness in nature. During the course of genetic mapping studies aimed at understanding the mechanisms underlying natural variation in ethanol-induced cross protection against H_2O_2, we found that a key H_2O_2 scavenging enzyme, cytosolic catalase T (Ctt1p), was absolutely essential for cross protection in a wild oak strain. This suggested the absence of other compensatory mechanisms for acquiring H_2O_2 resistance in that strain background under those conditions. In this study, we found surprising heterogeneity across diverse yeast strains in whether *CTT1* function was fully necessary for acquired H_2O_2 resistance. Some strains exhibited partial dispensability of *CTT1* when ethanol and/or salt were used as mild stressors, suggesting that compensatory peroxidases may play a role in acquired stress resistance in certain genetic backgrounds. We leveraged global transcriptional responses to ethanol and salt stresses in strains with different levels of *CTT1* dispensability, allowing us to identify possible regulators of these alternative peroxidases and acquired stress resistance in general. Ultimately, this study highlights how superficially similar traits can have different underlying molecular foundations and provides a framework for understanding the diversity and regulation of stress defense mechanisms.

Author Summary

Organisms in nature frequently experience environmental stress in predictable patterns. For example, during the summer months in temperate climates, the warmth of the morning sun gradually gives way to high afternoon temperatures. Organisms that can anticipate these predictable patterns to mobilize stress defenses would have an advantage in nature. One way organisms anticipate future stress is through 'cross protection,' where cells exposed to a mild dose of one stress gain the ability to survive an otherwise lethal dose of different stress. To better understand the molecular mechanisms that are responsible cross protection, we have been taking advantage of wild yeast strains that are either more resilient or more sensitive to stresses. During the course of this study, we found that strains with superficially similar levels of cross protection differ in the precise molecular mechanisms that underlie the trait. Our study suggests that different molecular strategies may be important for yielding similar stress resistances under different environmental constraints, and highlights the power of harnessing natural genetic diversity to understand the molecular mechanisms underlying differences in environmental responses.

Introduction

In nature, environmental stress often occurs in predictable patterns. Extremes of temperature and precipitation follow seasonal patterns, and stress gradients can also predictably occur over even shorter timescales. During the daytime, temperature predictably increases, while in estuary environments, and salinity predictably increases closer to seawater sources. In addition to single-stress gradients, some stresses tend to co-occur or happen in predictable succession. For example, high temperatures frequently coincide with low precipitation, so organisms often experience both heat and drought stresses simultaneously.

Because stresses can occur in predictable patterns, organisms that can anticipate these patterns would likely have a fitness advantage in nature. One such anticipatory strategy is a phenomenon called acquired stress resistance, where exposure to a mild dose of stress can enable organisms to survive an otherwise lethal dose of severe stress. Acquired stress resistance can occur when the mild and severe stresses are the same (same-stress protection) or are different (cross protection). Cross protection has been observed in diverse organisms ranging from prokaryotes to humans under a wide variety of circumstances. For example, mild stresses commonly used in food processing can cross protect against antibiotics in the foodborne pathogen *Listeria monocytogenes* [1]. Mild drought pretreatment protects several crop plants against severe heat stress [2-4], which has major implications for mitigating the effects of climate change. In addition to implications for biotechnology and agriculture, cross protection has a myriad of implications for human health. For instance, fever temperatures protect against oxidative stress, which may be important for immune cells experiencing increased generation of reactive oxygen species during infection [5]. Additionally, transient whole-body hyperthermia (i.e., a mild heat shock) cross protects against oxidative damage caused by reperfusion of oxygenated blood following surgery [6, 7]. Therefore, understanding the molecular mechanisms underlying acquired stress resistance and cross protection has wide-ranging applications.

While a large number of genetic screens have been performed to understand the genetic basis of the intrinsic resistance of unstressed cells [8-36], only a more limited number of studies have investigated acquired stress resistance [37-41]. Consequently, the cellular strategies and molecular mechanisms underlying acquired stress resistance and cross protection remain largely obscure. Additionally, while cross protection is widespread in yeast, it is not universal for all stress combinations. Berry and colleagues found that mild heat stress protected cells against all severe stresses tested (NaCl, H_2O_2, ethanol, and heat), while mild NaCl only protected against severe NaCl and severe H_2O_2 [42]. The observed patterns of stress cross protection likely mirror the sequential order in which organisms typically encounter stresses in their natural environments [43]. However, conflicting reports about which stresses can or cannot confer cross protection within a given species are common (reviewed in [44]). At least for the case of the yeast ethanol response, we have shown that cross protection phenotypes can differ dramatically depending on the genetic background of the strain [45-47]. While perhaps not surprising that the genetic context affects a complex trait such as cross protection, it does suggest that different genetically distinct populations of yeast may have different anticipatory responses that reflect specific selective pressures unique to their ecological niches.

Most of our limited knowledge about the mechanisms underlying acquired stress resistance in yeast is restricted to laboratory strains [37, 39, 40]. However, heterogeneity in acquired resistance phenotypes clearly occurs and can have broad impact. For example, cancer cells appear to have defective acquired stress resistance phenotypes, which can be targeted to enhance the effectiveness of chemotherapy while sparing healthy cells [48, 49]. We have been leveraging natural variation in acquired stress resistance phenotypes to better understand both

the genetic architecture of the trait and especially the stress defense genes ultimately responsible for the phenotype.

We previously reported that our commonly-used laboratory yeast strain (S288c) fails to acquire further H_2O_2 resistance when pre-treated with a mild dose of ethanol, while many wild yeast strains can [47]. To understand the genetic basis of this trait, we performed genetic mapping in a cross between the lab strain and a wild oak strain (YPS163). This analysis identified the causal polymorphism—a transposon insertion in the lab strain that partially inactivates the heme-activated transcription factor Hap1p, leading to decreased ethanol-responsive expression of a key H_2O_2 scavenging enzyme, cytosolic catalase T (Ctt1p) [47]. In that wild oak strain, a $ctt1\Delta$ mutation completely abolished ethanol-induced cross protection against H_2O_2, suggesting the absence of compensatory mechanisms for acquired H_2O_2 resistance under that condition.

In this study, we found surprising heterogeneity across diverse yeast strains in whether *CTT1* function was fully necessary for acquired H_2O_2 resistance. Some strains exhibited partial dispensability of *CTT1* when ethanol and/or salt were used as mild stressors, suggesting that compensatory peroxidases may play a role in acquired stress resistance in certain genetic backgrounds. We leveraged global transcriptional responses to ethanol and salt stresses in strains with different levels of *CTT1* dispensability, allowing us to identify possible regulators of these alternative peroxidases and acquired stress resistance in general. Ultimately, this study highlights how superficially similar traits can have different underlying molecular foundations and provides a framework for understanding diversity and regulation of stress defense mechanisms.

Results

Strain-specific catalase (Ctt1p) requirements for acquired oxidative stress resistance.

While a commonly-used S288c-derived lab strain of yeast is unable to acquire further H_2O_2 resistance following mild ethanol pretreatment, most wild yeast strains can [47]. Experiments designed to identify the mechanistic basis of this difference led to the discovery that cytosolic catalase T (Ctt1p)—a major H_2O_2-scavenging enzyme—plays a key role. Specifically, loss of *CTT1* function in a wild oak strain (YPS163) completely abolished ethanol-induced cross protection against H_2O_2 [47].

We initially suspected the involvement of *CTT1* in ethanol-induced cross protection because a previous study found that *CTT1* was necessary for salt-induced cross protection against H_2O_2 in the S288c lab strain [39]. Thus, when analyzing the phenotype of YPS163 $ctt1\Delta$ mutants for ethanol-induced cross protection, we included salt-induced cross protection as a "positive control." Surprisingly, YPS163 $ctt1\Delta$ mutants retained substantial acquired H_2O_2 resistance following mild salt pretreatment (~25% of wild-type resistance or an additional 1mM of resistance compared to the mock-treated control, Fig 1). In contrast, and consistent with previous studies [39], no residual acquired resistance was detected in the S288c background following mild salt pretreatment. Consistent with our previous studies [47], we did observe that mild ethanol pretreatment renders S288c even more sensitive to H_2O_2. Surprisingly, deletion of *CTT1* in S288c eliminated the sensitizing effects of ethanol pretreatment on H_2O_2 resistance. Overall, the results from this experiment suggested that *CTT1* was conditionally necessary for acquired H_2O_2 resistance depending on the identity of the primary stress and the genetic background (i.e., a gene-environment (GxE) interaction), which we sought to explore further.

To better understand the scope of *CTT1* conditional dependency, we deleted *CTT1* in a panel of yeast strains from diverse ecological niches (S1 Table), and then assayed acquired H_2O_2 resistance following either mild ethanol or salt pretreatments. With the exception of a wild coconut strain (Y10), lack of *CTT1* had little to no effect on intrinsic (basal) levels of H_2O_2

tolerance in the absence of pretreatment (Fig 2 and Fig S1). For cross protection however, we found that *CTT1* was partially dispensable in approximately half of tested strains (6/13), with half of those displaying *CTT1*-independent cross protection in only one condition (always salt-induced cross protection) (Fig 2). Hierarchical clustering of the tolerance scores showed that oak and vineyard strains partitioned together (Fig 2B), potentially providing another example of trait differentiation between oak and vineyard populations [50], though larger sample sizes are certainly needed to determine whether this observation is generalizable. To understand whether the lack of ethanol-induced cross protection against H_2O_2 was correlated with *CTT1* dependency for salt-induced cross protection, we included two strains that, similar to S288c, displayed low levels of acquired H_2O_2 resistance following mild ethanol-pretreatment [47]. *CTT1* was partially dispensable for salt-induced cross protection in one of the two strains, suggesting that the absence of ethanol-induced cross protection was uncorrelated with *CTT1* dispensability when salt was the primary stress. In summary, *CTT1* dispensability demonstrates complex patterns of GxE interactions across genetically diverse wild yeast strains, while suggesting the existence of multiple molecular pathways and/or mechanisms that can lead to superficially similar phenotypes (in this case maximal acquired stress resistance).

Extensive variation in the ethanol and salt responses across yeast strains.
        Stress-activated gene expression changes are necessary for acquired stress resistance [42, 45], and natural variation in cross protection has been linked to gene expression variation [47]. Therefore, we performed transcriptional profiling in a subset of strains with different levels of *CTT1* dispensability to identify differentially expressed genes and their potential transcriptional regulators that may be important for the phenotypic differences in cross protection. We included two each of strains with poor ethanol-induced acquisition (S288c and YJM1129), *CTT1* indispensability for both NaCl and ethanol pretreatments (M1 and Y10), *CTT1* dispensability for only NaCl pretreatment (YPS163 and YPS606), and *CTT1* dispensability for both NaCl and ethanol (M22 and YJM308). RNA-seq was performed on these strains comparing an unstressed control to cells treated with either 5% v/v ethanol for 30 minutes or 0.4 M NaCl for 45 minutes (timepoints which encompass the peak response for each stressor [42, 45]).
        Differential expression analysis was performed by comparing the $\log_2$ fold changes during stress for each strain to the mean expression for all 8 strains. Out of the 6,092 yeast genes expressed in at least one condition (cpm $\geq$ 1), 4,261 genes (69%) showed significant differences in ethanol-responsive expression in at least one strain compared to the mean of all strains, while 3,197 (54%) were differentially expressed in at least one strain during the NaCl response. To identify modules of co-regulated genes that differ in expression across strains, we performed hierarchical clustering on both strains and significantly differentially-expressed genes (see Methods). For ethanol stress, strains with poor overall cross protection (S288c and YJM1129) were clear outliers. S288c in particular had lower induction of stress defense-enriched clusters (Fig 3A). While YJM1129 was also an outlier compared to other strains, lower overall induction of stress defense genes was not generally observed. Moreover, while the lack of ethanol-induced cross protection in S288c is likely due to reduced *CTT1* mRNA expression (12-fold lower relative abundance during ethanol stress compared to the mean of all strains), this is likely not true of YJM1129 (which had only 1.5-fold lower relative mRNA abundance during ethanol stress compared to the mean of all strains), suggesting a different mechanistic basis for the absence of ethanol-induced cross protection in this strain (see Discussion).
        Strains did not generally cluster together based on their *CTT1* dispensability phenotype, suggesting a weak overall correlation between stress-responsive gene expression and these specific acquired resistance phenotypes. This is perhaps not surprising, considering the complexity of genomic responses to stress and their likely effects on multiple traits [51, 52]. However, strain-specific differences in stress-responsive expression were clearly apparent. For example, YPS163, YPS606, Y10, and M22 all had higher expression for salt-induced clusters

that were functionally enriched for stress related processes such as in trehalose metabolism (P = 9 x 10^{-6}), cell wall organization (P = 2 x 10^{-5}), and response to oxidative stress (P = 7 x 10^{-4}). Likewise, M1 had decreased expression for those same clusters, while having novel or amplified induction during salt stress for a cluster functionally enriched for genes involved in the cell cycle (P = 5 x 10^{-11}), protein glycosylation (P = 2 x 10^{-5}), mRNA transport (P = 8 x 10^{-4}), and regulation of the response to stress (P = 1 x 10^{-3}).

To explore strain-specific differences in stress responsive gene expression further, we identified genes in each strain that differed from the mean response of all strains (i.e., the 'consensus' response, see Methods), and separated them into classes (e.g., novel induction, amplified repression, Fig 4). As predicted from the clustering and consistent with our previous studies comparing S288c to a smaller panel of strains [45, 53], S288c had the largest number of genes with divergent expression compared to the consensus for ethanol stress, with ~600 genes being less induced than the consensus response (enriched for carbohydrate metabolism (P = 4 x 10^{-24}), response to chemical (P = 3 x 10^{-5}), and trehalose metabolism (P = 6 x 10^{-5})), and ~700 genes being less repressed than the consensus response (enriched for ribosome biogenesis (P = 9 x 10^{-151}) and cytoplasmic translation (P = 1 x 10^{-25})). Interestingly, YJM1129, which does not acquire further H$_2$O$_2$ resistance following ethanol pre-treatment, did not have a particularly high number of genes (107) with lower induction during ethanol stress, though there was a small enrichment for response to environmental stimulus (P = 2 x 10^{-4}). M1, despite not having a discernable defect in acquired stress resistance under the conditions tested, had a large number (~400) of salt-induced genes with lower induction than consensus, and these were enriched for carbohydrate metabolism (P = 2 x 10^{-17}), trehalose metabolism (P = 7 x 10^{-9}), and response to chemical (P = 9 x 10^{-3}).

We were particularly interested in strains with amplified stress-induced expression compared to all strains, as this may indicate a heightened response. Surprisingly, despite being deficient in ethanol-induced cross protection against H$_2$O$_2$, YJM1129 had the most genes (300) with amplified induction during ethanol stress, which were enriched for small molecule catabolism (P = 1 x 10^{-8}), cell wall organization (P = 3 x 10^{-4}), and respiration (P = 4 x 10^{-3}). For salt stress, the oak strains showed the most amplified induction (313 genes for YPS163 and 348 for YPS606), with the vast majority of those genes being shared between the two strains (258), which is unsurprising considering the similarity of their expression responses. Shared genes with amplified induction in YPS163 and YPS606 during salt stress were enriched for carbohydrate metabolism (P = 2 x 10^{-12}), trehalose metabolism (P = 1 x 10^{-8}) and response to chemical (P = 3 x 10^{-3}). We hypothesize that a subset of those genes with amplified induction are responsible for the residual salt-induced acquired H$_2$O$_2$ resistance in the absence of *CTT1* in the oak strain background (see below).

Identification of regulators necessary for ethanol or salt induced cross protection against H$_2$O$_2$.

To identify potential regulators of the salt and ethanol responses that may be responsible for acquired hydrogen peroxide resistance, we looked for regulatory enrichments for genes belonging to stress-induced clusters (Fig 3 and S6 Table). Prioritizing based on regulatory enrichment as well as transcription factors with known functions in relevant stressors (ethanol, osmotic, and/or oxidative stresses), we generated eight transcription factor gene deletions in the YPS606 strain background (*hot1Δ, HSF1/hsf1Δ, msn1Δ, msn2Δmsn4Δ, skn7Δ, sko1Δ, smp1Δ, yap1Δ*; see Table 1 for a brief description of each transcription factor), and tested whether those regulators were necessary for cross protection. We also tested the transcription factor mutations in combination with *ctt1Δ* mutations, to determine whether the tested transcription factors were potential regulators of the putative alternative peroxidases necessary for *CTT1*-independent acquisition. We chose YPS606 as a representative strain for further genetic dissection for two reasons. First, YPS606 possessed the highest level of residual

acquisition in the absence of *CTT1*, providing the largest dynamic range for detecting loss of *CTT1*-independent acquisition. Second, we reasoned that because *CTT1* is only partially dispensable for salt-induced cross protection and not for ethanol, comparison of expression profiles for potential compensatory peroxidases during both ethanol and salt stress may help implicate the responsible party.

With ethanol as the pre-treatment, the mutant with the largest defect in acquired resistance was the *msn2Δ msn4Δ* double mutant (Fig 5A), where cross protection was completely abolished. The *skn7Δ* mutant had an intermediate defect, as well as a slight defect for intrinsic (basal) H_2O_2 resistance. The *yap1Δ* mutant had a strong intrinsic H_2O_2 resistance defect, but was still able to fully acquire resistance, suggesting that low intrinsic stress resistance does not necessarily preclude maximal acquisition.

For salt as the mild pretreatment, none of the tested mutants showed a defect in acquired H_2O_2 resistance when *CTT1* was present (Fig 5 and S2 Fig), which was especially striking considering that *msn2Δ msn4Δ* double mutants have extremely low salt-induced cross protection against H_2O_2 in the S288c background [39, 42]. This is likely due to partial regulatory redundancy in the YPS606 strain background (see below). Intriguingly, both the *yap1Δ ctt1Δ* and *skn7Δ ctt1Δ* double mutants had significantly reduced levels of *CTT1*-independent acquisition (Fig 5B), suggesting that both Yap1p and Skn7p play a role in regulating those compensatory functions. Because low intrinsic stress resistance does not preclude the ability to acquire high levels of stress resistance, it was somewhat striking that both the *skn7Δ ctt1Δ* and especially the *yap1Δ ctt1Δ* had little acquisition beyond that of the intrinsic resistance of the single *skn7Δ* or *yap1Δ* mutants. This suggests that the peroxidases necessary for intrinsic resistance may overlap with those that partially compensate in the absence of Ctt1p. None of the other transcription factors tested (*HOT1, HSF1, MSN1, SKO1, or SMP1*) showed a defect in acquired resistance under any condition, though *HSF1* comes with the caveat that we were only able to test a heterozygous deletion due to *HSF1* being essential.

Transcriptional profiling reveals the overlapping and unique roles of the Msn2/4p, Skn7p, and Yap1p transcription factors during either ethanol or salt stress.

We next performed expression profiling on transcription factor mutants in the YPS606 background that had an acquired stress resistance defect (either in isolation or in combination with *ctt1Δ*) for both mild ethanol and mild salt stresses, as well as for an unstressed control: *msn2Δ msn4Δ*, *skn7Δ*, and *yap1Δ*. For the ethanol response, the *msn2Δ msn4Δ* mutant had the largest number of genes with significantly defective induction (i.e., reduced induction relative to wild type)—379 genes. In contrast, 95 genes showed defective induction in the *skn7Δ* mutant, while only 28 genes showed defective induction in the *yap1Δ* mutant (Fig 6 and S5 Table). We looked at the overlap between the induced ethanol response regulon for each transcription factor pair plus all three, as well as the unique regulons. Eight genes were part of the ethanol-induced regulon for all 3 transcription factors, and included four genes involved in protein folding (*AHA1, HSP78, SIS1, ERO1;* enrichment $P = 4 \times 10^{-4}$), as well as genes encoding an alcohol dehydrogenase (*ADH5*), a meiotic regulator *(WTM1)*, a hexokinase (*EMI2*), and stress-responsive protein of unknown function (*HOR7*). There were 22 genes in the overlapping *SKN7-YAP1* regulons, which were enriched for amino acid metabolism ($P = 1 \times 10^{-3}$). There was no significantly enriched functional category for the genes that were part of the overlapping *SKN7* and *MSN2/4* regulons, though *CTT1* was one of those genes. In fact, ethanol-responsive fold-changes in *CTT1* expression correlated well with the total (*msn2Δ msn4Δ*) and partial (*skn7Δ*) cross protection defects in those mutants—*CTT1* showed no induction (1.3-fold) in the *msn2Δ msn4Δ* mutant and 32-fold induction in the *skn7Δ* mutant (compared to 91-fold induction in the wild-type strain). In contrast, *CTT1* was 136-fold induced in the *yap1Δ* strain, which had wild-type levels of acquired resistance, suggesting that Yap1p plays no role in *CTT1* regulation during ethanol stress. Finally, there were 325 genes that were unique to the *MSN2/4* ethanol

regulon, and these were enriched for stress related functions such as metabolism of trehalose (P = 6 x 10^{-6}) and glycogen (P = 3 x 10^{-4}), and the response to oxidative stress (P = 5 x 10^{-3}).

In contrast with the ethanol response, *SKN7* had the largest salt-induced regulon (388 genes), while *MSN2/4* and *YAP1* had moderately sized regulons (162 and 132 genes, respectively). There were 43 genes with that were part of the salt-induced regulons for all three transcription factors, and these were enriched for response to stress (P = 5 x 10^{-3}), protein refolding (P = 1 x 10^{-9}), and trehalose metabolism (P = 9 x 10^{-9}). *CTT1* was part of this overlapping group, though the *CTT1* is much more strongly induced by salt than ethanol in the wild-type strain (340-fold vs. 91-fold) and still possessed relatively high induction in all three mutants (63-fold for *msn2Δ msn4Δ*, 94-fold for *skn7Δ*, and 63-fold for *yap1Δ*). *CTT1* expression thus appears combinatorially controlled by Msn2/4p, Skn7p, and Yap1p, which would explain why cells were still able to acquire resistance in the absence of any single transcription factor.

While the overlap in regulons was more substantial for salt, each transcription factor also regulated at least 20 genes uniquely. The 212 genes that were unique to the *SKN7* salt-induced regulon were enriched for cell wall organization (P = 8 x 10^{-8}), the 50 genes unique to the *MSN2/4* salt-induced regulon were enriched for responses to chemical (P = 1 x 10^{-3}) and oxidative (P = 2 x 10^{-3}) stresses, and the 21 genes unique to the *YAP1* salt regulon were enriched for proteasome assembly (P = 2 x 10^{-5}) and regulation of proteolysis (P = 4 x 10^{-4}). There were 110 genes in the overlapping *SKN7-MSN2* regulons, which were enriched for the response to stress (P = 3 x 10^{-3}), and glycogen (P = 6 x 10^{-7}) and trehalose (P = 1 x 10^{-14}) metabolism. There were 22 genes in the overlapping *SKN7-YAP1* regulons, which were enriched for proteasome assembly (P = 3 x 10^{-4}). Finally, there were no shared functionally enriched categories for the shared *MSN2-YAP1* regulons.

For intrinsic (basal) gene expression, there were 197 genes with differential expression compared to wild type in the *yap1Δ* mutant, 369 genes in the *skn7Δ* mutant, and 10 genes in the *msn2Δ msn4Δ* mutant (Table S5). For the *yap1Δ* strain, 64 genes had at least 1.5-fold higher expression relative to the wild-type strain (in other words, were de-repressed), and these genes were enriched for amino acid biosynthesis (P = 7 x 10^{-12}). Fifty-six genes had >1.5-fold lower expression in the *yap1Δ* strain (i.e., defective activation), which were importantly enriched for cellular oxidant detoxification (P = 3 x 10^{-8}). This includes genes that encode for proteins known to be involved in directly detoxifying H_2O_2 (*GPX2, PRX1, GRX2, AHP1, TSA1*) or genes necessary for proper function of those systems (*GSH1* involved in glutathione biosynthesis and *TRR1* encoding a thioredoxin reductase). Decreased expression of these genes likely explains the strong intrinsic H_2O_2 tolerance defect of the *yap1Δ* strain. The *skn7Δ* mutant, which also had an intrinsic H_2O_2 tolerance defect, had 44 genes with >1.5-fold lower expression than wild type, and these genes were also enriched for cellular oxidant detoxification (P = 4 x 10^{-4}), including those encoding direct H_2O_2 detoxifiers (*PRX1, GPX2, TSA1, AHP1*). There was strong and significant overlap between genes with either lower expression in both the *yap1Δ* and *skn7Δ* strains (44-fold over-enrichment and P = x 10^{-26}, Fisher's exact test) or higher expression (31-fold over-enrichment and P = 6 x 10^{-77}, Fisher's exact test), suggesting substantial overlap for the two regulons. Genes with >1.5-fold higher expression in both the *yap1Δ* and *skn7Δ* mutants relative to wild type were enriched for amino acid biosynthesis (P = 2 x 10^{-13}). The 99 genes with higher expression in the *skn7*mutant but not *yap1*mutant were enriched for those involved in iron ion transport (P = 3 x 10^{-5}).

CTT1-independent acquisition requires glutathione.

Finally, we sought to understand the mechanistic basis for partial *CTT1* dispensability. Leveraging our transcriptomic data, we focused on the expression of genes annotated to have any kind of peroxidase activity. We clustered just the expression of those genes (Fig 7), which revealed interesting patterns of differential expression across strains where *CTT1* was either dispensable or indispensable for acquisition. Clustering on strains for just this subset of genes

better matched acquired resistance phenotypes, particularly with ethanol, where S288c clearly showed reduced expression of most peroxidases in addition to *CTT1*, while the strains where *CTT1* was partially dispensable for ethanol (M22 and YJM308) clustered together and showed higher expression for most peroxidase genes. For salt, strains where *CTT1* was indispensable for acquisition (M1 and Y10, plus S288c and YJM1129) tended to have lower than average induction of peroxidase genes, with S288c and YJM1129 in particular clustering together.

For all non-*CTT1* peroxidase genes that were significantly induced at least 2-fold in any strain, expression was higher in strains where *CTT1* was dispensable (1.9-fold higher induction for ethanol, $P < 4 \times 10^{-4}$, t-test, and 1.6-fold higher for salt, $P < 4 \times 10^{-3}$, t-test), suggesting potentially redundant mechanisms. Because there were multiple, potentially redundant candidate genes that could be responsible for *CTT1*-independent acquisition, we sought to narrow the possible mechanisms. In addition to catalases, the major cellular peroxidases require reduction by either thioredoxin or glutathione to continue scavenging H_2O_2. Intriguingly, genes encoding glutathione-dependent peroxidases (*GRX2, GPX1, GPX2*) were among the those with the largest differences in expression between *CTT1*-dispensable and *CTT1*-indispensible strains during either ethanol or salt stresses. Additionally, the YPS606 *skn7Δ* mutant, which lacks *CTT1*-indepdent acquisition, also showed defective induction during salt stress for several glutathione-dependent peroxidase genes (*GPX1, GRX2, GRX1*). And while genes encoding glutathione-dependent peroxidases did not have lower expression in the *yap1Δ* mutant during salt stress, we did notice that expression of the gene that encodes the first enzymatic step of glutathione biosynthesis (*GSH1*) was 1.9-fold lower, suggesting that the *yap1Δ* mutant may have insufficient glutathione for those peroxidases to properly function.

Altogether, these data suggested that the peroxidase(s) responsible for *CTT1*-independent acquisition may be glutathione dependent. Thus, we tested the effects of loss of *GSH1* on acquired stress resistance, both alone and in combination with *ctt1Δ* in one strain that possesses *CTT1*-independent acquisition following either ethanol or salt pretreatments (M22) or two strains that only possess *CTT1*- independent acquisition following salt pretreatment (YPS163 and YPS606). Each *gsh1Δ* single mutant was able to fully acquire hydrogen peroxide resistance for either ethanol or salt pretreatments (Fig 8), suggesting that *GSH1* is not necessary for acquisition in the presence of *CTT1*. However, lack of *GSH1* significantly reduced *CTT1*-independent acquisition for M22 under both stress pretreatments (Fig 8), as well as for YPS163 and YPS606 during salt pretreatment (Fig 8B). For M22, the *gsh1Δ ctt1Δ* mutant also had an intrinsic H_2O_2 resistance defect in addition to the loss of *CTT1*- independent acquisition, suggesting that the compensatory peroxidases and *CTT1* may play a role in both processes. Altogether, these data support that glutathione-dependent peroxidases are at least partially responsible for the conditional dependency of *CTT1* in the tested wild strain backgrounds.

Discussion

In this study, we leveraged natural variation in acquired stress resistance to understand the mechanistic underpinnings of this important trait. We previously showed that cytosolic catalase T (Ctt1p) was completely essential for ethanol-induced cross protection in YPS163, a wild oak strain [47]. Here, we were surprised to find moderate residual salt-induced cross protection in the YPS163 *ctt1Δ* mutant, and that moreover, about half of the strains tested showed varied levels of *CTT1* dispensability. We leveraged transcriptional profiling in strains with varied levels of *CTT1* dispensability, which allowed us to both identify novel regulators important for acquired stress resistance, as well as potential compensatory mechanisms that lead to moderate acquisition in the absence of *CTT1*. This highlights the power of integrating comparative genomics with targeted mutational analysis to understand the mechanisms underlying stress defense traits.

Our analysis of regulatory mutants in the YPS606 oak strain background revealed interesting regulatory aspects for both intrinsic (basal) stress resistance and acquired stress resistance. For example, and consistent with the previous literature [54, 55], the YPS606 *yap1Δ* mutant had a severe defect in intrinsic H_2O_2 resistance. However, this mutant was able to fully acquire H_2O_2 with either ethanol or salt as pre-treatments, providing further support for the idea that the stress defense mechanisms underlying intrinsic and acquired stress resistance are distinct [37, 39, 40, 42, 45, 47]. The same was true of the YPS606 *skn7Δ* mutant, though its intrinsic H_2O_2 resistance defect was not as pronounced as that of the YPS606 *yap1Δ* mutant. In contrast, the YPS606 *msn2Δ msn4Δ* mutant had no observable defect for intrinsic H_2O_2 resistance, but had a strong defect for acquired H_2O_2 resistance when ethanol was the primary stress. Moreover, the intrinsic H_2O_2 resistance phenotypes of the tested transcription factor mutants correlated with differences in basal expression of genes annotated to the GO category of 'response to oxidative stress.' The YPS606 *msn2Δ msn4Δ* mutant, which had wild-type levels of intrinsic H_2O_2 resistance, also had no differentially expressed oxidative stress response genes under unstressed conditions. In contrast, the YPS606 *skn7Δ* mutant, with a moderate intrinsic H_2O_2 resistance defect, had four oxidative stress response genes with at least 2-fold lower basal expression than wild type: *AHP1, GPX2, PRX1*, and *TSA1*. And the YPS606 *yap1Δ* mutant, with a strong intrinsic H_2O_2 resistance defect, also possessed at least 2-fold lower expression of those same four genes, while also including three additional genes: *GSH1*, *HSP31*, and *SOD1*. Notable, *gsh1Δ* mutants in the M22, YPS163, YPS606, had little difference in intrinsic H_2O_2, suggesting that a combination of Yap1p-regulated thioredoxin- and glutathione-dependent peroxidases are responsible for intrinsic H_2O_2 tolerance in this particular yeast strain.

Our results also demonstrate that different yeast strains use different cellular strategies to yield superficially similar stress resistance phenotypes. For example, we showed that the paralogous general transcription factors Msn2/4p are essential in the YPS606 background for ethanol-induced cross protection against H_2O_2, but not salt-induced cross protection. This was somewhat surprising to us because Msn2/4p are necessary for salt-induced cross protection in the S288c background [39, 42], and highlights the importance of comparing different genetic backgrounds to obtain an integrated view of stress signaling within even a single species. While *MSN2* has been implicated in acquired ethanol (same stress) resistance [45], this is the first time it has been shown to play a role in ethanol-induced cross protection. We also found a partial defect in ethanol-induced cross protection in the YPS606 *skn7Δ* background, which has never been observed before, further highlighting the benefits of taking advantage of natural variation to identify novel biology even in well-studied model organisms [56]. Our results also provide hints about the complexity of regulation of stress responses beyond transcriptional responses. For example, YJM1129 is unable to acquire H_2O_2 resistance when ethanol is the primary stress, yet it had had only 1.5-fold lower relative *CTT1* mRNA abundance during ethanol stress compared to the mean of all strains. Because YJM1129 is able to acquire H_2O_2 resistance when salt is the primary stress, and that phenotype requires *CTT1*, it's unlikely that allelic variation resulting in defective Ctt1p protein is responsible. One possibility is that Ctt1p protein levels are translationally regulated during stress, with low levels of translation happening during ethanol stress in the YJM1129 background. There is precedent for potential translational regulation of Ctt1p abundance during stress, as Berry and colleagues found that while *CTT1* mRNA levels increase during DTT, heat shock, and NaCl stresses, only NaCl stress resulted in Ctt1p protein being detectable by western blot [39].

Our results implicated glutathione-dependent peroxidases as responsible for *CTT1*-independent acquisition in three tested strains that possess *CTT1*-independent acquisition. Catalases are among the most efficient and fastest-acting known enzymes [57], so it is not surprising that yeast cytosolic catalase T (Ctt1p) plays a major role in acquired stress resistance. What was somewhat surprising was that some yeast strains solely rely on Ctt1p, while others supplement Ctt1p with glutathione-dependent peroxidases. It is possible that these

different strategies might benefit strains in different ecological niches. For example, since catalases uses heme as a cofactor, wild strains in more-iron limited environments may supplement catalase with glutathione-dependent peroxidases. On the other hand, glutathione is biosynthesized from sulfur-containing cysteine, so glutathione-dependent peroxidases may be less favored in sulfur limited conditions. Overall, these results suggest that different molecular strategies may be favored depending on the selective pressures commonly encountered in different environmental niches, and highlight the power of incorporating natural variation to uncover these unique molecular mechanisms.

Materials and Methods

Strains and growth conditions

Strains and primers used in this study are listed in S1 and S2 Tables, respectively. Fully prototrophic diploids in the S288c (DBY8268) background were generated by repairing the *ura3* auxotrophic marker in both MATa (JL505) and MATα (JL506) haploids (whose construction is described in [46]) via transformation of a PCR product containing the M22 *URA3* allele, followed by mating the two prototrophic haploids to generate diploid strain JL518. Because wild yeast strains can be heterozygous, we generated isogenic homozygous diploid segregants of wild yeast strains prior to genetic manipulation. Sporulation was induced by growing cells for 2 days in YPD (1% yeast extract, 2% peptone, 2% dextrose), harvesting by centrifugation (1,500 x *g* for 3 minutes), resuspending in 1% potassium acetate, and incubating for 2–10 days at 25°C with orbital shaking (270 rpm). Tetrads were then dissected and streaked to obtain colonies consisting of 'selfed' homozygous diploids (which occurs because the wild strains are homothallic and thus capable of mating type switching), and diploidy was confirmed using mating-type tester strains [58]. Deletions were introduced into the different wild strains via homologous recombination following transformation of the relevant deletion::KanMX cassette amplified from the Yeast Knockout Collection [59], with the exception of *ctt1Δ*::KanMX, which was amplified from YPS163-derived JL143 [47]. To generate double or triple mutants, a KanMX marker from one mutant was first exchanged with NatMX or HygMX via homologous recombination [60], followed by transformation with the appropriate second deletion::KanMX PCR product. All deletions were verified by diagnostic PCR that included primer pairs that annealed to the MX cassette and a sequence upstream of the deletion (verifying the 'scar'), and primer pairs that annealed within the deleted region (verifying absence of the gene). All yeast strains were grown at 30°C with orbital shaking (270 rpm). When used for selection, antibiotic concentrations were 100 μg/ml for nourseothricin (Nat), 200 μg/ml for G418 (Kan), and 300 μg/ml for hygromycin B (Hyg).

Cross protection assays

Cross protection assays were performed as described [47]. Briefly, cells were grown in YPD to saturation and then sub-cultured for at least 8 generations (> 12 hours) to mid-exponential phase (OD_{600} 0.3 – 0.6 on a Unico spectrophotometer). The cultures were then split, with one sample receiving a mock treatment (diluted 1:1 into fresh media) and the other sample receiving a mild 'primary' dose of stress (diluted 1:1 into fresh media containing either 10% v/v EtOH (5% final concentration) or 0.8 M NaCl (0.4 M final concentration)). Following a 1-hour incubation at 30°C with orbital shaking (270 rpm), mock and pre-treated cells were collected by mild centrifugation at 1,500 x *g* for 3 min, resuspended in fresh media to an OD_{600} of 0.6, and then diluted 3-fold (150 μl total volume) into a microtiter plate containing a panel of 11 severe 'secondary' H_2O_2 doses ranging from 0.75 – 5 mM (0.75 mM, 1 mM, 1.25 mM, 1.5 mM, 1.75 mM, 2 mM, 2.5 mM, 3 mM, 3.5 mM, 4 mM, 5 mM). Microtiter plates were sealed with breathable Rayon films (VWR), and cells were incubated with secondary stress for 2 hours at 30°C with

800 rpm shaking in a VWR Symphony incubating microplate shaker. Cells were diluted 50-fold in YPD a microtiter plate, and then 4 µl of each well was spotted onto YPD agar plates and grown for 48 hours at 30°C. Viability at each dose was scored using a 4-point semi-quantitative scale comparing viability to a no secondary stress (YPD only) control: 100% viability = 3 pts, 50–90% viability = 2 pts, 10–50% viability = 1 pt, and <10% (3 or fewer colonies) = 0 pts. A detailed protocol can be found on protocols.io under doi dx.doi.org/10.17504/protocols.io.g7sbzne. Hierarchical clustering of tolerance scores was performed using Cluster 3.0 (http://bonsai.hgc.jp/~mdehoon/software/cluster/software.htm), using centered Pearson correlation as the similarity metric and centroid linkage with a weighted cutoff of 0.4 and an exponent value of 1 [61], and visualized using Java TreeView (https://sourceforge.net/projects/jtreeview/files/) [62]. Raw tabular data used to generate figures can be found in S3 Table.

RNA sequencing and analysis

Two separate RNA-seq experiments were performed: i) comparison of phenotypically diverse strains (M1, M22, S288c, Y10, YJM1129, YJM308, YPS163, YPS606) responding to ethanol or NaCl stress relative to an unstressed control (performed in biological triplicate, except for the Y10 ethanol-stressed samples, where one sample was found post-analysis to be a mix-up) and ii) comparison of YPS606 vs. YPS606 transcription factor mutants (*msn2Δ msn4Δ*, *skn7Δ*, *yap1Δ*) responding to ethanol or NaCl stress relative to an unstressed control (performed in biological quadruplicate). For both sets of experiments, cells were grown at least 8 generations in 80 ml YPD to mid-exponential phase (OD_{600} 0.3-0.6). Fifteen-ml of an unstressed control sample was taken, and then the culture was split, with 20 ml of culture being added to 20-ml media containing either 10% v/v ethanol (5% final concentration) or 0.8 M NaCl (0.4 M NaCl final concentration). Following incubation in stress media for either 30 minutes (ethanol) or 45 minutes (NaCl), 15 ml of cells were collected by centrifugation at 1,500 x *g* for 1 minute, flash frozen in liquid nitrogen, and stored at -80°C until processed.

RNA was extracted using the 'hot phenol' method [63], and a detailed protocol is available at protocols.io under DOI dx.doi.org/10.17504/protocols.io.inwcdfe. Phenol-extracted total RNA was treated with DNase I (Ambion), and then purified with a Quick-RNA MiniPrep Plus Kit (Zymo Research) including the optional on-column DNase digestion. RNA integrity for each sample was assessed using an Agilent 2200 TapeStation, and all samples had a RIN value above 9.0.

RNA-seq libraries were prepared using the KAPA mRNA HyperPrep Kit (Roche) and KAPA Single-Indexed Adapter Set A + B, according to manufacturer specifications with minor modifications. Briefly, 500 ng of total RNA was used as starting material for each sample, polyA-enrichment was performed, mRNA was fragmented to an average size of 200–300 nucleotides, and the cDNA libraries were amplified for 9 PCR cycles. A detailed a protocol is available on protocols.io under DOI dx.doi.org/10.17504/protocols.io.uueewte. cDNA libraries were pooled (24-plex) and sequenced on a HiSeq4000 at the University of Chicago Genomics Facility, generating 50-bp single-end reads. To minimize batch effects, a random block design was used for library construction for each biological replicate, and each biological replicate was sequenced together on a single lane.

Reads were trimmed of low-quality reads and adapter sequences using Trimmomatic (version 0.38) [64], with the following commands: ILLUMINACLIP:Kapa_indices.fa:2:30:10 LEADING:3 TRAILING:3 MAXINFO:40:0.4 MINLEN:40. To generate new reference genomes for each wild strain, RNA-seq reads for each individual strain were first mapped to the S288c genome (version Scer3) using Bowtie 2 (version 2.3.4.1) [65], variants were called using bcftools (version 1.9) [66], and then variant calls were used to generate new reference genomes using GATK3 (Version 3.8-1-0) [67]. Trimmed reads were then mapped to their new corresponding reference genomes using STAR (version 2.6.1) [68]. Expected counts for each

transcript were generated using RSEM (version 1.3.1) [69], and RSEM outputs are reported in S4 Table. Differential expression analysis was performed using edgeR (version 3.26.4), using TMM normalization on all genes with at least one count per million (CPM) in at least one strain-condition, and the quasi-likelihood (QL) model [70], using sample type (e.g., S288c unstressed, S288c ethanol stressed, S288c NaCl stressed…) and biological replicate as factors.

To identify strain-specific differences in stress responses, the response of each strain was compared to the mean response of all strains. To identify sub-classes of strain-specific responses, we generated a 'consensus' response to each stress by using only treatment (unstressed, ethanol stressed, or NaCl stressed) and biological replicate as factors. Genes with significant differential expression (FDR < 0.01) of at least $\pm$ 1.5-fold were considered to be induced or repressed by each stress. Then for each strain comparison, genes with significant differential expression (FDR < 0.01) of at least $\pm$ 1.5-fold compared to the mean of all strains were classified based on the direction of the difference (e.g., genes with >1.5-fold higher expression compared to the mean of all strains for 'consensus' induced genes were classified as having "amplified induction," while a sign shift from 'consensus' induction to >1.5-fold repression resulted in genes being classified as "novel repression"). All raw RNA-seq data are available through the National Institutes of Health Gene Expression Omnibus (GEO) database under accession no. GSE248219, and the edgeR outputs can be found in S5 Table.

Hierarchical clustering of gene expression $\log_2$ fold changes was performed with Cluster 3.0, using Euclidean distance as the similarity metric, and centroid linkage with weighted cutoff of 0.4 and an exponent value of 1, and visualized in Java TreeView. Functional enrichments of gene ontology (GO) categories were performed using Princeton's GO-TermFinder (https://go.princeton.edu/cgibin/GOTermFinder) [71], with Bonferroni-corrected *P*-values < 0.01 taken as significant. Significantly enriched regulatory associations (FDR-corrected *P* < 0.01) were identified using the YEAst Search for Transcriptional Regulators And Consensus Tracking (YEASTRACT) database [72], using documented DNA binding or expression evidence (with the transcription factor acting as an activator or unknown), filtered on conditions defined as stress. GO and regulatory enrichments can be found in S6 Table.

Acknowledgements

We thank Dr. Justin Fay and Dr. Audrey Gasch for wild yeast strains, Dr. Elizabeth Ruck for running samples on the TapeStation, and Dr. Andrew Alverson for the use of equipment. This work was supported in part by National Science Foundation Grants IOS-1656602 (JAL) and MCB-1941824 (JAL), the Arkansas Biosciences Institute (Arkansas Settlement Proceeds Act of 2000) (JAL), and a Research Assistantship provided through the University of Arkansas Cell and Molecular Biology Graduate Program (ANS). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

References

1. Al-Nabulsi AA, Osaili TM, Shaker RR, Olaimat AN, Jaradat ZW, Zain Elabedeen NA, et al. Effects of osmotic pressure, acid, or cold stresses on antibiotic susceptibility of *Listeria monocytogenes*. Food Microbiology. 2015;46:154-60. doi: 10.1016/j.fm.2014.07.015.
2. Wang X, Vignjevic M, Liu F, Jacobsen S, Jiang D, Wollenweber B. Drought priming at vegetative growth stages improves tolerance to drought and heat stresses occurring during grain filling in spring wheat. Plant Growth Regulation. 2014;75(3):677-87. doi: 10.1007/s10725-014-9969-x.
3. Yadav R, Juneja S, Kumar S. Cross priming with drought improves heat-tolerance in chickpea (*Cicer arietinum* L.) by stimulating small heat shock proteins and antioxidative defense. Environmental Sustainability. 2021;4(1):171-82. doi: 10.1007/s42398-020-00156-4.
4. Ru C, Hu X, Chen D, Wang W, Song T. Heat and drought priming induce tolerance to subsequent heat and drought stress by regulating leaf photosynthesis, root morphology, and antioxidant defense in maize seedlings. Environmental and Experimental Botany. 2022;202. doi: 10.1016/j.envexpbot.2022.105010.
5. Murapa P, Gandhapudi S, Skaggs HS, Sarge KD, Woodward JG. Physiological Fever Temperature Induces a Protective Stress Response in T Lymphocytes Mediated by Heat Shock Factor-1 (HSF1). The Journal of Immunology. 2007;179(12):8305-12. doi: 10.4049/jimmunol.179.12.8305.
6. Su C-Y, Chong K-Y, Chen J, Ryter S, Khardori R, Lai C-C. A Physiologically Relevant Hyperthermia Selectively Activates Constitutive hsp70 in H9c2 Cardiac Myoblasts and Confers Oxidative Protection. Journal of Molecular and Cellular Cardiology. 1999;31(4):845-55. doi: 10.1006/jmcc.1998.0923.
7. McCormick PH, Chen G, Tlerney S, Kelly CJ, Bouchier-Hayes DJ. Clinically relevant thermal preconditioning attenuates ischemia-reperfusion injury. Journal of Surgical Research. 2003;109(1):24-30. doi: 10.1016/s0022-4804(02)00035-5.
8. Fujita K, Matsuyama A, Kobayashi Y, Iwahashi H. The genome-wide screening of yeast deletion mutants to identify the genes required for tolerance to ethanol and other alcohols. FEMS yeast research. 2006;6(5):744-50. doi: 10.1111/j.1567-1364.2006.00040.x. PubMed PMID: 16879425.
9. García R, Botet J, Rodríguez-Peña JM, Bermejo C, Ribas JC, Revuelta JL, et al. Genomic profiling of fungal cell wall-interfering compounds: identification of a common gene signature. BMC genomics. 2015;16(1). doi: 10.1186/s12864-015-1879-4.
10. Hoepfner D, Helliwell SB, Sadlish H, Schuierer S, Filipuzzi I, Brachat S, et al. High-resolution chemical dissection of a model eukaryote reveals targets, pathways and gene functions. Microbiological Research. 2014;169(2-3):107-20. doi: 10.1016/j.micres.2013.11.004.
11. Kapitzky L, Beltrao P, Berens TJ, Gassner N, Zhou C, Wüster A, et al. Cross-species chemogenomic profiling reveals evolutionarily conserved drug mode of action. Molecular systems biology. 2010;6(1). doi: 10.1038/msb.2010.107.
12. Lum PY, Armour CD, Stepaniants SB, Cavet G, Wolf MK, Butler JS, et al. Discovering Modes of Action for Therapeutic Compounds Using a Genome-Wide Screen of Yeast Heterozygotes. Cell. 2004;116(1):121-37. doi: 10.1016/s0092-8674(03)01035-3.
13. Parsons AB, Brost RL, Ding H, Li Z, Zhang C, Sheikh B, et al. Integration of chemical-genetic and genetic interaction data links bioactive compounds to cellular target pathways. Nature biotechnology. 2003;22(1):62-9. doi: 10.1038/nbt919.

14. Yadav J, Muend S, Zhang Y, Rao R, Malhotra V. A Phenomics Approach in Yeast Links Proton and Calcium Pump Function in the Golgi. Molecular biology of the cell. 2007;18(4):1480-9. doi: 10.1091/mbc.e06-11-1049.

15. van Bakel H, Strengman E, Wijmenga C, Holstege FCP. Gene expression profiling and phenotype analyses of *S. cerevisiae* in response to changing copper reveals six genes with new roles in copper and iron metabolism. Physiological Genomics. 2005;22(3):356-67. doi: 10.1152/physiolgenomics.00055.2005.

16. Jiang L, Cao C, Zhang L, Lin W, Xia J, Xu H, et al. Cadmium-induced activation of high osmolarity glycerol pathway through its Sln1 branch is dependent on the MAP kinase kinase kinase Ssk2, but not its paralog Ssk22, in budding yeast. FEMS yeast research. 2014;14(8):1263-72. doi: 10.1111/1567-1364.12220.

17. Hwang G-W, Murai Y, Takahashi T, Naganuma A. The protein transportation pathway from Golgi to vacuoles via endosomes plays a role in enhancement of methylmercury toxicity. Scientific Reports. 2014;4(1). doi: 10.1038/srep05888.

18. Hernández RB, Moteshareie H, Burnside D, McKay B, Golshani A. Manganese-induced cellular disturbance in the baker's yeast, *Saccharomyces cerevisiae* with putative implications in neuronal dysfunction. Scientific Reports. 2019;9(1). doi: 10.1038/s41598-019-42907-2.

19. Bleackley MR, Young BP, Loewen CJR, MacGillivray RTA. High density array screening to identify the genetic requirements for transition metal tolerance in *Saccharomyces cerevisiae*. Metallomics. 2011;3(2). doi: 10.1039/c0mt00035c.

20. Mollapour M, Fong D, Balakrishnan K, Harris N, Thompson S, Schüller C, et al. Screening the yeast deletant mutant collection for hypersensitivity and hyper-resistance to sorbate, a weak organic acid food preservative. Yeast. 2004;21(11):927-46. doi: 10.1002/yea.1141.

21. Rieger K-J, El-Alama M, Stein G, Bradshaw C, Slonimski PP, Maundrell K. Chemotyping of yeast mutants using robotics. Yeast. 1999;15(10B):973-86. doi: 10.1002/(sici)1097-0061(199907)15:10b<973::Aid-yea402>3.0.Co;2-l.

22. Schmidt M, Akasaka K, Messerly JT, Boyer MP. Role of Hog1, Tps1 and Sod1 in boric acid tolerance of *Saccharomyces cerevisiae*. Microbiology. 2012;158(10):2667-78. doi: 10.1099/mic.0.060590-0.

23. Okada N, Ogawa J, Shima J. Comprehensive analysis of genes involved in the oxidative stress tolerance using yeast heterozygous deletion collection. FEMS yeast research. 2014;14(3):425-34. doi: 10.1111/1567-1364.12136.

24. Begley TJ, Rosenbach AS, Ideker T, Samson LD. Hot Spots for Modulating Toxicity Identified by Genomic Phenotyping and Localization Mapping. Molecular cell. 2004;16(1):117-25. doi: 10.1016/j.molcel.2004.09.005.

25. Dudley AM, Janse DM, Tanay A, Shamir R, Church GM. A global view of pleiotropy and phenotypically derived gene function in yeast. Molecular systems biology. 2005;1(1). doi: 10.1038/msb4100004.

26. van Voorst F, Houghton-Larsen J, Jonson L, Kielland-Brandt MC, Brandt A. Genome-wide identification of genes required for growth of *Saccharomyces cerevisiae* under ethanol stress. Yeast. 2006;23(5):351-9. doi: 10.1002/yea.1359. PubMed PMID: 16598687.

27. Teixeira MC, Raposo LR, Mira NP, Lourenco AB, Sa-Correia I. Genome-wide identification of *Saccharomyces cerevisiae* genes required for maximal tolerance to ethanol. Appl Environ Microbiol. 2009;75(18):5761-72. doi: 10.1128/AEM.00845-09. PubMed PMID: 19633105; PubMed Central PMCID: PMCPMC2747848.

28. Ruiz-Roig C, Vieitez C, Posas F, de Nadal E. The Rpd3L HDAC complex is essential for the heat stress response in yeast. Molecular microbiology. 2010;76:1049-62. Epub 2010/04/20. doi: 10.1111/j.1365-2958.2010.07167.x. PubMed PMID: 20398213.

29. Ando A, Tanaka F, Murata Y, Takagi H, Shima J. Identification and classification of genes required for tolerance to high-sucrose stress revealed by genome-wide screening of *Saccharomyces cerevisiae*. FEMS yeast research. 2006;6(2):249-67. doi: 10.1111/j.1567-1364.2006.00035.x.

30. Outten Caryn E, Falk Robert L, Culotta Valeria C. Cellular factors required for protection from hyperoxia toxicity in *Saccharomyces cerevisiae*. Biochemical Journal. 2005;388(1):93-101. doi: 10.1042/bj20041914.

31. Brown JA, Sherlock G, Myers CL, Burrows NM, Deng C, Wu HI, et al. Global analysis of gene function in yeast by quantitative phenotypic profiling. Molecular systems biology. 2006;2(1). doi: 10.1038/msb4100043.

32. Jarolim S, Ayer A, Pillay B, Gee AC, Phrakaysone A, Perrone GG, et al. *Saccharomyces cerevisiae* genes involved in survival of heat shock. G3 (Bethesda). 2013;3(12):2321-33. doi: 10.1534/g3.113.007971. PubMed PMID: 24142923; PubMed Central PMCID: PMCPMC3852394.

33. Giaever G, Chu AM, Ni L, Connelly C, Riles L, Veronneau S, et al. Functional profiling of the *Saccharomyces cerevisiae* genome. Nature. 2002;418(6896):387-91. doi: 10.1038/nature00935. PubMed PMID: 12140549.

34. Higgins VJ, Alic N, Thorpe GW, Breitenbach M, Larsson V, Dawes IW. Phenotypic analysis of gene deletant strains for sensitivity to oxidative stress. Yeast. 2002;19(3):203-14. doi: 10.1002/yea.811.

35. Yoshikawa K, Tanaka T, Furusawa C, Nagahisa K, Hirasawa T, Shimizu H. Comprehensive phenotypic analysis for identification of genes affecting growth under ethanol stress in *Saccharomyces cerevisiae.* FEMS yeast research. 2009;9:32-44. PubMed PMID: 19054128.

36. Auesukaree C, Damnernsawad A, Kruatrachue M, Pokethitiyook P, Boonchird C, Kaneko Y, et al. Genome-wide identification of genes involved in tolerance to various environmental stresses in *Saccharomyces cerevisiae*. Journal of Applied Genetics. 2009;50(3):301-10. doi: 10.1007/bf03195688.

37. Kelley R, Ideker T. Genome-wide fitness and expression profiling implicate Mga2 in adaptation to hydrogen peroxide. PLoS genetics. 2009;5(5):e1000488. Epub 2009/06/09. doi: 10.1371/journal.pgen.1000488. PubMed PMID: 19503593; PubMed Central PMCID: PMCPMC2676504.

38. Ng C-H, Tan S-X, Perrone GG, Thorpe GW, Higgins VJ, Dawes IW. Adaptation to hydrogen peroxide in Saccharomyces cerevisiae: The role of NADPH-generating systems and the *SKN7* transcription factor. Free Radical Biology and Medicine. 2008;44(6):1131-45. doi: 10.1016/j.freeradbiomed.2007.12.008.

39. Berry DB, Guan Q, Hose J, Haroon S, Gebbia M, Heisler LE, et al. Multiple means to the same end: the genetic basis of acquired stress resistance in yeast. PLoS genetics. 2011;7(11):e1002353. Epub 2011/11/22. doi: 10.1371/journal.pgen.1002353. PubMed PMID: 22102822; PubMed Central PMCID: PMCPMC3213159.

40. Gibney PA, Lu C, Caudy AA, Hess DC, Botstein D. Yeast metabolic and signaling genes are required for heat-shock survival and have little overlap with the heat-induced genes. Proceedings of the National Academy of Sciences of the United States of America. 2013;110(46):E4393-402. Epub 2013/10/30. doi: 10.1073/pnas.1318100110. PubMed PMID: 24167267; PubMed Central PMCID: PMCPMC3831991.

41. Mir SS, Fiedler D, Cashikar AG. Ssd1 Is Required for Thermotolerance and Hsp104-Mediated Protein Disaggregation in *Saccharomyces cerevisiae*. Molecular and cellular biology. 2023;29(1):187-200. doi: 10.1128/mcb.02271-07.

42. Berry DB, Gasch AP. Stress-activated genomic expression changes serve a preparative role for impending stress in yeast. Molecular biology of the cell. 2008;19(11):4580-7.

Epub 2008/08/30. doi: 10.1091/mbc.E07-07-0680. PubMed PMID: 18753408; PubMed Central PMCID: PMCPMC2575158.

43. Mitchell A, Romano GH, Groisman B, Yona A, Dekel E, Kupiec M, et al. Adaptive prediction of environmental changes by microorganisms. Nature. 2009;460(7252):220-4. doi: 10.1038/nature08112. PubMed PMID: 19536156.

44. Święciło A. Cross-stress resistance in *Saccharomyces cerevisiae* yeast—new insight into an old phenomenon. Cell Stress and Chaperones. 2016;21(2):187-200. doi: 10.1007/s12192-016-0667-7.

45. Lewis JA, Elkon IM, McGee MA, Higbee AJ, Gasch AP. Exploiting natural variation in *Saccharomyces cerevisiae* to identify genes for increased ethanol resistance. Genetics. 2010;186:1197-205. PubMed PMID: 20855568.

46. McDaniel EA, Stuecker TN, Veluvolu M, Gasch AP, Lewis JA. Independent Mechanisms for Acquired Salt Tolerance versus Growth Resumption Induced by Mild Ethanol Pretreatment in Saccharomyces cerevisiae. mSphere. 2018;3(6). Epub 2018/11/30. doi: 10.1128/mSphere.00574-18. PubMed PMID: 30487155; PubMed Central PMCID: PMCPMC6262259.

47. Stuecker TN, Scholes AN, Lewis JA. Linkage mapping of yeast cross protection connects gene expression variation to a higher-order organismal trait. PLoS genetics. 2018;14(4):e1007335. doi: 10.1371/journal.pgen.1007335. PubMed PMID: 29649251.

48. D'Orazi G, Cirone M. Interconnected Adaptive Responses: A Way Out for Cancer Cells to Avoid Cellular Demise. Cancers. 2022;14(11). doi: 10.3390/cancers14112780.

49. Raffaghello L, Lee C, Safdie FM, Wei M, Madia F, Bianchi G, et al. Starvation-dependent differential stress resistance protects normal but not cancer cells against high-dose chemotherapy. Proceedings of the National Academy of Sciences of the United States of America. 2008;105:8215-20. PubMed PMID: 18378900.

50. Clowers KJ, Heilberger J, Piotrowski JS, Will JL, Gasch AP. Ecological and Genetic Barriers Differentiate Natural Populations of *Saccharomyces cerevisiae*. Molecular biology and evolution. 2015;32(9):2317-27. Epub 2015/05/09. doi: 10.1093/molbev/msv112. PubMed PMID: 25953281; PubMed Central PMCID: PMCPMC4540968.

51. Gasch AP, Spellman PT, Kao CM, Carmel-Harel O, Eisen MB, Storz G, et al. Genomic expression programs in the response of yeast cells to environmental changes. Molecular biology of the cell. 2000;11(12):4241-57. PubMed PMID: 11102521; PubMed Central PMCID: PMC15070.

52. Causton HC, Ren B, Koh SS, Harbison CT, Kanin E, Jennings EG, et al. Remodeling of yeast genome expression in response to environmental changes. Molecular biology of the cell. 2001;12(2):323-37. PubMed PMID: 11179418; PubMed Central PMCID: PMC30946.

53. Lewis JA, Broman AT, Will J, Gasch AP. Genetic architecture of ethanol-responsive transcriptome variation in *Saccharomyces cerevisiae* strains. Genetics. 2014;198(1):369-82. doi: 10.1534/genetics.114.167429. PubMed PMID: 24970865; PubMed Central PMCID: PMC4174948.

54. Stephen DWS, Rivers SL, Jamieson DJ. The role of the *YAP1* and *YAP2* genes in the regulation of the adaptive oxidative stress responses of *Saccharomyces cerevisiae*. Molecular microbiology. 1995;16(3):415-23. doi: 10.1111/j.1365-2958.1995.tb02407.x.

55. Thorpe GW, Fong CS, Alic N, Higgins VJ, Dawes IW. Cells have distinct mechanisms to maintain protection against different reactive oxygen species: Oxidative-stress-response genes. Proceedings of the National Academy of Sciences. 2004;101(17):6564-9. doi: 10.1073/pnas.0305888101.

56. Gasch AP, Payseur BA, Pool JE. The Power of Natural Variation for Model Organism Biology. Trends in genetics : TIG. 2016;32(3):147-54. Epub 2016/01/19. doi:

10.1016/j.tig.2015.12.003. PubMed PMID: 26777596; PubMed Central PMCID: PMCPMC4769656.

57. Switala J, Loewen PC. Diversity of properties among catalases. Archives of Biochemistry and Biophysics. 2002;401(2):145-54. doi: 10.1016/s0003-9861(02)00049-8.

58. Julius D, Blair L, Brake A, Sprague G, Thorner J. Yeast α factor is processed from a larger precursor polypeptide: The essential role of a membrane-bound dipeptidyl aminopeptidase. Cell. 1983;32(3):839-52. doi: 10.1016/0092-8674(83)90070-3.

59. Winzeler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, Andre B, et al. Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. Science. 1999;285:901-6. Epub 1999/08/07. PubMed PMID: 10436161.

60. Goldstein AL, McCusker JH. Three new dominant drug resistance cassettes for gene disruption in *Saccharomyces cerevisiae*. Yeast. 1999;15(14):1541-53. doi: 10.1002/(sici)1097-0061(199910)15:14<1541::Aid-yea476>3.0.Co;2-k.

61. Eisen MB, Spellman PT, Brown PO, Botstein D. Cluster analysis and display of genome-wide expression patterns. Proceedings of the National Academy of Sciences of the United States of America. 1998;95:14863-8. PubMed PMID: 9843981.

62. Saldanha AJ. Java Treeview--extensible visualization of microarray data. Bioinformatics. 2004;20(17):3246-8. doi: 10.1093/bioinformatics/bth349.

63. Scholes AN, Lewis JA. Comparison of RNA isolation methods on RNA-Seq: implications for differential expression and meta-analyses. BMC genomics. 2020;21(1):249. Epub 2020/03/22. doi: 10.1186/s12864-020-6673-2. PubMed PMID: 32197587; PubMed Central PMCID: PMCPMC7082934.

64. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 2014;30(15):2114-20. doi: 10.1093/bioinformatics/btu170. PubMed PMID: 24695404; PubMed Central PMCID: PMCPMC4103590.

65. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:357-9. doi: 10.1038/nmeth.1923. PubMed PMID: 22388286; PubMed Central PMCID: PMC3322381.

66. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics. 2011;27(21):2987-93. doi: 10.1093/bioinformatics/btr509.

67. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. Genome research. 2010;20(9):1297-303. doi: 10.1101/gr.107524.110.

68. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. Epub 2012/10/30. doi: 10.1093/bioinformatics/bts635. PubMed PMID: 23104886; PubMed Central PMCID: PMCPMC3530905.

69. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC bioinformatics. 2011;12:323. Epub 2011/08/06. doi: 10.1186/1471-2105-12-323. PubMed PMID: 21816040; PubMed Central PMCID: PMCPMC3163565.

70. Chen Y, Lun ATL, Smyth GK. From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. F1000Research. 2016;5. doi: 10.12688/f1000research.8987.2.

71. Boyle EI, Weng S, Gollub J, Jin H, Botstein D, Cherry JM, et al. GO::TermFinder--open source software for accessing Gene Ontology information and finding significantly enriched Gene Ontology terms associated with a list of genes. Bioinformatics. 2004;20(18):3710-5. Epub 2004/08/07. doi: 10.1093/bioinformatics/bth456. PubMed PMID: 15297299; PubMed Central PMCID: PMC3037731.

72. Teixeira MC, Monteiro PT, Palma M, Costa C, Godinho CP, Pais P, et al. YEASTRACT: an upgraded database for the analysis of transcription regulatory networks in Saccharomyces cerevisiae. Nucleic acids research. 2018;46(D1):D348-D53. Epub 2017/10/17. doi: 10.1093/nar/gkx842. PubMed PMID: 29036684; PubMed Central PMCID: PMCPMC5753369.

73. Rep M, Reiser V, Gartner U, Thevelein JM, Hohmann S, Ammerer G, et al. Osmotic Stress-Induced Gene Expression in *Saccharomyces cerevisiae* Requires Msn1p and the Novel Nuclear Factor Hot1p. Molecular and cellular biology. 1999;19(8):5474-85. doi: 10.1128/mcb.19.8.5474.

74. Wiederrecht G, Seto D, Parker CS. Isolation of the gene encoding the *S. cerevisiae* heat shock transcription factor. Cell. 1988;54(6):841-53. doi: 10.1016/s0092-8674(88)91197-x.

75. Chandler M, Stanley GA, Rogers P, Chambers P. A genomic approach to defining the ethanol stress response in the yeast Saccharomyces cerevisiae. Annals of Microbiology. 2004;54(4):427-54. PubMed PMID: WOS:000226462900007.

76. Martinez-Pastor MT, Marchler G, Schuller C, Marchler-Bauer A, Ruis H, Estruch F. The *Saccharomyces cerevisiae* zinc finger proteins Msn2p and Msn4p are required for transcriptional induction through the stress response element (STRE). The EMBO journal. 1996;15:2227-35. PubMed PMID: 8641288.

77. Gorner W, Durchschlag E, Martinez-Pastor MT, Estruch F, Ammerer G, Hamilton B, et al. Nuclear localization of the C2H2 zinc finger protein Msn2p is regulated by stress and protein kinase A activity. Genes Dev. 1998;12:586-97. PubMed PMID: 9472026.

78. Raitt DC, Johnson AL, Erkine AM, Makino K, Morgan B, Gross DS, et al. The Skn7 Response Regulator of *Saccharomyces cerevisiae* Interacts with Hsf1 *In Vivo* and Is Required for the Induction of Heat Shock Genes by Oxidative Stress. Molecular biology of the cell. 2000;11(7):2335-47. doi: 10.1091/mbc.11.7.2335.

79. Hohmann S. Osmotic Stress Signaling and Osmoadaptation in Yeasts. Microbiology and Molecular Biology Reviews. 2002;66(2):300-72. doi: 10.1128/mmbr.66.2.300-372.2002.

80. Lee J, Godon C, Lagniel G, Spector D, Garin J, Labarre J, et al. Yap1 and Skn7 Control Two Specialized Oxidative Stress Response Regulons in Yeast. Journal of Biological Chemistry. 1999;274(23):16040-6. doi: 10.1074/jbc.274.23.16040.

81. Rep M, Proft M, Remize F, Tamas M, Serrano R, Thevelein JM, et al. The *Saccharomyces cerevisiae* Sko1p transcription factor mediates HOG pathway-dependent osmotic regulation of a set of genes encoding enzymes implicated in protection from oxidative damage. Molecular microbiology. 2001;40(5):1067-83. doi: 10.1046/j.1365-2958.2001.02384.x.

82. de Nadal E, Casadomé L, Posas F. Targeting the MEF2-Like Transcription Factor Smp1 by the Stress-Activated Hog1 Mitogen-Activated Protein Kinase. Molecular and cellular biology. 2023;23(1):229-37. doi: 10.1128/mcb.23.1.229-237.2003.

Tables

Table 1. Potential transcription factors involved in acquired stress resistance.

Transcription Factor (TF)	Description	References
Hot1p	Osmotic stress responsive TF	[72]
Hsf1p	Essential heat shock factor	[73, 74]
Msn1p	Osmotic stress responsive TF	[72]
Msn2/4p	Paralogous general stress-responsive TFs	[74-76]
Skn7p	Osmotic and oxidative stress-responsive TF	[77-79]
Sko1p	Osmotic stress-responsive TF	[80]
Smp1p	Osmotic stress responsive TF	[81]
Yap1p	Oxidative stress-responsive TF	[79]

Figures

Fig.1 *CTT1* is partially dispensable for salt-induced, but not ethanol-induced, cross protection against H_2O_2 in a wild oak strain. (A) Representative acquired H_2O_2 resistance assays for wild-type S288c (lab) and YPS163 (oak) strains and respective *ctt1Δ* mutants. Strains were exposed to a mild 'primary' stress pretreatment (5% ethanol or 0.4 M NaCl in culture media) or a mock control for 60 minutes, collected and resuspended in fresh media, exposed to a panel of 11 doses of severe 'secondary' doses of H_2O_2 for 2 hours, and then spot plated to score viability. (B) A single survival score was calculated from the viability at all H_2O_2 doses (see Materials and Methods). Each plot shows the mean and standard deviation of three independent biological replicates. The replicates for some strain conditions all had the same tolerance score and thus zero standard deviation (see S3 Table for raw numerical data). Asterisks represent acquired H_2O_2 resistance that was significantly higher in the YPS163 *ctt1Δ* mutant with NaCl as the pretreatment compared to the mock control (* $P < 0.02$, paired *t*-test).

Fig 2. Diverse yeast strains show complex patterns of conditional *CTT1* dependency. (A) Representative acquired H_2O_2 resistance assays of wild-type strains and their *ctt1Δ* mutant derivatives. (B) Hierarchical clustering of H_2O_2 survival scores for wild-type strains following either ethanol (E) or salt (N) pretreatments, or their *ctt1Δ* derivatives (ΔE and ΔE). Strain labels are color coded according to their origin as indicated in the key on the right. (C) Percent acquired H_2O_2 resistance in *ctt1Δ* mutants relative to that of the wild type revealed three classes of conditional *CTT1* dependency. Error bars denote the mean and standard deviation of biological duplicates, except for S288c and YPS163 which are in biological triplicate.

Fig 3. Extensive variation in stress responsive gene expression across diverse yeast isolates. Hierarchical clustering of 4,262 genes with significant (FDR < 0.01) differential ethanol-responsive expression in at least one strain relative to the mean of all strains (A), and 3,918 genes with significant (FDR < 0.01) differential salt-responsive expression in at least one strain relative to the mean of all strains (B). Genes are shown as rows and $\log_2$ fold changes for each strain's stress response (left) or difference in stress response (right) are shown as columns. Red indicates induced and blue indicates repressed expression in response to stress, while violet indicates higher and brown indicates lower expression relative to the mean stress response of all strains. Significant regulatory enrichments are annotated to the left and functional enrichments are annotated to the right (Bonferroni-corrected P < 0.01). Full lists of regulatory and functional enrichments are in S6 Table.

Fig 4. Strain-specific variation in different classes of stress-activated expression changes. Total number of genes for each strain with significantly (FDR < 0.01) different (> 1.5-fold) induction patterns (A) or repression patterns (B) relative to the mean or 'consensus' response (see Methods). S7 and S8 Tables contain the genes for ethanol-responsive and salt-responsive categories, respectively.

Fig 5. *msn2Δ msn4Δ*, *skn7Δ*, and *yap1Δ* mutants have varied effects on intrinsic and acquired H₂O₂ resistance. Survival score plots representing the mean and standard deviation for biological triplicates for depicted strains with either 5% ethanol (A) or 0.4 M NaCl (B) as the mild pre-treatment. Error bars denote the standard deviation. The replicates for NaCl-treated *yap1Δ* and *skn7Δ* mutants all had the same tolerance score and thus zero standard deviation (see Table S3 for raw numerical data). Asterisks represent significant differences in acquired resistance between denoted strains (* *P* < 0.05, ** *P* < 0.01, *** *P* < 0.001, *t*-test).

Fig 6. Msn2/4p, Skn7p, and Yap1p regulate shared and unique targets for the YPS606 ethanol and NaCl responses. The Venn diagrams represent genes with significant defective induction for each mutant's stress response (FDR < 0.01 and at least 1.5-fold reduced expression). Significant functional enrichments (Bonferroni-corrected P < 0.01) are shown below for all unique and shared genes.

Fig 7. Stress-responsive expression variation in genes involved in detoxifying reactive oxygen species. Hierarchical clustering of all yeast genes annotated as having peroxidase activity in response to either ethanol (A) or NaCl (B). Genes are shown as rows and $\log_2$ fold changes for each strain's stress response (left) or difference in stress response (right) are shown as columns. Red indicates induced and blue indicates repressed expression in response to stress, while violet indicates higher and brown indicates lower expression relative to the mean stress response of all strains.

Fig 8. *CTT1*-dispensible acquired H₂O₂ resistance requires glutathione biosynthesis.
Survival score plots indicating the mean and standard deviation for biological triplicates for
depicted strains with either 5% ethanol (A) or 0.4 M NaCl (B) as the mild pre-treatment. Error
bars denote the standard deviation. The replicates for NaCl-treated YPS606, YPS606 *gsh1Δ*,
M22, and M22 *gsh1Δ* strains all had the same tolerance score and thus zero standard deviation
(see Table S3 for raw numerical data). Asterisks represent significant differences in acquired
resistance between denoted strains (* *P* < 0.05, ** *P* < 0.01, ns = not significant (*P* > 0.05), *t*-
test).

Supporting Figures

S1 Figure. Representative acquired H₂O₂ resistance assays a panel of diverse yeast strains. Representative acquired H_2O_2 resistance assays are shown for all strains depicted in Fig 2.

S2 Figure. Acquired H₂O₂ resistance assays for a panel YPS606 transcription factor deletion mutants. Acquired H_2O_2 resistance assays were performed in homozygous YPS606 *sko1Δ*, *hot1Δ*, and *msn1Δ* mutants plus a heterozygous *HSF1/hsf1Δ* mutant (because *HSF1* is essential). Mutants were assayed in biological triplicate, while wild-type YPS606 was included as a control for each set of experiments and thus had 24 replicates. Error bars depict the standard deviation (note that the replicates some strain comparisons had the exact same tolerance score and thus zero standard deviation).

S3 Figure. Representative acquired H₂O₂ resistance assays for all transcription factor mutants. Representative acquired H_2O_2 resistance assays are shown for all strains depicted in Fig 6 and S2 Figure.

S4 Figure. Representative acquired H$_2$O$_2$ resistance assays for YPS606 *gsh1Δ* mutants. Representative acquired H$_2$O$_2$ resistance assays are shown for all strains depicted in Fig 8.

Supporting Tables

S1 Table: Strains used in this study.

S2 Table: Primers used in this study.

S3 Table: Raw data used to generate figures.

S4 Table: RSEM Output.

S5 Table: edgeR Output.

S6 Table: GO and TF binding-site enrichments for clusters in Figure 3.

S7 Table: Categories of differential ethanol-responsive expression in Figure 4.

S7 Table: Categories of differential salt-responsive expression in Figure 4.