Smell and Aftersmell: Fast Calcium Imaging Dynamics of Honey Bee Olfactory Coding

Marco Paoli[1], Antoine Wystrach[2], Brice Ronsin[2], Martin Giurfa[1,3]

[1] Neuroscience Paris-Seine – Institut de biologie Paris-Seine, Sorbonne Université, INSERM, CNRS, Paris, France

[2] Centre de Recherches sur la Cognition Animale, Centre de Biologie Intégrative, Université Paul Sabatier, CNRS, Toulouse, France

[3] Institut Universitaire de France (IUF), Paris, France

Abstract

Odour processing exhibits multiple parallels between vertebrate and invertebrate olfactory systems. Insects, in particular, have emerged as relevant models for olfactory studies because of the tractability of their olfactory circuits. Here, we used fast calcium imaging to track the activity of projection neurons in the honey bee antennal lobe (AL) during olfactory stimulation at high temporal resolution. We observed a heterogeneity of response profiles and an abundance of inhibitory activities, resulting in various response latencies and stimulus-specific post-odour neural signatures. Recorded calcium signals were fed to a mushroom body (MB) model constructed implementing the fundamental features of connectivity between olfactory projection neurons, Kenyon cells (KC), and MB output neurons (MBON). The model accounts for the increase of odorant discrimination in the MB compared to the AL and reveals the recruitment of two distinct KC populations that represent odorants and their aftersmell as two separate but temporally coherent neural objects. Finally, we showed that the learning-induced modulation of KC-to-MBON synapses can explain both the variations in associative learning scores across different conditioning protocols used in bees and the bees' response latency. Thus, it provides a simple explanation of how the time contingency between the stimulus and the reward can be encoded without the need for time tracking. This study broadens our understanding of olfactory coding and learning in honey bees. It demonstrates that a model based on simple MB connectivity rules and fed with real physiological data can explain fundamental aspects of odour processing and associative learning.

Introduction

The logic of olfactory coding and learning has been extensively studied at the behavioural and neural levels in various insect models **(Adam et al., 2022; Galizia, 2014; Giurfa, 2015; Jefferis et al., 2007)**. Among them, the honey bee *Apis mellifera* has played a pivotal role in our understanding of these processes due to its behavioural accessibility and tractability of its nervous system. In bees, olfactory perception and learning have been typically investigated using the proboscis extension reflex (PER) **(Giurfa, 2007; Menzel, 1999)**, a protocol that relies on pairing a neutral olfactory stimulus (the conditioned stimulus or CS) with a positive reinforcement of sugar solution (the unconditioned stimulus or US) **(Bitterman et al., 1983; Giurfa and Sandoz, 2012; Takeda, 1961)**. In the classical version of PER conditioning, the CS precedes and partially overlaps US presentation. This results in high levels of specific memory for the conditioned odorant. Conversely, when the reward precedes the stimulus (backward conditioning), no positive association can be established **(Felsenberg et al., 2014; Hellstern et al., 1998)**. It also enables the study of more sophisticated cognitive processes such as trace learning **(Paoli et al., 2023a; Szyszka et al., 2011)** and patterning discrimination **(Deisig et al., 2001; Devaud et al., 2015)**. In the former, CS and US are not overlapping but separated by a stimulus-free temporal gap, whereas in the latter, bees are trained to respond in opposite ways to a two-odorant mixture compared to its individual components **(Deisig et al., 2001; Devaud et al., 2015)**. Overall, the olfactory conditioning of PER provides a robust read-out for investigating the dynamics of olfactory memory formation **(Giurfa and Sandoz, 2012; Villar et al., 2020)** and olfactory perception **(Guerrieri et al., 2005)**.

Our understanding of the neural processing subtending olfactory coding and learning relies on decades of neuroanatomical and neurophysiological studies performed in honey bees **(Paoli and Galizia, 2021)**. Olfactory processing starts at the peripheral level, when volatile chemicals interact with the olfactory receptors expressed on the dendritic membrane of the

olfactory sensory neurons (OSNs). The biochemical nature of odorant-receptor interactions allows for a certain molecule to bind to multiple receptors with different affinities, resulting in the activation of an OSNs sub-population with odorant-specific response intensities and latencies **(Münch and Galizia, 2016)**. Olfactory sensory neurons innervate the first olfactory processing centre, the antennal lobe (AL), where the neurons expressing the same olfactory receptor converge onto one of ~160 glomeruli, the anatomical and functional units of the AL **(Flanagan and Mercer, 1989)**. Thus, odorant detection results in the stimulus-specific activation of a subset of glomeruli, creating a stereotypical map of odour-induced glomerular responses **(Galizia et al., 1999; Sachse et al., 1999)**. The signal processed in the AL is forwarded by ~800 output neurons - the projection neurons (PNs) - to higher-order brain centres: the mushroom bodies (MBs), central, paired structures dedicated to multisensory integration, memory storage and retrieval **(Heisenberg, 2003; Stopfer, 2014)**, and the lateral horn of the protocerebrum, a diffuse bilateral structure involved in valence coding of odorants **(Jeanne et al., 2018; Roussel et al., 2014; Strutz et al., 2014)**. The MB architecture is defined by the layout of its ~185 000 intrinsic neurons, the Kenyon cells (KCs). Each KC extends its dendritic arborisation within the MB input regions, termed the calyces, where it receives input from multiple PNs, and projects its axon to the MB output region, termed the pedunculus, where it bifurcates into the vertical (α and γ) and the medial (β) lobes **(Mobbs, 1982; Strausfeld, 2002)**. Kenyon cells integrate, among others, the excitatory input of olfactory PNs and the inhibitory input of recurrent GABAergic feedback neurons **(Ganeshina and Menzel, 2001; Grünewald, 1999; Rybak and Menzel, 1993; Zwaka et al., 2018)**. The latter plays a critical role in shaping KCs olfactory responsiveness and maintaining a sparse output over a wide range of odorants and concentrations **(Papadopoulou et al., 2011; Stopfer, 2014)**. Antennal lobe, mushroom body and lateral horn are innervated by the VUMmx1 neuron **(Hammer, 1993)**, an unpaired octopaminergic neuron conveying appetitive reward-related information to the olfactory circuit and mediating reward-based olfactory memory formation. The activity in the Kenyon cells is integrated by MB output neurons (MBONs). These neurons exhibit which show learning-related plasticity and provide valence-loaded information to pre-

motor areas **(Aso et al., 2014a; Okada et al., 2007; Schmalz et al., 2022; Strube-Bloss and**

Rössler, 2018).

The neural representation of odorants in the AL has been extensively described by

means of functional calcium imaging **(Paoli and Haase, 2018)**. *In vivo* imaging allowed

observing that each stimulus is represented with a specific pattern of excitatory and inhibitory

responses across the glomeruli of the AL, **(Galizia et al., 1999; Sachse and Galizia, 2002;**

Sachse et al., 1999), and that perceptually similar odorants elicit similar glomerular response

patterns **(Guerrieri et al., 2005)**. Calcium imaging of AL activity has been typically conducted

at low temporal resolutions (~100-200 ms) **(Locatelli et al., 2016; Mertes et al., 2021;**

Nouvian et al., 2018; Sachse and Galizia, 2003). This is partially justified by the slow

dynamics of fluorescent calcium sensors, which resulted - in general - in the compression of

olfactory representation into a spatial vector of glomerular response intensity, with a

concomitant loss of information on neural response dynamics. While calcium signal decay is

relatively slow (>100 ms), its onset is fast (<10 ms) **(Helassa et al., 2015; Moreaux and**

Laurent, 2007), and provides the possibility for investigating parameters such as glomerular

response latency **(Junek et al., 2010; Paoli et al., 2018)**, signal frequency components **(Paoli**

et al., 2016) and interglomerular information transfer **(Chen et al., 2023; Paoli et al., 2023b)**.

In this study, we significantly improved the temporal resolution of calcium imaging recordings

of olfactory neurons in the bee brain by means of a resonant scanning multiphoton microscope.

This allowed us to record calcium activity at a <10-ms resolution, preserving temporal

information of the olfactory code and yielding a more realistic representation of odour

trajectories in the AL. First, we observed that odour representation changes dynamically during

and after an olfactory stimulation, resulting in specific odour and after-odour images. Then, we

showed that most glomerular response profiles presented an inhibitory component, suggesting

that glomerular activity is strongly shaped by local inhibition, with only a minor influence - if any

- of local excitatory interneurons.

Calcium imaging analysis was combined with a modelling approach to investigate how

odorant representation evolves from the AL to the MB. Electrophysiology experiments have

provided insight on the phasic and sparse activity of Kenyon cells as well as into its oscillatory

nature **(Laurent and Davidowitz, 1994; Perez-Orive et al., 2002; Stopfer, 2014)**. However,

the unavailability of an imaging method allowing the visualisation of olfactory coding in a KC

ensemble prevented us from further understanding how odorants are represented within the

MB, *e.g.* how the neural representation of an odorant is transformed from the AL to the MB or

to what extent the similarity among odorant representation is maintained in the KC space. One

way to address these questions is by using neural network models constructed *via* the

abstraction of common features of MBs across insect species (*e.g.*, fruit fly, locust, honey bee)

to reproduce cognitive tasks such as stimulus discrimination or learning. Models can be built

by simulating multiple neurons interacting with each other according to physiological rules such

as the dynamics of action potentials or Hebbian synaptic plasticity **(Eschbach et al., 2020;**

Finelli et al., 2008; Gkanias et al., 2022; Huerta et al., 2004; Smith et al., 2008). Other

models simulate the MB neural network by considering the statistics of connectivity within the

neuropil, *e.g.* the ratio of PNs to KCs, the average number of synaptic connections, the

neuronal firing rate **(Ardin et al., 2016; Buehlmann et al., 2020; Le Moël et al., 2019; Peng**

and Chittka, 2017; Springer and Nawrot, 2021; Wystrach, 2023). Here, we followed the

second approach and built a simplified but realistic neural network model of the MB based on

the neuroanatomical and functional properties of the insect's olfactory circuit. The model

comprises three layers of neurons: (1) a MB input layer to provide the model experimentally

acquired time-series of PN activity; (2) a MB intrinsic layer, where the input signal is distributed

to a population of modelled KCs based on neuroanatomical and physiological data; (3) a MB

output layer, where one appetitive MB output neuron (MBON) receives input from the KC layer

and can be subject to learning-induced plasticity. Generally, the input for this type of model is

simulated based on the spatial and temporal statistics of the odour-induced glomerular activity

(Eschbach et al., 2020; Finelli et al., 2008; Gkanias et al., 2022; Huerta et al., 2004; Le

Moël and Wystrach, 2020; Peng and Chittka, 2017; Smith et al., 2008; Springer and

Nawrot, 2021). In this case, we fed the model with the time series of the PN responses

recorded via *in vivo* calcium imaging analysis from multiple individuals exposed to three

different odorants at the MB working frequency of 20 Hz **(Cassenaer and Laurent, 2007;**

Laurent and Naraghi, 1994).

Here, we show that an MB neural network model based solely on three simple

connectivity rules abstracted from insect studies - sparse connectivity, feedback inhibition, and

learning-induced synaptic modulation - is sufficient to explain fundamental MB processing

features such as improved olfactory discrimination and associative learning. The model also

predicts that the presence of post-stimulus inhibitory and excitatory activity leads to the

recruitment of a second pool of KC, resulting in a distinct after-odour representation in the MB.

Additionally, the model's performance in appetitive learning and across-stimuli generalisation

was coherent with empirical measurements obtained from behavioural protocols. Finally, using

real physiological datasets as input, we showed that such a model is robust to noise and

biological variability across multiple stimulus repetitions.

Results

154 Results

Response dynamics of projection neurons in high-temporal resolution

Response dynamics of projection neurons in high-temporal resolution

Antennal lobe PNs of eight bees were back-filled with the calcium sensor Fura-2 **(Paoli**

and Haase, 2018; Sachse and Galizia, 2002) to measure odour-induced glomerular activity

of (~25 glomeruli/AL) (**Fig. 1A**). Calcium signal was recorded upon stimulation with two

monomolecular odorants (1-hexanol, 1-heptanol) and a complex fragrance (peppermint oil) on

a 5s/25s ON/OFF conformation for 20 trials (**Fig. 1B-D**). Fast calcium imaging revealed the

high variability of odorant-elicited glomerular response profiles.

First, we assessed the stability of odorant representations across time and trials given

the duration of each imaging session (30 min) and the multiple stimulus repetitions. For each

individual, we calculated the Pearson's correlation coefficients across pairs of trials of the

glomerular response vectors before, during and after olfactory stimulation. **Figure 1E-E'** show

that across-trial correlation is elevated during odour arrival and remains around 0.5 after

stimulus termination. Moreover, a visual inspection of glomerular response profile dynamics

confirmed that, with our stimulation protocol, the temporal activation and inactivation of each

glomerulus appear to be conserved across stimulus repetitions (**Fig. S1**). This shows that, in

our experimental conditions, odour coding was stable throughout the entire imaging period.

All odorant response traces were pooled together (**Fig. 1F**) and clustered according to

their direction with respect to the pre-stimulus baseline (*i.e.*, excitatory or inhibitory) and

duration (**Fig. 2,** see **Materials and Methods**), allowing for a statistical description of

glomerular responses across individuals and odorants, and for the identification of classes of

recurrent profiles (**Fig. 2A,B**). The most common profiles comprised inhibitory (group 1) and

excitatory (group 4) responses lasting the entire olfactory stimulations and terminating after

odour offset. In some cases, they could result in prolonged inhibition (group 2) or excitation

(group 6) lasting after stimulus offset or could be followed by a post-stimulus excitatory (group

3) or inhibitory activity (group 5). Approximately 15% of responses consisted of short phasic

excitation followed by prompt signal termination (group 7) or even by an inhibitory response

(group 8). About 15% of all recorded glomerular traces showed no detectable odorant-induced

activity (group 9). Of all traces, 48% were excitatory, 37% inhibitory, and 15% unresponsive

(**Fig. 2C,D**). Moreover, 251 glomerular responses out of 546 (46%) displayed an inhibitory

component either during or after olfactory stimulation. Notably, the varied response types were

evenly distributed across the different combinations of bees and odorants, indicating an

absence of individual bias (**Fig. S2**).

The heterogeneity of PNs response profiles could be due to the diversified temporal

pattern of the ONSs input **(Kim et al., 2023)** and to second-order processing mediated by

lateral connections within the AL **(Girardin et al., 2013; Krofczik et al., 2009)**. We assessed

the contribution of these two components by measuring the latency of excitatory and inhibitory

glomerular responses (*i.e.* the timepoint t at which a response profile exceeds the threshold of

one standard deviation from the mean pre-stimulus activity). **Figure 2E** shows that excitatory

responses (groups 4 to 8) all share a similar onset (313 ± 22 ms, n = 263), which is shorter

than that of inhibitory profiles (groups 1 to 3; 351 ± 47 ms, n = 201) (Kruskal-Wallis test, $p<0.05$,

Tukey-Kramer multiple comparison correction). Moreover, the latency of short excitatory

responses' termination (groups 7 and 8; 346 ± 47 ms, n = 60) is coherent with the onset of

inhibitory profiles. These findings indicate that odour representation in the AL is shaped initially

by the excitatory input delivered by the OSNs and reshaped - approx. 40 ms later - by local

inhibition. Such an olfactory tuning is different from what is proposed in *Drosophila*, where both

excitatory and inhibitory local neurons contribute to moulding the neural correlate of an

olfactory input **(Chou et al., 2010; Olsen et al., 2007)**.

A mushroom body neural network model: Key principles

We constructed a simple but realistic MB neural network model based on the known

connectivity of this structure in the insect brain. The model architecture relies on three main

principles (**Fig. 3**): First, in each brain hemisphere, ~800 AL projection neurons (PNs) diverge

onto ~185,000 KCs **(Strausfeld, 2002)**. Neuroanatomical studies in bees **(Szyszka et al.,**

2005) and flies **(Ashok Litwin-Kumar et al., 2017; Caron et al., 2013)** suggest that each KC

is randomly innervated by approx. 7-10 PNs. Thus, we generated a MB network where ~25

PNs (*i.e.*, the average number of glomeruli imaged during a calcium imaging experiment)

diverge onto 1000 KCs, with each KC being innervated by ~8 PNs. Second, recurrent inhibitory

neurons such as the A3 feedback neurons in the bee **(Mobbs, 1982; Rybak and Menzel,**

1993; Szyszka et al., 2005; Zwaka et al., 2018), the APL neuron in fly **(Lin et al., 2014)**, and

the giant GABAergic neuron in the locust **(Papadopoulou et al., 2011)**, modulate MB KCs

firing rate so that less than 20% of KCs are active upon olfactory stimulation, and only ~5%

are stimulus-specific **(Honegger et al., 2011; Turner et al., 2008)**. Hence, we enriched the

model with a winner-takes-it-all feedback inhibition mechanism by forcing that only 10% of KCs

receiving the largest summed input generate an action potential. Third, as in bees and locusts,

MB have a 20-Hz oscillatory cycle **(Cassenaer and Laurent, 2007; Laurent and Naraghi,**

1994; Popov and Szyszka, 2020), we fed the model with experimentally recorded calcium

signals resampled at a 20 Hz-sampling frequency. This approach allowed testing if a simple neural architecture - so far challenged with simulated datasets **(Ardin et al., 2016; Buehlmann et al., 2020; Peng and Chittka, 2017)** - could be used for understanding how real odour activity recorded at the level of PNs is transformed in the MB, and if the rules governing the model were sufficient to support appetitive olfactory conditioning.

Model input and output: Simulating olfactory coding in the mushroom body

For each recorded individual AL ($n = 8$), an MB simulation was generated and fed with the time series of the recorded glomerular activity obtained for that individual. Thus, for each odorant and individual, we simulated the time course of the firing of a virtual KC ensemble (**Fig. 4A**). The analysis of KC population turnover (**Fig. 4B**) showed that, before odorant onset, the population varies randomly between adjacent time points while it stabilises during olfactory stimulation, with a turnover rate of ~10%. The dynamics of KC recruitment across bees and odorants (**Fig. 4C**) indicates that two odorant-specific sets of cells are recruited during an olfactory stimulation: the first one at stimulus onset, the second one at odour termination (see **Fig 4C**, $t = 5$ s). Both populations are rather stable, with a turnover rate below the pre-stimulus baseline for up to 15 s after stimulus offset (turnover rate$_{pre-stim}$ = 0.61 ± 0.14; turnover rate$_{ON\_0-5s}$ = 0.17 ± 0.05; turnover rate$_{OFF\_5-10s}$ = 0.25 ± 0.04; turnover rate$_{OFF\_10-15s}$ = 0.33 ± 0.08). Correlation analysis of stimulus-induced responses (**Fig. 5D**,**E**) showed a high correlation of neural activity across time points during olfactory stimulation and during the post-odour window both in the recorded PN population and in the simulated KCs. This indicates that the neural representation of an odorant is sufficiently stable to provide a neural image, which is coherent with itself during a 5-s stimulation window and during a ~10-s post-stimulation window. Furthermore, the across-odorants correlation is rather high in the PN space (Pearson's correlation coefficient, $r = 0.4$), but proximal to zero in the KC space, supporting the notion that the MB network increases stimulus discriminability, not only during but also after odour offset.

To further corroborate this idea, we compared the temporal relative trajectories obtained

for the three odorants in the measured PN space and in the modelled KC space (**Fig. 4F**). In

both cases, response trajectories diverged at stimulus onset and returned to the centre of

space after stimulus termination. However, the principal component analysis showed that

odorants are better separated from each other and that they remained separated for a longer

time in the KC space compared to the PN space. This indicates improved and prolonged

stimulus discriminability in the MB with respect to the AL (**Fig. S1**).

We next investigated the discriminatory power of the system across a larger set of odour

response profiles. For this, we pooled together all response profiles acquired during the

calcium imaging analysis (**Fig. 1F**) and we artificially combined them to simulate 100 odorant

response vectors, each comprising 30 different glomerular profiles (see **Methods**). Using

these artificially combined odorant response profiles, we simulated the relative KC response

time series and measured the correlation across all response vectors in the PNs (experimental

data) and KCs spaces (modelled data) (**Fig. 4G**). Notably, while response vectors calculated

from PN data were highly correlated ($r = 0.71 \pm 0.08$), such a correlation significantly decreased

in the modelled KC space ($r = 0.52 \pm 0.10$; Student's t-test between r distributions of PN and

KC data: $p \approx 0$, $n = 4774$), indicating that the signal transformation operated by the proposed

MB neural architecture induces a strong decorrelation among odour signatures.

Overall, the proposed MB network can process experimentally acquired input data and

produce physiologically plausible KC response patterns coherent with *in vivo* KC activity

measurements **(Lüdke et al., 2018)**. Moreover, it accounts for the expansion of the coding

space from the AL to the MB **(Lin et al., 2014; Papadopoulou et al., 2011; Stopfer, 2014)**,

enhancing inter-stimuli decorrelation and confirming previous theoretical works based on

artificial datasets **(Olshausen and Field, 2004; Peng and Chittka, 2017)**. Importantly, these

emergent properties of the model architecture are based on the known neuroanatomy and

physiology of the insect brain and have been obtained without any optimisation aiming at

reproducing specific features of olfactory coding.

 The neural network model predicts appetitive behaviour

Appetitive classical conditioning relies on the coincidental activation of the neural

elements representing the conditioned stimulus (CS) and the unconditioned stimulus (US). The

latter is mediated by various neuromodulators, such as specific dopaminergic, as shown in the

fly **(Burke et al., 2012; Liu et al., 2012)**, or octopaminergic neurons, as demonstrated in the

honey bee **(Hammer, 1993)**.

The coincidental activation of CS and US neural elements induces plastic modulations

in the MB output neurons (MBONs) **(Hige et al., 2015; Okada et al., 2007; Owald et al.,**

2015). Here, we assessed if the proposed MB model could reproduce empirical measurements

of appetitive learning by introducing a time window, during which the weights of the recruited

KC-to-MBON synapses could be downregulated from 1 to 0 **(Ardin et al., 2016)**. This rule

reflects MBON learning-induced plasticity, according to which they display a broadly tuned

response, which is reduced in presence of a learned stimulus **(Amin and Lin, 2019; Aso et**

al., 2014a; Aso et al., 2014b; Cognigni et al., 2018; Cohn et al., 2015; Hige et al., 2015;

Lyutova et al., 2019; Okada et al., 2007; Owald et al., 2015). To implement such a rule, we

introduce a synaptic plasticity threshold (*spt*) parameter, which defines the number of firing

events of a given KC upon which its synaptic output weight is reduced from 1 to 0. In other

words, it determines how active a synapse should be during the learning window before being

switched off.

It is well known that the inter-stimulus interval (ISI) - *i.e.* the time elapsed from CS to US

onsets - can influence the efficiency of Pavlovian learning **(Domjan, 2015; Holland, 1980)**.

The calcium signal dynamics and modelling analysis presented here (**Fig. 4A-F**) suggest that

such an ISI-dependent effect is the consequence of the strong temporal dynamics of the

odorant neural representation. As the neural representation of the CS changes in time, a

reward system recruited at different time points will interact with a different population of CS-

recruited KCs. Because learning results from the coincidental activation of CS and US

elements **(Pavlov, 1927)**, an individual should respond to the different time points of an

olfactory stimulation with different strengths, depending on the US arrival time experienced

during conditioning. To test this hypothesis, learning-related MBON plasticity was modelled for

four different CS/US temporal contingencies reflecting four conditioning paradigms, well-

studied in honey bee **(Giurfa and Sandoz, 2012)** (**Fig. 5A**). Having a fixed CS stimulation

interval from t = 0 to 5 s, and a learning window (US) of 3 s, we simulated MBON learning with

the following ISIs: *backward* pairing occurred when the US initiated at t = -2 s (*i.e.*, 2 s before

CS onset), *early* pairing when it started at t = 1 s (*1 s* after CS onset), *delay* pairing when it

started at t = 4 s (4 s after CS onset), and *trace* pairing when the US started at t = 7 s (2s after

CS offset). First, we trained the model with the experimental PN response profile recorded for

either 1-hexanol or peppermint oil (see **Fig. 1**). One odorant was used as CS and the other as

a novel odorant to test the specificity of the learning obtained under these experimental

conditions. After learning, we modelled the MBON action potential (AP) probability upon the

presentation of the CS (*e.g.*, 1-hexanol) and the novel odorant (*e.g.*, peppermint), and repeated

this operation for the response profiles of all individuals. To account for the biological variability

in the neural activities elicited by repetitions of the same stimulus, learning was assessed

against the KC response profiles resulting from five CS and five novel odorant presentations,

which were not included in the training dataset. As expected, we found that a trained MBON

showed a larger decrease in firing probability when presented with the CS than with the

unfamiliar stimulus, demonstrating learning (**Fig. 5B**). Interestingly, the model predicts weak

learning in the case of *backward* pairing - that is when US and CS are partially overlapping,

but the US begins before the CS - whereas both *early* and *delay* conditioning predict strong

learning, that is, a large CS-specific down-regulation of MBON firing probability (**Fig. 5B**).

Finally, in the case of *trace* conditioning, the model also produced a down-modulation of the

MBON response. However, this learnt response was shifted in time: it appeared only after the

CS offset, that is, around the US expected arrival time (**Fig. 5B**).

In addition, according to the model, high synaptic plasticity sensitivity (low *spt* values)

facilitates non-specific learning, resulting in a strong generalisation effect. Conversely, MBON

plasticity under high *spt* values requires a robust and stable KC activation. This scenario is not

compatible with a learning interval that is only partially overlapping with the stimulation window (note that measurements are provided at 20 Hz, and in this context, a *spt* = 40 requires a synapse to be activated during two whole seconds - although not continuously - to be switched off) (**Fig. S2**).

Stimulus representation is dynamic (see **Fig. 1** and **2**). Hence, our model predicts that an MBON exposed to an *early* protocol should learn to identify the early components of the stimulus neural signature. Indeed, the model shows that the learned response (a down-regulation of MBON firing probability) occurs earlier in the *early* than in the *delay* contingency (Wilcoxon test for non-parametric paired data, $p < 0.001$; $n = 16$) and with lower response latency variability (Barlett's test *early* vs *delay*, $p < 0.011$) (**Fig. 5D**). At the same time, it predicts that *trace* conditioning should rely on the identification of post-stimulus neural activity (**Fig. 5B**). As such, the learned response occurs after odour offset and with a variability comparable to the one detected for the *delay* protocol (Barlett's test *delay* vs *trace*, $p = 0.1808$) (**Fig. 5D**).

To test the model's predictions, we measured the timing of real behavioural responses (*i.e.*, the proboscis extension) of bees trained under the same CS/US conditions and tested 1 hour later with the CS and a novel odorant. Because the model's predictions were generated based on the odour response maps elicited in the AL by 1-hexanol and peppermint oil, the same odorants were used for the behavioural experiments. In agreement with the model's predictions and previous reports **(Felsenberg et al., 2014)**, *backward* conditioning led to scarce and unspecific responses (**Fig. 5E**). In contrast, positive ISIs of 1 or 4 seconds (*early* and *delay* protocols) induced specific memory formation (McNemar test, $p_{backward}$ = n.s., p_{early} = 0.003, p_{late} = 0.004), with ~90 to 70% of the trained bees showing conditioned responses to the CS and ~40% showing a generalised response to the unfamiliar stimulus during the test. Video tracking of the proboscis extension response showed that bees receiving the reward 1 s after CS onset extended the proboscis earlier and with less variability than those that received the reward towards the end of CS delivery (Kruskal-Wallis test, $p = 0.036$; Barlett's test for data variance comparison, $p = 5e10^{-6}$; n_{early} = 26, n_{delay} = 20) (**Fig. 5F**). Finally, bees exposed to *trace* conditioning (ISI = 7 s) showed weak and aspecific learning - highlighting the

complexity of the task **(Dylla et al., 2013; Ito et al., 2008; Paoli et al., 2023a; Szyszka et al., 2011)**. However, in opposition to the model's prediction, the learners showed a PER latency comparable to the *delay* group (Kruskal-Wallis test, $p = 0.992$) but with a broader distribution (Barlett's test for data variance comparison, $p = 0.007$; $n_{delay} = 20$, $n_{trace} = 31$).

Relative contribution of glomerular response dynamics to conditioned response latency

Early excitatory responses are confined to the initial part of the stimulus, while inhibitory responses - with a slower and delayed onset - show a delayed onset with respect to the excitatory responses. As such, they both contribute to the creation of an early (<1.5 s) and a delayed/stable (>1.5 s) odour signature (**Fig. 2**). Such a temporal structure might provide the neural basis for the difference in response latency observed upon *early* and *delay* conditioning both in behavioural and modelled data. If so, the absence of such dynamics might increase odour response stability and prevent the differential response observed in MBONs subject to the two conditioning protocols.

To test this hypothesis, we used the glomerular response database (**Fig. 1F**) to generate a set of simulated odorant responses, each comprising 32 glomerular response profiles selected among the different response groups (**Fig. 2A** and **B**) based on the average occurrence of each group across all recorded odour response maps (see **Methods** section). As we did with real odorants (**Fig. 5**), we used the simulated odorant responses to train modelled MBONs with *early* and *delay* protocols. Then, we assessed MBONs' firing rate upon stimulation with the conditioned and a novel stimulus. As with real odorant responses (**Fig. 5**), also when using artificially generated odorant response profiles, MBONs showed an anticipated response when trained with the *early* protocol (**Fig. 6A-C**, first row).

Next, we evaluated the relevance of the different glomerular response types in determining the different MBON response latency occurring upon training with the different CS/US contingencies. We selectively removed from the generated odorant response profiles all stable excitatory responses (groups 4 to 6; **Fig. 6A-C**, second row), inhibitory responses

(groups 1 to 3; **Fig. 6A-C**, third row), short excitatory responses (groups 7 and 8; **Fig. 6A-C**, fourth row), and both short excitatory and inhibitory responses together (groups 1 to 3, 7, 8; **Fig. 6A-C**, fifth and last row). These modified versions of the simulated odorants were used to model MBON response dynamics upon *early* and *delay* conditioning. Interestingly, removing stable excitatory or inhibitory response profiles or short excitatory responses alone did not affect the difference in response timing between protocols. On the contrary, the removal of stable components such as long excitatory and inhibitory profiles seemed to enhance such difference in response latency (**Fig. 6D**). Conversely, the combined removal of inhibitory and short excitatory responses from the odorant response patterns prevented the difference in MBON response latency between protocols.

In conclusion, reducing the richness in glomerular response dynamics by removing early excitatory responses and delayed inhibition increases the temporal stability of the neural representation of an olfactory stimulus and prevents the differential response latency observed between *early* and *delay* conditioning protocols.

Discussion

Our results show that a simple but biologically realistic MB neural network model simulating essential aspects of MB connectivity can process real neurophysiological input and generate robust predictions that agree with empirical observations. Our model was fed with experimental PN responses acquired at high temporal resolution **(Bestea et al., 2022; Paoli and Haase, 2018)**. This approach provided a better description of the temporal dynamics of olfactory coding and a more realistic modelling of odour signal transformation and learning.

 Olfactory trajectories in the glomerular space are shaped by local inhibition

Because of the high temporal resolution of the functional imaging recordings, we could

reliably measure odorant-induced latencies of excitatory and inhibitory PN responses.

Excitatory responses occurred earlier than inhibitory ones and exhibited a smaller temporal

variability. This suggests that excitation of PNs occurs in a single wave upon cholinergic input

from the OSNs to the glomeruli of the antennal lobe, with a minor contribution - if any - of

second-order excitatory local interneurons within the AL, which have been reported for

Drosophila **(Chou et al., 2010; Olsen et al., 2007)** but so far not for honey bees **(Girardin et**

al., 2013; Krofczik et al., 2009; Schäfer and Bicker, 1986). This finding also advocates that

odour responses of OSNs may not display onset-latency variability as shown in the locusts

(Kim et al., 2023). On the other hand, inhibitory PN responses and the termination of early

excitatory PN responses occurred within the same time window, approx. 40 ms after the onset

of excitatory responses, suggesting that lateral inhibition within the AL is responsible for

triggering PN inhibition and terminating early excitatory responses (see **Fig. 2**, groups 7 and

8). This is consistent with a significant presence of GABAergic local interneurons in the AL

(Schäfer and Bicker, 1986), which reshape the olfactory message conveyed to higher-order

brain structures such as the MBs and the lateral horn **(Paoli and Galizia, 2021)**. Notably, we

observed that almost 50% of PN responses recorded contain an inhibitory component (see

Fig. 2A, groups 1-3,5,7,8), underlining the dominant role of inhibitory AL interneurons in

shaping odour trajectories in the glomerular space. These observations are in contrast with

previous calcium imaging analyses, where only 2-10% of responses had an inhibitory nature

(Krofczik et al., 2009; Szyszka et al., 2005), but coherent with electrophysiological recordings

showing a rich response diversity and prominence of inhibitory activity **(Wilson et al., 2004)**.

Differently from *Drosophila*, where both local inhibitory and excitatory interneurons contribute

to the olfactory tuning **(Chou et al., 2010; Huang et al., 2010; Shang et al., 2007; Yaksi and**

Wilson, 2010), our analysis indicates that, in the honey bee AL, glomerular responses are

modulated mainly by a local inhibitory network. This conclusion is supported by a previous

study showing that pharmacological activation of a single glomerulus by topical acetylcholine

application provokes inhibitory but not excitatory responses in other glomeruli **(Girardin et al.,**

2013). Indeed, while the presence of excitatory local interneurons has been predicted **(Malaka**

et al., 1995), such prediction has never been supported by experimental evidence. A

population analysis suggests that the varied response profiles are equally distributed across

individuals and can be triggered by different olfactory stimuli. Still, it remains to be understood

to what extent the response profiles for each odour/glomerulus combination are conserved

across individuals.

Overall, these findings highlight possible differences between the ALs of bees and fruit

flies regarding the nature of the local interneuron populations modulating glomerular activity.

Thus, they cast caution in prioritising a single species as a representative general model of

insect olfaction.

Smell and aftersmell: An odorant-specific afterimage

Almost 50% of all PN responses in our database displayed either a prolonged post-

stimulus activity (see **Fig. 2**, groups 2 and 6) or a change in response direction after stimulus

termination (groups 3, 5, 7, 8). This results in the recruitment of an offset-specific population

of KC - both in our model and in the literature **(Ito et al., 2008; Lüdke et al., 2018)** - that

creates a neural image of the post-odour in the MB. Thus, the post-odour representation is

different from the neural image of the odorant itself and provides a stable, slowly decaying

after-image, which is stimulus-specific and temporally associated with the neural correlate of

the odorant itself. This is in agreement with the analysis conducted by Patterson *et al.*

(Patterson et al., 2013) on the olfactory representation in M/T cells in mice, which shows that

the post-odorant activity is rather conserved and stimulus-specific even across concentrations

and for about a 10-s window. Our modelling analysis predicts that the after-stimulus signature

of an odorant is more distinct and stable across time in the MB than in the AL (**Fig. S1**). This

could occur because the activity across the KC population is normalised by the MB feedback

neurons operating a gain control, which stabilises odorant-responses signature in the MB - including the post-odorant activity - while the input signal in the PN is fading.

The existence of a post-odour signature has important implications for understanding the neurobiological basis of trace learning **(Dylla et al., 2013; Paoli et al., 2023a)**, a conditioning protocol in which a stimulus-free time window separates CS and US. Previous studies reported that an odorant cannot be predicted by its post-odour signature **(Lüdke et al., 2018; Szyszka et al., 2011)**. Nevertheless, as shown in previous studies **(Galili et al., 2011; Lüdke et al., 2018; Patterson et al., 2013; Szyszka et al., 2011)**, the neural activity associated with the OFF response is stimulus-specific and reproducible, suggesting that the post-odour could be physiologically similar to the onset of a second odorant (in the same way as the back-taste of wine could remind us of an unrelated ingredient). If so, the odour/post-odour pattern could be processed as an individual neural object, effectively extending the duration of the nominal, conditioned odour. Alternatively, it is also possible that the stimulus specificity of the after-smell signature and its temporal contiguity with the CS may enable some form of second-order conditioning. Certainly, the behavioural analysis within this study indicates that honey bees subjected to trace conditioning display the conditioned reflex already during odour arrival. This indicates that bees do not form simply the association between the after-smell (as CS) and the sugar reward (US). Conversely, it supports the idea that trace learning requires some form of prolongation of the neural trace of the CS until US arrival **(Dylla et al., 2013; Paoli et al., 2023a)**. Interestingly, the duration of the post-odour neural signature reported by us and by previous studies **(Galili et al., 2011; Patterson et al., 2013; Szyszka et al., 2011)** is about 10 to 15 s. As such, it is compatible with the duration of the CS/US gap in trace learning experiments **(Galili et al., 2011; Paoli et al., 2023a; Szyszka et al., 2011; Wystrach et al., 2020)**, suggesting the possibility of its implication in this form of learning.

 Mushroom body modelling predicts associative learning scores and response latency for

 different conditioning protocols

To investigate if the proposed neural network model explains the temporal features of

associative learning, we simulated CS delivery by feeding the model with experimental PN

calcium imaging time series and applied a learning rule based on the downregulation of

recurrently recruited synapses in the MB. A learning time window of 3 s was defined according

to four well-studied protocols for associative conditioning, which correspond to four different

ISI situations (-2 s, +1 s, +4 and +7 s) **(Bitterman et al., 1983; Giurfa and Sandoz, 2012)**. A

simulation of the action potential probability of an MBON after olfactory conditioning revealed

that differentiation between the CS and a novel odorant was observable whenever the CS

anticipated the US (ISIs of +1 s and +4 s) but less successful in the *backward* (ISI of -2 s) and

trace protocols (ISI of 7 s). Generalisation errors were also observed in a similar way as in

behavioural experiments. Notably, in this study, we adopted a backward contingency with a 1-

second overlap between the US and the CS (**Fig. 5A**). This differs from the more traditional

configuration, where the CS is delivered after US termination, that is, without any temporal

overlap. In this case, the model would predict no learning at all, as observed in honey bee

behavioural experiments **(Felsenberg et al., 2014; Hellstern et al., 1998)**. Hence, we tested

the model's prediction in the form of *backward* conditioning where US and CS are partially

overlapping. In this case, the model predicts a weak appetitive association (see **Fig. 5B,C,E**).

This result is coherent with a previous behavioural report, where bees subject to a similar

protocol showed a weak response to the conditioned odorant **(Felsenberg et al., 2014)**. In the

early and *delay* configurations, a CS-specific associative memory was formed, both in the

model (**Fig. 5C**) and in the behavioural tests (**Fig. 5E**). However, the model predicts a

difference in response latency and latency variability depending on relative CS/US arrival (ISI

of +1 or +4 s): a learning window placed closer to stimulus onset (*early* protocol) resulted in

early and temporally precise MBON responses, whereas a delayed learning window produced

a similar learning score, but with higher response latencies and temporal uncertainty.

Experimental results based on the same odorants that were fed to the model confirmed the model's predictions. They showed that *early* and *delay* protocols yielded comparable results in terms of learning score and CS-specificity while differing in terms of CS latency and precision (**Fig. 5C-F**). These observations support the idea that associative learning occurs upon coincident detection of the CS and US neural elements. In fact, because the CS neural representation changes dynamically during the olfactory stimulation, a positive association with the initial odour signature will trigger an earlier and temporally precise conditioned response. Conversely, the association with the later part of the olfactory signature will elicit a delayed response with greater temporal variability.

Finally, both the model and data (**Fig. 5C** and **E**) show that *trace* conditioning yields lower and less specific learning scores. However, while the model predicts that learners should respond around the expected US arrival time (*i.e.*, during the after-smell component, **Fig. 5D**), empirical data show that the individuals who learned the association extend their proboscis during (and not after) the olfactory stimulation. Such a discrepancy in measured and predicted response latencies suggests that trace conditioning cannot be explained by the simple temporal coincidence of a reward system's activation and a pool of KC recruited at odour offset. In order to trigger a conditioned response anticipating the expected reward time, the network must have learned the nominal CS stimulus and not simply its after-smell. Thus, a higher-order cognitive process, possibly involving attention mediated by the serotonergic system, could contribute in prolonging the memory trace of the conditioned stimulus **(Paoli et al., 2023a; Zeng et al., 2023)**.

Vrontou *et al.* (2021) showed that activation of the fruit fly dopaminergic system at the end of a stimulus or after odour offset can regulate the MBON spiking rate already at odour onset **(Vrontou et al., 2021)**. This suggests that delayed reward system activation can modulate an olfactory stimulus's early neural coding. Consequently, the CS/US temporal contingencies used in studies on *delay* and *trace* protocols may trigger an earlier-than-expected behavioural response by inducing learning-related neural plasticity in the early components of a CS response.

Pavlov (1927) proposed that the timing of a conditioned response could be adjusted to

coincide with the expected arrival of the unconditioned stimulus **(Pavlov, 1927)**. In his view,

this capability involved two cognitive processes: learning to suppress the responses to the

earlier phases of the conditioned stimulus and - at the same time - learning to delay the

conditioned response until the US onset. Drew *et al.* showed that in goldfish, the timing of the

conditioned response is influenced by the CS/US inter-stimulus interval, and its precision

improves across learning trials **(Drew et al., 2005)**. Importantly, this improvement was due to

increased accurately timed responses rather than decreased off-time ones. Unlike in Pavlov's

hypothesis, these findings suggested that timing acquisition during associative conditioning

relies on learning to respond at the correct moment rather than to avoid responding at the

wrong time. Experimental evidence in mammals indicates that their ability to adjust the latency

of the conditioned response based on the US/CS relative onset depends on the cerebellar

neural network **(Perrett et al., 1993)**. Targeted lesions to the cerebellar cortex significantly

impair this ability, suggesting that the adaptive timing of conditioned responses relies on a

different mechanism than the one supporting the association. Although the neural mechanisms

governing timing in the cerebellum are not yet fully understood, the involvement of this brain

area in sensory-motor timing is well-established **(Kirkpatrick and Balsam, 2016; Paton and**

Buonomano, 2018).

While the influence of the inter-stimulus interval on learning is well documented also in

invertebrates **(Giurfa and Malun, 2004; Szyszka et al., 2011; Vogt et al., 2015)**, the ability of

relatively small insect brains to time the conditioned response based on the expected reward

arrival remains unclear. Previous studies have shown that the duration of the inter-stimulus

interval experienced during learning can influence the conditioned response's latency in

bumble bees **(Boisvert and Sherry, 2006)** and honey bees **(Szyszka et al., 2011)**. Both

studies indicate that bees trained with a longer ISI exhibited a larger latency in their response

to the conditioned stimulus. This provided clear evidence that CS-related information can

modulate response timing. However, the appetitive response largely anticipated the expected

reward time, and a clear interpretation of time tracking in insect brains is still missing.

Our model shows that this time contingency can be naturally encoded simply because

the olfactory representation in the KCs evolves through time in a deterministic fashion. Thus,

what is learnt during the US presentation is the KC configuration of the CS at this particular

time. In other words, the MBON learns which subpopulation of KC is active during reward

arrival rather than the expected reward time. Interestingly, because the KC representation of

the CS does not evolve linearly with time (**Fig. 4A-C**), the CS/US time contingency is not

encoded perfectly. Remarkably, the model captures the experimental measurements of the

proboscis response latency in bees, which indeed tend to respond earlier than the expected

reward (**Fig. 5D**,**F**). However, such reasoning may not apply to trace conditioning protocols -

when a temporal gap separates CS offset and US onset. In this case, higher-order neural

processes that are not captured by our model may contribute to extending the stimulus-related

information required for associative learning **(Paoli et al., 2023a)**.

As often in insect research, heuristics can provide simple solutions to apparently complex

cognitive problems. In this case, a mechanism based on the temporal dynamics of the

conditioned stimulus, explains the bees' systematic errors at encoding proper CS/US

contingency. Whether, in addition to this process, other mechanisms enable bees to track time

contingencies remains to be seen.

Conclusive remarks

In this study, we adopted a fast calcium imaging approach to investigate the dynamical

representation of odorants in the antennal lobe. Our findings revealed that about 50% of

glomerular responses had an inhibitory component and that local inhibition played a crucial

role in shaping the initial PN activity and defining odour trajectories in the glomerular space.

We observed that half of the PN response profiles prolonged their activity or changed response

direction after stimulus termination, generating a stable, odorant-specific after-smell. Although

the after-smell might be very different from the odour signature, it is specific and temporally

contiguous, thus providing the means to prolong the neural representation of the stimulus itself.

To investigate the logic of olfactory transformation and learning in the mushroom body,

we constructed an MB neural network model based on the known connectivity in the insect

brain. We showed that the model can process experimentally acquired input data and produce

realistic KC response patterns. Our analysis suggests that the mushroom bodies enhance

olfactory discrimination ability and stabilise the olfactory after-image, effectively prolonging the

stimulus signature after odour offset.

Finally, we implemented a learning rule to modulate MB output neurons firing rate based

on associative learning. We verified the model's predictions with behavioural measurements

on real honey bees subjected to different learning protocols. The coherence between

measured and predicted behavioural scores suggested that the rules behind this neural model

- *i.e.* random connectivity, feedback inhibition, and MBON plasticity - are sufficient to generate

relevant predictions for associative learning upon different learning protocols. In addition, by

testing the model against physiological recordings of different stimulations, we showed that

the model's predictions are robust to noise and biological variability.

Because the model incorporates not only neural assumptions pertaining to the honey

bee olfactory system but also to the fruit fly and locust olfactory systems, similar principles may

guide odour signal transformation at the level of the MB in appetitive associative learning

across insects. This generality does not apply, however, to the nature of odour reshaping by

local interneurons observed in the AL of bees, which seems to differ from that occurring in flies,

thus highlighting the fact that convergence, but also differences, can characterise the

functional architectures of olfactory systems in different insect species.

Methods

Experimental Model

Experiments were performed on honey bees *Apis mellifera* reared in outdoor hives at the experimental apiary of the Research Centre on Animal Cognition (CNRS, Toulouse, France) situated in the campus of the University Paul Sabatier. In all cases, honey bee foragers (>3-week old) were used. No institutional permission is required for experimental research on honeybees

Projection neurons labelling

Honey bee foragers were collected the day before the experiment at an artificial feeder, to which they were previously trained, and projection neurons were labelled for calcium imaging analysis as previously described **(Paoli et al., 2017; Sachse and Galizia, 2002)**. Shortly, bees were fixed in a 3D-printed holder with soft dental wax, antennae were blocked frontally with a drop of eicosane (Sigma Aldrich, CAS: 112-95-8). A small window was opened in the head cuticle, and glands and tracheas were displaced to expose the injection site. The tip of a borosilicate glass needle coated with Fura-2-dextran (Thermo Fisher Scientific Inc.) was inserted between the medial and lateral mushroom body calyces, where medial (m-ACTs) and lateral antenno-cerebral tracts (l-ACTs) cross. After dye injection, the head capsule was closed to prevent brain desiccation, and bees were fed *ad libitum* with a 50% sucrose/water solution. On the following day, antennal lobes were exposed to allow optical access, and the brain was covered in transparent two-component silicon (Kwik-Sil, WPI). Although the injection procedure may result in variable levels of PN labelling, the reproducibility of response amplitudes and the lack of labelling bias for specific glomeruli indicate that the loading procedure is reproducible.

 Calcium imaging analysis and signal processing

Undiluted solutions of 1-hexanol (Sigma-Aldrich, CAS:111-27-3, 7.98 M), 1-heptanol

(CAS:111-70-6, 7.05 M) and peppermint oil (CAS: 8006-90-4, 898 g/L) were delivered to the

bees using an Arduino Uno®-controlled automated olfactometer **(Bestea et al., 2022; Raiser**

et al., 2017). Briefly, 1 mL of pure odorant was placed in a 20-mL glass vial. During an olfactory

stimulation, the headspace concentration of volatile odorants is injected into a clean airflow,

where it is diluted approximately 10 times before reaching the honey bee antennae. The

odorous flow exiting the olfactometer is weak but detectable by a human nose, in the range of

honey bee olfactory sensitivity **(Carcaud et al., 2016; Gil-Guevara et al., 2022; Wright and**

Smith, 2004). Odorants were alternated and presented 20 times on a 5/25 second ON/OFF

configuration. Calcium imaging recordings were conducted with a straight Leica SP8 scanning

microscope (Leica Microsystems, Germany) equipped with a SpectraPhysics InSight X3

multiphoton laser tuned at 780 nm for Fura-2 excitation. All images were acquired with a water

immersion 16x objective (Leica HC FLUOTAR 16x/0.6 IMM CORR, Leica Microsystems,

Germany), at 64x64 pixel resolution and ~127 Hz.

Calcium imaging data were analysed with custom-made MATLAB (MathWorks Inc.)

scripts. The baseline signal, calculated as the mean fluorescence during the one second

before stimulus onset, was used to calculate baseline-subtracted and normalised stimulus-

induced glomerular activity ($\Delta F/F$). The normalised activity was multiplied by -1 to display

excitatory/inhibitory responses as positive/negative relative changes ($-\Delta F/F$). Time series of

glomerular activity were averaged across stimulus repetitions for response profile clustering

analysis. Glomeruli were hand-selected based on a morphological and functional map. To

facilitate glomerular identification, the odour response maps for the three stimuli were merged

into an RGB image: 1-hexanol was set as the red channel, 1-heptanol as the green, and

peppermint as the blue. Thus, glomeruli excited only by peppermint appear in blue, glomeruli

excited only by 1-hexanol appear in red, and glomeruli excited by a combination of the three

channels will result in different colour shades. Glomerular response profiles were calculated

as the mean intensity of an area of 5-by-5 pixels around the centre of the glomerulus across time. Stimulus representation stability (**Fig. 1E**) was assessed with a Person's correlation analysis between vectors of the mean glomerular response of different trials. Glomerular response vectors' stability was calculated before, during and after olfactory stimulation.

Mushroom Body Modelling

We used MatLab R2018a to investigate how the recorded PN activity would affect KC responses of a MB network model.

Projection neurons to Kenyon cells connectivity

Real projection neurons' activity acquired *via* calcium imaging analysis provided the input to a simple model of the mushroom body neural circuits. For each experimentally measured honey bee, approx. 26 glomeruli were detected. Considering that each glomerulus hosts multiple PNs, for each glomerular signal, we modelled 3 PNs, each with its own independent connections with a subset of the 1000 modelled KCs. Connections between PNs and KCs were modelled as pseudo-random, with each KC receiving connections 30% of the input PN population, as observed in bees **(Szyszka et al., 2005)** and flies **(Ashok Litwin-Kumar et al., 2017; Caron et al., 2013)**. PN-to-KC connections were assumed to be binary: either 0 (not connected), or 1 (connected), and the connectivity scheme is stable in a given MB model and not subject to plastic changes. This was implemented as a logic connectivity matrix (*W_PN_to_KC*) of size *number_of_PNs* ✕ *number_of_KCs*. Experimental PNs activity (*PNactivity*) was resampled at 20 Hz for modelling analysis (50 ms temporal resolution) to match the 20-Hz oscillatory cycle detected in the MB **(Cassenaer and Laurent, 2007; Laurent and Naraghi, 1994; Popov and Szyszka, 2020)**. For every time step *t,* the integrated input to each KCs (*KCepsp*) was calculated as the matrix product of the input neuron activity vector (*PNactivity*) and the logical connectivity matrix. Notably, we considered that a glomerulus is completely silent only at minimal inhibition. For this reason, at any time point, also glomeruli

with negative ∆F/F values contribute to determining a KC integrated input. This integrated input

of each KC is analogous to its excitatory post-synaptic potential (EPSP).

KCepsp(t) = PNactivity(t) × W_PN_to_KC.

Winner-takes-it-all inhibitory feedback element

Whether a KC will fire an action potential depends not only on the sum of its input signals

but also on the inhibitory activity of the MB feedback neurons **(Rybak and Menzel, 1993;**

Zwaka et al., 2018). These neurons act on the whole KC population, providing that the KCs

receiving the highest summed input will produce an action potential (AP). To account for such

effect, and based on known MB neurophysiology **(Honegger et al., 2011; Peng and Chittka,**

2017; Turner et al., 2008), we imposed that only the 10% most active *KCepsp* (*i.e.*, receiving

the strongest summed input) will fire an action potential.

If KC(i)epsp(t) > 10th percentil of KCepsp(t), then KC(i)activity(t) = 1; else KC(i)activity(t)

= 0

This outputs a binary vector of firing pattern across the whole KC population (*KCactivity*),

where ones correspond to action potential and zeroes to silent KC. This operation is repeated

for all epochs to obtain the time course of the activity of the KC population before, during and

after odour arrival. To ensure the robustness of the model and account for the variability of the

initial parameters, the MB simulation was run 10 times for each experimental AL input dataset.

The results were then averaged over the 10 MB network simulations. An exemplary data

subset and the main MatLab scripts used to model KC activity and MBON learning have been

uploaded to GitHub and are freely accessible at https://github.com/mp599/2024_eLife.git.

KC-to-MBON model and learning-induced plasticity

Evidence in *Drosophila* and in the honey bee suggests that the default state of an

appetitive MBON - *i.e.*, an MBON departing from an MB lobe innervated by a dopaminergic

neuron conveying reward information - is to be broadly responsive to neutral stimuli **(Okada**

et al., 2007; Owald et al., 2015). Under the assumption that similar principles are highly

conserved among insects, we modelled an MBON receiving connections from all KCs and with

all connections having an initial synaptic weight = 1 to reflect a generalised odour tuning. This

provides a broadly-tuned MBON that responds equally to any pattern of KC activity. Upon

learning, MBON synapses that are repeatedly recruited are switched off (= 0). As a

consequence, a trained MBON retains a high AP probability in the presence of an unfamiliar

stimulus but a lower one in response to the conditioned odorant (or to an odorant that is

represented by a similar KC population). Two parameters guide learning: a synaptic plasticity

threshold (*spt*) and a learning window. The variable *spt* defines the number of firing events

needed for a given KC to induce its output synaptic depression (*i.e.*, KC(*i*)_to_MBON synaptic

weight switch from 1 to 0). For the learning tests, a *spt* value of 15 was adopted. This value

falls in the middle of the range tested in our simulations (from 1 to 40). It is stringent enough

to hinder plastic modulation of rarely recruited KCs, while still requiring repetitive activation

(*i.e.* a synapse must be active at least 750 ms during a 3-s learning window) before allowing

learning-induced plasticity.

The learning window provides a temporal restriction of such a learning rule, analogous

to the reward/US delivery window in an appetitive conditioning protocol. Different CS/US

contingencies were tested. Inter-stimulus intervals (ISI) of -2, 1 and 4 were used, where such

values indicate the onset of a 3-s US window with respect to the onset of a 5-s CS window.

Throughout the manuscript, we referred to the three learning protocols as *backward* (ISI = -2),

early (ISI = 1), and *delay* (ISI = 4). The training was modelled using the mean glomerular

responses to the first 5 stimulations of 1-hexanol or peppermint oil; memory retention was

tested against the next 5 stimulations of both odorants (one acting as the conditioned stimulus,

the other one as the novel odorant).

Odorant response profiles simulation

The response profile database (**Fig. 1F**) was repeatedly subsampled to generate 100 odorant response maps, each comprising 30 glomerular responses. The proportion of response types (inhibitory, excitatory, non-responsive, *etc.*) was maintained constant and equal to the mean proportions across the entire population to generate a batch of similar glomerular response maps (representing perceptually similar odorants **(Guerrieri et al., 2005)**) (**Fig. 4G**). Projection neurons (or KC) response vectors used for the Pearson correlation analysis are constituted by the average response for each PN (or KC) during olfactory stimulation (t = 1 to 4 s). For **Fig. 6**, 16 odour response profiles were generated by assembling 32 glomerular response profiles from the database. Each simulated odorant contained the same number of glomerular profiles for each type: 5 inhibitory responses (group 1); 3 long inhibitory responses (group 2); 3 inhibitory-then-excitatory (group 3); 7 excitatory (group 4); 1 excitatory-then-inhibitory (group 5); 2 long excitatory (group 6); 2 short excitatory (group 7); 1 short excitatory then inhibited (group 8); 4 unclassified (group 9); 4 unresponsive profiles.

Proboscis extension response latency

Bees were collected in the afternoon at the institute's outdoor beehives, kept in custom-printed plastic cages in groups of 16 individuals, and provided with 240 microliters of 50% sugar/water solution (an average of 15 μl for each individual). On the next morning, they were harnessed in plastic tubes, blocked in place with tape, and fed 3 μl of sugar solution. Three hours later, they were exposed to an absolute olfactory conditioning protocol **(Villar et al., 2020)**. During conditioning, bees were placed in front of the odour delivery device (the same one used for the calcium imaging analysis) and exposed to clean air for 15 s (familiarisation phase), to the odorant for 5 s, and to clean air again for another 20 s. Three different intervals were selected for delivering the sugar reward: either from 2 s before odour onset to 1 s after (*backward conditioning*), from 1 to 4 s from odour onset (*early conditioning*), from 4 to 7 s from odour onset (*delay conditioning*), or from 7 to 10 s from odour onset (*trace conditioning*). Bees

were exposed to four rewarded trials, with a 10-minute inter-trial-interval. Memory was tested

1 hour after the last conditioning trial. During the memory test, videos were acquired with a

commercial video camera at 30 fps. A LED connected to the odour delivery device and in

synch with the olfactory stimulation was used as a marker for the starting timepoint for the

measurement of proboscis extension latency. Peppermint oil and 1-hexanol were used as

olfactory stimuli. They were used non-diluted and their role as conditioned stimulus (CS) or

novel odorant (NOd) was balanced between bees. During the test, half of the bees were

exposed first to the CS and then to the NOd, while the remaining half was tested in the opposite

order. A 50% sugar/water solution (w/w) was used as unconditioned stimulus (US).

Statistical Analysis

For MBON learning simulations, a total of 16 MB networks were generated and trained

with the PN response profiles to 1-hexanol ($n = 8$) or peppermint oil ($n = 8$) (**Fig. 5**), or with

computationally assembled odorant profiles (**Fig. 6**). After training, each MBON was tested

against the response profiles of the CS and of the novel odorant to assess AP firing probability

to the learned and the unfamiliar stimulus. For each simulated MBON, five memory tests

against the five repetitions of the CS and of the NOd were performed. To quantify stimulus

response to the conditioned/novel stimulus, the mean firing probability during the time-window

of stimulus arrival was calculated. The difference between the responses to the CS and to the

NOd was tested with a Kruskal-Wallis statistical test (**Fig. 5C**). The latency of MBON response

to the CS was calculated as the latency to reach the 90% of minimal firing probability upon

stimulation. The difference in latency between protocols was tested with a Wilcoxon signed

rank test (**Fig. 5D** and **6C**); the difference in latency variance was assessed with a Barlett's

test (**Fig. 5D**).

Proboscis extension response was used to assess memory retention 1 h after the last

absolute conditioning trial. Conditioned stimulus specificity was tested with a McNemar test.

The latency of CS-specific PER was measured as the first frame after stimulus onset, where

the proboscis trespasses the imaginary line between the open mandibles (**Fig. 4E**).

Differences between response latencies after early and delay conditioning were assessed with

a Kruskal-Wallis test as well as with Barlett's test for difference in variance among datasets

(**Fig. 4F**).

Acknowledgments

791   This work was supported by the MSCA long-term fellowship to M.P., the ERC Starting

792 Grant ('Emerg-Ant' 759817) to A.W., and the ERC Advanced Grant ('Cognibrains') to M.G.,

793 who also thanks the Institut Universitaire de France (IUF), the CNRS, and Sorbonne University

794 for generous support.

Bibliography

Adam, E., Hansson, B. S. and Knaden, M. (2022). Fast Learners: One Trial Olfactory
Learning in Insects. *Front. Ecol. Evol.* **10**, 876596.

Amin, H. and Lin, A. C. (2019). Neuronal mechanisms underlying innate and learned olfactory
processing in Drosophila. *Curr. Opin. insect Sci.* **36**, 9–17.

Ardin, P., Peng, F., Mangan, M., Lagogiannis, K. and Webb, B. (2016). Using an Insect
Mushroom Body Circuit to Encode Route Memory in Complex Natural Environments.
PLoS Comput. Biol. **12**, 1–22.

Ashok Litwin-Kumar, A., Decker Harris, K., Axel, R., Litwin-Kumar, A., Sompolinsky, H.,
Abbott, L. and Zuckerman, M. B. (2017). Optimal Degrees of Synaptic Connectivity.
Neuron **93**, 1153-1164.e7.

Aso, Y., Sitaraman, D., Ichinose, T., Kaun, K. R., Vogt, K., Belliart-Guérin, G., Plaçais, P.
Y., Robie, A. A., Yamagata, N., Schnaitmann, C., et al. (2014a). Mushroom body output
neurons encode valence and guide memory-based action selection in Drosophila. *Elife* **3**,
e04580.

Aso, Y., Hattori, D., Yu, Y., Johnston, R. M., Iyer, N. A., Ngo, T. T. B., Dionne, H., Abbott,
L. F., Axel, R., Tanimoto, H., et al. (2014b). The neuronal architecture of the mushroom
body provides a logic for associative learning. *Elife* **3**, e04577.

Bestea, L., Paoli, M., Arrufat, P., Ronsin, B., Carcaud, J., Sandoz, J.-C., Velarde, R.,
Giurfa, M. and de Brito Sanchez, M. G. (2022). The short neuropeptide F regulates
appetitive but not aversive responsiveness in a social insect. *iScience* **25**, 103619.

Bitterman, M. E., Menzel, R., Fietz, A. and Schäfer, S. (1983). Classical conditioning of
proboscis extension in honeybees (Apis mellifera). *J. Comp. Psychol.* **97**, 107–19.

Boisvert, M. J. and Sherry, D. F. (2006). Interval Timing by an Invertebrate, the Bumble Bee

819   Bombus impatiens. *Curr. Biol.* **16**, 1636–1640.

820 **Buehlmann, C., Wozniak, B., Goulard, R., Webb, B., Graham, P. and Niven, J. E.** (2020).
821   Mushroom Bodies Are Required for Learned Visual Navigation, but Not for Innate Visual
822   Behavior, in Ants. *Curr. Biol.* **30**, 3438-3443.e2.

823 **Burke, C. J., Huetteroth, W., Owald, D., Perisse, E., Krashes, M. J., Das, G., Gohl, D.,**
824   **Silies, M., Certel, S. and Waddell, S.** (2012). Layered reward signalling through
825   octopamine and dopamine in Drosophila. *Nature* **492**, 433–7.

826 **Carcaud, J., Giuria, M. and Sandoz, J. C.** (2016). Parallel olfactory processing in the honey
827   bee brain: Odor learning and generalization under selective lesion of a projection neuron
828   tract. *Front. Integr. Neurosci.* **9**, 171692.

829 **Caron, S. J. C., Ruta, V., Abbott, L. F. and Axel, R.** (2013). Random convergence of olfactory
830   inputs in the Drosophila mushroom body. *Nature* **497**, 113–7.

831 **Cassenaer, S. and Laurent, G.** (2007). Hebbian STDP in mushroom bodies facilitates the
832   synchronous flow of olfactory information in locusts. *Nature* **448**, 709–713.

833 **Chen, X., Ginoux, F., Carbo-Tano, M., Mora, T., Walczak, A. M. and Wyart, C.** (2023).
834   Granger causality analysis for calcium transients in neuronal networks, challenges and
835   improvements. *Elife* 1–25.

836 **Chou, Y.-H., Spletter, M. L., Yaksi, E., Leong, J. C. S., Wilson, R. I. and Luo, L.** (2010).
837   Diversity and Wiring Variability of Olfactory Local Interneurons in the Drosophila Antennal
838   Lobe. *Nat. Neurosci.* **13**, 439–449.

839 **Cognigni, P., Felsenberg, J. and Waddell, S.** (2018). Do the right thing: neural network
840   mechanisms of memory formation, expression and update in Drosophila. *Curr. Opin.*
841   *Neurobiol.* **49**, 51–58.

842 **Cohn, R., Morantte, I. and Ruta, V.** (2015). Coordinated and Compartmentalized
843   Neuromodulation Shapes Sensory Processing in Drosophila. *Cell* **163**, 1742–1755.

844 **Deisig, N., Lachnit, H., Giurfa, M. and Hellstern, F.** (2001). Configural Olfactory Learning in
845   Honeybees: Negative and Positive Patterning Discrimination. *Learn. Mem.* **8**, 70.

846 **Devaud, J.-M., Papouin, T., Carcaud, J., Sandoz, J.-C., Grünewald, B. and Giurfa, M.**
847   (2015). Neural substrate for higher-order learning in an insect: Mushroom bodies are
848   necessary for configural discriminations. *Proc. Natl. Acad. Sci.* **112**, E5854–E5862.

849 **Domjan, M.** (2015). Classical conditioning: Foundations. In *The principles of learning and*
850   *behavior* (ed. Domjan, M.), pp. 84–85. Stamford: Cengage Learning.

851 **Drew, M. R., Couvillon, P. A., Zupan, B., Cooke, A. and Balsam, P. D.** (2005). Temporal
852   control of conditioned responding in goldfish. *J. Exp. Psychol.* **31**, 31–39.

853 **Dylla, K. V., Galili, D. S., Szyszka, P. and Lüdke, A.** (2013). Trace conditioning in insects-
854   keep the trace! *Front. Physiol.* **4 AUG**, 1–12.

855 **Eschbach, C., Fushiki, A., Winding, M., Schneider-Mizell, C. M., Shao, M., Arruda, R.,**
856   **Eichler, K., Valdes-Aleman, J., Ohyama, T., Thum, A. S., et al.** (2020). Recurrent
857   architecture for adaptive regulation of learning in the insect brain. *Nat. Neurosci. 2020*
858   *234* **23**, 544–555.

859 **Felsenberg, J., Plath, J. A., Lorang, S., Morgenstern, L. and Eisenhardt, D.** (2014). Short-
860   and long-term memories formed upon backward conditioning in honeybees (Apis
861   mellifera). *Learn. Mem.* **21**, 37–45.

862 **Finelli, L. A., Haney, S., Bazhenov, M., Stopfer, M. and Sejnowski, T. J.** (2008). Synaptic
863   learning rules and sparse coding in a model sensory system. *PLoS Comput. Biol.* **4**,.

864 **Flanagan, D. and Mercer, A. R.** (1989). An atlas and 3-D reconstruction of the antennal lobes
865   in the worker honey bee, Apis mellifera L. (Hymenoptera : Apidae). *Int. J. Insect Morphol.*
866   *Embryol.* **18**, 145–159.

867 **Galili, D. S., Lüdke, A., Galizia, C. G., Szyszka, P. and Tanimoto, H.** (2011). Olfactory trace
868   conditioning in Drosophila. *J. Neurosci.* **31**, 7240–7248.

869 **Galizia, C. G.** (2014). Olfactory coding in the insect brain: data and conjectures. *Eur. J.*
870   *Neurosci.* **39**, 1784–1795.

871 **Galizia, C. G., Sachse, S., Rappert, A. and Menzel, R.** (1999). The glomerular code for odor
872   representation is species specific in the honeybee Apis mellifera. *Nat. Neurosci.* **2**, 473–
873   8.

Ganeshina, O. and Menzel, R. (2001). GABA-Immunoreactive Neurons in the Mushroom Bodies of the Honeybee: An Electron Microscopic Study Indexing terms: GABA; postembedding EM-immunocytochemistry; microcircuits. *J. Comp. Neurol* **437**, 335–349.

Gil-Guevara, O., Bernal, H. A. and Riveros, A. J. (2022). Honey bees respond to multimodal stimuli following the principle of inverse effectiveness. *J. Exp. Biol.* **225**,.

Girardin, C. C., Kreissl, S. and Galizia, C. G. (2013). Inhibitory connections in the honeybee antennal lobe are spatially patchy. *J. Neurophysiol.* **109**, 332–343.

Giurfa, M. (2007). Behavioral and neural analysis of associative learning in the honeybee: a taste from the magic well. *J. Comp. Physiol. A* **193**, 801–824.

Giurfa, M. (2015). Learning and cognition in insects. *Wiley Interdiscip. Rev. Cogn. Sci.* **6**, 383–395.

Giurfa, M. and Malun, D. (2004). Associative mechanosensory conditioning of the proboscis extension reflex in honeybees. *Learn. Mem.* **11**, 294–302.

Giurfa, M. and Sandoz, J.-C. (2012). Invertebrate learning and memory: Fifty years of olfactory conditioning of the proboscis extension response in honeybees. *Learn. Mem.* **19**, 54–66.

Gkanias, E., McCurdy, L. Y., Nitabach, M. N. and Webb, B. (2022). An incentive circuit for memory dynamics in the mushroom body of Drosophila melanogaster. *Elife* **11**,.

Grünewald, B. (1999). Morphology of feedback neurons in the mushroom body of the honeybee, Apis mellifera. *J. Comp. Neurol.* **404**, 114–26.

Guerrieri, F., Schubert, M., Sandoz, J. C. and Giurfa, M. (2005). Perceptual and neural olfactory similarity in honeybees. *PLoS Biol.* **3**, 0718–0732.

Hammer, M. (1993). An identified neuron mediates the unconditioned stimulus in associative olfactory learning in honeybees. *Nature* **366**, 59–63.

Heisenberg, M. (2003). Mushroom body memoir: From maps to models. *Nat. Rev. Neurosci.* **4**, 266–275.

Helassa, N., Zhang, X.-H., Conte, I., Scaringi, J., Esposito, E., Bradley, J., Carter, T., Ogden, D., Morad, M. and Török, K. (2015). Fast-Response Calmodulin-Based Fluorescent Indicators Reveal Rapid Intracellular Calcium Dynamics OPEN.

Hellstern, F., Malaka, R. and Hammer, M. (1998). Backward inhibitory learning in honeybees: A behavioral analysis of reinforcement processing. *Learn. Mem.* **4**, 429–444.

Hige, T., Aso, Y., Modi, M. N., Rubin, G. M. and Turner, G. C. (2015). Heterosynaptic Plasticity Underlies Aversive Olfactory Learning in Drosophila. *Neuron* **88**, 985–998.

Holland, P. C. (1980). CS-US interval as a determinant of the form of Pavlovian appetitive conditioned responses. *J. Exp. Psychol. Anim. Behav. Process.* **6**, 155–174.

Honegger, K. S., Campbell, R. A. A. and Turner, G. C. (2011). Cellular-Resolution Population Imaging Reveals Robust Sparse Coding in the Drosophila Mushroom Body. *J. Neurosci.* **31**, 11772–11785.

Huang, J., Zhang, W., Qiao, W., Hu, A. and Wang, Z. (2010). Functional Connectivity and Selective Odor Responses of Excitatory Local Interneurons in Drosophila Antennal Lobe. *Neuron* **67**, 1021–1033.

Huerta, R., Nowotny, T., García-Sanchez, M., Abarbanel, H. D. I. and Rabinovich, M. I. (2004). Learning classification in the olfactory system of insects. *Neural Comput.* **16**, 1601–1640.

Ito, I., Ong, R. C. Y., Raman, B. and Stopfer, M. (2008). Sparse odor representation and olfactory learning. *Nat. Neurosci.* **11**, 1177–1184.

Jeanne, J. M., Fişek, M. and Wilson, R. I. (2018). The organization of projections from olfactory glomeruli onto higher-order neurons. *Neuron* **98**, 1198-1213.e6.

Jefferis, G. S. X. E., Potter, C. J., Chan, A. M., Marin, E. C., Rohlfing, T., Maurer, C. R. and Luo, L. (2007). Comprehensive maps of Drosophila higher olfactory centers: spatially segregated fruit and pheromone representation. *Cell* **128**, 1187–1203.

Junek, S., Kludt, E., Wolf, F. and Schild, D. (2010). Olfactory Coding with Patterns of Response Latencies. *Neuron* **67**, 872–884.

Kim, B., Haney, S., Milan, A. P., Joshi, S., Aldworth, Z., Rulkov, N., Kim, A. T., Bazhenov, M. and Stopfer, M. A. (2023). Olfactory receptor neurons generate multiple response

motifs, increasing coding space dimensionality. *Elife* **12**,.

Kirkpatrick, K. and Balsam, P. D. (2016). Associative learning and timing. *Curr. Opin. Behav.*
Sci. **8**, 181–185.

Krofczik, S., Menzel, R. and Nawrot, M. P. (2009). Rapid odor processing in the honeybee
antennal lobe network. *Front. Comput. Neurosci.* **2**, 9.

Laurent, G. and Davidowitz, H. (1994). Encoding of olfactory information with oscillating
neural assemblies. *Science (80-.).* **265**, 1872–5.

Laurent, G. and Naraghi, M. (1994). Odorant-induced oscillations in the mushroom bodies of
the locust. *J. Neurosci.* **14**, 2993–3004.

Le Moël, F. and Wystrach, A. (2020). Towards a multi-level understanding in insect
navigation. *Curr. Opin. Insect Sci.* **42**, 110–117.

Le Moël, F., Stone, T., Lihoreau, M., Wystrach, A. and Webb, B. (2019). The central
complex as a potential substrate for vector based navigation. *Front. Psychol.* **10**, 380097.

Lin, A. C., Bygrave, A. M., De Calignon, A., Lee, T. and Miesenböck, G. (2014). Sparse,
decorrelated odor coding in the mushroom body enhances learned odor discrimination.
Nat. Neurosci. **17**, 559–568.

Liu, C., Plaçais, P.-Y., Yamagata, N., Pfeiffer, B. D., Aso, Y., Friedrich, A. B., Siwanowicz,
I., Rubin, G. M., Preat, T. and Tanimoto, H. (2012). A subset of dopamine neurons
signals reward for odour memory in Drosophila. *Nature* **488**, 512–516.

Locatelli, F. F., Fernandez, P. C. and Smith, B. H. (2016). Learning about natural variation
of odor mixtures enhances categorization in early olfactory processing. *J. Exp. Biol.* **219**,
2752–2762.

Lüdke, A., Raiser, G., Nehrkorn, J., Herz, A. V. M., Galizia, C. G. and Szyszka, P. (2018).
Calcium in Kenyon cell somata as a substrate for an olfactory sensory memory in
Drosophila. *Front. Cell. Neurosci.* **12**, 1–16.

Lyutova, R., Selcho, M., Pfeuffer, M., Segebarth, D., Habenstein, J., Rohwedder, A.,
Frantzmann, F., Wegener, C., Thum, A. S. and Pauls, D. (2019). Reward signaling in
a recurrent circuit of dopaminergic neurons and peptidergic Kenyon cells. *Nat. Commun.*
10,.

Malaka, R., Ragg, T. and Hammer, M. (1995). Kinetic models of odor transduction
implemented as artificial neural networks. Simulations of complex response properties of
honeybee olfactory neurons. *Biol. Cybern.* **73**, 195–207.

Menzel, R. (1999). Memory dynamics in the honeybee. *J. Comp. Physiol. - A* **185**, 323–340.

Mertes, M., Carcaud, J. and Sandoz, J. C. (2021). Olfactory coding in the antennal lobe of
the bumble bee Bombus terrestris. *Sci. Rep.* **11**, 1–14.

Mobbs, P. G. (1982). The brain of the honeybee Apis mellifera . I. The connections and spatial
organization of the mushroom bodies. *Philos. Trans. R. Soc. London. B, Biol. Sci.* **298**,
309–354.

Moreaux, L. and Laurent, G. (2007). Estimating firing rates from calcium signals in locust
projection neurons in vivo. *Front. Neural Circuits* **1**, 2.

Münch, D. and Galizia, C. G. (2016). DoOR 2.0 - Comprehensive mapping of Drosophila
melanogaster odorant responses. *Sci. Rep.* **6**, 1–14.

Nouvian, M., Deisig, N., Reinhard, J. and Giurfa, M. (2018). Seasonality, alarm pheromone
and serotonin: insights on the neurobiology of honeybee defence from winter bees. *Biol.*
Lett. **14**,.

Okada, R., Rybak, J., Manz, G. and Menzel, R. (2007). Learning-related plasticity in PE1 and
other mushroom body-extrinsic neurons in the honeybee brain. *J. Neurosci.* **27**, 11736–
11747.

Olsen, S. R., Bhandawat, V. and Wilson, R. I. (2007). Excitatory interactions between
olfactory processing channels in the Drosophila antennal lobe. *Neuron* **54**, 89–103.

Olshausen, B. A. and Field, D. J. (2004). Sparse coding of sensory inputs. *Curr. Opin.*
Neurobiol. **14**, 481–487.

Owald, D., Felsenberg, J., Talbot, C. B., Das, G., Perisse, E., Huetteroth, W. and Waddell,
S. (2015). Activity of defined mushroom body output neurons underlies learned olfactory
behavior in Drosophila. *Neuron* **86**, 417–427.

Paoli, M. and Galizia, G. C. (2021). Olfactory coding in honeybees. *Cell Tissue Res.* **383**, 35–
58.
Paoli, M. and Haase, A. (2018). In Vivo Two-Photon Imaging of the Olfactory System in
Insects. In *Methods in molecular biology (Clifton, N.J.)*, pp. 179–219. Methods Mol Biol.
Paoli, M., Weisz, N., Antolini, R. and Haase, A. (2016). Spatially resolved time-frequency
analysis of odour coding in the insect antennal lobe. *Eur. J. Neurosci.* **44**, 2387–2395.
Paoli, M., Andrione, M. and Haase, A. (2017). Imaging Techniques in Insects. In *Lateralized*
Brain Functions: Methods in Human and Non-Human Species (ed. Rogers, L. J.) and
Vallortigara, G.), pp. 471–519. New York, NY: Springer New York.
Paoli, M., Albi, A., Zanon, M., Zanini, D., Antolini, R. and Haase, A. (2018). Neuronal
response latencies encode first odor identity information across subjects. *J. Neurosci.* **38**,
9240 –9251.
Paoli, M., Macri, C. and Giurfa, M. (2023a). A cognitive account of trace conditioning in
insects. *Curr. Opin. Insect Sci.* **57**, 101034.
Paoli, M., Antonacci, Y., Albi, A., Faes, L. and Haase, A. (2023b). Granger Causality
Analysis of Transient Calcium Dynamics in the Honey Bee Antennal Lobe Network.
Insects 2023, Vol. 14, Page 539 **14**, 539.
Papadopoulou, M., Cassenaer, S., Nowotny, T. and Laurent, G. (2011). Normalization for
sparse encoding of odours by a wide-field interneurons. *Science (80-.).* **332**, 721–725.
Paton, J. J. and Buonomano, D. V. (2018). The Neural Basis of Timing: Distributed
Mechanisms for Diverse Functions. *Neuron* **98**, 687–705.
Patterson, M. A., Lagier, S. and Carleton, A. (2013). Odor representations in the olfactory
bulb evolve after the first breath and persist as an odor afterimage. *Proc. Natl. Acad. Sci.*
U. S. A. **110**,.
Pavlov, I. P. (1927). *Conditioned Reflexes: an Investigation of the Physiological Activity of the*
Cerebral Cortex. (ed. Dr. G. V. Anrep) Oxford University Press.
Peng, F. and Chittka, L. (2017). A Simple Computational Model of the Bee Mushroom Body
Can Explain Seemingly Complex Forms of Olfactory Learning and Memory. *Curr. Biol.*
27, 224–230.
Perez-Orive, J., Mazor, O., Turner, G. C., Cassenaer, S., Wilson, R. I. and Laurent, G.
(2002). Oscillations and sparsening of odor representations in the mushroom body.
Science (80-.). **297**, 359–365.
Perrett, S. P., Ruiz, B. P. and Mauk, M. D. (1993). Cerebellar cortex lesions disrupt learning-
dependent timing of conditioned eyelid responses. *J. Neurosci.* **13**, 1708–1718.
Popov, T. and Szyszka, P. (2020). Alpha oscillations govern interhemispheric spike timing
coordination in the honey bee brain. *Proc. R. Soc. B Biol. Sci.* **287**, 20200115.
Raiser, G., Galizia, C. G. and Szyszka, P. (2017). A High-Bandwidth Dual-Channel Olfactory
Stimulator for Studying Temporal Sensitivity of Olfactory Processing. *Chem. Senses* **42**,
141–151.
Roussel, E., Carcaud, J., Combe, M., Giurfa, M. and Sandoz, J. C. (2014). Olfactory coding
in the honeybee lateral horn. *Curr. Biol.* **24**, 561–567.
Rybak, J. and Menzel, R. (1993). Anatomy of the mushroom bodies in the honey bee brain:
The neuronal connections of the alpha-lobe. *J. Comp. Neurol.* **334**, 444–465.
Sachse, S. and Galizia, C. G. (2002). Role of inhibition for temporal and spatial odor
representation in olfactory output neurons: a calcium imaging study. *J. Neurophysiol.* **87**,
1106–17.
Sachse, S. and Galizia, C. G. (2003). The coding of odour-intensity in the honeybee antennal
lobe: Local computation optimizes odour representation. *Eur. J. Neurosci.* **18**, 2119–
2132.
Sachse, S., Rappert, A. and Galizia, C. G. (1999). The spatial representation of chemical
structures in the antennal lobe of honeybees: Steps towards the olfactory code. *Eur. J.*
Neurosci. **11**, 3970–3982.
Schäfer, S. and Bicker, G. (1986). Distribution of GABA-like immunoreactivity in the brain of
the honeybee. *Neuroscience* **246**, 287–300.
Schmalz, F., el Jundi, B., Rössler, W. and Strube-Bloss, M. (2022). Categorizing Visual

Information in Subpopulations of Honeybee Mushroom Body Output Neurons. *Front. Physiol.* **13**, 1–11.

Shang, Y., Claridge-Chang, A., Sjulson, L., Pypaert, M. and Miesenböck, G. (2007). Excitatory local circuits and their implications for olfactory processing in the fly antennal lobe. *Cell* **128**, 601–12.

Smith, D., Wessnitzer, J. and Webb, B. (2008). A model of associative learning in the mushroom body. *Biol. Cybern.* **99**, 89–103.

Springer, M. and Nawrot, M. P. (2021). A Mechanistic Model for Reward Prediction and Extinction Learning in the Fruit Fly. *eNeuro* **8**,.

Stopfer, M. (2014). Central processing in the mushroom bodies. *Curr. Opin. Insect Sci.* **6**, 99–103.

Strausfeld, N. J. (2002). Organization of the honey bee mushroom body: Representation of the calyx within the vertical and gamma lobes. *J. Comp. Neurol.* **450**, 4–33.

Strube-Bloss, M. F. and Rössler, W. (2018). Multimodal integration and stimulus categorization in putative mushroom body output neurons of the honeybee. *R. Soc. Open Sci.* **5**, 171785.

Strutz, A., Soelter, J., Baschwitz, A., Farhan, A., Grabe, V., Rybak, J., Knaden, M., Schmuker, M., Hansson, B. S. and Sachse, S. (2014). Decoding odor quality and intensity in the Drosophila brain. *Elife* **3**, e04147.

Szyszka, P., Ditzen, M., Galkin, A., Galizia, C. G. and Menzel, R. (2005). Sparsening and temporal sharpening of olfactory representations in the honeybee mushroom bodies. *J. Neurophysiol.* **94**, 3303–3313.

Szyszka, P., Demmler, C., Oemisch, M., Sommer, L., Biergans, S., Birnbach, B., Silbering, A. F. and Galizia, C. G. (2011). Mind the gap: Olfactory trace conditioning in honeybees. *J. Neurosci.* **31**, 7229–7239.

Takeda, K. (1961). Classical conditioned response in the honey bee. *J. Insect Physiol.* **6**, 168–179.

Turner, G. C., Bazhenov, M. and Laurent, G. (2008). Olfactory representations by Drosophila mushroom body neurons. *J. Neurophysiol.* **99**, 734–46.

Villar, M. E., Marchal, P., Viola, H. and Giurfa, M. (2020). Redefining single-trial memories in the honeybee. *Cell Rep.* **30**, 2603–2613.

Vogt, K., Yarali, A. and Tanimoto, H. (2015). Reversing Stimulus Timing in Visual Conditioning Leads to Memories with Opposite Valence in Drosophila. *PLoS One* **10**, e0139797.

Vrontou, E., Groschner, L. N., Szydlowski, S., Brain, R., Krebbers, A. and Miesenböck, G. (2021). Response competition between neurons and antineurons in the mushroom body. *Curr. Biol.* **31**, 4911-4922.e4.

Wilson, R. I., Turner, G. C. and Laurent, G. (2004). Transformation of Olfactory Drosophila Antennal Lobe. *Science (80-.).* **303**, 366–370.

Wright, G. A. and Smith, B. H. (2004). Different thresholds for detection and discrimination of odors in the honey bee (Apis mellifera). *Chem. Senses* **29**, 127–135.

Wystrach, A. (2023). Neurons from pre-motor areas to the Mushroom bodies can orchestrate latent visual learning in navigating insects. *bioRxiv* 2023.03.09.531867.

Wystrach, A., Buehlmann, C., Schwarz, S., Cheng, K. and Graham, P. (2020). Rapid Aversive and Memory Trace Learning during Route Navigation in Desert Ants. *Curr. Biol.* **30**, 1927-1933.e2.

Yaksi, E. and Wilson, R. I. (2010). Electrical coupling between olfactory glomeruli. *Neuron* **67**, 1034–47.

Zeng, J., Li, X., Zhang, R., Chu, J., Li, Y. and Li, Y. (2023). Local 5-HT signaling bi-directionally regulates the coincidence time window for associative learning Local 5-HT signaling bi-directionally regulates the coincidence time window for associative learning. 1–18.

Zwaka, H., Bartels, R., Grünewald, B. and Menzel, R. (2018). Neural organization of A3 mushroom body extrinsic neurons in the honeybee brain. *Front. Neuroanat.* **12**, 1–11.

Figures

1095

Figure 1. Projection Neurons Calcium Imaging Analysis

(A) For each AL, the odour response maps for the three stimuli were merged into an RGB image to highlight glomerular structures: 1-hexanol was set as the red channel, 1-heptanol as the green, and peppermint as the blue. Glomeruli were hand-labelled for time-series extraction. V, ventral; L, lateral.

(B) Exemplary AL odorant responses during olfactory stimulation with three odorants. Colorbar indicates the relative change of activity during olfactory stimulation with respect to the pre-stimulus baseline. Circular areas indicate identified glomeruli according to **(A)**

(C,D) Temporal profiles of the glomerular regions identified in **(A)** for the three odorants. Each line in **(C)**, labelled from 1 to 28, refers to the glomerular ID in **(A)**. Temporal profiles represent the average activity of 20 stimulations. Dashed lines in **(C,D)** limit the begin and the end of the olfactory stimulation.

(E-E') Pearson's correlations between pairs of glomerular response vectors across repetitions. **(left)** The upper right part of the matrix shows correlation scores among glomerular activity during olfactory stimulation (t = 1 to 5 s after odorant onset, ON vs ON) across trials; the bottom left part shows correlation scores among glomerular activity before stimulation (t = -1 to 0 s, PRE vs PRE) across trials. **(right)** The upper right part of the matrix shows correlation scores among glomerular activity after olfactory stimulation (t = 1 to 4 s after odorant offset, POST vs POST) across trials; the bottom left part shows correlation scores among glomerular activity during and after stimulation

1115   (ON *vs* POST) across trials. **(E')** Mean (± s.e.m.) across-trial correlation of the four combinations
1116   presented in the matrices in **E**.
1117   **(F)** All glomerular responses from 8 ALs to 3 odorants were pooled together to provide an overview
1118   of the complexity of response profiles *(n = 546).*

1119

Figure 2. Clustering of Projection Neurons' Response Profiles

(A) All glomerular responses were clustered with supervision according to their activity
(excitatory, inhibitory, non-responsive) during and after stimulus arrival.

(B) Mean ± s.e.m. of all curves of the relative groups in (**A**). The green patch indicates the
stimulus delivery interval.

(C) Average curves of all response groups from (**B**) are superimposed. The grey patch indicates
the stimulus delivery interval.

(D) Relative amount of all response categories.

(E) Latency of glomerular responses for excitatory and inhibitory profiles (groups 1 to 8). For
groups 7 and 8, the latency of short excitatory response's termination was calculated. Letters
refer to significative groups after Kruskal-Wallis statistical test and Tukey-Kramer correction. On
the left side, exemplary traces of phasic and tonic excitatory responses and an inhibitory
response are shown. The orange line indicates the latency of response onset or termination.

1136
1137 Figure 3. A Simple Neural Network Model for Olfactory Coding and Learning.
(A) The synaptic connectivity between AL PNs and KCs is built as a pseudorandom logic matrix
where each column represents KC dendritic arborisation and each row PN axon terminals. Each
KC receives synaptic contact – here represented as a white circle – from 30% of the PN
population.
(B) For each timepoint t of a calcium imaging recording, the measured activity level for each
glomerulus (represented by a PN in the model) is projected onto all its synaptic connections with
the KC population. Each KC, *i.e.* each column in the scheme, integrates all excitatory post-
synaptic potentials. Finally, recurrent inhibitory feedback is simulated by imposing that only the
10% most active KCs will generate an action potential (AP) (red cells), while all others remain
silent (grey cells).
(C) A single appetitive MBON receiving input from all KCs was modelled. Based on a synaptic
plasticity threshold parameter, all synapses from KCs to MBON that are activated with a
frequency greater than allowed by the defined threshold during the learning window will be
switched off. This phenomenon results in a decrease AP probability in the modelled MBON upon

stimulation with the learned stimulus. Abbreviations: action potential, AP; antennal lobe, AL;
excitatory post-synaptic potential, ePSP; Kenyon cells, KC; mushroom body, MB; mushroom
body output neuron, MBON; projection neurons, PN.

Figure 4. Olfactory Representation in Modelled Kenyon Cells
(A) The first row displays the recorded PN responses of one honey bee to three odorants. The second row shows the transformation of the PN activity operated by the model to simulate the activity of 1000 KCs. In each plot, KCs are ordered for response strength during the onset and offset to better visualise activity clusters.
(B) Turnover rate of recruited KCs across adjacent time points. Light blue traces refer to individual bees (mean of 10 MB simulations). Thick, dark blue traces indicate average curves across bees.
(C) Cumulative recruitment of KC shows two main recruitment events at stimulus onset and offset. To observe odorant-related KC recruitment dynamics, the cumulative sum was initiated at stimulus onset. Light blue traces refer to individual bees (mean of 10 MB simulations). Thick, dark blue traces indicate average curves across bees.
(D,E) Correlation matrix among time points of measured PN activity (**C**) and simulated KC activity (**E**). Mean responses to the three odorants were concatenated to allow observing within and across odorant correlations.
(F) PCA of odorants' trajectories in the measured PN space (top) and in the modelled KC space (bottom) for four exemplary bees. Trajectories comprise a 10-s interval, ranging from odour onset ($t = 0$ s, light) to 5 s after offset ($t = 10$ s, dark).
(G) The raster plot shows a set of 100 artificially combined glomerular responses (top), each simulating the neural representation of 100 similar odorants in the PN space. A second raster plot (middle) shows the representation of the same odorants in the modelled KC space. Note that

odorant representation was considered as the mean activity of PN (or KC) during stimulus arrival (0.5 to 5 s). The histogram shows the distributions of between-odorant correlations computed among all pairs of odour vectors in the PN and KC space. A significant difference between the two distributions was tested with a Student's t test ($p \approx 0$, $n = 4774$).

1181

Figure 5. Modeled Mushroom Body Output Neurons Can Predict Behavior
(A) Four experimental protocols differing for CS/US inter stimulus interval (ISI) were used for
training the model (**B**-**D**) and for behavioural measurements of the proboscis extension reflex
(PER) conditioning (**E**,**F**). Trained simulated MBONs, as well as conditioned honey bees were
tested against the conditioned stimulus (CS, blue) and a novel odorant (NOd, green).
(B) Time course of action potential (AP) probability of a trained MBON upon stimulation with the
CS (blue) or with the novel odorant (green). Olfactory learning was modeled according to
backward, *early*, *delay*, and *trace* conditioning protocols. Glomerular responses for 8 bees to 1-
hexanol and peppermint oil were used alternatively as CS and NOd. Data for responses to CS
and NOd were pooled together (*n* = 16 traces for each protocol). Thick trace: average AP
probability profile; dotted vertical lines: stimulus onset/offset; grey bar: modelled learning window.
(C) Distribution of the mean values of traces in (B) during CS and NOd stimulation. Because
appetitive learning produces a decrease in MBON firing rate, the complementary value of the
probability fraction provides a proxy for a learned appetitive response (Kruskal-Wallis statistical
test: $p_{backward} = 0.0545$; $p_{early} = 1.42*10^{-6}$; $p_{delay} = 1.04*10^{-5}$; $p_{trace} = 0.0595$; $n = 16$).
(D) Latency of 90% of minimal MBON activity in response to the CS for early and delay
conditioning protocols (Wilcoxon test early *vs* delay, $p = 0.0015$; delay *vs* trace, $p = 0.0005$.
Letters indicate statistical groups according to the Wilcoxon test. Barlett's test for variance
difference early *vs* delay, $p = 1.73*10^{-6}$; delay *vs* trace, $p = 0.1808$; early *vs* trace $p = 1.29*10^{-8}$).
(E) Memory retention test of honey bees 1 h after absolute conditioning. Bars indicate the
percentage of individuals showing proboscis extension reflex (PER) when presented with the
conditioned (blue) or the novel odorant (green). Error bars indicate 95% confidence intervals.

(Responses to the CS/NOd were compared with a McNemar test for binomial distribution; p_{early} =
0.003, p_{delay} = 0.004)
(F) Latency of the PER to the conditioned stimulus at the 1 h memory retention test (Kruskal-
Wallis statistical test: early *vs* delay, p = 0.036; early *vs* trace. p = 0.041; delay *vs* trace, p =
0.992. Letters indicate statistical groups according to the Kruskal-Wallis test. Barlett's test for
variance difference across protocols: early *vs* delay, p = 5.19*10^{-6}; early *vs* trace. p = 0.020; delay
vs trace, p = 0.007; n_{early} = 26; n_{delay} = 20; n_{trace} = 31).
1211

1212

Figure 6. Relative contribution of glomerular response types to MBON response latency.
(**A**) Two instances of odorant responses generated by artificially combining 32 glomerular
response profiles from all response groups from **Fig. 2** (first row). Response profiles were
manipulated eliminating stable excitatory responses (second row), inhibitory responses (third
row), short excitatory responses (fourth row), and short excitatory and inhibitory responses
together (last row). Dotted lines indicate stimulus onset and offset.
(**B**) Time course of action potential (AP) probability of a trained MBON upon stimulation with the
conditioned stimulus (CS, blue) or with the novel odorant (NOd, green). Olfactory learning was
modelled according to early and delay conditioning protocols and with the balanced (first row)
and modified simulated odorants (second to last row). Conditioned stimulus and novel odorant
are in blue and green, respectively *(n = 16)*. Thick trace: average AP probability profile; dotted
vertical lines: stimulus onset/offset; grey bar: modelled learning window.
(**C**) Latency of 90% of minimal MBON activity in response to the CS for *early* and *delay*
conditioning protocols (Wilcoxon signed rank test early *vs* delay; *n* = 16)
(**D**) Distribution of Wilcoxon signed rank test *p*-values of 100 simulations of (**C**). The red dashed
line indicates a significance level of 0.05; letters indicate significance groups.

1229

Supplementary Figures

Supplementary Figure S1. Sample of glomerular response profile dynamics across trials.
Each subplot displays the response profile across 20 repetitions of the same odorant for one glomerulus. For exemplificatory purpose, a subset of 10 glomerular response profiles from response groups 1 to 8 (80 out of 464) is shown. Vertical lines indicate stimulus onset and offset.

Supplementary Figure 2. Odorant response profiles for each bee and stimulus.
(**A**) Responses from 8 bees to three odorants. Each raster plot shows the average response profile
of 20 stimulus repetitions. Stimulation occurs between 0 and 5 s.
(**B**) Glomerular response profiles of all detected glomeruli (as in **A**) are shown as time traces. Green
bars indicate stimulus intervals. Each trace is the mean of 20 stimulus repetitions.
(**C**) Occurrence of the varied response profiles for each bee/odorant condition.

1243

1244

1245

Supplementary Figure S3. Odorant-Specific Activity Lasts Longer in the Kenyon Cells
Space.
Time traces of the normalised Euclidean distance from the centre of coordinates (where the centre
of coordinates was set to 0 and the maximum distance from it was set to 1) in the PN and KC space
for the three odorants. This transformation allowed performing a direct comparison between the
temporal dynamics of the representations of one odorant in the PN and KC space. Discriminability
– that is, distance from 0 – peaks during olfactory stimulation but is maintained longer in the KC
space. Notably, such a long-lasting odorant-specific signature is not achieved by maintaining onset-
recruited neurons activity for a longer period of time but by recruiting two pools of KCs (both
odorant-specific): one for the ON phase and one during the OFF phase (see also **Fig. 4**). Grey
patches indicate stimulation windows. Traces represent mean ± s.e.m. of 8 animals (PN data, red)
and relative MB simulations (KC data, blue).
1258

Supplementary Figure S4. Dependence of Simulated MBON Activity on Synaptic Plasticity Threshold.
Synaptic plasticity threshold (*spt*) dictates how many times a KC-to-MBON synapse has to be recruited during the learning window before being switched
off. The influence of *spt* on MBON learning was tested for all CS/US protocols. The extreme values of the parameters (*spt* = 1 and 40) indicate that a
certain synapse should be recruited only once or 40 times before being turned off (note that to be activated 40 times in a 3-s learning window and with
20 Hz sampling rate, a synapse should fire for two entire seconds). With the exceptions of the extreme parameters, the MBON AP probability upon CS
and NOd presentation is only weakly affected by *spt* value. Notably, a small *spt* results in higher generalised learning, while a large *spt* requires a stable
KC signal for synapses to be modulated and produces weaker learning scores.