Contiguous and complete assemblies of *Blastocystis* gut microbiome-associated

protists reveal evolutionary diversification to host ecology

Abigail L. Lind[1,2], Nathan A. McDonald[2,3], Elias R. Gerrick[4], Ami S. Bhatt[5,6], Katherine S.

Pollard[1,7,8]

1. Gladstone Institute for Data Science and Biotechnology, Gladstone Institutes, San

Francisco, CA.

2. School of Biological Sciences, Georgia Institute of Technology, Atlanta, Georgia.

3. Department of Biology, Stanford University, Stanford, CA.

4. Department of Microbiology, University of Chicago, Chicago, IL.

5. Department of Genetics, Stanford University, Stanford, California, USA

6. Department of Medicine (Hematology, Blood and Marrow Transplantation), Stanford

University, Stanford, California, USA

7. Department of Epidemiology & Biostatistics, University of California, San Francisco,

CA.

8. Chan Zuckerberg Biohub SF, San Francisco, CA.

Running title: Genomics of host adaptation in *Blastocystis*

Corresponding author contact information

katherine.pollard@gladstone.ucsf.edu

1700 Owens Street, San Francisco, CA 94158

Competing interests statement

The authors declare no competing interests.

Abstract

Blastocystis, an obligate host-associated protist, is the most common microbial eukaryote in the human gut and is widely distributed across vertebrate hosts. The evolutionary transition of *Blastocystis* from its free-living stramenopile ancestors to a radiation of host-associated organisms is poorly understood. To explore this, we cultured and sequenced eight strains representing the significant phylogenetic diversity of the genus using long-read, short-read, and Hi-C DNA sequencing, alongside gene annotation and RNA sequencing. Comparative genomic analyses revealed significant variation in gene content and genome structure across *Blastocystis.* Notably, three strains from herbivorous tortoises, phylogenetically distant from human subtypes, have markedly larger genomes with longer introns and intergenic regions, and retain canonical stop codons absent in the human-associated strains. Despite these genetic differences, all eight isolates exhibit gene losses linked to the reduced cellular complexity of *Blastocystis,* including losses of cilia and flagella genes, microtubule motor genes, and signal transduction genes. Isolates from herbivorous tortoises contained higher numbers of plant carbohydrate-metabolizing enzymes, suggesting that like gut bacteria, these protists ferment plant material in the host gut. We find evidence that some of these carbohydrate-metabolizing enzymes were horizontally acquired from bacteria, indicating that horizontal gene transfer is an ongoing process in *Blastocystis* that has contributed to host-related adaptation. Together, these results highlight substantial genetic and metabolic diversity within the *Blastocystis* genus, indicating different lineages of *Blastocystis* have varied ecological roles in the host gut.

Introduction

The vertebrate gut is host to a diverse ecosystem of bacteria, archaea, viruses, and microbial eukaryotes. This diverse microbial community has broad impacts on the physiology and development of the host. Gut microbes can metabolize compounds that are not digestible by the host, providing access to previously inaccessible nutrients (Yadav et al. 2018). Gut microbes also influence the host's immune system and impact host health through their metabolic byproducts (Sommer and Bäckhed 2013). The microbiome's composition, which is influenced by diet, physiology, and the host immune system, varies substantially across vertebrate animals (David et al. 2014; Thorman et al. 2023; Godoy-Vitorino et al. 2012).

Compared to bacteria, substantially less is understood about microbial eukaryote impacts on host physiology and the gut ecosystem, despite the prevalence and diversity of gut microbial eukaryotes across vertebrates (Campo et al. 2020; Nash et al. 2017). The protist *Blastocystis* is the most prevalent eukaryotic microorganism in the human gut, found in 10-50% of industrialized populations and at higher prevalence in non-industrialized communities (Khaled et al. 2020; Scanlan and Marchesi 2008; Scanlan et al. 2014; Asnicar et al. 2021; Jinatham et al. 2021; Scanlan and Stensvold 2013). Beyond humans, *Blastocystis* colonizes a wide range of animals including other mammals, birds, reptiles, fish, amphibians, and insects (Gantois et al. 2020; Boreham and Stenzel 1993; Yoshikawa et al. 2016). *Blastocystis* is a genetically diverse genus, with strains adapting to varied gut environments across different host species (Stensvold et al. 2007). Its presence is associated with positive health outcomes in humans, including lower levels of gut inflammation and metabolic health (Piperni et al.

2024; Asnicar et al. 2021; Nieves-Ramírez et al. 2018). However, very little is

understood about how *Blastocystis* contributes to the gut environment and interacts

with other gut microbiota.

Blastocystis belongs to the Opalinata lineage of stramenopiles, a group of

protists that are obligate animal gut commensals (Kostka 2017). Unlike most

stramenopiles, *Blastocystis* is non-motile, lacks flagella or surface hairs, and grows

predominantly as a round cell with large central vacuole (Zierdt 1988). The *Blastocystis*

genus contains numerous genetically distinct groups categorized into subtypes based

on polymorphism in the 18S rRNA gene. Thus far, 40 distinct subtypes have been

proposed, and this number is continuing to expand (Santin et al. 2024; Stensvold and

Clark 2020). To date, strains from three axenic lab-cultured subtypes (ST1, ST4, and

ST7), all of which are subtypes observed in humans, have been sequenced,

annotated, and deposited in public databases (Denoeud et al. 2011; Gentekaki et al.

2017). While these subtypes span a narrow range of the *Blastocystis* phylogenetic

diversity, they have very different genomes and gene contents (Gentekaki et al. 2017).

These genomes have either gapped scaffolds (ST7) or highly fragmented non-

contiguous genome assemblies (ST1, ST4), limiting the ability to determine synteny

between subtypes, and two genomes were annotated without consideration for the

unique *Blastocystis* phenomenon where a portion of genes lack DNA-encoded

termination codons. Thus, we lack highly complete and well annotated genomes

necessary for establishing cases of gene loss, horizontal transfer, and sequence

evolution so that functional repertoires can be associated with strain diversification.

Like most host-associated eukaryotes, *Blastocystis* grows best in the presence

of other microbiota. Establishing robust axenic cultures of intestinal protists including

Blastocystis is laborious and often either fails to remove all microbial contaminants or

results in the death of the protist (Zierdt and Williams 1974; Tan 2008). However,

removing sources of microbial contamination is important for eukaryotic genome

assembly via short-read sequencing (Gough et al. 2020). As a result, genome

sequencing for *Blastocystis* and other intestinal protists has been limited to axenic

cultures (Keeling et al. 2014; Gough et al. 2020). To overcome this barrier, we

establish high quality and contamination-free genomes from xenic cultures of

Blastocystis grown together with other microbes by using long-read Nanopore and Hi-

C sequencing technologies coupled with short reads and metagenomic assembly

techniques to establish genomes of *Blastocystis* strains that are grown together with

other microbes. The longer read length of Nanopore sequencing helps span repetitive

regions, resulting in greater contig length and fewer misassembles, and has been

successfully applied to assembling bacterial metagenomes (Moss et al. 2020). Hi-C

sequencing, which captures spatial interactions between DNA fragments, can further

improve genome assembly by correcting misassemblies and scaffolding contigs

(Šimková et al. 2024). By combining these technologies with metagenomic assembly

techniques, we were able to generate nearly complete *Blastocystis* genomes without

the need for axenic culturing. This approach allows us to investigate the full genetic

diversity of *Blastocystis*, even in the presence of other microbes, and to assemble

highly contiguous genomes that were previously difficult to obtain.

Here, we applied this hybrid sequencing strategy to eight *Blastocystis* strains,

five isolated from humans and three isolated from the herbivorous tortoise *Testudo*

horsfeldii. These isolates, five of which are xenic and three of which are axenic, span

the phylogenetic diversity of the *Blastocystis* genus, enabling us to investigate

Blastocystis evolution and determine how *Blastocystis* has adapted to thrive in animal

hosts with different diets, physiologies, and resident microbiota. Our highly contiguous

and complete genomes allowed us to investigate patterns of genome evolution within

and between subtypes, and to compare the genus to related protists. This work

demonstrates how high-quality genome assemblies can be created from xenic protist

culture by combining deep long-read sequencing with metagenomic assembly

techniques, enabling rigorous comparative genomic analyses.

Results

Capturing Blastocystis diversity

To investigate the evolution of the *Blastocystis* genus, from its initial divergence

through its later diversification, we sequenced eight diverse *Blastocystis* strains.

Strains were cultured anaerobically on egg media overlayed with a balanced salt

solution and 25% heat-inactivated horse serum. Five of these strains correspond to 3

different previously defined "subtypes" (genetically similar groups historically identified

in mammalian and avian species (Stensvold and Clark 2020). These five strains were

originally cultured from human fecal samples and were obtained from the American

Type Culture Collection. These human isolated strains correspond to three of the four

most prevalent *Blastocystis* "subtypes" (Beghini et al. 2017), genetically similar groups

historically identified in mammalian and avian species (Stensvold and Clark 2020): Two

subtype (ST) 1 (NandII, JDR), two ST3 (DL, NMH), and one ST4 (BT1). The remaining

three strains were isolated from companion animal tortoises: two from different

individuals of the domestic Russian tortoise species *Testudo horsfeldii* (Rus-B, Rus-S),

and one from a Hermann's tortoise *Testudo hermanni* (Hermann's). We performed

comparisons of the 18S rRNA gene of these our strains to other *Blastocystis* strains,

finding that the tortoise isolates were distantly related to the human isolates and do not

belong to a described mammalian or avian subtype (Figure 1A, Table S1). Analysis of

all eight cultures by light microscopy revealed predominantly vacuolar *Blastocystis*

cells, containing a large central vacuole and a thin peripheral ring of cytoplasm. While

Blastocystis have been observed to adopt a variety of morphologies, vacuolar

morphologies dominate cultured samples (Figure 1B).

To explore how the *Blastocystis* genus has emerged and diversified, we

compared the *Blastocystis* genomes generated in this work to those of two other

sequenced stramenopiles: *Cafeteria burkhardae*, a free-living heterotrophic marine

flagellate (Hackl et al. 2020), and the recently sequenced *Proteromonas lacertae*, a

reptile gut commensal closely related to *Blastocystis* (Záhonová et al. 2023). These

organisms are the closest relatives to *Blastocystis* that have sequenced genomes.

Genome comparisons allowed us to reconstruct three evolutionary transitions that

have occurred during stramenopile evolution: the transition from a free-living aquatic

lifestyle to an anaerobic lifestyle inside the animal gut (comparing *Cafeteria* to

Blastocystis/Proteromonas), the change in cell morphology from heterokont flagellate

to aflagellate, spherical cells (comparing *Cafeteria/Proteromonas* to *Blastocystis*), and

Figure 1.
(A) 18S rRNA maximum likelihood (ML) phylogeny of Blastocystis subtypes, with Proteromonas lacertae as an outgroup. Support values are 1000 replicates of ultrafast bootstrap and SH-aLRT tests (Guindon et al. 2010; Hoang et al. 2018). Strains in red were sequenced in this study. (B) 13-gene concatenation ML phylogeny for all strains and species with sequenced genomes used in this study. Important evolutionary transitions are highlighted. Images represent cellular morphology of Cafeteria burkhardae, Proteromonas lacertae, and Blastocystis.
(C) Assembly statistics of Blastocystis strains sequenced in this work compared with previously sequenced and annotated genomes. Stop-codon aware and not stop codon aware designations refer to gene annotation methodologies that take into consideration that Blastocystis genes can lack stop codons or do not consider this, respectively. The ST7 N50 is a gapped scaffold N50 generated with Sanger sequencing scaffolding.

160 adaptations to different animal hosts (comparing *Blastocystis* tortoise to *Blastocystis*

161 human) (Figure 1B).

163 Table 1. Genome assembly statistics for all genomes used in this study.

Species	Strain	Subtype	Source	Citation	Scaffolds (#)	Size (Mb)	N50 (Kb)	GC content (% GC)	BUSCO content[2]	Protein-coding genes
Blastocystis sp.	JDR	1	Human	**This study**	114	17.5	550	54.5%	61.2%	7891
Blastocystis sp.	NandII	1	Human	**This study**	54	16.0	939	54.7%	61.2%	7157
Blastocystis sp.	NMH	3	Human	**This study**	36	14.6	582	51.3%	58.0%	6571
Blastocystis sp.	DL	3	Human	**This study**	56	14.0	506	51.6%	55.7%	6154
Blastocystis sp.	BT1	4	Human	**This study**	34	15.3	882	39.4%	57.6%	7421
Blastocystis sp.	Rus-B	N/A	Russian tortoise	**This study**	33	33.3	1323	23.1%	43.2%	8547
Blastocystis sp.	Rus-S	N/A	Russian tortoise	**This study**	258	26.1	505	21.8%	42.0%	8501
Blastocystis sp.	Hermanns	N/A	Hermann's tortoise	**This study**	229	27.1	289	21.5%	42.4%	8349
Blastocystis sp.	WR1	4	Laboratory Wistar rat	No publication	1,301	12.9	30	39.5%	56.5%	5707
Blastocystis sp.	NandII	1	Human	Gentekaki et al. 2017	580	16.5	79	55.0%	61.1%	6544
Blastocystis sp.	Singapore	7	Human	Denoeud et al. 2011	54	18.8	901	45.0%	52.2%	6020
Proteromonas lacertae	LA	N/A	Sand lizard	Záhonová et al. 2023[2]	1,447	52.2	93	27.0%	62.7%	23189
Cafeteria burkhardae	BVI	N/A	Environmental	Hackl et al. 2020	169	36.3	465	70.0%	54.9%	8591

1. BUSCO version 5.6.1 with eukaryote_odb10 lineage, metaeuk genome mode

2. Gapped scaffold N50.

3. Citation for genome only. Annotations not publicly available and generated for comparative genomics in this work.

Contiguous and complete Blastocystis genomes

We prepared high quality, high molecular weight DNA, as well as RNA, from the

eight *Blastocystis* cultures. DNA was sequenced with Nanopore long-read technology.

Short-read DNA and RNA sequencing was also performed for assembly polishing and

gene annotation. Hi-C data was additionally generated to help resolve scaffolds for one

strain (ST4-BT1). Non-eukaryotic sequence was detected via multiple methods and

excluded from final assemblies (see Methods, Genome assembly and

decontamination). We additionally validated that our genomes were not contaminated

by bacterial sequence by determining that genes with high sequence similarity to

bacterial genes arose from polyadenylated eukaryotic transcripts (Table S2) (see

Methods, Gene contamination and horizontal gene transfer screens).

This combination of sequencing technologies enabled us to generate highly

contiguous genomes for all six subtypes. The N50 of these assemblies ranged from

289 Kb to 1.3 Mb, achieving a much higher assembly quality than currently available

Blastocystis genome datasets in NCBI (median: 37 Kb) (Table 1, Figure 1C). We used

the BUSCO pipeline to estimate genome completeness based on presence and

absence of single-copy universal eukaryotic genes (Manni et al. 2021). BUSCO

completeness estimates ranged from 42% to 61.2%, similar to BUSCO completeness

results of previously sequenced *Blastocystis* isolates and related species (Table 1,

Figure 1C). These values likely underestimate the true completeness, because BUSCO

is known to have poorer performance in protists relative to plants, animals, and fungi

(Saary et al. 2020). Nonetheless, the consistent recovery of a similar fraction of BUSCO

genes across stramenopile genomes indicates comparable completeness. It is

therefore striking that the sequenced *Blastocystis* strains showed substantial variability

in genome size, from 14.0 to 32.2 Mb. Strains in the same subtype exhibited similar

lengths: ST3 isolates were 14.0 and 15.3 Mb, and ST1 isolates were 17.2 and 16.0 Mb

in size. In contrast, the genomes of the three tortoise isolates were ~1.5 to 2 times

larger than human isolates, ranging from 26.0 Mb to 33.3 Mb.

Previous genome sequencing studies have raised questions about the ploidy of

Blastocystis and found evidence of segmentally duplicated blocks of genes. With the

highly contiguous genomes and multiple isolates of individual subtypes here, we

investigated questions of ploidy and conservation of synteny both within *Blastocystis*

subtypes and across the genus. *K*-mer frequency-based analysis using GenomeScope

(Ranallo-Benavidez et al. 2020) demonstrated a strong diploid signal in all *Blastocystis*

isolates, with limited evidence for aneuploidy or polyploidy. We found that strains

within ST3 and within ST1 were highly syntenic (Figure S1). However, synteny between

subtypes was much lower, broadly following the relatedness between strains based on

18S sequences.

We used a hybrid method of *de novo* gene prediction and RNA transcript

evidence to annotate *Blastocystis* genes (see Methods, Genome assembly and

decontamination). Because a substantial fraction of genes in *Blastocystis* ST1 and ST7

do not have DNA-encoded stop codons (Gentekaki et al. 2017; Klimeš et al. 2014)

(discussed below), open reading frame based gene predictions from DNA sequence

are inaccurate. Leveraging our RNA sequencing data, we were able to generate a

comprehensive gene annotation for each *Blastocystis* genome that was able to

accurately call genes lacking stop codons (see Methods, Gene annotation). Similar

approaches were taken for one previously sequenced ST1 genome (Gentekaki et al.

2017), but published *Blastocystis* genomes for ST4 and ST7 were not annotated with

these approaches, leading to incorrect gene annotations. For the *Blastocystis* strains

sequenced in this study, the number of genes per *Blastocystis* genome sequenced in

this work ranged from 6,154 to 8,547 (Table 1). Protein amino acid identity (AAI) across

753 single-copy orthologs was almost identical between strains of the same subtype

(99%-100% AAI for both ST1 and ST3), and decreased to 65-75% between human

Blastocystis isolates from different subtypes and to 40-42% between human and

tortoise isolates (Figure S2). Tortoise isolates had lower pairwise AAI than human

isolates, with the most distantly related strains in the tortoise-associated clade having

a protein identity of 56% AAI for the same 753 orthologs, suggesting substantial

genetic distance within this group.

Genomic diversity within Blastocystis

When we compared sequenced *Blastocystis* genomes, we found substantial

variation in the genomes between human *Blastocystis* isolates and the tortoise-

associated clade, including genome size. While the tortoise *Blastocystis* isolate

genomes are ~1.5x-2.4x larger than the human isolate genomes, this difference is not

reflected in protein-coding gene content, suggesting that non-genic regions are driving

the difference in genome size. We sought to determine why the genomes of

Blastocystis isolates in the tortoise-associated clade were larger than the genomes of

the *Blastocystis* human isolates.

We found little difference in transposable element presence across *Blastocystis*,

which ranges from 4.2%-6.7% of the genome in the human isolates and 5.6%-7.5% of

the genome in the tortoise isolates. However, we did observe an increase in low-

complexity regions in the tortoise isolates, comprising 0.02%-1.4% of the human

isolate genomes and 11.8%-12.7% of the tortoise isolate genomes (Table S3). This

increase in low complexity regions does not explain the difference in genome size; if

these regions are removed, the tortoise isolates remain 1.3x-2.1x larger than the

human isolates. These low complexity regions may be related to the overall lower GC

content of the tortoise isolates, which ranges from 21-23% in tortoise isolates to 39-

55% in human isolates (Table 1).

As repetitive elements did not explain differences in genome size, we

investigated differences in other genomic features. We found that the almost all introns

in human ST1, ST3, and ST4 *Blastocystis* are exactly 30 bp long, in agreement with

previous estimates of intron size in *Blastocystis* ST1 and ST7 (Gentekaki et al. 2017;

Denoeud et al. 2011). However, introns in the tortoise isolates are longer and more

variable in length (Figure 2A). We additionally found that the intergenic spaces between

genes are much longer in the tortoise *Blastocystis* isolate than in human isolates

(Figure 2B). The median intergenic length in the tortoise isolates ranges from 645-719

bp, while the median intergenic length in the human isolates ranges from 137-186 bp.

Intergenic lengths in the human isolates can frequently be extremely small, only

spanning tens of base pairs (Figure 2B). Together, the total amount of DNA spanned by

introns and intergenic regions is 13.9-16.5 Mb in the tortoise *Blastocystis* isolates,

whereas these regions span only 3.7-5.7 Mb in human *Blastocystis* isolates.

Considered together, the larger introns and intergenic distance in the tortoise isolates

explain the difference in their genome sizes relative to the human isolate.

Loss of stop codons is not universal in Blastocystis

Figure 2. (A) Intron lengths and (B) Intergenic lengths in tortoise isolates (left) and human isolates right). (C) Sequence motif enriched adjacent to transcript ends in ST1, ST3, and ST4. Motif generated with STREME (Bailey 2021). This motif is referred to as "downstream gene element" (DGE). (D) Enrichment (log(observed/expected)) of transcripts with downstream conserved element within 30 bp upstream or downstream of transcript end, normalized by expected DGE occurrence given GC content.

In addition to genome size and gene density, we found surprising differences in

how genes are encoded in the genome across *Blastocystis*. In *Blastocystis* ST1 and

ST7, a fraction of protein coding genes (15% for ST7 and 26% for ST1) lack canonical

DNA-encoded termination codons (Gentekaki et al. 2017; Klimeš et al. 2014). A

proposed mechanism for translation of these genes is that the translational machinery

instead uses a U or a UG sequence in mRNA, along with one or two adenines from the

poly-A tail, to generate UAA and UGA on-the-fly stop codons (Klimeš et al. 2014).

Additionally, these genomes have a highly conserved motif (downstream gene element,

or DGE) roughly 5 base pairs downstream of transcription stop sites of most genes.

These sequences are hypothesized to direct 3' mRNA cleavage and polyadenylation

factors to the pre-mRNA (Gentekaki et al. 2017; Klimeš et al. 2014). In a scenario where

the DGE is sufficient for 3' mRNA cleavage, the loss of DNA-encoded stop codons is

driven by relaxation of the selection pressure maintaining them.

We investigated all *Blastocystis* isolates for the DGE motif, finding enrichment

downstream of protein coding genes in human-isolated *Blastocystis* strains, but not

downstream of tortoise-isolated *Blastocystis* strains or other stramenopiles (Figure 2C-

D). We additionally found no evidence of transcripts lacking stop codons in the

tortoise-associated *Blastocystis* clade, suggesting that the unique transcriptional

cleavage machinery described in *Blastocystis* ST1, ST4, and ST7 (Klimeš et al. 2014;

Gentekaki et al. 2017) evolved after the divergence of the tortoise- and the human-

associated clades sequenced here.

Gene losses related to cell morphology and signaling unite the Blastocystis *genus*

Despite the substantial variation in genome diversity for all *Blastocystis* strains

sequenced here, the strains share the typical spherical, non-motile *Blastocystis* cell

morphology lacking flagella. We find that all *Blastocystis* isolates are united by

substantial gene loss related to this morphological change across multiple functional

categories. All *Blastocystis* isolates have lost nearly all cilia- and flagella- related genes

(Figure 3, Table S4-5). They have also undergone substantial reduction of dynein and

kinesin microtubule motor proteins, retaining what is likely the minimal amount of

motor proteins required for intracellular transport. This agrees with recent analyses

showing reduction in these gene families in human-associated *Blastocystis* subtypes

(Záhonová et al. 2023), and extends that observation by demonstrating that these

losses unite the genus. In addition, we find that *Blastocystis* isolates have lost

nucleotide cyclases and many genes related to signal transduction (Figure 3, Table S4-

5). As cilia are major mediators of cell signaling that use cyclic nucleotides as

intermediaries (Johnson and Leroux 2010), the loss of cilia in *Blastocystis* likely

corresponds to the loss of nucleotide cyclases and reduction in signal transduction

related genes. Thus, our sequencing of diverse *Blastocystis* strains provides strong

evidence that the flagella-free morphology and associated gene losses evolved in the

common ancestor of the *Blastocystis* genus.

Lineage-specific genes reveal specialization within the Blastocystis *genus*

We find multiple differences in gene content in between *Blastocystis* isolates

that likely reflect their ecological niche. All strains in the herbivorous tortoise-

associated *Blastocystis* clade have a larger number of genes involved in plant

Figure 3. Gene families lost or reduced in all Blastocystis strains.
See Table S2 for GO categories and InterPro domains used for defining categories.

carbohydrate degradation than the human-associated subtypes. We found five

carbohydrate-degrading enzyme families that were either uniquely present in the

tortoise isolates or in higher copy numbers, four of which are specific to degrading

plant carbohydrates. This includes polygalacturonases, β-D-xylosidases, α-L-

arabinofuranosidases, α-L-fucosidases, α-L-rhamnosidases, arabinogalactanases, and

β-hexosaminidases (Figure 4). Polygalacturonase is a pectinase that degrades plant

cell walls; it was originally characterized in plants, but is also present in

phytopathogenic microbes and in bacterial members of ruminant microbiomes (De

Lorenzo and Ferrari 2002; Deng et al. 2023). β-D-xylosidases and α-L-

arabinofuranosidases are both critical enzymes for degrading xylan, a carbohydrate

specific to plant hemicellulose (Yang et al. 2004; Lagaert et al. 2014). The GH154

arabinogalactanase gene family that is present only in the tortoise-associated clade

was recently characterized in the gut bacteria *Bacteroides thetaiotamicron* as acting on

plant-derived arabinogalactan glycans (Cartmell et al. 2018). Finally, α-L-

rhamnosidases are active on a wide variety of polysaccharide residues produced by

plants and bacteria (Guillotin et al. 2019). Together, the presence of multiple plant-

specific carbohydrate digesting enzymes in the herbivorous tortoise-associated clade

strongly suggests that, like the bacterial microbiota, *Blastocystis* utilize dietary plant

fiber as a carbon source and participate in the hindgut plant fiber fermentation that is

characteristic of herbivorous tortoises (Stevens and Hume 1995).

Ongoing horizontal gene transfer from bacteria in the Blastocystis genus

Figure 4. Clade-specific gene family expansion across *Blastocystis*, identified with InterProScan.

Horizontal gene transfer from bacteria has played an important role in

Blastocystis evolution, particularly with respect to survival in an anaerobic environment

(Eme et al. 2017). With high-quality genomes of *Blastocystis* isolates that span the

genus, in combination with the genomes of other stramenopiles, we can estimate times

of these transfer events. We re-analyzed HGT events into ST1 previously identified in

Eme et al 2017, and additionally screened all sequenced *Blastocystis* genomes for

evidence of additional transferred genes (see Methods, Screening for horizontally

transferred genes).

Previous analysis of *Blastocystis* ST1, ST4, and ST7 genomes identified 74

horizontal transfer events in *Blastocystis* ST1 from bacteria (Eme et al. 2017). We were

able to confidently identify orthologous genes for 67 of these genes (Table S6), of

which 27 genes had strong phylogenetic support for horizontal gene transfer from

bacteria to eukaryotes, 17 genes lacked strong phylogenetic support for horizontal

gene transfer, and the remaining 20 genes were ambiguous. In most ambiguous cases,

the *Blastocystis* genes were either sister to an entirely bacterial tree, there were too few

orthologs in the NR database to construct a gene tree, or the placing of the

Blastocystis sequences on the phylogeny lacked strong support.

We additionally screened all genomes sequenced for new cases of HGT. To

minimize the impact of potential genome contamination, we limited this screen to

genes that were expressed in our RNA-seq data, were shared in more than one

Blastocystis genome, and were flanked by unambiguously eukaryotic sequence to (see

Methods). We found 13 newly identified HGT cases, six which are unique to the

tortoise-associated *Blastocystis* lineage.

A

Human lineages
6 HGT

Blastocystis
16 HGT

Opalinata
18 HGT

Tortoise lineages
6 HGT

- ST3-DL
- ST3-NMH
- ST4-BT1
- ST1-JDR
- ST1-NandII
- Hermanns
- Rus-S
- Rus-B
- *Proteromonas lacertae*
- *Cafeteria burkhardae*

100/100 97.5/100
Verrucomicrobiota sequence — 305 other sequences (bacteria and eukaryotes)

99.6/100 — 55 Bacteroidota sequences
97.5/100 100/100 — 40 Bacteroidota sequences
100/100 — 15 Bacteroidota sequences
99.8/100 — 8 Planctomycetota sequences

99.4/100 — 65 gut Clostridia gut MAG sequences
96.7/100
97/100 100/100 94/100 — Clostridia gut isolate sequence
— Clostridia gut MAG sequence
— Clostridia gut MAG sequence
93/100 — Clostridia gut MAG sequence
100/100 — Clostridia gut MAG sequence
96.8/100 — Clostridia gut MAG sequence
— Clostridia gut isolate sequence

99.3/100 — Hermann's alpha-rhamnosidase 3
100/100 — Rus-S alpha-rhamnosidase 2
96.4/100 — Hermann's alpha-rhamnosidase 2
100/100 — Hermann's alpha-rhamnosidase 1
95/99 100/100 — Rus-S alpha-rhamnosidase 1
— Rus-B alpha-rhamnosidase 2
— Rus-B alpha-rhamnosidase 1

100/100 — Soil Bacteroidota sequence
— Soil Bacteroidota sequence
— Soil Bacteroidota sequence
— Soil Clostridia sequence

92.7/100 — 20 Actinomycetota sequences
100/100
98.2/100 — Bacillota sequence
— Chloroflexota sequence

91.1/100 — 17 Bacteroidota sequences
99.9/100
— Akkermansiacea sequence
96.8/100 — 11 other bacterial sequences
98.7/100 — 9 other bacterial sequences
99.9/100 — 6 Bacteroidota sequences
97.5/96 — 3 Planctomycetota sequences
93/100 — 3 other bacterial sequences
99.7/100 — 3 other bacterial sequences
100/100 — 3 other bacterial sequences
84.3/99 — 3 other bacterial sequences
— Bacteroidota sequence
— Myxococcota sequence
— Pseudomonadota sequence
— Euryarchaeota sequence
— Bacteroidota sequence

100/100

0.4

Figure 5. (A) Estimated time of transfer for 40 horizontally transferred genes in Blastocystis shared between more than one species or strain. Includes previously identified and validated cases originally identified in Eme et al. 2017 and newly-identified transfer events. (B) Gene phylogeny of tortoise isolate alpha-rhamnosidase genes and 588 homologous proteins. Support values indicate 1,000 replicates of SH-aLRT test and ultrafast bootstraps (UFboot). Branches with support values <95 for UFboot or <80 for SH-aLRT are collapsed.

Interestingly, two of the six tortoise-unique transfers are glycoside hydrolase

genes, alpha-L-rhamnosidase and arabinogalactanase, that are uniquely expanded in

this lineage and not present in the human *Blastocystis* isolates. The remaining transfers

are two GNAT family N-acetyltransferases, a cold-shock protein, and a SMI1/KNR4

family protein (Table S7).

The alpha-L-rhamnosidase gene family cleaves alpha-L-rhamnose from a large

diversity of molecules and is present across the tree of life. This gene is present in two

copies in Rus-B and Rus-S and three copies in Hermann's, and is not present in any

human *Blastocystis* isolates or in *P. lacertae*. Two genes with the alpha-L-

rhamnosidase Interpro domain were present in the *C. burkhardae* genome. We

constructed a gene phylogeny of the *C. burkhardae* genes and the tortoise genes

together with 588 homologous genes retrieved with a sequence search of the

Blastocystis and *C. burkhardae* sequences against the clustered NCBI NR database

(see Methods). The gene phylogeny was separated into two fully supported clades, one

with the *C. burkhardae* sequences, eukaryotic sequences from protists and rotifers,

and some bacterial sequences, and the other with bacterial sequences and the

Blastocystis sequences (Figure 5B). The tortoise sequences grouped together with

sequences from gut isolates and gut metagenome-assembled genomes in the

Clostridia bacterial class, which contains prevalent members of the animal gut

microbiota. Thus, the *Blastocystis* tortoise lineage likely acquired this carbohydrate

degrading enzyme from Clostridia bacteria present in the gut microbiome, suggesting a

gut-specific function.

The other horizontally transferred carbohydrate gene in the tortoise *Blastocystis*

lineage is within the recently described GH154 glycoside hydrolase gene family,

characterized in *Bacteroides thetaiotamicron* as acting on plant-derived

arabinogalactan glycans (Cartmell et al. 2018). This gene family is present exclusively in

the tortoise *Blastocystis* isolates and not found in any other *Blastocystis* isolates or any

other stramenopile genomes. A sequence search for the tortoise *Blastocystis* genes

against the clustered NCBI NR database did not return any eukaryotic hits. A gene

phylogeny of all bacterial homologs and the tortoise *Blastocystis* genes shows the

tortoise *Blastocystis* genes group together with bacteria in the *Bacteroidota* (syn.

Bacteroidetes) phylum (Figure S3).

We did not find strong evidence that the other three carbohydrate gene families

uniquely present or expanded in the tortoise *Blastocystis* lineage were horizontally

acquired from bacteria. Homologs to the polygalacturonase gene were all eukaryotic.

While the evolutionary history of this gene is likely complex as most homologs were

within Streptophyta, it was not horizontally acquired from bacteria. β-D-xylosidase and

α-L- arabinofuranosidase both appeared to have complex evolutionary histories but

without conclusive evidence of horizontal gene transfer. Specifically, several of the β-

D-xylosidase genes in the tortoise lineages had no eukaryotic homologs, but a gene

phylogeny of all β-D-xylosidase genes including the human isolates did not yield a

well-supported branch of the tortoise genes nested within bacteria. Similarly, the α-L-

arabinofuranosidase gene had both bacterial and eukaryotic homologs, but there was

not strong support for the placement of the *Blastocystis* sequences on this phylogeny

either with the eukaryotic sequences or with the bacterial sequences.

Together, these results indicate that horizontal gene transfer played a major role

in all evolutionary transitions investigated in this work: at the evolutionary transition

from free-living to gut symbiont, in the *Blastocystis* genus broadly, and is an ongoing

method of genetic diversification in this organism.

Discussion

The stramenopile protist *Blastocystis* is the most prevalent protist in the human

gut and is widespread throughout vertebrates. Here we present contiguous and

complete genomes of eight *Blastocystis* strains, five isolated from humans and three

from herbivorous tortoises. Long-read sequencing coupled with metagenomic

assembly techniques enabled us to assemble three of these genomes from xenic

cultures, which has been technically infeasible with standard short-read approaches.

Comparative genomic analyses between these *Blastocystis* genomes alongside other

closely related stramenopiles revealed evolutionary changes that the *Blastocystis*

genus has undergone to colonize a wide variety of animal guts.

The herbivorous tortoise isolates of *Blastocystis* diverged substantially from the

human isolated *Blastocystis,* and have many genomic features not seen in the human

isolates. We found substantial changes in genome size, with human isolates

possessing significantly shorter introns and intergenic regions than the tortoise

isolates, and substantially lower GC content in tortoise isolates. The GC content of the

tortoise isolates is substantially lower than the genomes of most sequenced

stramenopiles; of the 299 stramenopiles with sequenced genomes deposited in

GenBank, only three genomes have lower GC content.

It is not clear whether introns and intergenic regions have shrunk in size after the

split of the tortoise-human isolates, or if they have lengthened along the tortoise

lineage. Genomic streamlining is a frequently observed process in symbiotic and

parasitic organisms (McCutcheon and Moran 2012; Leckenby and Hall 2015) and some

parasitic eukaryotes have entirely or almost entirely lost introns (Wang et al. 2018;

Wilihoeft et al. 2001). The phenomenon of intergenic and intron size shrinking in

Blastocystis likely reflects this general process, but the unexpected differences

between different host-associated *Blastocystis* isolates highlights that this is an

ongoing evolutionary process that can proceed at different rates.

Despite these substantial differences in genomic streamlining, all *Blastocystis*

isolates sequenced here have undergone loss of the same genes associated with the

more complex flagellated cell morphology of other stramenopiles. *Blastocystis* isolates

have lost nearly all cilia- and flagella-related proteins, substantially reduced the number

of microtubule motor proteins, and have lost all nucleotide cyclases and many signal

transduction related genes. These changes unite all *Blastocystis* isolates,

demonstrating shared biology across the genus despite large evolutionary distances

and differences in genome structure between isolates.

We additionally find evidence consistent with host adaptation within

Blastocystis. The *Blastocystis* strain isolated from herbivorous tortoises contained

many carbohydrate digesting enzymes not present in other *Blastocystis* isolates (Figure

4). Two of these gene families were acquired via horizontal transfer from bacteria, and

in one case via horizontal transfer from bacteria that also colonize the animal gut. The

presence of plant-specific carbohydrate digesting enzymes suggests that, like some

ciliates and many gut bacteria, *Blastocystis* may ferment plant carbohydrates, a critical

element of energy homeostasis in predominantly plant-eating animals (Stevens and

Hume 1995).

While *Blastocystis* is widespread across vertebrates, there is limited evidence for

co-phylogeny of *Blastocystis* subtypes with their host (Figure 1). Instead, the

Blastocystis phylogeny more closely resembles that of spore-forming gut bacteria

(Moeller et al. 2016), with host switching as a common phenomenon. Because of the

lack of strong co-phylogenetic signal, open questions remain about *Blastocystis* host

association, including what the barriers to colonizing different hosts are and how

multiple strains or subtypes (commonly termed 'mixed infections') are maintained, will

require a denser sampling of genomes across the *Blastocystis* tree of life. As human

associated *Blastocystis* subtypes are not monophyletic, understanding the biology of

the protists that impact humans will require considering *Blastocystis* subtypes that are

present in the guts of diverse vertebrate hosts, and how these subtypes function in

different environments. Finally, the vast differences seen in the phylogenetically distinct

tortoise *Blastocystis* isolates sequenced here compared to previous sequenced

human-associated *Blastocystis* isolates highlights that substantial genetic diversity in

this genus has yet to be explored.

Five of the eight *Blastocystis* strains sequenced in this work were grown in the

presence of other microbes. Despite the challenges of generating highly contiguous

and contamination free genomes from mixed culture metagenomic sequencing, we

successfully combined advances in deep long read DNA sequencing and metagenomic

assembly techniques to acquire high-quality genomes from xenic cultures. To address

the challenge of annotating genes in genomes where stop codons are not universal, we

utilized RNA-sequencing and state-of-the-art annotation pipelines. These two technical

accomplishments were essential for enabling robust comparative genomic analyses of

Blastocystis and its closest sequenced relatives. The strategy established in this study

can be extended in future work to the many eukaryotic microbes that cannot be

cultured axenically. The resulting genomes will fill critical gaps in our understanding of

eukaryotic microbes and the evolution of this domain of life.

Methods

Strain cultivation

Blastocystis strains NandII (ATCC 50177), JDR (ATCC 50587), BT1 (ATCC

50608), DL (ATCC 50626), and NMH (ATCC 50754) were obtained from the American

Type Culture Collection. The JDR strain was labeled as subtype 3 based on rRNA

analysis, but 18S rRNA gene sequencing and subsequent genome sequencing

revealed in fact that this strain belongs to subtype 1. All strains were isolated from

human sources. Cultures were incubated under anaerobic conditions at 37C on

biphasic egg slant media overlaid with Locke's solution supplemented with 25% heat

inactivated horse serum (ATCC Medium 1671). Media was pre-reduced for 24 hours

before use. ST1 and ST4 strains were maintained in axenic culture, while ST3 and

tortoise isolates were grown in xenic culture alongside other microbes present in the

original samples. These xenic culture strains proved resistant to axenization.

Blastocystis strains from tortoises were isolated from fresh fecal material that

was processed within 8 hours of sampling. Fecal material was resuspended in sterile

PBS, strained using a 70µm cell strainer, and centrifuged at 700xg for 5 minutes at

room temperature. The pellet was then washed 3 times with PBS before being

resuspended in 80% Percoll. This was overlaid with 40% Percoll and centrifuged at

100xg for 10 minutes at room temperature. Protists were removed from the top of the

40% Percoll layer, washed in PBS, and resuspended in pre-reduced growth medium.

DNA extraction and sequencing

High molecular weight DNA was isolated from *Blastocystis* cultures with a

modified CTAB-based protocol (adapted from McLay 2017) that minimizes

polysaccharide contamination. *Blastocystis* cells were collected from cultures by

pelleting with centrifugation at 800xg for 5 minutes and snap frozen at -80° C. Frozen

pellets were gently resuspended in two volumes of a high salt CTAB lysis buffer (100

mM Tris pH 8, 1.2 M NaCl, 10 mM EDTA, 4% CTAB, 1% PVP-40). 200 µg/mL RNAse A

and 200 µg/mL proteinase K were added before incubation at 60° C for 90 minutes.

One volume of phenol:chloroform:isoamyl alcohol (25:24:1) was added and samples

were mixed end over end for 15 minutes. Samples were centrifuged at 17000g for 10

minutes and the upper aqueous layer was recovered. Two additional extractions were

performed with pure chloroform, recovering the upper aqueous layer each time. DNA

was next precipitated with 2 volumes of a low salt CTAB precipitation buffer (100 mM

Tris pH 8, 10 mM EDTA, 2% CTAB) by end over end rotation at 60° C for 2.5 hours.

DNA was pelleted by centrifugation at 17,000xg for 20 minutes. DNA pellets were

washed twice with 75% ethanol and resuspended in 10 mM Tris pH 8. Genomic DNA

was further cleaned with 1.8 volumes of Ampure XP beads according to the

manufacturer's protocol (Beckman Coulter, Brea, CA). RNA was isolated from

Blastocystis cultures with a Trizol extraction following manufacturer's protocols

(Invitrogen, Waltham, MA).

Nanopore sequencing libraries for all human isolates and for the Rus-B strain

were prepared using the Genomic DNA by Ligation protocol SQK-LSK109 (Oxford

Nanopore Technologies, Oxford, UK) following the manufacturer's instructions,

sequenced on a full FLO-MIN106D R9 flow cell on a MinION sequencer, and

basecalled with guppy version 5.0.16. Nanopore sequencing libraries for the Rus-S and

Hermann's isolates were prepared using the SQK-LSK110 kit LSK109 (Oxford

Nanopore Technologies, Oxford, UK) and sequenced on a full R10.4.1 flow cell, and

basecalled with dorado version 0.5.3 (https://github.com/nanoporetech/dorado).

Short read Illumina DNA and RNA libraries were prepared using the KAPA

HyperPlus PCR-free and KAPA mRNA capture kits (Roche, Basel, Switzerland),

respectively, following manufacturer's instructions. RNA integrity and library fragment

size distribution were quantified using an Bioanalyzer 2100 (Agilent, Santa Clara, CA).

Illumina DNA and RNA libraries for all human *Blastocystis* isolates and for the Rus-S

and Hermann's tortoise isolates were sequenced on an Illumina NextSeq 500 (Illumina,

San Diego, CA), and libraries for the Rus-B tortoise isolate were sequenced on an

Illumina MiSeq v3.

Genome assembly and decontamination

Basecalled nanopore reads were assembled using Flye version 2.9.2 with the --

meta option enabled (Kolmogorov et al. 2019, 2020). Assembly polishing was

performed using four rounds of nanopore-read polishing using racon version 1.4.3

(Vaser et al. 2017), three rounds of Illumina DNA polishing using PolyPolish version

0.5.0 (Wick and Holt 2022), and 1 round of short-read polishing using POLCA as

implemented in MaSuRCA version 4.0.9 (Zimin et al. 2013).

For the ST4 strain BT1, Hi-C data was generated using Phase Genomics

(Seattle, WA) Proximo Hi-C 2.0 Kit, and the Phase Genomics Proximo Hi-C genome

scaffolding platform was used to create chromosome scale scaffolds from the hybrid

assembled genome. The Hi-C scaffolding resulted only in modest improvements to the

original assembly quality; the number of scaffolds reduced from 45 to 34, and the

assembly N50 improved to 882 from 876.

We used three different methods to separate eukaryotic, *Blastocystis*-originating

sequence from other microbial sequences for xenic and axenic assemblies. For xenic

assemblies, sequences were binned using MetaBat2 version 2.17.60 as a binning

algorithm, taxonomically annotated using MetaEuk taxtocontig, and additionally

annotated as likely eukaryotic or prokaryotic using EukRep (Kang et al. 2019; Levy

Karin et al. 2020; West et al. 2018). For binning, short DNA sequencing reads were

aligned to the polished assembly using minimap2 version 2.26-r1175 and summarized

using the utility jgi_summarize_bam_contig_depths (Li 2018). Metaeuk taxonomy

prediction was performed by running version

1f7d7023bdac3d46964ca380bef1b2848685501e, metaeuk easy-predict with mmseqs

Swiss-Prot database build and metaeuk taxtocontig with the options '--majority 0.5 --

tax-lineage 1 --lca-mode 2'. EukRep v0.6.6. was run in 'balanced' mode.

Eukaryotic contigs were identified through a union of MetaEuk contigs

annotated as eukaryotic, removing any not identified as eukaryotic by EukRep. We

found that MetaEuk was not able to identify *Blastocystis* contigs reliably at lower

taxonomic levels than the superkingdom Eukaryota, which we validated by running the

pipeline on existing NCBI genomes. Instead, any contigs that were annotated as

eukaryotic were considered as potentially *Blastocystis* in origin. We cross-referenced

these eukaryotic contigs with metabat bins, finding bins corresponding to eukaryotic

sequence. In two cases, eukaryotic sequences that appeared to be *Blastocystis* in

origin were split across two metabat bins, and these bins were combined. We removed

any sequences from these bins that were annotated by MetaEuk as non-eukaryotic.

We verified that these final cleaned bins corresponded to *Blastocystis* by BLASTing the

NCBI RefSeq *Blastocystis* coding gene sequences against these bins. In several cases,

we found bins that were annotated as eukaryotic but did not have any hits to

Blastocysis coding sequences. These bins were discarded as likely originating from

non-*Blastocystis* eukaryotic organisms in the culture.

We found that contigs containing ribosomal RNA genes were often not binned

together with protein coding genes, likely due to their highly repetitive nature and

differences in DNA coverage. We searched for contigs containing 18S rRNA genes by

BLASTing 18S rRNA genes against the full assembly. Contigs with full length and high

percent identity hits to any Blastocystis 18S genes that were also annotated as

eukaryotic by MetaEuk were included in final assemblies.

For axenic assemblies, sequenced were not binned with MetaBat2, but MetaEuk

and EukRep were performed as described above to identify potential contaminating

sequences. This resulted in two contigs being removed from the BT1 assembly, and no

contigs being removed from Nand or JDR assemblies.

Genome completeness was determined with sequenced-based and gene-based

approaches. For the sequence based approach, we applied a *k*-mer frequency-based

approach with GenomeScope v2.0 (Ranallo-Benavidez et al. 2020). Gene-based

completeness was determined using BUSCO v5.6.1 with the lineage dataset

eukaryote_odb10 in the genome mode (Manni et al. 2021).

Annotating repetitive elements

Repetitive elements in *Blastocystis* were annotated with a combination of

RepeatMasker and RepeatModeler using the Dfam database version 3.8 (Smit et al.

2013; Flynn et al. 2020; Storer et al. 2021). Repetitive elements were predicted from all

genomes with RepeatModeler version 2.0.5 with LTRStruct enabled. The predicted

transposable elements from all *Blastocystis* genomes that had a putative predicted

annotation were combined, and repetitive elements were tabulated in each genome

with a combination of the custom repeat library created with RepeatModeler and the

Dfam v3.8 database with RepeatMasker 4.1.7-p1.

Gene annotation

Blastocystis genes occasionally do not encode stop codons, which confounds

prediction via open reading frame and requires RNA-seq based evidence (Klimeš et al.

2014; Gentekaki et al. 2017). Without consideration of lack of stop codons, common

gene annotation errors include insertion of erroneous introns and combining adjacent

genes. To accurately annotate the *Blastocystis* genomes, we created a pipeline that

used a combination of *de novo* gene prediction with Braker3 and transcriptome

assembly, combining all sets of gene models with Mikado (Venturini et al. 2018; Gabriel

et al. 2023).

RNA-seq reads were aligned to assemblies with STAR version 2.7.11b (Dobin et

al. 2013), with a minimum of 3 supporting reads required for each junction. Transcript

assemblies were generated with Stringtie version 2.2.3 (Pertea et al. 2015), which uses

RNA-seq alignments against a genome to assemble transcripts. Because *Blastocystis*

has very short intergenic regions, transcript assemblers can occasionally combine

multiple transcripts into one. To address this, assembled transcripts were split if an

exon had zero RNA-seq coverage. Assembled transcripts less than 100 bp long were

discarded. To generate *de novo* gene predictions, braker version 3.0.8 was run with

OrthoDB11 Eukaryota proteins as protein evidence and the aligned RNA-seq as

transcript evidence (Gabriel et al. 2023; Kuznetsov et al. 2022). The Mikado transcript

scoring file (available in code repository) was optimized to choose transcripts with

strictly RNA-seq supported introns, but to prioritize genes containing stop codons so

long as the intron boundaries were strictly supported.

Functional annotation and orthology inference

Proteins from *Cafeteria burkhardae* BVI (formerly *Cafeteria roenbergensis* BVI),

and predicted proteins from *Proteromonas lacertae* were used for comparative

genomics analyses (Záhonová et al. 2023; Hackl et al. 2020). The genome assembly,

transcripts, and proteins for *C. burkhardae* was downloaded from GenBank under

accession GCA_008330645.1. The genome sequence for *Proteromonas lacertae* was

downloaded from GenBank under accession GCA_002245135.1. As no predicted

proteins or RNA-seq data was available for *P. lacertae*, genes were predicted from this

genome using the Braker3 pipeline, using OrthoDB11 Eukaryota protein sequences as

evidence. Proteins from all *Blastocystis* assemblies, *C. burkhardae*, and *P. lacertae*

were annotated using a combination of InterProScan version 5.66 (Jones et al. 2014) ,

DeepLoc2 to predict protein localization (Thumuluri et al. 2022), and dbCAN version

4.0.0 to annotate carbohydrate active enzymes (Zhang et al. 2018). Orthologous

sequences were determined using OrthoFinder2 with diamond as the search algorithm

and mcl as the clustering algorithm (Emms and Kelly 2019; Buchfink et al. 2015; Li et

al. 2003).

Genes related to cilia and flagella, microtubule motor proteins, signal

transduction, ion channels, and nucleotide cyclase were identified using either GO

categories or Interpro domains corresponding to these categories, and are listed in

Table S4. Carbohydrate-degrading enzymes were identified based on the gene

annotation methods mentioned above; specifically, polygalacturonase enzymes were

identified with the PANTHER domain PTHR33928, beta-D-xylosidase with the Interpro

domain IPR044993, a-L-arabinofuranosidase with the Interpro domain IPR010720, a-L-

rhamnosidase with the PANTHER domain PTHR34987, and the GH154

arabinogalactanase with the PANTHER domain PTHR35339.

Identifying motifs adjacent to transcript ends

To identify motifs enriched downstream of transcript ends, we extracted a 60 bp

window up and downstream of all annotated transcript ends in all *Blastocystis*

genomes sequenced here, as well as *P. lacertae* and *C. burkhardae*. Motifs were

identified separately in each genome using STREME version 5.5.7 (Bailey 2021). A

motif similar to the one identified by Klimeš et al 2014 was identified in both ST1, both

ST3, and the ST4 human isolate, but no significant and informative motifs were

identified in the tortoise-isolated *Blastocystis* isolates or in the *P. lacertae* or *C.*

burkhardae genomes. We then combined the sequences from all human isolates and

ran STREME on this combined dataset to generate a sequence motif (Figure 2).

To determine enrichment of this motif in all genomes, we searched all transcript

ends for matches to a degenerate version of the core region of the sequence motif,

TGTTGTT, allowing up to two mismatches. As this motif is inherently an AT-rich

sequence, the probability of observing it by chance is a function of the GC content of

the sequence that is searched. We calculated the expected probability of seeing the 7-

base pair in each genome given the transcript window GC content and the number of

transcripts, and calculated the enrichment of this motif as the log of the observed motif

counts divided by the background expectation value (Figure 2).

Strain and species phylogenies

The *Blastocystis* genus phylogeny was constructed using 18S rRNA sequences

deposited in GenBank (Table S1) along with 18S rRNA sequences extracted from each

Blastocystis assembly and from the *Proteromonas lacertae* genome as an outgroup.

Sequences were aligned with MAFFT v7.505 (Katoh and Standley 2013) and a

maximum likelihood phylogeny constructed using IQ-TREE version 2.3.5 (Minh et al.

2020) using with the GTR sequence model, 1,000 ultrafast bootstrap and SH-aLRT

replicates, and rooted on *P. lacertae*.

To construct a species phylogeny of all species and strains with genomic data

used in this study, 13 conserved protein-coding genes identified with BUSCO version

5.6.1 as described above were extracted from *Cafeteria burkhardae*, *Proteromonas*

lacertae, and all *Blastocystis* strains sequenced here. Proteins were aligned individually

with MAFFT v7.505 and concatenated. Best-fit models for each partition were selected

using a relaxed hierarchical clustering algorithm as implemented by ModelFinder in IQ-

TREE version 2.3.5 (Chernomor et al. 2016; Minh et al. 2020; Kalyaanamoorthy et al.

2017). 1,000 ultrafast bootstrap and SH-aLRT replicates were performed.

Gene contamination and horizontal gene transfer screens

Blastocystis genomes were screened for bacterial contamination and for

possible horizontal gene transfer events using a modified version of an alien index

score (Gladyshev et al. 2008; Wisecaver et al. 2016). The alien index score compares

the best bit score of a query sequence to a group of related taxa versus to all other

unrelated groups, and normalizes this score by the maximum possible bit score of the

query sequence. It is determined by the formula $AI = \frac{bbsO}{bbsS} - \frac{bbsI}{bbs}$, where bbsO is the

bit score of the best hit to a species outside of the group lineage, bbsI is the bit score

of the best hit to a species within the group lineage (skipping self hits), and bbsS is the

bit score of the query aligned to itself. The alien index was calculated for each

696 *Blastocystis* protein by querying proteins against the NCBI nr database using

697 DIAMOND version 2.1.6 (Buchfink et al. 2015)(code availability:

698 https://github.com/allind/alienindex). The self-group was designated as *Blastocystis*

699 (taxid: 12967), and the group lineage was set as all eukaryotes (taxid: 2759).

700  As an additional control for whether our genomes had high levels of genome

701 contamination, we determined what fraction of genes with high alien index scores were

702 likely transcribed by *Blastocystis.* Our RNA-seq strategy involved poly-A purification,

703 which removes non-polyadenylated transcripts. As polyadenylation is unique to

704 eukaryotes, this removes bacterial transcripts. We quantified gene expression using

705 kallisto version v0.48.0 (Bray et al. 2016), and determined whether genes with alien

706 index scores greater than 0.2 were expressed and therefore likely truly integrated into

707 the *Blastocystis* genomes. The number of genes with alien index scores above 0.2

708 ranged from 59-81 genes across all sequenced genomes, and greater than 85% were

709 expressed in all genomes. Genes with alien index scores greater than 0.2 that were

710 expressed at TPM > 1 and had orthologs in one or more other *Blastocystis* genomes

711 were further examined for evidence of horizontal gene transfer with gene tree

712 reconstruction.

714 *Gene tree reconstruction for horizontal transfer*

715  Phylogenetic trees for putative horizontal gene transfer candidates were

716 constructed from all diamond blastp hits using the NCBI NR database clustered at

717 90% sequence identity and 90% sequence length (Buchfink et al. 2015). Sequence hits

718 that were outliers in length were excluded from downstream analyses. Protein

sequences were aligned with MAFFT v7.505 (Katoh and Standley 2013), and maximum

likelihood phylogenies were constructed using IQ-TREE version 2.3.5 using

ModelFinder to determine the best-fit substitution model, and with 1,000 ultrafast

bootstrap replicates and 1,000 replicates of the SH-like approximate likelihood ratio

test (Guindon et al. 2010). Branches with SH-aLRT support less than 70 and UF-boot

less than 80 were collapsed and trees were midpoint rooted before being examined for

horizontal transfer.

Acknowledgements

We thank members of the Pollard lab and the Bhatt lab for helpful discussions. This

work was supported NHLBI grant#R01-HL160862, Chan Zuckberg Biohub SF and

Gladstone Institutes to K.S.P, by the Benioff Center for Microbiome Medicine Trainee

Pilot Award and Georgia Institute of Technology to A.L.L. A.SB. is supported by NIH

R01AI143757 and R01AI148623, as well as a Stand Up 2 Cancer grant and a Paul Allen

Distinguished Investigator Award.

Author Contributions

Conceptualization – A.L.L, A.S.B, K.S.P.; Methodology – A.L.L., N.A.M, E.S.G, A.S.B,

K.S.P.; Investigation – A.L.L, K.S.P.; Resources – A.S.B., K.S.P.; Writing – original draft

– A.L.L., N.A.M, E.S.G, A.S.B, K.S.P; Writing – review & editing - A.L.L., N.A.M, E.S.G,

A.S.B, K.S.P; Visualization – A.L.L.; Supervision – K.S.P., A.S.B.; Funding acquisition –

A.L.L., K.S.P., A.S.B.

Competing Interests

The authors declare no competing interests.

DATA ACCESS

Genome assemblies and annotations are available on Genbank (Table 2) and are

additionally available on Figshare at

https://figshare.com/projects/Blastocystis_genomes/186136. As no gene annotations

were available for *Proteromonas lacertae*, BRAKER3 predictions were generated and

have been made publicly available in this repository as well. RNA sequencing for all

strains is available from the SRA under BioProject PRJNA1041441. Code to calculate

the alien index is available at https://github.com/allind/alienindex. All other code used

in this study is available at https://github.com/allind/blastannotate. All alignments and

phylogenies generated in this work are available at

https://doi.org/10.6084/m9.figshare.28143281.v1.

Table 2. NCBI accessions for genomes generated here.

Strain	NCBI Bioproject
Rus-B (also Blastoise)	PRJNA1084774
Rus-S	Processing on Genbank, currently available on Figshare
Hermann's	Processing on Genbank, currently available on Figshare
ST1-NandII	PRJNA1084796

ST1-JDR	PRJNA1084798
ST3-DL	PRJNA1084795
ST3-NMH	PRJNA1084799
ST4-BT1	PRJNA1084775

Figure legends

Figure 1. (A) 18S rRNA maximum likelihood (ML) phylogeny of Blastocystis subtypes, with *Proteromonas lacertae* as an outgroup. Support values are 1000 replicates of ultrafast bootstrap and SH-aLRT tests (Guindon et al. 2010; Hoang et al. 2018). Strains in red were sequenced in this study. (B) 13-gene concatenation ML phylogeny for all strains and species with sequenced genomes used in this study. Important evolutionary transitions are highlighted. Images represent cellular morphology of *Cafeteria burkhardae*, *Proteromonas lacertae*, and *Blastocystis*. (C) Assembly statistics of Blastocystis strains sequenced in this work compared with previously sequenced and annotated genomes. Stop-codon aware and not stop codon aware designations refer to gene annotation methodologies that take into consideration that Blastocystis genes can lack stop codons or do not consider this, respectively. The ST7 N50 is a gapped scaffold N50 generated with Sanger sequencing scaffolding.

Figure 2. (A) Intron lengths and (B) Intergenic lengths in tortoise isolates (left) and human isolates right). (C) Sequence motif enriched adjacent to transcript ends in ST1, ST3, and ST4. Motif generated with STREME (Bailey 2021). This motif is referred to as

"downstream gene element" (DGE). (D) Enrichment (log(observed/expected)) of

transcripts with downstream conserved element within 30 bp upstream or downstream

of transcript end, normalized by expected DGE occurrence given GC content.

Figure 3. Gene families lost or reduced in all Blastocystis strains. See Table S2 for GO

categories and InterPro domains used for defining categories.

Figure 4. Clade-specific gene family expansions across *Blastocystis,* identified with

InterProScan.

Figure 5. (A) Estimated time of transfer for 40 horizontally transferred genes

in *Blastocystis* shared between more than one species or strain. Includes

previously identified and validated cases originally identified in (Eme et al. 2017)

and newly-identified transfer events. (B) Gene phylogeny of tortoise isolate

alpha-rhamnosidase genes and 588 homologous proteins. Support values indicate

1,000 replicates of SH-aLRT test and ultrafast bootstraps (UFboot). Branches with

support values <95 for UFboot or <80 for SH-aLRT are collapsed.

Figure S1. Synteny of *Blastocystis* genomes based on reciprocal best blast hits. Plot

created with ODP (Schultz et al. 2023).

Figure S2. Median percent identity across 753 single-copy orthologs in *Blastocystis*.

Figure S3. Gene phylogeny of tortoise Blastocystis GH154 arabinogalactanase

enzyme. Support values indicate 1,000 replicates of SH-aLRT test and ultrafast

bootstrap. Branches with aLRT support <80 and UFboot support <90 are collapsed.

Tree is midpoint rooted.

Table S1. GenBank accessions of 18S rRNA sequences used to construct *Blastocystis*

phylogeny.

Table S2. Amount and percentages of genome masked per strain by RepeatMasker.

Table S3. GO and IPR categories used for gene loss described in Figure 3.

Table S4. Genes containing InterPro domains per genome assembly.

Table S5. Percentage of high alien index genes per genome with poly-A purified RNA-

seq expression support.

Table S6. Validation of previously reported ST1-NandII horizontal gene transfers.

Table S7. Newly identified HGT events in *Blastocystis*.

References

Asnicar F, Berry SE, Valdes AM, Nguyen LH, Piccinno G, Drew DA, Leeming E, Gibson R, Le Roy C, Khatib HA, et al. 2021. Microbiome connections with host metabolism and habitual diet from 1,098 deeply phenotyped individuals. *Nat Med* **27**: 321–332.

Bailey TL. 2021. STREME: accurate and versatile sequence motif discovery. *Bioinforma Oxf Engl* **37**: 2834–2840.

Beghini F, Pasolli E, Truong TD, Putignani L, Cacciò SM, Segata N. 2017. Large-scale comparative metagenomics of Blastocystis, a common member of the human gut microbiome. *ISME J* **11**: 2848–2863.

Boreham PFL, Stenzel DJ. 1993. Blastocystis in Humans and Animals: Morphology, Biology, and Epizootiology. In *Advances in Parasitology* (eds. J.R. Baker and R. Muller), Vol. 32 of, pp. 1–70, Academic Press https://www.sciencedirect.com/science/article/pii/S0065308X08602067 (Accessed October 24, 2023).

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**: 525–527.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND. *Nat Methods* **12**: 59–60.

Campo J del, Bass D, Keeling PJ. 2020. The eukaryome: Diversity and role of microeukaryotic organisms associated with animal hosts. *Funct Ecol* **34**: 2045–2054.

Cartmell A, Muñoz-Muñoz J, Briggs JA, Ndeh DA, Lowe EC, Baslé A, Terrapon N, Stott K, Heunis T, Gray J, et al. 2018. A surface endogalactanase in Bacteroides thetaiotaomicron confers keystone status for arabinogalactan degradation. *Nat Microbiol* **3**: 1314–1326.

Chernomor O, von Haeseler A, Minh BQ. 2016. Terrace Aware Data Structure for Phylogenomic Inference from Supermatrices. *Syst Biol* **65**: 997–1008.

David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin AS, Varma Y, Fischbach MA, et al. 2014. Diet rapidly and reproducibly alters the human gut microbiome. *Nature* **505**: 559–563.

De Lorenzo G, Ferrari S. 2002. Polygalacturonase-inhibiting proteins in defense against phytopathogenic fungi. *Curr Opin Plant Biol* **5**: 295–299.

Deng Q, Sun X, Gao D, Wang Y, Liu Y, Li N, Wang Z, Liu M, Wang J, Wang Q. 2023. Characterization of Two Novel Rumen-Derived Exo-Polygalacturonases: Catalysis and Molecular Simulations. *Microorganisms* **11**: 760.

Denoeud F, Roussel M, Noel B, Wawrzyniak I, Da Silva C, Diogon M, Viscogliosi E, Brochier-
Armanet C, Couloux A, Poulain J, et al. 2011. Genome sequence of the stramenopile
Blastocystis, a human anaerobic parasite. *Genome Biol* **12**: R29.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinforma Oxf Engl* **29**: 15–21.

Eme L, Gentekaki E, Curtis B, Archibald JM, Roger AJ. 2017. Lateral Gene Transfer in the
Adaptation of the Anaerobic Parasite Blastocystis to the Gut. *Curr Biol CB* **27**: 807–820.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol* **20**: 238.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020.
RepeatModeler2 for automated genomic discovery of transposable element families. *Proc
Natl Acad Sci* **117**: 9451–9457.

Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, Stanke M. 2023.
BRAKER3: Fully Automated Genome Annotation Using RNA-Seq and Protein Evidence
with GeneMark-ETP, AUGUSTUS and TSEBRA. *BioRxiv Prepr Serv Biol*
2023.06.10.544449.

Gantois N, Lamot A, Seesao Y, Creusy C, Li L-L, Monchy S, Benamrouz-Vanneste S,
Karpouzopoulos J, Bourgain J-L, Rault C, et al. 2020. First Report on the Prevalence and
Subtype Distribution of Blastocystis sp. in Edible Marine Fish and Marine Mammals: A
Large Scale-Study Conducted in Atlantic Northeast and on the Coasts of Northern
France. *Microorganisms* **8**: 460.

Gentekaki E, Curtis BA, Stairs CW, Klimeš V, Eliáš M, Salas-Leiva DE, Herman EK, Eme L,
Arias MC, Henrissat B, et al. 2017. Extreme genome diversity in the hyper-prevalent
parasitic eukaryote Blastocystis. *PLOS Biol* **15**: e2003769.

Godoy-Vitorino F, Goldfarb KC, Karaoz U, Leal S, Garcia-Amado MA, Hugenholtz P, Tringe
SG, Brodie EL, Dominguez-Bello MG. 2012. Comparative analyses of foregut and
hindgut bacterial communities in hoatzins and cows. *ISME J* **6**: 531–541.

Gough R, Barratt J, Stark D, Ellis J. 2020. Diversity profiling of xenic cultures of Dientamoeba
fragilis following systematic antibiotic treatment and prospects for genome sequencing.
Parasitology **147**: 29–38.

Guillotin L, Kim H, Traore Y, Moreau P, Lafite P, Coquoin V, Nuccio S, de Vaumas R,
Daniellou R. 2019. Biochemical Characterization of the α-l-Rhamnosidase DtRha from
Dictyoglomus thermophilum: Application to the Selective Derhamnosylation of Natural
Flavonoids. *ACS Omega* **4**: 1916–1922.

Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New
Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the
Performance of PhyML 3.0. *Syst Biol* **59**: 307–321.

Hackl T, Martin R, Barenhoff K, Duponchel S, Heider D, Fischer MG. 2020. Four high-quality
draft genome assemblies of the marine heterotrophic nanoflagellate Cafeteria
roenbergensis. *Sci Data* **7**: 29.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the
Ultrafast Bootstrap Approximation. *Mol Biol Evol* **35**: 518–522.

Jinatham V, Maxamhud S, Popluechai S, Tsaousis AD, Gentekaki E. 2021. Blastocystis One
Health Approach in a Rural Community of Northern Thailand: Prevalence, Subtypes and
Novel Transmission Routes. *Front Microbiol* **12**.
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2021.746340/f
ull (Accessed December 13, 2024).

Johnson J-LF, Leroux MR. 2010. cAMP and cGMP signaling: sensory systems with prokaryotic
roots adopted by eukaryotic cilia. *Trends Cell Biol* **20**: 435–444.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell
A, Nuka G, et al. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**: 1236–1240.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**: 587–589.

Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an adaptive
binning algorithm for robust and efficient genome reconstruction from metagenome
assemblies. *PeerJ* **7**: e7359.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol Biol Evol* **30**: 772–780.

Keeling PJ, Burki F, Wilcox HM, Allam B, Allen EE, Amaral-Zettler LA, Armbrust EV,
Archibald JM, Bharti AK, Bell CJ, et al. 2014. The Marine Microbial Eukaryote
Transcriptome Sequencing Project (MMETSP): Illuminating the Functional Diversity of
Eukaryotic Life in the Oceans through Transcriptome Sequencing. *PLOS Biol* **12**:
e1001889.

Khaled S, Gantois N, Ly AT, Senghor S, Even G, Dautel E, Dejager R, Sawant M, Baydoun M,
Benamrouz-Vanneste S, et al. 2020. Prevalence and Subtype Distribution of Blastocystis
sp. in Senegalese School Children. *Microorganisms* **8**: 1408.

Klimeš V, Gentekaki E, Roger AJ, Eliáš M. 2014. A Large Number of Nuclear Genes in the
Human Parasite Blastocystis Require mRNA Polyadenylation to Create Functional
Termination Codons. *Genome Biol Evol* **6**: 1956–1961.

Kolmogorov M, Bickhart DM, Behsaz B, Gurevich A, Rayko M, Shin SB, Kuhn K, Yuan J,
Polevikov E, Smith TPL, et al. 2020. metaFlye: scalable long-read metagenome assembly
using repeat graphs. *Nat Methods* **17**: 1103–1110.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using
repeat graphs. *Nat Biotechnol* **37**: 540–546.

Kostka M. 2017. Opalinata. In *Handbook of the Protists* (eds. J.M. Archibald, A.G.B. Simpson,
and C.H. Slamovits), pp. 543–565, Springer International Publishing, Cham
https://doi.org/10.1007/978-3-319-28149-0_4 (Accessed July 28, 2023).

Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, Zdobnov EM.
2022. OrthoDB v11: annotation of orthologs in the widest sampling of organismal
diversity. *Nucleic Acids Res* **51**: D445–D451.

Lagaert S, Pollet A, Courtin CM, Volckaert G. 2014. β-Xylosidases and α-l-
arabinofuranosidases: Accessory enzymes for arabinoxylan degradation. *Biotechnol Adv*
32: 316–332.

Leckenby A, Hall N. 2015. Genomic changes during evolution of animal parasitism in
eukaryotes. *Curr Opin Genet Dev* **35**: 86–92.

Levy Karin E, Mirdita M, Söding J. 2020. MetaEuk—sensitive, high-throughput gene discovery,
and annotation for large-scale eukaryotic metagenomics. *Microbiome* **8**: 48.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**: 3094–
3100.

Li L, Stoeckert CJ, Roos DS. 2003. OrthoMCL: identification of ortholog groups for eukaryotic
genomes. *Genome Res* **13**: 2178–89.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and
Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for
Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol Biol Evol* **38**: 4647–4654.

McCutcheon JP, Moran NA. 2012. Extreme genome reduction in symbiotic bacteria. *Nat Rev*
Microbiol **10**: 13–26.

McLay T. 2017. High quality DNA extraction protocol from recalcitrant plant tissues v1.
https://www.protocols.io/view/high-quality-dna-extraction-protocol-from-recalcit-i8jchun
(Accessed January 3, 2025).

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear
R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in
the Genomic Era. *Mol Biol Evol* **37**: 1530–1534.

Moeller AH, Caro-Quintero A, Mjungu D, Georgiev AV, Lonsdorf EV, Muller MN, Pusey AE,
Peeters M, Hahn BH, Ochman H. 2016. Cospeciation of gut microbiota with hominids.
Science **353**: 380–382.

Moss EL, Maghini DG, Bhatt AS. 2020. Complete, closed bacterial genomes from microbiomes
using nanopore sequencing. *Nat Biotechnol* **38**: 701–707.

Nash AK, Auchtung TA, Wong MC, Smith DP, Gesell JR, Ross MC, Stewart CJ, Metcalf GA,
Muzny DM, Gibbs RA, et al. 2017. The gut mycobiome of the Human Microbiome
Project healthy cohort. *Microbiome* **5**: 153.

Nieves-Ramírez ME, Partida-Rodríguez O, Laforest-Lapointe I, Reynolds LA, Brown EM,
Valdez-Salazar A, Morán-Silva P, Rojas-Velázquez L, Morien E, Parfrey LW, et al.
2018. Asymptomatic Intestinal Colonization with Protist Blastocystis Is Strongly
Associated with Distinct Microbiome Ecological Patterns. *mSystems* **3**.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015. StringTie
enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol*
33: 290–295.

Piperni E, Nguyen LH, Manghi P, Kim H, Pasolli E, Andreu-Sánchez S, Arrè A, Bermingham
KM, Blanco-Míguez A, Manara S, et al. 2024. Intestinal Blastocystis is linked to
healthier diets and more favorable cardiometabolic outcomes in 56,989 individuals from
32 countries. *Cell* **187**: 4554-4570.e18.

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. *Nat Commun* **11**: 1432.

Saary P, Mitchell AL, Finn RD. 2020. Estimating the quality of eukaryotic genomes recovered
from metagenomic analysis with EukCC. *Genome Biol* **21**: 244.

Santin M, Figueiredo A, Molokin A, George NS, Köster PC, Dashti A, González-Barrio D,
Carmena D, Maloney JG. 2024. Division of Blastocystis ST10 into three new subtypes:
ST42-ST44. *J Eukaryot Microbiol* **71**: e12998.

Scanlan PD, Marchesi JR. 2008. Micro-eukaryotic diversity of the human distal gut microbiota:
qualitative assessment using culture-dependent and -independent analysis of faeces.
ISME J **2**: 1183–1193.

Scanlan PD, Stensvold CR. 2013. Blastocystis: getting to grips with our guileful guest. *Trends*
Parasitol **29**: 523–529.

Scanlan PD, Stensvold CR, Rajilić-Stojanović M, Heilig HGHJ, De Vos WM, O'Toole PW,
Cotter PD. 2014. The microbial eukaryote Blastocystis is a prevalent and diverse member
of the healthy human gut microbiota. *FEMS Microbiol Ecol* **90**: 326–330.

Schultz DT, Haddock SHD, Bredeson JV, Green RE, Simakov O, Rokhsar DS. 2023. Ancient
gene linkages support ctenophores as sister to other animals. *Nature* **618**: 110–117.

Šimková H, Câmara AS, Mascher M. 2024. Hi-C techniques: from genome assemblies to
transcription regulation. *J Exp Bot* **75**: 5357–5365.

Smit A, Hubley R, Green P. 2013. Repeatmasker Open-4.0. http://www.repeatmasker.org
(Accessed January 10, 2015).

Sommer F, Bäckhed F. 2013. The gut microbiota--masters of host development and physiology.
Nat Rev Microbiol **11**: 227–238.

Stensvold CR, Clark CG. 2020. Pre-empting Pandora's Box: Blastocystis Subtypes Revisited.
Trends Parasitol **36**: 229–232.

Stensvold CR, Suresh GK, Tan KSW, Thompson RCA, Traub RJ, Viscogliosi E, Yoshikawa H,
Clark CG. 2007. Terminology for Blastocystis subtypes--a consensus. *Trends Parasitol*
23: 93–96.

Stevens CE, Hume ID. 1995. *Comparative physiology of the vertebrate digestive system*. 2nd ed.
Cambridge University Press, Cambridge
http://catdir.loc.gov/catdir/toc/cam026/95013969.html (Accessed October 16, 2023).

Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. 2021. The Dfam community resource of
transposable element families, sequence models, and genome annotations. *Mob DNA* **12**:
2.

Tan KSW. 2008. New Insights on Classification, Identification, and Clinical Relevance of
Blastocystis spp. *Clin Microbiol Rev* **21**: 639–665.

Thorman AW, Adkins G, Conrey SC, Burrell AR, Yu Y, White B, Burke R, Haslam D, Payne
DC, Staat MA, et al. 2023. Gut Microbiome Composition and Metabolic Capacity Differ
by FUT2 Secretor Status in Exclusively Breastfed Infants. *Nutrients* **15**: 471.

Thumuluri V, Almagro Armenteros JJ, Johansen AR, Nielsen H, Winther O. 2022. DeepLoc 2.0:
multi-label subcellular localization prediction using protein language models. *Nucleic*
Acids Res **50**: W228–W234.

Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo genome assembly from
long uncorrected reads. *Genome Res* **27**: 737–746.

Wang SE, Amir AS, Nguyen T, Poole AM, Simoes-Barbosa A. 2018. Spliceosomal introns in
Trichomonas vaginalis revisited. *Parasit Vectors* **11**: 607.

West PT, Probst AJ, Grigoriev IV, Thomas BC, Banfield JF. 2018. Genome-reconstruction for
eukaryotes from complex natural microbial communities. *Genome Res* **28**: 569–580.

Wick RR, Holt KE. 2022. Polypolish: Short-read polishing of long-read bacterial genome
assemblies. *PLOS Comput Biol* **18**: e1009802.

Wilihoeft U, Campos-Góngora E, Touzni S, Bruchhaus I, Tannich E. 2001. Introns of
Entamoeba histolytica and Entamoeba dispar. *Protist* **152**: 149–156.

Yadav M, Verma MK, Chauhan NS. 2018. A review of metabolic potential of human gut
microbiome in human nutrition. *Arch Microbiol* **200**: 203–217.

Yang JK, Yoon HJ, Ahn HJ, Lee BI, Pedelacq J-D, Liong EC, Berendzen J, Laivenieks M,
Vieille C, Zeikus GJ, et al. 2004. Crystal structure of beta-D-xylosidase from
Thermoanaerobacterium saccharolyticum, a family 39 glycoside hydrolase. *J Mol Biol*
335: 155–165.

Yoshikawa H, Koyama Y, Tsuchiya E, Takami K. 2016. Blastocystis phylogeny among various
isolates from humans to insects. *Parasitol Int* **65**: 750–759.

Záhonová K, Low RS, Warren CJ, Cantoni D, Herman EK, Yiangou L, Ribeiro CA, Phanprasert
Y, Brown IR, Rueckert S, et al. 2023. Evolutionary analysis of cellular reduction and
anaerobicity in the hyper-prevalent gut microbe Blastocystis. *Curr Biol CB* **33**: 2449-
2464.e8.

Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018.
dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic*
Acids Res **46**: W95–W101.

Zierdt CH. 1988. Blastocystis hominis, a long-misunderstood intestinal parasite. *Parasitol Today*
Pers Ed **4**: 15–17.

Zierdt CH, Williams RL. 1974. Blastocystis hominis: Axenic cultivation. *Exp Parasitol* **36**: 233–
243.

Zimin AV, Marcais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA
genome assembler. *Bioinformatics* **29**: 2669–2677.

1051

1052