Horizontally transferred cell-free chromatin particles function as autonomous satellite genomes and vehicles for transposable elements within host cells

Soumita Banerjee[1,2,†], Soniya Shende[1,2,†], Laxmi Kata[1,2, †], Relestina Lopes[1,2, †], Swathika Praveen[1,2], Ruchi Joshi[1,2], Naveen Kumar Khare[1,2], Gorantla V Raghuram[1,2], Snehal Shabrish[1,2], Indraneel Mittra[1,2*]

[1]Translational Research Laboratory
Advanced Centre for Treatment, Research and Education in Cancer
Tata Memorial Centre
Kharghar, Navi Mumbai 410210, India

[2]Homi Bhabha National Institute, Anushakti Nagar, Mumbai 400094, India

*Author for correspondence: Indraneel Mittra; imittra@actrec.gov.in

[†] These authors contributed equally to the project.

Abstract

Background:

Horizontal gene transfer (HGT) plays an important evolutionary role in prokaryotes, but it is thought to be less frequent in mammals. We previously reported that cell-free chromatin particles (cfChPs) - chromosomal fragments released from the billions of dying cells that circulate in human blood - are horizontally transferred to healthy cells with biological effects. However, the underlying mechanism and function of these effects remained unclear.

Methods:

We treated NIH3T3 mouse fibroblasts cells with cfChPs isolated from human serum and serially passaged the cells. The intracellular activities of cfChPs were analysed using chromatin fibre fluorography, cytogenetic analysis, immuno-fluorescence and fluorescent *in situ* hybridisation.

Results:

We discovered that the internalised cfChPs were almost exclusively comprised of non-coding DNA, and the disparate DNA sequences contained within them had randomly combined to form complex concatemers some of which were ostensibly multi-mega base pairs in size. The concatemers exhibited variable and bizarre spatial relationships with the host cell interphase DNA with many remaining in the cytoplasm and others aligning themselves with the mouse chromosomal DNA. The concatemers autonomously performed many functions attributable to the nuclear genome. Being associated with DNA polymerase, the concatemers could synthesize DNA and autonomously replicate themselves without heed to the mitotic cycle of the host cell. They could synthesise RNA,

RNA polymerase, ribosomal RNA, ribosomal proteins, and generate numerous human proteins, including oncogenes, within the mouse cells which manifested as complex and highly amplified multi-peptide fusion proteins. The concatemers harboured human LINE-1 and *Alu* elements, which being associated with DNA polymerase, reverse transcriptase and transposes could markedly amplify themselves and increase their copy number with time in culture with the potential to rearrange themselves within the mouse genome. The above findings were reproducible in four other cell lines derived from different species suggesting that horizontal transfer of cfChPs may be a universal phenomenon.

Conclusions:

Our results lead us to propose that: 1) a cell simultaneously harbours two autonomous genome forms: one that is inherited (hereditary genome) and numerous others that are acquired (satellite genomes); 2) satellite genomes may potentially have evolutionary functions given their ability to serve as vehicles for transposable elements and to generate a plethora of novel proteins; 3) transposable elements are "foreign" genetic elements that are acquired from dying cells via HGT; 4) non-coding DNA has many hidden biological functions that remain dormant but are activated following cellular apoptosis to become detectable in association with the cfChP concatemers; 5) "within-self" HGT occurs in mammals on a massive scale via the medium of cfChP concatemers that have undergone extensive and complex modifications resulting in their behaviour as "foreign" genetic elements.

Video Abstract

https://drive.google.com/file/d/1I6NbrYmT9BRM63ywYj47AU0O9WJ_N rcR/view?usp=drive_link

Introduction

Horizontal gene transfer (HGT) plays an important role in the adaptation of microorganisms to changing environments and in evolutionary processes (Bushman, 2002). However, defining the occurrence of HGT in mammals has been a challenge. Nonetheless, experimental studies on HGT have offered insight into the regulations and functions of exogenously introduced genes and facilitated the design of drug-resistant cells, transgenic plants, and animals (Hofmann et al., 2004; Nagler et al., 2011; Puonti-Kaerlas et al., 1990). Indeed, gene therapy is founded on the principle of HGT; and several studies have indicated that oncogenes can be horizontally transferred across mammalian cells (Anker et al., 1994; Bergsmedh et al., 2001; Dvořáková et al., 2013; García-Olmo et al., 2010; Trejo-Becerril et al., 2012). These observations support the hypothesis that HGT may occur naturally in mammalian cells and contributes to evolutionary, and potentially oncogenic, processes.

Several hundred billion to trillion cells die in the human body every day (Fliedner et al., 2002; Sender and Milo, 2021) and release chromosomal fragments in the form of cell-free chromatin particles (cfChPs), which enter the extracellular compartments, including the circulation (Holdenrieder and Stieber, 2009). We previously reported that cfChPs circulating in the blood, and those that are released locally from dying cells, are readily internalised by living cells through horizontal transfer followed by their association with the host cell genome (Mittra et al., 2015, 2017). The latter was confirmed by whole-genome sequencing and fluorescence *in situ* hybridisation (FISH) analysis, as well as by the detection of multiple human *Alu* elements in the mouse recipient cells. Cellular uptake of cfChPs has several biological ramifications such as induction of DNA damage and the

activation of inflammatory and apoptotic pathways (Mittra et al., 2015, 2017). Given the

high turnover of cells, it can be assumed that all cells in the body continually internalise

horizontally transferred cfChPs throughout their lifespan, leading to somatic mosaicism,

which may be associated with ageing, chronic diseases, and cancer (Raghuram et al.,

2019). However, the mechanisms and processes governing the intracellular

consequences of horizontally transferred cfChPs remain unclear.

In this study, we investigated the intracellular processes and functions that regulate the

various biological activities of cfChPs. We selected NIH3T3 mouse fibroblast cells for our

experiments as we have extensive experience using this cell line for studies relating to

cellular transformation; and these cells have also been widely used in various studies

relating to oncogene discovery (Parada et al., 1982 & Aaronson et al., 1982). In our

previous study (Mittra et al., 2015), we investigated the relative biological activities and

functions of healthy and cancerous cfChPs. We found that cfChPs isolated from cancer

patients had significantly greater activity in terms of DNA damage and activation of

apoptotic pathways than those isolated from healthy individuals. We have also reported

that cfChPs released locally from dying cells can horizontally transfer themselves into

healthy bystander cells leading to DNA damage and inflammation (Mittra et al., 2017).

Therefore, in the present study, we treated NIH3T3 cells with cfChPs isolated from the

sera of both cancer patients and healthy individuals, as well as with those released from

hypoxia induced dying MDA-MB-231 human breast cancer cells. The cells were serially

passaged following cfChPs treatment, and using chromatin fibre fluorography,

cytogenetic analysis, and treatment with appropriate antibodies and fluorescent *in situ*

hybridisation (FISH) probes, we analysed the intracellular dynamics and processes that

regulate the functional properties of the internalised cfChPs. The results of these

experiments are presented in this article which suggest that they may help to throw new

light on mammalian evolution, ageing and cancer.

120 Results

Our results are described under three separate headings depending on whether NIH3T3

cells were treated with cfChPs isolated from sera of cancer patients or healthy individuals

or with cfChPs that were released from hypoxia induced dying MDA-MB-231 breast

cancer cells. Most of the experiments were conducted using cfChPs isolated from cancer

patients because of our especial interest in cancer, and our earlier results (Mittra et al.,

2015) which had shown that cfChPs isolated from cancer patients had significantly

greater activity in terms of DNA damage and activation of apoptotic pathways than those

isolated from healthy individuals. The majority of experiments were performed more than

once with reproducible results.

130 1. Experiments using cfChPs from cancer patients

131 1.1 cfChPs are rapidly and abundantly internalised by NIH3T3 mouse fibroblast
132 cells

We isolated cfChPs from the sera of cancer patients by a protocol described by us earlier

(Mittra et al., 2015). A representative electron microscopy image of the cfChPs is given

in Supplementary Fig. S1. In order to track the intracellular activities of cfChPs, we

fluorescently dually labelled the isolated cfChPs in their DNA with Platinum Bright Red

550 and in their histones with ATTO-488 and applied them (10ng) to NIH3T3 mouse

fibroblast cells. A representative image of isolated cfChPs that had been fluorescently

dually labelled and used to treat the NIH3T3 cells is given in Fig. 1a. Examination of

NIH3T3 mouse fibroblasts treated with the dually fluorescently labelled cfChPs confirmed

our earlier observation (Mittra et al., 2015) that cfChPs can readily and rapidly horizontally

transfer themselves to healthy cells via phagocytosis (Mittra et al., 2017) to accumulate

in their cytoplasm and nuclei by 6 h (Fig. 1b). The fluorescent cfChPs appeared

prominent, ostensibly owing to the amalgamation of multiple cfChPs to form large

concatemers following their cellular entry (described in detail later). Chromatin fibres

prepared from similarly treated NIH3T3 cells at 6h revealed the presence of numerous

dually labelled cfChPs of varying sizes, which were present in both the cytoplasm and

overlapping with mouse chromatin fibres (Fig. 1c). Since chromatin fibres are derived

from interphase cells, we further investigated whether the internalised concatemers could

also be detected in metaphase cells. The detection of concatemers on metaphase

spreads was of especial interest to us in view of the resurgence of interest in extra-

chromosomal DNA (ecDNA) in human cancers and which have largely been conducted

on metaphase chromosomes (Wu et al., 2022; Yan et al., 2024). We were interested to

examine how the cfChP concatemers might relate to the reported characteristics of

cancer ecDNAs. Metaphase spreads prepared from cfChPs-treated NIH3T3 cells at 6 h

revealed many dually labelled cfChPs, some of which were associated with the

chromosomes while others were present in extrachromosomal spaces (Fig. 1d).

Fig. 1

158
Fig. 1. Abundant uptake of cfChPs by NIH3T3 cells at 6 h. The DNA and histones of cfChPs were
dually fluorescently labelled with Platinum Bright 550 and ATTO-488 respectively and applied (10ng)
to NIH3T3 cells. **a.** Representative images of dually-labelled cfChPs prior to their application to NIH3T3
cells. **b.** Confocal microscopy images of cfChPs-treated cells at 6 h showing many dually labelled
fluorescent signals in the cytoplasm and nuclei. **c.** Fluorescence microscopy images of chromatin
fibres prepared from similarly treated cells at 6 h showing numerous dually labelled fluorescent signals

of varying sizes in the cytoplasm and in association with the chromatin fibres. **d.** Fluorescence microscopy images of metaphase spreads prepared from cfChPs-treated cells at 6 h showing multiple dually labelled fluorescent signals which are either associated with the chromosomes or are present in extrachromosomal spaces. The latter are marked with arrows.

1.2 cfChPs randomly combine to form complex concatemers

We had earlier hypothesised that when the cfChPs are horizontally transferred to another cell, the latter perceives the dsDNA breaks present in their two ends as damaged 'self' DNA, and in an attempt to repair the 'perceived damage', activates proteins of the DDR pathway which links-up multiple disparate cfChPs as a part of the repair process leading to the formation of concatemers of variable sizes (Mittra et al., 2015). To test this hypothesis, we performed FISH analysis using multiple different pairs of human chromosome-specific FISH probes on chromatin fibres and metaphase spreads prepared from cfChPs treated NIH3T3 cells that were in continuous passage. Since these chromosome specific probes were custom synthesized, we ensured that the probes that we used were human specific and did not cross-react with mouse chromosomes (Supplementary Fig. S2a and S2b). We detected frequent co-localisation of red and green fluorescent signals suggesting that unrelated chromosomal fragments containing disparate DNA sequences had randomly amalgamated with each other to form highly complex concatemers (Fig. 2). Some of the concatemers appeared strikingly prominent, which is particularly evident from images presented in Figure 2a. This suggested that the components of the concatemer had been markedly amplified. Even the small fluorescent signals shown in Figure 2b appeared as concatemers. As the analytical resolution of a FISH signal under a fluorescent microscope is in the range of 100–200 Kb (Cui et al., 2016) even the small concatemers can be assumed to be at least of a similar size range. Surprisingly, we detected the fluorescence signals of chromosome 4 to co-localise with

those of centromeres and of chromosome 20 to co-localise with those of telomeres. This

finding highlighted the extent of genetic complexity and chaotic nature of the concatemers

(Fig. 2a, panels 4&5). Perhaps the ultimate evidence of the chaotic composition of the

concatemers comes from our detection of co-localising signals of telomeres and

centromeres (Fig. 2a, panel 6). The centromeric signal seen in Fig. 2a, panel 4 is strikingly

prominent, suggesting that the concatemers had not only incorporated centromeric DNA

sequences (171bp in size) within their folds but that they had undergone large-scale

amplification. Results of quantitative analysis of the degree of co-localisation of

fluorescent signals generated upon treatment with FISH probes specific to different

chromosome pairs are given in Supplementary Fig. S3a. The extent of co-localization

ranged between 32.4% and 39.8%. Concatemerisation could also be detected in

metaphase spreads prepared from serially passaged cfChPs-treated NIH3T3 cells (Fig.

2c). Negative control experiments performed on native NIH3T3 cells that had not been

exposed to cfChPs treatment did not react with human specific FISH probes against

chromosome 4 and chromosome 16 which were tested (Supplementary Fig. S4a).

Fig. 2

Scale bar = 20 μm

Scale bar = 20 μm

Fig. 2. Internalised cfChPs combine to form complex concatemers. a. FISH analysis of chromatin fibres prepared from cfChPs-treated NIH3T3 cells in continuous passage using different combinations of human chromosome-specific FISH probes, including probes specific for the human centromere and telomere, revealing co-localised fluorescent signals (arrows). **b.** Small cfChPs also show co-localised signals suggesting that they too are comprised of concatemers **c.** Similar co-localising signals are shown on metaphase spreads. Co-localised signals are marked with arrows.

1.3 Concatemers exhibit variable spatial relationships with mouse chromatin fibres

We next undertook immuno-FISH analysis using an antibody against histone H4 and a human whole genomic FISH probe on chromatin fibres prepared from cfChPs-treated NIH3T3 cells in continuous passage. We detected numerous dually labelled signals representing concatemers which exhibited extensive structural and size variability, as well as remarkably variable spatial relationships with mouse chromatin fibres (Fig. 3a). Many concatemers were detected in the cytoplasm or to have aligned with long stretches of mouse DNA, while others exhibited unusual conformations or were seen to be dangling from the mouse DNA. A similar peculiar conformation was found in clone D5 which had been developed several years earlier by treatment of NIH3T3 cells with cfChPs isolated from sera of cancer patients and which had undergone numerous passages and multiple freeze–thaw cycles (Mittra et al., 2015) (Fig. 3b). This finding suggested that concatemers persist across cell generations showing remarkable stability. The fluorescent signals of the concatemers were also strikingly prominent, suggesting that they greatly exceeded the threshold of detection of FISH signals (100–200 kb) (Cui et al., 2016). Control experiments were performed to confirm that the whole genomic DNA FISH probe used was human specific and did not cross-react with mouse (Supplementary Fig. S13).

Fig. 3

Fig. 3. Variable spatial relationships of concatemers with mouse chromatin fibres. a. Immuno-FISH analysis of chromatin fibres prepared from cfChPs-treated NIH3T3 cells in continuous passages using an antibody against histone H4 and a human whole-genome FISH probe. **b.** Immuno-FISH analysis of chromatin fibres prepared from clone D5 showing that the concatemers persisted even after numerous passages and freeze–thaw cycles.

1.4 Concatemers synthesise DNA polymerase and can replicate

Cells in continuous passage were pulse-labelled with BrdU followed by immuno-FISH analysis using an antibody against BrdU and a human genomic DNA probe. The analysis revealed co-localisation of the fluorescent signals of BrdU and DNA, indicating that the concatemers were actively synthesising DNA (Fig. 4a, upper panel). Immuno-FISH

experiments further revealed that the concatemers expressed human DNA polymerase γ

(Fig. 4a, middle panel), while dual immunofluorescence staining using antibodies against

BrdU and human specific DNA polymerase γ revealed co-localised signals, indicating that

the concatemers had the potential to autonomously replicate themselves (Fig. 4a, lower

panel). Since human DNA polymerases is well conserved in mouse and across mammals,

we ensured that the antibodies against DNA Polymerase γ was human specific and did

not cross-react with mouse (Supplementary Fig. S5). Results of quantitative analysis of

the degree of co-localisation of fluorescent signals of human DNA and BrdU; human DNA

and human DNA polymerase γ; and BrdU and human DNA polymerase γ are given in

Supplementary Fig. S3b. The extent of co-localization ranged between 82.0% and 88.8%.

These findings were confirmed on metaphase preparations from serially passaged cfChP-

treated NIH3T3 cells wherein we detected multiple co-localising signals of human DNA

and BrdU, human DNA and human DNA polymerase γ and BrdU and human DNA

polymerase γ in the extrachromosomal spaces (Fig. 4b). Negative control experiments

performed on native NIH3T3 cells that had not been exposed to cfChPs treatment did not

react with either to the human genomic DNA probe or the antibody against human DNA

polymerase γ (Supplementary Fig. S4b).

Fig. 4

a

Hoechst	Human DNA	BrdU	Merged
Hoechst	Human DNA	h-DNA Polymerase γ	Merged
Hoechst	BrdU	h-DNA Polymerase γ	Merged

Scale bar = 20 µm

b

DAPI	Human DNA	BrdU	Merged
DAPI	Human DNA	h-DNA Polymerase-γ	Merged
DAPI	BrdU	h-DNA Polymerase γ	Merged

Scale bar=10µm

Fig. 4. The concatemers synthesise DNA and express human DNA polymerase γ. **a.** Chromatin fibres were prepared from NIH3T3 cells in continuous passage were pulse-labelled for 24 h with BrdU (10µm). Immuno-FISH experiments using antibodies against BrdU, human-specific DNA polymerase γ, and a human whole-genome FISH probe show co-localised signals of human DNA and BrdU, human DNA and DNA polymerase γ, and BrdU and DNA polymerase γ. Co-localised signals are marked with arrows. **b.** Similar co-localising signals are shown on metaphase spreads. Co-localised signals are marked with arrows.

Given the above results that the concatemers expressed DNA polymerase which co-localised with BrdU signals, we investigated their potential for self-replication. Ten metaphase spreads were prepared from cells in each successive passage and the number of human FISH signals (representing concatemers) per chromosome was determined. The result showed that human FISH signals on mouse chromosomes increased progressively such that the number of signals at passage 198 was 4.07 times higher than those in passage 2 (Fig. 5a). It should be noted that this analysis is restricted to the proliferative capacity of concatemers that were associated with the chromosomes and did not take into account the replicative potential of those that are present in the cytoplasm. We also did a similar exercise to investigate whether the concatemers could amplify themselves with time by estimating the mean fluorescent intensity (MFI) per chromosome. We found that the increase in MFI between passage 2 and 198 was 237.2 fold (Fig. 5b). Taken together, these results indicated that the concatemers increased their copy number and extensively amplified themselves with time in culture.

Fig. 5

a

b

288 **Fig. 5. Concatemers proliferate and amplify themselves within the mouse genome over time.**
289 Metaphase spreads were prepared from cells in each progressively increasing passage and probed
290 with a human-specific DNA FISH probe. **a.** The total number of human DNA FISH signals on the DAPI

stained chromosomes was counted, and the mean number of human FISH signals per chromosome was calculated after analysing 10 metaphase spreads at each passage. A progressive increase in mean FISH signals per chromosomes is evident with increasing passage number (analysis of variance for linear trend p < 0.0001). The mean number of human FISH signals per chromosome between passage no. 2 and passage no. 198 increased by a factor of 4.07. **b.** a similar exercise was done as above except that mean fluorescent intensity (MFI) of human FISH signals per chromosome, indicative of amplification, was determined. A progressive increase in mean MFI per chromosomes is evident with increasing passage number (analysis of variance for linear trend p < 0.0001). The MFI per chromosome increased by a factor of 237.19 between passage no. 2 and passage no. 198. The insets represent partial metaphase images showing human DNA signals on the chromosomes. Blue and red signals represent DAPI and human DNA, respectively.

1.5 Concatemers are composed of open chromatin

We next examined the epigenetic constitution of the concatemers using antibodies against the histone markers representing trimethylation of histone H3 at lysine 4 (H3K4me3) and trimethylation of histone H3 at lysine 9 (H3K9me3), indicative of regions associated with active gene promoters and regions associated with long-term repression respectively (Barski et al., 2007). Chromatin fibres prepared from cfChPs-treated NIH3T3 cells in continuous passage were dually immune-stained with antibodies targeting H3K4me3 and H3K9me3, and the number of concatemers that reacted with the antibodies was counted. We found that the vast majority of the concatemers either reacted exclusively with antibodies against H3K4me3 (open chromatin) or were hybrids of H3K4me3 and H3K9me3, with only a small fraction of reacting exclusively with H3K9me3 (Fig. 6). The figure also shows that the concatemers contained open chromatin irrespective of whether the regions of the host mouse DNA reacted with antibodies against H3K4me3 or H3K9me3. Taken together, these data indicated that concatemers primarily contained nucleosome-depleted regions and could bind to protein factors that facilitate gene transcription (Thomas et al., 2011) and DNA replication (MacAlpine et al., 2010).

Fig. 6

Fig. 6. Concatemers largely comprise of open chromatin irrespective of the epigenetic constitution of the mouse DNA. Representative images of chromatin fibres immunostained with antibodies against H3k4me3 (red) and H3k9me3 (green). The hybrid concatemers are represented by co-localised yellow signals. The host mouse DNA in the upper image is seen to react with H3k4me3 antibody representing open chromatin while the host mouse DNA segment in the lower image is comprised of heterochromatin and seen to react with antibody against H3k9me3. Histograms represent quantitative estimates of H3k4me3, H3k9me3, and hybrid histones after counting 100 fluorescent signals. The values are expressed as mean ± SEM values. Statistical analysis was performed using two-tailed Student's t-test. $* p < 0.05$, $** p < 0.01$ and $**** p < 0.0001$.

1.6 Concatemers can synthesise RNA

As the above findings suggested that concatemers are largely composed of open
chromatin and potentially capable of active transcription, we investigated their potential
for RNA synthesis. Using an assay kit which detects global RNA transcription, we
detected abundant RNA in the cytoplasm of cfChP-treated passaged cells, which was
absent in the control NIH3T3 cells (Fig. 7). As RNA synthesis is normally restricted to the
nucleus, the detection of RNA in the cytoplasm indicated that DNA contained within the
concatemers were undergoing active transcription. Treatment of the cfChPs treated
passaged cells with actinomycin D or maintaining the cells at low temperature (31°C)
abolished RNA synthesis. These data indicated that the concatemers were actively
involved in RNA synthesis which is dependent on active cellular metabolism.

Fig. 7

Negative Control **NIH3T3 Control** **cfChPs treated cells**

cfChPs treated cells +
Actinomycin-D

cfChPs treated cells at
31°C

Fig. 7. Concatemers synthesise RNA which is dependent on active cellular metabolism. Images
showing cytoplasmic RNA synthesis by cfChP-treated cells in continuous passage which is absent in
control NIH3T3 cells. Treatment of cfChP-treated cells with Actinomycin D (0.0005 µg/mL) or
maintenance at low temperature (31°C) abolishes RNA synthesis indicating that the latter is dependent
on active cellular metabolism.

1.7 Concatemers synthesise their own protein synthetic machinery

The ability of the concatemers to synthesise RNA led us to investigate whether they were

involved in protein synthesis. We looked for three critical components of protein synthetic

machinery viz. ribosomal RNA, RNA polymerase and ribosomal protein. For detection of

ribosomal RNA we undertook dual-FISH using a human genomic FISH probe (to detect

the concatemers) and a FISH probe against human ribosomal RNA. We ensured that the

latter FISH probe was specific to human and did not cross-react with mouse

(Supplementary Fig. S6, upper panel). Dual FISH analysis on chromatin fibres prepared

from serially passaged cfChP-treated cells revealed strictly co-localised signals,

indicating that the concatemers were synthesising ribosomal RNA (Fig. 8a, upper panel).

We next investigated whether the concatemers could synthesise the other two

components of the protein synthetic machinery viz. RNA polymerase and ribosomal

protein after ensuring that the antibodies against them were specific to humans

(Supplementary Fig. S6, middle and lower panels). Immuno-FISH analysis revealed co-

localised fluorescent signals generated by a human-specific genomic DNA probe and

antibodies against RNA polymerase III and ribosomal protein (Fig. 8a, middle and lower

panels, respectively). These findings indicated that concatemers could autonomously

synthesise critical components of the protein synthetic machinery. Results of quantitative

analysis of the degree of co-localisation of fluorescent signals of human DNA and human

ribosomal RNA; human DNA and RNA polymerase III; and human DNA and ribosomal

protein are given in Supplementary Fig. S3c. The extent of co-localization ranged

between 77.2% and 81.4%. We confirmed these findings in metaphase preparations from

serially passaged cells for all three components of the protein synthetic machinery

mentioned above (Fig. 8b). Negative control experiments performed on native NIH3T3

cells that had not been exposed to cfChPs treatment did not react with the human specific

probes against DNA, ribosomal RNA and the human specific antibodies against RNA

polymerase III and ribosomal protein (Supplementary Fig. S4c).

Fig. 8

Scale bar = 20 µm

Scale bar = 10 µm

Fig. 8. Concatemers synthesise their own protein synthetic machinery. Chromatin fibres **(a)** and
metaphase spreads **(b)** were prepared from cfChPs treated cells in continuous passage. Dual-FISH
experiments were performed using a human specific genomic DNA probe and a probe against human
specific ribosomal RNA, and immune-FISH experiments using a human specific genomic DNA probe

384 and antibodies against human specific RNA polymerase III and human specific ribosomal protein. Co-
385 localised fluorescent signals of DNA and the components of the above protein synthetic machinery
386 are marked with arrows.

387
388
1.8 Concatemers synthesise a variety of human proteins in mouse cells

Having confirmed that concatemers are capable of synthesising RNA and assembling

their own protein synthetic machinery, we went on to investigate whether they were

capable of autonomously synthesising proteins. We conducted immune-FISH

experiments using a human specific genomic DNA probe (to detect the concatemers) and

antibodies against various proteins. We found that the fluorescent signals of the various

proteins frequently co-localized with those of human DNA suggesting that the

concatemers were capable of synthesizing proteins (Fig. 9a). Significantly, the newly

synthesised proteins consistently remained associated with the concatemers of their

origin (identified by fluorescent human DNA signals). This finding suggested that,

although the concatemers contained the critical components of a protein synthetic

machinery, they apparently lacked the machinery required for protein sorting. It should be

noted that all the proteins that we detected seemed to be over-expressed, confirming that

the gene segments corresponding to the proteins within the concatemers were amplified.

Results of quantitative analysis of the degree of co-localisation of fluorescent signals of

human DNA and of various human proteins are given in Supplementary Fig. S3d. The

extent of co-localization varied between 70.0% and 72.6%. We further confirmed these

findings in metaphase preparations from serially passaged cfChPs-treated NIH3T3 cells

(Fig. 9b). We also conducted extensive control experiments on native NIH3T3 cells that

had not been exposed to cfChPs treatment using a variety of human specific antibodies

and found that none of them showed any positive fluorescent signals (Supplementary Fig.

S4d). The above results indicated that the disparate DNA sequences that comprise the

concatemers are transcribed and translated to generate proteins corresponding to the

diverse DNA sequences that they contain apparently using their own protein synthetic

machinery. Results of control experiments confirming the human specificity of the

antibodies against all the proteins described above are given in Supplementary Fig. S7.

Fig. 9

a

Scale bar = 20 μm

b

Scale bar = 10 μm

Fig. 9. Concatemers synthesise a variety of human proteins. Immuno-FISH analysis using antibodies against various human proteins and a human whole genome FISH probe on chromatin fibres **(a)** and metaphase spreads **(b)** prepared from cfChPs treated cells in continuous passage show co-localizing signals. The co-localizing signals of human DNA and various proteins are marked with arrows.

1.9 The proteins that concatemers synthesize are complex fusion proteins

As the concatemers are formed as a result of amalgamation of widely disparate DNA sequences, we investigated whether the proteins that they synthesized might be fusion proteins. Dual-immunofluorescence experiments using pairs of antibodies against proteins, the corresponding genes of which were located on different chromosomes, revealed that the fluorescent signals frequently co-localised. This finding indicated that the proteins synthesised by the concatemers were fusion proteins with potentially novel functions (Fig. 10). Fusion proteins were detected both on chromatin fibres and metaphase preparations. Results of quantitative analysis of the degree of co-localisation of fluorescent signals indicative of fusion proteins are shown in Figure 10a. The extent of co-localising signals ranged between 33.0% and 36.3% (Supplementary Fig. S3e). Results of control experiments confirming the human specificity of the antibodies against the fusion proteins not included in Supplementary Fig. S7 are given in Supplementary Fig. S8.

Fig. 10

a

Scale bar = 20 µm

b

Scale bar = 10 µm

Fig. 10. Proteins synthesised by concatemers are fusion proteins. Dual immunofluorescence analysis using antibody pairs targeting diverse human-specific proteins in chromatin fibres **(a)** and metaphase spreads **(b)** prepared from cfChPs-treated NIH3T3 cells in continuous passage. Results show frequent co-localised signals indicative of fusion proteins (arrows). The numbers given in the parenthesis indicate the chromosomal location of the genes that correspond to the proteins.

1.10 Concatemers are vehicles for transposable elements

Among the fusion proteins shown in Figure 10 we accidently found reverse transcriptase

and transposase to co-localise with Bcl2 and cMyc, respectively. This finding raised the

possibility that the concatemers might harbour gene components related to transposable

elements. We used human specific LINE-1 and *Alu* DNA probes which had been custom

synthesised (Supplementary Table S2). Nonetheless, we checked for their specificity to

ensure that they reacted only to human and not to mouse transposable elements

(Supplementary Fig. S9a&b). Dual-FISH experiments using a human genomic DNA probe

and those targeting human LINE-1 and *Alu* showed that many of the fluorescent signals

had co-localised indicating that the concatemers harboured DNA sequences of retro-

transposable elements (Fig. 11a). Results of quantitative analysis of the degree of co-

localisation of fluorescent FISH signals of human DNA and human LINE-1 and Alu on

chromatin fibres are given in Supplementary Fig. S3f. The extent of co-localization was

70.4% and 72.8% for LINE-1 and Alu respectively. Similar co-localizing signals were also

seen on metaphase preparations (Fig. 11b). The fact that both LINE-1 and *Alu* signals

could be clearly detected by FISH analysis indicated that they were extensively amplified,

given that LINE-1 and Alu are approximately 6000 bp and 300 bp in size, respectively.

The issue of amplification of the transposable elements is discussed in detail under

section 1.12. Control experiments performed on native NIH3T3 cells that had not been

exposed to cfChPs treatment using human LINE-1 and Alu probes did not reveal any

positive signals (Supplementary Fig. S4e).

Fig. 11

a

Scale bar = 20 μm

b

Scale bar = 20 μm

Fig. 11. Concatemers harbour transposable elements. Dual-FISH analysis on chromatin fibres (**a)** and metaphase spreads **(b)** prepared from cfChPs-treated NIH3T3 cells in continuous passage using a human specific genomic DNA probe and those against human LINE-1 or human *Alu* show co-localised signals (marked with arrows).

1.11 LINE-1 and *Alu* elements are associated with reverse transcriptase, transposase and DNA polymerase

Immuno-FISH analysis using antibodies against human reverse transcriptase and transposase and FISH probes against LINE-1 and *Alu* revealed co-localised signals, indicating that the enzymes reverse transcriptase and transposase were frequently

associated with DNA sequences corresponding to LINE-1 and Alu (Fig. 12a and b). Such

an association could potentially allow the transposable elements to re-arrange

themselves on the mouse cell genome by translocating themselves from one location to

another. Results of quantitative analysis of the degree of co-localisation on chromatin

fibres of fluorescent FISH signals of human LINE-1 and human reverse transcriptase and

transposase; and Alu and human reverse transcriptase and transposase are given in

Supplementary Fig. S3g. The extent of co-localization ranged between 34.4% and 36.1%.

Control experiments performed on native NIH3T3 cells that had not been exposed to

cfChPs treatment using human specific antibodies against reverse transcriptase and

transposase and FISH probes against h-LINE-1 and h-ALU did not reveal any positive

signals (Supplementary Fig. S4f).

Fig. 12

Fig. 12. LINE-1 and Alu sequences are associated with reverse transcriptase and transposase. Immuno-FISH analysis of chromatin fibres **(a)** and metaphase spreads**(b)** prepared from cfChPs-treated NIH3T3 cells in continuous passage using human *Alu* or LINE-1 probes and antibodies against human reverse transcriptase and human transposase show co-localising signals (marked with arrows).

As transposable elements are known to increase their copy number with time (Li et al., 2013), we were curious to find out whether the LINE-1 and *Alu* elements would proliferate and increase their copy number with progressively increasing passages. Immuno-FISH

experiments using antibodies against human DNA polymerase γ and FISH probes against

LINE-1 and *Alu* detected co-localised signals, indicating that LINE-1 and *Alu* elements

were associated with DNA polymerase, raising the possibility that they may have the

potential to proliferate (Fig. 13a and b). This possibility was supported by the finding that

LINE-1 and *Alu* elements could synthesize DNA. Immuno-FISH analysis of cells in

continuous passage that had been pulse-labelled with BrdU showed co-localised signals

generated by antibodies against BrdU and FISH probes against LINE-1 and *Alu*. Taken

together these findings supported the idea that the DNA of both the transposable

elements had the potential to replicate. Results of quantitative analysis of the degree of

co-localisation of fluorescent FISH signals of human LINE-1 and human DNA polymerase

and BrdU; and Alu and human DNA polymerase and BrdU are given in Supplementary

Fig. S3h. The extent of co-localization varied between 35.8% and 36.9%.

Fig. 13

Scale bar = 10 μm

Fig. 13. LINE-1 and *Alu* elements are associated with DNA polymerase and can actively synthesise DNA. (a) Immuno-FISH images of cfChPs-treated cells in continuous passage showing co-localising signals of LINE-1 (green) and Alu (green) and DNA polymerase γ (orange) and BrdU (red) on chromatin fibres. **(b)** Similar co-localising signals are shown on metaphase spreads. Co-localised signals are marked with arrows.

1.12 LINE-1 and *Alu* elements increase their copy number and amplify themselves with time

Given the above findings, we were curious to find out whether LINE-1 and *Alu* elements could indeed replicate and increase their copy numbers within the mouse genome.

Metaphase spreads were prepared from cfChPs treated NIH3T3 cells in continuous

passage and probed with FISH probes against LINE-1 and *Alu* elements. Fifteen

metaphases were analysed at each passage and the number of LINE-1 and *Alu*

fluorescent signals per metaphase were determined (Fig. 14). Our results showed that

the numbers of human LINE-1 and *Alu* signals per chromosome increased steadily

between passage 2 and 198 resulting in a 7.6-fold increase in copy number in case of

LINE-1 and a 6.7-fold increase in case of *Alu* (Fig. 14a). These data indicated that LINE-

1 and *Alu* elements progressively increased their copy numbers by retrotransposition over

time in the mouse genome. It should be noted that this analysis is restricted to the

proliferative capacity of LINE-1 and *Alu* elements that are associated with chromosomes

and does not take into account the replicative potential of those elements that are present

in the cytoplasm. We also did a similar exercise to investigate whether the concatemers

could amplify themselves by estimating the mean fluorescent intensity (MFI) of LINE-1

and *Alu* signals per chromosome. We found that the increase in MFI between passage 2

and 198 was 151.3 fold in case of LINE-1 and 83.4 fold in case of *Alu* (Fig. 14b). Taken

together these data indicated that the transposable elements could not only proliferate

but also extensively amplify themselves thereby becoming increasingly effective in

modifying the host genome.

Fig. 14

a

LINE -1

Plot: +ve signals per chromosome (y-axis, 0.0 to 1.0) vs Passage no. (x-axis: 2, 11, 30, 60, 90, 120, 150, 180, 198)

Alu

Plot: +ve signals per chromosome (y-axis, 0.0 to 1.0) vs Passage no. (x-axis: 2, 11, 30, 60, 90, 120, 150, 180, 198)

b

LINE -1

Plot: MFI per chromosome(a.u) (y-axis, 100 to 1000000) vs Passage no. (x-axis: 2, 11, 30, 60, 90, 120, 150, 180, 198). Insets labelled P-2 and P-198, Scale bar=10µm

Alu

Plot: MFI per chromosome(a.u) (y-axis, 100 to 100000) vs Passage no. (x-axis: 2, 11, 30, 60, 90, 120, 150, 180, 198). Insets labelled P-2 and P-198, Scale bar=10µm

Fig. 14. LINE-1 and *Alu* elements proliferate and amplify themselves within the mouse genome over time. Metaphase spreads were prepared from cells in each progressively increasing passage and probed with a human-specific LINE-1 and *Alu* probes. **a.** The total number of human LINE-1 and *Alu* signals on the DAPI stained chromosomes were counted, and the mean number of LINE-1 and *Alu* signals per chromosome was calculated after analysing 15 metaphase spreads at each passage. A progressively increasing LINE-1 and *Alu* signals per chromosomes is evident with increasing passage number (analysis of variance for linear trend p < 0.0001). The number of human LINE-1 signals per chromosome increased by a factor of 7.6 between passage no. 2 and passage no. 198, and by a factor of 6.7 in case of *Alu.* **b.** a similar exercise was done as above except that mean fluorescent intensity (MFI) of human LINE-1 and *Alu* per chromosome, indicative of amplification, was determined. A progressively increasing in MFI per chromosomes is evident with increasing passage number (analysis of variance for linear trend p < 0.0001). Mean MFI per chromosome increased by a factor of 151.3 between passage no. 2 and passage no. 198 in case of LINE-1, and by a factor of 83.4

in case of *Alu.* The insets represent partial metaphase images showing human LINE-1 and Alu signals on the chromosomes. Blue and red signals represent DAPI and human DNA, respectively.

1.13 NIH3T3 cells are not unique in their ability to internalise cfChPs

We next investigated whether the ability to internalise cfChPs is a property that is unique

to NIH3T3 cells or whether other cells from different species also have similar ability to

internalise cfChPs. We treated 4 different cell lines namely Vero (monkey kidney cells;

ATCC Vero-CCL-81); Dolly (dog cells; spontaneous malignant round cell tumour in a dog

developed at our institute's animal facility); B/CMBA. OV (mouse ovary cells; ATCC RRID:

CVCL_5926) and HEK293 (human embryonic kidney cells; ATCC CRL-1573™) with

cfChPs isolated from human serum (10ng). The treated cells were harvested after the 5th

passage, and chromatin fibres prepared from them were probed with a human whole

genomic probe and a probe against human LINE-1. We detected the presence of human

DNA and human LINE-1 signals in all the above cell lines indicating that all of them had

the ability to internalise cfChPs (Supplementary Fig. S14a-d). This finding led us to the

conclusion that horizontal transfer of cfChPs is likely to be a universal phenomenon.

1.14 The concatemers are largely composed of non-coding DNA

Given that 99% of the human genome is comprised of non-coding DNA, and in view of

our detection of LINE-1 and Alu elements among the cfChP concatemers, we investigated

whether the majority of the cfChPs that had been internalised might have been derived

from non-coding DNA? To investigate this, we used a human long non-coding RNA probe

which comprised of a set of Stellaris RNA FISH probes containing a pool of up to 48

unique sequences, each labelled with a fluorophore, that collectively bind along an RNA

target transcript (Supplementary Table S2). When we applied this RNA probe to

chromatin fibres prepared from two human cell lines (HEK293 and MRC5), we found that

the RNA probe aligned with the entire length of the human DNA with almost 100%

coverage (Fig. 15a). The human specificity of the RNA probe was confirmed by the

demonstration that it did not bind to NIH3T3 mouse DNA (Fig. 15a). This finding

suggested that the long non-coding RNA probe could be used as a substitute to detect

human non-coding DNA. We, therefore, used the RNA probe to detect the presence of

non-coding human DNA in NIH3T3 cells treated with cfChPs isolated from human serum

(Fig. 15b). We discovered that the internalised cfChPs concatemers reacted with probes

against both human whole genomic DNA and human non-coding DNA and their

fluorescent signals co-localised to the extent of 98.2% (Fig. 15b). This finding was

confirmed on metaphase preparations of the cfChPs treated cells (Fig. 15c). This led us

to the conclusion that virtually all the cfChPs concatemers were composed of non-coding

DNA.

Fig. 15

a

	DAPI	Human DNA	Human Non-coding RNA	Merged
HEK 293 cells (Human)				
MRC5 cells (Human)				
NIH3T3 cells (Mouse)				

Scale bar = 10um

b

	DAPI	Human DNA	Human Non-coding DNA	Merged

Scale bar = 10um

% Co-localization

98.2

Human DNA n = 1000

c

DAPI	Human DNA	Human Non-coding DNA	Merged

Scale bar = 10um

Fig. 15. The concatemers are largely composed of non-coding DNA. a. The human long non-
coding RNA probe aligns with human DNA but not with mouse DNA. Chromatin fibres were prepared
from two human cell lines HEK293 and MRC5 and hybridised with long non-coding RNA probe which
shows that the probe has almost 100% coverage. **b.** The concatemers are largely composed of non-
coding DNA. cfChPs treated passaged cells were probed with a fluorescently labelled human whole
genomic as well as the human long non-coding RNA probe (which substituted for a non-coding DNA

probe). The degree of co-localisation of fluorescent signals is represented as a histogram, which was generated after counting 1000 human DNA signals, and which shows that the extent of co-localisation of the fluorescent signals was of the order of 98.2%. **c.** Confirmation of these findings on metaphase spreads.

1.15 Biological activities of concatemers are largely attributable to non-coding DNA

The above finding that the concatemers are largely composed of non-coding DNA led us

to investigate whether the biological activities of the concatemers that we described

earlier were attributable to non-coding DNA. We discovered this to be indeed the case.

We found that the fluorescent signals of a selected set of target enzymes viz. DNA

polymerase, RNA polymerase and reverse transcriptase co-localised with those of non-

coding DNA, as did BrdU signals representing DNA synthesis (Fig. 16). As anticipated,

the fluorescence signals of LINE-1 and Alu also co-localised with those of non-coding

DNA (Fig. 16). The degree of co-localisation of fluorescent signals of non-coding DNA

and those of the above biomarkers is given in Supplementary Fig. S15. The latter also

shows that the degree of co-localization with the biomarkers was similar irrespective of

whether we used a whole human genomic probe (results of which are shown earlier) or

the one against non-coding DNA. Collectively, these findings provided confirmation that

the biological activities of the concatemers are attributable to non-coding DNA, and

suggested that non-coding DNA has many unique biological functions which remain

dormant but are activated following cellular apoptosis to become detectable in association

with the cfChP cocatemers.

Fig. 16

Scale bar = 10um

Fig. 16. Biological activities of the concatemers are attributable to non-coding DNA. Immuno-FISH and dual-FISH analysis of chromatin fibres prepared from cfChPs treated cells in continuous passage. The chromatin fibres were probed with antibodies against various human proteins and a human non-coding DNA probe or FISH probes against LINE-1 and Alu and that against human non-coding DNA. Co-localisation of various signals is clearly seen. These data suggested that biological activities of the concatemers are attributable to non-coding DNA.

2. Experiments using cfChPs isolated from healthy individuals

The experiments described above were performed with NIH3T3 cells that had been

treated with cfChPs isolated from the sera of cancer patients and passaged continuously.

We simultaneously performed similar experiments with NIH3T3 cells that had been

treated with cfChPs isolated from the sera of healthy individuals. We found that cfChPs

from healthy individuals were also readily internalised by NIH3T3 cells wherein they

behaved in a manner similar to that of cfChPs derived from the cancer patients

(Supplementary Fig. S10).

642 3. Experiments using cfChPs released from hypoxia-induced dying
643   MDA-MB-231 cells

Finally, given our previous finding that cfChPs spontaneously released from dying cancer

cells were readily internalised by healthy cells (Mittra et al., 2017), we further investigated

the intracellular activities of cfChPs released from dying MDA-MB-231 breast cancer

cells. Hypoxia-induced cfChPs were collected in 1 ml of culture medium and added to

NIH3T3 cells (please see Materials and Methods). We observed that cfChPs that were

released from hypoxia-induced dying MDA-MB-231 cells could reciprocate all the

intracellular activities and functions that were observed using cfChPs isolated from serum

of both cancer patients and healthy individuals (Supplementary Fig. S11).

653 Discussion

654 The present study is founded on our earlier reports that cfChPs released from the billions

655 of cells that die in the body everyday can horizontally transfer themselves to healthy cells

656 resulting in activation of several biological processes (Mittra et al., 2015, 2017). Herein,

657 we have reported multiple additional activities and biological processes which can be

summarised as follows. The internalised cfChPs containing widely disparate DNA sequences randomly combined to form complex concatemers of variable sizes many of which were ostensibly multi-mega base pairs in size. The concatemers exhibited variable and bizarre spatial relationships with the host cell interphase DNA with many remaining in the cytoplasm and others being directly associated with the mouse chromosomal DNA. The concatemers once formed persisted across cell generations and showed remarkable stability. They exhibited the potential to perform many functions that are normally attributed to the nuclear genome. They could generate DNA polymerase which allowed them to synthesize DNA and independently replicate themselves without heed to the mitotic cycle of the host cell. The concatemers synthesised RNA as well as RNA polymerase, ribosomal RNA, ribosomal proteins, and numerous other human proteins in the mouse cells. The latter manifested as complex multi-peptide fusion proteins which were often highly overexpressed. Further analysis demonstrated that the concatemers harboured human LINE-1 and *Alu* retro-transposable elements which were associated with the enzyme DNA polymerase. The latter allowed them to increase their copy numbers and extensively amplify themselves with time in culture. The LINE-1 and Alu elements were also associated with reverse transcriptase and transposase, suggesting that they had the potential to rearrange themselves within the mouse genome leading to extensive modifications. Taken together, the above findings lead us to conclude that the concatemers that form following horizontal transfer of cfChPs act as autonomous satellite genomes which function independently of the nuclear genome. Being associated with transposable elements they have the potential to modify the host genome in unpredictable ways leading to somatic mosaicism - the hallmark of ageing and cancer (Vijg, 2014). Finally, the above findings could be reproduced in four other cell lines derived from

different species suggesting that horizontal transfer of cfChPs may be a universal

phenomenon.

Some of the summary points mentioned above call for further elaboration. We discovered

that the concatemers were primarily composed of non-coding DNA. The latter is not

normally known to perform the diverse biological activities that we detected to be

associated with the concatemers. This suggested that non-coding DNA has many hidden

biological functions which remain dormant but are activated following cellular apoptosis

to become detectable in association with the cfChP cocatemers. Our demonstration that

the concatemers synthesise ribosomal RNA, RNA polymerase and ribosomal protein

suggests that they are capable of creating their own autonomous protein synthetic

machinery without recourse to that of the host cell. The concatemers being composed of

open chromatin are apparently able to indiscriminately transcribe their DNA via RNA

polymerase; and the RNAs thus generated are translated into proteins with the help of

their own ribosomal RNA and ribosomal proteins. Significantly, the newly synthesised

proteins remained strictly associated with their concatemers of origin suggesting that,

although the concatemers contained the critical components of a protein synthetic

machinery, they lacked the machinery required for protein sorting. Taken together, these

findings suggest a novel mechanism of protein synthesis which function autonomously

without recourse to the host cell's own protein synthetic machinery. Clearly, such an

assertion, which defies existing knowledge, requires critical experimental scrutiny to

define the mechanism(s) that underlie such a unique biological process.

Our detection of the unusual intracellular characteristics and functions of cfChPs and their

concatemers was facilitated by several factors. First, our trans-species model comprising

of mouse recipient cells and cfChPs derived from human serum. This allowed us to track

the activities of internalised human cfChP concatemers within the mouse cells using

appropriate human-specific FISH probes and antibodies. Second, the chromatin fibre

technique facilitated the detection of spatial relationships between internalised cfChPs

concatemers and the mouse interphase DNA. Third, our use of human specific DNA FISH

probes and antibodies allowed the detection of short DNA sequences within the

concatemers and small peptides that they synthesized by binding to the corresponding

DNA sequences and peptide epitopes respectively. These would not have been readily

detectable by the standard techniques of molecular biology (discussed subsequently in

the manuscript). Finally, our discovery that the non-coding RNA probe could be

substituted to detect non-coding DNA allowed us to confirm that the concatemers were

primarily composed of non-coding DNA and that they(could perform many of the

functions that are normally attributed to the genomic DNA.

Much research has been devoted to transposable elements since the seminal discovery

of Barbara McClintock of "jumping genes" in maize. The sequencing of the human

genome has revealed that approximately 40% of the genome is composed of retro-

transposable elements (Lander et al., 2001), and the latter has been shown to numerically

increase with age (De Cecco et al., 2013). Although, many theories have been proposed

for this age related increase (Yushkova et al., 2023), our results would suggest that the

concatemers containing LINE-1 and Alu elements proliferate and amplify with time in

culture by virtue of their association with DNA polymerase. Furthermore, the association

of LINE-1 and Alu elements within the concatemers with reverse transcriptase and

transposase suggested that the amplified retro-transposable elements have the capability

to indiscriminately rearrange the host genome, potentially resulting in a degree of somatic

mosaicism which would prevent the assembly of a contiguous genome. We found that

this may indeed be the case; as we observed that the cfChPs treated NIH3T3 cells began

to die out beyond passage number ~250. It should be mentioned in this context that retro-

transposable elements are implicated in cancer due to their ability to create genomic

structural variation, alterations and instability (Burns, 2017). We have earlier

hypothesized that illegitimate and repeated genomic integration of cfChPs maybe the

underlying aetiological factor for cancer as well as ageing and age related chronic

diseases (Raghuram et. al. 2019). Taken together, the above discussion suggests that

horizontal transfer of cfChPs has profound effects on genome structure and function that

are critical to human health and disease. The results also raise the possibility of

developing potential therapies for prevention of cancer and retardation of ageing using

agents that could deactivate cfChPs before they entered into new host cells (Mittra, 2024;

Khanvilkar and Mittra, 2025; Pal et. al. 2022; Pilankar et. al. 2022; Bandiwadekar et. al.

2025).

Transposable elements carried by the concatemers may also play a significant role in

mammalian evolution by inserting themselves into unpredictable locations, disrupting or

restoring gene function, generating new regulatory elements, and potentially inducing

genomic rearrangements and instability (Senft, 2021; Pourrajab, 2021). However, the

origin of transposable elements has remained in the realm of speculation. Many

transposable elements are thought to represent remnants of ancient viruses that had

become integrated into host genomes millions of years earlier (Gilbert and Feschotte,

2018). Our demonstration that the concatemers carry LINE-1 and Alu elements suggest

transposable elements are "foreign" genetic elements that are acquired from dying cells

via HGT. It has been proposed that generation of novel proteins resulting from gene

duplication, sequence divergence, and gene combination may also have evolutionary

functions (Bashton et al., 2007; Zhang et al., 2019). We speculate that the plethora of

novel fusion proteins that we discovered may likewise play a role in the evolutionary

process.

Conventionally, HGT refers to the transfer of genetic material from one organism to

another. Although HGT occurs extensively in microorganisms, it has thus far been difficult

to detect HGT in mammals. Our results suggest that HGT occurs in mammalian cells via

cfChPs released from the billions of host cells that die on a daily basis. We previously

proposed that circulating cfChPs released from dying cells function as a new class of

mobile genetic elements (MGEs) that act intra-corporeally and transfer themselves

horizontally to the organism's own cells (Mittra, 2015). Thus, "within-self" HGT may occur

in mammals on a massive scale via the medium of cfChP concatemers that have

undergone extensive and complex modifications resulting in their behaviour as "foreign"

genetic elements. However, for concatemers to have evolutionary relevance in mammals,

they need to be horizontally transferred to the germ line. The latter is not unlikely given

the high turnover of daily cell death and release of cfChPs in the blood circulation which

could carry them to all cells and tissues of the body, including the germ line cells.

Numerous attempts on our part to characterise the concatemers by conventional real time

PCR were unsuccessful. Given the complex and chaotic amalgams of disparate DNA

fragments that comprise the concatemers, the primer sequences specific to a particular

gene could not find the appropriate DNA sequences to bind to. Advanced sequencing

techniques may help to define the complex nature of the DNA sequences that comprise

the concatemers. Likewise, multiple attempts to detect the components of the fusion

proteins by Western blotting also failed. Even if an antibody reacted to a peptide

component of a fusion protein, the band obtained would not correspond to the molecular

weight of the positive control band, and the possibility of it being an artefact could not be

excluded. Studies using liquid chromatography with tandem mass spectrometry may help

to delineate the composition of the fusion proteins.

Nevertheless, it is worth mentioning here that we have earlier reported results of whole

genome sequencing of NIH3T3 cells that had been treated with cfChPs isolated from

serum of cancer patients under identical conditions as used in the present study (Mittra I

et. al., 2015). Whole genome sequencing on Illumina GA IIX platform and bioinformatics

analysis of two single cell clones (D5 and B2) developed from the cfChPs treated cells

detected 25,000-28,000 strictly human reads in the mouse cell genomes. It is likely that

our highly stringent alignment criteria had greatly underestimated the number of human

sequence that we detected the mouse cells in that study. Computational analysis of the

two clones also detected 35-47 unique human Alu repeat sequences in the recipient cells

which were confirmed by PCR and Sanger sequencing. Taken together, these data

support our current results that human DNA in the form of cfChPs are present in large

numbers in the treated mouse cells and carry Alu elements within them.

The presence of extrachromosomal DNA (ecDNA) has been described in plants (Wong

and Wildman, 1972), yeast (Møller et al., 2015), *Caenorhabditis elegans* (Shoura et al.,

2017), *Drosophila* (Stanfield and Lengyel, 1979), mice, and humans (Møller et al., 2018;

Shibata et al., 2012). Recently, there has been a resurgence of interest in the role of

ecDNA in cancer (Wu et al., 2022; Yan et al., 2024). In particular, it has been suggested

that ecDNA can be a source of extreme gene amplification enabling a cell to harbour

multiple copies of oncogenes and can potentially promote tumour heterogeneity and

therapy resistance (Wu et al., 2022; Yan et al., 2024). It is believed that ecDNAs are

derived from the host cell's own genome resulting from chromothripsis, DNA damage, or

genomic instability (Wu et al., 2022). Our study which also detected multiple highly

amplified oncogene products, often appearing as fusion proteins, raises the possibility

that ecDNAs that have been detected in human cancers may represent concatemers

composed of cfChPs acquired from surrounding dying cancer cells via HGT. The origin

and biological properties of ecDNA call for further in-depth investigations in light of our

current findings.

Our study leaves many unanswered questions, and our findings will require to be critically

validated in future research. As such, our results should be viewed as representing an

expansive theory, wherein the activities of the concatemers formed by amalgamation of

horizontally transferred cfChPs are governed by unknown biological processes that are

distinct from those that govern the nuclear genome. Our results also suggest that non-

coding DNA may have many unique biological functions which remain dormant but are

activated following cellular apoptosis to become detectable in association with the cfChP

cocatemers. An understanding of these unexplored biological processes could offer a

new perspective on the process of mammalian evolution.

In summary, the present findings lead to a novel hypothesis that a cell simultaneously

harbours two genome forms that function independently of each other: one that is

inherited (hereditary genome) and numerous others that are acquired (satellite genomes).

The latter acting as autonomous genomic agents blur the line between inherited and

acquired genetic material, opening new research avenues in biology and medicine. The

satellite genomes may potentially perform evolutionary functions by acting as vehicles of

transposable elements and generating a plethora of novel proteins. Our findings suggest

that the biological life of the genome is not a co-terminus with the death of a cell, but

rather that the genome is recycled in a fragmented and extensively modified incarnation

to perform new functions in distant destinations acting as "within-self" foreign genetic

elements. This suggests that a cell's propensity to internalise cfChPs represents a natural

biological process designed to modify the hereditary genome, which in turn may promote

evolutionary change. We hypothesise that mammalian evolution is driven by "within-self"

HGT of cfChP concatemers that have undergone such extensive and complex

modifications that they behave as "foreign" genetic elements.

Materials and Methods

Ethics approval

This study was approved by the Institutional Ethics Committee (IEC) of the Advanced

Centre for Treatment, Research and Education in Cancer, Tata Memorial Centre for the

collection of blood (10 mL) from cancer patients and healthy individuals for the isolation

of cfChPs (approval no. 900520). A formal informed consent form that was approved by

the IEC was signed by each participant.

Isolation of cfChPs from human serum

cfChPs were isolated from sera of five patients suffering from cancer and five healthy

volunteers using the protocol described by us earlier (Mittra et al., 2015). Serum from five

donors of each group was pooled prior to a cfChPs isolation. The steps of the isolation

protocol can be summarized as follows: 1) centrifugation of the pooled serum samples (1

mL) at 700,000g for 16 h at 4°C; 2) treating the pellet obtained with lysis buffer; 3)

centrifugation of the lysate at 700,000g for 16 h at 4°C; 4) suspension of the pellet in 1 ml

PBS; 5) passing the suspension through an affinity column (ThermoFisher Scientific,

USA) containing biotinylated anti-histone H4 antibody (125 µg) bound to 2 mL of Pierce®

Streptavidin Plus Ultralink® Resin (ThermoFisher Scientific, USA); 6) elution of the

column with 1 mL 0.25 M NaCl; 7) ultra-centrifuging the elute as described above and

suspending the pellet containing cfChPs in 1 mL PBS. The presence of cfChPs was

confirmed using a nucleosome specific sandwich ELISA kit (Cell Death Detection

ELISA[PLUS] kit, Roche Diagnostics GmbH, Germany). A representative electron

microscopy image of the isolated cfChPs is given in Supplementary Fig. S1. The

concentrations of cfChPs in the isolates are expressed in terms of their DNA content, as

estimated using the PicoGreen dsDNA quantitation assay (ThermoFisher Scientific, USA)

(Mittra et al., 2015). The age, sex, and tumour types of the participants are given in

Supplementary Table S1.

Cell lines and culture

NIH3T3 mouse fibroblasts were obtained from the American Type Culture Collection

(ATCC; Manassas, VA, USA). The cells were maintained in Dulbecco's modified Eagle

medium (DMEM) containing 10% bovine calf serum and grown at 37°C in 35-mm culture

dishes in an atmosphere of 95% air, 5% CO_2, and 95% humidity. The cells (10×10^4)

were treated with 10 ng of cfChPs isolated from the serum of cancer patients or healthy

individuals and maintained in continuous culture by passaging every fourth day. We

chose to use 10ng based on our earlier report in which we had obtained robust biological

effects such as activation of DDR and apoptotic pathways using this concentration of

cfChPs (Mittra I et. al., 2015).

872 Collection of conditioned medium containing cfChPs released from dying MDA-
873 MB-231 human breast cancer cells

In addition to cfChPs isolated from the cancer patients and healthy individuals, we also

investigated the effects of treating NIH3T3 cells with cfChPs that were released into the

conditioned medium from hypoxia-induced dying MDA-MB-231 human breast cancer

cells. The protocol for collecting cfChPs from hypoxia induced dying cells has been

described by us in detail earlier (Raghuram et al., 2024). The only difference was that in

the earlier study we had used dying NIH3T3 cells instead of MDA-MB-231 cells for

collecting media containing hypoxia induced cfChPs. In the dual-chamber system used

in these experiments, the pore size of the filter separating the two chambers was 400μm.

Consequently, the cfChPs from the dying MDA-MB-231 cells that were collected in the

lower chamber were <400μm in size. NIH3T3 cells (1×10^5) were treated with 1 ml of the

hypoxic media containing cfChPs.

886 Fluorescent dual labelling of cfChPs

In some experiments, the DNA and histones of the isolated cfChPs from serum were

fluorescently dually labelled with Platinum Bright™ 550 Red Nucleic Acid Labelling Kit

(Kreatech Diagnostics, Cat # GLK-004) and ATTO 488 NHS-ester (ATTO-TEC GmbH,

Cat # AD488-35), respectively, according to our previously reported protocol (Mittra et al.,

2015). NIH3T3 cells were treated with 10 ng of dually labelled cfChPs for 6 h, washed

with PBS x 3 and examined under a confocal microscope (Carl-Zeiss, GmbH, Germany).

A representative image of dually labelled cfChPs is given in Fig. 1a.

Preparation of chromatin fibres

Chromatin fibres were prepared from NIH3T3 cells according to previously described

methods (Nieminuszczy et al., 2016; Quinet et al., 2017) with minor modifications.

Although the inventers of this technique had described the fibres as DNA fibres, when

stained with histone H4 antibody, we found the fibres to comprise of chromatin

(Supplementary Fig. S12). To prepare the chromatin fibres, 2 µl of the cfChPs treated

NIH3T3 cell suspension was spotted at one end of a glass slide and semi-evaporated for

10 min at room temperature (~25°C). The spotted cells were incubated with 7 µL of lysis

buffer (0.1% sodium dodecyl sulphate in phosphate-buffered saline) for 2 min with gentle

shaking, and the slides were tilted at an angle of 15°–25° to allow the cell lysate containing

the chromatin fibres to roll down along the slide surface. The slides were fixed with chilled

methanol for 10 min and processed for immunofluorescence and FISH analyses.

Metaphase spread preparation

909 Metaphase spreads from NIH3T3 cells treated with cfChPs in continuous culture were

910 prepared using a standard protocol (Mittra et al., 2015). The slides were processed for

911 immunofluorescence and FISH analyses.

Human specificity of antibodies and FISH probes

Sources and other details of the antibodies and FISH probes used in this study are

provided in Supplementary Table S2. Irrespective of the specifications of the antibodies

given in the vendors' data sheets, and the custom synthesised FISH probes, we

independently verified their human specificity by undertaking extensive positive and

negative control experiments to ensure that all antibodies and FISH probes were human

specific. The results of these control experiments are given in Supplementary Figs. S2

and S5 – S9 and S13.

Immunofluorescence and FISH

The expression of various proteins in the cells was evaluated using indirect

immunofluorescence, and the presence of human DNA in chromatin fibres and

metaphase spreads was detected using FISH, as described previously (Mittra et al.,

2015). Briefly, for immunofluorescence, the chromatin fibres were fixed in 4%

paraformaldehyde, blocking in saponin buffer containing 10% normal goat serum followed

by immune-staining with appropriate primary and secondary antibodies. In order to

remove unbound antibodies, 1X PBS containing 0.05% Tween20 washes were given x 3

before mounting with VectaShield DAPI for microscopic examination. In case of

immunofluorescence on metaphase spreads, the paraformaldehyde fixation step was

omitted. For FISH on both chromatin fibres and metaphase spreads, the slides were

dehydrated using alcohol series (70%, 80% and 100%) followed by hybridisation

overnight with appropriate FISH probes in a humidified chamber at 37ºC. In order to

remove the unbound probes, the slides were washed in 0.4X SSC at 70ºC for 1 minute

followed by 4X SSCT (4X SSC in 0.05% Tween20) washes at 45ºC for 5 minutes each.

The final wash was in 4X SSCT for 2 minutes at room temperature followed by mounting

with VectaShield DAPI for microscopic examination. For immuno-FISH experiments, the

slides were first stained for immunofluorescence as described above and fixed with 2%

paraformaldehyde for 10 minutes followed by FISH procedure as described above. For

microscopic examination, immunofluorescence slides were imaged at 400× magnification

while FISH slides were analysed at 600× under an Applied Spectral Bio-imaging System

(Applied Spectral Imaging, Israel). The immuno-FISH slides were analysed at 600×

magnification.

Detection of DNA synthesis

947 To detect DNA synthesis on chromatin fibres and metaphase spreads, cells were pulse-

948 labelled while in culture with 10 µM bromodeoxyuridine (BrdU) for 24h, washed in PBS x

949 3, trypsinized and used for preparing chromatin fibres and metaphase spreads. Newly

950 synthesized DNA was detected using anti-BrdU antibody.

Detection of RNA synthesis

RNA synthesis in cfChP-treated cells in continuous passage was detected using an

Abcam assay kit (Cambridge, UK; Catalogue No. ab228561) as per the manufacturer's

protocol. With the use of metabolic tagging with 5-ethynyluridine (EU), this assay offers a

reliable method for fluorescent staining of newly produced RNA with the click chemistry

facilitates viewing. In some experiments, the cfChPs treated cells were treated with

actinomycin D (0.0005 µg/mL) for 24 h or grown at low temperature (31°C) for 24 h.

Statistical analysis

961 Statistical analysis was done by Student's t-test and by Analysis of variance for linear

962 trend. Both analyses were done using GraphPad version 8.

Data availability

All data generated during this study are included in this published article and supplementary information.

Ethics Statement

The Institutional Ethics Committee (IEC) provided approval to collect blood samples from cancer patients and healthy participants. Consent forms approved by the IEC (Approval no. 900520) were used to obtain signed informed consent.

Author contributions

SB, SS†, LK, RL, SP, RJ, and NKK performed the experiments; GVR, SS and IM supervised the experiments and analysed the data; IM conceptualized and designed the project, interpreted the results, was responsible for overall supervision and acquisition of funding. GVR, SS and IM wrote the paper and IM approved the final draft.

Funding

This study was supported by the Department of Atomic Energy, Government of India, through its grant CTCTMC to the Tata Memorial Centre awarded to IM. The funding agency had no role in research design, collection, analysis, and interpretation of data, or manuscript writing.

Acknowledgements

We thank Mr. Ashish Pawar for his help in preparing this manuscript. We would also like to thank Editage (www.editage.com) for English language editing support.

Conflict of interests

The authors declare no competing interests.

References

Aaronson SA, Pulciani S & Barbacid M. 1982. T24 human bladder carcinoma oncogene is an activated form of the normal human homologue of BALB-and Harvey-MSV transforming genes. Nature, 298:343-347 doi: 10.1038/298343a0

Anker P, Lyautey J, Lefort F, Lederrey C, Stroun M. 1994. Transformation of NIH/3T3 cells and SW 480 cells displaying K-ras mutation. *C R Acad Sci III* **317**:869–874.

Bandiwadekar C, Devi N L, Moiyadi A V, Singh V, Shetty P, Epari S, ... & Mittra I. 2025. Downregulation of cancer hallmarks and immune check-points in patients with glioblastoma following a short course of the pro-oxidant combination of Resveratrol and Copper. *medRxiv*, 2025-04. doi: 10.1101/2025.02.19.25322384

Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z, Wei G, Chepelev I, Zhao K. 2007. High-Resolution Profiling of Histone Methylations in the Human Genome. *Cell* **129**. doi:10.1016/j.cell.2007.05.009

Bashton M, & Chothia C. 2007. The generation of new protein functions by the combination of domains. *Structure* **15**, 85-99. doi: 10.1016/j.str.2006.11.009

Bayles KW. 2014. Bacterial programmed cell death: making sense of a paradox. *Nat Rev Microbiol* **12**:63–69.

Bergsmedh A, Szeles A, Henriksson M, Bratt A, Folkman MJ, Spetz A-L, Holmgren L. 2001. Horizontal transfer of oncogenes by uptake of apoptotic bodies. *Proceedings of the National Academy of Sciences* **98**:6407–6411.

Bijlsma R, Loeschcke V. 2005. Environmental stress, adaptation and evolution: An overviewJournal of Evolutionary Biology. doi:10.1111/j.1420-9101.2005.00962.x

Burns, K. H. 2017. Transposable elements in cancer. *Nature Reviews Cancer* ***17:***415-424. doi: 10.1038/nrc.2017.35

Bushman F. 2002. Lateral DNA transfer. Cold Spring Harbor Laboratory Press.

Cui C, Shu W, Li P. 2016. Fluorescence in situ hybridization: cell-based genetic diagnostic and research applications. *Front Cell Dev Biol* **4**:89.

De Cecco, M., et al., 2013. Transposable elements become active and mobile in the genomes of aging mammalian somatic tissues. Aging (Albany. NY). doi: https://doi.org/10.18632/aging.100621.

Dvořáková M, Karafiát V, Pajer P, Kluzáková E, Jarkovská K, Peková S, Krutílková L, Dvořák M. 2013. DNA released by leukemic cells contributes to the disruption of the bone marrow microenvironment. *Oncogene* **32**. doi:10.1038/onc.2012.553

Fliedner TM, Graessle D, Paulsen C, Reimers K. 2002. Structure and function of bone marrow hemopoiesis: mechanisms of response to ionizing radiation exposure. *Cancer Biother Radiopharm* **17**:405–426.

García-Olmo DC, Domínguez C, García-Arranz M, Anker P, Stroun M, García-Verdugo JM, García-Olmo D. 2010. Cell-Free Nucleic Acids Circulating in the Plasma of Colorectal Cancer Patients Induce the Oncogenic Transformation of Susceptible Cultured CellsCirculating DNA Induces Oncogenic Transformation. *Cancer Res* **70**:560–567.

Gilbert C, Feschotte C. 2018. Horizontal acquisition of transposable elements and viral sequences: patterns and consequences. *Curr Opin Genet Dev* **49**:15–24. doi:10.1016/j.gde.2018.02.007

Häcker G. 2013. Is there, and should there be, apoptosis in bacteria? *Microbes Infect*.

Hoffmann AA, Parsons PA. 1997. Extreme environmental change and evolution.

Hofmann A, Zakhartchenko V, Weppert M, Sebald H, Wenigerkind H, Brem G, Wolf E, Pfeifer A. 2004. Generation of transgenic cattle by lentiviral gene transfer into oocytes. *Biol Reprod* **71**. doi:10.1095/biolreprod.104.028472

Holdenrieder S, Stieber P. 2009. Clinical use of circulating nucleosomes. *Crit Rev Clin Lab Sci* **46**:1–24. doi:10.1080/10408360802485875

Khanvilkar S & Mittra I. 2025. Copper Imparts a New Therapeutic Property to Resveratrol by Generating ROS to Deactivate Cell-Free Chromatin. *Pharmaceuticals* **18**: 132. doi: 10.3390/ph18010132

Lander, E.S., et al., 2001. Initial sequencing and analysis of the human genome. *Nature* **409**:860–921. doi: 10.1038/35057062

Li W, Prazak L, Chatterjee N, Grüninger S, Krug L, Theodorou D, Dubnau J. 2013. Activation of transposable elements during aging and neuronal decline in Drosophila. *Nat Neurosci* **16**:529–531. doi:10.1038/nn.3368

MacAlpine HK, Gordân R, Powell SK, Hartemink AJ, MacAlpine DM. 2010. Drosophila ORC localizes to open chromatin and marks sites of cohesin complex loading. *Genome Res* **20**:201–211.

Mittra I. 2015. Circulating nucleic acids: A new class of physiological mobile genetic
elements. *F1000Res*. doi:10.12688/f1000research.7095.1

Mittra I, Naveen Kumar K, Gorantla Venkata R, Rohan C, Fatema K, Deepika G,
Ashwini G. 2015. Circulating nucleic acids damage DNA of healthy cells by
integrating into their genomes. *J Biosci* **40**:91–111.

Mittra I, Samant U, Sharma S, Raghuram G V, Saha T, Tidke P, Pancholi N, Gupta D,
Prasannan P, Gaikwad A, Gardi N, Chaubal R, Upadhyay P, Pal K, Rane B, Shaikh
A, Salunkhe S, Dutt S, Mishra PK, Khare NK, Nair NK, Dutt A. 2017. Cell-free
chromatin from dying cancer cells integrate into genomes of bystander healthy cells
to induce DNA damage and inflammation. *Cell Death Discov* **3**:17015.
doi:10.1038/cddiscovery.2017.15

Mittra I. 2024. Exploiting the damaging effects of ROS for therapeutic use by
deactivating cell-free chromatin: the alchemy of resveratrol and copper. *Frontiers in*
Pharmacology **15**: 1345786. doi: 10.3389/fphar.2024.1345786

Møller HD, Mohiyuddin M, Prada-Luengo I, Sailani MR, Halling JF, Plomgaard P,
Maretty L, Hansen AJ, Snyder MP, Pilegaard H, Lam HYK, Regenberg B. 2018.
Circular DNA elements of chromosomal origin are common in healthy human
somatic tissue. *Nat Commun* **9**. doi:10.1038/s41467-018-03369-8

Møller HD, Parsons L, Jørgensen TS, Botstein D, Regenberg B. 2015.
Extrachromosomal circular DNA is common in yeast. *Proc Natl Acad Sci U S A*
112:E3114-22. doi:10.1073/pnas.1508825112

Nagler C, Hardt C, Zänker KS, Dittmar T. 2011. Co-cultivation of murine BMDCs with
67NR mouse mammary carcinoma cells give rise to highly drug resistant cells.
Cancer Cell Int **11**. doi:10.1186/1475-2867-11-21

Nevo E. 2011. Evolution under environmental stress at macro-and microscales.
Genome Biol Evol **3**:1039–1052.

Nieminuszczy J, Schwab RA, Niedzwiedz W. 2016. The DNA fibre technique – tracking
helicases at work. *Methods*. doi:10.1016/j.ymeth.2016.04.019

Pal K, Raghuram G V, Dsouza J, Shinde S, Jadhav V, Shaikh A, ... & Mittra I. 2022. A
pro-oxidant combination of resveratrol and copper down-regulates multiple biological
hallmarks of ageing and neurodegeneration in mice. *Scientific Reports* **12**: 17209.
doi: 10.1038/s41598-022-21388-w

Panaud O. 2016. Horizontal transfers of transposable elements in eukaryotes: the flying
genes. *C R Biol* **339**:296–299.

Parada LF, Tabin CJ, Shih C & Weinberg RA. 1982. Human EJ bladder carcinoma
oncogene is homologue of Harvey sarcoma virus ras gene. Nature, **297**:474-478.
doi: 10.1038/297474a0

Pilankar A, Singhavi H, Raghuram G V, Siddiqui S, Khare N K, Jadhav V, ... & Mittra I.
2022. A pro-oxidant combination of resveratrol and copper down-regulates hallmarks
of cancer and immune checkpoints in patients with advanced oral cancer: Results of
an exploratory study (RESCU 004). *Frontiers in Oncology* **12**: 1000957. doi:
10.3389/fonc.2022.1000957

Pourrajab F, & Hekmatimoghaddam S. 2021. Transposable elements, contributors in
the evolution of organisms (from an arms race to a source of raw materials). *Heliyon*:
7. doi: 10.1016/j.heliyon.2021.e06029

Puonti-Kaerlas J, Eriksson T, Engström P. 1990. Production of transgenic pea (Pisum
sativum L.) plants by Agrobacterium tumefaciens—mediated gene transfer.
Theoretical and Applied Genetics **80**:246–252.

Quinet A, Carvajal-Maldonado D, Lemacon D, Vindigni A. 2017. DNA fiber analysis:
mind the gap! *Methods Enzymol* **591**:55–82.

Raghuram GV, Gupta D, Subramaniam S, Gaikwad A, Khare NK, Nobre M, Nair NK,
Mittra I. 2017. Physical shearing imparts biological activity to DNA and ability to
transmit itself horizontally across species and kingdom boundaries. *BMC Mol Biol*
18. doi:10.1186/s12867-017-0098-8

Raghuram G v., Chaudhary S, Johari S, Mittra I. 2019. Illegitimate and repeated
genomic integration of cell-free chromatin in the aetiology of somatic mosaicism,
ageing, chronic diseases and cancer. *Genes (Basel).* doi:10.3390/genes10060407

Raghuram GV, Tripathy BK, Avadhani K, Shabrish S, Khare NK, Lopes R, ... & Mittra I.
2024. Cell-free chromatin particles released from dying cells inflict mitochondrial
damage and ROS production in living cells. Cell Death Discovery **10**, 30. doi:
10.1038/s41420-023-01728-z

Sender R, Milo R. 2021. The distribution of cellular turnover in the human body. *Nat*
Med **27**:45–48. doi:10.1038/s41591-020-01182-9

Senft, A.D., Macfarlan, T.S. 2021. Transposable elements shape the evolution of
mammalian development. *Nat Rev Genet* **22**:691–711. doi: 10.1038/s41576-021-
00385-1

Shibata Y, Kumar P, Layer R, Willcox S, Gagan JR, Griffith JD, Dutta A. 2012.
Extrachromosomal microDNAs and chromosomal microdeletions in normal tissues.
Science (1979) **335**. doi:10.1126/science.1213307

Shoura MJ, Gabdank I, Hansen L, Merker J, Gotlib J, Levene SD, Fire AZ. 2017.
Intricate and cell type-specific populations of endogenous circular DNA (eccDNA) in
Caenorhabditis elegans and Homo sapiens. *G3: Genes, Genomes, Genetics* **7**.
doi:10.1534/g3.117.300141

Stanfield SW, Lengyel JA. 1979. Small circular DNA of Drosophila melanogaster:
chromosomal homology and kinetic complexity. *Proc Natl Acad Sci U S A* **76**.
doi:10.1073/pnas.76.12.6142

Thomas S, Li X-Y, Sabo PJ, Sandstrom R, Thurman RE, Canfield TK, Giste E, Fisher
W, Hammonds A, Celniker SE. 2011. Dynamic reprogramming of chromatin
accessibility during Drosophila embryo development. *Genome Biol* **12**:1–17.

Trejo-Becerril C, Pérez-Cárdenas E, Taja-Chayeb L, Anker P, Herrera-Goepfert R,
Medina-Velázquez LA, Hidalgo-Miranda A, Pérez-Montiel D, Chávez-Blanco A,
Cruz-Velázquez J, Díaz-Chávez J, Gaxiola M, Dueñas-González A. 2012. Cancer
Progression Mediated by Horizontal Gene Transfer in an In Vivo Model. *PLoS One*
7:e52754. doi:10.1371/journal.pone.0052754

Vijg, J. 2014. Somatic mutations, genome mosaicism, cancer and aging. *Current*
opinion in genetics & development, **26**:141-149. doi: 10.1016/j.gde.2014.04.002

Wong FY, Wildman SG. 1972. Simple procedure for isolation of satellite DNA's from
tobacco leaves in high yield and demonstration of minicircles. *BBA Section Nucleic*
Acids And Protein Synthesis **259**. doi:10.1016/0005-2787(72)90468-6

Wu, S., Bafna, V., Chang, H. Y., & Mischel, P. S. 2022. Extrachromosomal DNA: an
emerging hallmark in human cancer. Annual Review of Pathology: *Mechanisms of*
Disease **17**: 367-386. doi: 10.1146/annurev-pathmechdis-051821-114223

Yan X, Mischel P, Chang H. 2024. Extrachromosomal DNA in cancer. *Nat Rev Cancer*
24:261–273. doi:10.1038/s41568-024-00669-8

Yushkova, E., & Moskalev, A. 2023. Transposable elements and their role in aging.
Ageing Research Reviews, **86**:101881. doi: 10.1016/j.arr.2023.101881

Zhang L, Ren Y, Yang T, Li G, Chen J, Gschwend A R, ... & Long M. 2019. Rapid
evolution of protein diversity by de novo origination in Oryza. Nature ecology &
evolution **3**: 679-690. doi: 10.1038/s41559-019-0822-5