1   **Stable long-term individual variation in chimpanzee technological**

2   **efficiency**

Berdugo, S.[1,2], Cohen, E.[1,3], Davis, A. J.[1,3], Matsuzawa, T.[4,5,6], & Carvalho, S.[2,7,9]

[1] Social Body Lab, Institute of Human Sciences, University of Oxford, Oxford

[2] Primate Models for Behavioural Evolution Lab, Institute of Human Sciences, University of

Oxford, Oxford

[3] Wadham College, Parks Road, Oxford, OX1 3PN, United Kingdom

[4] Chubu Gakuin University, Gifu, Japan

[5] Division of the Humanities and Social Sciences, California Institute of Technology,

Pasadena, CA, USA

[6] College of Life Science, Northwest University, Xi'an 710069, China

[7] ICArEHB, Interdisciplinary Center for Archaeology and Evolution of Human Behaviour

FCHS, Universidade do Algarve, Faro 8005, Portugal

[8] Gorongosa National Park, Sofala, Mozambique

Corresponding author

Sophie Berdugo, Institute of Human Sciences,

School of Anthropology and Museum Ethnography, University of Oxford,

Oxford, OX2 6PE, United Kingdom,

sophie.berdugo@anthro.ox.ac.uk

Abstract

Using tools to access hard-to-reach and high-quality resources, such as termites, honey, and nuts, initiated a fundamental adaptive shift in human and nonhuman primate cognitive and behavioural evolution. Variation in the efficiency of extracting calorie-rich and nutrient-dense resources directly impacts energy expenditure, and potentially has significant repercussions for cultural transmission where model selection biases are employed during skill acquisition. Assessing variation in efficiency is key to understanding the evolution of complex behavioural traits in primates. Yet, individual-level differences beyond age- and sex-class in primate extractive foraging efficiency have never been investigated. Here, we used 25 years (1992–2017) of video data of the Bossou chimpanzee community (Guinea), to investigate whether individual differences in nut-cracking efficiency exist across the life span of chimpanzees aged ≥ 6 years. Data from 3,882 oil-palm nut-cracking bouts from over 800 hours of observation were collected. We found long-term stable and reliable individual differences in four (out of five) measures of efficiency. We found no sex effect, challenging previous research on a female bias in chimpanzee tool use. These life-long differences in extractive foraging impacts daily energy budgets, which potentially have significant individual fitness and life history consequences. Additionally, the establishment of long-term individual variation in chimpanzee stone tool use has implications for interpreting archaeological records of hominins. Our findings highlight the importance of longitudinal data from long-term field sites when investigating underlying cognitive and behavioural diversity across individual lifespans and between populations.

Main

The importance of individual variation in cognition and behaviour is increasingly appreciated in research on nonhuman animals[1]. Such variation has ramifications at both the individual and population levels, with broader implications for life history[2–4], cultural evolution, and interpretations of the archaeological record[5] (see Figure 1). For example, a recent large-scale meta-analysis found individual variation in the migration timing of land-, water-, and seabirds, which impacts the breeding success and survival of a given migration[6].

Yet, although cognitive and behavioural variation has been reported, the research has mostly focused on one time-point for each individual or is limited to short time spans[1,7,8]. We argue that this approach does not reflect the true extent and patterns of variation within wild populations. Here, we aim to fill this knowledge gap and establish for the first time long-term stable and reliable individual differences in chimpanzee technological efficiency.

Firstly, persistent variation in the proficiency and efficiency with which individuals extract high quality resources from the environment directly impacts energy expenditure and individual fitness. An individual who is slow or inefficient at a given extractive-foraging technique relative to others in the population will have less time and energy for other fitness-enhancing tasks, thus incurring relative fitness costs[9]. This is particularly true for complex tool-assisted foraging tasks aimed at extracting high calorie resources, such as wood-boring beetle larvae extraction by New Caledonian crows (*Corvus moneduloides*)[10], oyster-cracking by Burmese long-tailed macaques (*Macaca fascicularis aurea*)[7], and honey-dipping, termite-fishing, and nut-cracking by chimpanzees (*Pan troglodytes spp.*)[8,11,12]. Understanding the extent and causes of individual differences in extractive-foraging tasks is key for identifying variation in factors known to be relevant to evolutionary fitness.

Secondly, there is growing recognition that variation in tool use produces variation in the traces left in the archaeological record, particularly for the signatures produced during percussive

behaviours[5]. For example, the number of unintentional flakes produced by wild bearded

capuchin monkeys (*Sapajus libidinosus*) whilst cracking nuts with stones will vary according

to the tool user, with the skill (or lack thereof) of the individual determining the frequency of

mishits and subsequent flakes[13]. Investigating the archaeological signatures of known

individuals can inform our understanding of hominin fossil sites and the context behind

technological traces[5]. However, there is currently no research on individual-level variation in

chimpanzee stone tool use, constituting a significant gap in our understanding of primate

technological behaviour and its potential implications for hominin tool use.

Thirdly, identifying systematic variation in technological behaviour is crucial for investigating

the factors influencing individual differences in behavioural acquisition via social learning.

Indeed, although there are species-typical forms of social learning for particular skills, there is

growing evidence that social learning mechanisms vary systematically across individuals[14].

Individual-level variation can develop from 'individual learning of social learning', whereby

differing experiences with previous social learning opportunities can result in different

strategies being employed in the future, or the rate of learning being altered. This phenotypic

plasticity can facilitate faster adaptation to environmental changes, potentially having profound

effects on evolutionary processes. This is particularly true where model selection and social

learning biases guide behavioural acquisition[15–18]. For example, female migrant chimpanzees

in the Taï Forest, Côte d'Ivoire, conform to the nut-cracking technique of their new community,

even when their previous technique was more efficient (in terms of strikes per nut and foraging

speed)[17]. Therefore, assessing variation contributes to our understanding of cultural

transmission, and the evolution of complex behavioural traits in primates[19].

Variation in technological behaviour in nonhuman primates is well established. For example,

only male bearded capuchins use sticks as probing tools[20]. In chimpanzees, inter- and intra-

population variation in tool use is found across all communities, including differences in

technological strategies and efficiency. For example, ant-dipping tool lengths differ between the two neighbouring communities in the Kalinzu Forest, Uganda[21]. Chimpanzees in Loango National Park, Gabon, use different grip types to perforate nests during honey extraction, which is unaffected by the soil hardness[8]. In Gombe National Park, Tanzania, females commence termite-dipping earlier, engage in fishing more frequently, and retrieve more termites per dip than males[22].

In nut-cracking, females in Taï crack *Coula edulis* and *Panda oleosa* nuts more efficiently than males[23,24]. In contrast, males in Bossou, Guinea, select, use, and transport tools more frequently than females[25], and spend a significantly greater proportion of their time cracking oil palm (*Elaeis guineensis*) nuts compared to adult females[26]. The number of hits required to successfully open the nutshell was previously found to reach an asymptote in adulthood[27], and the movement of adult crackers was considered to be stereotyped[28]. However, this finding is constrained to a measure of strikes per nut, which is only one measure of nut-cracking efficiency. Moreover, no research has sought to replicate this finding, and the presence and stability of variation in nut-cracking behaviour has not been investigated using long-term data. The combination of chimpanzees' long lifespans and the protracted learning periods (3–5 years for nut-cracking[28,29]) required to learn complex tool use necessitates longitudinal studies for assessing developmental trajectories and social learning. Although cross-sectional data can be useful for determining general developmental milestones, they do not allow for tests of individual variation across the lifespan. By obscuring potential variability, this may result in generalisations about population- or species-typical development of a behavioural trait. Tracking the development of a skill across an individual's life course is key for understanding the ontogenetic processes involved—and potential variation in—fitness-enhancing traits, including extractive technologies.

This study assessed five measures (see Table 1) of post-learning period nut-cracking efficiency

for 21 individuals present in a 25-year video archive of wild chimpanzees. Data were gathered

from all post-learning period individuals present in the archive for each year they were in the

footage (see Figure 2). 'Post-learning period' was defined as chimpanzees aged 6 and over,

based on the previously established critical learning period lasting until 5-years-old[28]. This

research is the first to take such a longitudinal approach to investigate the presence and stability

of variation in primate extractive foraging. To our knowledge, this research is also the first to

assess the reliability (whether individual differences are captured consistently across measures)

and stability (the relationship between measures and whether they hold over time)[1] of such

differences in a wild primate population.

134
Results

Nut-cracking bouts. A total of 3,882 oil-palm nut-cracking bouts were recorded across 21

chimpanzees (12 female; ages 6–60) over a 25-year period. Of these bouts, 318 (8.17%) ended

in failure, 336 (8.66%) ended with a smashed kernel (which results in only partial retrieval of

the kernel), and 3,228 (83.17%) ended in the successful retrieval of a whole kernel.

Individual variation. Analyses revealed that including random intercepts for individual

chimpanzees improved model fit for all measures: *bout duration* ($\chi^2(1) = 369.49$, $p < 0.0001$),

strikes per nut ($\chi^2(1) = 475.26$, $p < 0.0001$), *success rate* ($\chi^2(1) = 47.16$, $p < 0.0001$),

displacement rate ($\chi^2(1) = 87.022$, $p < 0.0001$), and *tool switch rate* ($\chi^2(1) = 4.17$, $p < 0.05$).

Age had a significant fixed effect on *bout duration* ($t = 7.724$, $p < 0.001$) and *strikes per nut* (z

$= 8.51$, $p < 0.001$), *displacement rate* ($z = -2.40$, $p < 0.05$), and *tool switch rate* ($z = -2.462$, p

< 0.05). There was a significant fixed effect of sex for *tool switch rate* ($z = -2.502$, $p < 0.05$),

but not for *bout duration* ($t = 0.238$, $p > 0.05$), *strikes per nut* ($z = 0.407$, $p > 0.05$), *success*

rate ($z = -0.1296$, $p > 0.05$), or *displacement rate* ($z = -1.029$, $p > 0.05$).

Stability of variation. Next, we ranked the random intercepts for each multilevel model;

random intercepts represent estimated individual-level effects on the outcome variable in the

model. Lower ranks represent greater relative nut cracking efficiency: fewer strikes per nut,

shorter bout durations, *etc*. Ranks ranged from 1 to 21 (the total number of individuals in the

dataset, see <u>Figure 3</u>). Individual ranks for *bout duration*, *strikes per nut*, *success rate*, and

displacement rate were moderately-strongly correlated (see Figure S1), meaning that when an

individual was ranked highly on one of these measures, they also ranked highly in the other

two. However, *tool switch rate* was only weakly–moderately correlated, suggesting it reflects

a different underlying construct.

Reliability of variation. The consistency of the relative rank of each chimpanzee's random

intercepts obtained in the *bout duration*, *strikes per nut*, *success rate*, and *displacement rate*

models indicates reliable individual differences in nut-cracking efficiency. A two-way intra-

class correlation (ICC) analysis also indicated good consistency, $ICC(C,1) = 0.718$, $F(20,60) =$

11.2, $0.543 < ICC < 0.857$, $p < 0.001$. Including *tool switch rate* in the two-way ICC decreased

the internal consistency of the efficiency measures ($ICC(C,1) = 0.558$, $F(20,80) = 7.32$, 0.366

$< ICC < 0.751$, $p < 0.001$), further suggesting it reflects a different underlying construct.

Inter-rater reliability. Two independent, hypothesis-blind coders reviewed 70 hours (8.41%)

of footage for inter-coder reliability analyses. Unweighted Cohen's κ and ICC analyses

indicated substantial–excellent agreement and consistency between coders (see Supplementary

Information).

Discussion

This research is the first to establish the presence of long-term stable and reliable systematic

individual differences in the technological efficiency of wild chimpanzees. This finding

highlights the necessity to move beyond exclusively using averages when investigating daily

energy expenditure and activity budgets—factors with large implications for life history and

foraging ecology in primates. Variation in time and energy expended for nut-cracking will

produce variation in resource allocation to other fitness-enhancing traits. Thus, the extent of

variation in nut-cracking efficiency may have large implications for other factors impacting

survival and reproduction at the individual-level. In turn, this will inform our understanding of

potential individual differences in social learning, which could directly impact the evolutionary

fitness of individuals as well as the evolution of cultural traits.

Moreover, this research is the first to demonstrate individual variation in chimpanzee stone tool

use. This finding reiterates the need to consider variation in the archaeological signatures left

by percussive technological behaviours, with some individuals potentially contributing more

to the record than others[5]. Research is now needed to ascertain the sources of this variation. A

particular focus on the nut-cracking learning period is key given the known developmental

drivers of variation in technological behaviour[22,30,31].

Each variable measured captured different aspects of 'efficiency', with *bout duration*, *strikes*

per nut, *success rate*, and *displacement rate* being closely correlated. Bout duration is an

indicator of energy intake, as shorter bouts allow more kernels to be retrieved over the course

of a feeding session. This may improve an individual's food security, as they will be at a

competitive advantage compared to conspecifics in terms of access to more nuts. Also, longer

bouts mean that the individual's attention is focused on the task for a relatively longer period

of time, such that there is reduced time to be allocated to fitness-enhancing behaviours such as

defence, socialisation, and grooming. Conversely, the number of strikes required to retrieve the

kernel and the success rate of the individual are indicators of energy expenditure relative to

energy intake. The number of strikes per nut also dictates how convenient it is for the individual

to use stones to open the nuts (rather than, for example, using their teeth or scrounging kernels),

with more convenient techniques being favoured due to their increased efficiency[31]. For

success rate, future research should investigate why chimpanzees choose to end a nut-cracking

bout before the kernel is extracted and assess whether certain individuals stop bouts earlier than

others (i.e., do not expend unnecessary energy and are hence more efficient).

Displacement rate measures the ability, or lack thereof, to judge the amount of kinetic energy

required to strike a nut. Displacing the nut not only expends more energy than is required, but

also elongates the bout duration, particularly if the individual has to travel to retrieve the nut.

This also reduces the convenience of using tools. Primate archaeological research should

investigate whether individuals with higher displacement rates are also more likely to fracture

their stone tools, resulting in potential unintentional flakes[33].

Our results also suggest that the number of times individuals adjust their tools during a bout

reflects a different underlying process. Tool switch rate is a proxy measure for the individual's

ability to select an efficient tool composite and position the stones in such a way to reduce the

energetic effort required to extract the kernel. Given the importance of tool properties in

determining the efficiency of the behaviour[17,32,34], learning to select the best tools is also an

important aspect of successful foraging though our results suggest that it does not relate to how

efficient the individual is at cracking nuts *per se*.

Unlike previous research, we found inconsistent effects of sex on technological efficiency. Sex

had no significant effect on bout duration, strikes per nut, success rate, or displacement rate,

but there was a significant effect on tool switch rate, with male chimpanzees switching tools

less frequently than female chimpanzees. This finding contradicts previous research

establishing a female bias in technological behaviour across the genus *Pan*[35]. The lack of sex

differences in this research may suggest that the mechanism(s) establishing a female bias in

nut-cracking efficiency (measured as the mean number of strikes per nut and nuts per minute)

in Taï[23,24] are not present in Bossou. However, it may be that the apparent female bias may be

an artefact of short-term, cross-sectional analyses, and that the effect does not hold

longitudinally. Further long-term research is required to determine whether this finding holds

for other populations of nut-cracking chimpanzees.

Age had a significant fixed effect on bout duration, strikes per nut, displacement rate, and tool

switch rate. Increasing age corresponds to longer bouts and more strikes per nut, which may

reflect greater muscle weakness and the need for more rests in old age. Indeed, there have been

many elderly chimpanzees in the Bossou population, with six individuals in the archive being

aged 45+. Conversely, increasing age also corresponds to fewer nut displacements and fewer

tool switches during bouts. This may reflect older individuals' greater proficiency at selecting

appropriate tools, and better ability at judging the amount of kinetic energy required for each

strike. More research is needed to assess the impact of old age on chimpanzee technological

capabilities.

The presence of variation in the different measures of nut-cracking efficiency may indicate

variation in the underlying cognitive and/or motor capacities. For example, longer bout

durations may signify that certain individuals are prone to distraction rather than an individual

being inherently less skilled. Therefore, the stable individual variation in nut-cracking

efficiency may suggest that there are stable cognitive differences between the individuals,

which potentially has implications for performance in other key fitness-enhancing behaviours.

Future research should assess whether variation exists in other behaviours in the community to

determine the extent of these cognitive differences.

The research is limited by two main points. First, despite the experimental nature of the outdoor

laboratory, there are risks of confounding variables. For example, there is no control for what

else each individual has consumed in a day or levels of physical activity, both of which could

influence motivations for cracking nuts[36]. However, footage selected for analysis was randomly

sampled, representing a period of many years, in an attempt to amplify the signal-to-noise ratio.

Second, the outdoor laboratory is a field experimental set up, with nuts and stones being

provisioned, which could be argued decreases the validity of the findings. However, the nut-

cracking that occurs in the outdoor laboratory is no different from the nut-cracking that the

Bossou chimpanzees perform at the natural cracking site of Moblim (07°38'20.7'' N;

008°30'39.2'' W)[25]. Moreover, the location of the outdoor laboratory was specifically selected

to be on Mont Gban's summit—the core of the home range for this chimpanzee community—

to optimise the frequency of chimpanzees visiting the site[37,38]. This suggests that nut-cracking

behaviour remains unchanged regardless of site location, reiterating the ecological validity of

the outdoor laboratory.

Conclusion

This study systematically assessed individual differences in nut-cracking in the Bossou

chimpanzees using a long-term video archive of 25 years to longitudinally investigate these

differences. Our results establish stable and reliable individual-level differences across four

measures of nut-cracking efficiency, shedding light on the underlying cognitive and

behavioural diversity in the Bossou chimpanzees. This research contributes to a growing body

of evidence finding stable cognitive abilities in great apes, and points to potential variation in

the development of this extractive foraging skill. Future research should seek to establish the

factors driving this individual variation.

Methods

Study site. Bossou is a village in south-eastern Guinea (7°38'71.7'N, 8°29'38.9'W), with a tropical wet seasonal climate and a predominant population of the Manon ethnic group[38]. The neighbouring chimpanzee community resides in primary and secondary forest, with a home range of 15 km^2, although their core area is 7 km^2. The chimpanzees in Bossou have been studied continuously since 1976, with 21 individuals being present[39]. Since then, the community has been declining in size[41]. The Bossou chimpanzees use a stone hammer-and-anvil composite to extract oil palm nuts[12,33,42], with individuals requiring complementary coordinated action of both hands to manoeuvre three objects (hammer, anvil, nut) during a nut-cracking bout.

Study materials. The Bossou chimpanzees have been recorded in every dry season (December–February) since 1988, resulting in a long-term video archive of their behaviour. Researchers observed and videoed the chimpanzees in the 'outdoor laboratory'[28,37]—a 7 m x 20 m clearing in the core of the community's home range on Mount Gban (7°38'41.5″ N, 8°29'50.0″ W), which is passed through daily[28,33,37].

The outdoor laboratory is experimental in nature, with many of the available raw materials (with established weights and dimensions) and nuts being provisioned by the researchers[32]. Observations of behaviour within the outdoor laboratory occur from behind a grass screen along one edge of the clearing. All observation sessions were recorded using at least two standardised camera angles (wide- and standard-angle lenses), optimising the viewing angles.

Data collection. Behavioural analysis was conducted using *Behavioural Observation Research Interactive Software* (BORIS, v. 7.11.1)[43]. Of the 1185 videos in the Bossou archive from 1992–2017, 966 videos contained visible nut-cracking bouts by at least one individual, amounting to 832 observation hours. Where possible, the videos from the standard-angle lens data were used for analysis. If behaviour was obscured in this footage, the equivalent video

with the wide-angle lens was uploaded to attempt to observe the behaviour. If the behaviour

remained unobservable, the bout(s) were excluded. No footage was collected in 2001 or 2011,

and so these are absent from the data set.

Data from each year each focal individual was present in the archive was collected. Multiple

bouts (up to 20) per individual per year were recorded to establish the degree of within-

individual variation in efficiency, while also producing more independent data points, allowing

between-individual variation to be assessed. The full data collection protocol can be found in

the Supplementary Information.

Statistical analysis. All bouts where an infant was clinging to the focal subject were removed

from analyses ($n = 210$), as this was exclusive to certain female chimpanzees (Fana, Fanle,

Fotaiu, Jire, Pili, Velu, Vuavua, and Yo) and so could reduce the internal validity of the findings

by altering the efficiency of these individuals. Removing these bouts ensured that all subjects

were compared under equal circumstances. This left 3,672 bouts for analysis.

Only data for bouts in which a kernel was retrieved were included in the analyses for *bout*

duration and *strikes per nut* ($n = 3,367$). Excluding the 'Failed' bouts here ensured that the

amount of time it took to access the energetic reward of the enclosed kernel was analysed. The

full data set ($n = 3,672$) was analysed for *success rate*, *displacement rate*, and *tool switch rate*.

All analyses were performed using R (v. 4.0.3)[44] and RStudio (v. 2022.02.3+492) for MacOS.

Multilevel models were constructed to test for individual differences in the five measures of

efficiency. Individual chimpanzees comprised the level-two random factor, while age and sex

were included as fixed effects. Simple models (without random effects) were constructed and

compared against the multilevel models (see Tables S2–S6); ANOVAs were used to determine

the model with the best fit. The ANOVAs assessed whether including a random intercept for

individual chimpanzees significantly decreased the model prediction error. The model with the

smallest prediction error (AIC and -2-log-likelihood values) was selected for each of the five

components of efficiency.

The linear mixed-effects model for *bout duration* was fitted using the *lme4*[45] and lmerTest[46]

packages. Initially, a simple linear multilevel model (subject as a random intercept) was

constructed, with age and sex as fixed effects. However, this model fit the data poorly, and so

the normality of the outcome variable *bout duration* was assessed using a histogram. The data

were found to be strongly right skewed, and so they were log transformed to normalise the

distribution and correct for heteroskedasticity[47].

Given that the count data for *strikes per nut* did not contain zeros, zero-truncated poisson

generalised linear (mixed) models (GL(M)Ms) were fitted using the truncated_poisson family

in the *glmmTMB* package[48], with age and sex as fixed effects. However, the check_dispersion

function from the *performance* package[49] detected overdispersion in the multilevel model

(subject as a random intercept). As such, zero-truncated negative binomial GL(M)Ms were

fitted with quadratic parameterisation.

As the data for *success rate* were ordinal, cumulative link models (CLMs) and cumulative link

mixed models (CLMMs) were fitted using the *ordinal* package[50]. The 'outcome' factor was

ordered to Failed, Smash, Successful, to account for the increasing degrees of efficiency with

each increasing level of outcome[31]. The Hessian matrices were calculated for all models to

allow the model summaries to be called. The number of quadrature points used in the adaptive

Gauss-Hermite quadrature approximation was set to 7. Age was originally included as a fixed

effect but was removed as the model did not converge. Sex remained as a fixed effect.

Negative binomial GL(M)Ms were fitted for *displacement rate* and *tool switch rate* using the

glmmTMB package. All models specified zero-inflation as being equal for all observations,

given the large number of zeros in the data where an individual did not displace the nut ($n =$

2909) or switch their tools ($n = 3308$) in a bout.

The individuals' random intercepts were ranked for the *bout duration*, *strikes per nut*, *success*

rate, *displacement rate*, and *tool switch rate* models, and a correlation matrix was constructed

to determine whether they represent the same underlying construct. The two-way ICC used to

assess the stability of individual variation was selected according to recommended guidelines[51].

Data availability

A sample dataset is provided in the public repository, below.

Code availability

All analysis code needed to reproduce the results and figures reported in the paper and the

Supplementary Information can be found in the following public repository:

https://github.com/arranjdavis/chimpanzee_nut_cracking_efficiency

References

1. Bohn, M. *et al.* Great ape cognition is structured by stable cognitive abilities and predicted by developmental conditions. *Nat. Ecol. Evol.* (2023) doi:10.1038/s41559-023-02050-8.

2. Pontzer, H. *et al.* Primate energy expenditure and life history. *Proc. Natl. Acad. Sci.* **111**, 1433–1437 (2014).

3. Kaplan, H. S., Hill, K., Lancaster, J. & Hurtado, A. M. A Theory of Human Life History Evolution: Diet, Intelligence, and Longevity. *Evol. Anthropol.* **9**, 156–185 (2000).

4. Kraft, T. S. *et al.* The energetics of uniquely human subsistence strategies. *Science.* **374**, (2021).

5. Pascual-Garrido, A., Carvalho, S. & Almeida-Warren, K. Primate archaeology 3.0. *Am. J. Biol. Anthropol.* 1–17 (2023) doi:10.1002/ajpa.24835.

6. Franklin, K. A. *et al.* Individual repeatability of avian migration phenology: A systematic review and meta-analysis. *J. Anim. Ecol.* **91**, 1416–1430 (2022).

7. Tan, A., Tan, S. H., Vyas, D., Malaivijitnond, S. & Gumert, M. D. There Is More than One Way to Crack an Oyster: Identifying Variation in Burmese Long-Tailed Macaque (*Macaca fascicularis aurea*) Stone-Tool Use. *PLoS One* **10**, e0124733 (2015).

8. Estienne, V., Stephens, C. & Boesch, C. Extraction of honey from underground bee nests by central African chimpanzees (*Pan troglodytes troglodytes*) in Loango National Park, Gabon: Techniques and individual differences. *Am. J. Primatol.* **79**, e22672 (2017).

9. Biro, D., Haslam, M. & Rutz, C. Tool use as adaptation. *Philos. Trans. R. Soc. B* **368**, (2013).

10. Rutz, C. *et al.* The Ecological Significance of Tool Use in New Caledonian Crows. *Science.* **329**, 1523–1526 (2010).

11. Bessa, J., Hockings, K. J. & Biro, D. First Evidence of Chimpanzee Extractive Tool Use in Cantanhez, Guinea-Bissau: Cross-Community Variation in Honey Dipping. *Front. Ecol. Evol.* **9**, 1–14 (2021).

12. Matsuzawa, T. Stone Tools for Nut-Cracking. in *The Chimpanzees of Bossou and Nimba* (eds. Matsuzawa, T., Humle, T. & Sugiyama, Y.) 73–84 (Springer, 2011).

398   13.   Luncz, L. V, Arroyo, A., Falótico, T., Quinn, P. & Proffitt, T. A primate model
399   for the origin of flake technology. *J. Hum. Evol.* **171**, 103250 (2022).

400   14.   Mesoudi, A., Chang, L., Dall, S. R. X. & Thornton, A. The Evolution of
401   Individual and Cultural Variation in Social Learning. *Trends Ecol. Evol.* **31**, 215–
402   225 (2016).

403   15.   Henrich, J. & Gil-White, F. J. The evolution of prestige: freely conferred
404   deference as a mechanism for enhancing the benefits of cultural transmission.
405   *Evol. Hum. Behav.* **22**, 165–196 (2001).

406   16.   Kendal, R. *et al.* Chimpanzees copy dominant and knowledgeable individuals:
407   implications for cultural diversity. *Evol. Hum. Behav.* **36**, 65–72 (2015).

408   17.   Luncz, L. V, Sirianni, G., Mundry, R. & Boesch, C. Costly culture: differences in
409   nut-cracking efficiency between wild chimpanzee groups. *Anim. Behav.* **137**, 63–
410   73 (2018).

411   18.   Tan, A., Hemelrijk, C. K., Malaivijitnond, S. & Gumert, M. D. Young macaques
412   (*Macaca fascicularis*) preferentially bias attention towards closer, older, and
413   better tool users. *Anim. Cogn.* **21**, 551–5563 (2018).

414   19.   Tan, A. From Play to Proficiency: The Ontogeny of Stone-Tool Use in Coastal-
415   Foraging Long-Tailed Macaques (*Macaca fascicularis*) From a Comparative
416   Perception-Action Perspective. *J. Comp. Psychol.* **131**, 89–114 (2017).

417   20.   Falótico, T., Bueno, C. Q. & Ottoni, E. B. Ontogeny and sex differences in object
418   manipulation and probe tool use by wild tufted capuchin monkeys (*Sapajus*
419   *libidinosus*). *Am. J. Primatol.* **83**, (2021).

420   21.   Koops, K., Schöning, C., Isaji, M. & Hashimoto, C. Cultural differences in ant-
421   dipping tool length between neighbouring chimpanzee communities at Kalinzu,
422   Uganda. *Sci. Rep.* **5**, (2015).

423   22.   Lonsdorf, E. V. Sex differences in the development of termite-fishing skills in the
424   wild chimpanzees, *Pan troglodytes schweinfurthii*, of Gombe National Park,
425   Tanzania. *Anim. Behav.* **70**, 673–683 (2005).

426   23.   Boesch, C. & Boesch, H. Sex Differences in the Use of Natural Hammers by Wild
427   Chimpanzees: A Preliminary Report. *J. Hum. Evol.* **10**, 585–593 (1981).

428   24.   Boesch, C. & Boesch, H. Possible Causes of Sex Differences in the Use of Natural
429   Hammers by Wild Chimpanzees. *J. Hum. Evol.* **13**, 415–440 (1984).

25. Carvalho, S., Cunha, E., Sousa, C. & Matsuzawa, T. Chaînes opératoires and resource-exploitation strategies in chimpanzee (*Pan troglodytes*) nut-cracking. *J. Hum. Evol.* **55**, 148–163 (2008).

26. Bain, M. *et al.* Automated audiovisual behavior recognition in wild primates. *Sci. Adv.* **7**, eabi4883 (2021).

27. Biro, D., Sousa, C. & Matsuzawa, T. Ontogeny and cultural propagation of tool use by wild chimpanzees at Bossou, Guinea: Case studies in nut-cracking and leaf-folding. in *Cognitive Development in Chimpanzees* (eds. Matsuzawa, T., Tomonaga, M. & Tanaka, M.) 476–506 (Springer, 2006).

28. Biro, D. *et al.* Cultural innovation and transmission of tool use in wild chimpanzees: evidence from field experiments. *Anim. Cogn.* **6**, 213–223 (2003).

29. Inoue-Nakamura, N. & Matsuzawa, T. Development of Stone Tool Use by Wild Chimpanzees (*Pan troglodytes*). *J. Comp. Psychol.* **111**, 159–173 (1997).

30. Lonsdorf, E. V, Eberly, L. E. & Pusey, A. E. Sex differences in learning in chimpanzees. *Nat. Br. Commun.* **428**, 715–716 (2004).

31. Humle, T., Snowdon, C. T. & Matsuzawa, T. Social influences on ant-dipping acquisition in the wild chimpanzees (*Pan troglodytes verus*) of Bossou, Guinea, West Africa. *Anim. Cogn.* **12**, S37–S48 (2009).

32. Sirianni, G., Luncz, L. V & Gratton, P. An energetic model of foraging optimization: wild chimpanzee hammer selection for nut-cracking. in *The Chimpanzees of the Taï Forest: 40 Years of Research* (eds. Boesch, C. & Wittig, R.) 104–124 (Cambridge University Press, 2019). doi:10.1017/9781108674218.009.

33. Carvalho, S., Biro, D., McGrew, W. C. & Matsuzawa, T. Tool-composite reuse in wild chimpanzees (*Pan troglodytes*): archaeologically invisible steps in the technological evolution of early hominins? *Anim. Cogn.* **12**, 103–114 (2009).

34. Schrauf, C., Call, J., Fuwa, K. & Hirata, S. Do Chimpanzees Use Weight to Select Hammer Tools? *PLoS One* **7**, 1–12 (2012).

35. Gruber, T., Clay, Z. & Zuberbühler, K. A comparison of bonobo and chimpanzee tool use: evidence for a female bias in the *Pan* lineage. *Anim. Behav.* **80**, 1023–1033 (2010).

461   36.   Koops, K., Gaspard Soumah, A., van Leeuwen, K. L., Didier Camara, H. &
462   Matsuzawa, T. Field experiments find no evidence that chimpanzee nut cracking
463   can be independently innovated. *Nat. Hum. Behav.* **6**, 487–494 (2022).

464   37.   Matsuzawa, T. Field experiments on use of stone tools by chimpanzees in the
465   wild. in *Chimpanzee Cultures* (eds. Wrangham, R. W., McGrew, W. C., De Waal,
466   F. B. M. & Heltne, P. G.) 351–370 (Harvard University Press, 1994).

467   38.   Sugiyama, Y., Fushimi, T., Sakura, O. & Matsuzawa, T. Hand Preference and
468   Tool Use in Wild Chimpanzees. *Primates* **34**, 151–159 (1993).

469   39.   Hockings, K. J., Anderson, J. R. & Matsuzawa, T. Socioecological adaptations
470   by chimpanzees, *Pan troglodytes verus*, inhabiting an anthropogenically
471   impacted habitat. *Anim. Behav.* **83**, 801–810 (2012).

472   40.   Sugiyama, Y. & Koman, J. Social Structure and Dynamics of Wild Chimpanzees
473   at Bossou, Guinea. *Primates* **20**, 323–339 (1979).

474   41.   Matsuzawa, T. & Humle, T. Bossou: 33 years. in *The Chimpanzees of Bossou and*
475   *Nimba* (eds. Matsuzawa, T., Humle, T. & Sugiyama, Y.) 3–10 (Springer, 2011).

476   42.   Sugiyama, Y. & Koman, J. Tool-Using and -Making Behaviour in Wild
477   Chimpanzees at Bossou, Guinea. *Primates* **20**, 513–524 (1979).

478   43.   Friard, O. & Gamba, M. BORIS: a free, versatile open-source event-logging
479   software for video/audio coding and live observations. *Methods Ecol. Evol.* **7**,
480   1325–1330 (2016).

481   44.   R Core Team. R: A Language and Environment for Statistical Computing. (2020).

482   45.   Bates, D., Maechler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects
483   Models Using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

484   46.   Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package:
485   Tests in Linear Mixed Effects Models. *J. Stat. Softw.* **82**, 1–26 (2017).

486   47.   Field, A., Miles, J. & Field, Z. *Discovering Statistics Using R*. (SAGE
487   Publications, 2012).

488   48.   Brooks, M. E. *et al.* glmmTMB Balances Speed and Flexibility Among Packages
489   for Zero-inflated Generalized Linear Mixed Modeling. *R J.* **9**, 378–400 (2017).

490   49.   Lüdecke, D., Ben-Shachar, M. S., Patil, I. & Makowski, D. performance: An R
491   Package for Assessment, Comparison and Testing of Statistical Models. *J. Open*
492   *Source Softw.* **6**, 3139 (2021).

50.   Christensen, R. H. B. ordinal - Regression Models for Ordinal Data. (2019).

51.   Koo, T. K. & Li, M. Y. A Guideline of Selecting and Reporting Intraclass
Correlation Coefficients for Reliability Research. *J. Chiropr. Med.* **15**, 155–163
(2016).

52.   Neufuss, J., Humle, T., Cremaschi, A. & Kivell, T. L. Nut-cracking behaviour in
wild-born, rehabilitated bonobos (*Pan paniscus*): a comprehensive study of hand-
preference, hand grips and efficiency. *Am. J. Primatol.* **79**, e22589 (2017).

Acknowledgements

We thank the Direction Générale de la Recherche Scientifique et de l'Innovation

Technologique (DGERSIT) and the Institut de Recherche Environnementale de Bossou (IREB)

in Guinea for research authorisation. The video data for this research was provided by Tetsuro

Matsuzawa and Kyoto University, Japan. The original video archive was digitised, organised,

and systematised by Daniel Schofield, Department of Engineering Science, University of

Oxford. Special thanks to all the field researchers and assistants at Bossou, specifically Henry

Didier Camara, Cé Vincent Mami, Gouanou Zogbila, Boniface Zogbila, Jules Doré, Guana

Goumy, Tino Camara, Pascal Goumy, Paquílé Cherif, Jiles Gondo Doré, and Marcel Doré.

Thank you to all the researchers who contributed to the Bossou archive since its inception.

Thank you to Alexander Mielke for help cataloguing the Bossou video archive. Thank you to

Dora Biro and Katarina Almeida Warren for feedback. Thank you to Zhangzhuoran Dai and

Victoire Martignac for being independent, hypothesis-blind coders for the inter-rater reliability

analyses.

Author contributions

S.B. conceived of the study, designed, and coordinated the study, collected data, participated

in the analysis and visualisation of the data, and wrote the manuscript. **E.C.** provided

supervision, participated in the design of the study, and commented on the manuscript. **A.J.D.**

analysed and visualised the data and commented on the manuscript. **T.M.** collected the original

data set and commented on the manuscript. **S.C.** collected the original data set, provided

supervision, participated in the design of the study, and commented on the manuscript.

Funding

This research was supported by the Clarendon Fund (University of Oxford, UK), Keble College

Sloane-Robinson Clarendon Scholarship (University of Oxford, UK), the Boise Trust Fund

(University of Oxford, UK) to **S.B.**, and by the Ministry of Education, Culture, Sports, Science,

and Technology (MEXT), Japan, (grant nos. 12002009, 16002001, 20002001, 24000001, and

16H06283) to **T.M.**. The funders had no role in study design, data collection and analysis,

decision to publish or preparation of the manuscript.

Competing interests

530 The authors declare no competing interests.

531 **Supplementary information** is available for this paper.

532 **Correspondence and requests for materials** should be addressed to Sophie Berdugo.

 Figures

535 ***Figure 1.*** Implications of long term stable individual variation in lithic technological
536 behaviour.

554 ***Figure 2.*** Stills taken from the Bossou video archive. On the left, a six-year-old female
555 chimpanzee, Fanle, cracks an oil palm nut in 2004, the year after finishing her learning period.
556 On the right, the same female chimpanzee cracks an oil palm nut in 2017 at the age of 20.
557 Image credit: Kyoto University.

559
560 ***Figure 3.*** Ranked random intercepts from each multilevel model for the four stable and reliable
561 measures of nut-cracking efficiency.

563   **Tables**

564   *Table 1.* Definitions of efficiency measures

Efficiency measure	Definition
Bout duration	The length of the continuous period (in seconds) of nut-cracking whereby the individual strikes a single nut on an anvil with a hammer stone involving the same hand, bodily posture, hand grip, and nut (the "bout")[52]. A bout starts in the frame when the hammer is lifted for the first strike of the nut on the anvil and ends in the frame when the hammer makes contact with the nut for the final time[7].
Strikes per nut	The number of hits required to open a single nut using a stone hammer and stone anvil.
Success rate	The outcome of the nut-cracking bout. A successful bout is one where the nut is cracked using the stone hammer and anvil composite and the full kernel is retrieved and eaten. A failed bout is one where the nut is not retrieved or eaten by any individual. A smashed nut is one where the nut is opened but the kernel has broken into multiple pieces such that each piece must be eaten separately.
Displacement rate	The number of times a hammer strike resulted in the nut being displaced from the anvil.
Tool switch rate	The number of times the focal individual alters their tools whilst cracking nuts[19].