Changes in length-at-first return of a sea trout

(*Salmo trutta*) population in northern France

Quentin Josset[1,4], Laurent Beaulaton[2,4], Atso Romakkaniemi[5], Marie

Nevoux[3,4]

[1] Service conservation et gestion durable des espèces exploitées, Direction de la recherche et de l'ap-

pui scientifique, OFB, U3E, 92 Chaussée de Picardie 76260 Eu, France

[2] Service conservation et gestion durable des espèces exploitées, Direction de la recherche et de l'ap-

pui scientifique, OFB, U3E, 65 Rue de Saint-Brieuc 35042 Rennes, France

[3] DECOD (Ecosystem Dynamics and Sustainability), INRAE, Institut Agro, IFREMER, 65 Rue de Saint-

Brieuc 35042 Rennes, France

[4] Management of Diadromous Fish in their Environment, OFB, INRAE, Institut Agro, UNIV PAU & PAYS

ADOUR/E2S UPPA, Rennes, France

[5] Natural Resources Institute Finland, Paavo Havaksen tie 3, 90570 Oulu, Finland

Corresponding author:

Quentin Josset

Office Français de la Biodiversité, 92 Chaussée de Picardie 76260 Eu, France

+33276375078, quentin.josset@ofb.gouv.fr

ABSTRACT
The resilience of sea trout populations is increasingly concerning, with evidence of major

demographic changes in some populations. Based on trapping data and related scale collec-

tion, we analysed long-term changes in body length of a sea trout population in the Bresle

River, France. From 1984-2022, the length of first-time returning individuals decreased by 1.18

mm.year-1 (SD = 0.03), which resulted in the loss of *c.* 8% of mean body length. This decrease

results from a decrease in the age at first return, with a gradual loss of the oldest individuals

and an increase in the proportion of the youngest. Timing of the return migration advanced

drastically, although shorter sea sojourn had little influence on body length. We found little

evidence of a decrease in length-at-age, to the exception of the oldest age class, suggesting

that growth conditions at sea might not have deteriorated greatly during the study period.

34
35 *Keywords:* age structure, body length, life history, marine growth, migration, sea trout

Introduction

The environmental conditions that individuals experience, in conjunction with their genotype

and epigenetic mechanisms (Ferguson et al., 2019), are susceptible to influence the expression

of their phenotype and their life history trajectories (Walsh and Reznick, 2011). Migration de-

cision, survival, growth, age at maturation or fecundity can thus be influenced by a series of

intrinsic and extrinsic factors, such as environmental conditions (Crozier and Hutchings, 2014;

Satterthwaite et al., 2012), genotype (McKinney et al., 2015), inter- and intra-specific compe-

tition (Helle et al., 2007), predator-prey relationships (Ohlberger et al., 2019), and anthropo-

genic activities (Hard et al., 2008; Law, 2000).

Recent environmental changes attributed to climate change and anthropogenic activities are

assumed to be responsible for the widespread decline in salmonid abundance and distribution

(Forseth et al., 2017; Kendall and Quinn, 2011; Kennedy and Crozier, 2010; Perrier et al., 2013).

Ongoing changes, such as an increase in sea temperature, ocean acidification and stratifica-

tion, rising sea levels, and eutrophication, are likely to influence marine trophic conditions

(Möllmann and Diekmann, 2012). These changes could greatly modify the quality, quantity,

and availability of trophic resources in the ocean (Bartolino et al., 2014; Brown et al., 2010;

Niiranen et al., 2013) and thus impact growth opportunities for salmonid populations at sea.

Reduced growth opportunities can influence individuals' life history trajectories and fitness

greatly by decreasing survival, fecundity, and competitive abilities (Perez and Munch, 2010;

Quinn et al., 2016) or by delaying the age at maturation (Ishida et al., 1993). These changes

can cause large variations in the length and age structure of populations. Such changes in

population composition have been recorded for many salmon populations around the world

(Bal et al., 2017; Jeffrey et al., 2017; Lewis et al., 2015; Morita and Fukuwaka, 2007; Ohlberger

et al., 2018; Oke et al., 2020), but also sea trout (Milner et al., 2017), and could indicate a

widespread decrease in marine growth and survival. In this context, identifying the ecological

mechanisms that most likely influence these changes is crucial to better understand and pre-

dict population dynamics. This knowledge is also necessary to provide recommendations for

effective management actions and to alleviate or compensate for pressures on sea trout pop-

ulations when and where possible.

The anadromous form of the brown trout (*Salmo trutta,* L.) (hereafter, "sea trout") (ICES,

2020) exhibits an extreme diversity of life histories across its wide distribution range (Jonsson

and L'Abée-Lund, 1993), which makes it an appropriate model species to study variations in

life histories in a changing environment. The age at seaward migration (*i.e.* smolt stage) ranges

from 1-3 years in France to 5-7 years in northern Norway (Nevoux et al., 2019). First reproduc-

tion occurs after 0-2 or 2-4 years at sea in populations in the English Channel (Richard, 1981)

and the Baltic Sea (Järvi, 1940), respectively.

In recent decades, alarming collapses in population abundance, as well as altered population

structure, have been recorded in Europe, such as in the Burrishoole River, Ireland (Gargan et

al., 2006), the Ewe River, Scotland (Butler and Walker, 2006), and the Vistula River, Poland

(Dębowski, 2018). Damming of rivers forbidding the access to spawning sites, overfishing, or

the development of offshore salmon aquaculture leading to sea lice epizootics, have been

pointed as major factors of decline for some of these populations. Consequently, it is neces-

sary to assess and understand possible changes in key life history traits, such as length at

spawning, age at maturation, and length-at-age, which drive the dynamics of European sea

trout populations.

In this study, we quantified the influence of key life history and phenological variables on the

length of sea trout returning to the Bresle River, France, using 39 years of data from long-term

population monitoring. Based on an intensive trapping protocol throughout the migration sea-

son, we obtained individual data on length, migration timing, and age, as inferred from scale

reading. We first examined the temporal trend in the mean length of all fish captured in the

trap to investigate a potential decrease in body length. We then focused on individuals return-

ing for the first time to test four non-exclusive hypotheses: whether such a decrease was due

to i) long-term effects of smaller length at seaward migration in smolts, ii) younger age at first

return, iii) a shorter growing season at sea and an advanced date of return, and vi) a decrease

in the intrinsic growth rate at sea and a smaller length-at-age. The results provide insights into

the ecology of sea trout in southern Europe and illustrate ongoing changes in life history strat-

egies.

Figure 1 - Geographical location of the Bresle River, France, and its catchment

 Materials and methods

 Study site

 The Bresle River, located in northern France, drains a 742 km² catchment into the English

 Channel at Le Tréport harbour (Figure 1). Mean annual discharge is 7.3 m^3 s^{-1}, and salmonids

 can access *c.* 2/3 of its total wetted length (137 km). Human activities strongly impact the

 Bresle estuary, and the Le Tréport harbour is the first obstacle to migration from the sea;

 however, the harbour is equipped with a fishway, which provides access to the main river

 (Euzenat et al., 2007). The Bresle River hosts a wild population of sea trout that has a life cycle

 of 1, 2, or 3 years in freshwater (hereafter, 1FW, 2FW, and 3FW, respectively) and 0, 1, or 2

 years at sea until first return (hereafter, 0SW, 1SW, and 2SW, respectively).

 Monitoring protocol

 Since 1981, an intensive sea trout trapping protocol has been conducted throughout the up-

 stream migration season (March-December) each year at Eu, in the lower section of the Bresle

 River (Josset et al., 2023). It follows the ethical review from the committee C2EA-07, from the

 National Committee for Ethical Reflection on Animal Experimentation (permit number:

 APAFIS#26834-2020080615584492 v3). The trap is located on an obstacle 3 km upstream from

 the estuary and is checked twice a day for the presence of fish. Each year, a mean of 995 (SD

 = 511) sea trout are caught in the trap on their return from the sea. Each individual is measured

 to the nearest millimetre (total length from 1981-1983; fork length – from the tip of the snout

 to the fork of the tail – since 1984) using a manual or electronic measuring board. Examination

 of secondary sexual characteristics, which are visible mainly from September-December, indi-

 cates the sex of certain individuals. However, visual sexing is imperfect and relies mainly on

the presence of an elongated lower jaw (*i.e.* kype) on males. Scales are collected from a sub-

sample of individuals to determine age, following (Baglinière et al., 2020)). Data on individual

biometry, age, and scale samples are stored in the French COLISA archive (COLlection of Ichty-

ological SAmples, (Marchand et al., 2019)), https://colisa.fr/) (in French) .

Data selection

For this analysis, we used data collected over a 39-year period (1984-2022). We focused only

on individuals that returned to the river after their first marine sojourn (hereafter, "first re-

turns"), to reduce heterogeneity between individuals as biological processes that influenced

individuals returning for the first time may have differed from those that influenced individ-

uals that had already returned to the river at least once. The returning status was defined

based on i) the fish characteristics at capture and ii) on the examination of scales, which were

available on a subset of 15,730 individuals out of the 41,412 individuals captured at the trap.

Individuals with low relative weight (Wr < 90) (Blackwell et al., 2000) captured before April

were considered late-running post-spawners migrating to the sea and were thus excluded

from the analysis. Then, we defined a first return as a fish with no a spawning mark on its

scales (n = 11844, 75.3% of available samples).

Data resampling

The subsample of fish which had their scales sampled, and age assessed, was unbalanced be-

tween length classes, with individuals in the more abundant length classes being generally

under-sampled. This bias skewed estimates of the mean length-at-age. To correct this bias,

we randomly resampled within the subsample of fish which had their scales sampled, a subset

of individuals with an even effort among length classes. We first calculated the proportion of

sampled individuals for each 50 mm length classes for a given year, determined the minimum

proportion of sampling for each year, and then used this minimum proportion as a target for

resampling individuals in each length classes for that given year. This resampling procedure

was repeated 1000 times to ensure robustness and stability of the results. The following se-

lection of first returns yielded datasets of varying lengths, with a mean sample size of 6060

individuals (SD = 23).

Statistical analyses

Long-term trend in the length of returning sea trout

To describe the long-term temporal trend in the length of returning sea trout, we modelled

the mean yearly length of all individuals captured in the trap as a linear function of time and

assessed the significance and direction of the trend. This was done using the "lm()" function

of the "stats" package of R software (version 4.0.3) (R Core Team, 2020), using the year of

return as a continuous variable.

Drivers of change in length-at-first return

We investigated the influence of key ecological variables on the length of first-time returning

sea trout using Gaussian generalized linear models (McCullagh and Nelder, 1989) with a log

link. The log-link function allows for multiplicative rather than additive effects, and the former

are generally considered more appropriate for modelling ecological relations. The Gaussian

distribution was selected because it is usually used when a linear relation between dependent

and independent variables is expected, assuming a normal distribution of the residuals. Model

selection followed stepwise inclusion of variables and relevant two-way interactions depend-

ing on the hypothesis selected.

Development of dimorphic sexual characteristics before reproduction, especially a kype at the end of the lower jaw (Witten and Hall, 2001), could bias length depending on the timing of return. This would be especially true in late-running males, although some females also have a small kype. To address this bias, we used the apparent sex recorded in the phenotypic data in the model as a corrective variable, with two levels: presence or absence of a kype.

Knowing that the longer the marine sojourn, the larger the trout is upon return (Thorstad et al., 2016), we included the "sea age" (*i.e.* the number of years spent at sea) as a categorical variable to explain length-at-first return. Similarly, we hypothesised that the number of years spent in the river influenced the length-at-first return, although less than the number of years spent at sea, due to substantial differences in growth potential in fresh and seawater (Gross et al., 1988; Nevoux et al., 2019). We thus used the "river age" (*i.e.* the number of years spent in freshwater before migrating to the sea) as another categorical variable in the model. We also considered an interaction between river age and sea age to capture effects of latent intrinsic differences among individuals depending on their life history (*e.g.* growth potential, metabolic rate, energy-use efficiency). Temporal variability in the effect of age on length was also tested by considering interactions between the year and river and sea ages.

The timing of migration can also influence the length of returning trout. Assuming that the timing of smolt migration to the sea remained unchanged, an early return in freshwater would mean shorter marine sojourn, reduced growth opportunities and potentially shorter bodies. Advanced migration timing can be driven by environmental conditions experienced by all individuals of a given sea-age class, as indicated by the mean day of year of return of each sea-

age class and year (ctrAvgDOY, centred on the inter-annual mean). Within a given sea-age

class, differences in the timing of migration among individuals can also influence body length.

This individual level was captured by the variable deltaDOY, which was calculated as an indi-

vidual's return day minus the mean day of return of the corresponding sea-age class and year.

We considered two-way interactions to test whether effects of phenology variables varied

over time or among sea ages. The increase in length for each additional day at sea may have

varied during the study period, and timing of return may have more influence on body length

of young individuals returning after a short sojourn, due to their faster growth (Davaine et al.,

1997).

Once the migration strategy and timing were considered, any remaining long-term trends in

the length of returning sea trout would indicate a gradual change in the intrinsic growth rate

at sea. To describe residual temporal variability in the length-at-first return, we used the year

of return to the river (Year) as a continuous variable, which grouped sea trout that had expe-

rienced the same environmental conditions during their last months at sea.

Implementing and selecting the model

Collinearity among variables was tested using Pearson correlation coefficients, and only vari-

ables with $|r| \leq 0.7$ were kept (Dormann et al., 2013). A bootstrap procedure of the "boot"

package (Canty and Ripley, 2021; Davison and Hinkley, 1997) was used to ensure stability of

the results by reducing effects of sampling variability. One thousand datasets were generated

using the bootstrap procedure, and candidate models were fitted to each dataset. Model se-

lection aimed to minimise Akaike's information criterion (Akaike, 1974), following guidelines

of Burnham et al. (2002). We identified the best model for each dataset and compared the

models' selection frequencies as a measure of support. Mean estimates and 95% confidence

intervals of the best models' effects and interactions were then calculated from the 1000 it-

erations of the bootstrap procedure. In addition, to assess temporal changes in the age struc-

ture, yearly age-class percentages were recorded for each of the 1000 iterations. Similarly,

slopes of a linear regression between the day of year and the year of return by sea age were

extracted from each iterations to assess variations in the timing of return.

Results

Long-term decrease in the length of returning sea trout

The modelling of body length from 1984-2022 revealed that sea trout in the Bresle River lost

in average 1.18 mm (SD = 0.03) per year, regardless of their age or the number of times they

had reproduced, which represented a loss of 46.02 mm (SD = 1.17) (*i.e. c.* 8% of the mean

body length from 1984-1988) over the 39 years studied (Figure 2).

Figure 2 - Long-term temporal trend in the mean fork-length of sea trout captured in the Bresle River from 1984-2022. Error

bars indicate 1 standard deviation. The shaded zone indicates the 95% confidence interval.

Model selection

Of the 1000 iterations of the bootstrap procedure and model selection, only two models (no. 21 and 19) were selected for having the lowest AIC (selected 77.3% and 22.7% of the time, respectively) (Table 1). The two models were similar, including all major variables, and thus had similar AIC and predictive power. They differed only in that model 21 included temporal interactions with the two phenology variables (*i.e.* ctrAvgDOY×Year and deltaDOY×Year). Although model selection based on the AIC balances likelihood and parsimony, two of the most complicated models were selected, which indicates overall complexity of the mechanisms that drive length-at-first return. The mean deviance calculated from the 1000 best models explained up to 66.00% (SD = 0.49) of the variability in the length-at-first return. Examination of the residuals showed slight deviation from the hypothesis of normality. As could be expected in a large dataset, wide tail distributions lead to the rejection of the normality hypothesis despite a quasi-normal distribution.

Decrease in the age of returning sea trout is a major driver of decrease in length-at-first return

We found that sea age influenced the length-at-first return the most. The more growth seasons an individual spent at sea, the greater was its length-at-first return. This result supports the hypothesis of a shift in the age structure, with a decrease in the number of growth seasons at sea, which would then drive the decrease in length. Indeed, the mean sea age structure of first returns changed greatly during the study period (Figure 3). In particular, 2SW individuals gradually disappeared and were not represented in 3 of the last 10 years of the study period.

264 **Table 1** - The models tested to explain length-at-first return and their frequencies of selection (out of 1000 iterations) based
265 on Akaike's information criterion. FL, fork length; Kype, presence/absence of a kype; Sea-age, number of years spent at sea;
266 River-age, number of years spent in the river; ctrAvgDOY, mean day of year of return to the river by year and sea age (centred
267 by sea age); deltaDOY, individual difference in the mean day of year of return; Year, year of return to the river.

Model number	Model	Selection frequency	Average explained deviance (%)	Standard deviation
1	Null	0	0	0
2	Kype	0	8.65	0.453
3	Kype + Sea-age	0	60.2	0.514
4	Kype + Sea-age + River-age	0	63.5	0.488
5	Kype + Sea-age + River-age + Sea-age×River-age	0	64.4	0.499
6	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY	0	64.6	0.498
7	Kype + Sea-age + River-age + Sea-age×River-age + deltaDOY	0	64.7	0.502
8	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY	0	64.9	0.501
9	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×Sea-age	0	65.0	0.501
10	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×River-age	0	64.9	0.500
11	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×Sea-age + Year×River-age	0	65.1	0.500
12	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Sea-age×ctrAvgDOY	0	65.2	0.497
13	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Sea-age×deltaDOY	0	65.5	0.498
14	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Sea-age×ctrAvgDOY + Sea-age×deltaDOY	0	65.7	0.494
15	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×ctrAvgDOY	0	64.9	0.502
16	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×deltaDOY	0	64.9	0.500
17	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×ctrAvgDOY + Year×deltaDOY	0	64.9	0.502
18	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×Sea-age + Year×River-age + Year×ctrAvgDOY + Year×deltaDOY	0	65.1	0.501
19	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×Sea-age + Year×River-age + Sea-age×ctrAvgDOY + Sea-age×deltaDOY	227	65.9	0.492
20	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×ctrAvgDOY + Year×deltaDOY + Sea-age×ctrAvgDOY + Sea-age×deltaDOY	0	65.8	0.493
21	Kype + Sea-age + River-age + Sea-age×River-age + ctrAvgDOY + deltaDOY + Year×Sea-age + Year×River-age + Year×ctrAvgDOY + Year×deltaDOY + Sea-age×ctrAvgDOY + Sea-age×deltaDOY	773	66.0	0.491

At the same time, the percentage of 0SW individuals increased in recent years to up to 60%

(in 2018) of all first returning sea trout. The percentage of 1SW individuals in the population

remained predominant and stable over time (mean = 82.3%; SD = 11.5%). This pattern indi-

cates a decrease in the mean sea age at first return of the Bresle sea trout population: 1.06

years (SD = 0.4) from 1984-1988, but 0.80 years (SD = 0.4) from 2018-2022. The age structure

differed greatly in 2001, when only 1SW individual was captured due to extreme flooding that

disrupted trapping and thus decreased its efficiency greatly.

Figure 3 - Temporal variability in the age structure of first-time returning sea trout captured in the Bresle. Average proportions
derived from the 1000 datasets.

281
Earlier life history determines the length of returning sea trout

River age had a long-term effect on the length-at-first return: individuals that smoltified after

only 1 year in the river were shorter on their return than those that smoltified after 2-3 years

(Figure 4). The younger the sea-age class the more pronounced this effect was, as indicated

by the interaction between river age and sea age. For a 0SW individual smolting at age 2 or 3,

returns length was on average respectively 83.3 and 96.7 mm larger than in FW1 returns. River

age was included in all of the best models, which highlights its high explanatory value.

Figure 4 - Estimated changes in the length of sea trout at first return over time as a function of life history strategy. Mean fitted values in the 'response' scale derived from predictions on each best models. Shaded zones indicate 95% confidence interval. FW: number of years spent in river, SW: number of years spent at sea.

Length of returning sea trout is influenced by the migration timing

The two variables for migration timing (ctrAvgDOY and deltaDOY) explained some of the vari-

ability in the length-at-first return. Effects of return timing depended on the sea age, as indi-

cated by the inclusion of the two-way interactions in the model. The effect of additional days

at sea, measured at both inter-annual (ctrAvgDOY) and intra-annual (deltaDOY) scales,

seemed to benefit 0SW individuals more than 1SW individuals. In contrast, this effect was

negative for 2SW individuals, as late-returning individuals in this age class tended to be smaller

than early-returning individuals. The inclusion of the temporal interaction indicated a poten-

tial change in effects of return timing over time, although support for it was weaker, as 22.7%

of the best models did not include it.

Mean slopes of temporal linear regressions of the day of return indicated that the capture

date of first returns advanced significantly (p < 0.001), with 0SW, 1SW, and 2SW being cap-

tured a mean of 1.38 (SD = 0.12), 1.21 (SD = 0.08), and 0.49 (SD = 0.26) days earlier each year,

respectively. Over the 39-year period, the date of return thus advanced by 53, 47, and 19 days

for 0SW, 1SW, and 2SW individuals, respectively. Despite this major change, the reduction in

time spent at sea due to advanced seasonal migration timing had only a small effect on the

length of sea trout at first return (Figure 5).

Figure 5 - Mean estimated effects of each predictor's modality, compared with a reference SW-1-FW-1 sea trout with a kype
for ctrAvgDOY and deltaDOY = 0 (black vertical line). Orange dot, red and blue crosses indicates respectively mean estimate,
lower and upper 95% Confidence Intervals (See Table 1 for the definition of variables.). Effects are presented in the 'response'
scale, i.e. back-transformed from log-scale. For example, length of a reference fish without a kype (L_{ref-N}) would be: L_{ref-N} =
Sex N x L_{ref}.

Weak and contrasting evidence of changes in growth

Interactions between year and both river age and sea age were included in the best models,

which supports the hypothesis of temporal changes in the length-at-first return as a function

of life history strategy. The effect of the year on length-at-first return remained small and

relatively stable for 1FW, but increased slightly over time for 2FW and 3FW, both of which had

similar variations (Figure 4). Furthermore, length-at-sea-age over time differed, being stable

or slightly positive for 0SW and 1SW individuals, but decreasing for 2SW individuals. Overall,

except for 2SW individuals, trends remained weak, with only slight changes in the length-at-

age for 0SW and 1SW individuals.

331   **Discussion**

Results indicate that individuals returning to the Bresle River for the first time became pro-

gressively smaller over the 39 years of the study period. This trend mainly resulted from

change in the age structure and contrasted variations in length-at-age. In the study popula-

tion, an increasing percentage of sea trout returned for the first time at age 0SW and the

return migration timing advanced. This decrease in duration of the marine sojourn can explain

the decrease in body length, while length-at-age remained constant or slightly increased over

time for 0SW and 1SW individuals. At the same time, the percentage and length-at-age of 2SW

individuals decreased.

Strengths and limitations of long-term datasets

Sampling biases are inevitable during long-term monitoring of wild populations. It is often

challenging to maintain sampling effort constant over long periods, whether due to changes

in the sampling protocol, a lack of resources, modifications of the observation device or ob-

jectives, or varying environmental conditions. In this study, the need to control for sampling

biases required rebalancing the dataset. The long duration of data collection on the Bresle

River ensured that the dataset remained large and deep enough for exploratory analysis and

modelling. Using the bootstrap procedure enabled multiple iterations of subsampling and

modelling, which increased the stability and robustness of the results. Ultimately, this ap-

proach is a useful example of how such a long-term dataset can be analysed, despite the lat-

ter's inherent flaws. One downside of long term datasets is that the amount of data is likely

to be important which can lead to the rejection of the assumption of normality in the residu-

als. In GLMs, this may be improved with the use of non-Gaussian distributions, such as the

Gamma distribution, which in the present case did not greatly enhanced the results of nor-

mality tests. However, McCullagh and Nelder (1989) reported that this assumption was not of

major importance in the case of "large datasets", where small deviations from normality will

most likely lead to rejection of the assumption. Everything kept equal, large deviations in the

residuals suggests either errors in the age determination, or significantly higher or lower per-

formances of some individuals. Considering the very high within strategy length variation that

can be observed in sea trout, it is likely to be a major factor here. Despite these difficulties,

long-term datasets are crucial for understanding highly complex ecological and evolutionary

processes (Eliott and Eliott, 2006; Hughes et al., 2017).

An additional case study of decreasing average length of salmonids

With a decrease of *c.* 8% in sea trout body length over the past four decades, the results add

to the growing corpus of literature that describes a decrease in the length of salmonid species

around the world (Bal et al., 2017; Jeffrey et al., 2017; Lewis et al., 2015; Morita and Fukuwaka,

2007; Ohlberger et al., 2018; Oke et al., 2020). The extent of the changes observed in the

Bresle sea trout population agrees with results of Oke et al. (2020) for four species of Pacific

salmon (Chinook, *Oncorhynchus tshawytscha*: -8.0%, coho, *Oncorhynchus kisutch*: -3.3%,

chum, *Oncorhynchus keta*: -2.4%, and sockeye, *Oncorhynchus nerka*: -2.1%), but in contrast,

we found little support for major changes in the length-at-age. Consistently with our results,

structural changes in the age at return have previously been reported by Milner et al., (2017)

in sea trout. Change in the proportion of low-weight fish in rod catches of four Welsh rivers

suggested of an increasing proportion of 0SW between 1977 and 2007.

The importance of considering the entire life history

We found that river age had a significant effect on length-at-first return, which indicates that

the increase in length acquired in the river had long-term consequences until the first return

to the river. This effect could have major implications for individuals' future life history, such

as the timing of maturation and duration of the sea sojourn, as length can strongly influence

when smolt reach key thresholds that drive life history transitions (Bohlin et al., 1996; Hutch-

ings and Jones, 1998; Tréhin et al., 2021). Several studies have described mechanisms of com-

pensatory growth in brown trout (Fahy, 1978; Johnsson and Bohlin, 2006; Marco-Rius et al.,

2012) or other salmonid species (Al-Chokhachy et al., 2019; Mortensen and Damsgård, 1993;

Nikki et al., 2004). In these studies, shorter smolts exhibited higher subsequent growth at sea

than larger smolts and thus reached similar lengths at return. In a review of biological charac-

teristics of 72 populations of sea trout in the British Isles, Fahy (1978) found no correlation

between smolt length and adult length. However, it seems likely that slower growth rates

there did not provide sufficient resolution to do so, as there were large overlaps in length

among age classes. In contrast, faster growth rates in chalk streams such as the Bresle likely

enabled us to more easily detect an influence of the river on the first return. A study from a

nearby area and similar period (Pratten and Shearer, 1983) found that the mean length of the

most common age classes (2FW and 3FW) in the North Esk River, Scotland, differed by a mean

of 1.32 cm (SD = 0.36) from 1976-1980. In comparison, the mean length of smolts in the two

most common age classes in the Bresle (1FW and 2FW) differed by 3.94 cm (SD = 1.08) (un-

published data). In contrast, results of Marco-Rius et al. (2012) and Johnsson and Bohlin (2006)

do not seem to contradict ours. Marco-Rius et al. determined growth trajectories from scales,

but studied only a single age class (2FW) and excluded all individuals that had not spent an

entire year at sea (*i.e.* 0SW). Our results suggest that length differences are most visible in

0SW and tend to decrease as time at sea increases. Similarly, Johnsson & Bohlin (2006) con-

sidered only 1FW individuals, in which they induced length differences by depriving some of

them of food. While small differences within an age class can be compensated for within a

few months, this may not be the case for larger differences between age classes. The current

study also covers a longer period, more individuals, and focuses on fish that were not subject

to experimentally induced length differences.

Effect of migration timing on length-at-first return

Variations of the return date had little influence on length, despite major changes in sea-age

classes 0SW and 1SW, which arrived earlier by 53 and 47 days, respectively. This observation

is consistent with, but larger than, the advance of 2.6 days per decade observed by Legrand

et al. (2021), in the migration timing of sea trout at 40 monitoring stations in France. It is also

concordant with data from the River Dee, where length at return of 0SW individuals was found

to be independent from the day of return, with a complex non-linear pattern (Celtic Sea Trout

Project, 2016). Effects of a longer sea sojourn were positive and stronger for sea trout that

returned after 0SW, which is consistent with brown trout growth curves, which have steeper

slopes for younger age classes before reaching a plateau (Davaine et al., 1997). This dynamic

illustrates that younger fish benefit more from additional days at sea than older counterparts.

Contribution of growth opportunities and survival to age-dependent changes

It was surprising to detect a clear negative temporal trend in the length-at-age only for 2SW first returns, while those of 0SW and 1SW remained relatively stable. Marine trophic webs are under tremendous pressure in the context of climate change. The general consensus is that ocean productivity has decreased in the past few decades and that the composition of its eco-systems has changed greatly (Behrenfeld et al., 2006; Gohin et al., 2019; Hoegh-Guldberg and Bruno, 2010; Möllmann and Diekmann, 2012). One would thus expect to observe a decrease in growth also in younger age classes, as observed in Atlantic salmon (*Salmo salar*) (Tréhin et al., 2021). However, our results suggest that growth conditions for 0SW and 1SW may not have been greatly degraded. In fact, Davidson et al., (2006) even reported increased post-smolt growth for these two age-class on the River Dee. This differing response among sea-age classes could indicate local changes in the conditions that sea trout encountered at sea, rather than a broad decrease in growth opportunities. During migration at sea, salmonids gain access to different feeding grounds and sets of prey (Knutsen et al., 2004, 2001; Rikardsen et al., 2006, 2007; Thorstad et al., 2016). Consequently, changes in food availability and quality on the feeding grounds used by older and larger sea trout may explain their lower growth. Addi-tionally, local variations in sea temperatures could also influence sea trout return size, as re-ported before by Milner et al., (2017) in the Irish sea.

The age-specific decrease in length-at-first return may have been due to length-specific selec-tive pressure that targeted mainly larger individuals (Reznick et al., 1990), which could explain the progressive disappearance of 2SW individuals in the Bresle population. Overfishing has been widely demonstrated to influence many characteristics of fish populations, such as length at reproduction (Hard et al., 2008; Neuheimer and Taggart, 2010; Ojuok et al., 2007).

Intensive net fishing occurred at sea near the Bresle, especially during the first half of the study

period, with up to 37% of the population being caught each year either by professional or

leisure net fishing at sea or rod-and-line fishing in the river (Fagard and Beaulaton, 2018).

More recently, the resurgence of marine apex predators with specific feeding behaviours has

been associated with a decrease in the mean length of Chinook salmon (Ohlberger et al.,

2019). However, although the Bresle River is located near the Bay of Somme, which hosts the

largest breeding colony of harbour seals (*Phoca vitulina* L.) in the English Channel, analysis of

86 scat samples from this colony revealed no evidence of salmonids in the seals' diet (Spitz et

al., 2015). Furthermore, most seal predation on salmonids seems to be opportunistic, with no

indication of length-dependent selection for larger fish (Suuronen and Lehtonen, 2012;

Thomas et al., 2017). Therefore, it seems unlikely that seal predation would have driven the

population-wide changes in sea trout age structure observed in this study.

Parasitism by sea lice (*Lepeophtheirus salmonis*, Krøyer, 1837) has also been demonstrated to

be a major factor of reduced survival in wild salmonids (Bjørn et al., 2001; Thorstad et al.,

2015). Sea lice may have been responsible for the collapse of the sea trout population in the

Burrishoole River (Gargan et al., 2006) and for profound changes in life history traits of that in

the Eriff River, Ireland (Gargan et al., 2016). In the Bresle River, most returning sea trout show

evidence of sea-lice parasitism, but the youngest and smallest individuals usually have the

highest parasite loads. Premature returns, in which individuals with heavy sea-lice loads return

to freshwater sooner than individuals that are not infested, have been well documented in

the literature (Birkeland, 1996; Birkeland and Jakobsen, 1997; Serra-Llinares et al., 2020).

Thus, parasitism could provide an explanation for the increase in the percentage of 0SW ob-

served in this study.

Future research and implications for management

Although length-at-age changed little overall, a decrease in the age-at-first-return appears to

be the main reason of the decrease in the length-at-first return observed for sea trout in the

Bresle River. A change in the fitness-driven trade-off between survival and growth can change

the duration of the sea sojourn or the prevalence of migratory or resident ecotypes in trout

(Archer et al., 2019; Thorstad et al., 2016). Accordingly, future research should explore tem-

poral changes in growth at sea, which would entail analysing fine growth increments in scales.

For instance, increased growth opportunities could explain the shift toward shorter sea so-

journs, as fish could reach the suitable condition to mature earlier. On the other hand, reduced

growth at sea, potentially associated with reduced survival, could also lead to selective pres-

sure against long marine sojourn. Indeed, age at maturation has been demonstrated to be a

partially heritable life history trait in sea trout (Ferguson et al., 2019; Reed et al., 2019). Thus,

natural selection and change in allele frequency could potentially explain the progressive shift

toward shorter sea sojourn observed in this study.

The potential carry-over effect of a trout's freshwater life on its length-at-first return could be

relevant for salmonid management. Growth potential at sea is much higher than that in fresh-

water, accordingly the latter often seems to be viewed as a mere transitional habitat between

hatching and smoltification. Nevertheless, the hypothesis that a mechanistic length threshold

drives the probability of maturing (Hutchings and Jones, 1998) predicts that smolts that are

larger when entering the sea will reach the critical maturation length sooner and thus return

after fewer years at sea. Thus, the freshwater and marine phases of the sea trout life cycle

could both influence the life history changes observed during the return migration (Russell et

al., 2012). Consequently, management actions could be aimed at alleviating pressures on ju-

veniles, by mitigating the effects of climate change and improving freshwater habitats. These

actions should especially include managing and protecting water resources (Waco and Taylor,

2010), restoring habitat connectivity (Forget et al., 2018), and creating thermal refuges (Blann

et al., 2002).

Changes such as the progressive loss of a sea-age class, as observed for 2SW individuals in this

study, can strongly influence a population's resilience (Erkinaro et al., 2019; Stewart, 2011).

Reduced diversity of life history strategies could weaken the "portfolio" effect (Markowitz,

1952), which could increase the risk of population collapse (Figge, 2004; Schindler et al., 2015,

2010). The loss of older age classes can also have an especially detrimental effect due to the

loss of "big, old, fat, fecund, female fish" (BOFFFFs). These older female fish contribute greatly

to reproduction as they produce more and often larger eggs than younger females do (Hixon

et al., 2014; Ohlberger et al., 2020; Pope et al., 1961). Therefore, management actions such

as specific fishing regulations to protect these older fish could be implemented in the Bresle

River.

The lack of knowledge about the reproductive status of first-time returning individuals is a

potential disadvantage for future studies of population dynamics, especially for the role of the

0SW life history strategy. The return status of the individuals included in this study was deter-

mined from the analysis of patterns on scale reading. However, it provides an imperfect record

of previous reproduction history (Baglinière et al., 2020). Assessing whether 0SW returns are

mature individuals which will spawn the following winter or immature individuals which will

overwinter in the river Birkeland, 1996) would help understand consequences of their recent

relative increase on population dynamics. However, the percentage of the Bresle population

that does so is unknown. In contrast, a study of 0SW trout in the nearby Calonne River showed

that most summer 0SW and all autumn 0SW individuals were mature (Maisse et al., 1991),

which bring support to the hypothesis that the immature-return strategy has been uncommon

in rivers in Normandy.

We showed that change in the length-at-first return of the Bresle sea trout population was

explained mainly by a change in the mean sea age of first returns, which decreased over time.

Other factors also had an influence on the long-term change in the length of returns, such as

a small decrease in length-at-age, an advanced mean date of return, as well as the carry-over

effect of the river age. Further studies investigating sea trout life and growth at sea would be

needed to better understand the response of this complex species to ongoing environmental

changes.

530   **Acknowledgements**

531 Long-term datasets that support studies such as the present work would not exist without the

532 continued work and support of successive generations of field workers, technicians, research-

533 ers, and managers, and we thank them all. We thank Fabien Quendo for the making of Figure

534 1.

535   Data availability

536 Data analysed in this study are available on the GBIF repository at:

537 https://doi.org/10.15468/jkpya8

540   Conflict of interest disclosure

541 The authors declare that they comply with the PCI rule of having no financial conflicts of in-

542 terest in relation to the content of the article.

544   Funding statement

545 This study was funded by the Pole "Management of Diadromous Fish in their Environment,

546 OFB, INRAE, Institut Agro, UNIV PAU, and PAYS ADOUR/E2S UPPA".

Author information

Authors' contributions

Q.J.: conceptualization, data curation, formal analysis, investigation, methodology, valida-

tion, visualization, writing the original draft, writing the review and editing;

L.B.: conceptualization, funding acquisition, methodology, project administration, resources,

supervision, validation, visualization, writing – original draft, writing the review and editing;

A.R.: conceptualization, project administration, supervision, validation, visualization, writing

the review and editing;

M.N.: conceptualization, methodology, project administration, supervision, validation, visu-

alization, writing – original draft, writing the review and editing.

559   Author ORCIDs

560 Quentin Josset https://orcid.org/0009-0005-4533-8341
561 Laurent Beaulaton https://orcid.org/0000-0002-9614-8602
562 Atso Romakkaniemi https://orcid.org/0000-0001-9696-117X
563 Marie Nevoux https://orcid.org/0000-0003-1451-7732
564
565
566

References

Akaike, H., 1974. A new look at the statistical model identification. IEEE Transactions on Automatic Control 19, 716–723.

Al-Chokhachy, R., Kovach, R.P., Sepulveda, A., Strait, J., Shepard, B.B., Muhlfeld, C.C., 2019. Compensatory growth offsets poor condition in native trout populations. Freshwater Biology 64, 2120–2130. https://doi.org/10.1111/fwb.13400

Archer, L.C., Hutton, S.A., Harman, L., O'Grady, M.N., Kerry, J.P., Poole, W.R., Gargan, P., McGinnity, P., Reed, T.E., 2019. The interplay between extrinsic and intrinsic factors in determining migration decisions in brown trout (Salmo trutta): An experimental study. Frontiers in Ecology and Evolution 7, 222. https://doi.org/10.3389/fevo.2019.00222

Baglinière, J.-L., Hamelet, V., Guéraud, F., Aymes, J.-C., Goulon, C., Richard, A., Josset, Q., Marchand, F., 2020. Guide to the interpretation of the scales and the estimation of the age of brown trout (Salmo trutta) from the French populations., Guides and. ed.

Bal, G., Montorio, L., Rivot, E., Prévost, E., Baglinière, J.-L., Nevoux, M., 2017. Evidence for long-term change in length, mass and migration phenology of anadromous spawners in French Atlantic salmon Salmo salar. Journal of Fish Biology 90, 2375–2393. https://doi.org/10.1111/jfb.13314

Bartolino, V., Margonski, P., Lindegren, M., Linderholm, H.W., Cardinale, M., Rayner, D., Wennhage, H., Casini, M., 2014. Forecasting fish stock dynamics under climate change: Baltic herring (Clupea harengus) as a case study. Fisheries Oceanography 23, 258–269. https://doi.org/10.1111/fog.12060

Behrenfeld, M.J., O'Malley, R.T., Siegel, D.A., McClain, C.R., Sarmiento, J.L., Feldman, G.C., Milligan, A.J., Falkowski, P.G., Letelier, R.M., Boss, E.S., 2006. Climate-driven trends in contemporary ocean productivity. Nature 444, 752–755. https://doi.org/10.1038/nature05317

Birkeland, K., 1996. Consequences of premature return by sea trout (Salmo trutta) infested with the salmon louse (Lepeophtheirus salmonis Kroyer): migration, growth, and mortality. Canadian Journal of Fisheries and Aquatic Sciences 53, 2808–2813. https://doi.org/10.1139/f96-231

Birkeland, K., Jakobsen, P.J., 1997. Salmon lice, Lepeophtheirus salmonis, infestation as a causal agent of premature return to rivers and estuaries by sea trout, Salmo trutta, juveniles. Environmental Biology of Fishes 49, 129–137. https://doi.org/10.1023/A:1007354632039

Bjørn, P.A., Finstad, B., Kristoffersen, R., 2001. Salmon lice infection of wild sea trout and Arctic char in marine and freshwaters: The effects of salmon farms. Aquaculture Research 32, 947–962. https://doi.org/10.1046/j.1365-2109.2001.00627.x

Blackwell, B.G., Brown, M.L., Willis, D.W., 2000. Relative Weight (Wr) Status and Current Use in Fisheries Assessment and Management. Reviews in Fisheries Science 8, 1–44. https://doi.org/10.1080/10641260091129161

Blann, K., Frost Nerbonne, J., Vondracek, B., 2002. Relationship of Riparian Buffer Type to Water Temperature in the Driftless Area Ecoregion of Minnesota. North American Journal of Fisheries Management 22, 441–451. https://doi.org/10.1577/1548-8675(2002)022<0441:rorbtt>2.0.co;2

Bohlin, T., Dellefors, C., Faremo, U., 1996. Date of smolt migration depends on body-size but not age in wild sea-run brown trout. Journal of Fish Biology 49, 157–164. https://doi.org/10.1006/jfbi.1996.0143

Brown, C.J., Fulton, E.A., Hobday, A.J., Matear, R.J., Possingham, H.P., Bulman, C., Christensen,
V., Forrest, R.E., Gehrke, P.C., Gribble, N.A., Griffiths, S.P., Lozano-Montes, H., Martin,
J.M., Metcalf, S., Okey, T.A., Watson, R., Richardson, A.J., 2010. Effects of climate-
driven primary production change on marine food webs: Implications for fisheries and
conservation. Global Change Biology 16, 1194–1212. https://doi.org/10.1111/j.1365-
2486.2009.02046.x
Burnham, K.P., Anderson, D.R., Burnham, K.P., 2002. Model selection and multimodel infer-
ence: a practical information-theoretic approach, 2nd ed. ed. Springer, New York.
Butler, J.R.A., Walker, A.F., 2006. Characteristics of the Sea Trout Salmo trutta (L.) Stock Col-
lapse in the River Ewe (Wester Ross, Scotland), in 1988–2001, in: Harris, G., Milner, N.
(Eds.), Sea Trout: Biology, Conservation and Management. Blackwell Science Ltd, pp.
45–59. https://doi.org/10.1002/9780470996027.ch4
Canty, A., Ripley, B., 2021. boot: Bootstrap R (S-Plus) Functions. R package version 1.3-28.
Celtic Sea Trout Project, 2016. Celtic Sea Trout Project - Technical Report to Ireland Wales
Territorial Co-operation Programme 2007-2013 (INTERREG 4A. Inland Fisheries Ire-
land, Dublin.
Crozier, L.G., Hutchings, J. a., 2014. Plastic and evolutionary responses to climate change in
fish. Evolutionary Applications 7, 68–87. https://doi.org/10.1111/eva.12135
Davaine, P., Beall, E., Guerri, O., Caraguel, J.M., 1997. Introduction de salmonidés en milieu
vierge (îles kerguelen, subantarctique): Enjeux, résultats, perspectives. Bulletin Fran-
çais de la Pêche et de la Pisciculture 93–110. https://doi.org/10.1051/kmae:1997013
Davidson, I.C., Cove, R.J., Hazlewood, M.S., 2006. Annual Variation in Age Composition,
Growth and Abundance of Sea Trout Returning to the River Dee at Chester, 1991–
2003, in: Harris, G., Milner, N. (Eds.), Sea Trout: Biology, Conservation and Manage-
ment. Proceedings of the First International Sea Trout Symposium, Cardiff, July 2004.
Blackwell Publishing Ltd, pp. 76–87.
Davison, A., Hinkley, D., 1997. Bootstrap Methods and Their Applications. Cambridge Univer-
sity Press, Cambridge.
Dębowski, P., 2018. The largest Baltic population of sea trout (Salmo trutta L.): Its decline,
restoration attempts, and current status. Archives of Polish Fisheries 26, 81–100.
https://doi.org/10.2478/aopf-2018-0010
Dodson, J.J., Aubin-Horth, N., Thériault, V., Páez, D.J., 2013. The evolutionary ecology of alter-
native migratory tactics in salmonid fishes. Biological Reviews 88, 602–625.
https://doi.org/10.1111/brv.12019
Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber,
B., Lafourcade, B., Leitão, P.J., Münkemüller, T., Mcclean, C., Osborne, P.E., Reineking,
B., Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S., 2013. Collinearity: A review
of methods to deal with it and a simulation study evaluating their performance. Ecog-
raphy 36, 27–46. https://doi.org/10.1111/j.1600-0587.2012.07348.x
Eliott, J.M., Eliott, J.A., 2006. A 35-year study of stock-recruitment relationships in a small pop-
ulation of sea trout: assumptions, implications and Limitations for predicting targets.,
in: Harris, G., Milner, N. (Eds.), Sea Trout: Biology, Conservation and Management.
Blackwell Publishing, Oxford, pp. 257–278.
Erkinaro, J., Czorlich, Y., Orell, P., Kuusela, J., Falkegård, M., Länsman, M., Pulkkinen, H., Prim-
mer, C.R., Niemelä, E., 2019. Life history variation across four decades in a diverse pop-
ulation complex of atlantic salmon in a large subarctic river. Canadian Journal of Fish-
eries and Aquatic Sciences 76, 42–55. https://doi.org/10.1139/cjfas-2017-0343

Euzenat, G., Fournel, F., Fagard, J.L., 2007. Population Dynamics and Stock-Recruitment Rela-
tionship of Sea Trout in the River Bresle, Upper Normandy, France. Sea Trout: Biology,
Conservation   and   Management   307–323.
https://doi.org/10.1002/9780470996027.ch20
Fagard, J.-L., Beaulaton, L., 2018. Éléments sur l'exploitation par pêche des salmonidés migra-
teurs en zone côtière et en rivière depuis 1978. Littoral proche de la rivière Bresle,
embouchures de cours d'eau de Seine-Maritime. Cours d'eau à salmonidés migrateurs
au nord de la Seine. Agence Française pour la Biodiversité.
Fahy, E., 1978. Variation in some biological characteristics of British sea trout, Salmo trutta L.
Journal   of   Fish   Biology   13,   123–138.   https://doi.org/10.1111/j.1095-
8649.1978.tb03420.x
Ferguson, A., Reed, T.E., Cross, T.F., McGinnity, P., Prodöhl, P.A., 2019. Anadromy, po-
tamodromy and residency in brown trout Salmo trutta: the role of genes and the en-
vironment. Journal of Fish Biology. https://doi.org/10.1111/jfb.14005
Figge, F., 2004. Bio-folio: applying portfolio thoery to biodiversity. Biodive 13, 827–849.
Forget, G., Baglinière, J.L., Marchand, F., Richard, A., Nevoux, M., Durif, C., 2018. A new
method to estimate habitat potential for Atlantic salmon (Salmo salar): Predicting the
influence of dam removal on the Sélune River (France) as a case study. ICES Journal of
Marine Science 75, 2172–2181. https://doi.org/10.1093/icesjms/fsy089
Forseth, T., Barlaup, B.T., Finstad, B., Fiske, P., Gjøsæter, H., Falkegård, M., Hindar, A., Mo,
T.A., Rikardsen, A.H., Thorstad, E.B., Vøllestad, L.A., Wennevik, V., 2017. The major
threats to Atlantic salmon in Norway. ICES Journal of Marine Science 74, 1496–1513.
https://doi.org/10.1093/icesjms/fsx020
Gargan, P., Kelly, F., Shephard, S., Whelan, K., 2016. Temporal variation in sea trout Salmo
trutta life history traits in the Erriff River, western Ireland. Aquaculture Environment
Interactions 8, 675–689. https://doi.org/10.3354/aei00211
Gargan, P.G., Poole, W.R., Forde, G.P., 2006. A review of the status of irish sea trout stocks.,
in: Harris, G., Milner, N. (Eds.), Sea Trout: Biology, Conservation and Management.
Blackwell Science Ltd, pp. 25–44. https://doi.org/10.1002/9780470996027.ch3
Gohin, F., Van der Zande, D., Tilstone, G., Eleveld, M.A., Lefebvre, A., Andrieux-Loyer, F.,
Blauw, A.N., Bryère, P., Devreker, D., Garnesson, P., Hernández Fariñas, T., Lamaury,
Y., Lampert, L., Lavigne, H., Menet-Nedelec, F., Pardo, S., Saulquin, B., 2019. Twenty
years of satellite and in situ observations of surface chlorophyll-a from the northern
Bay of Biscay to the eastern English Channel. Is the water quality improving? Remote
Sensing of Environment 233, 111343. https://doi.org/10.1016/j.rse.2019.111343
Gross, M.R., Coleman, R.M., McDowall, R.M., 1988. Aquatic productivity and the evolution of
diadromous fish migration. Science 239, 1291–1293. https://doi.org/10.1126/sci-
ence.239.4845.1291
Hard, J.J., Gross, M.R., Heino, M., Hilborn, R., Kope, R.G., Law, R., Reynolds, J.D., 2008. Evolu-
tionary consequences of fishing and their implications for salmon. Evolutionary Appli-
cations 1, 388–408. https://doi.org/10.1111/j.1752-4571.2008.00020.x
Helle, J., Martinson, E., Eggers, D., 2007. Influence of salmon abundance and ocean conditions
on body size of Pacific salmon. N. Pac. Anadr. Fish 289–298.
Hixon, M. a, Johnson, D.W., Sogard, S.M., 2014. BOFFFFs: on the importance of conserving
old-growth age structure in fishery populations. ICES Journal of Marine Science 71,
2171–2185. https://doi.org/10.1093/icesjms/fst200

Hoegh-Guldberg, O., Bruno, J.F., 2010. The impact of climate change on the world's marine
ecosystems. Science 328, 1523–1528. https://doi.org/10.1126/science.1189930
Hughes, B.B., Beas-Luna, R., Barner, A.K., Brewitt, K., Brumbaugh, D.R., Cerny-Chipman, E.B.,
Close, S.L., Coblentz, K.E., De Nesnera, K.L., Drobnitch, S.T., Figurski, J.D., Focht, B.,
Friedman, M., Freiwald, J., Heady, K.K., Heady, W.N., Hettinger, A., Johnson, A., Karr,
K.A., Mahoney, B., Moritsch, M.M., Osterback, A.M.K., Reimer, J., Robinson, J., Rohrer,
T., Rose, J.M., Sabal, M., Segui, L.M., Shen, C., Sullivan, J., Zuercher, R., Raimondi, P.T.,
Menge, B.A., Grorud-Colvert, K., Novak, M., Carr, M.H., 2017. Long-Term studies con-
tribute disproportionately to ecology and policy. BioScience 67, 271–278.
https://doi.org/10.1093/biosci/biw185
Hutchings, J. a, Jones, M.E.B., 1998. Life history variation and growth rate thresholds for ma-
turity in Atlantic salmon, Salmo salar. Canadian Journal of Fisheries and Aquatic Sci-
ences 55 (Suppl., 22–47. https://doi.org/10.1139/cjfas-55-S1-22
ICES, 2020. Working Group with the Aim to Develop Assessment Models and Establish Biolog-
ical Refer-ence Points for Sea Trout (Anadromous Salmo trutta) Populations
(WGTRUTTA; outputs from 2019 meeting). ICES Scientific Reports.
https://doi.org/10.17895/ices.pub.7431
Ishida, Y., Ito, S., Kaeriyama, M., McKinnell, S., Nagasawa, K., 1993. Recent changes in age and
size of chum salmon (Oncorhynchus keta) in the North Pacific Ocean and possible
causes. Canadian Journal of Fisheries and Aquatic Sciences 50, 290–295.
https://doi.org/10.1139/f93-033
Järvi, T.H., 1940. Sea trout in the Bothnian Bay. Acta Zoologica Fennica 1–29.
Jeffrey, K.M., Côté, I.M., Irvine, J.R., Reynolds, J.D., 2017. Changes in body size of Canadian
Pacific salmon over six decades. Canadian Journal of Fisheries and Aquatic Sciences 74,
191–201. https://doi.org/10.1139/cjfas-2015-0600
Johnsson, J.I., Bohlin, T., 2006. The cost of catching up: increased winter mortality following
structural growth compensation in the wild. Proceedings. Biological sciences / The
Royal Society 273, 1281–1286. https://doi.org/10.1098/rspb.2005.3437
Jonsson, B., L'Abée-Lund, J.H., 1993. Latitudinal clines in life-history variables of anadromous
brown trout in Europe. Journal of Fish Biology 43, 1–16.
https://doi.org/10.1111/j.1095-8649.1993.tb01175.x
Josset, Q., Flesselle, A., Bernardin, A., Macquet, T., Petit, L., Robin, E., Le Bras, R., Fagard, J.-L.,
Fournel, F., Euzenat, G., Beaulaton, L., Prévost, E., Marchand, F., 2023. Phenology and
biological traits of migrating trout and sea trout (Salmo trutta) sampled by trapping in
the survey in the Bresle river (France).
Kendall, N.W., Quinn, T.P., 2011. Length and age trends of Chinook salmon in the Nushagak
River, Alaska, related to commercial and recreational fishery selection and exploita-
tion. Transactions of the American Fisheries Society 140, 611–622.
https://doi.org/10.1080/00028487.2011.585575
Kennedy, R.J., Crozier, W.W., 2010. Evidence of changing migratory patterns of wild Atlantic
salmon Salmo salar smolts in the River Bush, Northern Ireland, and possible associa-
tions with climate change. Journal of Fish Biology 76, 1786–1805.
https://doi.org/10.1111/j.1095-8649.2010.02617.x
Knutsen, J.A., Knutsen, H., Gjøsæter, J., Jonsson, B., 2001. Food of anadromous brown trout
at sea. Journal of Fish Biology 59, 533–543. https://doi.org/10.1006/jfbi.2001.1662

Knutsen, J.A., Knutsen, H., Olsen, E.M., Jonsson, B., 2004. Marine feeding of anadromous
Salmo trutta during winter. Journal of Fish Biology 64, 89–99.
https://doi.org/10.1046/j.1095-8649.2003.00285.x
Law, R., 2000. Fishing, selection, and phenotypic evolution. ICES Journal of Marine Science 57,
659–668. https://doi.org/10.1006/jmsc.2000.0731
Legrand, M., Briand, C., Buisson, L., Besse, T., Artur, G., Azam, D., Baisez, A., Barracou, D.,
Bourré, N., Carry, L., Caudal, A.L., Corre, J., Croguennec, E., Der Mikaélian, S., Josset,
Q., Le Gurun, L., Schaeffer, F., Toussaint, R., Laffaille, P., 2021. Diadromous fish modi-
fied timing of upstream migration over the last 30 years in France. Freshwater Biology
66, 286–302. https://doi.org/10.1111/fwb.13638
Lewis, B., Grant, W.S., Brenner, R.E., Hamazaki, T., 2015. Changes in size and age of chinook
salmon oncorhynchus tshawytscha returning to Alaska. PLoS ONE 10, 1–17.
https://doi.org/10.1371/journal.pone.0130184
Maisse, G., Mourot, B., Breton, B., Fostier, A., Marcuzzi, O., Le Bail, P.Y., Baglinière, J.L., Ri-
chard, A., 1991. Sexual maturity in sea trout, Salmo trutta L., running up the river
Calonne (Normandy, France) at the "finnock" stage. Journal of Fish Biology 39, 705–
715. https://doi.org/10.1111/j.1095-8649.1991.tb04400.x
Marchand, F., Aymes, J.-C., Gueraud, F., Guillard, J., Goulon, C., Hamelet, V., Lange, F., Prévost,
E., Baglinière, J.L., Beaulaton, L., Penil, C., Azam, D., 2019. Colisa, the collection of ichty-
ological samples. https://doi.org/10.15454/D3ODJM
Marco-Rius, F., Caballero, P., Morán, P., Garcia de Leaniz, C., 2012. And the Last Shall Be First:
Heterochrony and Compensatory Marine Growth in Sea Trout (Salmo trutta). PLoS
ONE 7, 1–10. https://doi.org/10.1371/journal.pone.0045528
Markowitz, H.M., 1952. Portfolio selection. Journal of Finance 7, 77–91.
McCullagh, P., Nelder, J.A., 1989. Generalized Linear Models, Second Edition., Chapman&Hall.
ed, Monographs on Statistics and Applied Probability.
McKinney, G.J., Hale, M.C., Goetz, G., Gribskov, M., Thrower, F.P., Nichols, K.M., 2015. Onto-
genetic changes in embryonic and brain gene expression in progeny produced from
migratory and resident Oncorhynchus mykiss. Molecular Ecology 24, 1792–1809.
https://doi.org/10.1111/mec.13143
Milner, N., Potter, E., Roche, W., Tysklin, N., Davidson, I.C., King, J., Coyne, J., Davies, C., 2017.
Variation in sea trout (Salmo trutta) abundance and life histories in the Irish Sea., in:
Graeme Harris. Ed. (Ed.), Sea Trout: Science & Management. Proceedings of the 2nd
International Sea Trout Symposium, October 2015, Dundalk, Ireland.
Möllmann, C., Diekmann, R., 2012. Marine Ecosystem Regime Shifts Induced by Climate and
Overfishing. A Review for the Northern Hemisphere, Advances in Ecological Research.
https://doi.org/10.1016/B978-0-12-398315-2.00004-1
Morita, K., Fukuwaka, M.A., 2007. Why age and size at maturity have changed in Pacific
salmon. Marine Ecology Progress Series 335, 289–294.
https://doi.org/10.3354/meps335289
Mortensen, A., Damsgård, B., 1993. Compensatory growth and weight segregation following
light and temperature manipulation of juvenile Atlantic salmon (Salmo salar L.) and
Arctic charr (Salvelinus alpinus L.). Aquaculture 114, 261–272.
https://doi.org/10.1016/0044-8486(93)90301-E
Neuheimer, A.B., Taggart, C.T., 2010. Can changes in length-at-age and maturation timing in
Scotian Shelf haddock (Melanogrammus aeglefinus) be explained by fishing? Canadian

Journal of Fisheries and Aquatic Sciences 67, 854–865. https://doi.org/10.1139/F10-
025

Nevoux, M., Finstad, B., Davidsen, J.G., Finlay, R., Josset, Q., Poole, R., Höjesjö, J., Aarestrup,
K., Persson, L., Tolvanen, O., Jonsson, B., 2019. Environmental influences on life history
strategies in partially anadromous brown trout (Salmo trutta , Salmonidae). Fish and
Fisheries 1–32. https://doi.org/10.1111/faf.12396

Niiranen, S., Yletyinen, J., Tomczak, M.T., Blenckner, T., Hjerne, O., Mackenzie, B.R., Müller-
Karulis, B., Neumann, T., Meier, H.E.M., 2013. Combined effects of global climate
change and regional ecosystem drivers on an exploited marine food web. Global
Change Biology 19, 3327–3342. https://doi.org/10.1111/gcb.12309

Nikki, J., Pirhonen, J., Jobling, M., Karjalainen, J., 2004. Compensatory growth in juvenile rain-
bow trout, Oncorhynchus mykiss (Walbaum), held individually. Aquaculture 235, 285–
296. https://doi.org/10.1016/j.aquaculture.2003.10.017

Ohlberger, J., Schindler, D.E., Brown, R.J., Harding, J.M.S., Adkison, M.D., Munro, A.R., Horst-
mann, L., Spaeder, J., 2020. The reproductive value of large females: Consequences of
shifts in demographic structure for population reproductive potential in Chinook
Salmon. Canadian Journal of Fisheries and Aquatic Sciences 77, 1292–1301.
https://doi.org/10.1139/cjfas-2020-0012

Ohlberger, J., Schindler, D.E., Ward, E.J., Walsworth, T.E., Essington, T.E., 2019. Resurgence of
an apex marine predator and the decline in prey body size. Proceedings of the National
Academy of Sciences of the United States of America 116, 26682–26689.
https://doi.org/10.1073/pnas.1910930116

Ohlberger, J., Ward, E.J., Schindler, D.E., Lewis, B., 2018. Demographic changes in Chinook
salmon across the Northeast Pacific Ocean. Fish and Fisheries 19, 533–546.
https://doi.org/10.1111/faf.12272

Ojuok, J.E., Njiru, M., Ntiba, M.J., Mavuti, K.M., 2007. The effect of overfishing on the life-
history strategies of Nile tilapia, Oreochromis niloticus (L.) in the Nyanza Gulf of Lake
Victoria, Kenya. Aquatic Ecosystem Health and Management 10, 443–448.
https://doi.org/10.1080/14634980701708107

Oke, K.B., Cunningham, C.J., Westley, P.A.H., Baskett, M.L., Carlson, S.M., Clark, J., Hendry,
A.P., Karatayev, V.A., Kendall, N.W., Kibele, J., Kindsvater, H.K., Kobayashi, K.M., Lewis,
B., Munch, S., Reynolds, J.D., Vick, G.K., Palkovacs, E.P., 2020. Recent declines in
salmon body size impact ecosystems and fisheries. Nature Communications 11, 1–13.
https://doi.org/10.1038/s41467-020-17726-z

Perez, K.O., Munch, S.B., 2010. Extreme selection on size in the early lives of fish. Evolution
64, 2450–2457. https://doi.org/10.1111/j.1558-5646.2010.00994.x

Perrier, C., Guyomard, R., Bagliniere, J.L., Nikolic, N., Evanno, G., 2013. Changes in the genetic
structure of Atlantic salmon populations over four decades reveal substantial impacts
of stocking and potential resiliency. Ecology and Evolution 3, 2334–2349.
https://doi.org/10.1002/ece3.629

Pope, J.A., Mills, D.H., Shearer, W.M., 1961. The fecundity of Atlantic Salmon (Salmon salar)
Linn. Freshwater and Salmon Fisheries Research 26, 1–12.

Pratten, D.J., Shearer, W.M., 1983. The migrations of North Esk sea trout. Fisheries Manage-
ment 14, 99–113. https://doi.org/10.1111/j.1365-2109.1983.tb00056.x

Quinn, T.P., McGinnity, P., Reed, T.E., 2016. The paradox of "premature migration" by adult
anadromous salmonid fishes: Patterns and hypotheses. Canadian Journal of Fisheries
and Aquatic Sciences 73, 1015–1030. https://doi.org/10.1139/cjfas-2015-0345

R Core Team, 2020. R: A language and environment for statistical computing.

Reed, T.E., Prodöhl, P., Bradley, C., Gilbey, J., McGinnity, P., Primmer, C.R., Bacon, P.J., 2019.
Heritability estimation via molecular pedigree reconstruction in a wild fish population
reveals substantial evolutionary potential for sea age at maturity, but not size within
age classes. Can. J. Fish. Aquat. Sci. 76, 790–805. https://doi.org/10.1139/cjfas-2018-
0123

Reznick, D.A., Bryga, H., Endler, J.A., 1990. Experimentally induced life-history evolution in a
natural population. Nature 346, 357–359. https://doi.org/10.1038/346357a0

Richard, A., 1981. Observations préliminaires sur les populations de truite de mer (Salmo
trutta L.) en Basse-Normandie. Bulletin Français de la Pêche et de la Pisciculture 283,
114–124.

Rikardsen, a. H., Amundsen, P. a., Knudsen, R., Sandring, S., 2006. Seasonal marine feeding
and body condition of sea trout (Salmo trutta) at its northern distribution. ICES Journal
of Marine Science 63, 466–475. https://doi.org/10.1016/j.icesjms.2005.07.013

Rikardsen, A.H., Dempson, J.B., Amundsen, P.-A., Bjorn, P.A., Finstad, B., Jensen, A.J., 2007.
Temporal variability in marine feeding of sympatric Arctic charr and sea trout. Journal
of Fish Biology 70, 837–852. https://doi.org/10.1111/j.1095-8649.2007.01345.x

Russell, I.C., Aprahamian, M.W., Barry, J., Davidson, I.C., Fiske, P., Ibbotson, A.T., Kennedy, R.J.,
Maclean, J.C., Moore, A., Otero, J., Potter, T. (E. C.E.), Todd, C.D., 2012. The influence
of the freshwater environment and the biological characteristics of Atlantic salmon
smolts on their subsequent marine survival. ICES Journal of Marine Science 69, 1563–
1573. https://doi.org/10.1093/icesjms/fsr208

Satterthwaite, W.H., Mohr, M.S., O'Farrell, M.R., Wells, B.K., 2012. A Bayesian hierarchical
model of size-at-age in ocean-harvested stocks-quantifying effects of climate and tem-
poral variability. Canadian Journal of Fisheries and Aquatic Sciences 69, 942–954.
https://doi.org/10.1139/F2012-036

Schindler, D.E., Armstrong, J.B., Reed, T.E., 2015. The portfolio concept in ecology and evolu-
tion. Frontiers in Ecology and the Environment 13, 257–263.
https://doi.org/10.1890/140275

Schindler, D.E., Hilborn, R., Chasco, B., Boatright, C.P., Quinn, T.P., Rogers, L.A., Webster, M.S.,
2010. Population diversity and the portfolio effect in an exploited species. Nature 465,
609–612. https://doi.org/10.1038/nature09060

Serra-Llinares, R.M., Bøhn, T., Karlsen, Nilsen, R., Freitas, C., Albretsen, J., Haraldstad, T., Thor-
stad, E.B., Elvik, K.M.S., Bjørn, P.A., 2020. Impacts of salmon lice on mortality, marine
migration distance and premature return in sea trout. Marine Ecology Progress Series
635, 151–168. https://doi.org/10.3354/MEPS13199

Spitz, J., Dupuis, L., Becquet, V., Dubief, B., Trites, A.W., 2015. Diet of the harbour seal Phoca
vitulina: Implication for the flatfish nursery in the Bay of Somme (English Channel,
France). Aquatic Living Resources 28, 11–19. https://doi.org/10.1051/alr/2015001

Stewart, J., 2011. Evidence of age-class truncation in some exploited marine fish populations
in New South Wales, Australia. Fisheries Research 108, 209–213.
https://doi.org/10.1016/j.fishres.2010.11.017

Suuronen, P., Lehtonen, E., 2012. The role of salmonids in the diet of grey and ringed seals in
the Bothnian Bay, northern Baltic Sea. Fisheries Research 125–126, 283–288.
https://doi.org/10.1016/j.fishres.2012.03.007

Thomas, A.C., Nelson, B.W., Lance, M.M., Deagle, B.E., Trites, A.W., 2017. Harbour seals target
juvenile salmon of conservation concern. Canadian Journal of Fisheries and Aquatic
Sciences 74, 907–921. https://doi.org/10.1139/cjfas-2015-0558

Thomsen, D.S., Koed, A., Nielsen, C., Madsen, S.S., 2007. Overwintering of sea trout (Salmo
trutta) in freshwater: Escaping salt and low temperature or an alternate life strategy?
Canadian Journal of Fisheries and Aquatic Sciences 64, 793–802.
https://doi.org/10.1139/F07-059

Thorstad, E.B., Todd, C.D., Uglem, I., Bjørn, P.A., Gargan, P.G., Vollset, K.W., Halttunen, E.,
Kålås, S., Berg, M., Finstad, B., 2016. Marine life of the sea trout. Marine Biology 163,
1–19. https://doi.org/10.1007/s00227-016-2820-3

Thorstad, E.B., Todd, C.D., Uglem, I., Bjørn, P.A., Gargan, P.G., Vollset, K.W., Halttunen, E.,
Kålås, S., Berg, M., Finstad, B., 2015. Effects of salmon lice Lepeophtheirus salmonis on
wild sea trout salmo trutta - A literature review. Aquaculture Environment Interactions
7, 91–113. https://doi.org/10.3354/aei00142

Tréhin, C., Rivot, E., Lamireau, L., Meslier, L., Besnard, A., Gregory, Stephen, D., Nevoux, M.,
2021. Growth during the first summer at sea modulates sex-specific maturation sched-
ule in Atlantic salmon. Canadian Journal of Fisheries and Aquatic Sciences 1–40.
https://doi.org/10.1139/cjfas-2020-0236

Waco, K.E., Taylor, W.W., 2010. The influence of groundwater withdrawal and land use
changes on brook charr (Salvelinus fontinalis) thermal habitat in two coldwater tribu-
taries in Michigan, U.S.A. Hydrobiologia 650, 101–116.
https://doi.org/10.1007/s10750-010-0204-0

Walsh, M.R., Reznick, D.N., 2011. Experimentally induced life-history evolution in a killifish in
response to the introduction of guppies. Evolution 65, 1021–1036.
https://doi.org/10.1111/j.1558-5646.2010.01188.x

Witten, P.E., Hall, B.K., 2001. The salmon kype. How does it grow? What's its purpose? Does
it disappear after spawning? Atlantic Salmon Journal 36–39.