Bacillus velezensis HBXN2020 alleviates *Salmonella* Typhimurium infection in mice by improving intestinal barrier integrity and reducing inflammation

Linkang Wang,[1,2,3] Haiyan Wang,[1,2,3] Xinxin Li,[1,2,3] Mengyuan Zhu,[1,2,3] Dongyang Gao,[1,2,3] Dayue Hu,[1,2,3] Zhixuan Xiong,[1,2,3] Xiangmin Li[1,2,3*] and Ping Qian[1,2,3*]

[1]*National Key Laboratory of Agricultural Microbiology, Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan, Hubei, 430070, People's Republic of China.*

[2]*College of Veterinary Medicine, Huazhong Agricultural University, Wuhan, Hubei, People's Republic of China.*

[3]*Key Laboratory of Preventive Veterinary Medicine in Hubei Province, the Cooperative Innovation Center for Sustainable Pig Production, Wuhan, Hubei, People's Republic of China.*

* Corresponding author: Ping Qian: Tel: 86-27-87282608; Fax: 86-27-87282608; E-mail: qianp@mail.hzau.edu.cn. Address: No. 1, Shizishan Street, Hongshan District, Wuhan City, Hubei Province, China.

* Co-corresponding author: Xiang-min Li: Tel: 86-27-87282608; Fax: 86-27-87282608; E-mail: lixiangmin@mail.hzau.edu.cn. Address: No. 1, Shizishan

Street, Hongshan District, Wuhan City, Hubei Province, China.

Abstract

Bacillus velezensis is a species of *Bacillus* that has been widely

investigated because of its broad-spectrum antimicrobial activity. However,

most studies on *Bacillus velezensis* have focused on the biocontrol of plant

diseases, with few reports on antagonizing *Salmonella* Typhimurium infections.

In this investigation, it was discovered that *Bacillus velezensis* HBXN2020,

which was isolated from healthy black pigs, possessed strong anti-stress and

broad-spectrum antibacterial activity. Importantly, *Bacillus velezensis*

HBXN2020 did not cause any adverse side effects in mice when administered

at various doses (1×10^7, 1×10^8, and 1×10^9 CFU) for 14 d. Supplementing

Bacillus velezensis HBXN2020 spores, either as a curative or preventive

measure, dramatically reduced the levels of *Salmonella* Typhimurium

ATCC14028 in the mice's feces, ileum, cecum, and colon, as well as the

disease activity index (DAI), in a model of infection caused by this pathogen in

mice. Additionally, supplementing *Bacillus velezensis* HBXN2020 spores

significantly regulated cytokine levels (TNF-α, IL-1β, IL-6, and IL-10) and

maintained the expression of tight junction proteins and mucin protein. Most

importantly, adding *Bacillus velezensis* HBXN2020 spores to the colonic

microbiota improved its stability and increased the amount of beneficial

bacteria (*Lactobacillus* and *Akkermansia*). All together, *Bacillus velezensis*

HBXN2020 can improve intestinal microbiota stability and barrier integrity and

reduce inflammation to help treat infection by *Salmonella* Typhimurium.

Key words: *Bacillus velezensis*; antibacterial activity; *S.* Typhimurium

infection; infections; gut health

Introduction

Salmonella Typhimurium (*S.* Typhimurium) is a major foodborne zoonotic

pathogen that can cause diarrhea and colitis in humans and animals (Fabrega

and Vila, 2013; Yang et al., 2021). According to reports, *Salmonella* infections

are common in both industrialized and developing nations, posing a serious

risk to public health and resulting in significant financial losses (Katiyo et al.,

2019; Mohan et al., 2019). At present, antibiotics remain one of the most

effective treatment strategies for *Salmonella* infections. However, numerous

studies have reported that the prolonged use or misuse of antibiotics can lead

to environmental pollution, an increase in multi-drug-resistant (MDR) bacteria,

as well as gastrointestinal microbiota dysbiosis (Ferri et al., 2017; Paulson et

al., 2015; Reyman et al., 2022). In order to tackle *Salmonella* infections, a

hunt for novel antimicrobial drugs is now underway.

More research has recently demonstrated the critical role gut microbiota

plays in reducing intestinal inflammation (Cristofori et al., 2021) and mending

the intestinal barrier (Ohland and Macnaughton, 2010; Wang et al., 2020).

The use of probiotics is a popular approach to modulating intestinal microbiota

nowadays. Among all probiotics, *Bacillus* is one of the most popular probiotic

species because of its ability to form endospores that survive gastric transit,

storage and delivery conditions (Gu et al., 2015). Furthermore, *Bacillus*

species can generate extracellular enzymes and antimicrobial metabolites

that inhibit enteric pathogens, which lowers the risk of infection and enhances

nutrient utilization (Abdhul et al., 2015; Ouattara et al., 2017).

Recent studies have found that *Bacillus velezensis* (*B. velezensis*) has the

enormous potential to produce a variety of metabolites with broad-spectrum

antibacterial activity (Ye et al., 2018). Meanwhile, previous studies have

reported that *B. velezensis* exhibits varying degrees of beneficial effects on

plants, livestock, poultry and fish. For instance, the combined use of *B.*

velezensis SQR9 mutant and FZB42 improved root colonization and the

production of secondary metabolites that are beneficial to cucumber (Shao et

al., 2022). *B. velezensis* isolated from yaks was shown to enhance growth

performance and ameliorate blood parameters related to inflammation and

immunity in mice (Li et al., 2019). The dietary *B. velezensis* supplementation

can regulate the innate immune response in the intestine of crucian carp and

reduce the degree of intestinal inflammation damage induced by *A. veronii*

(Zhang et al., 2019). However, studies on *B. velezensis* in the prevention of *S.*

Typhimurium infection are rarely reported; instead, the majority of

investigations on *B. velezensis* focused on the biocontrol of fungal infections

in plants. In this study, a strain of *B. velezensis* HBXN2020 with broad-

spectrum antibacterial activity against *Salmonella* was selected from a vast number of *Bacillus* strains. Next, we evaluated its safety in mice by supplementing *B. velezensis* HBXN2020 and then explored the protective effects in STm-infected mice. Notably, *B. velezensis* HBXN2020 alleviated colon tissue damage caused by STm infection, as indicated by markers such as STm loads, TNF-α and ZO-1 levels. Moreover, supplementing *B. velezensis* HBXN2020 also increased the abundance of beneficial bacteria, specifically *Lactobacillus* and *Akkermansia*, within the colon microbiota. As a result, this research supports the creation of probiotic-based microbial products as a substitute method of preventing *Salmonella* infections.

Results

Isolation of *Bacillus*

Four strains of *Bacillus* with antibacterial activity were screened from a large amount of presumptive *Bacillus* isolates, based on spot-on-plate tests analysis (Fig. S1), and the four strains clearly displayed the properties of bacteria in the genus *Bacillus* such as colonies were crateriform, Gram-positive, rod shape, and endospore-forming ability. Next, a comparative analysis of the antibacterial spectrum of four strains of *Bacillus* found that one strain exhibited excellent antibacterial activity against common pathogens (Table S1). Therefore, this strain was selected in this study for further research, which was designated as the HBXN2020.

Growth curve and *in vitro* resistance against environmental assaults

The growth of HBXN2020 was assessed in flat-bottomed 100-well microtiter plates by measuring the values of OD_{600} every hour using an automatic growth curve analyzer. After measurement, we found that HBXN2020 entered the logarithmic growth phase after 2 h, and reached a plateau after 10 h of culture (Fig. 1A).

Next, we evaluated the effect of physical, chemical, and biological sterilization conditions such as high temperature, strong acidity, and enzyme preparation on the survival of HBXN2020. The survival rate of HBXN2020 showed a decreasing trend with increasing temperature (Fig. 1B), but this decrease was not obvious. In a strong acid environment (pH 2.0), HBXN2020 maintained a high survival rate (Fig. 1C), suggesting that HBXN2020 spores can survive under extreme conditions. Based on these results, we hypothesized that HBXN2020 spores might also exhibit improved survival in gastrointestinal tract environments. Therefore, we further evaluated the survival rate of HBXN2020 in bile salt (0.3%, v/v), simulated gastric fluid (pepsin 1 mg mL^{-1}, pH 1.2), and simulated intestinal fluid (trypsin 1 mg mL^{-1}, pH 6.8). As shown in Fig. 1D-F, HBXN2020 spores demonstrated significantly improved tolerance to these simulated gastrointestinal tract environments.

Antibiotic susceptibility of HBXN2020 and bacteriostasis effect *in vitro*

In order to assess HBXN2020's drug resistance, 19 antibiotics that are frequently used in clinical settings were chosen in this study. The results indicated that only polymyxin B had a relatively small inhibition zone diameter (less than 15 mm). Ampicillin, meropenem, minocycline, ofloxacin, and trimethoprim had the strongest inhibition on HBXN2020, with an inhibition zone diameter exceeding 30 mm (Fig. 2A). This indicated that HBXN2020 was extremely sensitive to β-lactams, tetracyclines, and quinolone drugs. We next utilized an in vitro antibacterial assay to evaluate the antibacterial activity of HBXN2020. The results showed that HBXN2020 had a similar inhibitory effect on standard and wild strains of *Escherichia coli* (*E. coli*), *Salmonella*, *Staphylococcus aureus* (*S. aureus*), and *Clostridium perfringens* (*C. perfringens*), as well as wild strains of *Streptococcus suis* (*S. suis*), *Pasteurella multocida* (*P. multocida*), and *Actinobacillus pleuropneumoniae* (*A. pleuropneumoniae*). Except for the wild strains of *S. suis* and *A. pleuropneumoniae*, the diameter of the inhibition zone of other strains was above 15 mm (Fig. 2B, S2), while the size of the inhibition zone of *S. suis* and *A. pleuropneumoniae* was also more than 12 mm.

Genomic Characteristics

HBXN2020's complete genome was sequenced using the Illumina Hiseq and PacBio platforms. The results showed that HBXN2020 has a circular chromosome of 3,929,792 bp (Fig. 3A) with a GC content of 46.5%, including

3,744 protein-coding genes (CDS), 86 tRNA genes, and 27 rRNA genes (Table S2). Furthermore, the HBXN2020 genome also contains 13 different clusters of secondary metabolic synthesis genes. such as Fengycin (genomic position:1,865,856) and Difficidin (genomic position:2,270,091) (Table S3).

A phylogenetic tree based on genome-wide data from all 14 *Bacillus* strains demonstrated that the HBXN2020 belongs to the *B. velezensis* species (Fig. 3B). To further understand the classification status of HBXN2020, the online tool JSpeciesWS was used to calculate the average nucleotide identity (ANI) based on the BLAST (ANIb) method. As shown in Fig. S3, HBXN2020 was found to be a member of the *B. velezensis* species due to the high percentage of ANIb (more than 97%).

Safety evaluation of *B. velezensis* HBXN2020

We assessed the *in vivo* effects of *B. velezensis* HBXN2020 in a mouse model in order to ascertain its safety. After gavage with *B. velezensis* HBXN2020 spores for two weeks, we observed no significant difference in the body weight of each group of mice (Fig. 4A). The gene expression levels of TNF-α, IL-1β, IL-6, and IL-10 of colon from all groups of mice exhibited no remarkable changes (Fig. 4B). However, in the colon, mRNA levels of the barrier proteins ZO-1 and occludin were trending towards an increase compared with the control group (Fig. 4C). Additionally, blood routine tests and serum biochemistry tests were performed for mice in the control group

and H-HBXN2020 group on the 14th d after oral administration of *B. velezensis* HBXN2020 spore multiple times. As shown in Fig. 4D and S4, the blood parameters of mice after *B. velezensis* HBXN2020 spores treatment, including red blood cells (RBC), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), hemoglobin (HGB), white blood cells (WBC), platelets (PLT), hematocrit (HCT), and mean corpuscular hemoglobin concentration (MCHC), were consistent with those of healthy mice. The serum biochemical parameters, including alanine aminotransferase (ALT), aspartate aminotransferase (AST), albumin (ALB), total bilirubin (TBIL), serum creatinine (CREA), and blood urea nitrogen (BUN), were also within normal limits (Fig. 4E). The corresponding histological analysis of colon tissue from mice receiving low, medium, and high doses of *B. velezensis* HBXN2020 spores (L-HBXN2020, M-HBXN2020, and H-HBXN2020 groups, respectively) is presented in Fig. 4F. The colon tissue sections of mice in the test groups showed no significant differences compared to the control group. Additionally, there were no observable differences in the major organ tissues (heart, liver, spleen, lung, and kidney) of mice treated with the high dose of *B. velezensis* HBXN2020 spores compared to healthy mice (Fig. S5). The results of this trial show that *B. velezensis* HBXN2020 is safe to use and has no negative side effects in mice.

Oral administration of *B. velezensis* HBXN2020 spores alleviated

infection by *S.* Typhimurium

Testing both solid and liquid co-culture methods to see if *B. velezensis* HBXN2020 might suppress *S.* Typhimurium ATCC14028 (STm), the findings indicated that *B. velezensis* HBXN2020 decreased the development of STm in a dose-dependent manner (Fig. 5A, S6). Next, the therapeutic potential of *B. velezensis* HBXN2020 was evaluated in an STm-infected mouse model. At days 1, 3, and 5 after STm infection, mice in the STm+HBXN2020 group and HBXN2020 group were orally administered *B. velezensis* HBXN2020 spores by gavage, STm+CIP group were given oral ciprofloxacin, while the control group and STm+PBS group were orally treated with sterile PBS (Fig. 5B). Also, the number of excreted STm in feces was counted daily at the designated time points. As shown in Fig. 5C, there was a significant and continuous reduction in the number of STm in feces following treatment with *B. velezensis* HBXN2020 spores or ciprofloxacin, while the number of *B. velezensis* HBXN2020 viable bacteria in feces is also gradually decreasing (Fig. S7A). The number of STm in the mice feces of STm+HBXN2020 group and STm+CIP group exhibited a reduction ranging from 0.12 to 1.18 logs and 0.76 to 1.81 logs respectively, compared with the STm+PBS group, over the period from day 2 to day 7 post-treatment. Moreover, treatment with *B. velezensis* HBXN2020 spore or ciprofloxacin, resulted in a 1.06, 1.69, 1.14 logs and 1.70, 2.15, 1.48 logs reduction in the number of STm in the cecum, colon and ileum, respectively (Fig. 5D-E, S7B). The weight loss ($p < 0.05$) and

Disease Activity Index (DAI) scores ($p < 0.01$) were significantly reduced in

mice after *B. velezensis* HBXN2020 spore treatment or ciprofloxacin

treatment compared to that in the PBS treatment group (Fig. 5F-G). In

addition, we also measured the colon length of the mice and found that the

STm+PBS group had a significantly shorter colon length ($p < 0.001$) than the

control group and HBXN2020 group. However, compared to the STm+PBS

group, the STm+HBXN2020 group ($p < 0.05$) and STm+CIP group exhibited a

longer colon (Fig. 5H-I).

Histological analysis further showed that in the control group and

HBXN2020 group, the colon epithelial cells and crypt structure were intact

with neat villi, while STm+PBS group showed significant histological damage,

including erosion or loss of the intestinal epithelium, crypt destruction, and

inflammatory cell infiltration in the colonic tissues (Fig. 6A). Compared with

the STm+PBS group, *B. velezensis* HBXN2020 and ciprofloxacin protected

the mucosal architecture and the loss of intestinal epithelial cells, and reduced

inflammatory cell infiltration (Fig. 6A).

To further assess the impact of *B. velezensis* HBXN2020 on intestinal

infammatory response, the mRNA levels of inflammatory cytokines in the

colonic tissues were measured. As shown in Fig. 6B-D, the mRNA levels of

TNF-α ($p < 0.05$), IL-1β ($p < 0.001$), IL-6 ($p <0.001$), and IL-10 ($p <0.001$) in

the colon tissue were significantly increased in the STm+PBS group

compared to the control group. *B. velezensis* HBXN2020 spore treatment

signifcantly decreased the mRNA levels of TNF-α ($p < 0.05$), IL-1β ($p < 0.001$),

IL-6 ($p < 0.01$), and increased the IL-10 level, compared with the STm+PBS

group. In contrast, ciprofloxacin treatment failed to signifcantly decrease the

mRNA levels of IL-6 in the colon of STm-infected mice (Fig. 6D). Similarly,

ciprofloxacin treatment drastically dampened the mRNA levels of IL-10 in the

colon of STm-infected mice (Fig. 6E).

   To assess the infuence of *B. velezensis* HBXN2020 on the intestinal barrier

integrity, the mRNA levels of tight junction proteins in the colonic tissues were

measured. As presented in Fig. 6, compared with control group, the mRNA

levels of tight junction protein in the STm+PBS group was significantly

reduced, including ZO-1 ($p < 0.05$), occludin ($p < 0.01$), claudin, and Muc2 ($p

< 0.05$) (Fig. 6F-I). Moreover, the levels of tight junction protein transcription

were remarkably elevated in the STm+HBXN2020, STm+CIP, and HBXN2020

group, including ZO-1 ($p < 0.001$, $p < 0.05$ and $p < 0.05$, respectively),

occludin ($p < 0.05$, $p < 0.05$ and $p < 0.01$, respectively), and claudin ($p <

0.001$, $p = 0.0857$ and $p < 0.01$, respectively), except for claudin ($p = 0.0857$)

in group STm+CIP, compared with that of the STm+PBS group (Fig. 6F-I). In

addition, we also detected the transcription level of Muc2 in colon tissue,

which mainly exists in and around goblet cells and is an important component

of the mucus layer. The transcription levels of Muc2 were dramatically

increased in the STm+HBXN2020 group compared with the STm+PBS group

(Fig. 6I). Conversely, the mRNA levels of Muc2 were significantly decreased

in the HBXN2020 group and STm+CIP group (Fig. 6I).

Next, we further explore the impact of *B. velezensis* HBXN2020 on the

intestinal microbiota composition of STm-treated mice by 16S rRNA gene high

throughout sequencing. As shown in Fig. 7, alpha-diversity analysis revealed

that both the richness and diversity (calculated in Sobs, Chao, Shannon, and

Simpson indexes) were lower in STm+PBS group than control group,

HBXN2020 group, STm+HBXN2020 group, and STm+CIP group (Fig. 7A-D).

Although an increasing trend of alpha diversity was observed in STm+CIP

group, but without statistical difference was achieved as compared with other

groups. Principal Components analysis (PCA) based on Bray-Curtis distance

showed that a separation in the gut microbiota structure among control and

STm+PBS group. When *B. velezensis* HBXN2020 spores were supplemented,

the gut microbiota composition of the STm+HBXN2020 group, HBXN2020

group and the control group were closer together ($R^2=0.1616$, $P=0.006$,

Fig. 7E).

We analyzed the community composition of colonic microbiota at the

phylum and genus level, and the results revealed that the composition of the

gut microbiota changed markedly after STm-infected mice. At the phylum

level, *Bacteroidetes*, *Firmicutes*, *Verrucomicrobiota* and *Proteobacteria* were

predominant phyla in the fecal microbiota (Fig. 7F, S8A). At the genus level,

infection with STm in the STm+PBS group dramatically reduced the relative

abundance of *norank_f_Muribaculaceae*, *Lactobacillus* and *Akkermansia* (Fig.

7G, H-J) and enhanced the abundance of *Enterococcus*, *Salmonella*,

Bacteroides, *Alloprevotella* and *Escherichia-Shigella* compared to the control

group (Fig. 7G, K, S8B). In contrast, the STm+HBXN2020 group and

HBXN2020 group significantly improved the relative abundance of

norank_f_Muribaculaceae ($p = 0.49$, $p < 0.05$,) and *Lactobacillus* ($p < 0.05$, p

< 0.001) compared with the STm+PBS group (Fig. 7H-I). However,

ciprofloxacin treatment significantly decreased the relative abundance of

Lactobacillus ($p < 0.05$) and *Akkermansia* ($p < 0.05$) compared with the

STm+HBXN2020 group (Fig. 7I-J). Differentially abundant fecal bacterial taxa

in STm-treated mice in response to *B. velezensis* HBXN2020 and

ciprofloxacin were identified by LEfSe analysis (Fig. 7L). Additionally, we

found that two bacterial genera including *Enterococcus* and *Streptococcus*

were enriched in the STm+PBS group, while the other three taxa were

enriched including *Akkermansia*, *Family_XIII_AD3011_group,* and

Corynebacterium in the STm+HBXN2020 group, *Parabacteroides*,

NK4A214_group, *Turicibacter*, *unclassified_o__Oscillospirales*, and

norank_f__Eubacterium_coprostanoligenes_group in the STm+CIP group,

and *norank_f_Muribaculaceae*, *Rikenellaceae_RC9_gut_group* and

Dubosiella in the HBXN2020 group (Fig. 7L).

Prophylactic *B. velezensis* HBXN2020 spores alleviated infection by *S.*

Typhimurium

Based on the improved therapeutic efficacy of *B. velezensis* HBXN2020 in

the treatment of STm-infected mice, we further explored its potential for

disease prevention by evaluating pretreatment. Mice in the PBS+STm group,

HBXN2020+STm group, and CIP+STm group were given the same amount of

sterile PBS, *B. velezensis* HBXN2020 spores or ciprofloxacin by gavage one

week in advance. As shown in Fig. 8, compared with the PBS+STm group,

oral administration of *B. velezensis* HBXN2020 spores or ciprofloxacin as a

preventive measure remarkably alleviated infection by STm, including weight

loss of mice ($p < 0.01$) (Fig. 8B), a significant reduction in DAI ($p < 0.001$) (the

comprehensive score of weight loss, stool consistency, and blood in the feces,

Fig. 8C), and the prevention of colon length shortening (Fig. 8D-E).

Meanwhile, the number of *B. velezensis* HBXN2020 viable bacteria in feces is

also gradually decreasing with time prolonging (Fig. S9A). Furthermore,

compared with the PBS+STm group, oral administration of *B. velezensis*

HBXN2020 spores or ciprofloxacin not only reduced the number of STm in

mouse feces (Fig. 8F) but also decreased STm colonization in the ileum,

cecum, and colon (Fig. 8G-H, S9B).

Histological analysis further revealed that prophylactic *B. velezensis*

HBXN2020 spore suppressed STm induced loss of the intestinal epithelium,

inflammatory cell infiltration and colonic mucosa damage (Fig. 9A), while

prophylactic ciprofloxacin failed to inhibit the loss of colonic epithelial cells.

We next measured the mRNA levels of inflammatory cytokines and tight

junction proteins in colon tissue in the prophylactic animal experiment. As with the treatment test, prophylactic *B. velezensis* HBXN2020 spores largely attenuated the mRNA levels of TNF-α ($p < 0.001$), IL-1β ($p < 0.001$) and IL-6 ($p < 0.001$), and significantly increased levels of IL-10 ($p < 0.05$), in the colon tissue of mice compared with the PBS+STm group (Fig. 9B-E). Similarly, prophylactic ciprofloxacin significantly reduced the mRNA levels of TNF-α ($p < 0.01$) and IL-1β ($p < 0.001$), and drastically elevated levels of IL-10 ($p < 0.001$) in the colon tissue of mice compared with the PBS+STm group (Fig. 9B-C, E). Also, prophylactic *B. velezensis* HBXN2020 spores signifcantly increased the transcription levels of ZO-1 ($p < 0.001$), occludin ($p < 0.001$), claudin ($p < 0.05$), and Muc2 ($p < 0.001$) compared with that of the PBS+STm group (Fig. 9F-I). Notably, the mRNA levels of ZO-1 ($p < 0.001$), occludin ($p < 0.05$), and Muc2 ($p < 0.05$) in the CIP+STm group were significantly lower than those in the HBXN2020+STm group. Overall, these results suggested that prophylactic *B. velezensis* HBXN2020 spores was capable of alleviating STm-induced colonic injury, and inflammation.

Next, we examined the impact of prophylactic *B. velezensis* HBXN2020 on the intestinal microbiota composition of STm-treated mice. Sobs, Chao and Shannon index were signifcantly increased in the HBXN2020, PBS+STm group, HBXN2020+STm group and CIP+STm group compared with the control group (Fig. 10A-C), while the Simpson index (not statistically different) significantly decreased (Fig. 10D). PCA analysis showed significant

separation between control and PBS+STm groups (R^2=0.3186, p=0.014, Fig. 10E). When *B. velezensis* HBXN2020 spores or ciprofloxacin were supplemented, the community clustering was significantly similar to that of the control group rather than the PBS+STm group (Fig. 10E). Meanwhile, the overall fecal bacterial composition of mice at the phylum and genus level in all groups was similar to that of the treatment experiment. The fecal microbiota was dominated by phyla *Firmicutes*, *Bacteroidetes*, *Verrucomicrobiota*, *Patescibacteria*, and *Actinobacteria* in all five groups (Fig. 10F, S10A). At the genus level, PBS+STm group exhibited lower relative abundances of *Lactobacillus* and *Akkermansia* (Fig. 10G, H-I), and higher relative abundances of *Alistipes*, *Bacteroides*, *Escherichia-Shigella*, *Enterococcus*, and *Alloprevotella*, compared with the control (Fig. 10G, J-M, S10B). Moreover, compared with the PBS+STm group, following supplementation with *B. velezensis* HBXN2020 spores or ciprofloxacin, *Lactobacillus* significantly increased, and the gut microbiota was restored to a composition similar to the control group. It was noteworthy that six bacterial genera including *Alistipes*, *Odoribacter*, and *norank_f__Oscillospiraceae* were enriched in PBS+STm group, while other three taxa (e.g., *Lactobacillus* and *Akkermansia*) were enriched in the control group (Fig. 10N). Moreover, we found that *Candidatus_Arthromitus* was enriched in the HBXN2020+STm group, *Dubosiella* was enriched in the CIP+STm group, and *Faecalibaculum*, *Rikenellaceae_RC9_gut_group*, and *unclassified_f__Lactobacillaceae* were

enriched in the HBXN2020 group (Fig. 10N). Above all, these results indicated

that prophylactic *B. velezensis* HBXN2020 spores regulated the gut

microbiota composition, leading to the potential to attenuate the STm-induced

dysbiosis.

Discussion

S. Typhimurium is an important intestinal pathogen that can cause invasive

intestinal diseases such as bacterial colitis (Herp et al., 2019; Schultz et al.,

2017). Antibiotics are frequently used to treat colitis caused by bacteria;

nevertheless, in recent years, their abuse has greatly increased bacterial

resistance, leading to serious environmental pollution. A recent study showed

that antibiotic-resistance genes in probiotics could be transmitted to the

intestinal microbiota, which may threaten human health (Crits-Christoph et al.,

2022). It is crucial to consider the source of probiotics. Strong environmental

adaptability and disease resistance make black pigs an outstanding

household breed in China (Yang et al., 2022). In this study, *B. velezensis*

HBXN2020 was isolated from the free-range feces of black piglets in a

mountain village in Xianning City (Hubei, China) and exhibited excellent

antibacterial activity. Otherwise, *in vitro* tolerance assays showed that *B.*

velezensis HBXN2020 spores have good tolerance to high temperature,

strong acids, bile salts, as well as simulated gastric and intestinal fluid, which

was similar to previous research results (Du et al., 2022). Furthermore,

studies for antibiotic susceptibility revealed that *B. velezensis* HBXN2020 is

not resistant to antibiotics.

Probiotics are distinct from food or medications in that they can cause

toxins or infections in the body when ingested, and they are living when

swallowed (Sanders et al., 2010; Snydman, 2008). Therefore, assessing the

biosafety of probiotics is crucial. In this study, the safety of *B. velezensis*

HBXN2020 was evaluated by measuring the body weight, intestinal barrier

proteins, and inflammatory cytokines of the experimented mice, as well as

conducting routine blood and biochemical tests. The results showed that the

expression levels of ZO-1 and occludin in the colon of mice increased after

treatment with different doses of *B. velezensis* HBXN2020 spores. It has been

reported that ZO-1, occludin and claudin are the three most important tight

junction proteins in intercellular connections, which play a crucial role in

maintaining the intestinal epithelial barrier (Bazzoni et al., 2000; Zihni et al.,

2016). Thus, it is possible to improve intestinal barrier function by upregulating

ZO-1 and occludin expression levels.

Important aminotransferases in animals, aspartate aminotransferase (AST)

and alanine aminotransferase (ALT), are regarded as critical metrics for

assessing liver function damage (Gou et al., 2022; Ozer et al., 2008). Under

normal circumstances, the levels of ALT and AST are in dynamic balance

without notable changes. However, when the liver is damaged or becomes

dysfunctional, the levels of aminotransferases (ALT and AST) increase

significantly (Tang et al., 2012). Previous studies have shown that CCL4-induced liver injury strongly increases plasma ALT/AST levels (Singhal et al., 2018). Zhang et al. reported that the addition of *Bacillus subtilis* to chicken diets significantly decreased the serum levels of ALT and AST (Zhang et al., 2017). In this present study, the levels of ALT and AST in the probiotic-treated group were slightly lower than those in the control group, indicating that *B. velezensis* HBXN2020 had no negative effect on liver health in mice. The *B. velezensis* HBXN2020 treatment group's blood indices were also found to be comparable to those of the control group, falling within the normal reference range, according to regular blood testing.

Mice were fed with *B. velezensis* HBXN2020 spores orally after being challenged with *S.* Typhimurium ATCC14028 in order to assess the spores' potential to alleviate infection caused by *S.* Typhimurium. Our experimental results demonstrated that oral treatment with *B. velezensis* HBXN2020 spores could alleviate infection by *S.* Typhimurium, as evidenced by decreased weight loss, DAI, and histological damage, which is similar to prior studies (Cao et al., 2019; Wu et al., 2022). Previous research showed that macrophages are the first line of host defense against bacterial infection. They release proinflammatory cytokines, which are critical in initiating adaptive immune responses (Cheng et al., 2014; Dinarello, 2000). Nevertheless, proinflammatory cytokines have immunological properties that can be beneficial for the host to resist the invasion of bacteria and other

microbes in the surrounding environment (Liu et al., 2021). For instance,

pretreatment with recombinant murine TNF-α was shown to protect mice

against lethal bacterial (*E. coli*) infection (Cross et al., 1989). PJ-34 exerted

protective effects on intestinal epithelial cells against invasive *Salmonella*

infection by upregulating IL-6 expression through the ERK and NF-κB signal

pathways (Huang, 2009).

On the other hand, some research has demonstrated that over-activation of

immune cells can result in tissue damage, organ failure, systemic or

persistent inflammation, and autoimmune diseases (Karki et al., 2021; Kotas

and Medzhitov, 2015). IL-10 is a recognized anti-inflammatory mediator that

plays a crucial role in maintaining intestinal microbe-immune homeostasis,

regulating the release of inflammatory mediators, and inhibiting

proinflammatory responses of innate and adaptive immunity (Maloy and

Powrie, 2011; Ouyang et al., 2011; Saraiva et al., 2020). For instance,

previous studies have shown that *Clostridium butyrate* can induce the

production of IL-10 in the intestine, thereby alleviating experimental colitis in

mice (Hayashi et al., 2013). Similar results were observed in our study, where

oral administration of *B. velezensis* HBXN2020 spores reduced the

expression levels of TNF-α, IL-1β, and IL-6 while increasing the levels of IL-10

in the colon of mice. Additionally, as demonstrated by the decreased

histological damage and increased expression levels of intestinal barrier

proteins, oral treatment of *B. velezensis* HBXN2020 spores reduced the

functional damage of the intestinal barrier induced by S. Typhimurium

infection.

Besides host itself, environmental factors such as diet and gut microbiota

have been associated with the development of STm-infected. Gut microbiota

constitutes a critical bridge between environmental factors and host health,

where the beneficial bacteria such as *Lactobacillus* in the microbiota may

exert repair and anti-infammatory function (Li et al., 2022; Liu et al., 2022).

Moreover, supplementation of probiotics has been shown to modulate

intestinal microbiota and reduce the risk of STm-infected (Buddhasiri et al.,

2021; Zhang et al., 2022). This is consistent with our research results

between the *B. velezensis* HBXN2020 treatment group and STm-infected

mice. Here, we found that oral *B. velezensis* HBXN2020 spores raised the

colonic microbiota's alpha diversity and modulated the composition of the

intestinal microbiota, namely, enriching the relative abundance of

Lactobacillus (known for enhancing the epithelial barrier by increasing mucus

secretion and upregulating the expression of tight junction proteins such as

claudin-1, occludin, and ZO-1) (Kaur et al., 2021) and *Akkermansia* (known

for modulating intestinal immune response by producing short chain fatty

acids and reducing the secretion of proinflammatory cytokines) (Cani et al.,

2022), reducing the relative abundance of *Enterococcus*, *Salmonella* and

Bacteroides (harmful to intestinal homeostasis) (Wang et al., 2023). Notably,

we also found that the relative abundance of *Lactobacillus* and *Akkermansia*

in the ciprofloxacin treatment group was significantly lower than that in the *B.*

velezensis HBXN2020 spores treatment group, indicating that excessive use

of antibiotics may interfere with the recovery of beneficial bacteria in the gut.

Collectively, infection with *S.* Typhimurium disrupts gut microbiota and the gut

barrier, leading to intestinal inflammation, while oral *B. velezensis* HBXN2020

spores protects the colon and reduce infection by increasing the abundance

of beneficial bacteria and and barrier integrity and decreasing inflammation.

We further assessed pretreatment in order to learn more about the potential

of *B. velezensis* HBXN2020 spores for disease prevention, given the

increased efficacy of treating infection caused by *S.* Typhimurium. Similarly to

the therapeutic effect of oral *B. velezensis* HBXN2020, prophylactic *B.*

velezensis HBXN2020 spores effectively alleviated the symptoms of STm-

infected mice, namely, reduced the mRNA levels of proinflammatory cytokines

and increased the anti-inflammatory cytokine level in the colon of STm-

infected mice. Prophylactic *B. velezensis* HBXN2020 spores also increased

the mRNA levels of gut barrier proteins in the colon of STm-infected mice.

Furthermore, the microbiota composition was also dramatically modulated by

prophylactic *B. velezensis* HBXN2020 spores in both healthy mice and STm-

infected mice. Also, the community richness revealed by the alpha diversity

index were not significantly impacted in STm-infected mice by prophylactic *B.*

velezensis HBXN2020 spores. Interestingly, we found that prophylactic *B.*

velezensis HBXN2020 spores signifcantly increased the community diversity

in healthy mice, suggesting the potentially benefcial efects of *B. velezensis*

HBXN2020 in gut microenvironment for animals in the future. Moreover,

prophylactic *B. velezensis* HBXN2020 spores significantly boosted SCFA-

producing bacteria including *Lactobacillus* and *Akkermansia*, which was

consistent with the effects of therapeutic oral *B. velezensis* HBXN2020 spores.

Meanwhile, harmful bacteria such as *Alistipes*, *Bacteroides*, *Escherichia-*

Shigella, and *Enterococcus* were significantly enriched in STm-infected mice,

reminding us that increased harmful bacteria might play a major role in

promoting the development of STm-infected. In summary, our results

demonstrated that *B. velezensis* HBXN2020-mediated microbial communities,

especially *Lactobacillus* and *Akkermansia* might play a crucial role in

alleviating the development of STm-infected.

Conclusion

In conclusion, *S*. Typhimurium ATCC14028 infection can result in gut

barrier disruption and intestinal inflammation and intestinal microbiota

dysbiosis. Supplementing *B. velezensis* HBXN2020 spores can improve

intestinal microbiota stability and gut barrier integrity and reduce inflammation

to help prevent or treat infection by *S*. Typhimurium. Thus, our results indicate

that supplementing *B. elezensis* HBXN2020 may represent a novel option for

preventing *Salmonella* infections.

Experimental procedures

Sample collection and strain isolation

Fresh fecal samples were collected from the farms of various locations in southern China including Hubei, Anhui, Hunan, Jiangxi, Guizhou, and Guangxi. All samples were placed in sterile plastic bags, sealed and placed on ice, and immediately transported to the laboratory. *Bacillus* was isolated following a previously described method with slight modification (Unban et al., 2020). The obtained strains were kept in Luria-Bertani (LB) media containing 25% (v/v) glycerol and stored at –80°C, and prepared in LB medium or LB-Agar medium before use.

Antibacterial activity of *Bacillus*

The antibacterial activity of *Bacillus* isolates was evaluated by using the spot-on-plate method on LB agar plates supplemented with indicator bacteria (*Salmonella*, *E. coli*, *S. aureus*). Briefly, an overnight culture of indicator bacteria was dipped by sterile cotton swabs and spread on LB agar plates. *Bacillus* solution was spotted onto the double-layer agar plates and incubated at 37°C for 12 h to measure the inhibition zone. A transparent zone of at least 1 mm around the spot was considered positive.

The antibacterial spectrum of *Bacillus* isolates was compared through the agar diffusion test, based on the results of the spot-on-plate test. In brief, an overnight culture of indicator strains was mixed with 10 mL of TSB soft agar (TSB broth containing 0.7% [wt/vol] agar) and poured into a sterile plate

covered with LB agar and Oxford cups, 8 mm diameter wells were prepared in the TSB agar after removing the cups. The wells were filled with 100 µL of *Bacillus* solution. The plates were incubated at 37°C for 14 h, and the inhibition zone was measured.

The indicator strains used in this study are listed in Table S4, and all the primers used in Table S5. *Clostridium perfringens* were cultured in a fluid thioglycollate medium (FTG) (Hopebio, Qingdao, China) at 45°C. *Streptococcus suis* and *Pasteurella multocida* were cultured in TSB medium supplemented with 5% (v/v) sheep serum (Solarbio, Beijing, China) at 37°C, and *Actinobacillus pleuropneumoniae* were cultured in TSB supplemented with 5% (v/v) sheep serum and 5mM nicotinamide adenine dinucleotide (NAD) at 37°C, while the other bacterial strains were cultured in Luria Bertani (LB) medium (Solarbio, Beijing, China) at 37°C. In addition, the Difco sporulation medium (DSM) was used for inducing the sporulation of *Bacillus* via the nutrient depletion method (Tang et al., 2017).

Growth curves of HBXN2020

The growth curve of HBXN2020 was recorded in flat-bottomed 100-well microtiter plates via detecting optical density at 600 nm (OD_{600}) at 1 h intervals using the automatic growth curve analyzer (Bioscreen, Helsinki, Finland).

***In vitro* resistance assay of HBXN2020**

HBXN2020 spores (100 µL) or vegetative cells (100 µL) were separately

resuspended in 900 µL of LB medium supplemented with different pH values

(2, 3, 4, 5 or 6), bile salts (0.85% NaCl, 0.3%), simulated gastric fluid (SGF,

HCl, pH 1.2) containing 10 g L^{-1} of pepsin in 0.85% NaCl solution, or

simulated intestinal fluid (SIF, NaOH, pH 6.8) containing 10 g L^{-1} of trypsin in

0.05 M KH2PO4 solution, and incubated at 37 °C. A normal LB medium (pH

7.0) was used as the control. At predetermined time points, 100 µL was taken

from each sample, serially diluted 10-fold with sterile PBS (pH 7.2), and then

spread onto LB agar plates. The plates were incubated overnight in a

constant temperature incubator at 37°C, and the bacterial colonies were

counted. The survival rate was calculated using the following formula: Survival

rate = (number of bacteria in the treatment group/number of bacteria in the

control group) × 100%.

One milliliter of HBXN2020 spores or vegetative cells were separately

placed in water baths at different temperatures (37°C, 45°C, 55°C, 65°C,

75°C, 85°C or 95°C) for 20 min, with a 37°C water bath used as the control.

The survival rate was calculated as described above.

Antibiotic susceptibility assays

Antimicrobial susceptibility testing was performed using the Kirby-Bauer

(KB) disk diffusion method in accordance with the Clinical Laboratory

Standards Institute (CLSI) guidelines (CLSI., 2018.). The antimicrobial agents

tested were: Ampicillin, Meropenem, Piperacillin, Gentamycin, Tetracycline,

Doxycycline, Minocycline, Erythromycin, Enrofloxacin, Ofloxacin,

Sulfamethoxazole, Trimethoprim-sulfamethoxazole, Polymyxin B, Teicoplanin,

Trimethoprim, Florfenicol, Spectinomycin, Nitrofurantoin, and Rifampicin. The

diameter of the inhibition zone was measured using a vernier caliper.

Antimicrobial Assays

The *in vitro* antagonistic activity of HBXN2020-CFS was tested using the

agar well-diffusion method against 18 indicator strains (pathogens), which

included 7 standard strains (*E. coli* ATCC 25922, *E. coli* ATCC 35150, *S.*

Typhimurium ATCC 14028 (STm), *S.* Typhimurium SL1344, *S. aureus* ATCC

29213, *S. aureus* ATCC 43300, and *C. perfringens* CVCC 2030) and 11

clinical isolates (*E. coli* EC024, *S.* Enteritidis SE006, *S. aureus* S21, *C.*

perfringens CP023, *C. perfringens* CP002, *S. suis* SC19, *S. suis* SS006, *P.*

multocida PM002, *P. multocida* PM008, *A. pleuropneumoniae* APP015, and *A.*

pleuropneumoniae APP017). All plates were cultured at 37°C for 16 h before

observing the inhibition zone, and the diameter of the inhibition zone was

measured using a vernier caliper. The presence of a clear zone indicated

antagonistic activity.

***In vitro* bacterial competition.**

To investigate whether HBXN2020 directly inhibited the growth of STm, we

performed spot-on lawn and agar-well diffusion assays, as well as co-culture

assays in liquid culture medium. In the spot-on lawn antimicrobial assays, we

prepared double layers of agar by first pouring LB agar into the plate as the

bottom layer. The top layer consisted of 10 mL TSB broth containing 0.7%

agar with STm overnight culture. Then, 10 µL of HBXN2020 overnight culture

and cell-free supernatant (CFS) were respectively spotted onto TSB agar and

incubated at 37°C for 12 h to measure the inhibition zone. A transparent zone

of at least 1 mm around the spot was considered positive.

The antagonistic effect of HBXN2020-CFS against STm was determined

using the agar-well diffusion assays. To collect HBXN2020-CFS, the culture

was centrifuged at 9,000g for 15 min at 4°C, and the supernatant was filtered

through a 0.22 µm membrane filter (Millipore, USA). The STm lawn medium

was prepared by mixing 10 mL of TSB broth containing 0.7% agar with STm

overnight culture and then poured into a sterile plate covered with LB agar

and Oxford cups, 8 mm diameter wells were prepared in the TSB agar after

removing the cups. The wells were filled with 100 µL of HBXN2020-CFS, LB

medium or ampicillin (100 µg mL^{-1}). The plates were incubated at 37°C for 14

h, and the inhibition zone was measured.

The co-culture assay was conducted by incubating HBXN2020 and *S.*

Typhimurium ATCC14028 (carry pET28a (+), kanamycin resistance) (Cao et

al., 2019) separately overnight at 37°C and diluting them to 10^4 CFU/mL.

Then, the two strains were mixed at different ratios (1:1, 1:10, 1:50 or 1:100)

and co-cultured at 37°C with shaking (180 rpm). At predetermined time points,

serial 10-fold dilutions were prepared for all samples and spread onto

selective (kanamycin 50 µg mL^{-1}) LB agar plates and cultured at 37°C for

12 h before bacterial counting. Viable colony counts ranged from 30 to 300

per plate.

Genome sequencing, annotation and analysis

The complete genome of strain was sequenced using the Illumina HiSeq

PE and PacBio RSII (SMRT) platforms (Shanghai Majorbio Bio-pharm

Technology Co., Ltd.). Briefly, the short reads from the Illumina HiSeq PE

were assembled into contigs using SPAdenovo (http://soap.genomics.org.cn/).

The long reads from the PacBio RSII and the Illumina contigs were then

aligned using the miniasm and Racon tools in Unicycler (version 0.4.8,

https://github.com/rrwick/Unicycler) to generate long-read sequences. During

the assembly process, sequence correction was performed using Pilon

(version 1.22, https://github.com/broadinstitute/pilon/wiki/Standard-Output).

Lastly, a complete genome with seamless chromosomes was obtained.

The coding sequences (CDs) of the strain genome were predicted using

Glimmer (version 3.02, http://ccb.jhu.edu/software/glimmer/index.shtml). The

tRNA and rRNA genes were predicted using TRNAscan-SE (version 2.0,

http://trna.ucsc.edu/software/)   and   Barrnap   (version   0.8,

https://github.com/tseemann/barrnap), respectively. The functional annotation

of all CDSs was performed using various databases, including Swiss-Prot

Database (https://web.expasy.org/docs/swiss-prot_guideline.html), Pfam

Database (http://pfam.xfam.org/), EggNOG Database (http://eggnog.embl.de/),

GO Database (http://www.geneontology.org/), and KEGG Database

(http://www.genome.jp/kegg/). The circular map of the strain genome was

generated using CGView (version 2, http://wishart.biology.ualberta.ca/cgview/)

(Stothard and Wishart, 2005).

Comparative genomic analysis

To elucidate the phylogenetic relationships of the strain from a whole-

genome perspective, a phylogenetic tree based on the whole genome was

constructed using the Type (Strain) Genome Server (TYGS) online

(https://ggdc.dsmz.de/) (Meier-Kolthoff et al., 2022). The average nucleotide

identity (ANI) values of the strain genome to the reference strain genome were

then calculated using the JSpeciesWS online service (Richter et al., 2016),

and a heatmap was generated using TBtools (version 1.113) (Chen et al.,

2020).

In vivo safety evaluation of *B. velezensis* HBXN2020

Female specific-pathogen-free (SPF) C57BL/6 mice aged 6 to 8 weeks

were purchased from the Animal Experimental Center of Huazhong

Agricultural University. All mice experiments were conducted in the standard

SPF facility of Huazhong Agricultural University, with 12 h of light and 12 h of

darkness at a temperature of 25°C and ad libitum access to food and water.

The use of animals in this experiment was approved by the Animal Care and

Ethics Committee of Huazhong Agricultural University (Ethics Approval

Number: HZAUMO-2023-0089).

Safety assay of *B. velezensis* HBXN2020 referred to the method described

by Zhou et al. (Zhou et al., 2022) with slight modifications. After a 7-day

acclimation period (free access to water and food), mice were randomly

divided into four treatment groups (n = 5): low-dose *B. velezensis* HBXN2020

spores group (L-HBXN2020 group, 10^7 CFU/mouse), medium-dose *B.*

velezensis HBXN2020 spores group (M-HBXN2020 group, 10^8 CFU/mouse),

high-dose *B. velezensis* HBXN2020 spores group (H-HBXN2020 group, 10^9

CFU/mouse), and a control group. During the experimental period (15 days),

mice were weighed and orally gavaged with their respective treatments once

every two days. On day 15, all mice were euthanized, and blood, heart, liver,

spleen, lung, kidney, ileum, cecum, and colon were collected. Blood samples

were used for routine blood and biochemistry tests. Major organ tissues (heart,

liver, spleen, lung, and kidney) and a 5-mm distal segment of the colon were

used for histopathology. The remaining colonic tissues were rapidly frozen in

liquid nitrogen and stored at –80°C for cytokine and tight junction protein

expression analysis.

Salmonella Typhimurium-infected mouse model

After 7 days of acclimation (free access to water and food), mice were randomly divided into five treatment groups (n = 6): control group, HBXN2020 group, STm+PBS group, STm+HBXN2020 group and STm+CIP group. The *S.* Typhimurium-infected model was performed as previously described (Stecher et al., 2005), with slight modifications. On the first day of the experiment, all mice in the STm+PBS group, STm+HBXN2020 group and STm+CIP group were orally inoculated with 200 μL ($5{\times}10^7$ CFU/mouse) of STm. On days 1, 3 and 5 following STm infection, each mouse in the HBXN2020 group, STm+HBXN2020 group and STm+CIP group received 200 μL ($1{\times}10^8$ CFU/mouse) of *B. velezensis* HBXN2020 spores or ciprofloxacin (1 mg/mL) via gavage administration, respectively. In contrast, the control group and STm+PBS group received 200 μL of sterile PBS by oral gavage. Fecal samples were collected daily following STm infection and resuspended in sterile PBS. The number of STm in mice feces from both the STm+PBS, STm+HBXN2020 and STm+CIP groups was then determined by spreading a serial 10-fold dilution on selective LB agar plates containing 50 μg mL^{-1} kanamycin. Throughout the entire experiment, the body weight, stool consistency and fecal occult blood of all mice were monitored daily. As shown in Table S6, disease activity index (DAI) was calculated by the sum of the scores from three parameters (Praveschotinunt et al., 2019). On day 7 after

STm infection, all mice were euthanized, their ileum, cecum and colon were

collected. The length of colon was measured, and a 5-mm distal segment of

the colon was fixed in 4% paraformaldehyde for further histopathology. The

remaining colon was then rapidly frozen in liquid nitrogen and stored at −80°C

for cytokine and tight junction protein expression analysis. Then, the number

of STm in the ileum, cecum, and colon was determined by spreading serial

10-fold dilutions on selective LB agar plates.

To investigate the prophylactic efficacy of *B. velezensis* HBXN2020 in

ameliorating STm-infected, another independent experiment was conducted

using six-week-old female C57BL/6 mice (SPF). After a seven-day

acclimation period, mice were randomly assigned to five groups (n = 6):

control group, HBXN2020 group, PBS+STm group, HBXN2020+STm group

and CIP+STm group. On days 1, 3, 5 and 7, each mouse in the HBXN2020

group, HBXN2020+STm group and CIP+STm group received 200 μL (1×10^8

CFU/mouse) of *B. velezensis* HBXN2020 spores or ciprofloxacin (1 mg/mL)

via gavage administration. Meanwhile, mice in the control group and

PBS+STm group received 200 μL of sterile PBS by oral gavage. On day 7, all

mice in the PBS+STm group, HBXN2020+STm group and CIP+STm group

were orally inoculated with 200 μL (5×10^7 CFU/mouse) of STm. Fecal

samples were collected daily following STm infection from both groups and

resuspended in sterile PBS. The number of STm in the feces of mice was

then determined by spreading serial 10-fold dilutions on selective LB agar

plates (50 µg mL^{-1} kanamycin). Throughout the entire experiment, the body

weight and DAI scores of all mice were monitored daily. On day 12, all mice

were euthanized, and the ileum, cecum, and colon were collected. The colon

length was measured, and a 5-mm distal segment of the colon was fixed in 4%

formalin for sectioning and staining. The remaining colon was stored at –80°C

for future analysis. Lastly, the number of STm in the ileum, cecum, and colon

was determined using a selective LB agar plate.

Determination of cytokines and tight junction protein expression in

colon tissue

Total RNA was extracted from colon tissues using TRIpure reagent (Aidlab,

China), and cDNA was obtained using HiScript$^®$ III RT SuperMix for qPCR

(+gDNA wiper) (Vazyme, China). RT-qPCR for each gene was performed in

triplicate using qPCR SYBR Green Master Mix (Yeasen Biotechnology,

Shanghai, China). The relative expression level of cytokine and tight junction

protein genes was calculated using the $2^{-\Delta\Delta Ct}$ method with β-actin and

GAPDH as reference genes. The primer sequences used in the RT-qPCR test

are listed in Table S5.

Histopathology analysis

Colon tissue samples (0.5 cm) were fixed in 4% paraformaldehyde for 24 h,

and the fixed tissues were then embedded in paraffin and sectioned. The

sections were stained with hematoxylin and eosin (H&E) and observed and

imaged using an optical microscope (Olympus Optical, Tokyo, Japan). The

histopathological score included the degree of inflammatory infiltration,

changes in crypt structures, and the presence or absence of ulceration and

edema. The scoring criteria were determined as previously described (Wu et

al., 2022).

16S rRNA gene sequencing and analysis

According to the manufacturer's instructions, colon microbial community

genomes were extracted using E.Z-N.A ® Stool DNA Kit (Omega; D4015-01),

and quality was detected by 1% agarose gel electrophoresis. The 16S rRNA

V3-V4 variable region was amplified by PCR using universal primers 338F (5

'-　　ACTCCTACGGGGGGCAG-3')　　and　　806R　　(5　　'-

GACTACHVGGGTWTCTAAT-3'). The PCR products were examined by

electrophoresis on 2% agarose gels and then purified with the AxyPrep DNA

gel extraction kit (Axygen Biosciences, USA). A sequencing library was

constructed using the NEXTFLEX® Rapid DNA-Seq Kit, and sequencing was

performed using the Illumina MiSeq platform. Raw reads were quality

evaluated and filtered by fastp (version 0.20.0) and merged using FLASH

(version 1.2.7). The optimized sequences were clustered into operational

taxonomic units (OTUs) based on 97% sequence similarity using UPARSE

(version 7.1). The representative sequences of each OTU was classified by

RDP classifier (version 2.2; confidence threshold value, 0.7). Alpha diversity

was assessed using the ACE, chao, Shannon, and Simpson indices. The β-

diversity analysis was performed using Bray-Curtis distances and visualized

through principal component analysis (PCA). Linear discriminant analysis

(LDA) effect size (LefSe) was used to identify differential microbiota between

groups.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism 8.3.0 (GraphPad

Software, San Diego, CA, USA) and Excel (Microsoft, Redmond, USA). Data

are presented as mean ± standard error of the mean (SEM). Differences

between two groups were evaluated using two-tailed unpaired Student's t-test,

and all other comparisons were conducted using one-way analysis of variance

(ANOVA) with Tukey's multiple comparisons test. For all analyses,

significance differences are denoted as: *, $p < 0.05$, **, $p < 0.01$, and ***, $p <$

0.001.

Data availability

The datasets supporting the conclusions of this article were deposited in the

NCBI Sequence Read Archive database under the accession numbers:

CP119399.1 (whole genome sequencing raw data) and PRJNA1150571

(microbiota raw sequencing data).

Conflict of interest: We declare no conflict of interest.

ACKNOWLEDGMENTS

This work was supported by grants from the National Program on Key Research Project of China (2021YFD1800300, 2022YFD1800800), "Yingzi Tech & Huazhong Agricultural University Intelligent Research Institute of Food Health" (No. IRIFH202209; IRIFH202301), and the Fundamental Research Funds for the Central Universities (2662016PY004).

Author contributions

Linkang Wang: Conceptualization, Methodology, Data curation, Formal analysis, Writing - original draft. **Haiyan Wang, Xinxin Li, Mengyuan Zhu**: Methodology, Data curation, Planned and performed the experiments. **Dongyang Gao, Dayue Hu, Zhixuan Xiong**: Methodology, Writing - Reviewing and Editing. **Xiangmin Li**: Supervision, Writing-reviewing and editing. **Ping Qian**: Conceptualization, Funding acquisition, Project administration, Supervision, Writing-Reviewing and Editing. All authors read and approved the submitted version of the paper.

Funding information

This work was supported by grants from the National Program on Key

Reference

Abdhul, K., Ganesh, M., Shanmughapriya, S., Vanithamani, S., Kanagavel, M., Anbarasu, K., and
Natarajaseenivasan, K. (2015). Bacteriocinogenic potential of a probiotic strain Bacillus coagulans
[BDU3] from Ngari. Int J Biol Macromol *79*, 800-806.
Bazzoni, G., Martinez-Estrada, O.M., Orsenigo, F., Cordenonsi, M., Citi, S., and Dejana, E. (2000).
Interaction of junctional adhesion molecule with the tight junction components ZO-1, cingulin, and
occludin. J Biol Chem *275*, 20520-20526.
Buddhasiri, S., Sukjoi, C., Kaewsakhorn, T., Nambunmee, K., Nakphaichit, M., Nitisinprasert, S., and
Thiennimitr, P. (2021). Anti-inflammatory Effect of Probiotic Limosilactobacillus reuteri KUB-AC5
Against Salmonella Infection in a Mouse Colitis Model. Front Microbiol *12*, 716761.
Cani, P.D., Depommier, C., Derrien, M., Everard, A., and de Vos, W.M. (2022). Akkermansia muciniphila:
paradigm for next-generation beneficial microorganisms. Nat Rev Gastroenterol Hepatol *19*, 625-637.
Cao, Z., Wang, X., Pang, Y., Cheng, S., and Liu, J. (2019). Biointerfacial self-assembly generates lipid
membrane coated bacteria for enhanced oral delivery and treatment. Nat Commun *10*, 5783.
Chen, C., Chen, H., Zhang, Y., Thomas, H.R., Frank, M.H., He, Y., and Xia, R. (2020). TBtools: An
Integrative Toolkit Developed for Interactive Analyses of Big Biological Data. Mol Plant *13*, 1194-1202.
Cheng, A., Yan, H., Han, C., Wang, W., Tian, Y., and Chen, X. (2014). Polyphenols from blueberries
modulate inflammation cytokines in LPS-induced RAW264.7 macrophages. Int J Biol Macromol *69*,
382-387.
CLSI. (2018.). Performance standards for antimicrobial disk susceptibility tests, M02, 13th ed. Clinical
and Laboratory Standards Institute, Wayne, PA.
Cristofori, F., Dargenio, V.N., Dargenio, C., Miniello, V.L., Barone, M., and Francavilla, R. (2021). Anti-
Inflammatory and Immunomodulatory Effects of Probiotics in Gut Inflammation: A Door to the Body.
Front Immunol *12*, 578386.
Crits-Christoph, A., Hallowell, H.A., Koutouvalis, K., and Suez, J. (2022). Good microbes, bad genes?
The dissemination of antimicrobial resistance in the human microbiome. Gut Microbes *14*, 2055944.
Cross, A.S., Sadoff, J.C., Kelly, N., Bernton, E., and Gemski, P. (1989). Pretreatment with recombinant
murine tumor necrosis factor alpha/cachectin and murine interleukin 1 alpha protects mice from
lethal bacterial infection. J Exp Med *169*, 2021-2027.
Dinarello, C.A. (2000). Proinflammatory cytokines. Chest *118*, 503-508.
Du, H., Yao, W., Kulyar, M.F., Ding, Y., Zhu, H., Pan, H., Li, K., Bhutta, Z.A., Liu, S., and Li, J. (2022). Effects
of Bacillus amyloliquefaciens TL106 Isolated from Tibetan Pigs on Probiotic Potential and Intestinal
Microbes in Weaned Piglets. Microbiol Spectr *10*, e0120521.

Fabrega, A., and Vila, J. (2013). Salmonella enterica serovar Typhimurium skills to succeed in the host:
virulence and regulation. Clin Microbiol Rev *26*, 308-341.

Ferri, M., Ranucci, E., Romagnoli, P., and Giaccone, V. (2017). Antimicrobial resistance: A global
emerging threat to public health systems. Crit Rev Food Sci Nutr *57*, 2857-2876.

Gou, S., Chen, N., Wu, X., Zu, M., Yi, S., Ying, B., Dai, F., Ke, B., and Xiao, B. (2022). Multi-responsive
nanotheranostics with enhanced tumor penetration and oxygen self-producing capacities for
multimodal synergistic cancer therapy. Acta Pharm Sin B *12*, 406-423.

Gu, S.B., Zhao, L.N., Wu, Y., Li, S.C., Sun, J.R., Huang, J.F., and Li, D.D. (2015). Potential probiotic
attributes of a new strain of Bacillus coagulans CGMCC 9951 isolated from healthy piglet feces. World
J Microbiol Biotechnol *31*, 851-863.

Hayashi, A., Sato, T., Kamada, N., Mikami, Y., Matsuoka, K., Hisamatsu, T., Hibi, T., Roers, A., Yagita, H.,
Ohteki, T., *et al.* (2013). A single strain of Clostridium butyricum induces intestinal IL-10-producing
macrophages to suppress acute experimental colitis in mice. Cell Host Microbe *13*, 711-722.

Herp, S., Brugiroux, S., Garzetti, D., Ring, D., Jochum, L.M., Beutler, M., Eberl, C., Hussain, S., Walter, S.,
Gerlach, R.G., *et al.* (2019). Mucispirillum schaedleri Antagonizes Salmonella Virulence to Protect Mice
against Colitis. Cell Host Microbe *25*, 681-694 e688.

Huang, F.C. (2009). Upregulation of Salmonella-induced IL-6 production in Caco-2 cells by PJ-34, PARP-
1 inhibitor: involvement of PI3K, p38 MAPK, ERK, JNK, and NF-kappaB. Mediators Inflamm *2009*,
103890.

Karki, R., Sharma, B.R., Tuladhar, S., Williams, E.P., Zalduondo, L., Samir, P., Zheng, M., Sundaram, B.,
Banoth, B., Malireddi, R.K.S., *et al.* (2021). Synergism of TNF-alpha and IFN-gamma Triggers
Inflammatory Cell Death, Tissue Damage, and Mortality in SARS-CoV-2 Infection and Cytokine Shock
Syndromes. Cell *184*, 149-168 e117.

Katiyo, S., Muller-Pebody, B., Minaji, M., Powell, D., Johnson, A.P., De Pinna, E., Day, M., Harris, R., and
Godbole, G. (2019). Epidemiology and Outcomes of Nontyphoidal Salmonella Bacteremias from
England, 2004 to 2015. J Clin Microbiol *57*.

Kaur, H., Gupta, T., Kapila, S., and Kapila, R. (2021). Protective effects of potential probiotic
Lactobacillus rhamnosus (MTCC-5897) fermented whey on reinforcement of intestinal epithelial
barrier function in a colitis-induced murine model. Food Funct *12*, 6102-6116.

Kotas, M.E., and Medzhitov, R. (2015). Homeostasis, inflammation, and disease susceptibility. Cell *160*,
816-827.

Li, A., Wang, Y., Li, Z., Qamar, H., Mehmood, K., Zhang, L., Liu, J., Zhang, H., and Li, J. (2019). Probiotics
isolated from yaks improves the growth performance, antioxidant activity, and cytokines related to
immunity and inflammation in mice. Microb Cell Fact *18*, 112.

Li, H., Cheng, S., Huo, J., Dong, K., Ding, Y., Man, C., Zhang, Y., and Jiang, Y. (2022). Lactobacillus
plantarum J26 Alleviating Alcohol-Induced Liver Inflammation by Maintaining the Intestinal Barrier
and Regulating MAPK Signaling Pathways. Nutrients *15*.

Liu, C., Chu, D., Kalantar-Zadeh, K., George, J., Young, H.A., and Liu, G. (2021). Cytokines: From Clinical
Significance to Quantification. Adv Sci (Weinh) *8*, e2004433.

Liu, Z., Zhao, J., Sun, R., Wang, M., Wang, K., Li, Y., Shang, H., Hou, J., and Jiang, Z. (2022). Lactobacillus
plantarum 23-1 improves intestinal inflammation and barrier function through the TLR4/NF-κB
signaling pathway in obese mice. Food Funct *13*, 5971-5986.

Maloy, K.J., and Powrie, F. (2011). Intestinal homeostasis and its breakdown in inflammatory bowel
disease. Nature *474*, 298-306.

Meier-Kolthoff, J.P., Carbasse, J.S., Peinado-Olarte, R.L., and Goker, M. (2022). TYGS and LPSN: a
database tandem for fast and reliable genome-based classification and nomenclature of prokaryotes.
Nucleic Acids Res *50*, D801-D807.

Mohan, A., Munusamy, C., Tan, Y.C., Muthuvelu, S., Hashim, R., Chien, S.L., Wong, M.K., Khairuddin,
N.A., Podin, Y., Lau, P.S., *et al.* (2019). Invasive Salmonella infections among children in Bintulu,
Sarawak, Malaysian Borneo: a 6-year retrospective review. BMC Infect Dis *19*, 330.

Ohland, C.L., and Macnaughton, W.K. (2010). Probiotic bacteria and intestinal epithelial barrier
function. Am J Physiol Gastrointest Liver Physiol *298*, G807-819.

Ouattara, H.G., Reverchon, S., Niamke, S.L., and Nasser, W. (2017). Regulation of the synthesis of pulp
degrading enzymes in Bacillus isolated from cocoa fermentation. Food Microbiol *63*, 255-262.

Ouyang, W., Rutz, S., Crellin, N.K., Valdez, P.A., and Hymowitz, S.G. (2011). Regulation and functions of
the IL-10 family of cytokines in inflammation and disease. Annu Rev Immunol *29*, 71-109.

Ozer, J., Ratner, M., Shaw, M., Bailey, W., and Schomaker, S. (2008). The current state of serum
biomarkers of hepatotoxicity. Toxicology *245*, 194-205.

Paulson, J.A., Zaoutis, T.E., Council on Environmental, H., and Committee on Infectious, D. (2015).
Nontherapeutic Use of Antimicrobial Agents in Animal Agriculture: Implications for Pediatrics.
Pediatrics *136*, e1670-1677.

Praveschotinunt, P., Duraj-Thatte, A.M., Gelfat, I., Bahl, F., Chou, D.B., and Joshi, N.S. (2019).
Engineered E. coli Nissle 1917 for the delivery of matrix-tethered therapeutic domains to the gut. Nat
Commun *10*, 5580.

Reyman, M., van Houten, M.A., Watson, R.L., Chu, M., Arp, K., de Waal, W.J., Schiering, I., Plotz, F.B.,
Willems, R.J.L., van Schaik, W., *et al.* (2022). Effects of early-life antibiotics on the developing infant gut
microbiome and resistome: a randomized trial. Nat Commun *13*, 893.

Richter, M., Rossello-Mora, R., Oliver Glockner, F., and Peplies, J. (2016). JSpeciesWS: a web server for
prokaryotic species circumscription based on pairwise genome comparison. Bioinformatics *32*, 929-
931.

Sanders, M.E., Akkermans, L.M., Haller, D., Hammerman, C., Heimbach, J., Hormannsperger, G., Huys,
G., Levy, D.D., Lutgendorff, F., Mack, D., *et al.* (2010). Safety assessment of probiotics for human use.
Gut Microbes *1*, 164-185.

Saraiva, M., Vieira, P., and O'Garra, A. (2020). Biology and therapeutic potential of interleukin-10. J Exp
Med *217*.

Schultz, B.M., Paduro, C.A., Salazar, G.A., Salazar-Echegarai, F.J., Sebastian, V.P., Riedel, C.A., Kalergis,
A.M., Alvarez-Lobos, M., and Bueno, S.M. (2017). A Potential Role of Salmonella Infection in the Onset
of Inflammatory Bowel Diseases. Front Immunol *8*, 191.

Shao, J., Liu, Y., Xie, J., Stefanic, P., Lv, Y., Fan, B., Mandic-Mulec, I., Zhang, R., Shen, Q., and Xu, Z.
(2022). Annulment of Bacterial Antagonism Improves Plant Beneficial Activity of a Bacillus velezensis
Consortium. Appl Environ Microbiol *88*, e0024022.

Singhal, M., Liu, X., Inverso, D., Jiang, K., Dai, J., He, H., Bartels, S., Li, W., Abdul Pari, A.A.,
Gengenbacher, N., *et al.* (2018). Endothelial cell fitness dictates the source of regenerating liver
vasculature. J Exp Med *215*, 2497-2508.

Snydman, D.R. (2008). The safety of probiotics. Clin Infect Dis *46 Suppl 2*, S104-111; discussion S144-
151.

Stecher, B., Macpherson, A.J., Hapfelmeier, S., Kremer, M., Stallmach, T., and Hardt, W.D. (2005).
Comparison of Salmonella enterica serovar Typhimurium colitis in germfree mice and mice pretreated

with streptomycin. Infect Immun *73*, 3228-3241.

Stothard, P., and Wishart, D.S. (2005). Circular genome visualization and exploration using CGView.
Bioinformatics *21*, 537-539.

Tang, Y., Gao, C., Xing, M., Li, Y., Zhu, L., Wang, D., Yang, X., Liu, L., and Yao, P. (2012). Quercetin
prevents ethanol-induced dyslipidemia and mitochondrial oxidative damage. Food Chem Toxicol *50*,
1194-1200.

Tang, Z., Sun, H., Chen, T., Lin, Z., Jiang, H., Zhou, X., Shi, C., Pan, H., Chang, O., Ren, P., *et al.* (2017).
Oral delivery of Bacillus subtilis spores expressing cysteine protease of Clonorchis sinensis to grass
carp (Ctenopharyngodon idellus): Induces immune responses and has no damage on liver and
intestine function. Fish Shellfish Immunol *64*, 287-296.

Unban, K., Kochasee, P., Shetty, K., and Khanongnuch, C. (2020). Tannin-Tolerant and Extracellular
Tannase Producing Bacillus Isolated from Traditional Fermented Tea Leaves and Their Probiotic
Functional Properties. Foods *9*.

Wang, L., Gou, X., Ding, Y., Liu, J., Wang, Y., Wang, Y., Zhang, J., Du, L., Peng, W., and Fan, G. (2023). The
interplay between herbal medicines and gut microbiota in metabolic diseases. Front Pharmacol *14*,
1105405.

Wang, Y., Xie, Q., Zhang, Y., Ma, W., Ning, K., Xiang, J.Y., Cui, J., and Xiang, H. (2020). Combination of
probiotics with different functions alleviate DSS-induced colitis by regulating intestinal microbiota, IL-
10, and barrier function. Appl Microbiol Biotechnol *104*, 335-349.

Wu, J., Lin, Z., Wang, X., Zhao, Y., Zhao, J., Liu, H., Johnston, L.J., Lu, L., and Ma, X. (2022).
Limosilactobacillus reuteri SLZX19-12 Protects the Colon from Infection by Enhancing Stability of the
Gut Microbiota and Barrier Integrity and Reducing Inflammation. Microbiol Spectr *10*, e0212421.

Yang, W., Liu, Z., Zhao, Q., Du, H., Yu, J., Wang, H., Liu, X., Liu, H., Jing, X., Yang, H., *et al.* (2022).
Population Genetic Structure and Selection Signature Analysis of Beijing Black Pig. Front Genet *13*,
860669.

Yang, W.H., Westman, J.S., Heithoff, D.M., Sperandio, M., Cho, J.W., Mahan, M.J., and Marth, J.D.
(2021). Neu3 neuraminidase induction triggers intestinal inflammation and colitis in a model of
recurrent human food-poisoning. Proc Natl Acad Sci U S A *118*.

Ye, M., Tang, X., Yang, R., Zhang, H., Li, F., Tao, F., Li, F., and Wang, Z. (2018). Characteristics and
Application of a Novel Species of Bacillus: Bacillus velezensis. ACS Chem Biol *13*, 500-505.

Zhang, D.X., Kang, Y.H., Zhan, S., Zhao, Z.L., Jin, S.N., Chen, C., Zhang, L., Shen, J.Y., Wang, C.F., Wang,
G.Q., *et al.* (2019). Effect of Bacillus velezensis on Aeromonas veronii-Induced Intestinal Mucosal
Barrier Function Damage and Inflammation in Crucian Carp (Carassius auratus). Front Microbiol *10*,
2663.

Zhang, L., Bai, K., Zhang, J., Xu, W., Huang, Q., and Wang, T. (2017). Dietary effects of Bacillus subtilis
fmbj on the antioxidant capacity of broilers at an early age. Poult Sci *96*, 3564-3573.

Zhang, R., Li, Z., Gu, X., Zhao, J., Guo, T., and Kong, J. (2022). Probiotic Bacillus subtilis LF11 Protects
Intestinal Epithelium Against Salmonella Infection. Front Cell Infect Microbiol *12*, 837886.

Zhou, J., Li, M., Chen, Q., Li, X., Chen, L., Dong, Z., Zhu, W., Yang, Y., Liu, Z., and Chen, Q. (2022).
Programmable probiotics modulate inflammation and gut microbiota for inflammatory bowel disease
treatment after effective oral delivery. Nat Commun *13*, 3432.

Zihni, C., Mills, C., Matter, K., and Balda, M.S. (2016). Tight junctions: from simple barriers to
multifunctional molecular gates. Nat Rev Mol Cell Biol *17*, 564-580.

Figure legends

Fig. 1 Growth curve of *B. velezensis* HBXN2020 and its *in vitro* resistance against environmental assaults. (**A**) Growth curves of *B. velezensis* HBXN2020 cultured in LB medium at 37°C, detection of OD_{600} values at 1 h intervals in microplate reader. (**B**) Survival rate of endospore and vegetative cells of *B. velezensis* HBXN2020 after 30 min at different temperatures (37°C, 45°C, 55°C, 65°C, 75°C, 85°C or 95°C). Equal amounts of endospore and vegetative cells of HBXN2020 were exposed to the following: (**C**) acid solution (pH 2 to 7), (**D**) 0.3% bile salts, (**E**) SGF (pH 1.2) supplemented with pepsin, and (**F**) SIF (pH 6.8) containing trypsin at 37°C. At predetermined time points, 100 μL was taken from each sample, and tenfold serially diluted with sterile PBS (pH 7.2), and then spread on LB agar plates and cultured at 37°C for 12 h before bacterial counting. Each group was repeated three times (n = 3).

Fig. 2 Antibiotic susceptibility of *B. velezensis* HBXN2020 and bacteriostasis assay *in vitro*. (**A**) The diameter of the antibacterial zone indicates the extent of sensitivity to antibiotics. (**B**) The diameter of the antibacterial zone indicates the extent of inhibition against pathogenic bacteria. The diameter of the antibacterial zone was measured with vernier caliper.

1028   Each group was repeated three times (n = 3). R, resistant; I, moderately

1029   sensitive; S, sensitive.

1030

Fig. 3 Genomic characteristics and phylogenetic relationships of *B.*

***velezensis* HBXN2020.** (**A**) The whole-genome map of *B. velezensis*

HBXN2020 with its genomic features. The map consists of 6 circles. From the

inner circle to the outer circle: (1) GC-skew, (2) GC content, (3) reverse

protein-coding genes, different colors represents different COG functional

classifications, (4) genes transcribed in reverse direction, (5) genes

transcribed in forward direction, (6) forward protein-coding genes, different

colors represents different COG functional classifications. (**B**) The whole

genome phylogenetic tree was constructed based on genome-wide data from

14 *bacillus* strains. *B. velezensis* HBXN2020 are indicated in bold.

1041

Fig. 4 *In vivo* safety evaluation of *B. velezensis* HBXN2020 in a mouse

model. (**A**) Body weights changes of mice during gavage with *B. velezensis*

HBXN2020 spores. Mice were treated with sterile PBS (control group) or low-

dose (L-HBXN2020 group), medium dose (M-HBXN2020 group), and high-

dose (H-HBXN2020 group) of *B. velezensis* HBXN2020 spores. Weighing and

gavage were performed once every two days during the experimental period

(15 days). Data were shown as mean values ± SEM (n = 5). (**B**) The mRNA

levels of inflammatory cytokines in the colon of mice measured by RT-qPCR.

Data were shown as mean values ± SEM (n = 5). (**C**) The mRNA levels of

barrier protein ZO-1, occludin, claudin and Muc2 in the colon of mice

measured by RT-qPCR. Data were shown as mean values ± SEM (n = 5). (**D**)

Major blood routine parameters and (**E**) serum biochemical parameters of

mice in the control group and H-HBXN2020 group. Data were shown as mean

values ± SEM (n = 3). (**F**) H&E stained colon sections in the different groups.

Scale bar: 200 μm.

Fig. 5 Oral *B. velezensis* HBXN2020 spores alleviated infection by *S.*

Typhimurium. (**A**) *In vitro* Bacterial competition between STm and *B.*

velezensis HBXN2020. STm were co-incubated with *B. velezensis* HBXN2020

at various ratios at 37°C with shaking. The growth of STm was reflected by

bacterial counting per hour. (**B**) Experimental design for treatment in this

study. Orally administrated with either sterile PBS, *B. velezensis* HBXN2020

spores or ciprofloxacin by gavage at days 1, 3, and 5 after STm (5×10^7

CFU/mouse) infection, respectively. All mice were euthanized at day 7 after

STm infection. (**C**) Bacterial count of STm in mouse feces. Fecal samples

were collected per day after STm infection and resuspended in sterile PBS

(0.1 g of fecal resuspended in 1 mL of sterile PBS). One hundred microliters

of each sample performed a serial of 10-fold dilutions and spread on selective

agar plates (50 μg mL^{-1} kanamycin) and incubated at 37°C for 12 h before

bacterial counting. The bacterial loads of STm in (**D**) cecum and (**E**) colon.

The cecum and colon were harvested and then homogenized. Statistical

signifcance was evaluated using Student's t-test (*, $p < 0.05$, **, $p < 0.01$, and

***, $p < 0.001$). (**F**) Daily body weight changes and (**G**) daily disease activity

index (DAI) scores of mice with different treatment groups. Data were shown

as mean values ± SEM (n = 6). Statistical signifcance was evaluated using

one-way analysis of variance (ANOVA) with Tukey's multiple comparisons

test (*, $p < 0.05$, **, $p < 0.01$, and ***, $p < 0.001$). (**H**) Colonic tissue images. (**I**)

The length of the colon from per group (n = 6). Statistical signifcance was

evaluated using one-way analysis of variance (ANOVA) with Tukey's multiple

comparisons test (*, $p < 0.05$, **, $p < 0.01$, and ***, $p < 0.001$).

1082

Fig. 6 Oral *B. velezensis* HBXN2020 spores attenuated colonic damage

and inflammatory reaction. (**A**) H&E stained colon tissue sections. Scale bar:

50 μm. The mRNA levels of (**B**) TNF-α, (**C**) IL-1β, (**D**) IL-6, and (**E**) IL-10 were

detected by RT-qPCR. Data were shown as mean values ± SEM (n = 5). The

mRNA levels of (**F**) ZO-1, (**G**) occludin, (**H**) claudin, and (**I**) Muc2 in colon

tissue were detected by RT-qPCR. Data were shown as mean values ± SEM

(n = 5). Statistical signifcance was evaluated using one-way analysis of

variance (ANOVA) with Tukey's multiple comparisons test (*, $p < 0.05$, **, $p <$

0.01, and ***, $p < 0.001$).

1092

1093 **Fig. 7 Oral *B. velezensis* HBXN2020 spores regulated the composition of**

intestinal microbiota. The α-diversity of the gut microbiota, determined by the (**A**) Sobs, (**B**) Chao, (**C**) Shannon, and (**D**) Simpson index. Data were shown as mean values ± SEM (n = 5). (**E**) The PCA plot showed the β-diversity of the gut microbiota based on Bray-Curtis distance at the OTU level. Heatmap of the community composition of colonic microbiota at the phylum (top 15 phyla) (**F**) and genus (top 30 genera) (**G**) level. Relative abundance of selected taxa (**H**) *norank_f_Muribaculaceae*, (**I**) *Lactobacillus*, (**J**) *Akkermansia*, and (**K**) *Enterococcus*. (**L**) Analysis of differences in the microbial communities by LEfSe (linear discriminant analysis, LDA score > 2) among different groups. Signifcance was evaluated by one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test (*, $p < 0.05$, **, $p < 0.01$, and ***, $p < 0.001$).

Fig. 8 Prophylactic *B. velezensis* HBXN2020 spores attenuated the symptoms of *S.* Typhimurium-infected mouse. (**A**) Experimental design for treatment in this study. At days 1, 3, 5, and 7, each mouse in the HBXN2020+STm group, CIP+STm group, and PBS+STm group were received 200 µL of *B. velezensis* HBXN2020 spores (1×10^8 CFU/mouse), ciprofloxacin or sterile PBS by gavage, respectively. Then, mice in PBS+STm group, HBXN2020+STm group and CIP+STm group were orally inoculated with 200 µL (5×10^7 CFU/mouse) of STm on day 7, respectively. On day 12, all mice were euthanized. (**B**) Daily body weight changes and (**C**) daily disease

activity index (DAI) scores of mice with different groups following STm

treatment. Data were shown as mean values ± SEM (n = 6). Statistical

signifcance was evaluated using one-way analysis of variance (ANOVA) with

Tukey's multiple comparisons test (*, $p < 0.05$, **, $p < 0.01$, and ***, $p < 0.001$).

(**D**) Colonic tissue images. (**E**) The length of the colon from per group (n = 6).

(**F**) Bacterial count of STm in mouse feces. Fecal samples were collected

every day after STm infection and resuspended in sterile PBS (0.1 g of fecal

resuspended in 1 mL of sterile PBS). One hundred microliters of each sample

performed a serial of 10-fold dilutions and spread on selective agar plates (50

µg mL^{-1} kanamycin) and incubated at 37°C for 12 h before bacterial counting.

The bacterial loads of STm in (**G**) cecum and (**H**) colon. The cecum and colon

were harvested and then homogenized. Statistical signifcance was evaluated

using Student's t-test (*, $p < 0.05$, **, $p < 0.01$, and ***, $p < 0.001$).

Fig. 9 Prophylactic *B. velezensis* HBXN2020 spores attenuated colonic

damage and inflammatory reaction. (**A**) H&E stained colon tissue sections.

Scale bar: 50 µm. The mRNA levels of (**B**) TNF-α, (**C**) IL-1β, (**D**) IL-6, and (**E**)

IL-10 were detected by RT-qPCR. Data were shown as mean values ± SEM

(n = 5). The mRNA levels of (**F**) ZO-1, (**G**) occludin, (**H**) claudin, and (**I**) Muc2

in colon tissue were detected by RT-qPCR. Data were shown as mean values

± SEM (n = 6). Statistical signifcance was evaluated using one-way analysis

of variance (ANOVA) with Tukey's multiple comparisons test (*, $p < 0.05$, **, p

1138 < 0.01, and [***], $p < 0.001$).

1139

Fig. 10 Prophylactic *B. velezensis* HBXN2020 spores regulated the

composition of gut microbiota. (**A** to **D**) Alpha diversity of the intestinal

microbiota. Data were shown as mean values ± SEM (n = 5). (**E**) The PCA

plot showed the β-diversity among different microbial community groups

based on Bray-Curtis distance at the OTU level. Heatmap of the community

composition of colonic microbiota at the phylum (top 15 phyla) (**F**) and genus

(top 30 genera) (**G**) level. Relative abundance of selected taxa (**H**)

Lactobacillus, (**I**) *Akkermansia*, (**J**) *Alistipes*, (**K**) *Bacteroides*, (**L**) *Escherichia-*

Shigella, and (**M**) *Enterococcus*. Data were shown as mean values ± SEM (n

= 5). (**N**) Analysis of differences in the microbial taxa by LEfSe (LDA score > 2)

in different groups. Signifcance was evaluated by one-way analysis of

variance (ANOVA) with Tukey's multiple comparisons test (*, $p < 0.05$, **, $p <$

0.01, and ***, $p < 0.001$).