Bacillus velezensis HBXN2020 alleviates Salmonella Typhimurium-induced colitis by improving intestinal barrier integrity and reducing inflammation

Linkang Wang,[1,2,3] Haiyan Wang,[1,2,3] Xinxin Li,[1,2,3] Mengyuan Zhu,[1,2,3] Dongyang Gao,[1,2,3] Dayue Hu,[1,2,3] Zhixuan Xiong,[1,2,3] Xiangmin Li[1,2,3*] and Ping Qian[1,2,3*]

[1]*National Key Laboratory of Agricultural Microbiology, Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan, Hubei, 430070, People's Republic of China.*

[2]*College of Veterinary Medicine, Huazhong Agricultural University, Wuhan, Hubei, People's Republic of China.*

[3]*Key Laboratory of Preventive Veterinary Medicine in Hubei Province, the Cooperative Innovation Center for Sustainable Pig Production, Wuhan, Hubei, People's Republic of China.*

* Corresponding author: Ping Qian: Tel: 86-27-87282608; Fax: 86-27-87282608; E-mail: qianp@mail.hzau.edu.cn. Address: No. 1, Shizishan Street, Hongshan District, Wuhan City, Hubei Province, China.

* Co-corresponding author: Xiang-min Li: Tel: 86-27-87282608; Fax: 86-27-87282608; E-mail: lixiangmin@mail.hzau.edu.cn. Address: No. 1, Shizishan

Street, Hongshan District, Wuhan City, Hubei Province, China.

Abstract

Bacillus velezensis is a species of *Bacillus* that has been widely

investigated because of its broad-spectrum antimicrobial activity. However,

most studies on *Bacillus velezensis* have focused on the biocontrol of plant

diseases, with few reports on antagonizing *Salmonella* Typhimurium infections.

In this investigation, it was discovered that *Bacillus velezensis* HBXN2020,

which was isolated from healthy black pigs, possessed strong anti-stress and

broad-spectrum antibacterial activity. Importantly, *Bacillus velezensis*

HBXN2020 did not cause any adverse side effects in mice when administered

at various doses (1×10^7, 1×10^8, and 1×10^9 CFU) for 14 d. Supplementing

Bacillus velezensis HBXN2020 spores, either as a curative or preventive

measure, dramatically reduced the levels of *Salmonella* Typhimurium

ATCC14028 in the mice's feces, ileum, cecum, and colon, as well as the

disease activity index (DAI), in a model of colitis caused by this pathogen in

mice. Additionally, supplementing *Bacillus velezensis* HBXN2020 spores

significantly regulated cytokine levels (TNF-α, IL-1β, IL-6, and IL-10) and

maintained the expression of tight junction proteins and mucin protein. Most

importantly, adding *Bacillus velezensis* HBXN2020 spores to the colonic

microbiota improved its homeostasis and increased the amount of beneficial

bacteria (*Lactobacillus* and *Akkermansia*). All together, *Bacillus velezensis*

HBXN2020 can improve intestinal microbiota homeostasis and gut barrier

integrity and reduce inflammation to help treat bacterial colitis.

Key words: *Bacillus velezensis*; antibacterial activity; *S.* Typhimurium-

induced colitis; infections; gut health

Introduction

Salmonella Typhimurium (*S.* Typhimurium) is a major foodborne zoonotic

pathogen that can cause diarrhea and colitis in humans and animals (Fabrega

and Vila, 2013; Yang et al., 2021). According to reports, *Salmonella* infections

are common in both industrialized and developing nations, posing a serious

risk to public health and resulting in significant financial losses (Katiyo et al.,

2019; Mohan et al., 2019). At present, antibiotics remain one of the most

effective treatment strategies for *Salmonella* infections. However, numerous

studies have reported that the prolonged use or misuse of antibiotics can lead

to environmental pollution, an increase in multi-drug-resistant (MDR) bacteria,

as well as gastrointestinal microbiota dysbiosis (Ferri et al., 2017; Paulson et

al., 2015; Reyman et al., 2022). In order to tackle *Salmonella* infections, a

hunt for novel antimicrobial drugs is now underway.

More research has recently demonstrated the critical role gut microbiota

plays in reducing intestinal inflammation (Cristofori et al., 2021) and mending

the intestinal barrier (Ohland and Macnaughton, 2010; Wang et al., 2020).

The use of probiotics is a popular approach to modulating intestinal microbiota

nowadays. Among all probiotics, *Bacillus* is one of the most popular probiotic

species because of its ability to form endospores that survive gastric transit,

storage and delivery conditions (Gu et al., 2015). Furthermore, *Bacillus*

species can generate extracellular enzymes and antimicrobial metabolites

that inhibit enteric pathogens, which lowers the risk of infection and enhances

nutrient utilization (Abdhul et al., 2015; Ouattara et al., 2017).

The recently identified *Bacillus velezensis* (*B. velezensis*) species of the

Bacillus genus has a large capacity to generate a variety of secondary

metabolites that possess broad-spectrum antibacterial activity (Ye et al.,

2018). Previous studies have reported that *B. velezensis* exhibits varying

degrees of probiotic effects on plants, livestock, poultry and fish. For instance,

the combined use of *B. velezensis* SQR9 mutant and FZB42 improved root

colonization and the production of secondary metabolites that are beneficial to

cucumber (Shao et al., 2022). *B. velezensis* isolated from yaks was shown to

enhance growth performance and ameliorate blood parameters related to

inflammation and immunity in mice (Li et al., 2019). The dietary *B. velezensis*

supplementation can regulate the innate immune response in the intestine of

crucian carp and reduce the degree of intestinal inflammation damage

induced by *A. veronii* (Zhang et al., 2019). Few studies have been conducted

on the treatment or prevention of colitis caused by *S.* Typhimurium; instead,

the majority of investigations on *B. velezensis* focused on the biocontrol of

fungal infections in plants.

A strain of *B. velezensis* HBXN2020 with broad-spectrum antibacterial activity was selected from a vast number of *Bacillus* strains for this study. We aimed to investigate the effectiveness of *B. velezensis* HBXN2020 in alleviating *S.* Typhimurium-induced mouse colitis and assessed the biological safety of *B. velezensis* HBXN2020. Notably, *B. velezensis* HBXN2020 can alleviate symptoms and colon tissue damage in experimental colitis, as indicated by markers such as *S.* Typhimurium loads, TNF-α and ZO-1 levels. Moreover, *B. velezensis* HBXN2020 also increases the abundance of beneficial bacteria, specifically *Lactobacillus* and *Akkermansia*, within the colon microbiota. As a result, this research supports the creation of probiotic-based microbial products as a substitute method of preventing *Salmonella* infections.

Results

Genomic Characteristics

HBXN2020's complete genome was sequenced using the Illumina Hiseq and PacBio platforms. The results showed that HBXN2020 has a circular chromosome of 3,929,792 bp (Fig. 1A) with a GC content of 46.5%, including 3,744 protein-coding genes (CDS), 86 tRNA genes, and 27 rRNA genes (Table S1). Furthermore, the HBXN2020 genome also contains 13 different clusters of secondary metabolic synthesis genes. such as Fengycin (genomic position:1,865,856) and Difficidin (genomic position:2,270,091) (Table S2).

A phylogenetic tree based on genome-wide data from all 14 *Bacillus* strains

demonstrated that the HBXN2020 belongs to the *B. velezensis* species (Fig.

1B). To further understand the classification status of HBXN2020, the online

tool JSpeciesWS was used to calculate the average nucleotide identity (ANI)

based on the BLAST (ANIb) method. As shown in Fig. S1, HBXN2020 was

found to be a member of the *B. velezensis* species due to the high percentage

of ANIb (more than 97%).

Growth curve and *in vitro* resistance against environmental assaults.

The growth of *B. velezensis* HBXN2020 was assessed in flat-bottomed 100-

well microtiter plates by measuring the values of OD_{600} every hour using an

automatic growth curve analyzer. After measurement, we found that *B.*

velezensis HBXN2020 entered the logarithmic growth phase after 2 h, and

reached a plateau after 10 h of culture (Fig. 2A).

Next, we evaluated the effect of physical, chemical, and biological

sterilization conditions such as high temperature, strong acidity, and enzyme

preparation on the survival of *B. velezensis* HBXN2020. The survival rate of *B.*

velezensis HBXN2020 showed a decreasing trend with increasing

temperature (Fig. 2B), but this decrease was not obvious. In a strong acid

environment (pH 2.0), *B. velezensis* HBXN2020 maintained a high survival

rate (Fig. 2C), suggesting that *B. velezensis* HBXN2020 spores can survive

under extreme conditions. Based on these results, we hypothesized that *B.*

velezensis HBXN2020 spores might also exhibit improved survival in gastrointestinal tract environments. Therefore, we further evaluated the survival rate of *B. velezensis* HBXN2020 in bile salt (0.3%, v/v), simulated gastric fluid (pepsin 1 mg mL^{-1}, pH 1.2), and simulated intestinal fluid (trypsin 1 mg mL^{-1}, pH 6.8). As shown in Fig. 2D-F, *B. velezensis* HBXN2020 spores demonstrated significantly improved tolerance to these simulated gastrointestinal tract environments.

Antibiotic susceptibility of *B. velezensis* HBXN2020 and bacteriostasis effect *in vitro*.

In order to assess *B. velezensis* HBXN2020's drug resistance, 19 antibiotics that are frequently used in clinical settings were chosen. The results indicated that only polymyxin B had a relatively small inhibition zone diameter (less than 15 mm). Ampicillin, meropenem, minocycline, ofloxacin, and trimethoprim had the strongest inhibition on *B. velezensis* HBXN2020, with an inhibition zone diameter exceeding 30 mm (Fig. 3A). In addition, the experimental results showed that *B. velezensis* HBXN2020 was extremely sensitive to β-lactams, tetracyclines, and quinolone drugs. Then, we performed an inhibition test of *B. velezensis* HBXN2020 against seven common clinical pathogenic bacteria. The results showed that *B. velezensis* HBXN2020 had a similar inhibitory effect on standard and wild strains of *Escherichia coli* (*E. coli*), *Salmonella*, *Staphylococcus aureus* (*S. aureus*), and *Clostridium perfringens* (*C.*

perfringens), as well as wild strains of *Streptococcus suis* (*S. suis*), *Pasteurella multocida* (*P. multocida*), and *Actinobacillus pleuropneumoniae* (*A. pleuropneumoniae*). Except for the wild strains of *S. suis* and *A. pleuropneumoniae*, the diameter of the inhibition zone of other strains was above 15 mm (Fig. 3B, S2), while the size of the inhibition zone of *S. suis* and *A. pleuropneumoniae* was also more than 12 mm.

Biosafety evaluation of *B. velezensis* HBXN2020

We assessed the biological safety of *B. velezensis* HBXN2020 in a mouse model in order to ascertain its safety. After gavage with *B. velezensis* HBXN2020 spores for two weeks, we observed no significant difference in the body weight of each group of mice (Fig. 4A). The gene expression levels of TNF-α, IL-1β, IL-6, and IL-10 of colon from all groups of mice exhibited no remarkable changes (Fig. 4B). However, in the colon, mRNA levels of the barrier proteins ZO-1 and occludin were trending towards an increase compared with the control group (Fig. 4C). Additionally, blood routine tests and serum biochemistry tests were performed for mice in the control group and H-HBXN2020 group on the 14th d after oral administration of *B. velezensis* HBXN2020 spores multiple times. As shown in Fig. 4D and S3, the blood parameters of mice after *B. velezensis* HBXN2020 treatment, including red blood cells (RBC), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), hemoglobin (HGB), white blood cells (WBC), platelets

(PLT), hematocrit (HCT), and mean corpuscular hemoglobin concentration (MCHC), were consistent with those of healthy mice. The serum biochemical parameters, including alanine aminotransferase (ALT), aspartate aminotransferase (AST), albumin (ALB), total bilirubin (TBIL), serum creatinine (CREA), and blood urea nitrogen (BUN), were also within normal limits (Fig. 4E). The corresponding histological analysis of colon tissue from mice receiving low, medium, and high doses of *B. velezensis* HBXN2020 spores (L-HBXN2020, M-HBXN2020, and H-HBXN2020 groups, respectively) is presented in Fig. 4F. The colon tissue sections of mice in the test groups showed no significant differences compared to the control group. Additionally, there were no observable differences in the major organ tissues (heart, liver, spleen, lung, and kidney) of mice treated with the high dose of *B. velezensis* HBXN2020 spores compared to healthy mice (Fig. S4). The results of this trial show that *B. velezensis* HBXN2020 is safe to use and has no negative side effects in mice.

Oral administration of *B. velezensis* HBXN2020 spores alleviated *S.* Typhimurium-induced colitis.

Testing both solid and liquid co-culture methods to see if *B. velezensis* HBXN2020 might suppress *S.* Typhimurium ATCC14028 (STm), the findings indicated that *B. velezensis* HBXN2020 decreased the development of STm in a dose-dependent manner (Fig. 5A, S5). Next, the therapeutic potential of *B.*

velezensis HBXN2020 was evaluated in an STm-induced mouse colitis model. At days 1, 3, and 5 after STm infection, mice in the STm + HBXN2020 group were orally administered *B. velezensis* HBXN2020 spores by gavage (Fig. 5B), while the control group and STm + PBS group were treated with sterile PBS. Also, the number of excreted STm in feces was counted daily at the designated time points. As shown in Fig. 5C, there was a significant and continuous reduction in the number of STm in feces following treatment with *B. velezensis* HBXN2020 spores, while the number of *B. velezensis* HBXN2020 viable bacteria in feces is also gradually decreasing (Fig. S6A). The number of STm in mouse feces was reduced by 10 to 50 times compared to the STm + PBS group from day 2 to 7 post-treatment. *B. velezensis* HBXN2020 spore treatment also resulted in a 5 to 10-fold reduction in the number of STm in the ileum, cecum, and colon, respectively (Fig. 5D-F). The increased inhibition of STm after *B. velezensis* HBXN2020 spore treatment also resulted in less weight loss and a marked reduction ($P < 0.001$) in DAI scores in STm + HBXN2020 group mice (Fig. 5G-H). In addition, we also measured the colon length of the mice and found that the STm + PBS group had a significantly shorter colon length ($P < 0.001$) than the control group. However, compared to the STm + PBS group, the STm + HBXN2020 group exhibited a longer colon ($P < 0.05$) (Fig. 5I-J).

To further assess the impact of *B. velezensis* HBXN2020 on the intestinal tract, we histologically analyzed their colonic tissues. The results showed that

mice treated with *B. velezensis* HBXN2020 spores exhibited a lower histology

score (total scoring) ($P < 0.05$) (Fig. 6A, S6B) than mice not treated with the

spores and were similar to that of healthy mice (control group). The colon

tissue of mice not treated with *B. velezensis* HBXN2020 spores displayed

higher histological scores, with significant crypt deformation, severe mucosal

damage, and obvious inflammatory cell infiltration. These findings suggested

that *B. velezensis* HBXN2020 can alleviate colon tissue damage. Moreover,

we measured the mRNA levels of inflammatory cytokines, including tumor

necrosis factor-α (TNF- α), interleukin-1 β (IL-1 β), interleukin-6 (IL-6), and

interleukin-10 (IL-10) in colon tissue using RT-qPCR analysis. As shown in

Fig. 6B-D, the mRNA levels of TNF-α ($P < 0.01$), IL-1β ($P < 0.05$), and IL-6 (P

< 0.05) in the colon tissue of STm-infected mice were significantly upregulated,

indicating the successful establishment of the STm-induced colitis model. As

expected, treatment with *B. velezensis* HBXN2020 spores led to a significant

reduction in the mRNA levels of these pro-inflammatory cytokines, while the

mRNA levels of the anti-inflammatory cytokine IL-10 were significantly

increased ($P < 0.01$) (Fig. 6E), suggesting that *B. velezensis* HBXN2020

spores can alleviate the inflammatory reaction of the colon. Moreover, the

mRNA expression levels of colonic epithelial barrier proteins, including ZO-1,

ocludin, claudin, and Muc2, were also measured by RT-qPCR. Compared

with the PBS-treated group, the *B. velezensis* HBXN2020 spores treatment

group exhibited significantly upregulated mRNA expression levels of ZO-1 (P

< 0.05), occludin (P < 0.05), claudin (P < 0.05), and Muc2 (P < 0.01) in the

colon tissue (Fig. 6F-I), further demonstrating the ability of *B. velezensis*

HBXN2020 to reduce damage to the colon epithelial barrier.

Next, we used 16S rRNA gene sequencing to investigate the impact of

HBXN2020 spores on the makeup of the intestinal microbiota in mice treated

with STm. The alpha diversity of colonic microbiota was evaluated using Sobs,

Chao1, Shannon, and Simpson indices (OTU levels). Compared with the

control and HBXN2020 group, the Sobs, Chao1, and Shannon indices of the

STm + PBS group clearly decreased (Fig. 7A-C), while the Simpson index

significantly increased (P < 0.05, Fig. 7D). Principle component analysis (PCA)

based on Bray-Curtis distance showed that compared with the STm + PBS

group, the gut microbiota composition of the STm + HBXN2020 group and the

control group were more closer together (R=0.3988, P=0.001, Fig. 7E). We

analyzed the community composition of colonic microbiota at the phylum and

genus level, and the results revealed that the composition of the gut

microbiota changed markedly after STm-induced mice. At the phylum level,

the STm + PBS group increased the abundance of *Verrucomicrobiota* and

Proteobacteria and decreased the abundance of *Firmicutes*, *Patescibacteria*,

Desulfobacterota, and *Actinobacteriota* compared to the Control group (Fig.

7F). The HBXN2020 group increased the abundance of *Bacteroidota* and

Firmicutes and decreased the abundance of *Verrucomicrobiota* compared

with the Control group. On the contrary, the relative abundance of *Firmicutes*,

Patescibacteria, *Desulfobacterota*, and *Actinobacteriota* in the STm + HBXN2020 group was significantly higher than that in the STm + PBS group. At the genus level, infection with *S.* Typhimurium ATCC14028 in the STm + PBS group dramatically reduced the relative abundance of *norank_f_Muribaculaceae*, *Desulfovibrio* and *Lactobacillus* (Fig. 7G, H-I) and enhanced the abundance of *Bacteroides*, *Escherichia-Shigella*, *Salmonella*, *Alistipes*, and *Enterococcus* compared to the Control group (Fig. 7G, J-K). The HBXN2020 group increased the abundance of *norank_f_Muribaculaceae* and *Alloprevotella* compared to the Control group. In contrast, the STm + HBXN2020 group significantly improved the relative abundance of *norank_f_Muribaculaceae* and *Lactobacillus* compared with the STm + PBS group (*P* < 0.01, *P* < 0.001, Fig. 7H-I). In addition, LEfSe analysis was also performed to identify specialized microbial communities between the STm + HBXN2020 group and the STm + PBS group and found that there was a clear difference in gut microbiota composition between them. The microbiota that played a major role in the STm + PBS group were harmful *Enterobacterales* (Fig. 7L), while the STm + HBXN2020 group was *Candidatus_Saccharimonas*.

Prophylactic *B. velezensis* HBXN2020 spores alleviated *S.* Typhimurium-induced colitis.

Based on the improved therapeutic efficacy of *B. velezensis* HBXN2020 in the treatment of STm-induced colitis in mice, we further explored its potential

for disease prevention by evaluating pretreatment. Mice in the PBS + STm

group and HBXN2020 + STm group were given the same amount of sterile

PBS and *B. velezensis* HBXN2020 spores by gavage one week in advance.

As shown in Fig. 8, compared with the PBS + STm group, oral administration

of *B. velezensis* HBXN2020 spores as a preventive measure remarkably

alleviated STm-induced colitis, including weight loss of mice ($P < 0.01$) (Fig.

8B), a significant reduction in DAI ($P < 0.001$) (the comprehensive score of

weight loss, stool consistency, and blood in the feces, Fig. 8C), and the

prevention of colon length shortening (Fig. 8D-E). However, the number of *B.*

velezensis HBXN2020 viable bacteria in feces is also gradually decreasing

with time prolonging (Fig. S8A). Furthermore, compared with the PBS + STm

group, oral administration of *B. velezensis* HBXN2020 spores not only

reduced the number of STm in mouse feces (Fig. 8F) but also decreased

STm colonization in the ileum, cecum, and colon (Fig. 8G-H, S8B).

Histological analysis further revealed that oral administration of *B.*

velezensis HBXN2020 significantly reduced the infiltration of inflammatory

cells, mucosal damage and the overall histological score of colon tissue (Fig.

9A, S8C). Additionally, the mRNA expression levels of proinflammatory

cytokines, including TNF-α ($P < 0.05$), IL-1β ($P < 0.05$) and IL-6 ($P < 0.01$), in

the colon tissue of mice pretreated with *B. velezensis* HBXN2020 spores were

significantly decreased (Fig. 9B-D), while the mRNA levels of anti-

inflammatory cytokine (IL-10) were significantly higher ($P < 0.05$) (Fig. 9E)

than mice pretreated with PBS. In addition, compared to the group pretreated

with PBS, the group pretreated with *B. velezensis* HBXN2020 spores showed

a significant increase in mRNA expression levels of ZO-1 ($P < 0.001$),

occludin ($P < 0.05$) and claudin ($P < 0.05$) in the colon tissue of mice (Fig. 9F-

H). While there was no significant difference in Muc2 expression ($p=0.252$)

(Fig. 9I), the results were consistent with those of the therapeutic experiments.

Overall, these findings suggest that *B. velezensis* HBXN2020 can effectively

alleviate the symptoms of STm-induced colitis and the associated damage to

colon tissue.

Next, we examined the impact of prophylactic *B. velezensis* HBXN2020 on

the intestinal microbiota composition of STm-treated mice. Sobs, Chao1 index

and Shannon index have changed to some extent between groups, but

differences were not significant (Fig. 10A-C). Compared with the Control

group, the Simpson index significantly decreased in the PBS + STm group (P

< 0.05, Fig. 10D). However, compared to the PBS + STm group, Simpson

index in the HBXN2020 + STm group did not change (Fig. 10D). PCA analysis

showed significant separation between control and PBS + STm groups, while

the HBXN2020 + STm group, HBXN2020 group, and the control group were

clustered together (Fig. 10E). Next, we analyzed microbial community

composition at the phylum and genus levels. At the phylum level, the PBS +

STm group increased the abundance of *Bacteroidota*, *Proteobacteria*,

Actinobacteriota, and *Deferribacterota* and reduced the abundance of

Verrucomicrobiota, *Firmicutes*, *Patescibacteria*, and *Desulfobacterota*

compared with the Control group (Fig. 10F). The HBXN2020 group increased

the abundance of *Bacteroidota* compared to the control group. In contrast,

compared to the PBS + STm group, the abundance of *Firmicutes* in the

HBXN2020+STm group was significantly increased ($P < 0.05$). At the genus

level, a depletion of *Lactobacillus* and *Akkermansia* (Fig. 10G, H-I) and an

enrichment of *Alistipes*, *Bacteroides*, *Escherichia-Shigella*, *Alloprevotella*,

Helicobacter, *Enterococcus*, and *Salmonella* were observed in PBS + STm

group compared with the control (Fig. 10G, J-M). The enrichment with

norank_f_Muribaculaceae and *Alloprevotella* was observed in the HBXN2020

group compared to the control. In contrast, an enrichment of *Lactobacillus*,

Akkermansia, and *Lachnospiraceae_NK4A136_group* and a sharp decrease

of *Alistipes*, *Bacteroides*, *Escherichia-Shigella*, *Salmonella*, and *Enterococcus*

were observed in HBXN2020 + STm group compared to the PBS + STm

group. LEfSe analysis showed that the abundance of *Lactobacillus* in the

intestinal microbiota of mice was significantly increased after *B. velezensis*

HBXN2020 treatment, while the abundance of harmful *Enterobacterales* in the

untreated group was clearly increased (Fig. 10N).

Discussion

S. Typhimurium is an important intestinal pathogen that can cause invasive

intestinal diseases such as bacterial colitis (Herp et al., 2019; Schultz et al.,

2017). Antibiotics are frequently used to treat colitis caused by bacteria; nevertheless, in recent years, their abuse has greatly increased bacterial resistance, leading to serious environmental pollution. A recent study showed that antibiotic-resistance genes in probiotics could be transmitted to the intestinal microbiota, which may threaten human health (Crits-Christoph et al., 2022). It is crucial to consider the source of probiotics. Strong environmental adaptability and disease resistance make black pigs an outstanding household breed in China (Yang et al., 2022). In this study, *B. velezensis* HBXN2020 was isolated from the free-range feces of black piglets in a mountain village in Xianning City (Hubei, China) and exhibited excellent antibacterial activity. Otherwise, *in vitro* tolerance assays showed that *B. velezensis* HBXN2020 spores have good tolerance to high temperature, strong acids, bile salts, as well as simulated gastric and intestinal fluid, which was similar to previous research results (Du et al., 2022). Furthermore, studies for antibiotic susceptibility revealed that *B. velezensis* HBXN2020 is not resistant to antibiotics.

Probiotics are distinct from food or medications in that they can cause toxins or infections in the body when ingested, and they are living when swallowed (Sanders et al., 2010; Snydman, 2008). Therefore, assessing the biosafety of probiotics is crucial. In this study, the safety of *B. velezensis* HBXN2020 was evaluated by measuring the body weight, intestinal barrier proteins, and inflammatory cytokines of the experimented mice, as well as

conducting routine blood and biochemical tests. The results showed that the

expression levels of ZO-1 and occludin in the colon of mice increased after

treatment with different doses of *B. velezensis* HBXN2020 spores. It has been

reported that ZO-1, occludin and claudin are the three most important tight

junction proteins in intercellular connections, which play a crucial role in

maintaining the intestinal epithelial barrier (Bazzoni et al., 2000; Zihni et al.,

2016). Thus, it is possible to improve intestinal barrier function by upregulating

ZO-1 and occludin expression levels.

Important aminotransferases in animals, aspartate aminotransferase (AST)

and alanine aminotransferase (ALT), are regarded as critical metrics for

assessing liver function damage (Gou et al., 2022; Ozer et al., 2008). Under

normal circumstances, the levels of ALT and AST are in dynamic balance

without notable changes. However, when the liver is damaged or becomes

dysfunctional, the levels of aminotransferases (ALT and AST) increase

significantly (Tang et al., 2012). Previous studies have shown that CCL4-

induced liver injury strongly increases plasma ALT/AST levels (Singhal et al.,

2018). Zhang et al. reported that the addition of *Bacillus subtilis* to chicken

diets significantly decreased the serum levels of ALT and AST (Zhang et al.,

2017). In this present study, the levels of ALT and AST in the probiotic-treated

group were slightly lower than those in the control group, indicating that *B.*

velezensis HBXN2020 had no negative effect on liver health in mice. The *B.*

velezensis HBXN2020 treatment group's blood indices were also found to be

comparable to those of the control group, falling within the normal reference

range, according to regular blood testing.

Mice were fed with *B. velezensis* HBXN2020 spores orally after being

challenged with *S.* Typhimurium ATCC14028 in order to assess the spores'

potential to alleviate colitis caused by S. Typhimurium. Our experimental

results demonstrated that oral treatment with *B. velezensis* HBXN2020 spores

could alleviate *S.* Typhimurium-induced colitis, as evidenced by decreased

weight loss, DAI, and histological damage, which is similar to prior studies

(Cao et al., 2019; Wu et al., 2022). Previous research showed that

macrophages are the first line of host defense against bacterial infection.

They release proinflammatory cytokines, which are critical in initiating

adaptive immune responses (Cheng et al., 2014; Dinarello, 2000).

Nevertheless, proinflammatory cytokines have immunological properties that

can be beneficial for the host to resist the invasion of bacteria and other

microbes in the surrounding environment (Liu et al., 2021). For instance,

pretreatment with recombinant murine TNF-α was shown to protect mice

against lethal bacterial (*E. coli*) infection (Cross et al., 1989). PJ-34 exerted

protective effects on intestinal epithelial cells against invasive *Salmonella*

infection by upregulating IL-6 expression through the ERK and NF-κB signal

pathways (Huang, 2009).

On the other hand, some research has demonstrated that over-activation of

immune cells can result in tissue damage, organ failure, systemic or

persistent inflammation, and autoimmune diseases (Karki et al., 2021; Kotas

and Medzhitov, 2015). IL-10 is a recognized anti-inflammatory mediator that

plays a crucial role in maintaining intestinal microbe-immune homeostasis,

regulating the release of inflammatory mediators, and inhibiting

proinflammatory responses of innate and adaptive immunity (Maloy and

Powrie, 2011; Ouyang et al., 2011; Saraiva et al., 2020). For instance,

previous studies have shown that *Clostridium butyrate* can induce the

production of IL-10 in the intestine, thereby alleviating experimental colitis in

mice (Hayashi et al., 2013). Similar results were observed in our study, where

oral administration of *B. velezensis* HBXN2020 spores reduced the

expression levels of TNF-α, IL-1β, and IL-6 while increasing the levels of IL-10

in the colon of mice. Additionally, as demonstrated by the decreased

histological damage and increased expression levels of intestinal barrier

proteins, oral treatment of *B. velezensis* HBXN2020 spores reduced the

functional damage of the intestinal barrier induced by S. Typhimurium

infection.

Subsequently, we conducted additional analysis on the colon microbiome.

Notably, oral *B. velezensis* HBXN2020 significantly raised the colonic

microbiota's alpha diversity. As reported in prior studies, elevated alpha

diversity is associated with enhanced anti-interference ability (Chen et al.,

2021). PCA analysis results showed that oral administration of *B. velezensis*

HBXN2020 significantly affected the structure of the intestinal microbiota.

Further analysis at the phylum/genus level revealed that *B. velezensis* HBXN2020 treatment reduced the increase of *Bacteroides*, *Escherichia-Shigella*, and *Salmonella* (harmful to intestinal homeostasis) in colitis (Wang et al., 2023) but enhanced the relative abundance of *Lactobacillus* (known for enhancing the epithelial barrier by increasing mucus secretion and upregulating the expression of tight junction proteins such as claudin-1, occludin, and ZO-1) (Kaur et al., 2021) and *Akkermansia* (known for modulating intestinal immune response by producing short chain fatty acids and reducing the secretion of proinflammatory cytokines) (Cani et al., 2022). *Bacteroides* are generally 'friendly' commensals in the intestine, but they can enter normal tissues through the intestinal mucosa when intestinal barrier function becomes compromised (Polk and Kasper, 1977), leading to inflammation, diarrhea, and abdominal abscesses (Pickard et al., 2017). *Escherichia-Shigella* is a common intestinal pathogen, and its relative abundance is associated with proinflammatory states (Morgan, 2013). *Salmonella* can secrete bacterial virulence proteins, which are directly delivered to the host cell cytosol through the Type III secretion system and induce the expression of pro-inflammatory cytokines (Hardt et al., 1998), leading to inflammation. Overall, the *S.* Typhimurium ATCC14028 infection altered the structure of the colon microbiota and raised the relative abundance of pathogenic bacteria, whereas the therapeutic *B. velezensis* HBXN2020 increased the abundance of beneficial bacteria, decreased inflammation, and

enhanced the function of the intestinal barrier.

We assessed pretreatment in order to learn more about the potential of *B.*

velezensis HBXN2020 spores for disease prevention, given the increased

efficacy of treating colitis caused by S. Typhimurium. Similarly to the

therapeutic effect of oral *B. velezensis* HBXN2020, pretreatment with *B.*

velezensis HBXN2020 spores effectively alleviated the symptoms of *S.*

Typhimurium-induced colitis and reduced the expression levels of

proinflammatory cytokines and increased anti-inflammatory cytokine levels in

the colon of *S.* Typhimurium-infected mice. Furthermore, pretreatment with *B.*

velezensis HBXN2020 spores also promoted the expression levels of

intestinal barrier proteins in the colon of *S.* Typhimurium-infected mice. In

addition, the microbiota composition was also dramatically modulated by

prophylactic *B. velezensis* HBXN2020 in mice with colitis, although the

community richness and diversity revealed by the alpha diversity index were

not significantly impacted in mice with colitis by prophylactic treatment.

Prophylactic *B. velezensis* HBXN2020 significantly boosted SCFA-producing

bacteria, including *Lactobacillus* and *Akkermansia*, according to additional

genus-level analysis, which was consistent with the effects of therapeutic *B.*

velezensis HBXN2020.

Similarly, harmful bacteria such as *Alistipes*, *Bacteroides*, *Escherichia-*

Shigella, and *Enterococcus* were significantly enriched in the PBS+STm

group. A facultative-pathogenic bacterium called Alistipes encourages the

growth of inflammation, obesity, and colorectal cancer (Parker et al., 2020).

Based on those results, we found that prophylactic *B. velezensis* HBXN2020

alleviates *S.* Typhimurium-induced colitis by increasing the abundance of

beneficial bacteria, attenuating proinflammatory responses of the intestine,

and increasing intestinal barrier function.

Conclusion

In conclusion, gut mucosal barrier disruption and intestinal inflammation

results from dysbiosis of the colonic microbiota caused by S. Typhimurium

ATCC14028 infection. Supplementing *Bacillus velezensis* HBXN2020 can

improve intestinal microbiota stability and gut barrier integrity and reduce

inflammation to help prevent or treat bacterial colitis.

Experimental procedures

Animals and bacterial strains.

Female specific-pathogen-free (SPF) C57BL/6 mice aged 6 to 8 weeks

were purchased from the Animal Experimental Center of Huazhong

Agricultural University. All mice experiments were conducted in the standard

SPF facility of Huazhong Agricultural University, with 12 h of light and 12 h of

darkness at a temperature of 25°C and ad libitum access to food and water.

The use of animals in this experiment was approved by the Animal Care and

Ethics Committee of Huazhong Agricultural University (Ethics Approval

Number: HZAUMO-2023-0089).

The strains used in this study are listed in Table S3, and all the primers

used in Table S4. *Clostridium perfringens* were cultured in a fluid thioglycollate

medium (FTG) (Hopebio, Qingdao, China) at 45°C. *Streptococcus suis* and

Pasteurella multocida were cultured in tryptic soy broth (TSB) (BD

Biosciences, MD, United States) medium supplemented with 5% (v/v) sheep

serum (Solarbio, Beijing, China) at 37°C, and *Actinobacillus*

pleuropneumoniae were cultured in TSB supplemented with 5% (v/v) sheep

serum and 5mM nicotinamide adenine dinucleotide (NAD) at 37°C, while the

other bacterial strains were cultured in Luria Bertani (LB) medium (Solarbio,

Beijing, China) at 37°C. In addition, the Difco sporulation medium (DSM) was

used for inducing the sporulation of *B. velezensis* HBXN2020 via the nutrient

depletion method (Tang et al., 2017).

Stain isolation, growth conditions and strain identification.

In this study, we isolated 362 *Bacillus* strains from the feces of healthy

piglets, and four *Bacillus* strains were screened through initial antibacterial

tests. Then, a strain with broad-spectrum antimicrobial activity was ultimately

screened out by the antibacterial spectrum assay (Table S5), and named as

HBXN2020. The strain was stored in 28% glycerol at –80°C. For culturing the

bacterial cells, Luria-Bertani (LB) media or agar plates supplemented with 1.5%

agar were used and incubated at 37°C.

A single bacterial colony was inoculated into 10 mL of LB medium and

incubated overnight at 37°C with shaking (180 rpm). The bacteria genomic

DNA was extracted using the TIANamp Genomic DNA Kit (TIANGEN, Beijing,

China) from bacteria. Then, 16S ribosomal sequence was amplified using

16S-specific primers and sequenced to identify the HBXN2020.

Genome sequencing, annotation and analysis

The complete genome of *B. velezensis* HBXN2020 was sequenced using

the Illumina HiSeq PE and PacBio RSII (SMRT) platforms (Shanghai Majorbio

Bio-pharm Technology Co., Ltd.). Briefly, the short reads from the Illumina

HiSeq PE were assembled into contigs using SPAdenovo

(http://soap.genomics.org.cn/). The long reads from the PacBio RSII and the

Illumina contigs were then aligned using the miniasm and Racon tools in

Unicycler (version 0.4.8, https://github.com/rrwick/Unicycler) to generate long-

read sequences. During the assembly process, sequence correction was

performed using Pilon (version 1.22,

https://github.com/broadinstitute/pilon/wiki/Standard-Output). Lastly, a

complete genome with seamless chromosomes was obtained.

The coding sequences (CDs) of the *B. velezensis* HBXN2020 genome were

predicted using Glimmer (version 3.02,

http://ccb.jhu.edu/software/glimmer/index.shtml). The tRNA and rRNA genes

were predicted using TRNAscan-SE (version 2.0,

http://trna.ucsc.edu/software/) and Barrnap (version 0.8,

https://github.com/tseemann/barrnap), respectively. The functional annotation

of all CDSs was performed using various databases, including Swiss-Prot

Database (https://web.expasy.org/docs/swiss-prot_guideline.html), Pfam

Database (http://pfam.xfam.org/), EggNOG Database (http://eggnog.embl.de/),

GO Database (http://www.geneontology.org/), and KEGG Database

(http://www.genome.jp/kegg/). The circular map of the *B. velezensis*

HBXN2020 genome was generated using CGView (version 2,

http://wishart.biology.ualberta.ca/cgview/) (Stothard and Wishart, 2005).

Comparative genomic analysis

To elucidate the phylogenetic relationships from a whole-genome

perspective, a phylogenetic tree based on the whole genome was constructed

using the Type (Strain) Genome Server (TYGS) online (https://ggdc.dsmz.de/)

(Meier-Kolthoff et al., 2022). The average nucleotide identity (ANI) values of

genome to genome were calculated using the JSpeciesWS online service

(Richter et al., 2016), and a heatmap was then generated using TBtools

(version 1.113) (Chen et al., 2020).

Growth curves of *B. velezensis* HBXN2020

The growth curve of *B. velezensis* HBXN2020 was recorded in flat-

bottomed 100-well microtiter plates via detecting optical density at 600 nm

(OD_{600}) at 1 h intervals using the automatic growth curve analyzer (Bioscreen,

Helsinki, Finland).

Antimicrobial Assays

The *in vitro* antagonistic activity of *B. velezensis* HBXN2020-CFS was

tested using the agar well-diffusion method against 18 indicator strains

(pathogens), which included 7 standard strains (*E. coli* ATCC 25922, *E. coli*

ATCC 35150, *S.* Typhimurium ATCC 14028 (STm), *S.* Typhimurium SL1344,

S. aureus ATCC 29213, *S. aureus* ATCC 43300, and *C. perfringens* CVCC

2030) and 11 clinical isolates (*E. coli* EC024, *S.* Enteritidis SE006, *S. aureus*

S21, *C. perfringens* CP023, *C. perfringens* CP002, *S. suis* SC19, *S. suis*

SS006, *P. multocida* PM002, *P. multocida* PM008, *A.* pleuropneumoniae

APP015, and *A.* pleuropneumoniae APP017). All plates were cultured at 37°C

for 16 h before observing the inhibition zone, and the diameter of the inhibition

zone was measured using a vernier caliper. The presence of a clear zone

indicated antagonistic activity.

Antibiotic susceptibility assays

Antimicrobial susceptibility testing was performed using the Kirby-Bauer

(KB) disk diffusion method in accordance with the Clinical Laboratory

Standards Institute (CLSI) guidelines (CLSI., 2018.). The antimicrobial agents

tested were: Ampicillin, Meropenem, Piperacillin, Gentamycin, Tetracycline,

Doxycycline, Minocycline, Erythromycin, Enrofloxacin, Ofloxacin,

Sulfamethoxazole, Trimethoprim-sulfamethoxazole, Polymyxin B, Teicoplanin,

Trimethoprim, Florfenicol, Spectinomycin, Nitrofurantoin, and Rifampicin. The

diameter of the inhibition zone was measured using a vernier caliper.

In vitro resistance assay of _B. velezensis_ HBXN2020

B. velezensis HBXN2020 spores (100 µL) or vegetative cells (100 µL) were

separately resuspended in 900 µL of LB medium supplemented with different

pH values (2, 3, 4, 5 or 6), bile salts (0.85% NaCl, 0.3%), simulated gastric

fluid (SGF, HCl, pH 1.2) containing 10 g L^{-1} of pepsin in 0.85% NaCl solution,

or simulated intestinal fluid (SIF, NaOH, pH 6.8) containing 10 g L^{-1} of trypsin

in 0.05 M KH2PO4 solution, and incubated at 37 °C. A normal LB medium (pH

7.0) was used as the control. At predetermined time points, 100 µL was taken

from each sample, serially diluted 10-fold with sterile PBS (pH 7.2), and then

spread onto LB agar plates. The plates were incubated overnight in a

constant temperature incubator at 37°C, and the bacterial colonies were

counted. The survival rate was calculated using the following formula: Survival

rate = (number of bacteria in the treatment group/number of bacteria in the

control group) × 100%.

One milliliter of _B. velezensis_ HBXN2020 spores or vegetative cells were

separately placed in water baths at different temperatures (37°C, 45°C, 55°C,

65°C, 75°C, 85°C or 95°C) for 20 min, with a 37°C water bath used as the

control. The survival rate was calculated as described above.

***In vitro* bacterial competition.**

To investigate whether *B. velezensis* HBXN2020 directly inhibited the

growth of STm, we performed spot-on lawn and agar-well diffusion assays, as

well as co-culture assays in liquid culture medium. In the spot-on lawn

antimicrobial assays, we prepared double layers of agar by first pouring LB

agar into the plate as the bottom layer. The top layer consisted of 10 mL TSB

broth containing 0.7% agar with STm overnight culture. Then, 10 μL of *B.*

velezensis HBXN2020 overnight culture and cell-free supernatant (CFS) were

respectively spotted onto TSB agar and incubated at 37°C for 12 h to

measure the inhibition zone. A transparent zone of at least 1 mm around the

spot was considered positive.

The antagonistic effect of *B. velezensis* HBXN2020-CFS against STm was

determined using the agar-well diffusion assays. To collect *B. velezensis*

HBXN2020-CFS, the culture was centrifuged at 9,000g for 15 min at 4°C, and

the supernatant was filtered through a 0.22 μm membrane filter (Millipore,

USA). The STm lawn medium was prepared by mixing 10 mL of TSB broth

containing 0.7% agar with STm overnight culture and then poured into a

sterile plate covered with LB agar and Oxford cups, 8 mm diameter wells were

prepared in the TSB agar after removing the cups. The wells were filled with

100 μL of *B. velezensis* HBXN2020-CFS, LB medium or ampicillin (100 μg

mL^{-1}). The plates were incubated at 37°C for 14 h, and the inhibition zone was measured.

The co-culture assay was conducted by incubating *B. velezensis* HBXN2020 and *S.* Typhimurium ATCC14028 (carry pET28a (+), kanamycin resistance) (Cao et al., 2019) separately overnight at 37°C and diluting them to 10^4 CFU/mL. Then, the two strains were mixed at different ratios (1:1, 1:10, 1:50 or 1:100) and co-cultured at 37°C with shaking (180 rpm). At predetermined time points, serial 10-fold dilutions were prepared for all samples and spread onto selective (kanamycin 50 µg mL^{-1}) LB agar plates and cultured at 37°C for 12 h before bacterial counting. Viable colony counts ranged from 30 to 300 per plate.

Biosafety assessment

Biosafety assay of *B. velezensis* HBXN2020 referred to the method described by Zhou et al.(Zhou et al., 2022) with slight modifications. After a 7-day acclimation period (free access to water and food), mice were randomly divided into four treatment groups (n = 5): low-dose *B. velezensis* HBXN2020 spores group (L-HBXN2020 group, 10^7 CFU/mouse), medium-dose *B. velezensis* HBXN2020 spores group (M-HBXN2020 group, 10^8 CFU/mouse), high-dose *B. velezensis* HBXN2020 spores group (H-HBXN2020 group, 10^9 CFU/mouse), and a control group. During the experimental period (15 days), mice were weighed and orally gavaged with their respective treatments once

every two days. On day 15, all mice were euthanized, and blood, heart, liver,

spleen, lung, kidney, ileum, cecum, and colon were collected. Blood samples

were used for routine blood and biochemistry tests. Major organ tissues (heart,

liver, spleen, lung, and kidney) and a 5-mm distal segment of the colon were

used for histopathology. The remaining colonic tissues were rapidly frozen in

liquid nitrogen and stored at –80°C for cytokine and tight junction protein

expression analysis.

Salmonella Typhimurium-induced mouse model of colitis.

After 7 days of acclimation (free access to water and food), mice were

randomly divided into three treatment groups (n = 8): Control group, STm +

PBS group, and STm + HBXN2020 group. The *S.* Typhimurium-induced colitis

was performed as previously described (Stecher et al., 2005), with slight

modifications. On the first day of the experiment, all mice in the STm + PBS

group and STm + HBXN2020 group were orally inoculated with 200 μL (5×10^{7}

CFU/mouse) of STm. On days 1, 3 and 5 following STm infection, each

mouse in the STm + HBXN2020 group received 200 μL (1×10^{8} CFU/mouse)

of *B. velezensis* HBXN2020 spores via gavage administration. In contrast, the

control group and STm + PBS group received 200 μL of sterile PBS by oral

gavage. Fecal samples were collected daily following STm infection and

resuspended in sterile PBS. The number of STm in mice feces from both the

STm + PBS and STm + HBXN2020 groups was then determined by

spreading a serial 10-fold dilution on selective LB agar plates containing 50 μg

mL^{-1} kanamycin. Throughout the entire experiment, the body weight, stool

consistency and fecal occult blood of all mice were monitored daily. As shown

in Table S6, disease activity index (DAI) was calculated by the sum of the

scores from three parameters (Praveschotinunt et al., 2019). On day 7 after

STm infection, all mice were euthanized, their ileum, cecum and colon were

collected. The length of colon was measured, and a 5-mm distal segment of

the colon was fixed in 4% paraformaldehyde for further histopathology. The

remaining colon was then rapidly frozen in liquid nitrogen and stored at –80°C

for cytokine and tight junction protein expression analysis. Then, the number

of STm in the ileum, cecum, and colon was determined by spreading serial

10-fold dilutions on selective LB agar plates.

To investigate the prophylactic efficacy of *B. velezensis* HBXN2020 in

ameliorating STm-induced colitis, another independent experiment was

conducted using six-week-old female C57BL/6 mice (SPF). After a seven-day

acclimation period, mice were randomly assigned to three groups (n = 8):

control group, PBS + STm group and HBXN2020 + STm group. On days 1, 3,

5 and 7, each mouse in the HBXN2020 + STm group received 200 μL (1×10^{8}

CFU/mouse) of *B. velezensis* HBXN2020 spores via gavage administration.

Meanwhile, mice in the control group and PBS + STm group received 200 μL

of sterile PBS by oral gavage. On day 7, all mice in the PBS + STm group and

HBXN2020 + STm group were orally inoculated with 200 μL (5×10^{7}

CFU/mouse) of STm. Fecal samples were collected daily following STm

infection from both groups and resuspended in sterile PBS. The number of

STm in the feces of mice was then determined by spreading serial 10-fold

dilutions on selective LB agar plates (50 μg mL^{-1} kanamycin). Throughout the

entire experiment, the body weight and DAI scores of all mice were monitored

daily. On day 12, all mice were euthanized, and the ileum, cecum, and colon

were collected. The colon length was measured, and a 5-mm distal segment

of the colon was fixed in 4% formalin for sectioning and staining. The

remaining colon was stored at −80°C for future analysis. Lastly, the number of

STm in the ileum, cecum, and colon was determined using a selective LB

agar plate.

Determination of cytokines and tight junction protein expression in colon tissue

Total RNA was extracted from colon tissues using TRIpure reagent (Aidlab,

China), and cDNA was obtained using HiScript$^®$ III RT SuperMix for qPCR

(+gDNA wiper) (Vazyme, China). RT-qPCR for each gene was performed in

triplicate using qPCR SYBR Green Master Mix (Yeasen Biotechnology,

Shanghai, China). The relative expression level of cytokine and tight junction

protein genes was calculated using the $2^{-\Delta\Delta Ct}$ method with β-actin and

GAPDH as reference genes. The primer sequences used in the RT-qPCR test

are listed in Table S4.

Histopathology analysis

Colon tissue samples (0.5 cm) were fixed in 4% paraformaldehyde for 24 h, and the fixed tissues were then embedded in paraffin and sectioned. The sections were stained with hematoxylin and eosin (H&E) and observed and imaged using an optical microscope (Olympus Optical, Tokyo, Japan). The histopathological score included the degree of inflammatory infiltration, changes in crypt structures, and the presence or absence of ulceration and edema. The scoring criteria were determined as previously described (Wu et al., 2022).

16S rRNA gene sequencing and analysis

According to the manufacturer's instructions, colon microbial community genomes were extracted using E.Z-N.A ® Stool DNA Kit (Omega; D4015-01), and quality was detected by 1% agarose gel electrophoresis. The 16S rRNA V3-V4 variable region was amplified by PCR using universal primers 338F (5 '- ACTCCTACGGGGGGGCAG-3') and 806R (5 '- GACTACHVGGGTWTCTAAT-3'). The PCR products were examined by electrophoresis on 2% agarose gels and then purified with the AxyPrep DNA gel extraction kit (Axygen Biosciences, USA). A sequencing library was constructed using the NEXTFLEX® Rapid DNA-Seq Kit, and sequencing was performed using the Illumina MiSeq platform. Raw reads were quality

evaluated and filtered by fastp (version 0.20.0) and merged using FLASH

(version 1.2.7). The optimized sequences were clustered into operational

taxonomic units (OTUs) based on 97% sequence similarity using UPARSE

(version 7.1). The representative sequences of each OTU was classified by

RDP classifier (version 2.2; confidence threshold value, 0.7). Alpha diversity

was assessed using the ACE, chao, Shannon, and Simpson indices. The β-

diversity analysis was performed using Bray-Curtis distances and visualized

through principal component analysis (PCA). Linear discriminant analysis

(LDA) effect size (LefSe) was used to identify differential microbiota between

groups.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism 8.3.0 (GraphPad

Software, San Diego, CA, USA) and Excel (Microsoft, Redmond, USA). Data

are presented as mean ± standard error of the mean (SEM). Differences

between two groups were evaluated using two-tailed unpaired Student's t-test,

and all other comparisons were conducted using one-way analysis of variance

(ANOVA) with Tukey's multiple comparisons test. For all analyses,

significance differences are denoted as: *, $P < 0.05$, **, $P < 0.01$, and ***, $P <$

0.001.

Data availability

The datasets supporting the conclusions of this article were deposited in the

NCBI Sequence Read Archive database under the accession numbers:

CP119399.1 (whole genome sequencing raw data) and PRJNA1045315

(microbiota raw sequencing data).

Conflict of interest: We declare no conflict of interest.

ACKNOWLEDGMENTS

This work was supported by grants from the National Program on Key

Research Project of China (2021YFD1800300, 2022YFD1800800), "Yingzi

Tech & Huazhong Agricultural University Intelligent Research Institute of Food

Health" (No. IRIFH202209; IRIFH202301), and the Fundamental Research

Funds for the Central Universities (2662016PY004).

Author contributions

Linkang Wang: Conceptualization, Methodology, Data curation, Formal

analysis, Writing - original draft. **Haiyan Wang, Xinxin Li, Mengyuan Zhu**:

Methodology, Data curation, Planned and performed the experiments.

Dongyang Gao, Dayue Hu, Zhixuan Xiong: Methodology, Writing -

Reviewing and Editing. **Xiangmin Li**: Supervision, Writing-reviewing and

editing. **Ping Qian**: Conceptualization, Funding acquisition, Project

administration, Supervision, Writing-Reviewing and Editing. All authors read

and approved the submitted version of the paper.

Funding information

This work was supported by grants from the National Program on Key

Research Project of China (2021YFD1800300, 2022YFD1800800), "Yingzi

Tech & Huazhong Agricultural University Intelligent Research Institute of Food

Health" (No. IRIFH202209; IRIFH202301), and the Fundamental Research

Funds for the Central Universities (2662016PY004).

Reference

Abdhul, K., Ganesh, M., Shanmughapriya, S., Vanithamani, S., Kanagavel, M., Anbarasu, K., and
Natarajaseenivasan, K. (2015). Bacteriocinogenic potential of a probiotic strain Bacillus coagulans
[BDU3] from Ngari. Int J Biol Macromol *79*, 800-806.
Bazzoni, G., Martinez-Estrada, O.M., Orsenigo, F., Cordenonsi, M., Citi, S., and Dejana, E. (2000).
Interaction of junctional adhesion molecule with the tight junction components ZO-1, cingulin, and
occludin. J Biol Chem *275*, 20520-20526.
Cani, P.D., Depommier, C., Derrien, M., Everard, A., and de Vos, W.M. (2022). Akkermansia muciniphila:
paradigm for next-generation beneficial microorganisms. Nat Rev Gastroenterol Hepatol *19*, 625-637.
Cao, Z., Wang, X., Pang, Y., Cheng, S., and Liu, J. (2019). Biointerfacial self-assembly generates lipid
membrane coated bacteria for enhanced oral delivery and treatment. Nat Commun *10*, 5783.
Chen, C., Chen, H., Zhang, Y., Thomas, H.R., Frank, M.H., He, Y., and Xia, R. (2020). TBtools: An
Integrative Toolkit Developed for Interactive Analyses of Big Biological Data. Mol Plant *13*, 1194-1202.
Chen, L., Wang, D., Garmaeva, S., Kurilshikov, A., Vich Vila, A., Gacesa, R., Sinha, T., Segal, E., Weersma,
R.K., Wijmenga, C., *et al.* (2021). The long-term genetic stability and individual specificity of the
human gut microbiome. Cell *184*, 2302-2315.e2312.
Cheng, A., Yan, H., Han, C., Wang, W., Tian, Y., and Chen, X. (2014). Polyphenols from blueberries
modulate inflammation cytokines in LPS-induced RAW264.7 macrophages. Int J Biol Macromol *69*,
382-387.
CLSI. (2018.). Performance standards for antimicrobial disk susceptibility tests, M02, 13th ed. Clinical
and Laboratory Standards Institute, Wayne, PA.
Cristofori, F., Dargenio, V.N., Dargenio, C., Miniello, V.L., Barone, M., and Francavilla, R. (2021). Anti-
Inflammatory and Immunomodulatory Effects of Probiotics in Gut Inflammation: A Door to the Body.
Front Immunol *12*, 578386.
Crits-Christoph, A., Hallowell, H.A., Koutouvalis, K., and Suez, J. (2022). Good microbes, bad genes?

The dissemination of antimicrobial resistance in the human microbiome. Gut Microbes *14*, 2055944.

Cross, A.S., Sadoff, J.C., Kelly, N., Bernton, E., and Gemski, P. (1989). Pretreatment with recombinant
murine tumor necrosis factor alpha/cachectin and murine interleukin 1 alpha protects mice from
lethal bacterial infection. J Exp Med *169*, 2021-2027.

Dinarello, C.A. (2000). Proinflammatory cytokines. Chest *118*, 503-508.

Du, H., Yao, W., Kulyar, M.F., Ding, Y., Zhu, H., Pan, H., Li, K., Bhutta, Z.A., Liu, S., and Li, J. (2022). Effects
of Bacillus amyloliquefaciens TL106 Isolated from Tibetan Pigs on Probiotic Potential and Intestinal
Microbes in Weaned Piglets. Microbiol Spectr *10*, e0120521.

Fabrega, A., and Vila, J. (2013). Salmonella enterica serovar Typhimurium skills to succeed in the host:
virulence and regulation. Clin Microbiol Rev *26*, 308-341.

Ferri, M., Ranucci, E., Romagnoli, P., and Giaccone, V. (2017). Antimicrobial resistance: A global
emerging threat to public health systems. Crit Rev Food Sci Nutr *57*, 2857-2876.

Gou, S., Chen, N., Wu, X., Zu, M., Yi, S., Ying, B., Dai, F., Ke, B., and Xiao, B. (2022). Multi-responsive
nanotheranostics with enhanced tumor penetration and oxygen self-producing capacities for
multimodal synergistic cancer therapy. Acta Pharm Sin B *12*, 406-423.

Gu, S.B., Zhao, L.N., Wu, Y., Li, S.C., Sun, J.R., Huang, J.F., and Li, D.D. (2015). Potential probiotic
attributes of a new strain of Bacillus coagulans CGMCC 9951 isolated from healthy piglet feces. World
J Microbiol Biotechnol *31*, 851-863.

Hardt, W.D., Chen, L.M., Schuebel, K.E., Bustelo, X.R., and Galán, J.E. (1998). S. typhimurium encodes
an activator of Rho GTPases that induces membrane ruffling and nuclear responses in host cells. Cell
93, 815-826.

Hayashi, A., Sato, T., Kamada, N., Mikami, Y., Matsuoka, K., Hisamatsu, T., Hibi, T., Roers, A., Yagita, H.,
Ohteki, T., *et al.* (2013). A single strain of Clostridium butyricum induces intestinal IL-10-producing
macrophages to suppress acute experimental colitis in mice. Cell Host Microbe *13*, 711-722.

Herp, S., Brugiroux, S., Garzetti, D., Ring, D., Jochum, L.M., Beutler, M., Eberl, C., Hussain, S., Walter, S.,
Gerlach, R.G., *et al.* (2019). Mucispirillum schaedleri Antagonizes Salmonella Virulence to Protect Mice
against Colitis. Cell Host Microbe *25*, 681-694 e688.

Huang, F.C. (2009). Upregulation of Salmonella-induced IL-6 production in Caco-2 cells by PJ-34, PARP-
1 inhibitor: involvement of PI3K, p38 MAPK, ERK, JNK, and NF-kappaB. Mediators Inflamm *2009*,
103890.

Karki, R., Sharma, B.R., Tuladhar, S., Williams, E.P., Zalduondo, L., Samir, P., Zheng, M., Sundaram, B.,
Banoth, B., Malireddi, R.K.S., *et al.* (2021). Synergism of TNF-alpha and IFN-gamma Triggers
Inflammatory Cell Death, Tissue Damage, and Mortality in SARS-CoV-2 Infection and Cytokine Shock
Syndromes. Cell *184*, 149-168 e117.

Katiyo, S., Muller-Pebody, B., Minaji, M., Powell, D., Johnson, A.P., De Pinna, E., Day, M., Harris, R., and
Godbole, G. (2019). Epidemiology and Outcomes of Nontyphoidal Salmonella Bacteremias from
England, 2004 to 2015. J Clin Microbiol *57*.

Kaur, H., Gupta, T., Kapila, S., and Kapila, R. (2021). Protective effects of potential probiotic
Lactobacillus rhamnosus (MTCC-5897) fermented whey on reinforcement of intestinal epithelial
barrier function in a colitis-induced murine model. Food Funct *12*, 6102-6116.

Kotas, M.E., and Medzhitov, R. (2015). Homeostasis, inflammation, and disease susceptibility. Cell *160*,
816-827.

Li, A., Wang, Y., Li, Z., Qamar, H., Mehmood, K., Zhang, L., Liu, J., Zhang, H., and Li, J. (2019). Probiotics
isolated from yaks improves the growth performance, antioxidant activity, and cytokines related to

immunity and inflammation in mice. Microb Cell Fact *18*, 112.

Liu, C., Chu, D., Kalantar-Zadeh, K., George, J., Young, H.A., and Liu, G. (2021). Cytokines: From Clinical

Significance to Quantification. Adv Sci (Weinh) *8*, e2004433.

Maloy, K.J., and Powrie, F. (2011). Intestinal homeostasis and its breakdown in inflammatory bowel

disease. Nature *474*, 298-306.

Meier-Kolthoff, J.P., Carbasse, J.S., Peinado-Olarte, R.L., and Goker, M. (2022). TYGS and LPSN: a

database tandem for fast and reliable genome-based classification and nomenclature of prokaryotes.

Nucleic Acids Res *50*, D801-D807.

Mohan, A., Munusamy, C., Tan, Y.C., Muthuvelu, S., Hashim, R., Chien, S.L., Wong, M.K., Khairuddin,

N.A., Podin, Y., Lau, P.S., *et al.* (2019). Invasive Salmonella infections among children in Bintulu,

Sarawak, Malaysian Borneo: a 6-year retrospective review. BMC Infect Dis *19*, 330.

Morgan, K.L. (2013). Infection, illness, and fenugreek seeds: was illness a good enough proxy for

infection in the 2011 outbreak of shiga toxin-producing Escherichia coli O104:H4 in France? Clin Infect

Dis *56*, 1055-1056.

Ohland, C.L., and Macnaughton, W.K. (2010). Probiotic bacteria and intestinal epithelial barrier

function. Am J Physiol Gastrointest Liver Physiol *298*, G807-819.

Ouattara, H.G., Reverchon, S., Niamke, S.L., and Nasser, W. (2017). Regulation of the synthesis of pulp

degrading enzymes in Bacillus isolated from cocoa fermentation. Food Microbiol *63*, 255-262.

Ouyang, W., Rutz, S., Crellin, N.K., Valdez, P.A., and Hymowitz, S.G. (2011). Regulation and functions of

the IL-10 family of cytokines in inflammation and disease. Annu Rev Immunol *29*, 71-109.

Ozer, J., Ratner, M., Shaw, M., Bailey, W., and Schomaker, S. (2008). The current state of serum

biomarkers of hepatotoxicity. Toxicology *245*, 194-205.

Parker, B.J., Wearsch, P.A., Veloo, A.C.M., and Rodriguez-Palacios, A. (2020). The Genus Alistipes: Gut

Bacteria With Emerging Implications to Inflammation, Cancer, and Mental Health. Front Immunol *11*,

906.

Paulson, J.A., Zaoutis, T.E., Council on Environmental, H., and Committee on Infectious, D. (2015).

Nontherapeutic Use of Antimicrobial Agents in Animal Agriculture: Implications for Pediatrics.

Pediatrics *136*, e1670-1677.

Pickard, J.M., Zeng, M.Y., Caruso, R., and Núñez, G. (2017). Gut microbiota: Role in pathogen

colonization, immune responses, and inflammatory disease. Immunol Rev *279*, 70-89.

Polk, B.F., and Kasper, D.L. (1977). Bacteroides fragilis subspecies in clinical isolates. Ann Intern Med *86*,

569-571.

Praveschotinunt, P., Duraj-Thatte, A.M., Gelfat, I., Bahl, F., Chou, D.B., and Joshi, N.S. (2019).

Engineered E. coli Nissle 1917 for the delivery of matrix-tethered therapeutic domains to the gut. Nat

Commun *10*, 5580.

Reyman, M., van Houten, M.A., Watson, R.L., Chu, M., Arp, K., de Waal, W.J., Schiering, I., Plotz, F.B.,

Willems, R.J.L., van Schaik, W., *et al.* (2022). Effects of early-life antibiotics on the developing infant gut

microbiome and resistome: a randomized trial. Nat Commun *13*, 893.

Richter, M., Rossello-Mora, R., Oliver Glockner, F., and Peplies, J. (2016). JSpeciesWS: a web server for

prokaryotic species circumscription based on pairwise genome comparison. Bioinformatics *32*, 929-

931.

Sanders, M.E., Akkermans, L.M., Haller, D., Hammerman, C., Heimbach, J., Hormannsperger, G., Huys,

G., Levy, D.D., Lutgendorff, F., Mack, D., *et al.* (2010). Safety assessment of probiotics for human use.

Gut Microbes *1*, 164-185.

Saraiva, M., Vieira, P., and O'Garra, A. (2020). Biology and therapeutic potential of interleukin-10. J Exp
Med *217*.

Schultz, B.M., Paduro, C.A., Salazar, G.A., Salazar-Echegarai, F.J., Sebastian, V.P., Riedel, C.A., Kalergis,
A.M., Alvarez-Lobos, M., and Bueno, S.M. (2017). A Potential Role of Salmonella Infection in the Onset
of Inflammatory Bowel Diseases. Front Immunol *8*, 191.

Shao, J., Liu, Y., Xie, J., Stefanic, P., Lv, Y., Fan, B., Mandic-Mulec, I., Zhang, R., Shen, Q., and Xu, Z.
(2022). Annulment of Bacterial Antagonism Improves Plant Beneficial Activity of a Bacillus velezensis
Consortium. Appl Environ Microbiol *88*, e0024022.

Singhal, M., Liu, X., Inverso, D., Jiang, K., Dai, J., He, H., Bartels, S., Li, W., Abdul Pari, A.A.,
Gengenbacher, N., *et al.* (2018). Endothelial cell fitness dictates the source of regenerating liver
vasculature. J Exp Med *215*, 2497-2508.

Snydman, D.R. (2008). The safety of probiotics. Clin Infect Dis *46 Suppl 2*, S104-111; discussion S144-
151.

Stecher, B., Macpherson, A.J., Hapfelmeier, S., Kremer, M., Stallmach, T., and Hardt, W.D. (2005).
Comparison of Salmonella enterica serovar Typhimurium colitis in germfree mice and mice pretreated
with streptomycin. Infect Immun *73*, 3228-3241.

Stothard, P., and Wishart, D.S. (2005). Circular genome visualization and exploration using CGView.
Bioinformatics *21*, 537-539.

Tang, Y., Gao, C., Xing, M., Li, Y., Zhu, L., Wang, D., Yang, X., Liu, L., and Yao, P. (2012). Quercetin
prevents ethanol-induced dyslipidemia and mitochondrial oxidative damage. Food Chem Toxicol *50*,
1194-1200.

Tang, Z., Sun, H., Chen, T., Lin, Z., Jiang, H., Zhou, X., Shi, C., Pan, H., Chang, O., Ren, P., *et al.* (2017).
Oral delivery of Bacillus subtilis spores expressing cysteine protease of Clonorchis sinensis to grass
carp (Ctenopharyngodon idellus): Induces immune responses and has no damage on liver and
intestine function. Fish Shellfish Immunol *64*, 287-296.

Wang, L., Gou, X., Ding, Y., Liu, J., Wang, Y., Wang, Y., Zhang, J., Du, L., Peng, W., and Fan, G. (2023). The
interplay between herbal medicines and gut microbiota in metabolic diseases. Front Pharmacol *14*,
1105405.

Wang, Y., Xie, Q., Zhang, Y., Ma, W., Ning, K., Xiang, J.Y., Cui, J., and Xiang, H. (2020). Combination of
probiotics with different functions alleviate DSS-induced colitis by regulating intestinal microbiota, IL-
10, and barrier function. Appl Microbiol Biotechnol *104*, 335-349.

Wu, J., Lin, Z., Wang, X., Zhao, Y., Zhao, J., Liu, H., Johnston, L.J., Lu, L., and Ma, X. (2022).
Limosilactobacillus reuteri SLZX19-12 Protects the Colon from Infection by Enhancing Stability of the
Gut Microbiota and Barrier Integrity and Reducing Inflammation. Microbiol Spectr *10*, e0212421.

Yang, W., Liu, Z., Zhao, Q., Du, H., Yu, J., Wang, H., Liu, X., Liu, H., Jing, X., Yang, H., *et al.* (2022).
Population Genetic Structure and Selection Signature Analysis of Beijing Black Pig. Front Genet *13*,
860669.

Yang, W.H., Westman, J.S., Heithoff, D.M., Sperandio, M., Cho, J.W., Mahan, M.J., and Marth, J.D.
(2021). Neu3 neuraminidase induction triggers intestinal inflammation and colitis in a model of
recurrent human food-poisoning. Proc Natl Acad Sci U S A *118*.

Ye, M., Tang, X., Yang, R., Zhang, H., Li, F., Tao, F., Li, F., and Wang, Z. (2018). Characteristics and
Application of a Novel Species of Bacillus: Bacillus velezensis. ACS Chem Biol *13*, 500-505.

Zhang, D.X., Kang, Y.H., Zhan, S., Zhao, Z.L., Jin, S.N., Chen, C., Zhang, L., Shen, J.Y., Wang, C.F., Wang,
G.Q., *et al.* (2019). Effect of Bacillus velezensis on Aeromonas veronii-Induced Intestinal Mucosal

Barrier Function Damage and Inflammation in Crucian Carp (Carassius auratus). Front Microbiol *10*,
2663.
Zhang, L., Bai, K., Zhang, J., Xu, W., Huang, Q., and Wang, T. (2017). Dietary effects of Bacillus subtilis
fmbj on the antioxidant capacity of broilers at an early age. Poult Sci *96*, 3564-3573.
Zhou, J., Li, M., Chen, Q., Li, X., Chen, L., Dong, Z., Zhu, W., Yang, Y., Liu, Z., and Chen, Q. (2022).
Programmable probiotics modulate inflammation and gut microbiota for inflammatory bowel disease
treatment after effective oral delivery. Nat Commun *13*, 3432.
Zihni, C., Mills, C., Matter, K., and Balda, M.S. (2016). Tight junctions: from simple barriers to
multifunctional molecular gates. Nat Rev Mol Cell Biol *17*, 564-580.

Figure legends

Fig. 1 Genomic characteristics and phylogenetic relationships of *B.*

***velezensis* HBXN2020.** (**A**) The whole-genome map of *B. velezensis*

HBXN2020 with its genomic features. The map consists of 6 circles. From the

inner circle to the outer circle: (1) GC-skew, (2) GC content, (3) reverse

protein-coding genes, different colors represents different COG functional

classifications, (4) genes transcribed in reverse direction, (5) genes

transcribed in forward direction, (6) forward protein-coding genes, different

colors represents different COG functional classifications. (**B**) The whole

genome phylogenetic tree was constructed based on genome-wide data from

14 bacillus strains. *B. velezensis* HBXN2020 are indicated in bold.

Fig. 2 Growth curve of *B. velezensis* HBXN2020 and its *in vitro*

resistance against environmental assaults. (**A**) Growth curves of *B.*

velezensis HBXN2020 cultured in LB medium at 37°C, detection of OD_{600}

values at 1 h intervals in microplate reader. (**B**) Survival rate of endospore

and vegetative cells of *B. velezensis* HBXN2020 after 30 min at different

temperatures (37°C, 45°C, 55°C, 65°C, 75°C, 85°C or 95°C). Equal amounts

of endospore and vegetative cells of HBXN2020 were exposed to the

following: (**C**) acid solution (pH 2 to 7), (**D**) 0.3% bile salts, (**E**) SGF (pH 1.2)

supplemented with pepsin, and (**F**) SIF (pH 6.8) containing trypsin at 37°C. At

predetermined time points, 100 µL was taken from each sample, and tenfold

serially diluted with sterile PBS (pH 7.2), and then spread on LB agar plates

and cultured at 37°C for 12 h before bacterial counting. Each group was

repeated three times (n = 3).

Fig. 3 Antibiotic susceptibility of *B. velezensis* HBXN2020 and

bacteriostasis assay *in vitro*. (**A**) The diameter of the antibacterial zone

indicates the extent of sensitivity to antibiotics. (**B**) The diameter of the

antibacterial zone indicates the extent of inhibition against pathogenic bacteria.

The diameter of the antibacterial zone was measured with vernier caliper.

Each group was repeated three times (n = 3). R, resistant; I, moderately

sensitive; S, sensitive.

1002

Fig. 4 *In vivo* safety evaluation of *B. velezensis* HBXN2020 in a mouse

model. (**A**) Body weights changes of mice during gavage with *B. velezensis*

HBXN2020 spores. Mice were treated with sterile PBS (Control group) or low-

dose (L-HBXN2020 group), medium dose (M-HBXN2020 group), and high-

dose (H-HBXN2020 group) of *B. velezensis* HBXN2020 spores. Weighing and

gavage were performed once every two days during the experimental period

(15 days). Data were shown as mean values ± SEM (n = 5). (**B**) The relative

gene expression levels of inflammatory cytokines in the colon of mice

measured by RT-qPCR. Data were shown as mean values ± SEM (n = 5). (**C**)

The relative gene expression levels of barrier protein ZO-1, occludin, claudin

and Muc2 in the colon of mice measured by RT-qPCR. Data were shown as

mean values ± SEM (n = 5). (**D**) Major blood routine parameters and (**E**)

serum biochemical parameters of mice in the control group and H-HBXN2020

group. Data were shown as mean values ± SEM (n = 3). (**F**) H&E stained

colon sections in the different groups. Scale bar: 200 μm.

1018

Fig. 5 Oral *B. velezensis* HBXN2020 spores alleviated *S.* Typhimurium-

induced experimental mouse colitis. (**A**) *In vitro* Bacterial competition

between STm and *B. velezensis* HBXN2020. STm were co-incubated with *B.*

velezensis HBXN2020 at various ratios at 37°C with shaking. The growth of

STm was reflected by bacterial counting per hour. (**B**) Experimental design for

treatment in this study. Orally administrated with either PBS, or *B. velezensis*

HBXN2020 spores by gavage at days 1, 3, and 5 after STm infection (5×10^7

CFU/mouse), respectively. All mice were euthanized at day 7 after STm

infection. (**C**) Bacterial count of STm in mouse feces. Fecal samples were

collected per day after STm infection and resuspended in sterile PBS (0.1 g of

fecal resuspended in 1 mL of sterile PBS). One hundred microliters of each

sample performed a serial of 10-fold dilutions and spread on selective agar

plates (50 µg mL^{-1} kanamycin) and incubated at 37°C for 12 h before

bacterial counting. The bacterial loads of STm in (**D**) ileum, (**E**) cecum, and (**F**)

colon. The cecum and colon were harvested and then homogenized.

Statistical signifcance was evaluated using Student's t-test (*, $P < 0.05$, **, $P <$

0.01, and ***, $P < 0.001$). (**G**) Daily body weight changes and (**H**) daily disease

activity index (DAI) scores of mice with different treatment groups. Data were

shown as mean values ± SEM (n = 8). Statistical signifcance was evaluated

using one-way analysis of variance (ANOVA) with Tukey's multiple

comparisons test. *, $P < 0.05$, **, $P < 0.01$, and ***, $P < 0.001$ relative to Control

group; $^{#}$, $P < 0.05$, $^{##}$, $P < 0.01$, $^{###}$, $P < 0.001$ relative to STm + HBXN2020

group. (**I**) Colonic tissue images. (**J**) The length of the colon from per group (n

= 8). Statistical signifcance was evaluated using one-way analysis of variance

(ANOVA) with Tukey's multiple comparisons test (*, $P < 0.05$, **, $P < 0.01$, and

***, $P < 0.001$).

Fig. 6 Oral *B. velezensis* HBXN2020 spores attenuated colonic damage

and inflammatory reaction. (**A**) H&E stained colon tissue sections. Scale bar:

200 µm. The relative gene expression levels of TNF-α (**B**), IL-1β (**C**), IL-6 (**D**),

and IL-10 (**E**) were detected by RT-qPCR. Data were shown as mean values

± SEM (n = 6). The relative gene expression levels of ZO-1 (**F**), ocludin (**G**),

claudin (**H**), and Muc2 (**I**) in colon tissue were detected by RT-qPCR. Data

were shown as mean values ± SEM (n = 6). Statistical signifcance was

evaluated using one-way analysis of variance (ANOVA) with Tukey's multiple

comparisons test (*, $P < 0.05$, **, $P < 0.01$, and ***, $P < 0.001$).

Fig. 7 Oral *B. velezensis* HBXN2020 spores regulated the composition of

intestinal microbiota. The α-diversity of the gut microbiota, determined by

the (**A**) Sobs, (**B**) Chao1, (**C**) Shannon, and (**D**) Simpson diversity index. Data

were shown as mean values ± SEM (n = 5). (**E**) The PCA plot showed the β-

diversity of the gut microbiota based on Bray-Curtis distance at the OTU level.

Heatmap of the community composition of colonic microbiota at the phylum

(top 10 phyla) (**F**) and genus (top 30 genera) (**G**) level. Relative abundance of

selected taxa (**H**) *norank_f_Muribaculaceae*, (**I**) *Lactobacillus*, (**J**) *Bacteroides*,

and (**K**) *Escherichia-Shigella*. (**L**) Analysis of differences in the microbial

communities by LEfSe (linear discriminant analysis (LDA score > 3.5) among

different groups. Signifcance was evaluated by the Kruskal-Wallis test or

ANOVA with Tukey's multiple comparisons test (*, P < 0.05, **, P < 0.01, and

***, P < 0.001).

Fig. 8 Prophylactic *B. velezensis* HBXN2020 spores attenuated the

symptoms of *S.* Typhimurium-induced mouse colitis. (**A**) Experimental

design for treatment in this study. At days 1, 3, 5, and 7, each mouse in the

HBXN2020 + STm group and PBS + STm group were received 200 μL (1×10^8

CFU/mouse) of *B. velezensis* HBXN2020 spores or sterile PBS by gavage.

Then, mice in PBS + STm group and HBXN2020 + STm group were orally

inoculated with 200 μL (5×10^7 CFU/mouse) of STm on day 7. On day 12, all

mice were euthanized. (**B**) Daily body weight changes and (**C**) daily disease

activity index (DAI) scores of mice with different groups following STm

treatment. Data were shown as mean values ± SEM (n = 8). Statistical

signifcance was evaluated using one-way analysis of variance (ANOVA) with

Tukey's multiple comparisons test. *, $P < 0.05$, **, $P < 0.01$, and ***, $P < 0.001$

relative to Control group; $^\#$, $P < 0.05$, $^{\#\#}$, $P < 0.01$, $^{\#\#\#}$, $P < 0.001$ relative to

HBXN2020 + STm group. (**D**) Colonic tissue images. (**E**) The length of the

colon from per group (n = 8). (**F**) Bacterial count of STm in mouse feces.

Fecal samples were collected every day after STm infection and resuspended

in sterile PBS (0.1 g of fecal resuspended in 1 mL of sterile PBS). One

hundred microliters of each sample performed a serial of 10-fold dilutions and

spread on selective agar plates (50 μg mL^{-1} kanamycin) and incubated at

37°C for 12 h before bacterial counting. The bacterial loads of STm in (**G**)

cecum and (**H**) colon. The cecum and colon were harvested and then

homogenized. Statistical signifcance was evaluated using Student's t-test (*, P

< 0.05, **, $P < 0.01$, and ***, $P < 0.001$).

Fig. 9 Prophylactic *B. velezensis* HBXN2020 spores attenuated colonic damage and inflammatory reaction. (**A**) H&E stained colon tissue sections. Scale bar: 200 μm. The relative gene expression levels of TNF-α (**B**), IL-1β (**C**), IL-6 (**D**), and IL-10 (**E**) were detected by RT-qPCR. Data were shown as mean values ± SEM (n = 6). The relative gene expression levels of ZO-1 (**F**), ocludin (**G**), claudin (**H**), and Muc2 (**I**) in colon tissue were detected by RT-qPCR. Data were shown as mean values ± SEM (n = 6). Statistical signifcance was evaluated using one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test (*, $P < 0.05$, **, $P < 0.01$, and ***, $P < 0.001$).

Fig. 10 Prophylactic *B. velezensis* HBXN2020 spores regulated the composition of gut microbiota. (**A** to **D**) Alpha diversity of the intestinal microbiota. Data were shown as mean values ± SEM (n = 5). (**E**) The PCA plot showed the β-diversity among different microbial community groups based on Bray-Curtis distance at the OTU level. Heatmap of the community composition of colonic microbiota at the phylum (top 10 phyla) (**F**) and genus (top 30 genera) (**G**) level. Relative abundance of selected taxa (**H**) *Lactobacillus*, (**I**) *Akkermansia*, (**J**) *Alistipes*, (**K**) *Bacteroides*, (**L**) *Escherichia-Shigella*, and (**M**) *Enterococcus*. Data were shown as mean values ± SEM (n = 5). (**N**) Analysis of differences in the microbial taxa by LEfSe (LDA score > 3.5) in different groups. Signifcance was evaluated by the Kruskal-Wallis test or ANOVA with Tukey's multiple comparisons test (*, $P < 0.05$, **, $P < 0.01$,

1116   and ***, P < 0.001).

A

A RNA processing and modification	
B Chromatin structure and dynamics	
C Energy production and conversion	
D Cell cycle control, cell division, chromosome partitioning	
E Amino acid transport and metabolism	
F Nucleotide transport and metabolism	
G Carbohydrate transport and metabolism	
H Coenzyme transport and metabolism	
I Lipid transport and metabolism	
J Translation, ribosomal structure and biogenesis	
K Transcription	
L Replication, recombination and repair	
M Cell wall/membrane/envelope biogenesis	
N Cell motility	
O Posttranslational modification, protein turnover, chaperones	
P Inorganic ion transport and metabolism	
Q Secondary metabolites biosynthesis, transport and catabolism	
R General function prediction only	
S Function unknown	
T Signal transduction mechanisms	
U Intracellular trafficking, secretion, and vesicular transport	
V Defense mechanisms	
W Extracellular structures	
Y Nuclear structure	
Z Cytoskeleton	
CDS	
tRNA	
rRNA	
Other	
GC content	
GC skew+	
GC skew-	

Bacillus velezensis
HBXN2020
3,929,792 bp

500 kbp
1000 kbp
1500 kbp
2000 kbp
2500 kbp
3000 kbp
3500 kbp

B

B. tequilensis NCTC13306
B. vallismortis DV1-F-3
B. rugosus SPB7
B. spizizenii TU-B-10
B. stercoris D7XPN1
B. subtilis NCIB 3610
B. subtilis ATCC 6051
B. nakamurai NRRL B-41091
B. amyloliquefaciens DSM 7
B. vanillea XY18
B. siamensis KCTC 13613
B. velezensis FZB42
B. velezensis NRRL B-41580
B. velezensis KACC 13105
**B. velezensis* HBXN2020*

Tree scale: 0.02