Bacillus velezensis HBXN2020 alleviates Salmonella Typhimurium-induced colitis by improving intestinal barrier integrity and reducing inflammation

Linkang Wang,[1,2,3] Haiyan Wang,[1,2,3] Xinxin Li,[1,2,3] Mengyuan Zhu,[1,2,3] Dongyang Gao,[1,2,3] Dayue Hu,[1,2,3] Zhixuan Xiong,[1,2,3] Xiangmin Li[1,2,3]* and Ping Qian[1,2,3]*

[1]*National Key Laboratory of Agricultural Microbiology, Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan, Hubei, 430070, People's Republic of China.*

[2]*College of Veterinary Medicine, Huazhong Agricultural University, Wuhan, Hubei, People's Republic of China.*

[3]*Key Laboratory of Preventive Veterinary Medicine in Hubei Province, the Cooperative Innovation Center for Sustainable Pig Production, Wuhan, Hubei, People's Republic of China.*

* Corresponding author: Ping Qian: Tel: 86-27-87282608; Fax: 86-27-87282608; E-mail: qianp@mail.hzau.edu.cn. Address: No. 1, Shizishan Street, Hongshan District, Wuhan City, Hubei Province, China.

* Co-corresponding author: Xiang-min Li: Tel: 86-27-87282608; Fax: 86-27-87282608; E-mail: lixiangmin@mail.hzau.edu.cn. Address: No. 1, Shizishan

Street, Hongshan District, Wuhan City, Hubei Province, China.

Abstract

Bacillus velezensis is a novel species of Bacillus that has been widely

investigated because of its broad-spectrum antimicrobial activity. However,

most studies on *Bacillus velezensis* have focused on biocontrol of plant

diseases, with few reports on antagonizing *Salmonella* Typhimurium infections.

In this study, *Bacillus velezensis* HBXN2020 was isolated from healthy black

piglets and was found to exhibit broad-spectrum antibacterial activity and

robust anti-stress capabilities. Importantly, *Bacillus velezensis* HBXN2020 did

not cause any adverse side effects in mice when administered at various

doses (1×10^7, 1×10^8, and 1×10^9 CFU) for 14 days. In a *Salmonella*

Typhimurium ATCC14028-induced mice colitis model, either curative or

prophylactic, supplementing *Bacillus velezensis* HBXN2020 spores

significantly lowered the levels of *Salmonella* Typhimurium ATCC14028 in

their feces, ileum, cecum, and colon, and the disease activity index (DAI).

Importantly, supplementing *Bacillus velezensis* HBXN2020 spores

significantly regulated cytokine levels (TNF-α, IL-1β, IL-6, and IL-10) and

maintained the expression of tight junction proteins and mucin protein. More

importantly, supplementing *Bacillus velezensis* HBXN2020 spores also

significantly enhanced the homeostasis of colonic microbiota and the

abundance of beneficial bacteria. Collectively, *Bacillus velezensis* HBXN2020

can alleviate bacterial colitis by enhancing intestinal homeostasis and gut

barrier integrity and reducing inflammation.

Introduction

Salmonella Typhimurium (*S.* Typhimurium) is a major foodborne zoonotic

pathogen that can cause diarrhea and colitis in humans and animals (1, 2). It

has been reported that *Salmonella* infections often occur in both developing

and developed countries, which poses a substantial risk to public health and

causes huge economic loss (3, 4). At present, antibiotic therapies is one of

the most effective strategies for *Salmonella* infections. However, numerous

studies have reported that the prolonged use or misuse of antibiotics can lead

to environmental pollution, an increase in multi-drug-resistant (MDR) bacteria,

as well as gastrointestinal microbiota dysbiosis (5-7). Therefore, there is an

emergent search for novel antimicrobial agents to combat *Salmonella*

infections.

Recently, an increasing number of studies have shown that gut microbiota

plays an important role in alleviating intestinal inflammation (8) and repairing

the intestinal barrier (9, 10). The use of probiotics is a popular approach to

modulate intestinal microbiota nowadays. Among all probiotics, Bacillus is one

of the most popular probiotic species because of their ability to form

endospores that survive gastric transit, storage and delivery conditions (11).

In addition, Bacillus species can also produce antimicrobial metabolites

against enteric pathogens and provide extracellular enzymes, thereby

reducing infection risk and improving nutrient utilization (12, 13).

Bacillus velezensis (*B. velezensis*)*,* a newly discovered species of the

Bacillus genus, can produce an abundance of secondary metabolites with

broad-spectrum antibacterial activity (14). Previous studies have reported that

B. velezensis exhibits varying degrees of probiotic effects on plants, livestock,

poultry and fish. For instance, the combined use of *B. velezensis* SQR9

mutant and FZB42 improved root colonization and the production of

secondary metabolites that are beneficial to cucumber (15). *B. velezensis*

isolated from yaks was shown to enhance growth performance and ameliorate

blood parameters related to inflammation and immunity in mice (16). The

dietary *B. velezensis* supplementation can regulate the innate immune

response in the intestine of crucian carp and reduce the degree of intestinal

inflammation damage induce by *A. veronii* (17). However, most studies on *B.*

velezensis were mainly biocontrol of plant fungal diseases, with few reports

on the treatment or prevention of *S.* Typhimurium-induced colitis.

In this study, we have screened a strain of *B. velezensis* HBXN2020 with

broad-spectrum antibacterial activity from a large number of *Bacillus* strains.

We aimed to investigate the effectiveness of *B. velezensis* HBXN2020 in

alleviating *S.* Typhimurium-induced mouse colitis and assessed the biological

safety of *B. velezensis* HBXN2020. Notably, *B. velezensis* HBXN2020 can

alleviate symptoms and colon tissue damage in experimental colitis, as

indicated by markers such as *S.* Typhimurium loads, TNF-α and ZO-1 levels.

Moreover, *B. velezensis* HBXN2020 was also enhanced the homeostasis of

colonic microbiota and the abundance of beneficial bacteria (*Lactobacillus*

and *Akkermansia*). Therefore, this study provides support for the development

of probiotic-based microbial products as an alternative prevention strategy for

Salmonella infection.

Results

Genomic Characteristics

The Illumina Hiseq and PacBio platforms were used to perform whole

genome sequencing of HBXN2020, yielding a total of 8,194,986 Illumina clean

reads (1,237,442,886 bases) and 225,593 PacBio reads (1,807,288,759

bases). Following a quality assessment via k-mer and a GC-depth analysis,

the whole genome was assembled *de novo* using Unicycler (Version 0.4.8).

The results showed that HBXN2020 has a circular chromosome of 3,929,792

bp (Fig. 1A) with a GC content of 46.5%. The genome of HBXN2020 was

predicted to contain 3,857 genes, including 3,744 protein-coding genes (CDS),

86 tRNA genes, and 27 rRNA genes (Table S1).

A phylogenetic tree based on genome-wide data from all 14 Bacillus strains

demonstrated that the HBXN2020 belongs to the *B. velezensis* species (Fig.

1B). To further understand the classification status of HBXN2020, the online tool JSpeciesWS was used to calculate the average nucleotide identity (ANI) based on the BLAST (ANIb) method. As shown in Fig. 1C, HBXN2020 was found to be a member of the *B. velezensis* species due to the high percentage of ANIb (more than 97%).

Growth curve of *B. velezensis* HBXN2020 and its *in vitro* resistance against environmental assaults.

The growth of *B. velezensis* HBXN2020 was assessed in flat-bottomed 100-well microtiter plates by measuring the values of OD_{600} every hour using an automatic growth curve analyzer. The OD_{600} value of *B. velezensis* HBXN2020 showed slight fluctuations within the first 2 h, followed by a logarithmic growth phase that reached a stable phase after 10 h (Fig. 2A). Despite environmental restrictions such as limited space and nutrients, the OD_{600} value of the bacteria was not significantly reduced even after 22 h, indicating that the growth of *B. velezensis* HBXN2020 was relatively stable.

Next, we evaluated the effect of physical, chemical, and biological sterilization conditions such as high temperature, strong acidity, and enzyme preparation on the survival of *B. velezensis* HBXN2020. The survival rate of *B. velezensis* HBXN2020 showed a decreasing trend with increasing temperature. At 65°C, the survival rate of *B. velezensis* HBXN2020 was less than 1%, while the spores of *B. velezensis* HBXN2020 exceeded 90% (Fig.

2B). The survival rate of *B. velezensis* HBXN2020 spores was also above 80% even under high-temperature conditions (95°C). Similar results were observed following exposure to a strong acid environment (pH 2.0) (Fig. 2C), whereby *B. velezensis* HBXN2020 spores were found to effectively tolerate strong acidic conditions and exhibited improved survival rate. Based on these results, we hypothesized that *B. velezensis* HBXN2020 spores might also exhibit improved survival in gastrointestinal tract environments. Therefore, we further evaluated the survival rate of *B. velezensis* HBXN2020 in bile salt (0.3%, v/v), simulated gastric fluid (pepsin 1 mg mL^{-1}, pH 1.2), and simulated intestinal fluid (trypsin 1 mg mL^{-1}, pH 6.8). As shown in Fig. 2D-F, *B. velezensis* HBXN2020 spores demonstrated significantly improved tolerance to these simulated gastrointestinal tract environments.

Antibiotic susceptibility of *B. velezensis* HBXN2020 and bacteriostasis effect *in vitro*.

To evaluate the drug resistance of *B. velezensis* HBXN2020, 19 commonly used antibiotics in clinical practice were selected. The results indicated that only polymyxin B had a relatively small inhibition zone diameter (less than 15 mm). Ampicillin, meropenem, minocycline, ofloxacin, and trimethoprim had the strongest inhibition on *B. velezensis* HBXN2020, with an inhibition zone diameter exceeding 30 mm (Fig. 3A). In addition, the experimental results showed that *B. velezensis* HBXN2020 was extremely sensitive to β-lactams,

tetracyclines, and quinolone drugs. Then, we performed an inhibition test of *B.*

velezensis HBXN2020 against seven common clinical pathogenic bacteria.

The results showed that *B. velezensis* HBXN2020 had a similar inhibitory

effect on standard and wild strains of *E. coli*, *Salmonella*, *S. aureus*, and *C.*

perfringens, as well as wild strains of *S. suis*, *P. multocida*, and *A.*

pleuropneumoniae. Except for the wild strains of *S. suis* and *A.*

pleuropneumoniae, the diameter of the inhibition zone of other strains was

above 15 mm (Fig. 3B, S3), while the size of the inhibition zone of *S. suis* and

A. pleuropneumoniae was also more than 12 mm.

Biosafety evaluation of *B. velezensis* HBXN2020

To determine the safety of *B. velezensis* HBXN2020, we evaluated its

biological safety in a mouse model. After gavage with *B. velezensis*

HBXN2020 spores for two weeks, we observed no significant difference in the

body weight of each group of mice (Fig. 4A). The gene expression levels of

TNF-α, IL-1β, IL-6, and IL-10 of colon from all groups of mice exhibited no

remarkable changes (Fig. 4B). However, in the colon, mRNA levels of the

barrier proteins ZO-1 and occludin were trending towards an increase

compared with the control group (Fig. 4C). Additionally, blood routine tests

and serum biochemistry tests were performed for mice in the control group

and H-HBXN2020 group on the 14th day after oral administration of *B.*

velezensis HBXN2020 spores multiple times. As shown in Fig. 4D and S1, the

blood parameters of mice after *B. velezensis* HBXN2020 treatment, including red blood cells (RBC), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), hemoglobin (HGB), white blood cells (WBC), platelets (PLT), hematocrit (HCT), and mean corpuscular hemoglobin concentration (MCHC), were consistent with those of healthy mice. The serum biochemical parameters, including alanine aminotransferase (ALT), aspartate aminotransferase (AST), albumin (ALB), total bilirubin (TBIL), serum creatinine (CREA), and blood urea nitrogen (BUN), were also within normal limits (Fig. 4E). The corresponding histological analysis of colon tissue from mice receiving low, medium, and high doses of *B. velezensis* HBXN2020 spores (L-HBXN2020, M-HBXN2020, and H-HBXN2020 groups, respectively) is presented in Fig. 4F. The colon tissue sections of mice in the test groups showed no significant differences compared to the control group. Additionally, there were no observable differences in the major organ tissues (heart, liver, spleen, lung, and kidney) of mice treated with the high dose of *B. velezensis* HBXN2020 spores compared to healthy mice (Fig. S2). Collectively, these experimental results indicate that *B. velezensis* HBXN2020 is safe for use and does not cause any adverse side effects in mice.

Oral administration of *B. velezensis* HBXN2020 spores alleviated *S. Typhimurium*-induced colitis.

To investigate whether *B. velezensis* HBXN2020 could inhibit *S.*

Typhimurium ATCC14028 (STm), solid and liquid co-culture tests were conducted, and the results showed that *B. velezensis* HBXN2020 inhibited the growth of STm in a dose-dependent manner (Fig. 5A, S4A-B, S4D). Next, the therapeutic potential of *B. velezensis* HBXN2020 was evaluated in an STm-induced mouse colitis model. At days 1, 3, and 5 after STm infection, mice in the STm + HBXN2020 group were orally administered *B. velezensis* HBXN2020 spores by gavage (Fig. 5B), while the control group and STm + PBS group were treated with sterile PBS. Also, the number of excreted STm in feces was counted daily at the designated time points. As shown in Fig. 5C, there was a significant and continuous reduction in the number of STm in feces following treatment with *B. velezensis* HBXN2020 spores, while the number of *B. velezensis* HBXN2020 viable bacteria in feces is also gradually decreasing (Fig. S5A). The number of STm in mouse feces was reduced by 10 to 50 times compared to the STm + PBS group from day 2 to 7 post-treatment. *B. velezensis* HBXN2020 spore treatment also resulted in a 5 to 10-fold reduction in the number of STm in the ileum, cecum, and colon, respectively (Fig. 5D-E, S5B). The increased inhibition of STm after *B. velezensis* HBXN2020 spore treatment also resulted in less weight loss and a marked reduction ($P < 0.001$) in DAI scores in STm + HBXN2020 group mice (Fig. 5F-G). In addition, we also measured the colon length of the mice and found that the STm + PBS group had a significantly shorter colon length ($P < 0.001$) than the control group. However, compared to the STm + PBS group,

the STm + HBXN2020 group exhibited a longer colon ($P < 0.05$) (Fig. 5H-I).

To further assess the impact of *B. velezensis* HBXN2020 on the intestinal

tract, we histologically analyzed their colonic tissues. The results showed that

mice treated with *B. velezensis* HBXN2020 spores exhibited a lower histology

score (total scoring) ($P < 0.05$) (Fig. S5C) than mice not treated with the

spores and were similar to that of healthy mice (control group). The colon

tissue of mice not treated with *B. velezensis* HBXN2020 spores displayed

higher histological scores, with significant crypt deformation, severe mucosal

damage, and obvious inflammatory cell infiltration. These findings suggest

that *B. velezensis* HBXN2020 can alleviate colon tissue damage. Moreover,

we measured the mRNA levels of inflammatory cytokines, including tumor

necrosis factor-α (TNF- α), interleukin-1 β (IL-1 β), interleukin-6 (IL-6), and

interleukin-10 (IL-10) in colon tissue using RT-qPCR analysis. As shown in

Fig. 6B-D, the mRNA levels of TNF-α ($P < 0.01$), IL-1β ($P < 0.05$), and IL-6 (P

< 0.05) in the colon tissue of STm-infected mice were significantly upregulated,

indicating the successful establishment of the STm-induced colitis model. As

expected, treatment with *B. velezensis* HBXN2020 spores led to a significant

reduction in the mRNA levels of these pro-inflammatory cytokines, while the

mRNA levels of the anti-inflammatory cytokine IL-10 were significantly

increased ($P < 0.01$) (Fig. 6E), suggesting that *B. velezensis* HBXN2020

spores can alleviate the inflammatory reaction of the colon. Moreover, the

mRNA expression levels of colonic epithelial barrier proteins, including ZO-1,

ocludin, claudin, and Muc2, were also measured by RT-qPCR. Compared

with the PBS-treated group, the *B. velezensis* HBXN2020 spores treatment

group exhibited significantly upregulated mRNA expression levels of ZO-1 (P

< 0.05), occludin (P < 0.05), claudin (P < 0.05), and Muc2 (P < 0.01) in the

colon tissue (Fig. 6F-I), further demonstrating the ability of *B. velezensis*

HBXN2020 to reduce damage to the colon epithelial barrier.

Next, we further explored the effect of HBXN2020 spores on the intestinal

microbiota composition of STm-treated mice via 16S rRNA gene sequencing.

The alpha diversity of colonic microbiota was evaluated using Sobs, Chao1,

Shannon, and Simpson indices (OTU levels). Compared with the control and

HBXN2020 group, the Sobs, Chao1, and Shannon indices of the STm + PBS

group clearly decreased (Fig. 7A-C), while the Simpson index significantly

increased (P < 0.05, Fig. 7D). Principle component analysis (PCA) based on

Bray‑Curtis distance showed that compared with the STm + PBS group, the

gut microbiota composition of the STm + HBXN2020 group and the control

group were more closer together (R=0.3988, P=0.001, Fig. 7E). We analyzed

the significance of differences among groups at different classification levels

(Fig. 7F-G), and the results revealed that the composition of the gut

microbiota changed markedly after STm-induced mice. At the phylum level,

the STm + PBS group increased the abundance of *Verrucomicrobiota* and

Proteobacteria and decreased the abundance of *Firmicutes*, *Patescibacteria*,

Desulfobacterota, and *Actinobacteriota* compared to the Control group (Fig.

7F). The HBXN2020 group increased the abundance of *Bacteroidota* and

Firmicutes and decreased the abundance of *Verrucomicrobiota* compared

with the Control group. On the contrary, the relative abundance of *Firmicutes*,

Patescibacteria, *Desulfobacterota*, and *Actinobacteriota* in the STm +

HBXN2020 group was significantly higher than that in the STm + PBS group.

At the genus level, infection with *S.* Typhimurium ATCC14028 in the STm +

PBS   group   dramatically   reduced   the   relative   abundance   of

norank_f_Muribaculaceae and *Lactobacillus* (Fig. 7H-I) and enhanced the

abundance of *Bacteroides*, *Escherichia-Shigella*, *Salmonella*, *Alistipes*, and

Enterococcus compared to the Control group (Fig. 7J-K, S6B). The

HBXN2020 group increased the abundance of *norank_f_Muribaculaceae* and

Alloprevotella compared to the Control group. In contrast, STm + HBXN2020

group   significantly   improved   the   relative   abundance   of

norank_f_Muribaculaceae and *Lactobacillus* compared with the STm + PBS

group (*P* < 0.01, *P* < 0.001, Fig. 7H-I). In addition, LEfSe analysis was also

performed to identify specialized microbial communities between the STm +

HBXN2020 group and the STm + PBS group, and found that there was a

clear difference in gut microbiota composition between them. The microbiota

that played a major role in the STm + PBS group were harmful

Enterobacterales (Fig. 7L), while the STm + HBXN2020 group was

Candidatus_Saccharimonas (Fig. 7L).

Prophylactic *B. velezensis* HBXN2020 spores alleviated *S.* Typhimurium-induced colitis.

Based on the improved therapeutic efficacy of *B. velezensis* HBXN2020 in the treatment of STm-induced colitis in mice, we further explored its potential for disease prevention by evaluating pre-treatment. Mice in the PBS + STm group and HBXN2020 + STm group were given the same amount of sterile PBS and *B. velezensis* HBXN2020 spores by gavage one week in advance. As shown in Fig. 8, compared with the PBS + STm group, oral administration of *B. velezensis* HBXN2020 spores as a preventive measure remarkably alleviated STm-induced colitis, including weight loss of mice ($P < 0.01$) (Fig. 8B), a significant reduction in DAI ($P < 0.001$) (the comprehensive score of weight loss, stool consistency, and blood in the feces, Fig. 8C), and the prevention of colon length shortening (Fig. 8D-E). However, the number of *B. velezensis* HBXN2020 viable bacteria in feces is also gradually decreasing with the time prolonging (Fig. S7A). Furthermore, compared with the PBS + STm group, oral administration of *B. velezensis* HBXN2020 spores not only reduced the number of STm in mouse feces (Fig. 8F) but also decreased STm colonization in the ileum, cecum, and colon (Fig. 8G-H, S7B).

Histological analysis further revealed that oral administration of *B. velezensis* HBXN2020 significantly reduced the infiltration of inflammatory cells, mucosal damage and the overall histological score of colon tissue (Fig. 9A, S7C). Additionally, the mRNA expression levels of pro-inflammatory

cytokines, including TNF-α ($P < 0.05$), IL-1β ($P < 0.05$) and IL-6 ($P < 0.01$), in

the colon tissue of mice pretreated with *B. velezensis* HBXN2020 spores were

significantly decreased (Fig. 9B-D), while the mRNA levels of anti-

inflammatory cytokine (IL-10) were significantly higher ($P < 0.05$) (Fig. 9E)

than mice pretreated with PBS. In addition, compared to the group pretreated

with PBS, the group pretreated with *B. velezensis* HBXN2020 spores showed

a significant increase in mRNA expression levels of ZO-1 ($P < 0.001$), ocludin

($P < 0.05$), and claudin ($P < 0.05$) in the colon tissue of mice (Fig. 9F-H).

While there was no significant difference in Muc2 expression ($p=0.252$) (Fig.

9I), the results were consistent with those of the therapeutic experiments.

Overall, these findings suggest that *B. velezensis* HBXN2020 can effectively

alleviate the symptoms of STm-induced colitis and the associated damage to

colon tissue.

Next, we examined the impact of prophylactic *B. velezensis* HBXN2020 on

the intestinal microbiota composition of STm-treated mice. Sobs, Chao1 index

and Shannon index have changed to some extent between groups, but

differences were not significant (Fig. 10A-C). Compared with the Control

group, the Simpson index significantly decreased in the PBS+STm group (P <

0.05, Fig. 10D). However, compared to the PBS+STm group, Simpson index

in the HBXN2020+STm group did not change (Fig. 10D). PCA analysis

showed a significant separation between control and PBS+STm groups, while

the HBXN2020+STm group, HBXN2020 group, and the control group were

clustered together (Fig. 10E). Next, we analyzed microbial community

composition at the phylum and genus levels (Fig. 10F-G). At the phylum level,

the PBS + STm group increased the abundance of *Bacteroidota*,

Proteobacteria, *Actinobacteriota*, and *Deferribacterota* and reduced the

abundance of *Verrucomicrobiota*, *Firmicutes*, *Patescibacteria*, and

Desulfobacterota compared with the Control group(Fig. 10F, S8A). The

HBXN2020 group increased the abundance of *Bacteroidota* compared to the

control group. In contrast, compared to the PBS+STm group, the abundance

of *Firmicutes* in the HBXN2020+STm group was significantly increased (*P* <

0.05). At the genus level, a depletion of *Lactobacillus* and *Akkermansia* (Fig.

10H-I) and an enrichment of *Alistipes*, *Bacteroides*, *Escherichia-Shigella*,

Alloprevotella, *Helicobacter*, *Enterococcus*, and *Salmonella* were observed in

PBS + STm group compared with the control (Fig. 10J-M, 8SB). The

enrichment with *norank_f_Muribaculaceae* and *Alloprevotella* were observed

in HBXN2020 group compared to the control. In contrast, an enrichment of

Lactobacillus, *Akkermansia*, and *Lachnospiraceae_NK4A136_group* and a

sharp decrease of *Alistipes*, *Bacteroides*, *Escherichia-Shigella*, and

Enterococcus were observed in HBXN2020 + STm group compared to the

PBS + STm group. LEfSe analysis showed that the abundance of

Lactobacillus in the intestinal microbiota of mice was significantly increased

after *B. velezensis* HBXN2020 treatment, while the abundance of harmful

Enterobacterales in the untreated group was clearly increased (Fig. 10N).

Discussion

S. Typhimurium is an important intestinal pathogen that can cause invasive

intestinal diseases such as bacterial colitis (18, 19). Although antibiotics are

commonly used to treat bacterial-induced colitis, their overuse has

significantly increased bacterial resistance in recent years, resulting in serious

environmental pollution. A recent study showed that antibiotic-resistance

genes in probiotics could be transmitted to the intestinal microbiota, which

may threaten human health (20). Therefore, the source of probiotics is

particularly important. Black pigs are an excellent domestic breed in China,

with strong environmental adaptability and disease resistance (21). In this

study, *B. velezensis* HBXN2020 was isolated from the free-range feces of

black piglets in a mountain village in Xianning City (Hubei, China) and

exhibited excellent antibacterial activity. Previous studies have demonstrated

that the difficidins isolated from *B. subtilis* can inhibit bacterial protein

synthesis, giving it broad-spectrum antibacterial activity against *Gram positive*

and *Gram negative* bacteria. Therefore, we speculate that *B. velezensis*

HBXN2020 has the ability to synthesize Difficidin or its derivatives (22).

Otherwise, *in vitro* tolerance assays showed that *B. velezensis* HBXN2020

spores have good tolerance to high temperature, strong acids, bile salts, and

simulated gastric and intestinal fluid, which was similar to previous research

results (23). Furthermore, antibiotic susceptibility tests showed that *B.*

velezensis HBXN2020 does not exhibit antibiotic resistance.

Probiotics are unique in that they are alive when consumed, and unlike food

or drug, they have the potential to produce toxins or infections in the body (24,

25). Therefore, assessing the biosafety of probiotics is crucial. In this study,

the safety of *B. velezensis* HBXN2020 was evaluated by measuring the body

weight, intestinal barrier proteins, and inflammatory cytokines of the

experimented mice, as well as conducting routine blood and biochemical tests.

The results showed that the expression levels of ZO-1 and occludin in the

colon of mice increased after treatment with different doses of *B. velezensis*

HBXN2020 spores. It has been reported that ZO-1, occludin and claudin are

the three most important tight junction proteins in intercellular connections,

which play a crucial role in maintaining the intestinal epithelial barrier (26, 27).

Therefore, upregulating the expression levels of ZO-1 and Occludin might

enhance intestinal barrier function.

Alanine aminotransferase (ALT) and aspartate aminotransferase (AST) are

important aminotransferases in animals and are considered key parameters

for evaluating liver function injury (28, 29). Under normal circumstances, the

levels of ALT and AST are in dynamic balance without notable changes.

However, when the liver is damaged or becomes dysfunctional, the levels of

aminotransferases (ALT and AST) increase significantly (30). Previous

studies have shown that CCL4-induced liver injury strongly increases plasma

ALT/AST levels (31). Zhang et al. reported that the addition of *Bacillus subtilis*

to chicken diets significantly decreased the serum levels of ALT and AST (32).

In this present study, the levels of ALT and AST in the probiotic-treated group

were slightly lower than those in the control group, indicating that *B.*

velezensis HBXN2020 had no negative effect on liver health in mice.

Furthermore, routine blood tests found that the blood indices of the *B.*

velezensis HBXN2020 treatment group were within the normal reference

range, similar to that of the control group.

To further evaluate the potential of *B. velezensis* HBXN2020 spores in

alleviating *S.* Typhimurium-induced colitis, mice were challenged with *S.*

Typhimurium ATCC14028 and then treated with *B. velezensis* HBXN2020

spores via oral administration. Our experimental results demonstrated that

oral treatment with *B. velezensis* HBXN2020 spores could alleviate *S.*

Typhimurium-induced colitis, as evidenced by decreased weight loss, DAI,

and histological damage, which is similar with prior studies (33, 34). Previous

research showed that macrophages are the first line of host defense against

bacterial infection. They release pro-inflammatory cytokines, which are critical

in initiating adaptive immune responses (35, 36). Nevertheless, pro-

inflammatory cytokines have immunological properties that can be beneficial

for the host to resist the invasion of bacteria and other microbes in the

surrounding environment (37). For instance, pre-treatment with recombinant

murine TNF-α was shown to protect mice against lethal bacterial (*E. coli*)

infection (38). PJ-34 exerted protective effects on intestinal epithelial cells

against invasive *Salmonella* infection by upregulating IL-6 expression through

the ERK and NF-κB signal pathways (39). However, some studies have

shown that excessive immune cell activation can lead to tissue damage, multi-

organ malfunction, systemic or chronic inflammation, and autoimmune

disorders (40, 41). IL-10 is a recognized anti-inflammatory mediator that plays

a crucial role in maintaining intestinal microbe-immune homeostasis,

regulating the release of inflammatory mediators, and inhibiting pro-

inflammatory responses of innate and adaptive immunity (42-44). For instance,

previous studies have shown that *Clostridium butyrate* can induce the

production of IL-10 in the intestine, thereby alleviating experimental colitis in

mice (45). Similar results were observed in our study, where oral

administration of *B. velezensis* HBXN2020 spores reduced the expression

levels of TNF-α, IL-1β and IL-6 while increasing the levels of IL-10 in the colon

of mice. Furthermore, oral administration of *B. velezensis* HBXN2020 spores

improved the functional damage and integrity of the intestinal barrier caused

by *S.* Typhimurium infection, as evidenced by the upregulated expression

levels of intestinal barrier proteins and reduced histological damage.

Subsequently, we further analyzed the colon microbiome and it was

noteworthy that oral *B. velezensis* HBXN2020 markedly increased alpha

diversity of the colonic microbiota. As reported in prior studies, elevated alpha

diversity associates with enhanced anti-interference ability (46). PCA analysis

results showed that oral administration of *B. velezensis* HBXN2020

significantly affected the structure of the intestinal microbiota. Further analysis at the phylum/genus level revealed that *B. velezensis* HBXN2020 treatment reduced the increase of *Bacteroides*, *Escherichia-Shigella*, and *Salmonella* (harmful to intestinal homeostasis) in colitis (47), but enhanced the relative abundance of *Lactobacillus* (known for enhancing the epithelial barrier by increasing mucus secretion and upregulating the expression of tight junction proteins such as claudin-1, occludin, and ZO-1) (48) and *Akkermansia* (known for modulating intestinal immune response by producing short chain fatty acids and reducing the secretion of pro-inflammatory cytokines) (49). *Bacteroides* is generally 'friendly' commensals in the intestine, but they can enter normal tissues through the intestinal mucosa when intestinal barrier function becomes compromised (50), leading to inflammation, diarrhea, and abdominal abscess (51). *Escherichia-Shigella* is a common intestinal pathogen, and its relative abundance is associated with pro-inflammatory states (52). *Salmonella* can secrete bacterial virulence proteins, which are directly delivered to the host cell cytosol through the Type III secretion system and induce the expression of pro-inflammatory cytokines (53), leading to inflammation. Overall, the structure of the colon microbiota was disrupted by *S.* Typhimurium ATCC14028 infection and increased the relative abundance of harmful bacteria, while therapeutic *B. velezensis* HBXN2020 enhanced the abundance of beneficial bacteria, reduced inflammation, and improved intestinal barrier function.

Due to the improved efficacy of treating *S.* Typhimurium-induced colitis, we

further investigated the potential of *B. velezensis* HBXN2020 spores for

disease prevention by evaluating pre-treatment. Similarly to the therapeutic

effect of oral *B. velezensis* HBXN2020, pre-treatment with *B. velezensis*

HBXN2020 spores effectively alleviated the symptoms of *S.* Typhimurium-

induced colitis and reduced the expression levels of pro-inflammatory

cytokines and increased anti-inflammatory cytokine levels in the colon of *S.*

Typhimurium-infected mice. Furthermore, pre-treatment with *B. velezensis*

HBXN2020 spores also promoted the expression levels of intestinal barrier

proteins in the colon of *S.* Typhimurium-infected mice. In addition, the

microbiota composition was also dramatically modulated by prophylactic *B.*

velezensis HBXN2020 in mice with colitis, although the community richness

and diversity revealed by alpha diversity index was not significantly impacted

in mice with colitis by prophylactic-treatment. Further analysis at the genus

level revealed that prophylactic *B. velezensis* HBXN2020 clearly increased

SCFAs-producing bacteria including *Lactobacillus* and *Akkermansia*, which

was consistent with the results of therapeutic *B. velezensis* HBXN2020.

Similarly, harmful bacteria such as *Alistipes*, *Bacteroides*, *Escherichia-*

Shigella, and *Enterococcus* were significantly enriched in the PBS+STm

group. *Alistipes* is a facultative‐pathogenic bacteria and promotes the

development of inflammation, obesity, and colorectal cancer (54). Based on

those results, we found that prophylactic *B. velezensis* HBXN2020 alleviates

S. Typhimurium induced colitis by increasing the abundance of beneficial bacteria, attenuating pro-inflammatory response of the intestine, and increasing intestinal barrier function.

Conclusion

In conclusion, infection with *S.* Typhimurium ATCC14028 disrupts the intestinal mucosal barrier, and leading to intestinal inflammation. Both therapeutic and prophylactic, supplementing *B. velezensis* HBXN2020 spores alleviates *S.* Typhimurium-induced colonic injury by reducing Salmonella colonization, and enhancing the abundance of beneficial bacteria (especially *Lactobacillus* and *Akkermansia*) and intestinal barrier integrity, and decreasing inflammation.

Experimental procedures

Animals and bacterial strains.

Female specific-pathogen-free (SPF) C57BL/6 mice aged 6 to 8 weeks were purchased from the Animal Experimental Center of Huazhong Agricultural University. All mice experiments were conducted in the standard SPF facility of Huazhong Agricultural University, with 12 h of light and 12 h of darkness at a temperature of 25°C and ad libitum access to food and water. The use of animals in this experiment was approved by the Animal Care and

Ethics Committee of Huazhong Agricultural University (Ethics Approval
Number: HZAUMO-2023-0089).

The strains used in this study are listed in Table S2, and all the primers
used in Table S3. *Clostridium perfringens* were cultured in a fluid thioglycollate
medium (FTG) (Hopebio, Qingdao, China) at 45°C. *Streptococcus suis*,
Pasteurella multocida, and *Actinobacillus pleuropneumoniae* were cultured in
tryptic soy broth (TSB) (BD Biosciences, MD, United States) medium
supplemented with 5% (v/v) sheep serum (Solarbio, Beijing, China) at 37°C,
while the other bacterial strains were cultured in Luria Bertani (LB) medium
(Solarbio, Beijing, China) at 37°C. In addition, the Difco sporulation medium
(DSM) was used for inducing the sporulation of *B. velezensis* HBXN2020 via
the nutrient depletion method (55).

Stain isolation, growth conditions and strain identification.

In this study, we isolated 362 *Bacillus* strains from the feces of healthy
black piglets that were free-ranging in a mountain village in Xianning City,
Hubei Province, China, and 4 *Bacillus* strains were screened through initial
antibacterial tests. Then, a strain with broad-spectrum antimicrobial activity
was ultimately screened out by the antibacterial spectrum assay (Table S4),
and named as *B. velezensis* HBXN2020. The strain was stored in 28%
glycerol at −80°C. For culturing the bacterial cells, Luria-Bertani (LB) media or
agar plates supplemented with 1.5% agar were used and incubated at 37°C.

A single bacterial colony was inoculated into 10 mL of LB medium and incubated overnight at 37°C with shaking (180 rpm). Then, the 16S rRNA sequence was amplified by the polymerase chain reaction (PCR) using forward primer 27F and reverse primer 1492R and similarity analysis with the National Center for Biotechnology Information (NCBI) alignment tool (BLAST) was used to identify the species.

Genome sequencing, annotation and analysis

The complete genome of *B. velezensis* HBXN2020 was sequenced using the Illumina HiSeq PE and PacBio RSII (SMRT) platforms (Shanghai Majorbio Bio-pharm Technology Co., Ltd.). Briefly, the short reads from the Illumina HiSeq PE were assembled into contigs using SPAdenovo (http://soap.genomics.org.cn/). The long reads from the PacBio RSII and the Illumina contigs were then aligned using the miniasm and Racon tools in Unicycler (version 0.4.8, https://github.com/rrwick/Unicycler) to generate long-read sequences. During the assembly process, sequence correction was performed using Pilon (version 1.22, https://github.com/broadinstitute/pilon/wiki/Standard-Output). Lastly, a complete genome with seamless chromosomes was obtained.

The coding sequences (CDs) of the *B. velezensis* HBXN2020 genome were predicted using Glimmer (version 3.02, http://ccb.jhu.edu/software/glimmer/index.shtml). The tRNA and rRNA genes

were predicted using TRNAscan-SE (version 2.0, http://trna.ucsc.edu/software/) and Barrnap (version 0.8, https://github.com/tseemann/barrnap), respectively. The functional annotation of all CDSs was performed using various databases, including Swiss-Prot Database (https://web.expasy.org/docs/swiss-prot_guideline.html), Pfam Database (http://pfam.xfam.org/), EggNOG Database (http://eggnog.embl.de/), GO Database (http://www.geneontology.org/), and KEGG Database (http://www.genome.jp/kegg/). The circular map of the *B. velezensis* HBXN2020 genome was generated using CGView (version 2, http://wishart.biology.ualberta.ca/cgview/) (56).

Comparative genomic analysis

To elucidate the phylogenetic relationships from a whole-genome perspective, a phylogenetic tree based on the whole genome was constructed using the Type (Strain) Genome Server (TYGS) online (https://ggdc.dsmz.de/) (57). The average nucleotide identity (ANI) values of genome to genome were calculated using the JSpeciesWS online service (58), and a heatmap was then generated using TBtools (version 1.113) (59).

Growth curves of *B. velezensis* HBXN2020

The growth curve of *B. velezensis* HBXN2020 was recorded in flat-bottomed 100-well microtiter plates via detecting optical density at 600 nm (OD_{600}) at 1 h intervals using the automatic growth curve analyzer (Bioscreen,

Helsinki, Finland).

Antimicrobial Assays

The *in vitro* antagonistic activity of *B. velezensis* HBXN2020 was tested

using the agar well-diffusion method against 18 indicator strains (pathogens),

which included 7 standard strains (*E. coli* ATCC 25922, *E. coli* ATCC 35150,

S. Typhimurium ATCC 14028 (STm), *S.* Typhimurium SL1344, *S. aureus*

ATCC 29213, *S. aureus* ATCC 43300, and *C. perfringens* CVCC 2030) and

11 clinical isolates (*E. coli* EC024, *S.* Enteritidis SE006, *S. aureus* S21, *C.*

perfringens CP023, *C. perfringens* CP002, *S. suis* SC19, *S. suis* SS006, *P.*

multocida PM002, *P. multocida* PM008, *A.* pleuropneumoniae APP015, and *A.*

pleuropneumoniae APP017). All plates were cultured at 37°C for 16 h before

observing the inhibition zone, and the diameter of the inhibition zone was

measured using a vernier caliper. The presence of a clear zone indicated

antagonistic activity.

Antibiotic susceptibility assays

Antimicrobial susceptibility testing was performed using the Kirby-Bauer

(KB) disk diffusion method in accordance with the Clinical Laboratory

Standards Institute (CLSI) guidelines (60). The antimicrobial agents tested

were: Ampicillin, Meropenem, Piperacillin, Gentamycin, Tetracycline,

Doxycycline, Minocycline, Erythromycin, Enrofloxacin, Ofloxacin,

Sulfamethoxazole, Trimethoprim-sulfamethoxazole, Polymyxin B, Teicoplanin,

Trimethoprim, Florfenicol, Spectinomycin, Nitrofurantoin, and Rifampicin. The

diameter of the inhibition zone was measured using a vernier caliper.

***In vitro* resistance assay of *B. velezensis* HBXN2020**

B. velezensis HBXN2020 spores (100 µL) or vegetative cells (100 µL) were

separately resuspended in 900 µL of LB medium supplemented with different

pH values (2, 3, 4, 5 or 6), bile salts (0.85% NaCl, 0.3%), simulated gastric

fluid (SGF, HCl, pH 1.2) containing 10 g L^{-1} of pepsin in 0.85% NaCl solution,

or simulated intestinal fluid (SIF, NaOH, pH 6.8) containing 10 g L^{-1} of trypsin

in 0.05 M KH2PO4 solution, and incubated at 37 °C. A normal LB medium (pH

7.0) was used as the control. At predetermined time points, 100 µL was taken

from each sample, serially diluted 10-fold with sterile PBS (pH 7.2), and then

spread onto LB agar plates. The plates were incubated overnight in a

constant temperature incubator at 37°C, and the bacterial colonies were

counted. The survival rate was calculated using the following formula: Survival

rate = (number of bacteria in the treatment group/number of bacteria in the

control group) × 100%.

One milliliter of *B. velezensis* HBXN2020 spores or vegetative cells were

separately placed in water baths at different temperatures (37°C, 45°C, 55°C,

65°C, 75°C, 85°C or 95°C) for 20 min, with a 37°C water bath used as the

control. The survival rate was calculated as described above.

In vitro bacterial competition.

To investigate whether _B. velezensis_ HBXN2020 directly inhibited the growth of STm, we performed spot-on lawn and agar-well diffusion assays, as well as co-culture assays in liquid culture medium. In the spot-on lawn antimicrobial assays, we prepared double layers of agar by first pouring LB agar into the plate as the bottom layer. The top layer consisted of 10 mL TSB broth containing 0.7% agar with STm overnight culture. Then, 10 μL of _B. velezensis_ HBXN2020 overnight culture and cell-free supernatant (CFS) were respectively spotted onto TSB agar and incubated at 37°C for 12 h to measure the inhibition zone. A transparent zone of at least 1 mm around the spot was considered positive.

The antagonistic effect of _B. velezensis_ HBXN2020-CFS against STm was determined using the agar-well diffusion assays. To collect _B. velezensis_ HBXN2020-CFS, the culture was centrifuged at 9,000g for 15 min at 4°C, and the supernatant was filtered through a 0.22 μm membrane filter (Millipore, USA). The STm lawn medium was prepared by mixing 10 mL of TSB broth containing 0.7% agar with STm overnight culture and then poured into a sterile plate covered with LB agar and Oxford cups, 8 mm diameter wells were prepared in the TSB agar after removing the cups. The wells were filled with 100 μL of _B. velezensis_ HBXN2020-CFS, LB medium or ampicillin (100 μg mL^{-1}). The plates were incubated at 37°C for 14 h, and the inhibition zone was

measured.

The co-culture assay was conducted by incubating *B. velezensis*

HBXN2020 and *S.* Typhimurium ATCC14028 (carry pET28a (+), kanamycin

resistance) (33) separately overnight at 37°C and diluting them to 10^4

CFU/mL. Then, the two strains were mixed at different ratios (1:1, 1:10, 1:50

or 1:100) and co-cultured at 37°C with shaking (180 rpm). At predetermined

time points, serial 10-fold dilutions were prepared for all samples and spread

onto selective (kanamycin 50 μg mL^{-1}) LB agar plates and cultured at 37°C for

12 h before bacterial counting. Viable colony counts ranged from 30 to 300

per plate.

Biosafety assessment

After a 7-day acclimation period (free access to water and food), mice were

randomly divided into four treatment groups (n = 5): low-dose *B. velezensis*

HBXN2020 spores group (L-HBXN2020 group, 10^7 CFU/mouse), medium-

dose *B. velezensis* HBXN2020 spores group (M-HBXN2020 group, 10^8

CFU/mouse), high-dose *B. velezensis* HBXN2020 spores group (H-

HBXN2020 group, 10^9 CFU/mouse), and a control group. During the

experimental period (15 days), mice were weighed and orally gavaged with

their respective treatments once every two days. On day 15, all mice were

euthanized, and blood, heart, liver, spleen, lung, kidney, ileum, cecum, and

colon were collected. Blood samples were used for routine blood and

biochemistry tests. Major organ tissues (heart, liver, spleen, lung, and kidney)

and a 5-mm distal segment of the colon were used for histopathology. The

remaining colonic tissues were rapidly frozen in liquid nitrogen and stored at

−80°C for cytokine and tight junction protein expression analysis.

***Salmonella* Typhimurium-induced mouse model of colitis.**

After 7 days of acclimation (free access to water and food), mice were

randomly divided into three treatment groups (n = 8): Control group, STm +

PBS group, and STm + HBXN2020 group. The *S.* Typhimurium-induced colitis

was performed as previously described (61), with slight modifications. On the

first day of the experiment, all mice in the STm + PBS group and STm +

HBXN2020 group were orally inoculated with 200 µL (5×10^7 CFU/mouse) of

STm. On days 1, 3 and 5 following STm infection, each mouse in the STm +

HBXN2020 group received 200 µL (1×10^8 CFU/mouse) of *B. velezensis*

HBXN2020 spores via gavage administration. In contrast, the control group

and STm + PBS group received 200 µL of sterile PBS by oral gavage. Fecal

samples were collected daily following STm infection and resuspended in

sterile PBS. The number of STm in mice feces from both the STm + PBS and

STm + HBXN2020 groups was then determined by spreading a serial 10-fold

dilution on selective LB agar plates containing 50 µg mL^{-1} kanamycin.

Throughout the entire experiment, the body weight, stool consistency and

fecal occult blood of all mice were monitored daily. As shown in Table S5,

disease activity index (DAI) was calculated by the sum of the scores from

three parameters (62). On day 7 after STm infection, all mice were euthanized,

their ileum, cecum and colon were collected. The length of colon was

measured, and a 5-mm distal segment of the colon was fixed in 4%

paraformaldehyde for further histopathology. The remaining colon was then

rapidly frozen in liquid nitrogen and stored at −80°C for cytokine and tight

junction protein expression analysis. Then, the number of STm in the ileum,

cecum, and colon was determined by spreading serial 10-fold dilutions on

selective LB agar plates.

To investigate the prophylactic efficacy of *B. velezensis* HBXN2020 in

ameliorating STm-induced colitis, another independent experiment was

conducted using six-week-old female C57BL/6 mice (SPF). After a seven-day

acclimation period, mice were randomly assigned to three groups (n = 8):

control group, PBS + STm group and HBXN2020 + STm group. On days 1, 3,

5 and 7, each mouse in the HBXN2020 + STm group received 200 μL ($1×10^8$

CFU/mouse) of *B. velezensis* HBXN2020 spores via gavage administration.

Meanwhile, mice in the control group and PBS + STm group received 200 μL

of sterile PBS by oral gavage. On day 7, all mice in the PBS + STm group and

HBXN2020 + STm group were orally inoculated with 200 μL ($5×10^7$

CFU/mouse) of STm. Fecal samples were collected daily following STm

infection from both groups and resuspended in sterile PBS. The number of

STm in the feces of mice was then determined by spreading serial 10-fold

dilutions on selective LB agar plates (50 μg mL^{-1} kanamycin). Throughout the

entire experiment, the body weight and DAI scores of all mice were monitored

daily. On day 12, all mice were euthanized, and the ileum, cecum, and colon

were collected. The colon length was measured, and a 5-mm distal segment

of the colon was fixed in 4% formalin for sectioning and staining. The

remaining colon was stored at −80°C for future analysis. Lastly, the number of

STm in the ileum, cecum, and colon was determined using a selective LB

agar plate.

Determination of cytokines and tight junction protein expression in

colon tissue

Total RNA was extracted from colon tissues using TRIpure reagent (Aidlab,

China), and cDNA was obtained using HiScript$^®$ III RT SuperMix for qPCR

(+gDNA wiper) (Vazyme, China). RT-qPCR for each gene was performed in

triplicate using qPCR SYBR Green Master Mix (Yeasen Biotechnology,

Shanghai, China). The relative expression level of cytokine and tight junction

protein genes was calculated using the $2^{-\Delta\Delta Ct}$ method with β-actin and

GAPDH as reference genes. The primer sequences used in the RT-qPCR test

are listed in Table S3.

Histopathology analysis

Colon tissue samples (0.5 cm) were fixed in 4% paraformaldehyde for 24 h,

and the fixed tissues were then embedded in paraffin and sectioned. The

sections were stained with hematoxylin and eosin (H&E) and observed and

imaged using an optical microscope (Olympus Optical, Tokyo, Japan). The

histopathological score included the degree of inflammatory infiltration,

changes in crypt structures, and the presence or absence of ulceration and

edema. The scoring criteria were determined as previously described (34).

16S rRNA gene sequencing and analysis

According to the manufacturer's instructions, colon microbial community

genomes were extracted using E.Z-N.A ® Stool DNA Kit (Omega; D4015-01),

and quality was detected by 1% agarose gel electrophoresis. The 16S rRNA

V3-V4 variable region was amplified by PCR using universal primers 338F (5

'- ACTCCTACGGGGGGCAG-3') and 806R (5 '-

GACTACHVGGGTWTCTAAT-3'). The PCR products were examined by

electrophoresis on 2% agarose gels and then purified with the AxyPrep DNA

gel extraction kit (Axygen Biosciences, USA). A sequencing library was

constructed using the NEXTFLEX® Rapid DNA-Seq Kit, and sequencing was

performed using the Illumina MiSeq platform. Raw reads were quality

evaluated and filtered by fastp (version 0.20.0) and merged using FLASH

(version 1.2.7). The optimized sequences were clustered into operational

taxonomic units (OTUs) based on 97% sequence similarity using UPARSE

(version 7.1). The representative sequences of each OTU was classified by

RDP classifier (version 2.2; confidence threshold value, 0.7). Alpha diversity

was assessed using the ACE, chao, Shannon, and Simpson indices. The β-

diversity analysis was performed using Bray-Curtis distances and visualized

through principal component analysis (PCA). Linear discriminant analysis

(LDA) effect size (LefSe) was used to identify differential microbiota between

groups.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism 8.3.0 (GraphPad

Software, San Diego, CA, USA) and Excel (Microsoft, Redmond, USA). Data

are presented as mean ± standard error of the mean (SEM). Differences

between two groups were evaluated using two-tailed unpaired Student's t-test,

and all other comparisons were conducted using one-way analysis of variance

(ANOVA) with Tukey's multiple comparisons test. For all analyses,

significance differences are denoted as: *, $P < 0.05$, **, $P < 0.01$, and ***, $P <$

0.001.

Data availability

The complete genome sequence of *Bacillus velezensis* HBXN2020 has

been deposited in GenBank and assigned accession numbers: CP119399.1

Conflict of interest: We declare no conflict of interest.

ACKNOWLEDGMENTS

This work was supported by grants from the National Program on Key Research Project of China (2021YFD1800300, 2022YFD1800800), "Yingzi Tech & Huazhong Agricultural University Intelligent Research Institute of Food Health" (No. IRIFH202209; IRIFH202301), and the Fundamental Research Funds for the Central Universities (2662016PY004).

Author contributions

Linkang Wang: Conceptualization, Methodology, Data curation, Formal analysis, Writing - original draft. **Haiyan Wang, Xinxin Li, Mengyuan Zhu**: Methodology, Data curation, Planned and performed the experiments. **Dongyang Gao, Dayue Hu, Zhixuan Xiong**: Methodology, Writing - Reviewing and Editing. **Xiangmin Li**: Supervision, Writing-reviewing and editing. **Ping Qian**: Conceptualization, Funding acquisition, Project administration, Supervision, Writing-Reviewing and Editing. All authors read and approved the submitted version of the paper.

Funding information

This work was supported by grants from the National Program on Key Research Project of China (2021YFD1800300, 2022YFD1800800), "Yingzi Tech & Huazhong Agricultural University Intelligent Research Institute of Food

795 Reference

1. Fabrega A, Vila J. 2013. Salmonella enterica serovar Typhimurium skills to succeed in the host:
virulence and regulation. Clin Microbiol Rev 26:308-41.

2. Yang WH, Westman JS, Heithoff DM, Sperandio M, Cho JW, Mahan MJ, Marth JD. 2021. Neu3
neuraminidase induction triggers intestinal inflammation and colitis in a model of recurrent
human food-poisoning. Proc Natl Acad Sci U S A 118.

3. Katiyo S, Muller-Pebody B, Minaji M, Powell D, Johnson AP, De Pinna E, Day M, Harris R,
Godbole G. 2019. Epidemiology and Outcomes of Nontyphoidal Salmonella Bacteremias from
England, 2004 to 2015. J Clin Microbiol 57.

4. Mohan A, Munusamy C, Tan YC, Muthuvelu S, Hashim R, Chien SL, Wong MK, Khairuddin NA,
Podin Y, Lau PS, Ng DC, Ooi MH. 2019. Invasive Salmonella infections among children in
Bintulu, Sarawak, Malaysian Borneo: a 6-year retrospective review. BMC Infect Dis 19:330.

5. Ferri M, Ranucci E, Romagnoli P, Giaccone V. 2017. Antimicrobial resistance: A global
emerging threat to public health systems. Crit Rev Food Sci Nutr 57:2857-2876.

6. Paulson JA, Zaoutis TE, Council on Environmental H, Committee on Infectious D. 2015.
Nontherapeutic Use of Antimicrobial Agents in Animal Agriculture: Implications for Pediatrics.
Pediatrics 136:e1670-7.

7. Reyman M, van Houten MA, Watson RL, Chu M, Arp K, de Waal WJ, Schiering I, Plotz FB,
Willems RJL, van Schaik W, Sanders EAM, Bogaert D. 2022. Effects of early-life antibiotics on
the developing infant gut microbiome and resistome: a randomized trial. Nat Commun
13:893.

8. Cristofori F, Dargenio VN, Dargenio C, Miniello VL, Barone M, Francavilla R. 2021. Anti-
Inflammatory and Immunomodulatory Effects of Probiotics in Gut Inflammation: A Door to
the Body. Front Immunol 12:578386.

9. Wang Y, Xie Q, Zhang Y, Ma W, Ning K, Xiang JY, Cui J, Xiang H. 2020. Combination of
probiotics with different functions alleviate DSS-induced colitis by regulating intestinal
microbiota, IL-10, and barrier function. Appl Microbiol Biotechnol 104:335-349.

10. Ohland CL, Macnaughton WK. 2010. Probiotic bacteria and intestinal epithelial barrier
function. Am J Physiol Gastrointest Liver Physiol 298:G807-19.

11. Gu SB, Zhao LN, Wu Y, Li SC, Sun JR, Huang JF, Li DD. 2015. Potential probiotic attributes of a
new strain of Bacillus coagulans CGMCC 9951 isolated from healthy piglet feces. World J
Microbiol Biotechnol 31:851-63.

12. Ouattara HG, Reverchon S, Niamke SL, Nasser W. 2017. Regulation of the synthesis of pulp
degrading enzymes in Bacillus isolated from cocoa fermentation. Food Microbiol 63:255-262.

13. Abdhul K, Ganesh M, Shanmughapriya S, Vanithamani S, Kanagavel M, Anbarasu K,
Natarajaseenivasan K. 2015. Bacteriocinogenic potential of a probiotic strain Bacillus
coagulans [BDU3] from Ngari. Int J Biol Macromol 79:800-6.

832 14. Ye M, Tang X, Yang R, Zhang H, Li F, Tao F, Li F, Wang Z. 2018. Characteristics and Application
833   of a Novel Species of Bacillus: Bacillus velezensis. ACS Chem Biol 13:500-505.

834 15. Shao J, Liu Y, Xie J, Stefanic P, Lv Y, Fan B, Mandic-Mulec I, Zhang R, Shen Q, Xu Z. 2022.
835   Annulment of Bacterial Antagonism Improves Plant Beneficial Activity of a Bacillus velezensis
836   Consortium. Appl Environ Microbiol 88:e0024022.

837 16. Li A, Wang Y, Li Z, Qamar H, Mehmood K, Zhang L, Liu J, Zhang H, Li J. 2019. Probiotics
838   isolated from yaks improves the growth performance, antioxidant activity, and cytokines
839   related to immunity and inflammation in mice. Microb Cell Fact 18:112.

840 17. Zhang DX, Kang YH, Zhan S, Zhao ZL, Jin SN, Chen C, Zhang L, Shen JY, Wang CF, Wang GQ,
841   Shan XF, Qian AD. 2019. Effect of Bacillus velezensis on Aeromonas veronii-Induced Intestinal
842   Mucosal Barrier Function Damage and Inflammation in Crucian Carp (Carassius auratus).
843   Front Microbiol 10:2663.

844 18. Herp S, Brugiroux S, Garzetti D, Ring D, Jochum LM, Beutler M, Eberl C, Hussain S, Walter S,
845   Gerlach RG, Ruscheweyh HJ, Huson D, Sellin ME, Slack E, Hanson B, Loy A, Baines JF, Rausch P,
846   Basic M, Bleich A, Berry D, Stecher B. 2019. Mucispirillum schaedleri Antagonizes Salmonella
847   Virulence to Protect Mice against Colitis. Cell Host Microbe 25:681-694 e8.

848 19. Schultz BM, Paduro CA, Salazar GA, Salazar-Echegarai FJ, Sebastian VP, Riedel CA, Kalergis AM,
849   Alvarez-Lobos M, Bueno SM. 2017. A Potential Role of Salmonella Infection in the Onset of
850   Inflammatory Bowel Diseases. Front Immunol 8:191.

851 20. Crits-Christoph A, Hallowell HA, Koutouvalis K, Suez J. 2022. Good microbes, bad genes? The
852   dissemination of antimicrobial resistance in the human microbiome. Gut Microbes
853   14:2055944.

854 21. Yang W, Liu Z, Zhao Q, Du H, Yu J, Wang H, Liu X, Liu H, Jing X, Yang H, Shi G, Zhou L, Liu J.
855   2022. Population Genetic Structure and Selection Signature Analysis of Beijing Black Pig.
856   Front Genet 13:860669.

857 22. Kaspar F, Neubauer P, Gimpel M. 2019. Bioactive Secondary Metabolites from Bacillus subtilis:
858   A Comprehensive Review. J Nat Prod 82:2038-2053.

859 23. Du H, Yao W, Kulyar MF, Ding Y, Zhu H, Pan H, Li K, Bhutta ZA, Liu S, Li J. 2022. Effects of
860   Bacillus amyloliquefaciens TL106 Isolated from Tibetan Pigs on Probiotic Potential and
861   Intestinal Microbes in Weaned Piglets. Microbiol Spectr 10:e0120521.

862 24. Sanders ME, Akkermans LM, Haller D, Hammerman C, Heimbach J, Hormannsperger G, Huys
863   G, Levy DD, Lutgendorff F, Mack D, Phothirath P, Solano-Aguilar G, Vaughan E. 2010. Safety
864   assessment of probiotics for human use. Gut Microbes 1:164-85.

865 25. Snydman DR. 2008. The safety of probiotics. Clin Infect Dis 46 Suppl 2:S104-11; discussion
866   S144-51.

867 26. Bazzoni G, Martinez-Estrada OM, Orsenigo F, Cordenonsi M, Citi S, Dejana E. 2000. Interaction
868   of junctional adhesion molecule with the tight junction components ZO-1, cingulin, and
869   occludin. J Biol Chem 275:20520-6.

870 27. Zihni C, Mills C, Matter K, Balda MS. 2016. Tight junctions: from simple barriers to
871   multifunctional molecular gates. Nat Rev Mol Cell Biol 17:564-80.

872 28. Gou S, Chen N, Wu X, Zu M, Yi S, Ying B, Dai F, Ke B, Xiao B. 2022. Multi-responsive
873   nanotheranostics with enhanced tumor penetration and oxygen self-producing capacities for
874   multimodal synergistic cancer therapy. Acta Pharm Sin B 12:406-423.

875 29. Ozer J, Ratner M, Shaw M, Bailey W, Schomaker S. 2008. The current state of serum

biomarkers of hepatotoxicity. Toxicology 245:194-205.

30.   Tang Y, Gao C, Xing M, Li Y, Zhu L, Wang D, Yang X, Liu L, Yao P. 2012. Quercetin prevents
ethanol-induced dyslipidemia and mitochondrial oxidative damage. Food Chem Toxicol
50:1194-200.

31.   Singhal M, Liu X, Inverso D, Jiang K, Dai J, He H, Bartels S, Li W, Abdul Pari AA, Gengenbacher
N, Besemfelder E, Hui L, Augustin HG, Hu J. 2018. Endothelial cell fitness dictates the source
of regenerating liver vasculature. J Exp Med 215:2497-2508.

32.   Zhang L, Bai K, Zhang J, Xu W, Huang Q, Wang T. 2017. Dietary effects of Bacillus subtilis fmbj
on the antioxidant capacity of broilers at an early age. Poult Sci 96:3564-3573.

33.   Cao Z, Wang X, Pang Y, Cheng S, Liu J. 2019. Biointerfacial self-assembly generates lipid
membrane coated bacteria for enhanced oral delivery and treatment. Nat Commun 10:5783.

34.   Wu J, Lin Z, Wang X, Zhao Y, Zhao J, Liu H, Johnston LJ, Lu L, Ma X. 2022. Limosilactobacillus
reuteri SLZX19-12 Protects the Colon from Infection by Enhancing Stability of the Gut
Microbiota and Barrier Integrity and Reducing Inflammation. Microbiol Spectr 10:e0212421.

35.   Dinarello CA. 2000. Proinflammatory cytokines. Chest 118:503-8.

36.   Cheng A, Yan H, Han C, Wang W, Tian Y, Chen X. 2014. Polyphenols from blueberries
modulate inflammation cytokines in LPS-induced RAW264.7 macrophages. Int J Biol
Macromol 69:382-7.

37.   Liu C, Chu D, Kalantar-Zadeh K, George J, Young HA, Liu G. 2021. Cytokines: From Clinical
Significance to Quantification. Adv Sci (Weinh) 8:e2004433.

38.   Cross AS, Sadoff JC, Kelly N, Bernton E, Gemski P. 1989. Pretreatment with recombinant
murine tumor necrosis factor alpha/cachectin and murine interleukin 1 alpha protects mice
from lethal bacterial infection. J Exp Med 169:2021-7.

39.   Huang FC. 2009. Upregulation of Salmonella-induced IL-6 production in Caco-2 cells by PJ-34,
PARP-1 inhibitor: involvement of PI3K, p38 MAPK, ERK, JNK, and NF-kappaB. Mediators
Inflamm 2009:103890.

40.   Karki R, Sharma BR, Tuladhar S, Williams EP, Zalduondo L, Samir P, Zheng M, Sundaram B,
Banoth B, Malireddi RKS, Schreiner P, Neale G, Vogel P, Webby R, Jonsson CB, Kanneganti TD.
2021. Synergism of TNF-alpha and IFN-gamma Triggers Inflammatory Cell Death, Tissue
Damage, and Mortality in SARS-CoV-2 Infection and Cytokine Shock Syndromes. Cell 184:149-
168 e17.

41.   Kotas ME, Medzhitov R. 2015. Homeostasis, inflammation, and disease susceptibility. Cell
160:816-827.

42.   Ouyang W, Rutz S, Crellin NK, Valdez PA, Hymowitz SG. 2011. Regulation and functions of the
IL-10 family of cytokines in inflammation and disease. Annu Rev Immunol 29:71-109.

43.   Saraiva M, Vieira P, O'Garra A. 2020. Biology and therapeutic potential of interleukin-10. J Exp
Med 217.

44.   Maloy KJ, Powrie F. 2011. Intestinal homeostasis and its breakdown in inflammatory bowel
disease. Nature 474:298-306.

45.   Hayashi A, Sato T, Kamada N, Mikami Y, Matsuoka K, Hisamatsu T, Hibi T, Roers A, Yagita H,
Ohteki T, Yoshimura A, Kanai T. 2013. A single strain of Clostridium butyricum induces
intestinal IL-10-producing macrophages to suppress acute experimental colitis in mice. Cell
Host Microbe 13:711-22.

46.   Chen L, Wang D, Garmaeva S, Kurilshikov A, Vich Vila A, Gacesa R, Sinha T, Segal E, Weersma

RK, Wijmenga C, Zhernakova A, Fu J. 2021. The long-term genetic stability and individual
specificity of the human gut microbiome. Cell 184:2302-2315.e12.

47.   Wang L, Gou X, Ding Y, Liu J, Wang Y, Wang Y, Zhang J, Du L, Peng W, Fan G. 2023. The
interplay between herbal medicines and gut microbiota in metabolic diseases. Front
Pharmacol 14:1105405.

48.   Kaur H, Gupta T, Kapila S, Kapila R. 2021. Protective effects of potential probiotic Lactobacillus
rhamnosus (MTCC-5897) fermented whey on reinforcement of intestinal epithelial barrier
function in a colitis-induced murine model. Food Funct 12:6102-6116.

49.   Cani PD, Depommier C, Derrien M, Everard A, de Vos WM. 2022. Akkermansia muciniphila:
paradigm for next-generation beneficial microorganisms. Nat Rev Gastroenterol Hepatol
19:625-637.

50.   Polk BF, Kasper DL. 1977. Bacteroides fragilis subspecies in clinical isolates. Ann Intern Med
86:569-71.

51.   Pickard JM, Zeng MY, Caruso R, Núñez G. 2017. Gut microbiota: Role in pathogen colonization,
immune responses, and inflammatory disease. Immunol Rev 279:70-89.

52.   Morgan KL. 2013. Infection, illness, and fenugreek seeds: was illness a good enough proxy for
infection in the 2011 outbreak of shiga toxin-producing Escherichia coli O104:H4 in France?
Clin Infect Dis 56:1055-6.

53.   Hardt WD, Chen LM, Schuebel KE, Bustelo XR, Galán JE. 1998. S. typhimurium encodes an
activator of Rho GTPases that induces membrane ruffling and nuclear responses in host cells.
Cell 93:815-26.

54.   Parker BJ, Wearsch PA, Veloo ACM, Rodriguez-Palacios A. 2020. The Genus Alistipes: Gut
Bacteria With Emerging Implications to Inflammation, Cancer, and Mental Health. Front
Immunol 11:906.

55.   Tang Z, Sun H, Chen T, Lin Z, Jiang H, Zhou X, Shi C, Pan H, Chang O, Ren P, Yu J, Li X, Xu J,
Huang Y, Yu X. 2017. Oral delivery of Bacillus subtilis spores expressing cysteine protease of
Clonorchis sinensis to grass carp (Ctenopharyngodon idellus): Induces immune responses and
has no damage on liver and intestine function. Fish Shellfish Immunol 64:287-296.

56.   Stothard P, Wishart DS. 2005. Circular genome visualization and exploration using CGView.
Bioinformatics 21:537-9.

57.   Meier-Kolthoff JP, Carbasse JS, Peinado-Olarte RL, Goker M. 2022. TYGS and LPSN: a database
tandem for fast and reliable genome-based classification and nomenclature of prokaryotes.
Nucleic Acids Res 50:D801-D807.

58.   Richter M, Rossello-Mora R, Oliver Glockner F, Peplies J. 2016. JSpeciesWS: a web server for
prokaryotic species circumscription based on pairwise genome comparison. Bioinformatics
32:929-31.

59.   Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, Xia R. 2020. TBtools: An Integrative
Toolkit Developed for Interactive Analyses of Big Biological Data. Mol Plant 13:1194-1202.

60.   CLSI. 2018. Performance standards for antimicrobial disk susceptibility tests, M02, 13th ed.
Clinical and Laboratory Standards Institute, Wayne, PA.

61.   Stecher B, Macpherson AJ, Hapfelmeier S, Kremer M, Stallmach T, Hardt WD. 2005.
Comparison of Salmonella enterica serovar Typhimurium colitis in germfree mice and mice
pretreated with streptomycin. Infect Immun 73:3228-41.

62.   Praveschotinunt P, Duraj-Thatte AM, Gelfat I, Bahl F, Chou DB, Joshi NS. 2019. Engineered E.

coli Nissle 1917 for the delivery of matrix-tethered therapeutic domains to the gut. Nat
Commun 10:5580.

Figure legends

Fig. 1 Genomic characteristics and phylogenetic relationships of _B._

velezensis HBXN2020. (**A**) The whole-genome map of _B. velezensis_

HBXN2020 with its genomic features. The map consists of 6 circles. From the

inner circle to the outer circle: (1) GC-skew, (2) GC content, (3) reverse

protein-coding genes, different colors represents different COG functional

classifications, (4) genes transcribed in reverse direction, (5) genes

transcribed in forward direction, (6) forward protein-coding genes, different

colors represents different COG functional classifications. (**B**) The whole

genome phylogenetic tree was constructed based on genome-wide data from

14 bacillus strains. _B. velezensis_ HBXN2020 are indicated in bold. (**C**) The

heatmap based on the ANIb value of strain HBXN2020 and other Bacillus

species. The _B. velezensis_ HBXN2020 was labeled in red letters.

Fig. 2 Growth curve of _B. velezensis_ HBXN2020 and its _in vitro_

resistance against environmental assaults. (**A**) Growth curves of _B._

velezensis HBXN2020 cultured in LB medium at 37°C, detection of OD_{600}

values at 1 h intervals in microplate reader. (**B**) Survival rate of endospore

and vegetative cells of _B. velezensis_ HBXN2020 after 30 min at different

temperatures (37°C, 45°C, 55°C, 65°C, 75°C, 85°C or 95°C). Equal amounts

of endospore and vegetative cells of HBXN2020 were exposed to the

following: (**C**) acid solution (pH 2 to 7), (**D**) 0.3% bile salts, (**E**) SGF (pH 1.2)

supplemented with pepsin, and (**F**) SIF (pH 6.8) containing trypsin at 37°C. At

predetermined time points, 100 μL was taken from each sample, and tenfold

serially diluted with sterile PBS (pH 7.2), and then spread on LB agar plates

and cultured at 37°C for 12 h before bacterial counting. Each group was

repeated three times (n = 3).

Fig. 3 Antibiotic susceptibility of *B. velezensis* **HBXN2020 and**

bacteriostasis assay *in vitro.* (**A**) The diameter of the antibacterial zone

indicates the extent of sensitivity to antibiotics. (**B**) The diameter of the

antibacterial zone indicates the extent of inhibition against pathogenic bacteria.

The diameter of the antibacterial zone was measured with vernier caliper.

Each group was repeated three times (n = 3). R, resistant; I, moderately

sensitive; S, sensitive.

1003

1004 **Fig. 4** *In vivo* **safety evaluation of** *B. velezensis* **HBXN2020 in a mouse**

1005 **model.** (**A**) Body weights changes of mice during gavage with *B. velezensis*

1006 HBXN2020 spores. Mice were treated with sterile PBS (Control group) or low-

1007 dose (L-HBXN2020 group), medium dose (M-HBXN2020 group), and high-

1008 dose (H-HBXN2020 group) of *B. velezensis* HBXN2020 spores. Weighing and

gavage were performed once every two days during the experimental period

(15 days). Data were shown as mean values ± SEM (n = 5). (**B**) The relative

gene expression levels of inflammatory cytokines in the colon of mice

measured by RT-qPCR. Data were shown as mean values ± SEM (n = 5). (**C**)

The relative gene expression levels of barrier protein ZO-1, occludin, claudin

and Muc2 in the colon of mice measured by RT-qPCR. Data were shown as

mean values ± SEM (n = 5). (**D**) Major blood routine parameters and (**E**)

serum biochemical parameters of mice in the control group and H-HBXN2020

group. Data were shown as mean values ± SEM (n = 3). (**F**) H&E stained

colon sections in the different groups. Scale bar: 200 μm.

Fig. 5 Oral *B. velezensis* HBXN2020 spores alleviated *S.* Typhimurium-induced experimental mouse colitis. (**A**) *In vitro* Bacterial competition

between STm and *B. velezensis* HBXN2020. STm were co-incubated with *B.*

velezensis HBXN2020 at various ratios at 37°C with shaking. The growth of

STm was reflected by bacterial counting per hour. (**B**) Experimental design for

treatment in this study. Orally administrated with either PBS, or *B. velezensis*

HBXN2020 spores by gavage at days 1, 3, and 5 after STm infection (5× 10^7

CFU/mouse), respectively. All mice were euthanized at day 7 after STm

infection. (**C**) Bacterial count of STm in mouse feces. Fecal samples were

collected per day after STm infection and resuspended in sterile PBS (0.1 g of

fecal resuspended in 1 mL of sterile PBS). One hundred microliters of each

sample performed a serial of 10-fold dilutions and spread on selective agar

plates (50 µg mL^{-1} kanamycin) and incubated at 37°C for 12 h before bacterial

counting. The bacterial loads of STm in (**D**) cecum, and (**E**) colon. The cecum

and colon were harvested and then homogenized. One hundred microliters of

each sample performed a serial of 10-fold dilutions and spread on selective

agar plates (50 µg mL^{-1} kanamycin) and incubated at 37°C for 12 h before

bacterial counting. Statistical signifcance was evaluated using Student's t-test

(*, P < 0.05, **, P < 0.01, and ***, P < 0.001). (**F**) Daily body weight changes

and (**G**) daily disease activity index (DAI) scores of mice with different

treatment groups. Data were shown as mean values ± SEM (n = 8). Statistical

signifcance was evaluated using one-way analysis of variance (ANOVA) with

Tukey's multiple comparisons test. *, P < 0.05, **, P < 0.01, and ***, P < 0.001

relative to Control group; $^{#}$, P < 0.05, $^{##}$, P < 0.01, $^{###}$, P < 0.001 relative to

STm + HBXN2020 group. (**H**) Colonic tissue images. (**I**) The length of the

colon from per group (n = 8). Statistical signifcance was evaluated using one-

way analysis of variance (ANOVA) with Tukey's multiple comparisons test (*,

P < 0.05, **, P < 0.01, and ***, P < 0.001).

1048

1049 **Fig. 6 Oral *B. velezensis* HBXN2020 spores attenuated colonic damage**

1050 **and inflammatory reaction.** (**A**) H&E stained colon tissue sections. Scale bar:

1051 200 µm. The relative gene expression levels of TNF-α (**B**), IL-1β (**C**), IL-6 (**D**),

1052 and IL-10 (**E**) were detected by RT-qPCR. Data were shown as mean values

± SEM (n = 6). The relative gene expression levels of ZO-1 (**F**), ocludin (**G**), claudin (**H**), and Muc2 (**I**) in colon tissue were detected by RT-qPCR. Data were shown as mean values ± SEM (n = 6). Statistical signifcance was evaluated using one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test (*, $P < 0.05$, **, $P < 0.01$, and ***, $P < 0.001$).

Fig. 7 Oral *B. velezensis* HBXN2020 spores regulated the composition of intestinal microbiota. (**A** to **D**) The α-diversity of the gut microbiota, determined by the (**A**) Sobs, (**B**) Chao1, (**C**) Shannon, and (**D**) Simpson diversity index. Data were shown as mean values ± SEM (n = 5). (**E**) The PCA plot showed the β-diversity of the gut microbiota based on Bray-Curtis distance at the OTU level. (**F** and **G**) The relative abundance of colonic microbiota at the phylum (**F**) and genus (**G**) levels. (**H-K**) Relative abundance of selected taxa (**H**) *norank_f_Muribaculaceae*, (**I**) *Lactobacillus*, (**J**) *Bacteroides*, and (**K**) *Escherichia-Shigella*. Data were shown as mean values ± SEM (n = 5). (**L**) Analysis of differences in the microbial communities by LEfSe (linear discriminant analysis (LDA) score > 3.5) among different groups. Signifcance was evaluated by the Kruskal-Wallis test or ANOVA with Tukey's multiple comparisons test (*, P < 0.05, **, P < 0.01, and ***, P < 0.001).

Fig. 8 Prophylactic *B. velezensis* HBXN2020 spores attenuated the symptoms of *S.* Typhimurium-induced mouse colitis. (**A**) Experimental

design for treatment in this study. At days 1, 3, 5, and 7, each mouse in the HBXN2020 + STm group and PBS + STm group were received 200 μL (1×10^8 CFU/mouse) of *B. velezensis* HBXN2020 spores or sterile PBS by gavage. Then, mice in PBS + STm group and HBXN2020 + STm group were orally inoculated with 200 μL (5×10^7 CFU/mouse) of STm on day 7. On day 12, all mice were euthanized. (**B**) Daily body weight changes and (**C**) daily disease activity index (DAI) scores of mice with different groups following STm treatment. Data were shown as mean values ± SEM (n = 8). Statistical signifcance was evaluated using one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test. *, $P < 0.05$, **, $P < 0.01$, and ***, $P < 0.001$ relative to Control group; $^\#$, $P < 0.05$, $^{\#\#}$, $P < 0.01$, $^{\#\#\#}$, $P < 0.001$ relative to HBXN2020 + STm group. (**D**) Colonic tissue images. (**E**) The length of the colon from per group (n = 8). (**F**) Bacterial count of STm in mouse feces. Fecal samples were collected every day after STm infection and resuspended in sterile PBS (0.1 g of fecal resuspended in 1 mL of sterile PBS). One hundred microliters of each sample performed a serial of 10-fold dilutions and spread on selective agar plates (50 μg mL^{-1} kanamycin) and incubated at 37°C for 12 h before bacterial counting. The bacterial loads of STm in (**G**)cecum and (**H**)colon. The cecum and colon were harvested and then homogenized. One hundred microliters of each sample performed a serial of 10-fold dilutions and spread on selective agar plates (50 μg mL^{-1} kanamycin) and incubated at 37°C for 12 h before bacterial counting. Statistical

1097 signifcance was evaluated using Student's t-test ([*], $P < 0.05$, [**], $P < 0.01$, and

1098 [***], $P < 0.001$).

1099

Fig. 9 Prophylactic *B. velezensis* HBXN2020 spores attenuated colonic

damage and inflammatory reaction. (**A**) H&E stained colon tissue sections.

Scale bar: 200 μm. The relative gene expression levels of TNF-α (**B**), IL-1β

(**C**), IL-6 (**D**), and IL-10 (**E**) were detected by RT-qPCR. Data were shown as

mean values ± SEM (n = 6). The relative gene expression levels of ZO-1 (**F**),

ocludin (**G**), claudin (**H**), and Muc2 (**I**) in colon tissue were detected by RT-

qPCR. Data were shown as mean values ± SEM (n = 6). Statistical

signifcance was evaluated using one-way analysis of variance (ANOVA) with

Tukey's multiple comparisons test ([*], $P < 0.05$, [**], $P < 0.01$, and [***], $P < 0.001$).

1109

Fig. 10 Prophylactic *B. velezensis* HBXN2020 spores regulated the

composition of gut microbiota. (**A** to **D**) Alpha diversity of the intestinal

microbiota. Data were shown as mean values ± SEM (n = 5). (**E**) The PCA

plot showed the β-diversity among different microbial community groups

based on Bray-Curtis distance at the OTU level. (**F** and **G**) The relative

abundance of colonic microbiota at the phylum (**F**) and genus (**G**) levels. (**H-M**)

Relative abundance of selected taxa (**H**) *Lactobacillus*, (**I**) *Akkermansia*, (**J**)

Alistipes, (**K**) *Bacteroides*, (**L**) *Escherichia-Shigella*, and (**M**) *Enterococcus*.

Data were shown as mean values ± SEM (n = 5). (**N**) Analysis of differences

in the microbial taxa by LEfSe (LDA score > 3.5) in different groups.

Signifcance was evaluated by the Kruskal-Wallis test or ANOVA with Tukey's

multiple comparisons test (*, $P < 0.05$, **, $P < 0.01$, and ***, $P < 0.001$).

1122

A

B