<u>Title</u>

Human leukocyte antigen HLA-DR-expressing fibroblast-like synoviocytes are

inducible antigen presenting cells that present autoantigens in Lyme arthritis

<u>Authors</u>

Joseph R Rouse[1], Rebecca Danner[1], Amanda Wahhab[1], Michaela Pereckas[2], Mecaila

E McClune[3,4], Allen C Steere[5], Klemen Strle[6], Brandon L Jutras[3,4], Robert B

Lochhead[1,7]

<u>Affiliations</u>

[1] Department of Microbiology and Immunology, Medical College of Wisconsin,

Milwaukee, WI, USA

[2] Department of Biochemistry, Medical College of Wisconsin, Milwaukee, WI, USA

[3] Department of Biochemistry, Virginia Tech, Blacksburg, VA, USA

[4] Center for Emerging, Zoonotic and Arthropod-borne Pathogens, Virginia Tech,

Blacksburg, VA, USA

[5] Center for Immunology and Inflammatory Diseases, Division of Rheumatology, Allergy,

and Immunology, Massachusetts General Hospital, Harvard Medical School, Boston,

MA, USA

[6] Department of Molecular Biology and Microbiology, Tufts University School of

Medicine, Boston, MA, USA

[7] Division of Rheumatology, Department of Medicine, Medical College of Wisconsin,

Milwaukee, WI, USA

Send correspondence to: Robert B Lochhead, PhD, Medical College of Wisconsin

Department of Microbiology and Immunology BSB-2850, 8701 Watertown Plank Road,

Milwaukee, Wisconsin, 53226, USA

Email address: rlochhead@mcw.edu

Telephone number: 414-955-4882

AUTHORS' SUMMARY

This study demonstrates that IFNγ-activated MHC-II+ fibroblast-like synoviocytes (FLS)

stimulated with *Borrelia burgdorferi* present foreign and self MHC-II antigens, including

Lyme autoantigens. Furthermore, IFNγ-activated MHC-II+ FLS stimulated with *B.*

burgdorferi peptidoglycan can induce activation and proliferation of naïve CD4+ T cells

in an MHC-II antigen-dependent manner, demonstrating that activated MHC-II+ FLS are

inducible antigen presenting cells.

ABSTRACT

Background: HLA-DR-expressing fibroblast-like synoviocytes (FLS) are a prominent

cell type in synovial tissue in chronic inflammatory forms of arthritis. We recently

showed that peptides from several extracellular matrix (ECM) proteins, including

fibronectin-1 (FN1), contained immunogenic CD4+ T cell epitopes in patients with

postinfectious Lyme arthritis (LA). However, the role of FLS in presentation of these T

cell epitopes remains uncertain.

Methods: Primary LA FLS and primary murine FLS stimulated with interferon gamma

(IFNγ), *Borrelia burgdorferi*, and/or *B. burgdorferi* peptidoglycan (PG) were assessed for

properties associated with antigen presentation. HLA-DR-presented peptides from

stimulated LA FLS were identified by immunopeptidomics analysis. OT-II T cells were

cocultured with stimulated murine FLS in the presence of cognate ovalbumin antigen to

determine the potential of FLS to act as inducible antigen presenting cells (APC).

Results: FLS expressed HLA-DR molecules within inflamed synovial tissue and

tendons from patients with post-infectious LA patients *in situ.* MHC class II and

costimulatory molecules were expressed by FLS following *in vitro* stimulation with IFNγ

and *B. burgdorferi* and presented both foreign and self MHC-II peptides, including T cell

epitopes derived from two Lyme autoantigens fibronectin-1 (FN1) and endothelial cell

growth factor (ECGF). Stimulated murine FLS induced proliferation of naïve OT-II CD4+

T cells, particularly when FLS were stimulated with both IFNγ and PG.

Conclusions: MHC-II+ FLS are inducible APCs that can induce CD4+ T cell activation

and can present Lyme autoantigens derived from ECM proteins, thereby amplifying

tissue-localized autoimmune CD4+ T cell responses in LA.

INTRODUCTION

Human leukocyte antigen DR-expressing fibroblast-like synoviocytes (HLA-DR+

FLS) are believed to be key drivers of synovial inflammation (1-4). We have recently

demonstrated that synovial extracellular matrix (ECM) proteins, including fibronectin-1

(FN1), are targets of autoimmune CD4+ T cell responses in over half of patients with

postinfectious Lyme arthritis (LA), defined as patients who fail to fully resolve their

arthritis after 2-3 months of appropriate antibiotic therapy and apparent killing of the

Lyme disease spirochete *Borrelia burgdorferi*. FLS are major producers of ECM

proteins, and HLA-DR+ FLS are highly abundant in postinfectious LA synovial tissue

(5), and we have hypothesized that FLS are both producing and presenting Lyme

autoantigens (1, 2). However, this hypothesis has not yet been fully tested.

FLS are synovial tissue resident cells that are important in joint health and

disease. FLS are 1) the primary cell type within joint synovial tissue; 2) responsible for

production and maintenance of extracellular matrix; 3) critical for tissue repair; and 4)

required to achieve homeostasis following an inflammatory insult (6). Under chronic

inflammatory conditions, however, FLS play a critical role in driving pathogenesis. In

RA, the prototypic autoimmune joint disease, activated HLA-DR+ FLS accumulate

within the synovial lining and sub-lining, where they interact with CD4+ T cells and other

immune cells (7-11). Furthermore, abundance of activated HLA-DR+ FLS positively

correlates with RA disease activity (12).

Lyme disease (LD) is caused by the tick-borne spirochete *Borrelia burgdorferi*

and occurs primarily in temperate zones of North America, Europe, and Asia. Each

year, an estimated 500,000 new cases occur in the U.S. and the infection is epidemic in

affected regions (13, 14). LA is the most common late-stage LD manifestation in the

U.S. and is usually effectively treated with 1-3 months of oral and, if necessary, IV

antibiotic therapy, called antibiotic-responsive LA (15). However, in a subset (~5%) of

patients, arthritis can persist or worsen, even after apparent spirochetal killing by

appropriate antibiotic therapy, called post-infectious LA (15). Post-infectious LA is

characterized by severe synovial hyperplasia and accumulation of large numbers of

HLA-DR+ FLS within inflamed joints (2, 5, 16-18), and is frequently accompanied by

autoimmune T and B cell responses targeting self-proteins associated with vascular

inflammation and ECM damage (1, 2, 15, 19-22). In addition, peptidoglycan (PG), a

pathogen-associated molecular pattern (PAMP) that is an essential cell wall component

in most bacteria required for protection from osmotic pressure, is a key *B. burgdorferi*

antigen that can persist in joints of post-infectious LA patients up to several years

following oral and/or IV antibiotic therapy. This spirochetal remnant may contribute to

persistent arthritis after the infection itself is cleared (23, 24).

Bacterial PG has also been detected in the inflamed synovium of patients with

RA (25) and osteoarthritis (OA) (26), suggesting that PG accumulates in synovial tissue,

particularly within the context of joint inflammation following an infection, such as in

postinfectious LA. Moreover, PG has been reported to have an immune adjuvant effect

on FLS (27, 28). In this study, we demonstrate that HLA-DR+ FLS can present MHC

class II (MHC-II) epitopes, including a Lyme autoantigen derived from ECM protein FN1,

and that primary murine FLS stimulated with IFNγ and *B. burgdorferi* PG can induce

proliferation and activation of CD4+ T cells in an antigen-dependent manner.

MATERIALS AND METHODS

Ethics statement

Mice were housed in the Medical College of Wisconsin Biomedical Resource Center

(Milwaukee, WI), following strict adherence to the guidelines according to the National

Institutes of Health for the care and use of laboratory animals, as described in the Guide

for the Care and Use of Laboratory Animals, 8th Edition. Protocols conducted in this

study were approved and carried out in accordance with the Medical College of

Wisconsin Institutional Animal Care and Use Committee (IACUC protocol number

AUA00006528). Mouse experiments were performed under isoflurane anesthesia, and

every effort was made to minimize suffering.

The study "Immunity in Lyme Arthritis" was approved by the Human

Investigations Committee at Massachusetts General Hospital (MGH), according to

principles for medical research involving human subjects expressed in the World

Medical Association Declaration of Helsinki. Written informed consent was obtained

from all participants 18 years of age or older, or from patient and a parent or guardian of

those between the ages of 12 and 17.

Patients

All patients with Lyme disease met the Centers for Disease Control and Prevention

criteria for *B. burgdorferi* infection (29). All patients had swollen knees and high IgG

antibody responses to *B. burgdorferi,* as determined by ELISA and Western blot. LA

patients received treatment according to an algorithm (15), as mentioned in the

guidelines of the Infectious Diseases Society of America (30).

Human LA FLS analysis and LC-MS/MS

Human LA tissue biopsies and FLS were analyzed analyzed as described (5) and as

detailed in the Supplemental Methods. HLA-DR peptide isolation was performed as

described (31), with modifications detailed in the Supplemental Methods. MS data were

analyzed using Proteome Discoverer 2.4 (Thermo) platform. SequestHT was used as a

search algorithm. Fixed Value (FV) and Target-Decory (TD) peptide-spectrum match

(PSM) validators and Protein FDR validator were used to identify potential PSM-peptide

matches. For FV validation, PSMs with cross-correlation scores (Xcorr) lower than 2.5

were excluded as potential matches. Proteome databases searched for this study were

Borrelia burgdorferi (N40 and B31) and Human (Swissprot, with isoforms).

Bacteria and mouse strains

Mouse strains C57B6/J (B6) and B6.Cg-Tg(TcraTcrb)425Cbn/J (OT-II) were

acquired from Jackson Laboratory. The class II major histocompatibility complex trans-

activator (CIITA) EGFP-reporter mouse was generated as described (32) and back-

crossed onto the C57BL/6J background for 7 generations. For the *in vitro* stimulation

experiments using live bacteria, we used *B. burgdorferi* strain B31-A2.

Primary cell stimulation and analysis

Bone marrow-derived macrophages (BMDM) and murine FLS were isolated from

bone marrow and ankle joints as described (33). FLS were passaged at least 3 times to

remove contaminating cells with 10% DMEM. Cells were stimulated for 48 hours with

IFNγ (20 ng/ mL), *B. burgdorferi* (1×10^6 cells/ mL), *B. burgdorferi* PG (10 µg/ mL). *B.*

burgdorferi PG was purified as previously described (34). Cells and secreted cytokines

were analyzed by flow cytometry, as recommended by the manufacturer (Biolegend).

For a complete list of commercial antibodies and reagents used, see Supplemental

Methods.

IncucyteTM imaging of CIITA EGFP-reporter FLS

CIITA EGFP-reporter FLS (passage four), were stimulated with 0, 0.2, 2, or 20 ng/mL IFNγ and compared to non-GFP-tagged controls. Cells were plated at 70,000 cells per well in 100 µl of medium in a 96-well plate and were allowed to rest overnight in a 32°C incubator with 5% CO_2. Green fluorescence (ex:440-480 nm, em:504-544 nm) was used to detect the EGFP reporter and analyzed by IncuCyte imaging with images taken every two hours for 96 hours total at 10X objective. Statistical analysis was performed using Tukey's multiple comparisons test ($p < 0.05$). Error bars represent standard deviation.

T cell proliferation assay and flow cytometric analysis

FLS were cultured from frozen stock for 24h in 24-well plates (1×10^5 cells/ well) then stimulated for 48 hours with IFNγ, *B. burgdorferi* PG, or both in 2% DMEM at 37°C. CD3+ T cells were isolated from spleens of 6-to-8-week-old OT-II mice by MojoSort mouse CD3+ T cell isolation kit (Biolegend 480024). Stimulated FLS were washed with PBS and co-cultured with CD3+ T cells at a concentration of 5-10×10^5 cells/mL in 10% RPMI for 5 days with either the cognate $OVA_{323\text{-}339}$ antigen (2μg/mL, a non-reactive $OVA_{257\text{-}264}$ peptide (2μg/mL), or ovalbumin (10ug/mL). IL-2 (0.5 μg/10^5 cells) was used to maintain T cell viability.

Statistical analysis

Statistically significant differences between groups were determined by one-way ANOVA and Tukey's multiple comparisons using GraphPad Prism (v.9). For gene

expression analysis, copies per million reads (CPMR) was used to normalize gene

expression.

RESULTS

Human LA FLS express HLA-DR within synovial tissue and inflamed tendons

Synovial hyperplasia and accumulation of activated FLS expressing HLA-DR

MHC-II molecules are hallmarks of human post-infectious LA (2, 5). We examined HLA-

DR expression of vimentin+ stromal cells in synovial tissue and tendon biopsies from

patients with post-infectious LA (Figure 1A). HLA-DR+/Vimentin+ double-positive

stromal cells (e.g., HLA-DR+ FLS) were primarily localized within the synovial sub-lining

(Figure 1A, patients 1 & 2), consistent with a similar study examining a larger number of

synovial tissue biopsies (5). In two post-infectious LA patients with tenosynovitis for

which tendons were available, double-positive cells were also observed primarily along

the lining of inflamed joint tendons (Figure 1A, patients 3 & 4). This would mark the first

report, to our knowledge, of HLA-DR+ stromal cells observed in post-infectious LA

patient tendons, which is consistent with reports of ligamentous tissue as a reservoir of

B. burgdorferi during chronic Lyme borreliosis (35) and significant tendinopathy seen in

LA during active infection (36) and in post-infectious LA (37).

Human LA FLS express genes associated with antigen presentation when

stimulated with IFNγ

FLS do not express the canonical B7 costimulatory molecules B7-1 (CD80) or

B7-2 (CD86), although FLS express other B7-family costimulatory molecules that may

play a role in antigen presentation (38). We examined expression of potential co-

stimulatory molecules using a human LA FLS RNAseq dataset generated as part of an

earlier study (5). Primary LA FLS were stimulated with either *B. burgdorferi,* IFNγ, or *B.*

burgdorferi + IFNγ together, and expression of genes associated with antigen

processing and presentation was assessed (Figure 1B). As expected, expression of co-

stimulatory molecules B7-1 and B7-2 were not consistently expressed at levels above

zero (Figure 1B). However, other Signal 2 molecules were expressed by FLS. PD-L1

and CD40 were both strongly upregulated in response to IFNγ stimulation, expression of

ICOS-LG trended up in response to *B. burgdorferi* + IFNγ stimulation, and expression of

B7-H3 trended down with *B. burgdorferi* + IFNγ stimulation (Figure 1B). Expression of

genes associated with regulation of the MHC-II locus (e.g., CIITA), Class II antigen

loading (e.g., CD74/invariant chain, HLA-DM, and HLA-DO) and presentation to CD4+ T

cells (e.g., HLA-DR) was increased in FLS stimulated with IFNγ or IFNγ + *B. burgdorferi*.

Human LA FLS present MHC-II peptides derived from Lyme autoantigens

222   Post-infectious LA is frequently accompanied by autoimmune T and B cell

reactivity against certain HLA-DR-presented self-antigens, which were previously

identified in patients' synovial tissue and/or synovial fluid mononuclear cells (SFMCs)

using an unbiased immunopeptidomics approach (39). A subset of these Lyme

autoantigens is derived from extracellular matrix (ECM) proteins, which include

fibronectin 1 (FN1), laminin B2, collagen Vα1, and matrix metalloprotease 10 (1, 20).

Another subset is derived from proteins involved in vascular damage, including and

ECGF (21), annexin A2 (22), and apolipoprotein B-100 (19). We and others have

previously speculated that HLA-DR+ FLS may present some of these autoantigens

within inflamed synovial tissue, thereby contributing to tissue-localized CD4+ T cell

autoimmunity (1, 2, 5). Furthermore, we presume that autoimmunity is initiated during

infection with *B. burgdorferi*.

We used an *in vitro* immunopeptidomics approach using primary human LA FLS

to test whether activated HLA-DR+ FLS were able to present MHC-II antigens, including

potential Lyme autoantigens and foreign antigens derived from *B. burgdorferi*. FLS were

isolated from synovial tissue collected from an LA patient during the post-infectious

period (same patient as patient 1 shown in Figure 1A), whose HLA-DRB type is HLA-

DRB1*0701/DRB1*1454. Cells were passaged over six times prior to use to remove any

potential antigens retained from the time of initial sample collection. FLS were

stimulated with IFNγ and *B. burgdorferi* (10:1 MOI) for 48 hours, following which HLA-

DR peptides were isolated, purified, and identified by LC-MS/MS as described (39)

(Figure 2A).

Fixed Value (FV) validator analysis, with a minimum cross-correlation score

(Xcorr) of 2.5, was used to identify 695 total peptides (Supplemental Table S1),

including 10 peptides derived from *B. burgdorferi* proteins (Supplemental table S2). Of

the *B. burgdorferi* proteins, four are predicted to be localized to the inner membrane,

three are cytosolic, and two with unknown localization or function. Three source proteins

were involved in chemotaxis/motility, three in sugar transport, two in transcription and

translation. Surprisingly, none of these *B. burgdorferi* peptides were derived from

surface lipoproteins, which are immunodominant B cell antigens (40).

We also identified 685 peptides derived from 367 unique human proteins using

FV validation analysis (Supplemental Table S1). The most abundant source protein of

peptides identified was the Invariant chain, suggesting FLS are engaging in

endogenous formation and transport of MHC class II peptide complexes, and is

consistent with CD74 gene expression data from Figure 1B. Proteins associated with

the ECM were also abundantly represented in our dataset, with the top three KEGG

Pathway functional annotation terms being 1) focal adhesion; 2) alanine, aspartate, and

glutamate metabolism; and 3) ECM-receptor interactions. Interestingly, the peptide size

had a binomial distribution, with 433 (63%) of the peptides greater than 25 amino acids

long (median length=42), and 262 (38%) of the peptides 9-25 amino acids long (median

value=17), the latter being typical of fully processed MHC-II-bound peptides (41).

A second, more stringent Target Decoy (TD) validation analysis was also

performed, using both a 1% false discovery cutoff rate and an Xcorr cutoff of 2.5 to

minimize the potential number of false positive peptide-spectrum matches. Using this

highly stringent method, we identified 73 unique predicted peptides derived from 50

human proteins (Supplemental Table S1), nearly all of which were 13-18 amino acids

long, typical of fully processed MHC-II peptides (Figure 2B). Although all 73 peptides

were also identified using FV analysis, no *B. burgdorferi*-derived peptides were

identified using this stringent TD analysis method. Of the 73 human peptides identified,

14 peptides were predicted to bind to the peptide binding groove of DRB1*0701, and 46

peptides were predicted to bind to the peptide binding groove of DRB1*1454 (Figure

2B), matching the HLA-DRB type of LA Patient 1. WIKI Pathway, KEGG Pathway, and

Gene Ontology cellular localization analyses were used to further characterize source

proteins of identified HLA-DR peptides (Figure 2C). Peptides from our TD dataset were

enriched for proteins involved in ECM interactions, including focal adhesion, ECM-

receptor interactions, TGF-beta receptor signaling, and CCL18 signaling, which is

induced by collagen and fibronectin produced by FLS (42). Furthermore, source

proteins included 4 ECM, 11 secreted, 16 membrane-associated, and 21 cytosolic

proteins (Figure 2C).

Among the self-peptides identified by both FV and TD analysis were two derived

from LA autoantigens fibronectin (FN1) and endothelial cell growth factor (ECGF) (Table

1). The FN1 peptide identified, FN1$_{1996-2010}$, is predicted to bind to *0701; and the ECGF

peptide, ECGF$_{457-467}$, is predicted to bind to *1454. Interestingly, both the FN1$_{1996-2010}$

and the ECGF$_{457-467}$ epitopes from our screen are identical or nearly identical to HLA-

DR-bound peptides previously isolated from post-infectious LA synovial tissue (39), and

match several epitopes from the Immune Epitope Database (IEDB). Notably, the

FN1$_{1996-2010}$ peptide we identified is nearly identical to an autoreactive CD4+ T cell

epitope in a subset of patients with post-infectious LA (1), demonstrating that FLS are

able to present Lyme autoantigens.

IFNγ induces MHC-II expression in murine FLS

In C3H/HeJ and C57BL/6 mice infected with *B. burgdorferi*, FLS are major

sources of arthritogenic cytokines and chemokines in inflamed joints (33, 43).

Furthermore, *B. burgdorferi* PG, which persists in joints of LA patients for months to

several years following antibiotic therapy, is arthritogenic in BALB/c mice (24). We

hypothesized that in post-infectious LA, IFNγ produced by dysregulated T cells and

persistent *B. burgdorferi* PG may induce FLS into an antigen presenting cell phenotype,

driving CD4+ T cell activation in synovial tissue even after the infection itself is resolved

using antibiotic therapy (2).

Consistent with similar studies using primary human LA FLS (5), stimulation of

C57BL/6 FLS with IFNγ (20ng/mL) induced upregulation of MHC-II I-ab ~1.5-fold after

short-term stimulation (18h), and ~2-fold after long-term stimulation (72h) with IFNγ. In

comparison, long-term stimulation with both IFNγ and *B. burgdorferi* PG (10 µg/mL)

significantly induced upregulation of I-ab ~2.5-fold (Figure 3A). MHC-II I-ab expression

in IFNγ-stimulated BMDM was also significantly increased ~5-fold at 72 hours post-

stimulation, but PG did not affect I-ab expression (Figure 3A). Expression of FLS

activation marker Intercellular Adhesion Molecule 1 (ICAM-1), which is important for T

cell attachment to APCs during antigen presentation and formation of immune

synapses, was also upregulated in FLS stimulated with IFNγ or IFNγ+PG for 24 hours,

but not PG alone (Figure 3B). This confirms that IFNγ, but not PG, is required for

immune activation of FLS. When stimulated with IFNγ, FLS also upregulated production

of IL-6, CCL2, and TNFα, approximately 2-fold compared to unstimulated controls and

stimulation with *B. burgdorferi* PG alone (Supplemental Figure S1). Interestingly,

stimulation with both IFNγ and *B. burgdorferi* PG dampened production of IL-6 in FLS,

compared to IFNγ alone, but both stimuli were required for BMDM upregulation of IL-6.

Stimulation with *B. burgdorferi* PG alone did not affect the production of CCL2 or TNFα

in FLS or BMDM.

MHC-II expression is regulated by the γ-interferon-induced transcriptional

coactivator MHC class II transactivator (CIITA) (44). To determine whether IFNγ induces

MHC-II expression in FLS in an endogenous, CIITA-dependent manner, FLS from

CIITA-GFP reporter mice stimulated with IFNγ at various concentrations. FLS from

CIITA-GFP reporter mice had significantly increased fluorescence over controls,

particularly when stimulated with 20ng/mL IFNγ for 72-96 hours (Figure 3C).

Murine MHC-II+ FLS activate CD4+ T cells in an antigen-dependent manner

To evaluate whether MHC-II+ FLS can activate CD4+ T cells, we conducted a T

cell proliferation assay using activated MHC-II+ FLS co-cultured with CD4+ T cells from

mice that express a transgenic T cell receptor (TCR) specific for chicken ovalbumin

MHC-II antigen OVA$_{323-339}$ as a model system to study antigen-specific CD4+ T cell

responses (OT-II mouse model, Figure 4A). Co-culture of OT-II CD4+ T cells with FLS

primed with IFNγ and with the OT-II OVA$_{323-339}$ epitope had ~25% increased levels of

CD4+ T cell proliferation, compared to ~15% seen using unstimulated FLS (Figure 4B).

However, FLS primed with both IFNγ and *B. burgdorferi* PG induced significantly

increased (~37%) levels of CD4+ T cell proliferation, compared with the other two

conditions (Figure 4B). T cells co-cultured with IFNγ-primed FLS given the OVA$_{257-264}$

control peptide, an alternative MHC-II binding epitope not recognized by the OT-II

transgenic TCR, failed to induce T cell proliferation under any conditions (Figure 4B),

indicating that T cell proliferation was MHC-II epitope-dependent.

Priming of activated MHC-II+ FLS with PG had a pronounced adjuvant effect on

the ability of FLS to induce multiple rounds of T cell proliferation. An average of 30.6%

of CD4+ T cells underwent two or more rounds of proliferation when co-cultured with

IFNγ+PG-activated MHC-II+ FLS, compared to a significantly lower average of 17.3% of

CD4+ T cells that underwent two or more rounds of proliferation when cocultured with

IFNγ-activated MHC-II+ FLS without PG (Figure 4B). Similar results were observed

when full-length ovalbumin was added to the coculture medium (Figure 4C), indicating

that FLS are able to take up, process, and load MHC-II antigen.

To characterize the CD4+ T cell responses induced by activated MHC-II+ FLS,

we measured cytokines in supernatants collected from FLS-T cell co-culture

experiments using the OT-II OVA$_{323-339}$ epitope as antigen (Figure 4D). Supernatants

from T cells co-cultured with FLS primed with IFNγ alone had an average of 15.4 pg/mL

of IL-10, which was significantly higher than unstimulated FLS (6.4 pg/mL), or FLS

primed with IFNγ+PG (6.2 pg/mL). In contrast, supernatants from T cells co-cultured

with FLS primed with both IFNγ and PG had elevated levels of IFNγ (369.5 pg/mL),

compared with unprimed FLS (69.2 pg/mL) or FLS primed with IFNγ alone (143.1

pg/mL). As a result, the IFNγ/IL-10 ratio was ~10-fold higher in supernatants from T

cells co-cultured with FLS primed with IFNγ+PG, compared with the other two groups,

indicating a skew towards a T_H1 effector phenotype. FLS primed with either IFNγ or

IFNγ+PG induced production of similarly elevated levels of IL-17A in supernatants of co-

culture experiments, compared with unprimed FLS (Figure 4D), suggesting that addition

of PG to FLS did not enhance IL-17 production.

DISCUSSION

This study shows that activated MHC-II+ FLS are able to process and present

MHC-II antigens, including antigens associated with LA autoimmunity, and can induce

CD4+ T cell activation in an antigen-dependent manner, particularly when primed with

PG and IFNγ, which partially recapitulates elements of the inflammatory

microenvironment within the post-infectious LA synovial lesion. These data also support

earlier speculation that MHC-II+ FLS are likely major contributors of HLA-DR-presented

peptides derived from ECM proteins that accumulate within the post-infectious LA

inflammatory lesion (1, 2, 5). We propose that under pathogenic conditions, localized

IFNγ responses during *B. burgdorferi* infection stimulate FLS into an activated MHC-II+

immune effector phenotype. If IFNγ levels within the joint microenvironment become

excessive due to autoimmunity and/or accumulation of *B. burgdorferi* PG, activated

MHC-II+ FLS can prolong pathogenic T cell reactivity, perpetuating tissue-localized

inflammation even in the absence of active infection.

We report that IFNγ induces murine FLS to upregulate numerous genes involved

in uptake, processing, and presentation of MHC-II antigen. By utilizing an

immunopeptidomics approach, we characterized peptides presented by human LA

MHC-II+ FLS and found that these cells are able to present peptides derived from LA

autoantigen FN1, which raises the possibility that these activated MHC-II+ FLS may

drive ECM autoimmunity following *B. burgdorferi* infection. We also showed that the

adjuvant effects of *B. burgdorferi* PG can fundamentally alter T cell responses induced

by MHC-II+ FLS. We show that MHC-II+ FLS, induced by IFNγ, can take up antigen

from the environment and induce T cell activation in an antigen-dependent manner,

which is exacerbated by *B. burgdorferi* PG to a robust T_H1-like response. Yet, in the

absence of PG, T cell proliferation is attenuated and is accompanied by elevated

production of the immunomodulatory cytokine IL-10. Overall, these findings support

observations in previous studies of post-infectious LA (5, 24, 45), validating our *in vitro*

model of LA. Together, these data show that activated MHC-II+ FLS function as

inducible APCs to promote ongoing recruitment and activation of IFNγ-producing

lymphocytes.

393   Other studies have demonstrated that tissue-localized antigen presentation in the

joint is fundamental for driving T cell-mediated autoimmune inflammation against self-

peptides derived from proteins at sites of infection associated with synovial tissue

damage (1, 19-22). Identification of FN1 as an FLS-presented peptide was particularly

noteworthy, since autoreactive CD4+ T cells targeting FN1 and other FLS-derived ECM

proteins have a marked T_H1 phenotype, and are detected in over 50% of post-infectious

LA patients (1). Our study suggests that retained *B. burgdorferi* PG plays a critical role

in perpetuating T_H1 reactivity against autoreactive ECM proteins in post-infectious LA

(24).

402   Our data indicate that activated MHC-II+ FLS phenotypes are plastic, and can be

dramatically altered by presence or absence of PG. This finding may have broad

implication in other chronic inflammatory joint diseases, particularly when an infection is

suspected as an autoimmune trigger. For example, previous reports have identified PG

in inflamed synovia from patients with RA (46) and from patients undergoing total knee

arthroplasty (26). This idea merits a reexamination of the role of PG and other PAMPS

in RA autoimmunity. Our study also has potential clinical and therapeutic implications. A

recent study showed that part of the therapeutic effect of JAK inhibitors is reversing the

pro-inflammatory effects of activated MHC-II+ FLS by interfering with IFNγ signaling

(11).

There are some limitations with our study. While we were able to isolate HLA-

DR+ FLS-presented peptides from one patient after *in vitro* stimulation, we are unable to

obtain sufficient numbers of FLS directly from patients' tissue to determine what

peptides are presented by FLS *ex vivo*. Furthermore, we and others (38) have shown

that FLS lack expression of canonical CD80 and CD86 costimulatory molecules. Thus,

the source of signal 2 for CD4+ T cell activation is still unknown. As shown in this and

other studies, FLS express an altered repertoire of costimulatory molecules such as

CD40, ICOS-L, PD-L1, B7-H3 (38, 47), which may provide both pro-inflammatory and

anti-inflammatory signals to T cells, depending on the FLS activation state. Further

analysis of Signal 2 will be needed to understand the nature of FLS-mediated T cell

activation more fully.

In conclusion, this study demonstrates that activated MHC-II+ FLS can present

HLA-DR-presented Lyme autoantigens and are inducible APCs that acquire a potent T

cell-activating phenotype, particularly when exposed to IFNγ and PG. These findings

add to a growing body of evidence that MHC-II+ stromal cells play critical roles in

shaping tissue-localized autoimmune CD4+ T cell responses in disease pathogenesis.

For example, pancreatic cancer-associated fibroblasts induce activation of autoimmune,

pro-tumorigenic regulatory T cell responses in an MHC-II antigen-dependent manner

(48). Additionally, IFNγ-activated cardiac fibroblasts are capable of becoming inducible

APCs that promote cardiac fibrosis and dysfunction (49). Thus, MHC-II expressing

fibroblasts can have profound effects on human health and disease.

ACKNOWLEDGMENTS

We extend our sincerest gratitude to Catherine Costello, PhD, for her invaluable

assistance with the immunopeptidomics study. We would also like to acknowledge Mark

Wooten, PhD, for providing the CIITA EGFP-reporter mouse and for reviewing the

manuscript; Savannah Neu at the Versiti Flow Cytometry Core for flow cytometry

assistance; John Corbett, PhD, Director of the MCW Center for Biomedical Mass

Spectrometry Research for providing us with mass spectrometry resources and

expertise; and Michael Dwinell, PhD, Director of the MCW Center for Immunology for

providing equipment and scientific resources. We would also like to thank Jenifer

Coburn, PhD, for generously providing the *B. burgdorferi* strains used in this study.

FUNDING SOURCES

RBL and BLJ were supported by awards from the National Institutes of Health

(1R21AI148982-01, 1R01AI173256-01 and 1R01AI178711-01). RBL and BLJ were also

supported by the Assistant Secretary of Defense for Health Affairs through the Tick-

Borne Disease Research Program, endorsed by the Department of Defense under

Award No. W81XWH-22-1- 0728. Opinions, interpretations, conclusions, and

recommendations are those of the author and are not necessarily endorsed by the

National Institutes of Health or the Department of Defense.

TABLES

Table 1. HLA-DR presented peptides derived from LA autoantigens isolated from

LA FLS.

Superscripts indicate position of the peptide in the parent protein sequence. Underlined

amino acids indicate the sequence corresponding to the HLA-DR binding groove. Bold

numbers indicate predicted HLA-DR binding alleles corresponding to the HLA type of

LA patient 1. IEDB (Immune Epitope Database) epitope ID numbers were retrieved from

iedb.org. Abbreviations: FN1 (fibronectin 1), ECGF (endothelial cell growth factor, also

called TYMP, thymidine phosphorylase).

Protein	Sequence	Source	Predicted HLA-DR binding (NetMHCIIpan4.0)
FN1	[1996]APSNLRFLATTPNSL[2010]	This study	HLA-DRB1*0101, *0102, *0103, *0401, *0403, *0404, *0405, *0408, ***0701**, *0803, *0901, *1001, *1601, DRB3*0202
FN1	[1996]APSNLRFLATTPNSL[2010]	Wang et al. 2017	
FN1	[1996]APSNLRFLATTPNSLLVSW[2014]	Kanjana et al. 2023	
FN1	[1996]APSNLRFLATTPNSL[2010]	IEDB (ID 144206)	
ECGF	[457]EALVLSDRAPF[467]	This study	HLA-DRB1*0301, *0305, *1104, *1301, *1302, *1303, *1401, *1402, ***1454**
ECGF	[462]SDRAPFAAPSPFAE[475]	Wang et al. 2017	
ECGF	[455]LQEALVLSDRAPF[467]	IEDB (ID 1253447)	
ECGF	[458]ALVLSDRAPFAAPSPFAE[467]	IEDB (ID 758118)	

465 **FIGURES AND FIGURE LEGENDS**

Figure 1. FLS expression of MHC class II and costimulatory molecules is induced

by IFNɣ

Characterization of synovial and tendon tissue samples from four different LA patients.

Representative images of co-localization of HLA-DR and vimentin in two LA synovial

and two tendon tissue biopsies (A). Bulk RNA sequencing analysis of one LA patient's

(Patient 1) FLS stimulated *ex vivo* with IFNɣ, *Borrelia burgdorferi (Bb)*, or both (B).

Statistical analysis was performed using Tukey's multiple comparisons test (p values

indicated in figure). Error bars represent standard errors from mean. CPMR: copies per

million reads.

Figure 2. Human LA FLS present MHC-II peptides derived from Lyme

autoantigens Immunopeptidomics

screen of unique LA patient FLS HLA-DR presented peptides (A). Peptide length and

predicted HLA binding motifs of peptides predicted by both Fixed Value (FV) and Target

Decoy (TD) analyses corresponding with LA patient 1 HLA-DR type (B). WIKI

Pathways, KEGG Pathways, and Gene Ontology cellular localization analyses showing

enriched pathways and cellular localization associated with source proteins (C).

Figure 3. Murine FLS upregulate MHC-II in response to IFNɣ

Surface expression of MHC-II by primary B6 murine FLS or BMDMs stimulated with *B. burgdorferi* PG (10 µg/ml), IFNɣ (20 ng/ ml), or both for 18 or 72 hours (A). A separate experiment demonstrating expression of ICAM by FLS in response to IFNɣ and *B. burgdorferi* PG stimulation for 24 hours (B). Data pooled from two independent experiments with three technical replicates (A-B). CIITA EGFP-reporter mouse FLS (passage 4) were stimulated with 0, 0.2, 2, or 20 ng/mL IFNɣ and compared to non-EGFP expressing control B6 FLS. Fluorescence was analyzed by IncuCyte imaging with images taken every 2 hours for 96 hours total at 10X objective (C). All statistical

497 analysis performed using Tukey's multiple comparisons test (p<0.05). Error bars

498 represent standard deviation.

Figure 4. Murine activated MHC-II+ FLS induce CD4+ T cell activation in an

antigen-dependent manner

MHC-II+ FLS were induced with either IFNɣ alone or IFNɣ and *B. burgdorferi* PG for 48

hours and washed prior to isolation and coculture of OT-II mouse T cells for 5 days and

subsequent analysis by flow cytometry (A). Percentage of proliferating CD4+ T cells and

cells that completed two or more rounds of replication, induced by MHC-II+ FLS when

loaded with ovalbumin peptides (B). Percentage of proliferating CD4+ T cells induced by

MHC-II+ FLS when loaded with full-length ovalbumin (C). Co-culture cytokine analysis

of IFNɣ, IL-10, and IL-17a production (D). All data were pooled from two independent

experiments with three technical replicates and statistical analysis was performed using

Tukey's multiple comparisons test (p<0.05). Error bars represent standard deviation.

Supplemental Figure S1. Murine MHC-II+ FLS upregulate proinflammatory mediators in response to IFNɣ

Production of IL-6, TNFα, and CCL2 by primary B6 murine FLS (A) and BMDM (B) stimulated with *B. burgdorferi* PG (10 µg/ml), IFNɣ (20 ng/ ml), or both for ~24 hrs. All data pooled from two independent experiments and statistical analysis was performed using Tukey's multiple comparisons test (p<0.05). Error bars represent standard deviation.

<u>References</u>

1.   Kanjana K, Strle K, Lochhead RB, Pianta A, Mateyka LM, Wang Q, et al. Autoimmunity to

synovial extracellular matrix proteins in patients with post-infectious lyme arthritis. J Clin

Invest. 2023.

2.   Lochhead RB, Strle K, Arvikar SL, Weis JJ, Steere AC. Lyme arthritis: linking infection,

inflammation and autoimmunity. Nat Rev Rheumatol. 2021;17(8):449-61.

3.   Nygaard G, Firestein GS. Restoring synovial homeostasis in rheumatoid arthritis by

targeting fibroblast-like synoviocytes. Nat Rev Rheumatol. 2020;16(6):316-33.

4.   Wei K, Nguyen HN, Brenner MB. Fibroblast pathology in inflammatory diseases. J Clin

Invest. 2021;131(20).

5.   Lochhead RB, Ordonez D, Arvikar SL, Aversa JM, Oh LS, Heyworth B, et al. Interferon-

gamma production in Lyme arthritis synovial tissue promotes differentiation of fibroblast-like

synoviocytes into immune effector cells. Cell Microbiol. 2019;21(2):e12992.

6.   Plikus MV, Wang X, Sinha S, Forte E, Thompson SM, Herzog EL, et al. Fibroblasts: Origins,

definitions, and functions in health and disease. Cell. 2021;184(15):3852-72.

7.   Carmona-Rivera C, Carlucci PM, Moore E, Lingampalli N, Uchtenhagen H, James E, et al.

Synovial fibroblast-neutrophil interactions promote pathogenic adaptive immunity in

rheumatoid arthritis. Sci Immunol. 2017;2(10).

8.   Croft AP, Campos J, Jansen K, Turner JD, Marshall J, Attar M, et al. Distinct fibroblast

subsets drive inflammation and damage in arthritis. Nature. 2019;570(7760):246-51.

9.   Mizoguchi F, Slowikowski K, Wei K, Marshall JL, Rao DA, Chang SK, et al. Functionally

distinct disease-associated fibroblast subsets in rheumatoid arthritis. Nat Commun.

2018;9(1):789.

10.   Tran CN, Davis MJ, Tesmer LA, Endres JL, Motyl CD, Smuda C, et al. Presentation of

arthritogenic peptide to antigen-specific T cells by fibroblast-like synoviocytes. Arthritis Rheum.

2007;56(5):1497-506.

11.   Zhao S, Grieshaber-Bouyer R, Rao DA, Kolb P, Chen H, Andreeva I, et al. Effect of JAK

Inhibition on the Induction of Proinflammatory HLA-DR+CD90+ Rheumatoid Arthritis Synovial

Fibroblasts by Interferon-gamma. Arthritis Rheumatol. 2022;74(3):441-52.

12.   Alivernini S, MacDonald L, Elmesmari A, Finlay S, Tolusso B, Gigante MR, et al. Distinct

synovial tissue macrophage subsets regulate inflammation and remission in rheumatoid

arthritis. Nat Med. 2020;26(8):1295-306.

13.   Dong Y, Zhou G, Cao W, Xu X, Zhang Y, Ji Z, et al. Global seroprevalence and

sociodemographic characteristics of Borrelia burgdorferi sensu lato in human populations: a

systematic review and meta-analysis. BMJ Glob Health. 2022;7(6).

14.   Kugeler KJ, Schwartz AM, Delorey MJ, Mead PS, Hinckley AF. Estimating the Frequency

of Lyme Disease Diagnoses, United States, 2010-2018. Emerg Infect Dis. 2021;27(2):616-9.

15.   Arvikar SL, Steere AC. Lyme Arthritis. Infect Dis Clin North Am. 2022;36(3):563-77.

16.   Snydman DR, Schenkein DP, Berardi VP, Lastavica CC, Pariser KM. Borrelia burgdorferi in

joint fluid in chronic Lyme arthritis. Ann Intern Med. 1986;104(6):798-800.

17.   Steere AC. Lyme disease. N Engl J Med. 1989;321(9):586-96.

18.   Steere AC, Schoen RT, Taylor E. The clinical evolution of Lyme arthritis. Ann Intern Med.

1987;107(5):725-31.

19.   Crowley JT, Drouin EE, Pianta A, Strle K, Wang Q, Costello CE, et al. A Highly Expressed

Human Protein, Apolipoprotein B-100, Serves as an Autoantigen in a Subgroup of Patients With

Lyme Disease. J Infect Dis. 2015;212(11):1841-50.

20.   Crowley JT, Strle K, Drouin EE, Pianta A, Arvikar SL, Wang Q, et al. Matrix

metalloproteinase-10 is a target of T and B cell responses that correlate with synovial pathology

in patients with antibiotic-refractory Lyme arthritis. J Autoimmun. 2016;69:24-37.

21.   Drouin EE, Seward RJ, Strle K, McHugh G, Katchar K, Londono D, et al. A novel human

autoantigen, endothelial cell growth factor, is a target of T and B cell responses in patients with

Lyme disease. Arthritis Rheum. 2013;65(1):186-96.

22.   Pianta A, Drouin EE, Crowley JT, Arvikar S, Strle K, Costello CE, et al. Annexin A2 is a

target of autoimmune T and B cell responses associated with synovial fibroblast proliferation in

patients with antibiotic-refractory Lyme arthritis. Clin Immunol. 2015;160(2):336-41.

23.   Bockenstedt LK, Gonzalez DG, Haberman AM, Belperron AA. Spirochete antigens persist

near cartilage after murine Lyme borreliosis therapy. J Clin Invest. 2012;122(7):2652-60.

24.   Jutras BL, Lochhead RB, Kloos ZA, Biboy J, Strle K, Booth CJ, et al. Borrelia burgdorferi

peptidoglycan is a persistent antigen in patients with Lyme arthritis. Proc Natl Acad Sci U S A.

2019;116(27):13498-507.

25.   Schrijver IA, Melief MJ, Markusse HM, Van Aelst I, Opdenakker G, Hazenberg MP, et al.

Peptidoglycan from sterile human spleen induces T-cell proliferation and inflammatory

mediators in rheumatoid arthritis patients and healthy subjects. Rheumatology (Oxford).

2001;40(4):438-46.

26.   Holub MN, Wahhab A, Rouse JR, Danner R, McClune MM, Dressler JM, et al.

Peptidoglycan in osteoarthritis synovial tissue is associated with joint inflammation. Res Sq.

2023.

27.   Chiu YC, Lin CY, Chen CP, Huang KC, Tong KM, Tzeng CY, et al. Peptidoglycan enhances

IL-6 production in human synovial fibroblasts via TLR2 receptor, focal adhesion kinase, Akt, and

AP-1- dependent pathway. J Immunol. 2009;183(4):2785-92.

28.   Kyburz D, Rethage J, Seibl R, Lauener R, Gay RE, Carson DA, et al. Bacterial

peptidoglycans but not CpG oligodeoxynucleotides activate synovial fibroblasts by toll-like

receptor signaling. Arthritis Rheum. 2003;48(3):642-50.

29.   Centers for Disease C, Prevention. Recommendations for test performance and

interpretation from the Second National Conference on Serologic Diagnosis of Lyme Disease.

MMWR Morb Mortal Wkly Rep. 1995;44(31):590-1.

30.   Lantos PM, Rumbaugh J, Bockenstedt LK, Falck-Ytter YT, Aguero-Rosenfeld ME,

Auwaerter PG, et al. Clinical Practice Guidelines by the Infectious Diseases Society of America

(IDSA), American Academy of Neurology (AAN), and American College of Rheumatology (ACR):

2020 Guidelines for the Prevention, Diagnosis and Treatment of Lyme Disease. Clin Infect Dis.

2021;72(1):1-8.

31.   Seward RJ, Drouin EE, Steere AC, Costello CE. Peptides presented by HLA-DR molecules

in synovia of patients with rheumatoid arthritis or antibiotic-refractory Lyme arthritis. Mol Cell

Proteomics. 2011;10(3):M110 002477.

32.   Boes M, Cerny J, Massol R, Op den Brouw M, Kirchhausen T, Chen J, et al. T-cell

engagement of dendritic cells rapidly rearranges MHC class II transport. Nature.

2002;418(6901):983-8.

33.   Lochhead RB, Sonderegger FL, Ma Y, Brewster JE, Cornwall D, Maylor-Hagen H, et al.

Endothelial cells and fibroblasts amplify the arthritogenic type I IFN response in murine Lyme

disease and are major sources of chemokines in Borrelia burgdorferi-infected joint tissue. J

Immunol. 2012;189(5):2488-501.

34.   Davis MM, Brock AM, DeHart TG, Boribong BP, Lee K, McClune ME, et al. The

peptidoglycan-associated protein NapA plays an important role in the envelope integrity and in

the pathogenesis of the lyme disease spirochete. PLoS Pathog. 2021;17(5):e1009546.

35.   Haupl T, Hahn G, Rittig M, Krause A, Schoerner C, Schonherr U, et al. Persistence of

Borrelia burgdorferi in ligamentous tissue from a patient with chronic Lyme borreliosis. Arthritis

Rheum. 1993;36(11):1621-6.

36.   Steere AC. Musculoskeletal manifestations of Lyme disease. Am J Med.

1995;98(4A):44S-8S; discussion 8S-51S.

37.   Miller JB, Albayda J, Aucott JN. The Value of Musculoskeletal Ultrasound for Evaluation

of Postinfectious Lyme Arthritis. J Clin Rheumatol. 2022;28(2):e605-e8.

38.   Tran CN, Thacker SG, Louie DM, Oliver J, White PT, Endres JL, et al. Interactions of T cells

with fibroblast-like synoviocytes: role of the B7 family costimulatory ligand B7-H3. J Immunol.

2008;180(5):2989-98.

39.   Wang Q, Drouin EE, Yao C, Zhang J, Huang Y, Leon DR, et al. Immunogenic HLA-DR-

Presented Self-Peptides Identified Directly from Clinical Samples of Synovial Tissue, Synovial

Fluid, or Peripheral Blood in Patients with Rheumatoid Arthritis or Lyme Arthritis. J Proteome

Res. 2017;16(1):122-36.

40.   Barbour AG, Jasinskas A, Kayala MA, Davies DH, Steere AC, Baldi P, et al. A genome-wide

proteome array reveals a limited set of immunogens in natural infections of humans and white-

footed mice with Borrelia burgdorferi. Infect Immun. 2008;76(8):3374-89.

41.   Pishesha N, Harmand TJ, Ploegh HL. A guide to antigen processing and presentation. Nat

Rev Immunol. 2022;22(12):751-64.

42.   Schraufstatter IU, Zhao M, Khaldoyanidi SK, Discipio RG. The chemokine CCL18 causes

maturation of cultured monocytes to macrophages in the M2 spectrum. Immunology.

2012;135(4):287-98.

43.   Helble JD, Walsh MJ, McCarthy JE, Smith NP, Tirard AJ, Arnold BY, et al. Single-cell RNA

sequencing of murine ankle joints over time reveals distinct transcriptional changes following

Borrelia burgdorferi infection. iScience. 2023;26(11):108217.

44.   Mach B, Steimle V, Martinez-Soria E, Reith W. Regulation of MHC class II genes: lessons

from a disease. Annu Rev Immunol. 1996;14:301-31.

45.   Lochhead RB, Arvikar SL, Aversa JM, Sadreyev RI, Strle K, Steere AC. Robust interferon

signature and suppressed tissue repair gene expression in synovial tissue from patients with

postinfectious, Borrelia burgdorferi-induced Lyme arthritis. Cell Microbiol. 2019;21(2):e12954.

46.   van der Heijden IM, Wilbrink B, Tchetverikov I, Schrijver IA, Schouls LM, Hazenberg MP,

et al. Presence of bacterial DNA and bacterial peptidoglycans in joints of patients with

rheumatoid arthritis and other arthritides. Arthritis Rheum. 2000;43(3):593-8.

47.   Sempowski GD, Chess PR, Phipps RP. CD40 is a functional activation antigen and B7-

independent T cell costimulatory molecule on normal human lung fibroblasts. J Immunol.

1997;158(10):4670-7.

48.   Huang H, Wang Z, Zhang Y, Pradhan RN, Ganguly D, Chandra R, et al. Mesothelial cell-

derived antigen-presenting cancer-associated fibroblasts induce expansion of regulatory T cells

in pancreatic cancer. Cancer Cell. 2022;40(6):656-73 e7.

49.   Ngwenyama N, Kaur K, Bugg D, Theall B, Aronovitz M, Berland R, et al. Antigen

presentation by cardiac fibroblasts promotes cardiac dysfunction. Nat Cardiovasc Res.

2022;1(8):761-74.