Microbiome-based correction of nutrient profiles derived from self-reported dietary assessments

Tong Wang[1], Yuanqing Fu[2,3,4], Menglei Shuai[2,3], Ju-Sheng Zheng[2,3,4], Lu Zhu[5], Qi Sun[1,6,7], Frank B. Hu[1,6,7], Scott T. Weiss[1], Yang-Yu Liu[1,8,*]

[1]*Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02115, USA*
[2]*Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, China*
[3]*Westlake Intelligent Biomarker Discovery Lab, Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, China*
[4]*Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, China*
[5]*Department of Epidemiology, University of Iowa College of Public Health, Iowa City, IA 52242, USA*
[6]*Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA*
[7]*Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA*
[8]*Center for Artificial Intelligence and Modeling, The Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA*
* Correspondence: yyl@channing.harvard.edu

Abstract

Since dietary intake is hard to directly measure in large-scale cohort studies, we often rely on self-reported instruments (e.g., food frequency questionnaires, diet recall surveys, and diet diary methods) developed in nutritional epidemiology. Those self-reported instruments are prone to measurement errors. The measurement errors eventually lead to inaccuracies in the calculation of nutrient profiles. Currently, there is a lack of computational methods to address this problem. To fill the gap, we introduce a deep-learning approach --- **M**icrobiom**e**-based nu**t**rient p**r**of**i**le **c**orrector (METRIC), which leverages gut microbial compositions to correct the errors in nutrient profiles due to measurement errors in self-reported dietary assessments. We demonstrate the excellent performance of METRIC in minimizing the simulated random errors in both synthetic and three real-world datasets. METRIC has the potential to significantly improve the accuracy of self-reported dietary assessments and hence facilitate the research of nutritional epidemiology.

Introduction

An unhealthy diet can increase the risk of many diseases[1,2,3]. For instance, consuming too much sugar or saturated fat could elevate the risk of coronary heart disease[4–6]. The investigation of the association between poor dietary habits and chronic diseases requires an accurate assessment of dietary intake in large population samples. Usually, we rely on self-reported instruments such as food frequency questionnaires (FFQ)[7], ASA24 (Automated Self-Administered 24-hour Dietary Assessment Tool)[8], and 7-Day Dietary Record (7DDR)[9], to collect diet data in these studies. However, self-reported instruments are subject to random measurement errors[10–13]. In particular, recall methods (e.g., FFQ and ASA24) are prone to recall bias[14] and individuals' intentional misreporting of certain food consumptions[15]. Such measurement errors in the assessed food intake will be naturally propagated forward to the computation of the nutrient profile, causing inaccuracies in the assessed nutrient profile (Fig. 1a). Correcting such errors in the nutrient profile is crucial for improving the quality of nutritional epidemiology research. Yet, we still lack a computational method to achieve that.

Signal reconstruction from corrupted or incomplete measurements is a common problem in various fields of research. For example, image denoising (i.e., removing noise from a noisy image to restore the true image) is an important topic in computer vision. The recent advance in deep neural networks has led to new methods that learn to map corrupted images to unobserved clean ones[16,17]. However, training those methods typically requires clean images, which are difficult to obtain in many cases. To overcome this challenge, a new class of methods that only leverages noisy images in training has been developed[18,19]. One noticeable example is Noise2Noise[18], which restores clean images by only being trained on corrupted ones. When the noise is random with a mean of zero, its performance is comparable and sometimes even better than other methods trained using clean images[18].

Inspired by the success of Noise2Noise[18], we hypothesize that we can correct the error in the assessed nutrient profile derived from the self-reported dietary assessment without using clean data (i.e., the ground truth dietary intake). We focused on correcting the nutrient profile instead of the food profile (or the original dietary assessment) because the prevalence of zero values in the food profile (i.e., no consumption for many food items) renders a big challenge for machine learning tasks, while the derived nutrient profile usually has non-zero values.

Our key idea is to incorporate gut microbial compositions as part of the input of a deep-learning model and to infer the true nutrient profile based on the assessed nutrient profile and the measured gut microbial composition (Fig. 1b). This idea is based on the fact that many

dietary components reaching the large intestine fuel the growth of gut microbes[20–22]. One
example is the growth of gut microbes enabled by anaerobic fermentation of indigestible
polysaccharides[20,21]. As a result, the gut microbial composition is linked to dietary intake.
Advances in sequencing technologies have made it possible to determine the gut microbial
composition quickly and accurately. It has been shown that fecal bacteria and metabolites can
be used as biomarkers for predicting the presence of food items ingested by healthy adults[23,24].
Hence, using objective microbial biomarkers to infer dietary intake might complement the self-
reported dietary assessment and thus reduce measurement error in dietary intake.

In this work, we developed a deep-learning method: **M**icrobiom**e**-based nu**t**rient p**r**of**i**le
corrector (METRIC). We demonstrated the effectiveness of METRIC in removing randomly
added noise to both synthetic and real data.

Results

Overview of METRIC

We aim to infer the true nutrient profile based on the assessed one and the measured gut
microbial composition. A naive way to do this is to train a machine-learning model with the
assessed nutrient profiles and microbial compositions as the input and the true nutrient profiles
as the output. However, this is not feasible because such training requires the true nutrient
profiles that are not easily available. To address this issue, we developed METRIC that does
not rely on the true nutrient profile during its training but still can remove measurement errors
in the assessed nutrient profile during testing (Fig. 2). During the training, we generated the
corrupted nutrient profiles by adding random noise to the assessed nutrient profiles, and then
trained METRIC to remove the added noise by taking the corrupted nutrient profiles and the
measured microbial compositions as its input and generating the assessed nutrient profiles as
its output (Fig. 2a). We introduced the corrupted nutrient profiles to avoid METRIC copying
the assessed nutrient profiles directly to the output, thus forcing METRIC to remove the added
noise. Note that the training of METRIC did not use true nutrient profiles.

METRIC comprises stacked neural networks with three hidden layers (Fig. 2b). One
unique feature is the skip connection that adds the corrupted nutrient profiles directly to the
final output of neural networks, which previously has been shown to enhance the training and
predictive performance for deep neural networks[25]. Similar to the existence of the skip
connection in Noise2Noise[18], the introduction of the skip connection enables the neural
networks to adjust the prediction based on the corrupted nutrient profile, ensuring that output

variables would not deviate too much from the corrupted nutrient profile. More details about
the architecture can be found in the Methods section. Overall, METRIC is a generic "denoiser"
that learns to remove any random noise added to the nutrient profile. With this generic ability
to remove random noise, the well-trained METRIC should be able to remove random
measurement errors by generating predictions closer to the true nutrient profiles when it takes
the assessed nutrient profiles and microbial compositions as its input (Fig. 2c).

We split each dataset into two non-overlapping parts: a training and a test set. METRIC
was trained on the training set and then used to generate predictions for the test set. The
correction performance is measured by comparing the predicted nutrient profiles with the true
nutrient profiles in the test set (how we obtain the true nutrient profiles for each scenario will
be explained in separate sections). To measure the predictive performance of each nutrient,
we adopted the Spearman Correlation Coefficient (SCC) ρ between its predicted and true
corrected values.

METRIC can reduce measurement errors in nutrient profiles of synthetic data

We first validated METRIC using synthetic data for which we know the ground truth. We used
the Microbial Consumer-Resource Model (MiCRM)[26] to generate three types of data: (1) *true*
nutrient profiles (i.e., the ground-truth nutrient consumption), (2) *assessed* nutrient profiles (i.e.,
the true nutrient profiles with random noise added to mimic measurement errors), and (3)
corrupted nutrient profiles (i.e., the assessed nutrient profiles with artificially added random
noise). MiCRM simulates the process of nutrient consumption by microbes and the following
microbial growth[26]. Note that the random noise added to assessed and corrupted nutrient
profiles are different. For simplicity, we only considered the nutrient consumption in MiCRM
and did not model the nutrient production because most dietary nutrients cannot be produced
by microbes. We created different samples by randomly sampling nutrient fluxes and then
running the community assembly until we achieved a steady state. We consider sampled
ground-truth nutrient fluxes as true nutrient profiles and the steady-state microbial abundances
as microbial compositions. Gaussian noise $N(0, \sigma^2)$ with the mean of zero and standard
deviation σ is added to true nutrient profiles to create assessed nutrient profiles. More details
about MiCRM, the generation of synthetic data, and added noise can be found in the Methods
section.

We generated 250 samples by simulating the assembly process for 250 independent
communities with 20 nutrients and 20 bacterial species. We trained METRIC on 200 samples

and tested it on the remaining 50 samples. We measured how assessed values and corrected

values (i.e., predicted values on the test set) respectively correlate with true values. For each

nutrient, we measured ρ between its corrected values from predictions and its true

concentration (denoted as ρ_c). Similarly, we calculated ρ between its assessed concentrations

and its true concentration (denoted as ρ_a). As the standard deviation σ of the Gaussian noise

increases, METRIC starts to correct the introduced noise, represented by the switch from

negative values of $(\rho_c - \rho_a)$ to positive values in Fig. 3a. It is natural to expect that the

correction of nutrient profiles with weaker noises does not work ($\rho_c < \rho_a$ for small σ in Fig. 3a),

because the correction is not necessary when the measurement error is small (e.g., when $\rho_a >$

0.8). We focused on the case of $\sigma = 1.5$ from now on. The difference between the two metrics

(i.e., $\rho_c - \rho_a$) reflects the correction performance. The nutrient in Figs. 3d&e is slightly

corrected ($\rho_c - \rho_a = 0.02$), while the nutrient in Figs. 3f&g is strongly corrected ($\rho_c - \rho_a = 0.10$).

The majority of nutrients have a better alignment between their corrected values and true

values (Fig. 3e).

METRIC mitigates measurement errors added to nutrient profiles in three real-

world datasets

Next, we tested METRIC on three real-world datasets. The first real dataset MCTs

(MiCrobiome dieT study) comes from a previous study that investigated the influence of diets

on gut microbial composition[27]. It is unique because it has a large number of samples (n=210)

of paired nutrient profiles and microbial compositions. The nutrient profiles were calculated

from ASA24 (see Methods). Since the true nutrient profiles are not available, we treated the

nutrient profiles derived from ASA24 as the "true" nutrient profiles and added random noise

(Gaussian noise with the mean of zero and standard deviations σ) to them as "assessed"

nutrient profiles. As σ increases, METRIC starts to better correct the introduced noise (Fig. 4a).

We focused on the case of $\sigma = 1.0$ from now on. We found that carotene has a large ρ_a and is

barely improved by METRIC ($\rho_a = 0.91$ versus $\rho_c = 0.91$ in Figs. 4b-c). By contrast, fiber has

a small ρ_a and is strongly improved ($\rho_a = 0.36$ versus $\rho_c = 0.63$ for Figs. 4f-g). We believe a

large correction in the total fiber intake because most fibers are digested by gut microbes[20,21].

Overall, nutrients with smaller ρ_a have better correction performance, and nutrients with large

ρ_a rarely improved (Fig. 4e). The mean correction performance averaged over all nutrients

$(\overline{\rho_c} - \overline{\rho_a}) = 0.069$.

We also tried to run METRIC without including the microbial composition, finding that

the correction performance ($\overline{\rho_c} - \overline{\rho_a} = 0.060$) is worse than that with included microbial

composition (Supplementary Fig. 1). More specifically, ρ_c of fiber decreases from 0.63 (Fig. 4e)

to 0.52 (Supplementary Fig. 1f). We also computed the sensitivity which is defined as the ratio

between the reduction in ρ of nutrient α and the perturbation amount of species i

(Supplementary Fig. 2), which could detect some possible interactions between nutrients and

species. For example, the sensitivity of riboflavin towards *Faecalibacterium prausnitzii* is

relatively large (~0.0074), supported by the previously observed growth effect of riboflavin on

Faecalibacterium prausnitzii[28–30]. Finally, to check the stability of METRIC, we applied METRIC

iteratively to the test data. The predictions of METRIC converge quickly in less than three

iterations, represented by saturating ρ of nutrient profiles between two neighboring iterations

(Supplementary Fig. 3).

Then we applied METRIC to the second dataset MLVS (Men's Lifestyle Validation

Study)[31,32]. Specifically, we utilized the composition of gut microbiomes and the one-day dietary

assessment of the 7-day dietary records (7DDRs). The 7DDRs are widely recognized to be the

most reliable estimation of dietary intake because participants were required to measure and

report gram weights for foods before and after eating to compute actual intake[33]. To guarantee

the usefulness of gut microbial composition in correcting dietary assessment, we need to find

paired microbial compositions and dietary assessments with matching dates. In MLVS, a total

of 599 paired samples with matching dates were found. Similarly, as we did for MCTs dataset,

we regarded the nutrient profile derived from 7DDRs as the "true" nutrient profile and added

varying levels of Gaussian noise to it as the "assessed" nutrient profile. Then we trained

METRIC on 80% of the data and tested on the remaining 20%. Consistent with our previous

findings, the trained METRIC exhibits an ability to correct the nutrient profile (Fig. 5a),

especially for large σ. For the case of $\sigma = 1.0$, the mean correction performance ($\overline{\rho_c} - \overline{\rho_a}$) =

0.078 (Fig. 5e). Across all nutrients, dietary fiber is a strongly corrected nutrient (Figs. 5f-g).

Finally, we applied METRIC to the third real-life dataset, WE-MACNUTR (Westlake N-

of-1 Trials for Macronutrient Intake)[34]. WE-MACNUTR is a dietary intervention study that

implemented a 'complete feeding' strategy, providing three isocaloric meals per day to 28

participants over a span of 72 days. Each participant completed high-fat, low-carbohydrate

(HF-LC) and low-fat, high-carbohydrate (LF-HC) diets in a randomized sequence, with a 6-day

wash-out period between them. Since the diets were completely controlled and well-

documented, the nutrient profile derived from this dataset closely reflects the true nutrient

profile. In WE-MACNUTR, we found 317 paired samples with both microbial compositions and

nutrient profiles. Considering the nutrient profile from the complete feeding as the true nutrient profile, we introduced varying levels of noise (Gaussian noise $N(0, \sigma^2)$ with different standard deviations σ) to create the assessed nutrient profile. Like our earlier results, METRIC can correct the nutrient profile when σ is large (Fig. 6a). When $\sigma = 1.0$, the mean correction performance $(\overline{\rho_c} - \overline{\rho_a}) = 0.125$ (Fig. 6e) and dietary fiber again exhibits a substantial correction (Figs. 6f-g).

Discussion

We presented a deep-learning method METRIC to correct random measurement errors in the nutrient profile. The method relies on the assessed nutrient profile and microbial composition to learn how to infer the true nutrient profile. First, we validated its performance on synthetic data generated by MiCRM where we directly modeled true and assessed nutrient profiles. Then we applied METRIC to three distinct real clinical datasets with added noise to the nutrient profile and found that it can correct the nutrient profile amount very well, especially for nutrients with large errors or metabolized by gut microbes. Based on our results, we believe it is possible to correct the nutrient profile using the microbial composition.

Similar to the class of computational methods that denoise images even without clean targets[18,19], METRIC offers a significant advantage by being capable of correcting the random measurement error without using the true nutrient profiles during the training. This attribute makes our method particularly useful in real-life scenarios where only assessed nutrient profiles are available, without access to true nutrient profiles. Yet, denoising methods without using clean targets such as Noise2Noise[18] cannot remove noise with a non-zero mean. Similarly, METRIC does have a limitation in that it cannot correct the systematic bias/error with a non-zero mean. Effectively correcting the systematic bias requires both assessed and true nutrient profiles to discern the consistent deviation between them. If both types of nutrient profiles are available, it becomes feasible to design a regression model that predicts true nutrient profiles based on assessed nutrient profiles, thereby achieving the task of removing the systematic bias.

In the future, some studies can be designed to improve the accuracy of dietary intake than self-reported approaches. For instance, digital documentation of meals through taking photos and direct observation by professionals can be leveraged to reduce measurement errors. Such data with more accurate dietary documentation may enable us to better investigate the feasibility of correcting nutrient profiles. Alternatively, some researchers made

234   some preliminary attempts to use nutritional biomarkers such as DNA barcodes for plants[35]

235   and metabolite biomarkers[36,24] for assessing food intake.

236   A more accurate reflection of nutrient consumption may be achieved by using more

objective measurements of microbial composition or metabolomic profile, instead of relying on

self-reported dietary recall. Currently, an active research direction in the field of precision

nutrition is to find microbial and metabolite biomarkers to help us to achieve a more complete

understanding of nutrient intakes. There are several attempts to predict the presence of food

items based on fecal bacteria and metabolites[23,24]. However, they were limited to several food

items and no connection to nutrient profiles was examined.

Methods

Datasets. The first dataset we used comes from a study that investigated the association

between diet and microbiome[27]. The study has paired 24-h food records and fecal shotgun

metagenomes from 34 healthy human subjects collected daily over 17 days, with 210 paired

samples in total. The second dataset is from MLVS[31,32] with 599 paired gut microbial

compositions and 7DDRs. The third dataset is from WE-MACNUTR[34], a dietary intervention

study that provides three meals per day for all participants. The study involved 30 participants,

with 2 participants withdrawing from the trial early and not included in the final analysis. It has

317 paired gut microbial compositions and true nutrient profiles of complete feeding with

matching dates. For all machine learning tasks, the same five random 80/20 train-test splits

were utilized to guarantee a fair comparison of methods.

The generation of nutrient profiles based on dietary assessments. The nutrient profiles

generated from dietary assessments involve several steps and rely on comprehensive food

composition databases such as USDA's Food and Nutrient Database for Dietary Studies

(FNDDS)[37] or Harvard Food Composition Database (HFDB)[38], etc. Each food item consumed

is matched with a corresponding item in a food composition database. These databases

provide detailed information on the nutrient content of foods, including macronutrients (like fats,

proteins, carbohydrates) and micronutrients (like vitamins and minerals). By multiplying the

amount of each food consumed by its nutrient content, the total intake of each nutrient is

calculated.

The generation of synthetic data. Similar to the traditional consumer-resource model[39,40]

and MiCRM (Microbial Consumer-Resource Model)[26], we simulated the process of nutrient

consumption by microbes and the following microbial growth. For simplicity, we assumed a

global pool with 20 microbial species and 20 nutrients in total. To mimic the difference in
nutrient consumption profiles across individuals, we randomly assigned flux for each sample:
to create a unique sample, the supply rate of each nutrient is randomly drawn from a uniform
distribution $\mathcal{U}[0,1]$. We recorded the assigned supply rates of all nutrients as the true nutrient
profile. The Gaussian noise with a mean of 0 and standard deviation of 0.5 is added to true
nutrient profile to create assessed nutrient profiles with measurement errors. After the
assignment of nutrient fluxes, we simulated the community assembly dynamics until the
system reaches a steady state. The steady-state microbial relative abundances are recorded
as the microbial composition for the sample. Then we repeated this procedure of generating
samples 250 times, forming the synthetic dataset we used in this study. The 250 samples are
split in the 80/20 ratio as the training/test set.

The overall population dynamics for the concentration of nutrient N_α and abundance of
microbial species M_i can be written as follows:

280
$$\frac{dN_\alpha}{dt} = -\sum_i a_{i\alpha} M_i N_\alpha - \delta N_\alpha,$$

281
$$\frac{dM_i}{dt} = \frac{\sum_\gamma a_{i\gamma} M_i N_\gamma}{Y} - \delta M_i,$$

where $a_{i\alpha}$ is the consumption rate of nutrient α by the species i, δ is the dilution rate, and Y is
the yield. For simplicity, we assumed $\delta = 0.1$ and $Y = 1$. The $a_{i\alpha}$ is assumed to be non-zero
with a probability of 50%. If $a_{i\alpha}$ is non-zero, its value is drawn from the uniform distribution
$\mathcal{U}[0,10]$. Eventually, each $a_{i\alpha}$ is divided by the number of nutrients that can be consumed by
the species i, avoiding the outgrowth of generalists.

METRIC. The core of METRIC is the MLP (Multilayer Perceptrons).

- Data processing: The CLR (Centered Log-Ratio) transformation is applied to microbial
relative abundances, and the log transformation is applied to the nutrient profiles.

- Model detail: METRIC has 3 hidden layers in the middle, sandwiched by input and
output variables. Each hidden layer has a fixed hidden layer dimension of 256. A skip
connection from corrupted nutrient profiles is introduced to add to the final layer of MLP.
Specifically, the final prediction is the sum of (1) the corrupted input from the skip
connection multiplied by a weight parameter α and (2) the final layer of MLP multiplied
by $(1 - \alpha)$. The optimal value for α is chosen based on the five-fold cross-validation
results on the training set.

- Training method: The Adam optimizer[41] is used for the gradient descent. The training
stops when the mean SCC (Spearman Correlation Coefficient) of nutrients $\bar{\rho}$ on the
validation/test set starts to decrease within the past 10 epochs.

•   Activation function: ReLU (Rectified Linear Unit).

Statistics. To calculate correlation throughout the study, we used Spearman's correlation

coefficient. Wherever we used *P* values, we explained in the Methods how we calculated them,

since for all such measurements in the study, we calculated the associated null distributions

from scratch. All statistical tests were performed using standard numerical and scientific

computing libraries in the Python programming language (version 3.7.1) and Jupyter Notebook

(version 6.1).

Data and code availability. All code for simulations used in this manuscript can be found at

https://github.com/wt1005203/METRIC.

Acknowledgements. We acknowledge grants from the National Institutes of Health

(R01AI141529, R01HD093761, RF1AG067744, UH3OD023268, U19AI095219,

U01HL089856, U01-152905, and U01-167552). We thank Walter Willett, Eric Rimm, and

Lorelei Mucci for valuable discussions.

Author contributions. T.W. and Y.-Y.L. designed the project. T.W. performed all the

numerical calculations and data analysis. T.W. processed the real data with assistance from

Y.Fu. All authors analyzed the results. T.W. and Y.-Y.L. wrote the paper. All authors edited

and approved the paper.

Correspondence. Correspondence and requests for materials should be addressed to Y.-Y.

L. (email: yyl@channing.harvard.edu).

Competing Interests. The authors declare no competing interests.

References

1.   Afshin, A. *et al.* Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet* **393**, 1958–1972 (2019).

2.   Mozaffarian, D. *et al.* Global Sodium Consumption and Death from Cardiovascular Causes. *New England Journal of Medicine* **371**, 624–634 (2014).

3.   Donaldson, M. S. Nutrition and cancer: A review of the evidence for an anti-cancer diet. *Nutrition Journal* **3**, 19 (2004).

4.   DiNicolantonio, J. J., Lucan, S. C. & O'Keefe, J. H. The Evidence for Saturated Fat and for Sugar Related to Coronary Heart Disease. *Progress in Cardiovascular Diseases* **58**, 464–472 (2016).

5.   Mensink, R. P., Zock, P. L., Kester, A. D. & Katan, M. B. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials. *The American Journal of Clinical Nutrition* **77**, 1146–1155 (2003).

6.   Hu, F. B. & Willett, W. C. Optimal Diets for Prevention of Coronary Heart Disease. *JAMA* **288**, 2569–2578 (2002).

7.   McNutt, S., Zimmerman, T. P. & Hull, S. G. Development of food composition databases for food frequency questionnaires (FFQ). *Journal of Food Composition and Analysis* **21**, S20–S26 (2008).

8.   Sharpe, I. *et al.* Automated Self-Administered 24-H Dietary Assessment Tool (ASA24) recalls for parent proxy-reporting of children's intake (> 4 years of age): a feasibility study. *Pilot and Feasibility Studies* **7**, 123 (2021).

9.   Hebert, J. R. *et al.* Development and testing of a seven-day dietary recall. *Journal of Clinical Epidemiology* **50**, 925–937 (1997).

10.   Westerterp, K. R. & Goris, A. H. C. Validity of the assessment of dietary intake: problems of misreporting. *Current Opinion in Clinical Nutrition & Metabolic Care* **5**, 489 (2002).

11.   Poslusna, K., Ruprich, J., Vries, J. H. M. de, Jakubikova, M. & Veer, P. van't. Misreporting of energy and micronutrient intake estimated by food records and 24 hour recalls, control and adjustment methods in practice. *British Journal of Nutrition* **101**, S73–S85 (2009).

12.   Dao, M. C. *et al.* Dietary assessment toolkits: an overview. *Public Health Nutrition* **22**, 404–418 (2019).

13.   Ravelli, M. N. & Schoeller, D. A. Traditional Self-Reported Dietary Instruments Are Prone to Inaccuracies and New Approaches Are Needed. *Frontiers in Nutrition* **7**, (2020).

14.   Gibson, R. S., Charrondiere, U. R. & Bell, W. Measurement Errors in Dietary Assessment Using Self-Reported 24-Hour Recalls in Low-Income Countries and Strategies for Their Prevention. *Advances in Nutrition* **8**, 980–991 (2017).

15.   Lissner, L. *et al.* OPEN about obesity: recovery biomarkers, dietary reporting errors and BMI. *Int J Obes* **31**, 956–961 (2007).

16.   Schmidt, U. & Roth, S. Shrinkage Fields for Effective Image Restoration. in 2774–2781 (2014).

17.   Chen, Y. & Pock, T. Trainable Nonlinear Reaction Diffusion: A Flexible Framework for Fast and Effective Image Restoration. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **39**, 1256–1272 (2017).

18.   Lehtinen, J. *et al.* Noise2Noise: Learning Image Restoration without Clean Data. Preprint at https://doi.org/10.48550/arXiv.1803.04189 (2018).

19.   Krull, A., Buchholz, T.-O. & Jug, F. Noise2Void - Learning Denoising From Single Noisy Images. in 2129–2137 (2019).

376 20.   Donia, M. S. & Fischbach, M. A. Small molecules from the human microbiota. *Science* **349**, 1254766 (2015).
377
21.   Koh, A., De Vadder, F., Kovatcheva-Datchary, P. & Bäckhed, F. From Dietary Fiber to
Host Physiology: Short-Chain Fatty Acids as Key Bacterial Metabolites. *Cell* **165**, 1332–1345
(2016).
22.   Santhiravel, S. *et al.* The Impact of Plant Phytochemicals on the Gut Microbiota of
Humans for a Balanced Life. *International Journal of Molecular Sciences* **23**, 8124 (2022).
23.   Shinn, L. M. *et al.* Fecal Bacteria as Biomarkers for Predicting Food Intake in Healthy
Adults. *The Journal of Nutrition* **151**, 423–433 (2021).
24.   Shinn, L. M. *et al.* Fecal Metabolites as Biomarkers for Predicting Food Intake by
Healthy Adults. *The Journal of Nutrition* nxad95 (2022) doi:10.1093/jn/nxad95.
25.   He, K., Zhang, X., Ren, S. & Sun, J. Deep Residual Learning for Image Recognition.
in *2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)* 770–778
(IEEE, 2016). doi:10.1109/CVPR.2016.90.
26.   Marsland, R. *et al.* Available energy fluxes drive a transition in the diversity, stability,
and functional structure of microbial communities. *PLoS Comput Biol* **15**, e1006793 (2019).
27.   Johnson, A. J. *et al.* Daily Sampling Reveals Personalized Diet-Microbiome
Associations in Humans. *Cell Host & Microbe* **25**, 789-802.e5 (2019).
28.   Clifford, M. N. Diet-Derived Phenols in Plasma and Tissues and their Implications for
Health. *Planta Med* **70**, 1103–1114 (2004).
29.   Rigottier-Gois, L. Dysbiosis in inflammatory bowel diseases: the oxygen hypothesis.
ISME J **7**, 1256–1261 (2013).
30.   Steinert, R. E., Sadaghian Sadabad, M., Harmsen, H. J. M. & Weber, P. The prebiotic
concept and human health: a changing landscape with riboflavin as a novel prebiotic
candidate? *Eur J Clin Nutr* **70**, 1348–1353 (2016).
31.   Mehta, R. S. *et al.* Stability of the human faecal microbiome in a cohort of adult men.
Nat Microbiol **3**, 347–355 (2018).
32.   Li, J. *et al.* Interplay between diet and gut microbiome, and circulating concentrations
of trimethylamine N-oxide: findings from a longitudinal cohort of US men. *Gut* **71**, 724–733
(2022).
33.   Yue, Y. *et al.* Reproducibility and validity of diet quality scores derived from food-
frequency questionnaires. *The American Journal of Clinical Nutrition* **115**, 843–853 (2022).
34.   Ma, Y. *et al.* Individual Postprandial Glycemic Responses to Diet in n-of-1 Trials:
Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR). *The Journal of Nutrition*
151, 3158–3167 (2021).
35.   Reese, A. T. *et al.* Using DNA Metabarcoding To Evaluate the Plant Component of
Human Diets: a Proof of Concept. *mSystems* **4**, e00458-19 (2019).
36.   Guasch-Ferré, M., Bhupathiraju, S. N. & Hu, F. B. Use of Metabolomics in Improving
Assessment of Dietary Intake. *Clinical Chemistry* **64**, 82–98 (2018).
37.   FNDDS : USDA ARS. https://www.ars.usda.gov/northeast-area/beltsville-md-
bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/fndds/#.
38.   Colditz, G. A., Manson, J. E. & Hankinson, S. E. The Nurses' Health Study: 20-Year
Contribution to the Understanding of Health Among Women. *Journal of Women's Health* **6**,
49–62 (1997).
39.   MacArthur, R. Species packing and competitive equilibrium for many species.
Theoretical Population Biology **1**, 1–11 (1970).
40.   Chesson, P. MacArthur's consumer-resource model. *Theoretical Population Biology*
37, 26–38 (1990).
41.   Kingma, D. P. & Ba, J. Adam: A Method for Stochastic Optimization. Preprint at
https://doi.org/10.48550/arXiv.1412.6980 (2017).
426

Figure 1: **The idea of inferring the true nutrient profile based on the assessed nutrient**
profile and microbial composition. a, The typical dietary assessment such as ASA24
(Automated Self-Administered 24-hour Dietary Assessment Tool) often induces a
measurement error in the food intake due to unreliable mental recall. Such a random
measurement error in the food intake is carried to the nutrient profile when the food profile is
converted to the nutrient profile. **b,** Our goal in this study is to remove the measurement error.
More specifically, we would like to infer true nutrient profiles based on assessed nutrient
profiles and microbial compositions.

Figure 2: **The architecture and workflow of METRIC to infer true nutrient profiles.** For simplicity, we used a hypothetical example with n=3 training samples and 2 samples in the test set. For each sample, there are N_s microbial species and N_n nutrients. Across panels, microbial species and their relative abundances are colored blue. Nutrients and their amounts are colored red. The corrupted nutrient profiles are created by adding different types of random noise (i.e., Gaussian, Uniform, etc.) to the assessed nutrient profiles. Icons associated with assessed/corrupted nutrient profiles are bounded by solid black/dashed lines. Icons associated with true nutrient profiles are bounded by solid green lines. **a,** During the training of METRIC, the method takes corrupted nutrient profiles and microbial compositions as the input and learns to infer assessed nutrient profiles. **b,** Similar to MLP (MultiLayer Perceptrons), METRIC has several hidden layers in the middle. The skip connection provides the corrupted nutrient profile directly to one layer before the final output, enabling it to skip the propagation through the hidden layers. The skipped corrupted nutrient profile multiplied by the weight parameter α and the final hidden layer (the bottom grey nodes) multiplied by $(1 - \alpha)$ add up as the final output (the bottom red nodes). **c,** The well-trained METRIC is applied to the test set to generate predictions for nutrient profiles whose values are compared to true nutrient profiles.

Figure 3: **METRIC can correct the measurement error in assessed nutrient profiles on**
synthetic data from MiCRM[26]**.** The Spearman's Rank Correlation Coefficient ρ is adopted to
evaluate the correlation across various types of nutrient profiles. All corrected/true values
shown are the log of nutrient concentrations. **a,** ρ_c (i.e., ρ between corrected and true values)
and ρ_a (i.e., ρ between assessed and true values) decrease as the standard deviation of
added Gaussian noise σ increases. All following panels focus on the case of σ=1.5. **b,** The
correlation between assessed values and true values of log concentrations of one nutrient
among different samples. **c,** The correlation between corrected values (predictions of METRIC)
and true values of log concentrations of the same nutrient among different samples. Similar
comparisons for another two nutrients are shown in **d-e** and **f-g**. **e,** The correction performance
of all nutrients is measured by ($\rho_c - \rho_a$).

Figure 4: **METRIC can correct the measurement error in assessed nutrient profiles on**
real data from MCTs[27]**.** The Spearman's Rank Correlation Coefficient ρ is adopted to evaluate
the correlation across various types of nutrient profiles. All nutrient concentrations are in the
unit of grams. All corrected/true values shown are the log of nutrient concentrations. **a,** ρ_c (i.e.,
ρ between corrected and true values) and ρ_a (i.e., ρ between assessed and true values)
decrease as the standard deviation of added Gaussian noise σ increases. All following panels
focus on the case of σ=1.0. **b,** The correlation between assessed values and true values of log
concentrations of carotene among different samples. **c,** The correlation between corrected
values (predictions of METRIC) and true values of log concentrations of carotene among
different samples. **d-e,** The similar comparison for octadecaonoic acid shows a modest
correction. **f-g,** The similar comparison for fiber shows a strong correction. **e,** The correction
performance for all nutrients is measured by $(\rho_c - \rho_a)$.

482
Figure 5: **METRIC can correct the measurement error in assessed nutrient profiles from**
MLVS[31,32]**.** The Spearman's Rank Correlation Coefficient ρ is adopted to evaluate the
correlation across various types of nutrient profiles. All nutrient concentrations are in the unit
of grams. All corrected/true values shown are the log of nutrient concentrations. **a,** ρ_c (i.e., ρ
between corrected and true values) and ρ_a (i.e., ρ between assessed and true values)
decrease as the standard deviation of added Gaussian noise σ increases. All following panels
focus on the case of σ=1.0. **b,** The correlation between assessed values and true values of log
concentrations of fructose among different samples. **c,** The correlation between corrected
values (predictions of METRIC) and true values of log concentrations of fructose among
different samples. **d-e,** The similar comparison for monounsaturated fatty acids shows a
modest correction. **f-g,** The similar comparison for dietary fiber shows a strong correction. **e,**
The correction performance for all nutrients is measured by ($\rho_c - \rho_a$). Nutrient names are not
added due to lack of space.

496
Figure 6: **METRIC can correct the measurement error in assessed nutrient profiles from**
WE-MACNUTR[34]**.** The Spearman's Rank Correlation Coefficient ρ is adopted to evaluate the

correlation across various types of nutrient profiles. **a,** ρ_c (i.e., ρ between corrected and true
values) and ρ_a (i.e., ρ between assessed and true values) decrease as the standard deviation
of added Gaussian noise σ increases. All nutrient concentrations are in the unit of grams. All
corrected/true values shown are the log of nutrient concentrations. All following panels focus
on the case of σ=1.0. **b,** The correlation between assessed values and true values of log
concentrations of magnesium among different samples. **c,** The correlation between corrected
values (predictions of METRIC) and true values of log concentrations of magnesium among
different samples. **d-e,** The similar comparison for threonine shows a modest correction. **f-g,**
The similar comparison for dietary fiber shows a strong correction. **e,** The correction
performance for all nutrients is measured by $(\rho_c - \rho_a)$. Nutrient names are not added due to
lack of space.